

New models and Big Bang Nucleosynthesis constraints in $f(Q)$ gravity

Fotios K. Anagnostopoulos,^{1,*} Viktor Gakis,^{2,†} Emmanuel N. Saridakis,^{3,4,5,‡} and Spyros Basilakos^{3,6,7,§}

¹*Department of Informatics and Telecommunications,
University of Peloponnese, Karaiskaki 70, 22100, Tripoli, Greece*

²*Department of Physics, National Technical University of Athens, Zografou Campus GR 157 73, Athens, Greece*

³*National Observatory of Athens, Lofos Nymfon, 11852 Athens, Greece*

⁴*Department of Astronomy, School of Physical Sciences,
University of Science and Technology of China, Hefei, Anhui 230026, China*

⁵*CAS Key Laboratory for Research in Galaxies and Cosmology,*

University of Science and Technology of China, Hefei, Anhui 230026, China

⁶*Academy of Athens, Research Center for Astronomy and Applied Mathematics, Soranou Efessiou 4, 11527, Athens, Greece*

⁷*School of Sciences, European University Cyprus, Diogenes Street, Engomi, 1516 Nicosia, Cyprus*

The $f(Q)$ theories of modified gravity arise from the consideration of non-metricity as the basic geometric quantity, and have been proven to be very efficient in describing the late-time Universe. We use the Big Bang Nucleosynthesis (BBN) formalism and observations in order to extract constraints on various classes of $f(Q)$ models. In particular, we calculate the deviations that $f(Q)$ terms bring on the freeze-out temperature T_f in comparison to that of the standard Λ CDM evolution, and then we impose the observational bound on $\left| \frac{\delta T_f}{T_f} \right|$ to extract constraints on the involved parameters of the considered models. Concerning the polynomial model, we show that the exponent parameter should be negative, while for the power-exponential model and the new hyperbolic tangent - power model we find that they pass the BBN constraints trivially. Finally, we examine two DGP-like $f(Q)$ models, and we extract the bounds on their model parameters. Since many gravitational modifications, although able to describe the late-time evolution of the Universe, produce too-much modification at early times and thus fall to pass the BBN confrontation, the fact that $f(Q)$ gravity can safely pass the BBN constraints is an important advantage of this modified gravity class.

I. INTRODUCTION

The accelerated expansion of the late Universe is commonly attributed to the existence of an extra cosmic constituent known collectively as “Dark Energy”. In the context of the concordance Λ CDM scenario, dark energy is just the standard cosmological constant and this scenario is very successful in interpreting observations such as Cosmic Microwave Background (CMB) [1], Supernovae Ia (SNIa) [2], Large Scale Structure (LSS) [3], etc.

However, recently various observational issues have gain attention, for example the Hubble constant tension, that is the discrepancy between the value of the Hubble constant from the CMB and the corresponding one from Cepheids up to 5σ (see [4] for a thorough and recent review). Another example is the difference of the root-mean-square amplitude of matter over-density perturbation σ_8 , between the value extracted from the CMB (imposing Λ CDM) and the corresponding value extracted by fitting on the Large Scale power spectrum [5]. Furthermore, the value of the cosmological constant in the late Universe seems to be inconsistent with the corresponding value at the early Universe [6].

On the other hand, the current gravitational theory, namely General Relativity (GR), cannot be re-

normalized and thus it is difficult to be consistent with a quantum description [7]. Hence, a large portion of the literature follows the direction of modified gravity, which seems promising to solve the latter problem [8], and in addition it is very efficient in describing the two phases of accelerated expansion of the universe [9, 10]. Finally, note that modified gravity can be efficient in alleviating the two aforementioned tensions of Λ CDM cosmology, namely the H_0 and the σ_8 ones [11].

Along these lines and due to the fact that the Einstein-Hilbert Lagrangian is just one of many possible Lagrangians that could lead to the same field equations, the community has developed many frameworks of modified gravity. A class of gravitational modification arises from the extension of the Einstein-Hilbert Lagrangian with extra terms, however maintaining its geometrical structure, i.e. its Riemannian formulation, resulting for instance in $f(R)$ gravity [12], in Gauss-Bonnet and $f(G)$ gravity [13, 14], in Weyl gravity [15], etc. A more radical approach to follow is to modify the building blocks of General Relativity, that is the underlying geometrical structure. An instance of this kind of modification arises from the torsional formulation of gravity, namely the modified teleparallel theories, such as the $f(T)$ gravity [16, 17], the $f(T, T_G)$ gravity [18], the scalar-torsion theories [19, 20] etc. In contrast to the standard GR and the corresponding $f(R)$ extensions, which are based on curvature, in modified teleparallel formulation gravity is manifested via the torsion tensor, with curvature being zero.

There is yet another possibility, which is formulated us-

* fotis-anagnostopoulos@hotmail.com

† vgakis@central.ntua.gr

‡ msaridak@phys.uoa.gr

§ svasil@academyofathens.gr

ing the non-metricity Q of the connection. In this setup, only the non-metricity is non-zero while curvature and torsion both vanish. Hence, non-metricity can be used in order to describe gravity through geometry, leading to the so-called Symmetric Teleparallel Equivalent to General Relativity (STEGR) [21]. It is interesting to note that although Teleparallel Equivalent of General Relativity (TEGR) as well as STEGR, are completely equivalent to General Relativity at the level of equations, when we modify them the resulting theories $f(R)$, $f(T)$ and $f(Q)$ are different theories. These types of modifications have in general more degrees of freedom than the two of standard GR, unless specific constraints are imposed in the functional forms. Thus, during the last years $f(Q)$ gravity, its application to cosmology and generalizations has attracted the interest of the literature [22–42].

A very strong constraint on modified gravity models in the early Universe comes from the era of Big Bang Nucleosynthesis (BBN) [43–48]. In the present work we are interested in imposing BBN constraints on $f(Q)$ gravity, within an analytical approach, following [49–53]. Specifically, we use observational data on primordial abundances of ${}^4\text{He}$ to constrain all known $f(Q)$ models along some that are proposed for the first time inspired by the successful $f(T)$ models. The plan of the manuscript is the following. In Section II we provide a formal description of Symmetric Teleparallel gravity and its $f(Q)$ extension, while in Section III we present the cosmological solutions of the theory alongside the specific $f(Q)$ models that are going to be studied. In Section IV we briefly review the formalism of BBN constraints, and then in Section V we apply it in the framework of $f(Q)$ gravity. Finally, in Section VI we derive our conclusions and we point out further directions.

II. SYMMETRIC TELEPARALLEL GRAVITY

General Relativity is built upon Lorentzian Geometry which is specified by choosing a connection that is symmetric and metric compatible. The latter is the Levi-Civita connection and due to its properties it only produces non-zero curvature, whilst torsion and non-metricity are zero [54]. Nevertheless, a different kind of connection can be utilized when using geometrodynamics as the mathematical framework for gravity. As a matter of fact the most general connection is called metric-affine it is described by the formula

$$\Gamma^\alpha{}_{\mu\nu} = \overset{\circ}{\Gamma}{}^\alpha{}_{\mu\nu} + K^\alpha{}_{\mu\nu} + L^\alpha{}_{\mu\nu}, \quad (1)$$

where $\overset{\circ}{\Gamma}{}^\alpha{}_{\mu\nu}$ represents the Christoffel symbols of the Levi-Civita connection, $K^\alpha{}_{\mu\nu}$ is the contorsion tensor and $L^\alpha{}_{\mu\nu}$ is the *disformation* tensor related to the non-metricity [55]

$$Q_{\alpha\mu\nu} := \nabla_\alpha g_{\mu\nu}. \quad (2)$$

The disformation tensor can then be expanded as [21]

$$L^\alpha{}_{\mu\nu} := \frac{1}{2}g^{\alpha\beta}(-Q_{\mu\beta\nu} - Q_{\nu\beta\mu} + Q_{\beta\mu\nu}), \quad (3)$$

and encapsulates all the information coming from the non-metricity tensor of the general affine connection. In summary, these three types of geometric deformation $\overset{\circ}{\Gamma}{}^\alpha{}_{\mu\nu}$, $K^\alpha{}_{\mu\nu}$ and $L^\alpha{}_{\mu\nu}$ form a kind of “trinity of gravity”, which encompasses all components of the general connection $\Gamma^\alpha{}_{\mu\nu}$ [56]. This practically implies that it is possible to express the geometry of a gravitational theory in either curvature, non-metricity or torsion. A connection which admits only non-metricity, whilst curvature and torsion are zero, is called Symmetric Teleparallel Gravity (STG) connection, while connections which have only zero curvature are called teleparallel connections [57, 58].

Similarly to TEGR, which is based on a connection which has only torsion [59], the STG connection has only non-metricity and hence STEGR is equivalent to GR at the level of field equations, since its action differs by a boundary term from the Einstein-Hilbert. In particular, defining the non-metricity scalar as [21]

$$Q := -\frac{1}{4}Q_{\alpha\mu\nu}Q^{\alpha\mu\nu} + \frac{1}{2}Q_{\alpha\mu\nu}Q^{\mu\alpha\nu} + \frac{1}{4}Q_\alpha Q^\alpha - \frac{1}{2}\bar{Q}_\alpha Q^\alpha, \quad (4)$$

with

$$Q_\mu := Q_\mu{}^\alpha{}_\alpha, \quad \bar{Q}_\mu := Q_\alpha{}^\alpha{}_\mu, \quad (5)$$

one has the relation

$$\overset{\circ}{R} = Q + B, \quad (6)$$

where $\overset{\circ}{R}$ is the Ricci scalar of the Levi-Civita connection, and the boundary term is

$$B := \overset{\circ}{\nabla}_\alpha(\bar{Q}^\alpha - Q^\alpha). \quad (7)$$

Therefore, General Relativity, which uses $\overset{\circ}{R}$ as the Lagrangian density, and STEGR, which uses Q , lead to exactly the same equations.

One can extend STEGR by extending the Lagrangian to an arbitrary function, resulting to $f(Q)$ gravity, with action [21, 60]

$$S_G = \int d^4x \sqrt{-g} \left[-\frac{1}{16\pi G} f(Q) + \mathcal{L}_m \right], \quad (8)$$

where G is the gravitational constant and \mathcal{L}_m the matter Lagrangian. The corresponding field equations are

$$\begin{aligned} \sqrt{-g} \left(\frac{1}{2} f_{g^{\mu\nu}} - \frac{\partial f}{\partial g^{\mu\nu}} \right) - 2\nabla_\alpha (\sqrt{-g} P^\alpha{}_{\mu\nu}) \\ = 8\pi G \sqrt{-g} T_{\mu\nu}, \end{aligned} \quad (9)$$

with

$$\begin{aligned}
\frac{\partial f}{\partial g^{\mu\nu}} &= -\frac{f_Q}{\sqrt{-g}} \left(\frac{\partial(\sqrt{-g}Q)}{\partial g^{\mu\nu}} - \frac{1}{2}\sqrt{-g}Qg_{\mu\nu} \right) \\
&= -f_Q \left[c_1 (2Q_{\alpha\beta\mu}Q^{\alpha\beta}{}_{\nu} - Q_{\mu\alpha\beta}Q^{\nu}{}_{\alpha\beta}) \right. \\
&\quad + c_2 Q_{\alpha\beta\mu}Q^{\beta\alpha}{}_{\nu} + c_3 (2Q_{\alpha}Q^{\alpha}{}_{\mu\nu} - Q_{\mu}Q_{\nu}) \\
&\quad \left. + c_4 \tilde{Q}_{\mu}\tilde{Q}_{\nu} + c_5 \tilde{Q}_{\mu}Q^{\alpha}{}_{\mu\nu} \right], \quad (10)
\end{aligned}$$

and where the conjugate to $f(Q)$ is defined as [21]

$$\begin{aligned}
P^{\alpha}{}_{\mu\nu} &:= \frac{1}{2\sqrt{-g}} \frac{\partial(\sqrt{-g}f(Q))}{\partial Q_{\alpha}{}^{\mu\nu}} \\
&= -f_Q \left\{ c_1 Q^{\alpha}{}_{\mu\nu} + c_2 Q_{(\mu}{}^{\alpha}{}_{\nu)} + c_3 g_{\mu\nu} Q^{\alpha} \right. \\
&\quad \left. + c_4 \delta^{\alpha}{}_{(\mu} \tilde{Q}_{\nu)} + \frac{c_5}{2} [\tilde{Q}^{\alpha} g_{\mu\nu} + \delta^{\alpha}{}_{(\mu} Q_{\nu)}] \right\}, \quad (11)
\end{aligned}$$

while as usual the matter energy momentum tensor is defined as

$$T_{\mu\nu} := -\frac{2}{\sqrt{-g}} \frac{\delta(\sqrt{-g}\mathcal{L}_m)}{\delta g^{\mu\nu}}. \quad (12)$$

The field equations (9) are quite general with respect to both their normalizations described by G and also the defined constants c_i of Q . For instance, by choosing Q as in Eq. (4) then we retrieve the standard modified class of $f(Q)$ gravity [21]. In the rest of this work we deal with this standard modified class of $f(Q)$ theories.

III. COSMOLOGICAL APPLICATION OF $f(Q)$ GRAVITY

In this section we proceed to the cosmological application of $f(Q)$ gravity. We impose a spatially flat Friedmann-Robertson-Walker (FLRW) metric of the form

$$ds^2 = -dt^2 + a^2(t)\delta_{ij}dx^i dx^j, \quad (13)$$

with $a(t)$ the scale factor. In this case Eq. (9) gives rise to the two Friedmann equations [22]

$$6f_Q H^2 - \frac{1}{2}f = 8\pi G(\rho_m + \rho_r), \quad (14)$$

$$(12H^2 f_{QQ} + f_Q)\dot{H} = -4\pi G(\rho_m + p_m + \rho_r + p_r), \quad (15)$$

where $H = \dot{a}/a$ is the the Hubble function and with ρ_m , ρ_r and p_m , p_r the energy densities and pressures of the matter and radiation perfect fluids respectively. Additionally, note that the non-metricity scalar Q in an FRW background becomes $Q = 6H^2$. Finally, the equations constitute a close system by considering the matter and radiation conservation equations, namely

$$\dot{\rho}_m + 3H(\rho_m + p_m) = 0, \quad (16)$$

$$\dot{\rho}_r + 3H(\rho_r + p_r) = 0. \quad (17)$$

We can re-write Eqs. (14)-(15) as

$$H^2 = \frac{8\pi G}{3} (\rho_m + \rho_r + \rho_{DE}), \quad (18)$$

$$\dot{H} = -4\pi G (\rho_m + p_m + \rho_r + p_r + \rho_{DE} + p_{DE}), \quad (19)$$

where we have defined the energy density and pressure of the effective dark energy sector as

$$\rho_{DE} = \frac{1}{16\pi G} [6H^2 (1 - 2f_Q) + f], \quad (20)$$

$$\begin{aligned}
p_{DE} &= \frac{1}{16\pi G} [4(f_Q - 1)\dot{H} - f \\
&\quad + 6H^2 (8f_{QQ}\dot{H} + 2f_Q - 1)]. \quad (21)
\end{aligned}$$

Finally, it proves convenient to introduce the density parameters for the various sectors us

$$\Omega_i = \frac{8\pi G\rho_i}{3H^2}, \quad (22)$$

where ‘‘i’’ stands for matter, radiation and dark energy.

Let us now focus on particular $f(Q)$ forms that are of interest for the late Universe description. In general, any model that exhibits identical Hubble rate with the concordance one (see [28]) can pass the BBN constraints trivially, hence we do not examine models of this kind. Note also that we consider versions of our models in their ‘‘bare’’ form, i.e. without explicitly containing the cosmological constant, in order to avoid re-introducing the cosmological constant problem and also to keep the smallest possible set of free parameters. One should in principle combine any selection of the following models, however the final model could have a large number of free parameters.

1. Polynomial model

The following model was introduced in [22]

$$f(Q) = Q - 6\lambda M^2 \left(\frac{Q}{6M^2} \right)^{\alpha}, \quad (23)$$

where λ and α are dimensionless parameters. The parameter M corresponds to a mass scale that should be of the order $\sqrt{\Lambda}$, with Λ the standard cosmological constant. The case of $\alpha = 0$ gives STEGR plus a cosmological constant equal to $6\lambda M^2$, while the case $\alpha = 1$ corresponds to STEGR with $G \rightarrow G/(1 - \lambda)$. More generally, $\alpha > 1$ is mostly relevant to early Universe, while $\alpha < 1$ is able to describe dark energy and thus it is relevant to late Universe. Moreover, in the latter case the existence of an asymptotic GR limit at early times is apparent. Finally, note that the case $\alpha = -1$ has been confronted with late universe observations in [34].

In this model for the effective dark-energy density (20) we obtain

$$\rho_{DE} = \frac{1}{16\pi G} [6^{-\alpha} \Omega_{F0} H_0^{-2\alpha} Q^{\alpha+1}], \quad (24)$$

where $\Omega_{F0} = 1 - \Omega_{m0} - \Omega_{r0}$, and with the subscript denoting the value of a quantity at present time. Calculating the first Friedmann (14) at present time we find that the free parameter λ can be eliminated in terms of α and M as

$$\lambda = \frac{1}{12\alpha + 6} (\Omega_{F0} H_0^{-2\alpha} M^{2\alpha-2}). \quad (25)$$

2. Power - Exponential model

This model has proposed in [41] and has been proven to provide slightly better fits to observational data than the concordance model [41]. It is characterized by

$$f(Q) = Q e^{\lambda \frac{Q_0}{Q}}. \quad (26)$$

As it is apparent, in case $\lambda = 0$ GR without a cosmological constant is recovered. In the past, where the term Q_0/Q decreases, since the Hubble function increases, the model at hand effectively reduces to the power-law model, thus alleviating the cosmological constant problem. In a sense, this model behaves as infinite number of different power-law models that coexist, and at each moment of the cosmic history one of them becomes dominant. Another advantage of the model is that it has the same number of free parameters with the concordance one, while additionally containing rich phenomenology.

In this case (20) gives

$$\rho_{DE} = \frac{1}{16\pi G} \left[Q - e^{\frac{\lambda Q_0}{Q}} (Q - 2\lambda Q_0) \right]. \quad (27)$$

The parameter λ is expressed from the first Friedmann Eq. (14) at present as

$$\lambda = \frac{1}{2} + W \left(\frac{\Omega_{F0} - 1}{2\sqrt{e}} \right), \quad (28)$$

where W is the Lambert function.

3. Log-square-root model

Let us now propose a new $f(Q)$ model. In particular, following the approach of [61], we introduce a logarithmic-square-root $f(Q)$ model as:

$$f(Q) = Q + nQ_0 \sqrt{\frac{Q}{\lambda Q_0}} \ln \left(\lambda \frac{Q_0}{Q} \right), \quad (29)$$

with n and $\lambda > 0$ the model parameters. The corresponding effective energy density (20) gives

$$\rho_{DE} = \frac{1}{16\pi G} \Omega_{F0} \sqrt{Q_0 Q}, \quad (30)$$

and the first Friedmann equation at present leads to

$$n = \frac{\Omega_{F0}}{2} \sqrt{\lambda}. \quad (31)$$

4. Hyperbolic tangent - power model

We consider the hyperbolic tangent power model, in similar lines with [62], namely we choose

$$f(Q) = Q + \lambda Q_0 \left(\frac{Q}{Q_0} \right)^n \tanh \left(\frac{Q_0}{Q} \right). \quad (32)$$

In this case we have

$$\rho_{DE} = \frac{1}{16\pi G} Q_0^{1-n} Q^{n-1} \cdot \left[(1-2n)Q \tanh \left(\frac{Q_0}{Q} \right) + 2Q_0 \operatorname{sech}^2 \left(\frac{Q_0}{Q} \right) \right], \quad (33)$$

while

$$n = \frac{1}{2} [-\coth(1)\Omega_{F0} + 1 + 4\operatorname{csch}(2)]. \quad (34)$$

5. DGP-like $f(Q)$ model - I

In the recent work [63] two new $f(Q)$ models were introduced, by requiring the extra terms in the modified Friedman equation to be of the form $\sim H$. It was argued that a term proportional to Q is responsible for the H^2 term, thus adding a term proportional to \sqrt{Q} leads to a term $\sim H$ in the Friedman equation. These two models resemble the Dvali-Gabadadze-Porrati cosmology [64] at the background level, and this is where they acquire their names from. We mention here that this resemblance is only at the background level, since the perturbations in the usual DGP model and $f(Q)$ gravity are fundamentally different, and hence the present scenario does not share the known problem of usual DGP model in fitting the perturbation-related data (such as LSS and CMB temperature/polarization) [65, 66].

The first model of this kind reads as

$$f(Q) = \alpha \sqrt{Q} \log Q + 2\beta Q, \quad (35)$$

where α, β are free parameters. We mention that in the original parametrization the free parameter α is not dimensionless, and we use the form (35) in order to maintain compatibility with the results of [63] (although it would be more convenient to impose a re-scaling $\alpha \rightarrow \sqrt{Q_0}\alpha$ to ensure that α is dimensionless). Nevertheless, we stress that inside the logarithm one should explicitly include a constant value of dimension H_0^{-2} , and for consistency in the present work we do apply this modification.

In this case (20) gives

$$\rho_{DE} = \frac{1}{16\pi G} [\sqrt{6}H_0\sqrt{Q}(2\beta + \Omega_{F0} - 1) - 2\beta Q + Q], \quad (36)$$

while the first Friedmann equation gives

$$\alpha = -\sqrt{\frac{3}{2}}H_0(2\beta + \Omega_{F0} - 1). \quad (37)$$

Finally, note that near the current time, the term $\sim H$ ceases to be negligible in comparison to H^2 , thus giving rise to new phenomenology. However, in the past this term is negligible and the model is very close to the concordance one.

6. DGP-like $f(Q)$ model - II

The second model introduced in [63] reads as

$$f(Q) = \frac{Q\sqrt{u(Q)}\left[\sqrt{u(Q)} - \sqrt{\gamma}\operatorname{arctanh}\left(\frac{\sqrt{u(Q)}}{\sqrt{\gamma}}\right)\right]}{8\pi G\sqrt{Qu(Q)}}, \quad (38)$$

where $u(Q) = \gamma + \beta^2 Q$, and β, γ are free parameters, different than zero. The particular choice of u is motivated from the previous case and includes also a mixing between H^2 and H terms [63]. In this case we acquire

$$\rho_{DE} = \frac{1}{16\pi G}\left[Q - \sqrt{Q}\sqrt{\gamma + \beta^2 Q}\right], \quad (39)$$

while

$$\gamma = -6H_0^2(\beta - \Omega_{F0} + 1)(\beta + \Omega_{F0} - 1). \quad (40)$$

Similarly to the previous case, in order to maintain dimensional consistency within the square root, we include an H_0^{-2} normalization (i.e. we apply the re-scaling $\gamma \rightarrow Q_0\gamma$).

IV. BIG BANG NUCLEOSYNTHESIS CONSTRAINTS

In this Section we review the Big Bang Nucleosynthesis (BBN) formalism following [43–46]. The BBN takes place during the radiation era and thus the energy density of relativistic particles needs to be taken into account, namely

$$\rho_r = \frac{\pi^2}{30}g_*T^4, \quad (41)$$

where

$$g_* \sim 10 \quad (42)$$

is the effective number of degrees of freedom and T is the temperature (for more details of the BBN framework used in this work see the appendix of [47]). The calculation of the neutron abundance is realized by taking into account the protons-neutron conversion rate

$$\lambda_{pn}(T) = \lambda_{(n+\nu_e \rightarrow p+e^-)} + \lambda_{(n+e^+ \rightarrow p+\bar{\nu}_e)} + \lambda_{(n \rightarrow p+e^-+\bar{\nu}_e)}, \quad (43)$$

and its inverse $\lambda_{np}(T)$, thus the total rate is

$$\lambda_{tot}(T) = 4AT^3(4!T^2 + 2 \times 3!QT + 2!Q^2), \quad (44)$$

where $Q = m_n - m_p = 1.29 \times 10^{-3}\text{GeV}$ is the neutron-proton mass difference and $A = 1.02 \times 10^{-11}\text{GeV}^{-4}$.

Regarding the primordial mass fraction of ${}^4\text{He}$, it can be estimated [44] as

$$Y_p := \lambda \frac{2x(t_f)}{1 + x(t_f)}, \quad (45)$$

with $\lambda = e^{-(t_n - t_f)/\tau}$, t_f the freeze-out time of the weak interactions, t_n the corresponding freeze-out time of nucleosynthesis, τ the neutron mean lifetime and $x(t_f) = e^{-Q/T(t_f)}$ the neutron-to-proton equilibrium ratio. The role of the function $\lambda(t_f)$ is to account for the fraction of neutrons that decay into protons during the time interval $t \in [t_f, t_n]$.

In case of modified gravity models, in general the Friedmann equations will contain extra terms than the standard GR ones. The BBN is realized in the radiation epoch and according to observations these extra contributions have to be small compared to the radiation sector in the Standard Model of particles physics in the framework of General Relativity, whilst we can safely neglect the matter sector as we are deep in the radiation era. Thus, the first Friedmann equation can be approximated as

$$H^2 \approx \frac{8\pi G}{3}\rho_r =: H_{GR}^2, \quad (46)$$

where the scale factor evolves as $a \sim t^{1/2}$, with t the cosmic time. Consequently, temperature and time are related by $\frac{1}{t} \simeq \left(\frac{32\pi^3 g_*}{90}\right)^{1/2} \frac{T^2}{M_P}$ (or $T(t) \simeq (t/\text{sec})^{-1/2}\text{MeV}$), which can further lead to

$$H \approx \left(\frac{4\pi^3 g_*}{45}\right)^{1/2} \frac{T^2}{M_P}, \quad (47)$$

where

$$M_P = (8\pi G)^{-1} = 1.22 \times 10^{19}\text{GeV}, \quad (48)$$

is the Planck mass.

Assuming that the expansion time is much smaller than the interaction time, then the interaction rate $\lambda_{tot}(T)$ given in (44) satisfies $\frac{1}{H} \ll \lambda_{tot}(T)$, which means that all processes can be approximated as being in thermal equilibrium [44, 67]. In contrast, if the particles do not have the necessary time intervals to interact then $\frac{1}{H} \gg \lambda_{tot}(T)$ and thus they decouple. The temperature at which the particles decouple is called freeze-out temperature and is denoted as T_f , and it is defined through

$H = \lambda_{tot}(T)|_{T=T_f}$. Using $H \approx \left(\frac{4\pi^3 g_*}{45}\right)^{1/2} \frac{T^2}{M_P}$, while $\lambda_{tot}(T) \approx qT^5$, with

$$q = 4A4! \simeq 9.8 \times 10^{-10}\text{GeV}^{-4}, \quad (49)$$

the freeze-out temperature is provided as

$$T_f = \left(\frac{4\pi^3 g_*}{45 M_p^2 q^2} \right)^{1/6} \sim 0.0006 \text{ GeV}. \quad (50)$$

In the realm of modified gravity, the Hubble function H will, in general, deviate from H_{GR} . This in turn means that the corresponding freeze-out temperatures will also deviate by an amount of δT_f . This deviation induces a difference δY_p of the fractional mass Y_p

$$\delta Y_p = Y_p \left[\left(1 - \frac{Y_p}{2\lambda} \right) \ln \left(\frac{2\lambda}{Y_p} - 1 \right) - \frac{2t_f}{\tau} \right] \frac{\delta T_f}{T_f}, \quad (51)$$

where $\delta T(t_n) = 0$ was imposed due to the fact that T_n is fixed by the binding energy of deuterium [49–53]. Thus, the observational imprints of the mass fraction Y_p of baryons that convert to ${}^4\text{He}$ during the BBN epoch, are [68–74]

$$Y_p = 0.2476, \quad |\delta Y_p| < 10^{-4}. \quad (52)$$

By replacing these into (51), the upper bound of $\frac{\delta T_f}{T_f}$ is obtained as

$$\left| \frac{\delta T_f}{T_f} \right| < 4.7 \times 10^{-4}, \quad (53)$$

which quantifies the allowed deviation from the cosmology of GR.

As we mentioned above, the effective dark energy ρ_{DE} of modified gravity is present during the BBN times too, and compared to ρ_r , ρ_{DE} should be smaller and thus it can be considered as a first-order deviation. Hence, the Hubble function is

$$H = H_{GR} \sqrt{1 + \frac{\rho_{DE}}{\rho_r}} = H_{GR} + \delta H, \quad (54)$$

where

$$\delta H = \left(\sqrt{1 + \frac{\rho_{DE}}{\rho_r}} - 1 \right) H_{GR}. \quad (55)$$

The deviation δH from standard H_{GR} will induce the deviation δT_f from T_f , and according to the relation $H_{GR} = \lambda_{tot} \approx q T^5$ it turns out that

$$\left(\sqrt{1 + \frac{\rho_{DE}}{\rho_r}} - 1 \right) H_{GR} = 5q T_f^4 \delta T_f. \quad (56)$$

Since $\rho_{DE} \ll \rho_r$ we finally obtain

$$\frac{\delta T_f}{T_f} \simeq \frac{\rho_{DE}}{\rho_r} \frac{H_{GR}}{10q T_f^5}. \quad (57)$$

We have now all the information needed to proceed to the investigation of the BBN bounds on the parameters of the $f(Q)$ models introduced above.

V. BBN CONSTRAINTS ON $f(Q)$ GRAVITY

In this section, the models introduced in Sec. III will be tested against the constraint (53). This will be realized by calculating (57) for each of the models. In general, we are interested in the regions of the parameter space where the constraint (57) is satisfied. Obviously, all models are expected to satisfy the constraint trivially for the case $\Omega_{F0} = 0$, i.e. in the case where there is no effective dark energy. In the following, in all figures the drawn parts correspond to satisfaction of the aforementioned BBN constraint. For the standard free parameters we impose the ranges $h \in [0.6, 0.9]$ and $\Omega_{F0} \in [0, 1]$. For the case of free model parameters we use the mathematically allowed ranges.

1. Polynomial model

Substituting Eq. (25) along with Eq. (24) in Eq. (57) we obtain

$$\frac{\delta T_f}{T_f} \simeq \frac{1}{q} 2^{4\alpha+3} 3^{\frac{1}{2}-\alpha} 5^{-\alpha-\frac{3}{2}} \pi^{3\alpha+2} H_0 \Omega_{F0} \cdot g_*^{\alpha+\frac{1}{2}} T_f^{4\alpha-3} (H_0 M_p)^{-2\alpha-1}. \quad (58)$$

Imposing the constraint (53) on the parameter space of the polynomial model using (58), we obtain Fig. 1. Note that we imposed the relation $H_0 = 2.12 \cdot 10^{-42} h \text{ GeV}$ and we have also replaced the numerical values (49), (42), (50) and (48).

From Fig. 1 we deduce that the model parameter α should be in the range $\alpha \lesssim 0.88$ in order for the model to satisfy the BBN constraints. However, if we additionally desire the model to be able to describe dark energy in the late-time Universe, then we obtain the combined constraint $\alpha < 0$.

2. Power-Exponential model

Substituting Eq. (28) alongside with (27) into (57) we acquire

$$\frac{\delta T_f}{T_f} \simeq \frac{8\sqrt{\frac{3}{5}}\pi^2\sqrt{g}T_0^4 \left\{ e^{\lambda\left(\frac{T_0}{T_f}\right)^4} \left[2\lambda - \left(\frac{T_f}{T_0}\right)^4 \right] + \left(\frac{T_f}{T_0}\right)^4 \right\}}{5qT_f^7 M_p}. \quad (59)$$

Inserting this expression into (53) we find that for the range of values $0.5 \leq h \leq 0.9$ and $0 \leq \Omega_{F0} \leq 1$, the constraint (53) is satisfied trivially for all parameter values. This result is intuitively expected, since as we mentioned above the past asymptotic behavior of the model recovers pure GR at early times, and thus during the BBN epoch [41]. This is a great advantage of this particular model.




FIG. 1. The parameter space of the polynomial model (23) that is consistent with the BBN constraints. If we also desire the model to be able to describe dark energy in the late-time Universe we obtain the combined constraint $\alpha < 0$.

3. Log-square-root

In this case, inserting (31) and (30) in (57) we find

$$\frac{\delta T_f}{T_f} \simeq 8\pi^2 \sqrt{\frac{3g}{5}} \frac{\Omega_{F0} T_0^2}{5qT_f^5 M_p}. \quad (60)$$

Therefore, substituting (60) in (53) we observe that for the range of values $0.5 \leq h \leq 0.9$ and $0 \leq \Omega_{F0} \leq 1$, the constraint (53) is always satisfied. This is an advantage of the model, and supports its viability. Definitely, one should confront the model with late-time observational data, too.

4. Hyperbolic tangent - power model

In the case of the hyperbolic tangent - power model (32), substituting (34) and (33) into (57) we obtain

$$\frac{\delta T_f}{T_f} \simeq \sqrt{\frac{3}{5}} \frac{8\pi^2 \sqrt{g}}{5qM_p T_f^7} \left(\frac{T_f}{T_0}\right)^{4n-4} \cdot \left\{ (1-2n)T_f^4 \tanh\left[\left(\frac{T_0}{T_f}\right)^4\right] + 2T_0^4 \operatorname{sech}^2\left[\left(\frac{T_0}{T_f}\right)^4\right] \right\}. \quad (61)$$

In Fig. 2 we depict the constraint (53) by using (61). The procedure is the same as in case of Fig. 1 above. As we observe, the BBN constraints are satisfied for $n \lesssim 1.88$.




FIG. 2. The parameter space of the hyperbolic tangent - power model (32) that is consistent with the BBN constraints. If we also desire the model to be able to describe dark energy in the late-time Universe we obtain the combined constraint $n \lesssim 1.88$.

5. DGP-like model - I

For the DGP-like model - I (35), if we insert (37) and (36) into (57), we acquire

$$\frac{\delta T_f}{T_f} \simeq \frac{8\sqrt{\frac{3}{5}}\pi^2 \sqrt{g} T_0^2 \left[2\beta + \Omega_{F0} - 2\beta \left(\frac{T_f}{T_0}\right)^2 + \left(\frac{T_f}{T_0}\right)^2 - 1 \right]}{5qM_p T_f^5}. \quad (62)$$

Furthermore, numerical exploration of this equation for the standard parameter ranges $0.5 \leq h \leq 0.9$ and $0 \leq \Omega_{F0} \leq 1$, shows that the constraint (53) is satisfied only in the tuned interval $0.499984 < \beta < 0.500016$.

6. DGP-like model - II

Finally, let us come to the DGP-like model - II of (38). The constraint (57), after inserting (40) and (39), reduces

to

$$\frac{\delta T_f}{T_f} \simeq \sqrt{\frac{3}{5} \frac{8\pi^2 \sqrt{g} T_0^2}{5qM_p T_f^5}} \cdot \left\{ \left(\frac{T_f}{T_0} \right)^2 - \sqrt{(\Omega_{F0} - 2)\Omega_{F0} + \beta^2 \left[\left(\frac{T_f}{T_0} \right)^4 - 1 \right]} + 1 \right\}. \quad (63)$$

Examining this expression for the range of values $0.5 \leq h \leq 0.9$ and $0 \leq \Omega_{F0} \leq 1$, we deduce that the constraint (53) is satisfied in the tuned window $0.999969 < |\beta| < 1.00003$.

VI. CONCLUSIONS

We use the Big Bang Nucleosynthesis formalism and observations in order to extract constraints on $f(Q)$ gravity. The latter is a modified gravitational theory arising from the consideration of non-metricity as the basic geometric quantity. We investigated various classes of $f(Q)$ models, and we additionally introduced a number of new ones, inspired by other modified gravities, that are capable of describing the late-time Universe evolution. We applied the semi-analytic approach of [50–52] for the physics of the BBN epoch, and we calculated the deviations that $f(Q)$ terms bring on the freeze-out temperature T_f in comparison to that of the standard Λ CDM evolution. We then imposed the observational bound on $\left| \frac{\delta T_f}{T_f} \right|$ to extract constraints on the involved parameters of the considered models.

Concerning the polynomial model, we showed that the exponent parameter α should be in the interval $\alpha < 0$ if we desire the model to simultaneously pass the BBN constraints and be able to describe the late-time Universe acceleration. On the other hand, the power-exponential model is found to pass the BBN constraints trivially, which was expected since this model recovers general rel-

ativity at early times. This is a significant advantage, since this particular model is known to be compatible with late-time observational datasets slightly more efficiently than Λ CDM scenario [41], and thus it can be considered as a good candidate for the description of Nature.

In the case of the new proposed Log-square-root models we showed that BBN constraints are also trivially satisfied. However, in the new hyperbolic tangent - power model we showed that the combined BBN and late-time constraints require the model parameter to lie in the range $n \lesssim 1.88$. Finally, we examined the two DGP-like $f(Q)$ models that have recently appeared in the literature. In both cases we found that the BBN constraints are satisfied for a very narrow window for the model parameter β .

In summary, we showed that $f(Q)$ gravity can safely pass the BBN constraints, and in some cases this is obtained trivially. This is an important advantage, since many gravitational modifications, although able to describe the late-time evolution of the Universe, fall to pass the BBN confrontation since they produce too-much modification at early times. Hence, $f(Q)$ gravity proves to be a class of modified gravity that deserves further investigation. One could try to proceed beyond the semi-analytical approach of this work, using numerical codes like Parthenope [75] and AlterBBN [76]. Such a detailed investigation lies beyond the scope of the present work and it is left for a future project.

ACKNOWLEDGMENTS

The authors acknowledge participation in the COST Association Action CA18108 “*Quantum Gravity Phenomenology in the Multimessenger Approach (QG-MM)*”.

-
- [1] **Planck** Collaboration, N. Aghanim *et al.*, “Planck 2018 results. VI. Cosmological parameters,” [arXiv:1807.06209](https://arxiv.org/abs/1807.06209) [[astro-ph.CO](https://arxiv.org/abs/1807.06209)]. **1**
- [2] D. M. Scolnic, D. Jones, A. Rest, Y. Pan, R. Chornock, R. Foley, M. Huber, R. Kessler, G. Narayan, A. Riess, *et al.*, “The complete light-curve sample of spectroscopically confirmed sne ia from pan-starrs1 and cosmological constraints from the combined pantheon sample,” *The Astrophysical Journal* **859** (2018) no. 2, 101. The numerical data of the full Pantheon Smla sample are available at <http://dx.doi.org/10.17909/T95Q4X>, <https://archive.stsci.edu/prepds/ps1cosmo/index.html>. **1**
- [3] H. Gil-Marín *et al.*, “The clustering of galaxies in the SDSS-III Baryon Oscillation Spectroscopic Survey: RSD measurement from the LOS-dependent power spectrum of DR12 BOSS galaxies,” *Mon. Not. Roy. Astron. Soc.* **460** (2016) no. 4, 4188–4209, [arXiv:1509.06386](https://arxiv.org/abs/1509.06386) [[astro-ph.CO](https://arxiv.org/abs/1509.06386)]. **1**
- [4] E. Di Valentino, L. A. Anchordoqui, O. Akarsu, Y. Ali-Haimoud, L. Amendola, N. Arendse, M. Asgari, M. Ballardini, S. Basilakos, E. Battistelli, *et al.*, “Cosmology intertwined ii: the hubble constant tension,” *arXiv preprint arXiv:2008.11284* (2020). **1**
- [5] E. Di Valentino, L. A. Anchordoqui, Ö. Akarsu, Y. Ali-Haimoud, L. Amendola, N. Arendse, M. Asgari, M. Ballardini, S. Basilakos, E. Battistelli, *et al.*, “Cosmology intertwined iii: $f\sigma_8$ and s_8 ,” *Astroparticle Physics* **131** (2021) 102604. **1**
- [6] S. Weinberg, “The cosmological constant problem,” *Reviews of modern physics* **61** (1989) no. 1, 1. **1**
- [7] A. Addazi *et al.*, “Quantum gravity phenomenology at the dawn of the multi-messenger era—A review,” *Prog.*

- Part. Nucl. Phys.* **125** (2022) 103948, [arXiv:2111.05659 \[hep-ph\]](#). I
- [8] K. S. Stelle, “Renormalization of higher-derivative quantum gravity,” *Physical Review D* **16** (1977) no. 4, 953. I
- [9] CANTATA Collaboration, E. N. Saridakis *et al.*, “Modified Gravity and Cosmology: An Update by the CANTATA Network,” [arXiv:2105.12582 \[gr-qc\]](#). I
- [10] S. Capozziello and M. De Laurentis, “Extended Theories of Gravity,” *Phys. Rept.* **509** (2011) 167–321, [arXiv:1108.6266 \[gr-qc\]](#). I
- [11] E. Abdalla *et al.*, “Cosmology intertwined: A review of the particle physics, astrophysics, and cosmology associated with the cosmological tensions and anomalies,” *JHEAp* **34** (2022) 49–211, [arXiv:2203.06142 \[astro-ph.CO\]](#). I
- [12] A. De Felice and S. Tsujikawa, “f(R) theories,” *Living Rev. Rel.* **13** (2010) 3, [arXiv:1002.4928 \[gr-qc\]](#). I
- [13] I. Antoniadis, J. Rizos, and K. Tamvakis, “Singularity-free cosmological solutions of the superstring effective action,” *Nucl. Phys. B* **415** (1994) 497–514, [arXiv:hep-th/9305025](#). I
- [14] S. Nojiri and S. D. Odintsov, “Modified Gauss-Bonnet theory as gravitational alternative for dark energy,” *Phys. Lett. B* **631** (2005) 1–6, [arXiv:hep-th/0508049](#). I
- [15] P. D. Mannheim and D. Kazanas, “Exact Vacuum Solution to Conformal Weyl Gravity and Galactic Rotation Curves,” *Astrophys. J.* **342** (1989) 635–638. I
- [16] G. R. Bengochea and R. Ferraro, “Dark torsion as the cosmic speed-up,” *Phys. Rev. D* **79** (2009) 124019, [arXiv:0812.1205 \[astro-ph\]](#). I
- [17] Y.-F. Cai, S. Capozziello, M. De Laurentis, and E. N. Saridakis, “f(T) teleparallel gravity and cosmology,” *Rept. Prog. Phys.* **79** (2016) no. 10, 106901, [arXiv:1511.07586 \[gr-qc\]](#). I
- [18] G. Kofinas and E. N. Saridakis, “Teleparallel equivalent of Gauss-Bonnet gravity and its modifications,” *Phys. Rev. D* **90** (2014) 084044, [arXiv:1404.2249 \[gr-qc\]](#). I
- [19] C.-Q. Geng, C.-C. Lee, E. N. Saridakis, and Y.-P. Wu, ““Teleparallel” dark energy,” *Phys. Lett. B* **704** (2011) 384–387, [arXiv:1109.1092 \[hep-th\]](#). I
- [20] M. Hohmann, L. Järv, and U. Ualikhanova, “Covariant formulation of scalar-torsion gravity,” *Phys. Rev. D* **97** (2018) no. 10, 104011, [arXiv:1801.05786 \[gr-qc\]](#). I
- [21] J. Beltrán Jiménez, L. Heisenberg, and T. Koivisto, “Coincident General Relativity,” *Phys. Rev. D* **98** (2018) no. 4, 044048, [arXiv:1710.03116 \[gr-qc\]](#). I, II, II, II, II
- [22] J. Beltrán Jiménez, L. Heisenberg, T. S. Koivisto, and S. Pekar, “Cosmology in $f(Q)$ geometry,” *Phys. Rev. D* **101** (2020) no. 10, 103507, [arXiv:1906.10027 \[gr-qc\]](#). I, III, 1
- [23] Y. Xu, G. Li, T. Harko, and S.-D. Liang, “ $f(Q, T)$ gravity,” *Eur. Phys. J. C* **79** (2019) no. 8, 708, [arXiv:1908.04760 \[gr-qc\]](#).
- [24] J. B. Jiménez, L. Heisenberg, D. Iosifidis, A. Jiménez-Cano, and T. S. Koivisto, “General Teleparallel Quadratic Gravity,” [arXiv:1909.09045 \[gr-qc\]](#).
- [25] R. Percacci and E. Sezgin, “New class of ghost- and tachyon-free metric affine gravities,” *Phys. Rev. D* **101** (2020) no. 8, 084040, [arXiv:1912.01023 \[hep-th\]](#).
- [26] A. Delhom, I. P. Lobo, G. J. Olmo, and C. Romero, “Conformally invariant proper time with general non-metricity,” *Eur. Phys. J. C* **80** (2020) no. 5, 415, [arXiv:2001.10633 \[gr-qc\]](#).
- [27] A. Delhom, “Minimal coupling in presence of non-metricity and torsion,” *Eur. Phys. J. C* **80** (2020) no. 8, 728, [arXiv:2002.02404 \[gr-qc\]](#).
- [28] B. J. Barros, T. Barreiro, T. Koivisto, and N. J. Nunes, “Testing $F(Q)$ gravity with redshift space distortions,” *Phys. Dark Univ.* **30** (2020) 100616, [arXiv:2004.07867 \[gr-qc\]](#). III
- [29] J. Beltrán Jiménez, L. Heisenberg, and T. Koivisto, “The coupling of matter and spacetime geometry,” *Class. Quant. Grav.* **37** (2020) no. 19, 195013, [arXiv:2004.04606 \[hep-th\]](#).
- [30] A. Jiménez-Cano, “New metric-affine generalizations of gravitational wave geometries,” *Eur. Phys. J. C* **80** (2020) no. 7, 672, [arXiv:2005.02014 \[gr-qc\]](#).
- [31] F. D’Ambrosio, M. Garg, and L. Heisenberg, “Non-linear extension of non-metricity scalar for MOND,” *Phys. Lett. B* **811** (2020) 135970, [arXiv:2004.00888 \[gr-qc\]](#).
- [32] D. Rubiera-Garcia, “From fundamental physics to tests with compact objects in metric-affine theories of gravity,” *Int. J. Mod. Phys. D* **29** (2020) no. 11, 2041007, [arXiv:2004.00943 \[gr-qc\]](#).
- [33] Y. Xu, T. Harko, S. Shahidi, and S.-D. Liang, “Weyl type $f(Q, T)$ gravity, and its cosmological implications,” *Eur. Phys. J. C* **80** (2020) no. 5, 449, [arXiv:2005.04025 \[gr-qc\]](#).
- [34] I. Ayuso, R. Lazkoz, and V. Salzano, “Observational constraints on cosmological solutions of $f(Q)$ theories,” *Phys. Rev. D* **103** (2021) no. 6, 063505, [arXiv:2012.00046 \[astro-ph.CO\]](#). 1
- [35] F. Cabral, F. S. N. Lobo, and D. Rubiera-Garcia, “Fundamental Symmetries and Spacetime Geometries in Gauge Theories of Gravity—Prospects for Unified Field Theories,” *Universe* **6** (2020) no. 12, 238, [arXiv:2012.06356 \[gr-qc\]](#).
- [36] K. Flathmann and M. Hohmann, “Post-Newtonian limit of generalized symmetric teleparallel gravity,” *Phys. Rev. D* **103** (2021) no. 4, 044030, [arXiv:2012.12875 \[gr-qc\]](#).
- [37] N. Frusciante, “Signatures of $f(Q)$ -gravity in cosmology,” *Phys. Rev. D* **103** (2021) no. 4, 044021, [arXiv:2101.09242 \[astro-ph.CO\]](#).
- [38] J.-Z. Yang, S. Shahidi, T. Harko, and S.-D. Liang, “Geodesic deviation, Raychaudhuri equation, Newtonian limit, and tidal forces in Weyl-type $f(Q, T)$ gravity,” *Eur. Phys. J. C* **81** (2021) no. 2, 111, [arXiv:2101.09956 \[gr-qc\]](#).
- [39] Q.-M. Fu, L. Zhao, and Q.-Y. Xie, “Thick braneworld model in nonmetricity formulation of general relativity and its stability,” [arXiv:2103.03618 \[gr-qc\]](#).
- [40] W. Khylllep, A. Paliathanasis, and J. Dutta, “Cosmological solutions and growth index of matter perturbations in $f(Q)$ gravity,” [arXiv:2103.08372 \[gr-qc\]](#).
- [41] F. K. Anagnostopoulos, S. Basilakos, and E. N. Saridakis, “First evidence that non-metricity $f(Q)$ gravity could challenge Λ CDM,” *Phys. Lett. B* **822** (2021) 136634, [arXiv:2104.15123 \[gr-qc\]](#). 2, V, VI
- [42] I. S. Albuquerque and N. Frusciante, “A designer approach to $f(Q)$ gravity and cosmological implications,” *Phys. Dark Univ.* **35** (2022) 100980, [arXiv:2202.04637 \[astro-ph.CO\]](#). I

- [43] J. Bernstein, L. S. Brown, and G. Feinberg, “COSMOLOGICAL HELIUM PRODUCTION SIMPLIFIED,” *Rev. Mod. Phys.* **61** (1989) 25. I, IV
- [44] E. W. Kolb and M. S. Turner, *The Early Universe*, vol. 69. 1990. IV, IV
- [45] K. A. Olive, G. Steigman, and T. P. Walker, “Primordial nucleosynthesis: Theory and observations,” *Phys. Rept.* **333** (2000) 389–407, [arXiv:astro-ph/9905320](#).
- [46] R. H. Cyburt, B. D. Fields, K. A. Olive, and T.-H. Yeh, “Big Bang Nucleosynthesis: 2015,” *Rev. Mod. Phys.* **88** (2016) 015004, [arXiv:1505.01076 \[astro-ph.CO\]](#). IV
- [47] J. D. Barrow, S. Basilakos, and E. N. Saridakis, “Big Bang Nucleosynthesis constraints on Barrow entropy,” *Phys. Lett. B* **815** (2021) 136134, [arXiv:2010.00986 \[gr-qc\]](#). IV
- [48] P. Asimakis, S. Basilakos, N. E. Mavromatos, and E. N. Saridakis, “Big bang nucleosynthesis constraints on higher-order modified gravities,” *Phys. Rev. D* **105** (2022) no. 8, 084010, [arXiv:2112.10863 \[gr-qc\]](#). I
- [49] D. F. Torres, H. Vucetich, and A. Plastino, “Early universe test of nonextensive statistics,” *Phys. Rev. Lett.* **79** (1997) 1588–1590, [arXiv:astro-ph/9705068](#). [Erratum: *Phys.Rev.Lett.* 80, 3889 (1998)]. I, IV
- [50] G. Lambiase, “Lorentz invariance breakdown and constraints from big-bang nucleosynthesis,” *Phys. Rev. D* **72** (2005) 087702, [arXiv:astro-ph/0510386](#). VI
- [51] G. Lambiase, “Constraints on massive gravity theory from big bang nucleosynthesis,” *JCAP* **10** (2012) 028, [arXiv:1208.5512 \[gr-qc\]](#).
- [52] G. Lambiase, “Dark matter relic abundance and big bang nucleosynthesis in Horava’s gravity,” *Phys. Rev. D* **83** (2011) 107501. VI
- [53] S. Capozziello, G. Lambiase, and E. N. Saridakis, “Constraining $f(T)$ teleparallel gravity by Big Bang Nucleosynthesis,” *Eur. Phys. J. C* **77** (2017) no. 9, 576, [arXiv:1702.07952 \[astro-ph.CO\]](#). I, IV
- [54] M. Nakahara, *Geometry, topology and physics*. Taylor and Francis, 2003. II
- [55] J. M. Nester and H.-J. Yo, “Symmetric teleparallel general relativity,” *Chin. J. Phys.* **37** (1999) 113, [arXiv:gr-qc/9809049 \[gr-qc\]](#). II
- [56] J. B. Jiménez, L. Heisenberg, and T. S. Koivisto, “The Geometrical Trinity of Gravity,” *Universe* **5** (2019) no. 7, 173, [arXiv:1903.06830 \[hep-th\]](#). II
- [57] T. Ortín, *Gravity and Strings*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 2007. <https://books.google.com/mt/books?id=HDMucsxABZyc>. II
- [58] J. Beltrán Jiménez, L. Heisenberg, and T. S. Koivisto, “Teleparallel Palatini theories,” *JCAP* **1808** (2018) no. 08, 039, [arXiv:1803.10185 \[gr-qc\]](#). II
- [59] R. Aldrovandi and J. Pereira, *Teleparallel Gravity: An Introduction*. Fundamental Theories of Physics. Springer Netherlands, 2012. II
- [60] T. Harko, T. S. Koivisto, F. S. N. Lobo, G. J. Olmo, and D. Rubiera-Garcia, “Coupling matter in modified Q gravity,” *Phys. Rev. D* **98** (2018) no. 8, 084043, [arXiv:1806.10437 \[gr-qc\]](#). II
- [61] K. Bamba, C.-Q. Geng, C.-C. Lee, and L.-W. Luo, “Equation of state for dark energy in $f(T)$ gravity,” *JCAP* **1101** (2011) 021, [arXiv:1011.0508 \[astro-ph.CO\]](#). 3
- [62] P. Wu and H. W. Yu, “ $f(T)$ models with phantom divide line crossing,” *Eur. Phys. J. C* **71** (2011) 1552, [arXiv:1008.3669 \[gr-qc\]](#). 4
- [63] I. Ayuso, R. Lazkoz, and J. P. Mimoso, “DGP and DGPish cosmologies from $f(Q)$ actions,” [arXiv:2111.05061 \[astro-ph.CO\]](#). 5, 5, 6, 6
- [64] G. Dvali, G. Gabadadze, and M. Porrati, “4d gravity on a brane in 5d minkowski space,” *Physics Letters B* **485** (2000) no. 1-3, 208–214. 5
- [65] M. Fairbairn and A. Goobar, “Supernova limits on brane world cosmology,” *Phys. Lett. B* **642** (2006) 432–435, [arXiv:astro-ph/0511029](#). 5
- [66] W. Fang, S. Wang, W. Hu, Z. Haiman, L. Hui, and M. May, “Challenges to the dgp model from horizon-scale growth and geometry,” *Physical Review D* **78** (2008) no. 10, 103509. 5
- [67] J. Bernstein, L. S. Brown, and G. Feinberg, “Cosmological helium production simplified,” *Reviews of Modern physics* **61** (1989) no. 1, 25. IV
- [68] A. Coc, E. Vangioni-Flam, P. Descouvemont, A. Adahchour, and C. Angulo, “Updated Big Bang nucleosynthesis confronted to WMAP observations and to the abundance of light elements,” *Astrophys. J.* **600** (2004) 544–552, [arXiv:astro-ph/0309480](#). IV
- [69] K. A. Olive, E. Skillman, and G. Steigman, “The Primordial abundance of He-4: An Update,” *Astrophys. J.* **483** (1997) 788, [arXiv:astro-ph/9611166](#).
- [70] Y. I. Izotov and T. X. Thuan, “The Primordial abundance of 4-He revisited,” *Astrophys. J.* **500** (1998) 188.
- [71] B. D. Fields and K. A. Olive, “On the evolution of helium in blue compact galaxies,” *Astrophys. J.* **506** (1998) 177, [arXiv:astro-ph/9803297](#).
- [72] Y. I. Izotov, F. H. Chaffee, C. B. Foltz, R. F. Green, N. G. Guseva, and T. X. Thuan, “Helium abundance in the most metal-deficient blue compact galaxies: I zw 18 and sbs 0335-052,” *Astrophys. J.* **527** (1999) 757–777, [arXiv:astro-ph/9907228](#).
- [73] D. Kirkman, D. Tytler, N. Suzuki, J. M. O’Meara, and D. Lubin, “The Cosmological baryon density from the deuterium to hydrogen ratio towards QSO absorption systems: D/H towards Q1243+3047,” *Astrophys. J. Suppl.* **149** (2003) 1, [arXiv:astro-ph/0302006](#).
- [74] Y. I. Izotov and T. X. Thuan, “Systematic effects and a new determination of the primordial abundance of He-4 and dY/dZ from observations of blue compact galaxies,” *Astrophys. J.* **602** (2004) 200–230, [arXiv:astro-ph/0310421](#). IV
- [75] S. Gariazzo, P. de Salas, O. Pisanti, and R. Consiglio, “Parthenope revolutions,” *arXiv preprint arXiv:2103.05027* (2021) . VI
- [76] A. Arbey, J. Auffinger, K. P. Hickerson, and E. S. Jansen, “AlterBBN v2: A public code for calculating Big-Bang nucleosynthesis constraints in alternative cosmologies,” *Comput. Phys. Commun.* **248** (2020) 106982, [arXiv:1806.11095 \[astro-ph.CO\]](#). VI