

# Low-Rank Univariate Sum of Squares Has No Spurious Local Minima

Benoît Legat

Chenyang Yuan

Pablo A. Parrilo

May 24, 2022

## Abstract

We study the problem of decomposing a polynomial  $p$  into a sum of  $r$  squares by minimizing a quadratically-penalized objective  $f_p(\mathbf{u}) = \|\sum_{i=1}^r u_i^2 - p\|^2$ . This objective is non-convex and is equivalent to the rank- $r$  Burer-Monteiro factorization of a semidefinite program (SDP) encoding the sum of squares decomposition. We show that for all univariate polynomials  $p$ , if  $r \geq 2$  then  $f_p(\mathbf{u})$  has no spurious second-order critical points, showing that all local optima are also global optima. This is in contrast to previous work showing that for general SDPs, in addition to genericity conditions,  $r$  has to be roughly the square root of the number of constraints (the degree of  $p$ ) for there to be no spurious second-order critical points. Our proof uses tools from computational algebraic geometry and can be interpreted as constructing a certificate using the first and second order necessary conditions. We also show that by choosing a norm based on sampling equally-spaced points on the circle, the gradient  $\nabla f_p$  can be computed in nearly linear time using fast Fourier transforms. Experimentally we demonstrate that this method has very fast convergence using first-order optimization algorithms such as L-BFGS, with near-linear scaling to million-degree polynomials.

## 1 Introduction

Burer-Monteiro factorization [BM03] is a methodology to solve large-scale semidefinite programs (SDPs) by replacing positive semidefinite (PSD) variables  $X \succeq 0$  with a factorization  $X = UU^\top$ . This automatically enforces the PSD constraint and lets us find low-rank solutions by choosing the rank of the new variable  $U$ . In addition, this factorization results in a nonlinear optimization problem that can be solved with first-order methods with fast per-iteration times especially when  $\text{rank}(U)$  is small. However, the resulting problem is non-convex so these methods may get stuck in local optima. We show that this will not happen to the SDP finding the sum-of-squares decomposition of univariate polynomials; in this setting all local optima are also global.

In this work we study SDPs arising from Sum-of-Squares optimization [Par00]. The ability to represent the cone of sum of squares polynomials as a SDP enables many applications in polynomial optimization, control, and relaxations of combinatorial problems [Lau09, BPT13]. To determine if a polynomial  $p(x) \in \mathbb{R}[x]_{2d}$  is a sum of squares, it suffices to find a feasible solution to the following SDP:

$$\begin{aligned} p(x) &= b(x)^\top X b(x) \\ X &\succeq 0 \end{aligned} \tag{1}$$

where  $b(x)$  is a suitable polynomial basis of  $\mathbb{R}[x]_d$ . The constraint  $p(x) = b(x)^\top X b(x)$  defines an affine subspace of the space of symmetric matrices, which can be expressed by matching the coefficients of  $p(x)$  with the corresponding coefficients of the polynomial  $b(x)^\top X b(x)$ .

Given the factorization  $X = UU^\top$  where  $\bar{u}_1, \dots, \bar{u}_r$  are the column vectors of  $U$ , we have the explicit sum of squares decomposition

$$p(x) = \sum_{i=1}^r u_i(x)^2, \quad (2)$$

where  $u_i(x) = b(x)^\top \bar{u}_i$ . In other words, (2) is a coordinate-free formulation for the Burer-Monteiro factorization of (1). Instead of solving a SDP to find  $u_i(x)$ , we apply a quadratic penalty to the equality constraint (2) to arrive at the following non-convex objective:

$$f_p(\mathbf{u}) = \left\| \sum_{i=1}^r u_i(x)^2 - p(x) \right\|^2, \quad (3)$$

where  $\mathbf{u} = [u_1 \ \dots \ u_r]$  is a vector of  $r$  degree- $d$  polynomials and the norm is induced by any inner product on polynomials of degree- $2d$ . Then a degree- $2d$  polynomial  $p(x)$  is a sum of  $r$  squares if and only if

$$\min_{\mathbf{u} \in \mathbb{R}[x]_d^r} f_p(\mathbf{u}) = 0.$$

We say that  $\mathbf{u}$  is a *first-order critical point* (FOCP) of  $f_p(\mathbf{u})$  when  $\nabla f_p(\mathbf{u}) = 0$ , where the derivatives are taken with respect to the variables in  $\mathbf{u}$ . Since for every  $p$  we can find spurious FOCPs (for example taking  $\mathbf{u} = 0$ ), it is essential to consider second-order necessary conditions. If  $\mathbf{u}$  also satisfies  $\nabla^2 f_p(\mathbf{u}) \succeq 0$ , then it is a *second-order critical point* (SOCP) of  $f_p(\mathbf{u})$ .

Every local minima of a function is a SOCP, but the converse is not true when the function is non-convex<sup>1</sup>. If  $\mathbf{u}$  is not a SOCP, then we can produce a descent direction using the gradient or Hessian. This leads to efficient first-order methods [GHJY15, JGN<sup>+</sup>17] that converge to SOCPs. Thus if we can show that at every SOCP  $f_p(\mathbf{u}) = 0$ , these algorithms will always converge to a global minimum. A recent line of work [BVB20, BBJN18] has shown that for general SDPs under smoothed analysis or genericity conditions, when the rank of the factorization is above the Barvinok-Pataki bound (roughly the square root of the number of constraints), there are no spurious SOCPs. Moreover, the smoothed analysis or genericity conditions are necessary, as [BBJN18] constructed a SDP where only a full-rank factorization can guarantee no spurious SOCPs.

In this paper we consider the setting of univariate polynomial optimization, which is a class of problems with applications in signal processing, control [RV06, Dum07] as well as computing equilibria of polynomial games [Par06]. The main result of our paper shows that without any additional assumptions, the rank-2 quadratic penalized Burer-Monteiro factorization of the SDP describing the sum of squares decomposition (3) of a univariate polynomial has no spurious SOCPs.

**Theorem 1.1.** *For all nonnegative univariate polynomials  $p(x) \in \mathbb{R}[x]_{2d}$  and any  $r \geq 2$ , if  $\mathbf{u} \in \mathbb{R}[x]_d^r$  satisfies  $\nabla f_p(\mathbf{u}) = 0$  and  $\nabla^2 f_p(\mathbf{u}) \succeq 0$ , then  $f_p(\mathbf{u}) = 0$ .*

In particular, the rank bound in our result matches the Pythagoras number for univariate polynomials [CLR95, Example 2.13]<sup>2</sup>. In comparison, applying rank bounds for general SDPs to this setting require  $r \gtrsim \sqrt{d}$  (see Section 2 for more details).

<sup>1</sup>Consider, for example,  $f(x) = x^3$  at  $x = 0$ .

<sup>2</sup>The Pythagoras number for  $\Sigma[x]_{2d}$  is the smallest  $r$  such that all polynomials in  $\Sigma[x]_{2d}$  can be written as a sum of  $r$  squares of polynomials in  $\mathbb{R}[x]_d$ .

Theorem 1.1 is proved in Section 4, by deriving a series of increasingly stronger sufficient conditions ((C1), (C2) and (C3)) implying  $f_p(\mathbf{u}) = 0$  for increasingly larger classes of  $\mathbf{u} = (u_1, u_2)$ , eventually proving the result for all  $\mathbf{u} \in \mathbb{R}[x]^2$ . To illustrate this, we first show that when  $r = 2$  and  $u_1, u_2$  are coprime,  $\nabla f_p(\mathbf{u}) = 0$  implies that  $f_p(\mathbf{u}) = 0$  (the precise statement and proof of this case is in Section 4.1). Note that in this simplified setting only the first-order gradient condition is needed. By computing the gradient,  $\nabla f_p(\mathbf{u}) = 0$  is equivalent to

$$\nabla f_p(\mathbf{u})(\mathbf{v}) = \langle u_1 v_1 + u_2 v_2, u_1^2 + u_2^2 - p \rangle = 0,$$

for all  $\mathbf{v} = (v_1, v_2) \in \mathbb{R}[x]^2$ . Since  $u_1, u_2$  are coprime, Bézout's identity (Lemma 3.2) implies that we can find  $\mathbf{v}' = (v'_1, v'_2)$  so that

$$u_1 v'_1 + u_2 v'_2 = u_1^2 + u_2^2 - p, \tag{4}$$

thus showing that  $f_p(\mathbf{u}) = \|u_1^2 + u_2^2 - p\|^2 = 0$ . However, we cannot assume a priori that  $u_1, u_2$  are coprime as we are only given  $p$  as the input. The main technical contribution of our proof is how to handle the more involved case when  $u_1, u_2$  share a common factor.

We can also interpret our proof as a certificate. When we choose  $\mathbf{v}'$  satisfying (4), we obtain the identity

$$\nabla f_p(\mathbf{u})(\mathbf{v}') = f_p(\mathbf{u}).$$

This implies that  $f_p(\mathbf{u}) = 0$  when  $\nabla f_p(\mathbf{u}) = 0$ . In Section 5 we generalize this example to our full proof of Theorem 1.1, showing for all  $\mathbf{u}$  and  $p$  how to find  $\mathbf{v}'$  and  $Q \succeq 0$  satisfying the following identity:

$$\nabla f_p(\mathbf{u})(\mathbf{v}') + \langle Q, \nabla^2 f_p(\mathbf{u}) \rangle = -f_p(\mathbf{u}).$$

From this identity it is clear that if  $\mathbf{u}$  is a SOCP then  $f_p(\mathbf{u}) = 0$ . This compact form of our proof allows us to easily extend our result to other problems in Section 6.

Since Theorem 1.1 holds for any inner product, we can choose one that enables efficient computation of  $\nabla f_p(\mathbf{u})$ . When  $p$  is a degree- $2d$  univariate polynomial, an equivalent way of ensuring the constraint  $p(x) = b(x)^\top X b(x)$  in (1) is to write  $2d + 1$  constraints  $p(\hat{x}_i) = b(\hat{x}_i)^\top X b(\hat{x}_i)$ , where  $\hat{x}_0, \dots, \hat{x}_d$  are distinct sample points. This formulation can be cast into the following least-squares objective,

$$\min_{U \in \mathbb{R}^{(2d+1) \times r}} f_p(U) = \frac{1}{2d+1} \sum_{i=1}^{2d+1} \left( \left\| U^\top b(\hat{x}_i) \right\|_2^2 - p(\hat{x}_i) \right)^2. \tag{5}$$

This is equivalent to choosing an inner product in (3) that evaluates the polynomial on  $2d+1$  points. If we choose  $2d + 1$  points on the complex unit circle, we can compute  $\nabla f_p(U)$  in  $O(d \log d)$  time using the Fast Fourier Transform (FFT). In Section 7, we show that this method exhibits linear convergence experimentally using unconstrained optimization such as L-BFGS, and has near-linear scaling to million-degree polynomials.

## 1.1 Contributions

In summary, our main contributions in this paper are:

1. Proving that the quadratic penalty form Burer-Monteiro factorization for univariate polynomial sum of squares decomposition has no spurious SOCPs (Theorem 1.1). Our result holds where the rank of the factorization is at least 2, matching the Pythagoras number for univariate polynomials. This is in contrast to previous work requiring rank  $r \gtrsim \sqrt{d}$  in addition to genericity conditions or smoothed analysis for general SDPs, or showing that no spurious local minima exist in statistical problems.
2. Developing a new framework for proving that there are no spurious SOCPs for a quadratic-penalized factorized SDPs, by constructing a certificate (20) using the first and second order necessary conditions. This certificate representation helps us extend our results to projection onto the sum of squares cone (Corollary 6.1), certifying non-negativity on intervals (Corollary 6.2) and sum of squares optimization.
3. Showing that by choosing a special norm (based on the evaluation of the polynomial on points on the unit circle), the full gradient of the objective can be computed in near-linear time using FFTs. It enables us to efficiently scale first-order methods to instances with millions of variables (Table 1). This is possible because our result (1.1) is independent of the penalty function.

## 2 Background and Related Work

Let  $\mathbb{S}_n$  be the space of  $n \times n$  symmetric matrices. Given  $A_i \in \mathbb{S}_n$ , we consider the standard-form semidefinite feasibility problem with variable  $X \in \mathbb{S}_n$ :

$$\begin{aligned} \langle A_i, X \rangle &= b_i \\ X &\succeq 0 \end{aligned} \tag{SDP}$$

**Non-convex formulation** Burer and Monteiro [BM03] introduced the non-convex reformulation  $X = UU^\top$  to enforce the semidefinite constraint, where  $U \in \mathbb{R}^{n \times r}$ .

$$\langle A_i, UU^\top \rangle = b_i \tag{NSDP}_r$$

This motivates the following least-squares formulation:

$$\min_{U \in \mathbb{R}^{n \times r}} \sum_i \|\langle A_i, UU^\top \rangle - b_i\|^2 \tag{SDPLS}_r$$

If in addition to the equality constraints in (SDP) one wishes to minimize the objective  $\langle C, X \rangle$ , the works [BM05, BBJN18, CM19] formulate an augmented Lagrangian problem, where the term  $\lambda \langle C, UU^\top \rangle$  is added to the objective of (SDPLS<sub>r</sub>).

**General SDP rank bounds** Burer and Monteiro subsequently showed in [BM05] that when  $r \geq n$ , there are no spurious SOCPs to (SDPLS<sub>r</sub>). This result is in fact tight and [BBJN18] constructed an explicit instance where if  $r = n - 1$ , one can find a SOCP that is not a global minimum. Thus for general SDP feasibility, additional conditions on the objective or

analysis must be imposed. Then the rank bound can be improved to the maximum rank of extreme points of the section of the PSD cone with  $m$  affine constraints. This maximum rank is  $O(\sqrt{m})$ , also known as the Barvinok-Pataki bound ([Bar01, Pat98]). In the same work, Burer and Monteiro [BM05] showed that if  $r \gtrsim \sqrt{m}$ , any local minimum of  $\text{NSDP}_r$  is also a local minimum of SDP with an additional rank- $r$  constraint. Then they showed that such a local minimum is either an optimal extreme point, or contained within the relative interior of a face of the feasible set of SDP which is constant with respect to the objective function.

Subsequent work [BBJN18] then showed that if  $C$  is generic enough, all local minimum of  $\text{NSDP}_r$  are global minimal of SDP (see Cifuentes and Moitra [CM19] for more references). In summary this line of work either requires smoothness of the constraint set ([BVB16, BVB20]), generic constraints and/or smoothed analysis ([BBJN18, CM19, Cif21]). In addition, [WW20] showed that when  $r$  is smaller than  $\sqrt{m}$ , SOCPs are not generically optimal.

**Structured SDPs** Problems such as matrix completion and matrix sensing can be expressed as instances of  $\text{SDPLS}_r$ . There has been a lot of recent interest in studying the global landscape of matrix sensing problems ([GLM16, BNS16, GJZ17]). A recent line of work ([GJZ17, GLM16, BBV16, SQW18, LZT19]) shows that for certain planted statistical problems where the SDP has a rank-1 solution in the noiseless setting, there are no spurious SOCPs when the noise level is low enough. See [CLC19] for a survey of these problems. In summary, for a wide range of statistical problems, local minima are also global minima. These results either assume a planted solution, or that the sensing operators  $A_i$  satisfy the Restricted Isometry Property (RIP).

**Sampling basis** The sampling or interpolation basis for sum of squares optimization is studied in [LP04] and [CP17]. Informally, they showed that if the sampled points are “generic” enough, the problem expressed in the sampled basis is equivalent to the original problem.

**Univariate polynomials** Univariate/trigonometric polynomial optimization and their applications are studied in [WBV99, RV06, Dum07] and contained references. The decomposition of a nonnegative univariate polynomial into two squares is equivalent to the spectral factorization of a nonnegative trigonometric polynomial. Previous methods for spectral factorization either require finding all  $n$  roots of the polynomial, solving linear systems of order  $n$ , or use an approximate  $O(N \log N)$  FFT-based algorithm by sampling  $N \gg n$  points [WBV99, Dum07]. Design problems involving constraints on nonnegative trigonometric polynomials can be formulated as SDPs. Due to their special structure, [RV06] used FFTs to speed up per-iteration complexity for interior point methods solving these SDPs to  $O(n^3)$ .

The set of all  $X$  satisfying (1) is known as the Gram spectrahedron of  $p(x)$ . A recent work by Scheiderer [Sch22] showed that if  $p(x)$  is a sufficiently general positive univariate polynomial of degree  $d$ , its Gram spectrahedron has extreme points of all ranks up to the Barvinok-Pataki bound. Thus applying the general SDP rank bounds give  $r \gtrsim \sqrt{d}$ .

## 2.1 Notation

Let  $\mathbb{R}[x]_d$  be the space of univariate polynomials of degree at most  $d$  in  $n$  variables. Let  $\mathbf{u}(x) \in \mathbb{R}[x]_d^r$  be a vector of  $r$  polynomials where each  $u_i(x)$  is a polynomial in  $\mathbb{R}[x]_d$ . Let  $\sigma : \mathbb{R}[x]_d^r \rightarrow \mathbb{R}[x]_{2d}$  be the quadratic map defined by  $\mathbf{v}(x) \mapsto \sum_{i=1}^r v_i(x)^2$ , and  $\Sigma[x]_{2d} := \text{cone}(\sigma(\mathbb{R}[x]_d^r)) \subseteq \mathbb{R}[x]_{2d}$  be

the cone of sum of squares univariate polynomials of degree- $2d$ . Correspondingly, for homogeneous polynomials let  $\mathbb{R}[x]_{\mathbf{d}}$  be the space of binary forms of degree- $d$  and  $\Sigma[x]_{2\mathbf{d}} \subseteq \mathbb{R}[x]_{2\mathbf{d}}$  be the space of sum of squares binary forms of degree- $2d$ .

## 2.2 Univariate Polynomials

Any monic univariate polynomial  $p \in \mathbb{R}[x]_d$  can be uniquely factored as  $p(x) = \prod_{i=1}^d (x - \alpha_i)$ , where  $\alpha_i \in \mathbb{C}$  are the roots of  $p$ . Given univariate polynomials  $p, g$  and  $q$ , we define an equivalence relation  $p \equiv q \pmod{g}$  if there exists  $w \in \mathbb{R}[x]$  such that  $p = q + wg$ . We say that  $g$  is a *divisor* of  $p$  if  $p \equiv 0 \pmod{g}$ . In addition, if  $g$  is also a divisor of  $q$ , then we say that  $g$  is a *common divisor* of  $p$  and  $q$ . Let  $\gcd(p, q)$  be the *greatest common divisor* of  $p$  and  $q$ , which is the common divisor with the highest degree. By the unique factorization of  $p$  and  $q$ ,  $\gcd(p, q)$  is unique up to multiplication by a scalar. We say that  $p$  and  $q$  are *coprime* if  $\gcd(p, q) = 1$ .

Any binary form  $p \in \mathbb{R}[x]_{\mathbf{d}}$  can be factored as  $p(x_1, x_2) = \prod_{i=1}^d (\alpha_i x_1 - \beta_i x_2)$ , where  $(\alpha_i, \beta_i) \in \mathbb{C}^2$ . This factorization is unique up to multiplication of  $(\alpha_i, \beta_i)$  by a scalar. The equivalence relation  $p \equiv q \pmod{g}$  and  $\gcd$  on binary forms are defined analogously to those on univariate polynomials.

## 3 Preliminary Results

Since the polynomials we are optimizing over have fixed degrees, to better keep track of this degree we will work with homogeneous polynomials (forms). In Section 3.1 we derive expressions for the first and second order necessary conditions ((8) and (9)). Next in Sections 3.2 and 3.3 we review some results on binary forms and prove our main Lemma 3.6. Particularly important is our decomposition of  $\mathbf{u}$  in Proposition 3.3.

### 3.1 First and Second Order Necessary Conditions

First we derive expressions for the gradient and Hessian of  $f_p(\mathbf{u})$ . In addition to the binary forms we consider in this paper, the derivations in this section also hold for general multivariate forms. For any inner product on forms  $\langle \cdot, \cdot \rangle : \mathbb{R}[x]_{2\mathbf{d}} \times \mathbb{R}[x]_{2\mathbf{d}} \rightarrow \mathbb{R}$  and its associated norm  $\|\cdot\| : \mathbb{R}[x]_{2\mathbf{d}} \rightarrow \mathbb{R}$ , the objective function is written as

$$f_p(\mathbf{u}(x)) = \left\| \sum_{j=1}^r u_j(x)^2 - p(x) \right\|^2.$$

To find the gradient and Hessian of the objective, we compute the first and second order terms of  $f_p(\mathbf{u}(x) + \epsilon \mathbf{v}(x))$  to obtain:

$$\frac{1}{4} \nabla f_p(\mathbf{u}(x))(\mathbf{v}(x)) = \left\langle \sum_{j=1}^r u_j(x) v_j(x), \sum_{j=1}^r u_j(x)^2 - p(x) \right\rangle, \quad (6)$$

$$\frac{1}{4} \nabla^2 f_p(\mathbf{u}(x))(\mathbf{v}(x), \mathbf{v}(x)) = \left\langle \sum_{j=1}^r v_j(x)^2, \sum_{j=1}^r u_j(x)^2 - p(x) \right\rangle + 2 \left\| \sum_{j=1}^r u_j(x) v_j(x) \right\|^2. \quad (7)$$

Given vectors of polynomials  $\mathbf{u}(x) \in \mathbb{R}[x]_{\mathbf{d}_1}^r$  and  $\mathbf{v}(x) \in \mathbb{R}[x]_{\mathbf{d}_2}^r$ , we define the linear map  $\mathcal{A}_{\mathbf{u}(x)} : \mathbb{R}[x]_{\mathbf{d}_2}^r \rightarrow \mathbb{R}[x]_{\mathbf{d}_1 + \mathbf{d}_2}$  as

$$\mathcal{A}_{\mathbf{u}(x)} : \mathbf{v}(x) \mapsto \sum_{i=1}^r u_i(x) v_i(x).$$

For example, we can write  $\sigma(\mathbf{u}) = \mathcal{A}_{\mathbf{u}}(\mathbf{u})$ . When  $n = 2$ ,  $\mathcal{A}_{\mathbf{u}}$  is up to a constant the map induced by the *Sylvester* matrix [Stu02]. If  $d_1 = d_2$ , the determinant of  $\mathcal{A}_{\mathbf{u}}(\mathbf{u})$  is up to a constant the *resultant* of  $u_1$  and  $u_2$ ; it vanishes if and only if  $u_1$  and  $u_2$  share a common divisor. We say that  $\mathbf{u} \in \mathbb{R}[x]_{\mathbf{d}}^r$  is a second-order critical point of  $f_p(\mathbf{u})$  if its gradient is zero and Hessian is positive semidefinite. Using our new notation, for all  $\mathbf{v} \in \mathbb{R}[x]_{\mathbf{d}}$  we must have

$$\frac{1}{4} \nabla f_p(\mathbf{u})(\mathbf{v}) = \langle \mathcal{A}_{\mathbf{u}}(\mathbf{v}), \sigma(\mathbf{u}) - p \rangle = 0, \quad (8)$$

$$\frac{1}{4} \nabla^2 f_p(\mathbf{u})(\mathbf{v}, \mathbf{v}) = \langle \sigma(\mathbf{v}), \sigma(\mathbf{u}) - p \rangle + 2 \|\mathcal{A}_{\mathbf{u}}(\mathbf{v})\|^2 \geq 0. \quad (9)$$

**Remark 3.1.** *The first-order condition (8) alone is insufficient to guarantee global optimality, even when  $p$  is generic. Suppose  $p \in \Sigma[x]_{2d}$  cannot be written as a sum of less than  $\hat{r}$  squares. Then any FOCP  $\hat{\mathbf{u}} = (u_1, \dots, u_{\hat{r}})$  of  $f_p(\hat{\mathbf{u}})$  is spurious as  $f_p$  is always non-zero. In fact we can lift this result to any  $r > \hat{r}$  by choosing  $\mathbf{u} = (u_1, \dots, u_{\hat{r}}, 0, \dots, 0)$ . Since we can always construct nontrivial spurious FOCPs, even when  $p$  is generic, we need to consider the second-order conditions.*

### 3.2 Pairs of Binary Forms

In this section we present some results on pairs of binary forms that will be helpful for characterizing the sets  $\text{Im}(\mathcal{A}_{\mathbf{u}})$  and  $\text{cone}(\sigma(\ker(\mathcal{A}_{\mathbf{u}})))$  and proving Theorem 1.1 in Section 4. From now on we assume that  $r = 2$  and  $\mathbf{u} = (u_1, u_2)$ .

The following lemma is a restatement of Bézout’s lemma for univariate polynomials using our notation. It states that the map  $\mathcal{A}_{\mathbf{u}}$  can generate any form in  $\mathbb{R}[x]_{2\mathbf{d}}$  as long as  $u_1$  and  $u_2$  are coprime. We provide a proof below for completeness.

**Lemma 3.2.** *Given  $\mathbf{u} = (u_1, u_2) \in \mathbb{R}[x]_{\mathbf{d}_1}^2$  and  $d_2 \geq d_1 - 1$ , consider the map  $\mathcal{A}_{\mathbf{u}} : \mathbb{R}[x]_{\mathbf{d}_2}^2 \rightarrow \mathbb{R}[x]_{\mathbf{d}_1 + \mathbf{d}_2}$ . Then  $u_1$  and  $u_2$  are coprime if and only if  $\text{Im}(\mathcal{A}_{\mathbf{u}}) = \mathbb{R}[x]_{\mathbf{d}_1 + \mathbf{d}_2}$ .*

*Proof.* For the “only if” direction, when  $u_1$  and  $u_2$  are not coprime, then they share a common factor  $w$  with degree greater than 1, and so does every form in  $\text{Im}(\mathcal{A}_{\mathbf{u}})$ . Thus  $\text{Im}(\mathcal{A}_{\mathbf{u}})$  is not equal to  $\mathbb{R}[x]_{\mathbf{d}_1 + \mathbf{d}_2}$ .

Next we prove the “if” direction. Since  $\mathcal{A}_{\mathbf{u}}(\mathbf{v}) = v_1 u_1 + v_2 u_2$ , for every  $\mathbf{v} \in \ker(\mathcal{A}_{\mathbf{u}})$  we have

$$u_1 v_1 = -u_2 v_2. \quad (10)$$

If  $d_2 \leq d_1$ , since  $u_1$  and  $u_2$  are coprime, by evaluating (10) on the roots of  $u_1$  and  $u_2$  we can conclude that  $v_1 = v_2 = 0$  and  $\dim(\ker(\mathcal{A}_{\mathbf{u}})) = 0$ . Otherwise if  $d_2 > d_1$ , this implies that  $u_1$  is a divisor of  $v_2$  and  $u_2$  is a divisor of  $v_1$ . So there exists  $w_1, w_2 \in \mathbb{R}[x]_{\mathbf{d}_2 - \mathbf{d}_1}$  such that  $v_1 = w_1 u_2, v_2 = w_2 u_1$ . Then (10) becomes  $(w_1 + w_2)u_1 u_2 = 0$ , implying that  $w_2 = -w_1$ . Thus  $\ker(\mathcal{A}_{\mathbf{u}}) = \{(w u_2, -w u_1) \mid w \in \mathbb{R}[x]_{\mathbf{d}_2 - \mathbf{d}_1}\}$  and  $\dim(\ker(\mathcal{A}_{\mathbf{u}})) = \dim(\mathbb{R}[x]_{\mathbf{d}_2 - \mathbf{d}_1})$ . In summary,  $d_2 \geq d_1 - 1$  implies that  $\dim(\ker(\mathcal{A}_{\mathbf{u}})) = d_2 + 1 - d_1$ . Therefore by the rank-nullity theorem,

$$\dim(\text{Im}(\mathcal{A}_{\mathbf{u}})) = 2 \dim(\mathbb{R}[x]_{\mathbf{d}_2}) - \dim(\ker(\mathcal{A}_{\mathbf{u}})) = \dim(\mathbb{R}[x]_{\mathbf{d}_1 + \mathbf{d}_2}).$$

Since  $\text{Im}(\mathcal{A}_{\mathbf{u}}) \subseteq \mathbb{R}[x]_{\mathbf{d}_1 + \mathbf{d}_2}$  and these two sets have the same dimensions, they must be equal.  $\square$

In particular, Lemma 3.2 motivates the decomposition  $u_1 = u'_1 \hat{g}$  and  $u_2 = u'_2 \hat{g}$ , where  $u'_1$  and  $u'_2$  are coprime and  $\hat{g} = \gcd(u_1, u_2)$ . Indeed when  $\hat{g} = 1$  we can apply the sufficient condition

(C1) in Section 4.1 to show that  $\mathbf{u}$  is not a SOCP. When  $\gcd(\hat{g}, \sigma(\mathbf{u}')) = 1$  we can apply (C2) in Section 4.2. Otherwise we need to decompose  $\hat{g} = gh$  where  $h$  do not share common roots with  $\sigma(\mathbf{u}')$ , and apply (C3) in Section 4.3.

**Proposition 3.3.** *Given  $\mathbf{u} = (u_1, u_2) \in \mathbb{R}[x]_{\mathbf{d}}^2$ , we can always find  $g \in \mathbb{R}[x]_{\mathbf{m}}$ ,  $h \in \mathbb{R}[x]_{\mathbf{k}}$  and  $\mathbf{u}' = (u'_1, u'_2) \in \mathbb{R}[x]_{\mathbf{d}-\mathbf{m}-\mathbf{k}}^2$  so that*

$$\begin{aligned} (u_1, u_2) &= (u'_1 gh, u'_2 gh) \\ \gcd(u_1, u_2) &= gh \\ \gcd(u'_1, u'_2) &= 1 \\ \gcd(\sigma(\mathbf{u}'), g) &= \gcd(u_1'^2 + u_2'^2, g) = 1 \\ r \text{ is a (possibly complex) root of } h &\Rightarrow r \text{ is a root of } \sigma(\mathbf{u}'). \end{aligned}$$

Moreover, this decomposition is unique up to multiplication by constants.

*Proof.* Let  $\hat{g} = \gcd(u_1, u_2)$  so that  $u'_1 = u_1/\hat{g}$  and  $u'_2 = u_2/\hat{g}$ . By partitioning the roots of  $\hat{g}$  we can decompose  $\hat{g} = gh$ , where  $g$  has no common roots with  $\sigma(\mathbf{u}')$  and every root of  $h$  is also a root of  $\sigma(\mathbf{u}')$ . The uniqueness of this decomposition follows from the unique factorization theorem for binary forms.  $\square$

We demonstrate this decomposition with an example.

**Example 3.4.** *Let  $u_1(x_1, x_2) = (x_1^2 + x_2^2)^2 x_1^2$  and  $u_2(x_1, x_2) = (x_1^2 + x_2^2)^2 x_1 x_2$ . Then the decomposition in Proposition 3.3 gives  $u'_1 = x_1$ ,  $u'_2 = x_2$ ,  $g = x_1$  and  $h = (x_1^2 + x_2^2)^2$ .*

The following observation about the roots of  $\sigma(\mathbf{u}')$  and  $h$  is useful in our proofs.

**Proposition 3.5.** *In the decomposition of Proposition 3.3, both  $\sigma(\mathbf{u}')$  and  $h$  have no real roots and  $\deg(h)$  is even.*

*Proof.* If  $\sigma(\mathbf{u}')$  has a real root  $\hat{x}$ , then  $u'_1(\hat{x})^2 + u'_2(\hat{x})^2 = 0$ . Thus  $\hat{x}$  is a root of  $u'_1$  and  $u'_2$ , contradicting the fact that  $\gcd(u'_1, u'_2) = 1$ . Thus every root of  $h$  is complex, and  $h$  must have even degree.  $\square$

### 3.3 Main Lemma

In our proof of Theorem 1.1 we will use following main result, which may be of independent interest.

**Lemma 3.6.** *Given binary forms  $g \in \mathbb{R}[x]_{\mathbf{m}}$ ,  $q \in \Sigma[x]_{2\mathbf{d}-2\mathbf{m}}$  and  $p \in \Sigma[x]_{2\mathbf{d}}$ , if  $g$  and  $q$  are coprime then there exists a sum of squares binary form  $s \in \Sigma[x]_{2\mathbf{m}}$  such that*

$$p \equiv sq \pmod{g}.$$

The main ingredient of the proof of Lemma 3.6 is a result stating that any univariate polynomial  $a(x)$  strictly positive on the real zeros of  $g(x)$  can be written as a single square<sup>3</sup> modulo  $g(x)$ .

<sup>3</sup>The number of squares is not important in our proof of Lemma 3.6, we only need the property that  $a(x)$  can be written as a sum of squares modulo  $g(x)$ .

**Proposition 3.7.** *Let  $g(x)$  and  $a(x)$  be coprime univariate polynomials where  $\deg(g) = m$ . If  $a(x) > 0$  for all  $\{x \in \mathbb{R} \mid g(x) = 0\}$  then there exists a polynomial  $t \in \mathbb{R}[x]_m$  such that*

$$a \equiv t^2 \pmod{g}.$$

This result is related to Schmüdgen's certificate [Sch91], which states that if a polynomial is strictly positive on a compact semialgebraic set, then it has a Positivstellensatz certificate in terms of the equations describing the set. We prove Proposition 3.7 using Hermite interpolation on the series expansion of  $\sqrt{a(x)}$  around the roots of  $g$ .

*Proof of Proposition 3.7.* Let  $r_i$  be the roots (possibly complex) of  $g(x)$ , each with multiplicity  $n_i$ , so that  $\sum_i n_i = m$ . Consider the Taylor series expansion of  $f(x) = \sqrt{a(x)}$  centered at  $r_i$ . Since  $a$  and  $g$  do not share any common roots as they are coprime, this Taylor series is well defined around any root of  $g$ . Let the polynomials  $\gamma_i(x)$  be the first  $n_i$  terms of the Taylor expansion of  $f(x)$  centered at  $r_i$ . The polynomials  $\gamma_i$  have real coefficients if  $r_i$  is real, and if  $r_i$  and  $r_j$  are a pair of conjugate roots,  $\gamma_i = \bar{\gamma}_j$ . We can then use the Chinese Remainder Theorem [DF03, Section 7.6] to construct the unique polynomial  $t(x)$  with real coefficients and  $\deg(t) < m$  such that

$$\forall i, \quad t(x) \equiv \gamma_i(x) \pmod{(x - r_i)^{n_i}}.$$

By construction, for all roots  $r_i$  of  $g$  and any  $k = 0, \dots, n_i - 1$ , we have

$$\frac{d^k}{dx^k} f(r_i) = \frac{d^k}{dx^k} \gamma_i(r_i) = \frac{d^k}{dx^k} t(r_i).$$

For each root  $r_i$ , we have

$$\sqrt{a(r_i)} = f(r_i) = \gamma_i(r_i) = t(r_i).$$

and

$$\frac{d}{dx} a(r_i) = 2f(r_i) \frac{d}{dx} f(r_i) = 2t(r_i) \frac{d}{dx} t(r_i) = \frac{d}{dx} t(r_i)^2.$$

By induction we get  $\frac{d^k}{dx^k} a(r_i) = \frac{d^k}{dx^k} t(r_i)^2$  for  $k = 0, \dots, n_i - 1$ . This is a generalization of Hermite interpolation for a variable number of consecutive derivatives at each point [Ham73, section 17.6]<sup>4</sup>. Since  $a(x)$  and  $t(x)^2$  match at  $m$  points in total (counting derivatives), we have shown that  $a \equiv t^2 \pmod{g}$ .  $\square$

Then we prove the affine version of Lemma 3.6.

**Lemma 3.8.** *Given univariate polynomials  $g, p$  and  $q$  where  $\deg(g) = m$ ,  $\deg(p) = 2d$  and  $\deg(q) = 2d - 2m$ ,  $p$  and  $q$  are sum of squares,  $g$  and  $q$  are coprime. Then there exists a sum of squares polynomial  $s \in \Sigma[x]_{2m}$  such that*

$$p \equiv sq \pmod{g}.$$

---

<sup>4</sup>This is referred to as Birkhoff interpolation in [Ham73], and the existence of a unique interpolating polynomial crucially depends on the use of consecutive derivatives.

*Proof.* Since  $q$  is coprime with  $g$ , Bézout's lemma guarantees that there exists a polynomial  $a \in \mathbb{R}[x]_m$  such that

$$aq \equiv 1 \pmod{g}. \quad (11)$$

Since  $q$  is nonnegative and coprime with  $g$ , for all  $x \in \mathbb{R}$  where  $g(x)$  vanishes,  $q(x) > 0$  and thus  $a(x) > 0$ . Since  $a(x)q(x) = 1$  for all roots  $x$  of  $g$ ,  $a$  is also coprime with  $g$ . Then we can apply Proposition 3.7 to find  $t \in \mathbb{R}[x]_m$  so that  $a \equiv t^2 \pmod{g}$ . Multiplying both sides of (11) by  $p$ , we get

$$p \equiv t^2 pq \pmod{g}.$$

Since  $t^2 p$  is a sum of squares, we can reduce each squared polynomial modulo  $g$  to get  $s \in \Sigma[x]_{2m}$ .  $\square$

Finally we prove Lemma 3.6, which is the projective version of Lemma 3.8.

*Proof of Lemma 3.6.* We first apply a linear change of coordinates so that  $(0, 1)$  is not a root of  $g$ ,  $p$  or  $q$ . Then let  $g'(x) = g(x, 1)$ ,  $p'(x) = p(x, 1)$  and  $q'(x) = q(x, 1)$ . Since this dehomogenization procedure preserves the degree of  $g, p$  and  $q$ , we can apply Lemma 3.8 to find polynomials  $s' \in \Sigma[x]_{2m}$  and  $t' \in \mathbb{R}[x]_{2d-m}$  so that

$$p' = s'q' + t'g'.$$

We can then homogenize by letting  $s(x_1, x_2) = x_2^{2m}s'(x_1/x_2)$  and  $t(x_1, x_2) = x_2^{2d-m}t'(x_1/x_2)$ . Thus  $s$  is also a sum of squares and

$$p = sq + tg.$$

$\square$

## 4 Main Theorem and Proof

In this section we prove Theorem 1.1, which states that for univariate polynomials, a rank-2 decomposition has no spurious second-order critical points. Using the decomposition in Proposition 3.3, we first prove simplified versions of Theorem 1.1 in Sections 4.1 and 4.2, before proving the full version in Section 4.3.

### 4.1 Coprime Case: $g = 1, h = 1$

This is the case explained in the introduction. In the decomposition of Proposition 3.3,  $g = h = 1$  implies that  $u_1$  and  $u_2$  are coprime. This happens generically and implies that for a fixed  $p$ , the gradient condition (8) is sufficient for almost all  $\mathbf{u}$ .

**Proposition 4.1.** *Suppose  $\mathbf{u} \in \mathbb{R}[x]_{\mathbf{d}}^2$  and  $p \in \Sigma[x]_{2\mathbf{d}}$  satisfies  $\nabla f_p(\mathbf{u}) = 0$ . If*

$$p \in \text{Im}(\mathcal{A}_{\mathbf{u}}), \quad (C1)$$

*then we have  $f_p(\mathbf{u}) = 0$ .*

*Proof.* Since  $p \in \text{Im}(\mathcal{A}_{\mathbf{u}})$ , we can find  $\mathbf{v} \in \mathbb{R}[x]_{\mathbf{d}}^2$  so that  $\mathcal{A}_{\mathbf{u}}(\mathbf{v}) = \sigma(\mathbf{u}) - p$ . But this combined with the gradient condition (8) implies that  $f_p(\mathbf{u}) = \|\sigma(\mathbf{u}) - p\|^2 = 0$ .  $\square$

Since Lemma 3.2 implies that  $\text{Im}(\mathcal{A}_{\mathbf{u}}) = \mathbb{R}[x]_{2\mathbf{d}}$  if and only if  $u_1$  and  $u_2$  are coprime, we have shown that when  $g = h = 1$ , we always have  $p \in \text{Im}(\mathcal{A}_{\mathbf{u}})$  and there are no spurious FOCPs and SOCPs.

## 4.2 Special Case: $h = 1$

When  $u_1$  and  $u_2$  are not coprime,  $\sigma(\mathbf{u}) - p$  might not be in  $\text{Im}(\mathcal{A}_{\mathbf{u}})$  and we cannot use the argument in Proposition 4.1. Thus we need to use make use of the Hessian condition (9).

**Proposition 4.2.** *Suppose  $\mathbf{u} \in \mathbb{R}[x]_{\mathbf{d}}^2$  and  $p \in \Sigma[x]_{2\mathbf{d}}$  satisfies  $\nabla f_p(\mathbf{u}) = 0$  and  $\nabla^2 f_p(\mathbf{u}) \succeq 0$ . If*

$$p \in \text{Im}(\mathcal{A}_{\mathbf{u}}) + \text{cone}(\sigma(\ker(\mathcal{A}_{\mathbf{u}}))), \quad (\text{C2})$$

*then we have  $f_p(\mathbf{u}) = 0$ .*

This means that if for all  $\mathbf{u}$  we can decompose  $p = q + r$  where  $q \in \text{Im}(\mathcal{A}_{\mathbf{u}})$  and  $r \in \text{cone}(\sigma(\ker(\mathcal{A}_{\mathbf{u}})))$ , then  $f_p(\mathbf{u})$  has no spurious SOCPs. In particular, similar to how  $\text{Im}(\mathcal{A}_{\mathbf{u}})$  is related to the gradient condition (8) in Proposition 4.1,  $\text{cone}(\sigma(\ker(\mathcal{A}_{\mathbf{u}})))$  is related to the Hessian condition (9). The following result states that (C2) is satisfied if  $h = 1$ .

**Lemma 4.3.** *Given  $\mathbf{u} \in \mathbb{R}[x]_{\mathbf{d}}^2$ , if in the decomposition of Proposition 3.3 we have  $h = 1$  then for every  $p \in \Sigma[x]_{2\mathbf{d}}$ ,*

$$p \in \text{Im}(\mathcal{A}_{\mathbf{u}}) + \text{cone}(\sigma(\ker(\mathcal{A}_{\mathbf{u}}))).$$

*Proof.* We want to show that any  $p(x) \in \Sigma[x]_{2\mathbf{d}}$  can be written as the sum of polynomials in  $\text{Im}(\mathcal{A}_{\mathbf{u}})$  and  $\sigma(\ker(\mathcal{A}_{\mathbf{u}}))$ . Therefore it is useful to have a characterization of these sets. Lemma 3.2 tells us that

$$\begin{aligned} \text{Im}(\mathcal{A}_{\mathbf{u}}) &= \{\mathcal{A}_{\mathbf{u}'}(\mathbf{v})g \mid \mathbf{v} \in \mathbb{R}[x]_{\mathbf{d}}\} \\ &= \{wg \mid w \in \mathbb{R}[x]_{2\mathbf{d}-\mathbf{m}}\}. \end{aligned}$$

Since for all  $t \in \mathbb{R}[x]_{\mathbf{m}}$  we have  $(-tu'_2, tu'_1) \in \ker(\mathcal{A}_{\mathbf{u}})$ , thus

$$\begin{aligned} \{t^2\sigma(\mathbf{u}') \mid t \in \mathbb{R}[x]_{\mathbf{m}}\} &\subseteq \sigma(\ker(\mathcal{A}_{\mathbf{u}})) \\ \{s\sigma(\mathbf{u}') \mid s \in \Sigma[x]_{2\mathbf{m}}\} &\subseteq \text{cone}(\sigma(\ker(\mathcal{A}_{\mathbf{u}}))). \end{aligned} \quad (12)$$

Since  $\sigma(\mathbf{u}')$  is coprime with  $g$  by assuming  $h = 1$ , we can apply Lemma 3.6 to show that there exists  $w \in \mathbb{R}[x]_{2\mathbf{d}-\mathbf{m}}$  and  $s \in \Sigma[x]_{2\mathbf{m}}$  such that  $p = s\sigma(\mathbf{u}') + wg$ .  $\square$

Finally we prove Proposition 4.2.

*Proof of Proposition 4.2.* The condition (C2) implies that there exist  $\mathbf{v} \in \mathbb{R}[x]_{\mathbf{d}}^2$ ,  $\mathbf{w}^{(i)} \in \ker(\mathcal{A}_{\mathbf{u}})$  such that

$$p = \mathcal{A}_{\mathbf{u}}(\mathbf{v}) + \sum_i \sigma(\mathbf{w}^{(i)}).$$

Since  $\nabla f_p(\mathbf{u}) = 0$ , (8) implies that

$$\langle \mathcal{A}_{\mathbf{u}}(\mathbf{v}), \sigma(\mathbf{u}) - p \rangle = 0. \quad (13)$$

Since  $\nabla^2 f_p(\mathbf{u}) \succeq 0$  and  $\mathbf{w}^{(i)} \in \ker(\mathcal{A}_{\mathbf{u}})$ , (9) implies that

$$\langle \sigma(\mathbf{w}^{(i)}), \sigma(\mathbf{u}) - p \rangle \geq 0. \quad (14)$$

Combining (13) and (14) gives

$$\langle p, \sigma(\mathbf{u}) - p \rangle \geq 0. \quad (15)$$

Since  $\nabla f_p(\mathbf{u}) = 0$  implies that

$$\langle \mathcal{A}_{\mathbf{u}}(\mathbf{u}), \sigma(\mathbf{u}) - p \rangle = \langle \sigma(\mathbf{u}), \sigma(\mathbf{u}) - p \rangle = 0,$$

we have

$$f_p(\mathbf{u}) = \|\sigma(\mathbf{u}) - p\|^2 = \langle \sigma(\mathbf{u}) - p, \sigma(\mathbf{u}) - p \rangle = -\langle p, \sigma(\mathbf{u}) - p \rangle.$$

This together with (15) implies that  $f_p(\mathbf{u}) \leq 0$ . However  $f_p(\mathbf{u})$  is always non-negative, thus it must be 0.  $\square$

### 4.3 General Case

Lemma 4.3 alone is insufficient to prove Theorem 1.1. It is possible for  $\hat{g} = \gcd(u_1, u_2)$  to share complex roots with  $\sigma(\mathbf{u}')$  (recall from Proposition 3.5 that all roots of  $\sigma(\mathbf{u}')$  are complex), as seen in Example 3.4. Hence the argument in the proof of Lemma 4.3 fails as  $\hat{g}$  is not coprime with  $\sigma(\mathbf{u}')$ . To get around this issue, we will derive the sufficient condition (C3) in Proposition 4.4, a stronger version of (C2), by carefully examining the Hessian condition (9). Roughly speaking, Proposition 4.4 shows that we can replace every root of  $h$  (which must be complex) with any real root<sup>5</sup>. Since  $\sigma(\mathbf{u}')$  has no real roots,  $\hat{g}x_1^k/h$  is now coprime with  $\sigma(\mathbf{u}')$ , and we can then complete the proof by following the argument in the previous section.

**Proposition 4.4.** *Suppose  $\mathbf{u} \in \mathbb{R}[x]_{\mathbf{d}}^2$  and  $p \in \Sigma[x]_{2\mathbf{d}}$  satisfies  $\nabla f_p(\mathbf{u}) = 0$  and  $\nabla^2 f_p(\mathbf{u}) \succeq 0$ , with the decomposition in Proposition 3.3 where  $k = \deg(h)$ . If*

$$p \in \text{Im} \left( \mathcal{A}_{\mathbf{u}x_1^k/h} \right) + \text{cone} \left( \Sigma \left( \ker(\mathcal{A}_{\mathbf{u}}) \right) \right), \quad (C3)$$

then  $f_p(\mathbf{u}) = 0$ .

*Proof.* We first prove that if  $r \in \mathbb{R}[x]_{\ell}$  is a common divisor of  $\sigma(\mathbf{u}')$  and  $\hat{g} = \gcd(u_1, u_2)$ , then for all  $\mathbf{b} = (b_1, b_2) \in \mathbb{R}[x]_{\mathbf{d}}^2$  we have

$$\langle \mathcal{A}_{\mathbf{u}}(\mathbf{b})x_1^{\ell}/r, \sigma(\mathbf{u}) - p \rangle = 0. \quad (16)$$

---

<sup>5</sup>Without loss of generality we choose this real root to be  $x_1$ .

Given any  $b_1, b_2 \in \mathbb{R}[x]_{\mathbf{d}}$  and  $\eta \in \mathbb{R}$ , let  $v_1 = \eta x_1^\ell u_2/r + b_2$  and  $v_2 = -\eta x_1^\ell u_1/r - b_1$ . We have

$$\begin{aligned}\mathcal{A}_{\mathbf{u}}(\mathbf{v}) &= u_1(\eta x_1^\ell u_2/r + b_2) - u_2(\eta x_1^\ell u_1/r + b_1) = u_1 b_2 - u_2 b_1 \\ \sigma(\mathbf{v}) &= \eta^2 x_1^{2\ell} (u_1^2 + u_2^2)/r^2 + 2\eta x_1^\ell (b_1 u_1 + b_2 u_2)/r + (b_1^2 + b_2^2).\end{aligned}$$

Since  $r$  is a divisor of both  $\sigma(\mathbf{u}')$  and  $\hat{g}$ ,  $x_1^{2\ell}(u_1^2 + u_2^2)/r^2 = x_1^{2\ell}\sigma(\mathbf{u}')\hat{g}^2/r^2$  is a multiple of  $\hat{g}$ . Thus  $x_1^{2\ell}(u_1^2 + u_2^2)/r^2 \in \text{Im}(\mathcal{A}_{\mathbf{u}})$  and we have

$$\left\langle \eta^2 x_1^{2\ell} (u_1^2 + u_2^2)/r^2, \sigma(\mathbf{u}) - p \right\rangle = 0.$$

Therefore, the Hessian condition (9) implies that for all  $\eta \in \mathbb{R}$ ,

$$2\eta \left\langle (b_1 u_1 + b_2 u_2) x_1^\ell / r, \sigma(\mathbf{u}) - p \right\rangle + \langle b_1^2 + b_2^2, \sigma(\mathbf{u}) - p \rangle + 2 \|u_1 b_2 - u_2 b_1\|^2 \geq 0. \quad (17)$$

This implies the identity (16) otherwise there exists  $\eta$  such that (17) is negative.

Since  $\sigma(\mathbf{u}')$  and  $h$  have no real roots (Proposition 3.5), we can write  $h = \prod_{i=1}^{k/2} r_i$ , where each  $r_i \in \mathbb{R}[x]_{\mathbf{2}}$  is a quadratic form corresponding to the product of a pair of complex roots. We then iteratively apply the same argument from the above proof to show that for every  $1 \leq k' \leq k/2$  and  $\mathbf{b} \in \mathbb{R}[x]_{\mathbf{d}}^2$ ,

$$\left\langle \mathcal{A}_{\mathbf{u}}(\mathbf{b}) x_1^{2k'} / \prod_{i=1}^{k'} r_i, \sigma(\mathbf{u}) - p \right\rangle = 0.$$

This is because each  $r_i$  is a divisor of  $\sigma(\mathbf{u}')$  and  $\prod_{i=1}^{k'} r_i$  divides  $\hat{g}$ . From here we can finish our proof by following the same steps as in the proof of Proposition 4.2.  $\square$

With Proposition 4.4 we can prove Theorem 1.1, by showing that every  $p \in \Sigma[x]_{\mathbf{2d}}$  has the required decomposition.

*Proof of Theorem 1.1.* Since  $\langle \mathcal{A}_{\mathbf{u}'}(\mathbf{b}) g x_1^k, \sigma(\mathbf{u}) - p \rangle = 0$  for all  $\mathbf{b} \in \mathbb{R}[x]_{\mathbf{d}}^2$  and  $u'_1, u'_2$  are coprime, Lemma 3.2 implies that  $\langle w g x_1^k, \sigma(\mathbf{u}) - p \rangle = 0$  for all  $w \in \mathbb{R}[x]$ . Since  $\sigma(\mathbf{u}')$  has no real roots (Proposition 3.5), it is coprime with  $x_1^k$ . As  $\sigma(\mathbf{u}')$  is coprime with  $g$ , it is also coprime with  $g x_1^k$ . Thus Lemma 3.6 tells us that there exists a sum of squares polynomial  $s$  such that  $p \equiv s \sigma(\mathbf{u}') \pmod{g x_1^k}$ . Since  $s \sigma(\mathbf{u}') \in \text{cone}(\sigma(\ker(\mathcal{A}_{\mathbf{u}})))$  by (12), we are done.  $\square$

## 5 Geometric Interpretation and Certificates

In this section, we provide a geometric interpretation of our proof of Theorem 1.1, which allows us to turn the proof into a certificate. In order to prove that there is no spurious second-order critical points when minimizing  $f_p(\mathbf{u}) = \|\sigma(\mathbf{u}) - p\|^2$ , we have to show that for all  $\mathbf{u} \in \mathbb{R}[x]^r$  and for all  $p \in \Sigma[x]$ ,  $\nabla f_p(\mathbf{u}) = 0$  and  $\nabla^2 f_p(\mathbf{u}) \succeq 0$  implies that  $f_p(\mathbf{u}) = 0$  and  $p = \sigma(\mathbf{u})$ . One way to tackle this problem is to fix  $p$  then characterize the set of  $\mathbf{u}$  satisfying the second-order critical point conditions. This is the approach taken by [BBJN18] and related works, where they used an argument based on the dimension of the subspace generated by the constraints of the SDP. However the SOCP conditions are non-convex in  $\mathbf{u}$ . In order to do better than a dimension-counting argument, our proof takes a different approach. If we fix  $\mathbf{u}$ , the set of all  $p$  satisfying the gradient condition (8) is an affine subspace, whereas the set of all  $p \in \Sigma[x]$  satisfying the Hessian condition (9) is a convex




Figure 1: The geometric interpretation

semidefinite-representable set. We need to show that these two sets intersect at only one point,  $p = \sigma(\mathbf{u})$  (see picture below).

Our proof can be interpreted as constructing a certificate to show that these two sets only intersect at one point. This is true if and only if the following optimization problem has a zero optimal objective value:

$$\max_{p \in \Sigma[x]} \min_{Q \succeq 0, \boldsymbol{\lambda} \in \mathbb{R}[x]^r} \|\sigma(\mathbf{u}) - p\|^2 + \nabla f_p(\mathbf{u})(\boldsymbol{\lambda}) + \langle Q, \nabla^2 f_p(\mathbf{u}) \rangle \quad (18)$$

Expanding the gradient and Hessian, we get

$$\begin{aligned} \nabla f_p(\mathbf{u})(\boldsymbol{\lambda}) &= \langle \mathcal{A}_{\mathbf{u}}(\boldsymbol{\lambda}), \sigma(\mathbf{u}) - p \rangle \\ \langle Q, \nabla^2 f_p(\mathbf{u}) \rangle &= \sum_i \langle \sigma(\mathbf{v}^{(i)}), \sigma(\mathbf{u}) - p \rangle + 2 \left\| \mathcal{A}_{\mathbf{u}}(\mathbf{v}^{(i)}) \right\|^2 \end{aligned}$$

where  $Q = \sum_i \mathbf{v}^{(i)} \mathbf{v}^{(i)\top}$ . If for every  $p \in \Sigma[x]$  we can find  $\boldsymbol{\lambda}$  and  $Q$  such that

$$\nabla f_p(\mathbf{u})(\boldsymbol{\lambda}) + \langle Q, \nabla^2 f_p(\mathbf{u}) \rangle = -\|\sigma(\mathbf{u}) - p\|^2, \quad (19)$$

then the objective of (18) is at most 0 and cannot be positive, showing that  $p = \sigma(\mathbf{u})$  is the only point satisfying the gradient and Hessian conditions. Since  $\mathcal{A}_{\mathbf{u}}(\mathbf{u}) = \sigma(\mathbf{u})$ , this is equivalent to finding  $\boldsymbol{\lambda}$  and  $Q$  such that  $\nabla f_p(\mathbf{u})(\boldsymbol{\lambda}) + \langle Q, \nabla^2 f_p(\mathbf{u}) \rangle = \langle p, \sigma(\mathbf{u}) - p \rangle$ .

## 5.1 Warmup

As a warmup, we construct such a certificate if  $h = 1$  in the decomposition of  $\mathbf{u}$  in Proposition 3.3. Recall that in this case  $g = \gcd(u_1, u_2)$ ,  $u_1 = gu'_1$ ,  $u_2 = gu'_2$  and  $\sigma(\mathbf{u}')$  is coprime with  $g \in \mathbb{R}[x]_{\mathbf{m}}$ . Therefore, by Lemma 3.6, there exists  $s \in \Sigma[x]_{2\mathbf{m}}$  such that  $p \equiv s\sigma(\mathbf{u}') \pmod{g}$ . Let

$$Q = s \begin{bmatrix} u_2'^2 & -u_1' u_2' \\ -u_1' u_2' & u_1'^2 \end{bmatrix}.$$

As both  $\sigma(\mathbf{u})$  and  $p - s\sigma(\mathbf{u}')$  are divisible by  $g$ , Lemma 3.2 implies that there exists  $\boldsymbol{\lambda}$  such that:

$$\mathcal{A}_{\mathbf{u}}(\boldsymbol{\lambda}) = -\sigma(\mathbf{u}) + p - s\sigma(\mathbf{u}').$$

These values of  $\boldsymbol{\lambda}$  and  $Q$  give

$$\begin{aligned}\nabla f_p(\mathbf{u})(\boldsymbol{\lambda}) &= -\|\sigma(\mathbf{u}) - p\|^2 - \langle s\sigma(\mathbf{u}'), \sigma(\mathbf{u}) - p \rangle \\ \langle Q, \nabla^2 f_p(\mathbf{u}) \rangle &= \langle s\sigma(\mathbf{u}'), \sigma(\mathbf{u}) - p \rangle,\end{aligned}$$

hence taking the sum we have the identity (19).

## 5.2 Certificate

Now we can present the proof of Theorem 1.1 in the form of a certificate. First we decompose  $\mathbf{u} \in \mathbb{R}[x]_{\mathbf{d}}^2$  as in Proposition 3.3. Since  $g$  is coprime with  $\sigma(\mathbf{u}')$  and  $\sigma(\mathbf{u}')$  has no real roots,  $gx_1^k$  is also coprime with  $\sigma(\mathbf{u}')$ . Then by Lemma 3.6, there exists  $s \in \Sigma[x]_{2(\mathbf{m}+\mathbf{k})}$  such that  $p \equiv s\sigma(\mathbf{u}') \pmod{gx_1^k}$ . As  $p - s\sigma(\mathbf{u}')$  is divisible by  $gx_1^k$ , Lemma 3.2 implies that there exists  $\mathbf{b}^0 \in \mathbb{R}[x]_{\mathbf{d}}^2$  so that:

$$2\frac{x_1^k}{h}\mathcal{A}_{\mathbf{u}}(\mathbf{b}^0) = 2gx_1^k\mathcal{A}_{\mathbf{u}'}(\mathbf{b}^0) = p - s\sigma(\mathbf{u}').$$

As in the proof of Proposition 4.4, we write  $h = \prod_{i=1}^{k/2} r_i$ . For every  $1 \leq j \leq k/2$ , since  $r_j$  divides  $\sigma(\mathbf{u}')$ , by Lemma 3.2 there exists  $\mathbf{b}^j \in \mathbb{R}[x]_{\mathbf{d}}^2$  so that

$$2gx_1^{k-2j}\mathcal{A}_{\mathbf{u}'}(\mathbf{b}^j) \prod_{i=1}^j r_i = -g^2x_1^{2k-4(j-1)}\sigma(\mathbf{u}') \prod_{i=1}^{j-1} r_i^2.$$

Given any  $\mathbf{a} = (a_1, a_2) \in \mathbb{R}[x]_{\mathbf{d}}^2$ , we define  $\bar{\mathbf{a}} := (a_2, -a_1)$ . Given a parameter  $\eta \in \mathbb{R}$ , for every  $0 \leq j \leq k/2$  let  $\eta_j = \eta^{3^j}$  and

$$\mathbf{v}^j = \eta_j^{3/2}gx_1^{k-2j}\bar{\mathbf{u}}' \prod_{i=1}^j r_i + \eta_j^{-1/2}\bar{\mathbf{b}}^j.$$

Then define

$$\begin{aligned}Q &= s\bar{\mathbf{u}}'\bar{\mathbf{u}}'^{\top} + \sum_{j=0}^{k/2} \mathbf{v}^j \mathbf{v}^{j\top} \\ \boldsymbol{\lambda} &= -(1 + \eta_{k/2+1})\mathbf{u}.\end{aligned}$$

Since  $\mathcal{A}_{\mathbf{u}}(\bar{\mathbf{u}}') = 0$ ,  $\sigma(\bar{\mathbf{u}}') = \sigma(\mathbf{u})$ ,  $\sigma(\bar{\mathbf{b}}^j) = \sigma(\mathbf{b}^j)$ ,  $\mathcal{A}_{\bar{\mathbf{u}}'}(\bar{\mathbf{b}}^j) = \mathcal{A}_{\mathbf{u}'}(\mathbf{b}^j)$  and  $\eta_j^3 = \eta_{j+1}$ , we have

$$\begin{aligned}\sum_{j=0}^{k/2} \sigma(\mathbf{v}^j) &= \sum_{j=0}^{k/2} \left( \eta_j^3 g^2 x_1^{2k-4j} \prod_{i=1}^j r_i^2 \sigma(\mathbf{u}') + 2\eta_j g x_1^{k-2j} \prod_{i=1}^j r_i \mathcal{A}_{\mathbf{u}'}(\mathbf{b}^j) + \eta_j^{-1} \sigma(\mathbf{b}^j) \right) \\ &= p - s\sigma(\mathbf{u}') + \eta_{k/2+1} \sigma(\mathbf{u}) + \sum_{j=0}^{k/2} \eta_j^{-1} \sigma(\mathbf{b}^j),\end{aligned}$$

$$\mathcal{A}_{\mathbf{u}}(\mathbf{v}^j) = \eta_j^{-1/2} \mathcal{A}_{\mathbf{u}}(\bar{\mathbf{b}}^j).$$

So we have proven the identity

$$\nabla f_p(\mathbf{u})(\boldsymbol{\lambda}) + \langle Q, \nabla^2 f_p(\mathbf{u}) \rangle = -\|\sigma(\mathbf{u}) - p\|^2 + \sum_{j=0}^{k/2} \frac{1}{\eta_j} \left( \langle \sigma(\mathbf{b}^j), \sigma(\mathbf{u}) - p \rangle + \left\| \mathcal{A}_{\mathbf{u}}(\bar{\mathbf{b}}^j) \right\|^2 \right). \quad (20)$$

This implies that for every  $\eta > 0$  and every  $\mathbf{u}$  that satisfies  $\nabla f_p(\mathbf{u}) = 0$  and  $\nabla^2 f_p(\mathbf{u}) \succeq 0$ ,

$$\|\sigma(\mathbf{u}) - p\|^2 \leq \sum_{j=0}^{k/2} \eta^{-3j} \left( \langle \sigma(\mathbf{b}^j), \sigma(\mathbf{u}) - p \rangle + \left\| \mathcal{A}_{\mathbf{u}}(\bar{\mathbf{b}}^j) \right\|^2 \right).$$

Since  $\mathbf{b}^j$  does not depend on  $\eta$ , we can make the right hand side arbitrarily small by taking the limit  $\eta \rightarrow \infty$ . Thus we can conclude that  $\|\sigma(\mathbf{u}) - p\| = 0$ .

## 6 Extensions and Generalizations

The certificate interpretation discussed in the previous section allows us to generalize Theorem 1.1 to other settings, such as projecting onto the sum of squares cone, certifying non-negativity on intervals and imposing linear constraints on coefficients of univariate sum of squares polynomials.

### 6.1 Projection Onto the Sum-of-Squares Cone

A natural question to consider is what happens to the optimization landscape of  $f_p(\mathbf{u})$  when  $p$  cannot be expressed as a sum of squares. In this case the objective  $f_p(\mathbf{u})$  can never be zero, but we show that all SOCPs have the same objective value, which is the projection of  $p$  to the sum of squares cone.

**Corollary 6.1.** *For all  $\mathbf{u} \in \mathbb{R}[x]_{\mathbf{d}}^2$  where  $\nabla f_p(\mathbf{u}) = 0$  and  $\nabla^2 f_p(\mathbf{u}) \succeq 0$ ,  $\sigma(\mathbf{u})$  is the projection of  $p$  to the sum-of-squares cone with respect to the inner product used to define  $f_p$ . In other words,  $f_p(\mathbf{u}) = \|\sigma(\mathbf{u}) - p\|^2 \leq \|q - p\|^2$  for all  $q \in \Sigma[x]_{2\mathbf{d}}$ .*

*Proof.* Corollary 6.1 can be proved by a simple modification of the certificate (20). Although  $p$  is no longer a sum of squares, we can use (20) to show that for all  $q \in \Sigma[x]_{2\mathbf{d}}$ ,

$$\langle \sigma(u) - q, \sigma(u) - p \rangle \leq 0.$$

This is exactly the variational characterization of projection onto the convex cone  $\Sigma[x]_{2\mathbf{d}}$ .  $\square$

### 6.2 Certifying Non-negativity on Intervals

Suppose we wish to certify that a univariate polynomial  $p(x)$  is nonnegative in a union of intervals  $I = \bigcup_{i=1}^m I_i$  where  $I_i = \{x \in \mathbb{R} \mid \alpha_i \leq x \leq \beta_i\}$ . This can be accomplished by finding a decomposition

$$p(x) = \sum_i^m a_i(x)q_i(x),$$

where  $a_i(x)$  are fixed polynomials depending on the intervals  $I_i$  and  $q_i(x)$  are sum of squares polynomials (see e.g. [BPT13, Theorem 3.72]). This objective can also be written as a non-convex optimization problem by the decomposition  $q_i(x) = \sum_{j=1}^r u_{ij}(x)^2 = \sigma(\mathbf{u}_i)$ . If we let

$$s(x) = \sum_{i=1}^m a_i(x)\sigma(\mathbf{u}_i),$$

then the objective  $f_p^I(\mathbf{u})$ , its gradient and Hessian can be written as

$$\begin{aligned} f_p^I(\mathbf{u}) &= \left\| \sum_{i=1}^m a_i \sigma(\mathbf{u}_i) - p \right\|^2 = \|s - p\|^2 \\ \nabla f_p^I(\mathbf{u})(\mathbf{v}) &= \left\langle \sum_{i=1}^m a_i \mathcal{A}_{\mathbf{u}_i}(\mathbf{v}_i), s - p \right\rangle \\ \nabla^2 f_p^I(\mathbf{u})(\mathbf{v}, \mathbf{v}) &= \left\langle \sum_{i=1}^m a_i \sigma(\mathbf{v}_i), s - p \right\rangle + 2 \left\| \sum_{i=1}^m a_i \mathcal{A}_{\mathbf{u}_i}(\mathbf{v}_i) \right\|^2 \end{aligned}$$

**Corollary 6.2.** *Suppose  $r \geq 2$  and we are given  $p = \sum_{i=1}^m a_i q_i \in \mathbb{R}[x]_{2d+k}$  where  $a_i \in \mathbb{R}[x]_k$ ,  $q_i \in \Sigma[x]_{2d}$ . For all  $\mathbf{u} \in \mathbb{R}[x]_d^{m \times r}$  such that  $\nabla f_p^I(\mathbf{u}) = 0$  and  $\nabla^2 f_p^I(\mathbf{u}) \succeq 0$ ,  $f_p^I(\mathbf{u}) = 0$ .*

*Proof.* We can prove this by constructing a certificate of the form (20). For each  $i$  we can choose  $\mathbf{v}_j = 0$  for all  $j \neq i$ , then follow the reasoning in Section 5.2 to find  $\lambda_i$  and  $Q_i$  such that for all  $\eta_i > 0$ ,

$$\nabla f_p^I(\mathbf{u})(\lambda_i) + \langle Q_i, \nabla^2 f_p^I(\mathbf{u}) \rangle = \langle a_i \sigma(\mathbf{u}_i), s - p \rangle + C_i, \quad (21)$$

where  $C_i$  is a value that can be made arbitrarily small by taking a limit. Since  $\langle s, s - p \rangle = \langle \mathbf{u}, \nabla f_p^I(\mathbf{u}) \rangle$ , we can sum (21) for all  $i$  to conclude  $f_p^I(\mathbf{u}) = 0$ .  $\square$

### 6.3 Sum-of-Squares Optimization

More generally, we can consider the problem of finding a feasible point in the intersection of the cone  $\Sigma[x]_{2d}$  with any affine subspace. This allows us to solve Sum-of-Squares optimization problems involving univariate polynomials. Let  $\mathcal{B} : \mathbb{R}[x]_{2d} \rightarrow \mathbb{R}^m$  be a linear map. Given  $b \in \mathbb{R}^m$ , we want to find  $p \in \Sigma[x]_{2d}$  so that  $\mathcal{B}(p) = b$ . This is equivalent to minimizing the quadratic-penalized problem

$$f_{\mathcal{B}}(\mathbf{u}) = \|\mathcal{B}(\sigma(\mathbf{u})) - b\|^2. \quad (22)$$

**Corollary 6.3.** *Suppose there exists  $p \in \Sigma[x]_{2d}$  such that  $\mathcal{B}(p) = b$ . Then  $\nabla f_{\mathcal{B}}(\mathbf{u}) = 0$  and  $\nabla^2 f_{\mathcal{B}}(\mathbf{u}) \succeq 0$  implies that  $f_{\mathcal{B}}(\mathbf{u}) = 0$ .*

*Proof.* The gradient and Hessian of the objective (22) can be written as

$$\begin{aligned} \frac{1}{4} \nabla f_{\mathcal{B}}(\mathbf{u})(\mathbf{v}) &= \langle \mathcal{B}(\mathcal{A}_{\mathbf{u}}(\mathbf{v})), \mathcal{B}(\sigma(\mathbf{u}) - p) \rangle \\ \frac{1}{4} \nabla^2 f_{\mathcal{B}}(\mathbf{u})(\mathbf{v}, \mathbf{v}) &= \langle \mathcal{B}(\sigma(\mathbf{v})), \mathcal{B}(\sigma(\mathbf{u}) - p) \rangle + 2 \|\mathcal{B}(\mathcal{A}_{\mathbf{u}}(\mathbf{v}))\|^2. \end{aligned}$$

Thus by the linearity of  $\mathcal{B}$  we can use the same construction as in the certificate (20) to show that  $f_{\mathcal{B}}(\mathbf{u}) = 0$ .  $\square$

## 7 Implementation and Experiments

In this section we describe an efficient implementation of finding a sum of squares decomposition of trigonometric polynomials. A trigonometric polynomial of degree- $d$  is defined by  $2d + 1$  coefficients and has the form

$$p(t) = a_0 + \sum_{k=1}^d (a_k \cos(kt) + a_{-k} \sin(kt)).$$

By the substitution  $\cos(t) = \frac{1-x^2}{1+x^2}$  and  $\sin(t) = \frac{2x}{1+x^2}$ ,  $p(t)$  a rational function with the denominator a power of  $1+x^2$  and the numerator a degree- $2d$  polynomial in  $x$ . Thus certifying the non-negativity of the numerator is equivalent to certifying the non-negativity of  $p(t)$ . By this correspondence the result of Theorem 1.1 also applies to trigonometric polynomials.

Since the proof of Theorem 1.1 does not depend on the norm used for  $f_p(\mathbf{u}) = \|p - \sum_i u_i^2\|^2$ , we can choose one most suitable for the gradient computation. For the rest of this section we assume that  $d$  is even for simplicity of notation; a similar decomposition exists for odd  $d$  by choosing “half-angles” (see [LP04] for more details). We then choose the inner product defined by evaluation at  $2d + 1$  points on the circle,

$$\langle p, q \rangle = \frac{1}{2d+1} \sum_{k=1}^{2d+1} p(x_k)q(x_k), \quad x_k = \frac{2k\pi}{2d+1}.$$

Since a trigonometric polynomial of degree- $d$  is uniquely defined by evaluation on  $2d + 1$  unique points,  $\|p(x)\|^2 = 0$  if and only if  $p$  is identically zero.

Let  $U \in \mathbb{R}^{(d+1) \times r}$  be a matrix with column  $U_i$  representing the coefficients of  $u_i(x)$ , and  $B \in \mathbb{R}^{(d+1) \times (2d+1)}$  be the evaluation map on  $2d + 1$  points with columns

$$B_k = [1 \quad \cos(x_k) \quad \cdots \quad \cos(\frac{d}{2}x_k) \quad \sin(x_k) \quad \cdots \quad \sin(\frac{d}{2}x_k)]^\top,$$

so that  $B_k^\top U_i = u_i(x_k)$ . Let  $\bar{p}$  be the vector of coefficients of  $p(x)$ , so that  $B_k^\top \bar{p} = p(x_k)$ . Then we can write

$$f_p(U) = \frac{1}{2d+1} \sum_{k=1}^{2d+1} \left( \|U^\top B_k\|^2 - p(x_k) \right)^2$$

$$\nabla f_p(U) = \frac{4}{2d+1} U^\top B \text{Diag} \left( \|U^\top B_k\|^2 - p(x_k) \right) B^\top,$$

where  $\text{Diag} \left( \|U^\top B_k\|^2 - p(x_k) \right)$  is a diagonal matrix with  $\|U^\top B_k\|^2 - p(x_k)$  as the  $k$ -th diagonal entry. Since matrix-vector multiplication by  $B$  is equivalent to a Discrete Fourier Transform,  $\nabla f_p(U)$  can be computed in  $O(rd \log d)$  time using the FFT. Theorem 1.1 shows that spurious local minima do not exist when  $r \geq 2$ , so we can pick  $r$  to be a constant and obtain a near-linear iteration complexity.

We implemented our for finding the sum of squares decomposition of trigonometric polynomials in Julia, using the `FFTW.jl` package for FFTs to compute  $\nabla f_p(U)$  and the `NLopt.jl` package to minimize  $f_p(U)$  using a first-order algorithm (L-BFGS). The test polynomials are generated with

coefficients drawn from a standard normal distribution, with a constant coefficient added so that they all have a small positive minimum value.  $U$  is initialized with a small random value depending on the size of the problem. The algorithm is terminated when the relative error for each entry of  $U$  is on the order of  $10^{-7}$ . Although  $r = 2$  is sufficient, we observe that setting  $r = 4$  in our experiments improves the convergence rate with a minimal slowdown in iteration cost as the matrix-vector products can be easily parallelized across multiple threads. Figure 2 plots the convergence rate of 20 instances, and we can see that they achieve a linear convergence rate.




Figure 2: Value of  $f_p(\mathbf{u})$  against iterations of L-BFGS for computing the sum of squares decomposition of 20 random nonnegative trigonometric polynomials ( $2d = 100,000$ ), showing a linear convergence rate.

We performed the timing experiments on Intel Xeon Platinum 8260 processors, allocating 5 cores to each run, using polynomials of degree- $2d$  ranging from 2000 to 1,000,000. The results are summarized in Table 1.

Degree	Time in seconds	Iterations
2,000	2 (1 – 2)	340 (306 – 384)
10,000	6 (5 – 6)	530 (497 – 592)
20,000	9 (8 – 10)	632 (587 – 695)
100,000	53 (46 – 59)	1126 (980 – 1248)
200,000	160 (139 – 174)	1375 (1212 – 1532)
1,000,000	1461 (1212 – 1532)	2303 (1934 – 2437)

Table 1: Timing results for sum of squares decomposition of 50 random nonnegative trigonometric polynomials of varying degrees. Median time and iterations to termination due to stopping criterion (with range based on 25-th and 75-th percentile) are tabulated for different values of polynomial degree- $2d$ .

## 8 Conclusion

When does it make sense to solve non-convex formulations of convex problems? In this paper we addressed this question for sum of squares decomposition and optimization of univariate polynomials, showing that solving the nonconvex formulation can provide a large computational speedup while still maintaining provable guarantees on the convergence to the global optima. Key to our approach is retaining polynomial structure in the non-convex formulation. This enables us to use algebraic methods to construct a certificate showing that all SOCPs are global minima.

Our approach for finding sum of squares decompositions generalizes to multivariate polynomials, although the guarantees do not. However, we conjecture that a version of Theorem 1.1 is true for ternary quartics and matrix polynomials, cases where non-negativity is equivalent to the existence of a sum of squares decomposition. Another direction for future work is to apply our methods to other structured semidefinite programs or polynomial-valued objectives such as symmetric tensor decomposition.

## References

- [Bar01] A. Barvinok, *A remark on the rank of positive semidefinite matrices subject to affine constraints*, Discrete & Computational Geometry **25** (2001), no. 1, 23–31 (en).
- [BBJN18] Srinadh Bhojanapalli, Nicolas Boumal, Prateek Jain, and Praneeth Netrapalli, *Smoothed analysis for low-rank solutions to semidefinite programs in quadratic penalty form*, Proceedings of the 31st Conference On Learning Theory (Sébastien Bubeck, Vianney Perchet, and Philippe Rigollet, eds.), Proceedings of Machine Learning Research, vol. 75, PMLR, 06–09 Jul 2018, pp. 3243–3270.
- [BBV16] Afonso S Bandeira, Nicolas Boumal, and Vladislav Voroninski, *On the low-rank approach for semidefinite programs arising in synchronization and community detection*, Conference on learning theory, PMLR, 2016, pp. 361–382.
- [BM03] Samuel Burer and Renato D.C. Monteiro, *A nonlinear programming algorithm for solving semidefinite programs via low-rank factorization*, Mathematical Programming **95** (2003), no. 2, 329–357 (en).
- [BM05] ———, *Local minima and convergence in low-rank semidefinite programming*, Mathematical Programming **103** (2005), no. 3, 427–444 (en).
- [BNS16] Srinadh Bhojanapalli, Behnam Neyshabur, and Nathan Srebro, *Global optimality of local search for low rank matrix recovery*, Proceedings of the 30th International Conference on Neural Information Processing Systems (Red Hook, NY, USA), NIPS’16, Curran Associates Inc., 2016, p. 3880–3888.
- [BPT13] Grigoriy Blekherman, Pablo A. Parrilo, and Rekha R. Thomas, *Semidefinite optimization and convex algebraic geometry*, vol. 13, SIAM, 2013.
- [BVB16] Nicolas Boumal, Vlad Voroninski, and Afonso Bandeira, *The non-convex Burer-Monteiro approach works on smooth semidefinite programs*, Advances in Neural Information Processing Systems (D. Lee, M. Sugiyama, U. Luxburg, I. Guyon, and R. Garnett, eds.), vol. 29, Curran Associates, Inc., 2016.

- [BVB20] Nicolas Boumal, Vladislav Voroninski, and Afonso S Bandeira, *Deterministic guarantees for Burer-Monteiro factorizations of smooth semidefinite programs*, Communications on Pure and Applied Mathematics **73** (2020), no. 3, 581–608.
- [Cif21] Diego Cifuentes, *On the Burer-Monteiro method for general semidefinite programs*, Optimization Letters (2021), 1–11.
- [CLC19] Y. Chi, Y. M. Lu, and Y. Chen, *Nonconvex optimization meets low-rank matrix factorization: An Overview*, IEEE Transactions on Signal Processing **67** (2019), no. 20, 5239–5269.
- [CLR95] Man-Duen Choi, Tsit Yuen Lam, and Bruce Reznick, *Sums of squares of real polynomials*, Proceedings of Symposia in Pure mathematics, vol. 58, American Mathematical Society, 1995, pp. 103–126.
- [CM19] Diego Cifuentes and Ankur Moitra, *Polynomial time guarantees for the Burer-Monteiro method*, arXiv:1912.01745 [math, stat] (2019) (en).
- [CP17] Diego Cifuentes and Pablo A. Parrilo, *Sampling algebraic varieties for sum of squares programs*, SIAM Journal on Optimization **27** (2017), no. 4, 2381–2404 (en).
- [DF03] D.S. Dummit and R.M. Foote, *Abstract algebra*, Wiley, 2003.
- [Dum07] Bogdan Dumitrescu, *Positive trigonometric polynomials and signal processing applications*, vol. 103, Springer, 2007.
- [GHJY15] Rong Ge, Furong Huang, Chi Jin, and Yang Yuan, *Escaping from saddle points—online stochastic gradient for tensor decomposition*, Conference on learning theory, PMLR, 2015, pp. 797–842.
- [GJZ17] Rong Ge, Chi Jin, and Yi Zheng, *No spurious local minima in nonconvex low rank problems: A unified geometric analysis*, Proceedings of the 34th International Conference on Machine Learning - Volume 70, ICML’17, JMLR.org, 2017, p. 1233–1242.
- [GLM16] Rong Ge, Jason D. Lee, and Tengyu Ma, *Matrix completion has no spurious local minimum*, Proceedings of the 30th International Conference on Neural Information Processing Systems (Red Hook, NY, USA), NIPS’16, Curran Associates Inc., 2016, p. 2981–2989.
- [Ham73] Richard W. Hamming, *Numerical methods for scientists and engineers*, McGraw-Hill, Inc., USA, 1973.
- [JGN<sup>+</sup>17] Chi Jin, Rong Ge, Praneeth Netrapalli, Sham M Kakade, and Michael I Jordan, *How to escape saddle points efficiently*, International Conference on Machine Learning, PMLR, 2017, pp. 1724–1732.
- [Lau09] Monique Laurent, *Sums of squares, moment matrices and optimization over polynomials*, Emerging applications of algebraic geometry, Springer, 2009, pp. 157–270.

- [LP04] J. Lofberg and P.A. Parrilo, *From coefficients to samples: A new approach to SOS optimization*, 2004 43rd IEEE Conference on Decision and Control (CDC) (IEEE Cat. No.04CH37601) (Nassau), IEEE, 2004, pp. 3154–3159 (en).
- [LZT19] Qiuwei Li, Zhihui Zhu, and Gongguo Tang, *The non-convex geometry of low-rank matrix optimization*, Information and Inference: A Journal of the IMA **8** (2019), no. 1, 51–96.
- [Par00] Pablo A. Parrilo, *Structured semidefinite programs and semialgebraic geometry methods in robustness and optimization*, PhD thesis, California Institute of Technology, 2000.
- [Par06] Pablo A Parrilo, *Polynomial games and sum of squares optimization*, Proceedings of the 45th IEEE Conference on Decision and Control, IEEE, 2006, pp. 2855–2860.
- [Pat98] Gábor Pataki, *On the rank of extreme matrices in semidefinite programs and the multiplicity of optimal eigenvalues*, Mathematics of Operations Research (1998) (en).
- [RV06] Tae Roh and Lieven Vandenberghe, *Discrete transforms, semidefinite programming, and sum-of-squares representations of nonnegative polynomials*, SIAM Journal on Optimization **16** (2006), no. 4, 939–964.
- [Sch91] Konrad Schmüdgen, *The  $k$ -moment problem for compact semi-algebraic sets*, Mathematische Annalen **289** (1991), no. 1, 203–206.
- [Sch22] Claus Scheiderer, *Extreme points of gram spectrahedra of binary forms*, Discrete & Computational Geometry (2022), 1–17.
- [SQW18] Ju Sun, Qing Qu, and John Wright, *A geometric analysis of phase retrieval*, Foundations of Computational Mathematics **18** (2018), no. 5, 1131–1198.
- [Stu02] Bernd Sturmfels, *Solving systems of polynomial equations*, no. 97, American Mathematical Soc., 2002.
- [WBV99] Shao-Po Wu, Stephen Boyd, and Lieven Vandenberghe, *FIR filter design via spectral factorization and convex optimization*, Applied and computational control, signals, and circuits, Springer, 1999, pp. 215–245.
- [WW20] Irene Waldspurger and Alden Waters, *Rank optimality for the Burer-Monteiro factorization*, SIAM journal on Optimization **30** (2020), no. 3, 2577–2602.