SYK does not Transit Gloria Mundi just yet

D. V. Khveshchenko

Department of Physics and Astronomy, University of North Carolina, Chapel Hill, NC 27599

This note discusses examples of 0 + 1-dimensional Liouvillean dynamics instigated by the various deformations of the Sachdev-Ye-Kitaev (SYK) model. In reference to such deformations the main focus is on the regions of parameter space where the competing SYK couplings are of comparable strength and can not be treated as each others perturbations in the vicinity of the conformal fixed points corresponding to the pure SYK_q models with different values of q. Crossovers between such fixed points ('SYK transits') can be efficiently studied in the equivalent framework of single-particle quantum mechanics.

I. THE RISE OF SYK

The glorious rise of the celebrated SYK model^{1–8} into one of the central themes in modern interdisciplinary theoretical studies was due to a rare confluence of such precious properties as its elegant solubility, maximally chaotic behavior, asymptotic conformal symmetry, and more. The numerous in-depth analyses of the SYK model revealed a number of important connections between such seemingly disjoint subjects as random matrices, quantum black holes, disordered quantum dots, and, possibly, strange metallic behaviors in the various condensed matter systems.

One of these novel connections may have already contributed towards a resolution of the long-standing black hole information paradox by demonstrating that the properly (re)defined Hawking radiation entropy can be unitary, following the previously predicted Page curve^{9,10}.

In the condensed matter context, the SYK model has served as a powerful inspiration for a great many proposed non-Fermi-liquid (NFL) scenarios^{11–26}. However, the very existence of numerous plausible explanations of, e.g., the ubiquitous linear temperature dependence of resistivity in bad metals^{27–43} may seem to suggest that its ultimate explanation is yet to be found.

Nevertheless, alongside the renewed interest in hydrodynamics inspired by the holographic ideas, the SYK scenaria have been particularly important for pursuing the ad hoc field of 'bottom-up' (a.k.a.'non-AdS/non-CFT') holography which purports to describe a variety of (allegedly) strongly correlated condensed matter systems^{44–50}. Indeed, with the once abundant and defiantly upbeat claims of 'explaining' high- T_c materials, heavy fermions, graphene, etc. by virtue of some uncontrolled calculations in the conveniently chosen (and/or previously studied) classical gravity theories all but gone, the SYK model has remained a rather unique theoretical playground for obtaining rigorous results.

In that regard, the SYK model would be often referred to as a genuine example of low-dimensional holographic correspondence - even despite the fact that, both being effectively one-dimensional, the low-energy sector of SYK and its dual (formally, two-dimensional) Jackiw-Teitelboim (JT) gravity present a form of equivalence between different realizations of the quantized co-adjoint orbit of the (chiral) Virasoro group.

Such equivalence does not quite rise to the level of full-fledged holographic duality, as the JT bulk dual is non-dynamical and determined by the boundary degrees of freedom. By contrast, in order to qualify as a true holographic scenario the bulk theory would have to have some non-trivial bulk dynamics that gets quenched and turns classical only in a certain ('large-N') limit^{44–50}.

Moreover, similar remarks can also be made about the (historically, somewhat less extensively discussed) correspondence between the 3d gravity with BTZ-like black hole backgrounds and the various (KdV and alike) families of solvable 1 + 1-dimensional systems (see, e.g., Ref.⁵¹ and references therein).

II. THE SYK DEFORMATIONS

Since the beginning of the SYK era there have been attempts to explore deviations from the original SYK_4 model in order to assess the generality (or, conversely, uniqueness) of the behavior that it represents. In particular, there has been much discussion of the conjectured NFL-Fermi liquid (FL) transition in the hybrid $SYK_4 - SYK_2 \mod^{1-8,11-26}$.

A renormalization flow between the two fixed points has been mostly studied by means of perturbation theory operating in terms of the propagator $G(\tau_1, \tau_2)$ of $N \gg 1$ spaceless Majorana fermions^{11–26}. In the conformal limit of the generalized SYK_q of order $q \ge 4$ the latter exhibits the fermion dimension $\Delta = 1/q$, thus making the perturbation proportional to $G^{q/2}$ strongly relevant (dimension one) in the near vicinity of the UV SYK_q fixed point. Conversely, the formerly leading term G^q becomes strongly irrelevant (dimension four) near the IR fixed point of $SYK_{q/2}$. A unique feature of the q = 4 case, though, is that the transition occurs not between two different NFLs but the SYK_4 NFL and the disordered FL.

Notably, in the course of crossing over between the different fixed point regimes the value of q of the dominant term plays a role akin to that of the central charge in 2dconformal field theories. As far as the potential physical applications are concerned, some of the previous analyses^{11–26,52–55} suggest that the putative phase transition may take place at critical couplings vanishing as powers of 1/N - which value would be practically indistinguishable from zero in a macroscopic system - while others yield values that remain finite in the $N \to \infty$ limit.

The common approach to a SYK-type model starts out by integrating the fermions out, thereby arriving at the action in terms of the bi-local fields $G(\tau_1, \tau_2)$ and the corresponding self-energy $\Sigma(\tau_1, \tau_2)$

$$S = \frac{N}{2} Tr \ln(\partial_{\tau} - \Sigma) + \frac{N}{2} \int \int d\tau_1 d\tau_2 (\Sigma(\tau_1, \tau_2) G(\tau_1, \tau_2) - F[G(\tau_1, \tau_2)])$$
(1)

where the functional F[G] results from averaging over the Gaussian-correlated random amplitudes of all-to-all q-body entanglement. Moreover, such entangling couplings can be made non-uniform, thus introducing a notion of spatial dimensions and further extending the class of attainable models to include those with 'distance'dependent entanglement^{56,57}.

Solving for the self-energy, the Schwinger-Dyson equation derived from (1) takes the form

$$\partial_{\tau_1} G(\tau_1, \tau_2) - \int d\tau_3 < \tau_1 |\frac{\delta F}{\delta G}| \tau_3 > G(\tau_3, \tau_2) = \delta(\tau_1 - \tau_2)$$
(2)

In the original SYK_q model with $F(G) = J^2G^q$ the equation (2) remains invariant under any diffeomorphisms $\tau \to f(\tau)$ of the thermodynamic time variable τ subject to the boundary condition $f(\tau + \beta) = f(\tau) + \beta$ as long as the derivative term is neglected and provided that Gand Σ transform as

$$G(\tau_1, \tau_2) \to G_f = (f'(\tau_1)f'(\tau_2))^{\Delta}G(f(\tau_1), f(\tau_2)) \quad (3)$$

$$\Sigma(\tau_1, \tau_2) \to \Sigma_f = (f'(\tau_1)f'(\tau_2))^{1-\Delta}\Sigma(f(\tau_1), f(\tau_2))$$

At $J\tau \gg 1$ a representative power-law solution to the equation (2) reads $G_0(\tau_1, \tau_2) \sim sgn\tau/(J\tau)^{2\Delta}$ (hereafter $\tau = \tau_1 - \tau_2$).

Choosing a particular mean-field solution reduces the invariance under arbitrary diffeomorphisms down to the subgroup of the Mobius transformations SL(2, R). Correspondingly, a gradient expansion of the logarithm in (1) yields the (approximately) local effective action which describes finite temperature dynamics of the reparametrization mode^{58–79}

$$S_0(f) = -O(1)\frac{N}{Jq^2} \int Sch\{\tan\frac{\pi f}{\beta},\tau\}d\tau \qquad (4)$$

where Sch stands for the Schwarzian derivative $Sch\{f,x\} = \frac{f'''}{f'} - \frac{3}{2}(\frac{f''}{f'})^2$ obeying the differential 'chain rule' $Sch\{F(f),x\} = Sch\{F(f),f\}f'^2 + Sch\{f,x\}$ and operating on the manifold of (nearly) degenerate states related by virtue of the transformations (3).

III. LIOUVILLEAN QUANTUM MECHANICS

Under the customary parametrization $f'(\tau) = e^{\phi(\tau)}$ the Schwarzian action (4) assumes the (pseudo-)free form $S_0(\phi) \sim \int d\tau (\phi')^2$. In the process of averaging the products of propagators

$$< G_{f}(\tau_{1}, \tau_{2}) \dots G_{f}(\tau_{2p-1}, \tau_{2p}) > =$$

$$= \int D\phi e^{-S_{0}(\phi)} \prod_{i=1}^{p} \frac{e^{\Delta(\phi(\tau_{2i-1}) + \phi(\tau_{2i}))}}{(\int_{\tau_{2i-1}}^{\tau_{2i}} d\tau e^{\phi})^{2\Delta}}$$
(5)

over the fluctuations of ϕ the action $S_0(\phi)$ gets augmented by the Liouville term $\Delta S_2(\phi) = h_2 \int d\tau e^{2\phi(\tau)}$ with $h_2 \sim J$. Technically, upon promoting the denominator in (5) to the exponent with the help of some auxiliary integration a la Feynman the overall effective potential acquires a piece-wise Liouville term acting during the time intervals between 2p consecutive insertions of the operator $e^{i\Delta\phi 81,82}$.

The resulting action $S_0 + \Delta S_2$ can then be quantized by considering the corresponding (rescaled) Schroedinger equation^{81,82}

$$\left(-\frac{1}{2}\partial_{\phi}^{2}+h_{2}e^{2\phi}\right)\psi=E\psi\tag{6}$$

whose scattering states $\psi_k(z) \sim K_{2ik}(2\sqrt{z})$ (here $z = \lambda e^{\phi}$) belong to the continuum with the spectrum $E_k = k^2$ and density of states $\rho(E) \sim \sinh 2\pi \sqrt{2NE/J^{58-79,81,82}}$.

These exact expressions can be used to compute the matrix elements $< 0|e^{\Delta\phi}|k>$ exactly. Such calculations reveal the universal limit of an arbitrary power of G_f averaged over the soft mode fluctuations in the late-time, $\tau > N/J$ - or, at finite temperatures, in the strong coupling, $J\beta/N \gtrsim 1$ - regime^{81,82},

$$\langle G_f^p \rangle \propto \frac{1}{(J\tau)^{3/2}} \tag{7}$$

This behavior is markedly different from the (nonuniversal) mean-field one $G_0^p \propto 1/\tau^{2p/q}$ at shorter times $(\tau > N/J)$ or weak couplings $(J\beta/N \lesssim 1)$.

In that regard, the presence of the exponential term $\Delta S_2(\phi)$ in the overall effective action is instrumental. In its absence the Gaussian fluctuations of the field ϕ governed by $S_0(\phi)$ would have caused non-algebraic decay, thus being unable to deliver the universal power-law (7). In fact, such a behavior could have never emerged out of the purely Gaussian ϕ fluctuations even if the correlator $<\phi(\tau)\phi(0) >$ were logarithmic, as $\ln < G_f^p >$ would still depend on p and Δ (in both cases, quadratically).

The effective action might also include the various intrinsically non-local terms

$$\Delta S_n = h_n \int \int d\tau_1 d\tau_2 (\frac{f'(\tau_1) f'(\tau_2)}{(f(\tau_1) - f(\tau_2))^2})^{\Delta_n}$$
(8)

which can dominate over (4) for $\Delta_n \leq 3/2$. In the previous analyses, such terms would be routinely substituted

with the local operators $\Delta S_n = h_n \int d\tau e^{\Delta_n \phi(\tau)}$ thus further modifying the equivalent quantum mechanical Hamiltonian in (6).

IV. BI-QUADRATIC SYK DEFORMATION

This important example of the deformed SYK model has been extensively discussed in the context of random tunneling between two different SYK systems. For example, it arises in such, at first sight, unrelated fields as theoretical cosmology ('traversable wormhole')^{58–79} and coupled quantum dots^{83–93}.

In most analyses, the perturbed propagator would be taken in the form (3) of a 'gauge-transformed' meanfield solution G_0 , thereby accounting for the 'soft' reparametrization mode $f(\tau)$ while ignoring any potential changes to the mean-field background field configuration itself.

In particular, adding the SYK_2 ('tunneling') term with the amplitude Γ replaces the Liouville potential in the SYK_4 action (written in the Euclidean signature) with the Morse-type one⁹⁴⁻⁹⁷

$$S = \int d\tau (\frac{1}{2J} (\phi')^2 + J e^{2\phi} + \frac{\Gamma^2}{J} e^{\phi} + \frac{1}{\beta^2 J} e^{4\phi}) \qquad (9)$$

The corresponding Schroedinger equation with the (properly rescaled) Hamiltonian

$$H = -\frac{1}{2}\partial_{\phi}^2 + e^{2\phi} + \lambda e^{\phi} + \lambda' e^{4\phi}$$
(10)

which can be solved exactly in terms of the wave functions (here $z = 2\lambda e^{\phi}$) $\psi_k(z) \sim e^{-\phi/2} W_{\lambda,ik}(z)$ with the continuous spectrum $E_k = k^2 + 1/4 + \lambda^2$ if the last subdominant at low temperatures (or large negative ϕ) term in (10) can be dropped^{94–98}.

Besides, for $\lambda < 0$ the Hamiltonian (10) appears to possess a finite number of bound states

$$\psi_n(z) \sim z^{\lambda - n - 1/2 - z/2} L_n^{2\lambda - 2n - 1}(z)$$
 (11)

at the discrete energies $E_n = -(\lambda - n + 1/2)^2$, $n = 0, \ldots, [\lambda - 1/2]$. Near its minimum this spectrum can be approximated by the oscillator one.

A somewhat different path leading up to the Morsetype action (9) was taken in Ref.⁹⁸. The effect of the tunneling term with $\Delta_1 = 1/2$ was argued to be twofold: first, it contributes to (and/or refines) the purely Schwarzian (or 'hard' mode) saddle-point solution and, second, controls the pseudo-Goldstone (or 'soft' mode) fluctuations. These roles would be separately played by the 'longitudinal' (or radial, $e^{\xi} = 1 - f'$, in the holographically dual JT picture) and 'transverse' (or angular, ϕ) fluctuations, respectively. The former were argued to be strongly non-Gaussian and the effect of such fluctuations was claimed in Ref.⁹⁸ to strengthen (somewhat unexpectedly) the SYK_4 conformal mean-field behavior over a broader range of parameters.

More specifically, the strong coupling Schwarzian regime was argued to sustain the SYK_2 perturbation at all couplings $\gamma \equiv \Gamma/J$ below $\gamma_c \sim 1/N$ while at its higher values the propagator was found to crossover to the q = 2 FL fixed point. This was argued to be suggestive of a zero-temperature phase transition taking place at γ_c , rather than at a much larger value of order $1/N^{1/2}$, as per the naive estimate. Such parametric reduction of γ_c was claimed to manifest a stabilizing effect of the SYK_2 coupling on the mean-field conformal solution against the Schwarzian fluctuations due to the formation.

Correspondingly, the earlier perturbative analysis by the same authors revealed that a weak SYK_2 coupling does not alter the Schwarzian asymptotic (7) up to the values of γ of order γ_c^{55} .

Such observations appear to be generally consistent with those of Refs.^{99,100} which conjectured the existence of a chaotic-integrable transition in the $SYK_q - SYK_2$ model at finite temperatures. Above the transition temperature the system was found to behave chaotically while below it the chaos-related Lyapunov exponent (see below) dropped to zero, thus hinting at the FL nature of the underlying ground state.

On the technical side, upon first introducing two Lagrange multipliers λ and Λ and then voluntarily relaxing the corresponding constraints by fixing their mean-field values, Ref.⁹⁸ arrived at the effective action

$$S = \int d\tau (\frac{1}{2} (\phi')^2 + \Lambda (e^{2\phi} - f') + \lambda (e^{\phi} - \chi)) - \frac{1}{(\beta J)^2} e^{4\phi} - \frac{1}{2} \int \int d\tau_1 d\tau_2 \frac{\chi(\tau)\chi(\tau')}{|f(\tau_1) - f(\tau_2)|}$$
(12)

In this (perhaps, somewhat excessive) parametrization, functional integration about the mean-field SYK_4 fixed point factorizes into first taking a quantum mechanical expectation value over the exact ground state $\psi_0(\xi)$ of the Hamiltonian (10) and then additionally averaging over the Gaussian (perturbative) ϕ fluctuations. In Ref⁹⁸ neither mechanism was found to have any profound effect on the correlators, though.

In particular, an arbitrary power of the mean-field propagator would still retain its bare mean-field form provided that the ϕ -fluctuations were controlled by a large parameter λ . Likewise, averaging over the ground state of (10) adds the square of a non-singular expectation value $< 0|e^{p\Delta\phi}|0> = \int d\phi e^{p\Delta\phi}\psi_0^2(\phi)$ which does not give rise to any decaying power-law factor either. In that sense, the largely negligible effect of, both, the Gaussian fluctuations and the ground state averaging may indeed be viewed as increased stability of the mean-field regime in the presence of even a small SYK_2 coupling.

It should be noted, though, that under the assumption of $\lambda < 0$ the Morse potential in (10) appears to differ from that of Refs.^{52–54} which is strictly repulsive,

monotonic ($\lambda > 0$), and lacks any bound states. It might also be concerning that if the potential in (10) were to support any bound states with $E_n < 0$ then the fluctuation-averaged two-point correlator $\langle G_f(\tau) \rangle =$ $\sum_n e^{-E_n\tau} N(E_n)$ would be receiving - on top of the universal term (7) that stems from the continuum of scattering states with $E_k > 0$ - a non-unitary (exponential) contribution whose potential divergence could only be arrested by the squared matrix element $N(E_n < 0) = | < 0|e^{\Delta\phi}|n > |^2$.

Interestingly, for $J = \Gamma$ the aforementioned monotonic and non-monotonic Morse potentials represent two super-partners fitting into one super-symmetric pair $W_{\pm}(\phi) = V^2 \pm dV/d\phi$ with $V(\phi) \propto e^{\phi}$. The ground state of the binding potential then takes the form $\psi_0(\phi) \propto \exp(-\int V d\phi)$.

Conceivably, the effective action $S(\phi)$ may develop other interesting regimes at the points of still higher symmetry. One such example would be provided by the Hulten potential

$$W(\phi) = \lambda \frac{e^{\phi}}{1 - e^{\phi}} \tag{13}$$

whose first three terms of the expansion in powers of e^{ϕ} coincide with the 'hyper-symmetric' (or 'tri-critical') point $J = \Gamma = 1/\beta$ in (9). On the other hand, the $1/\phi$ -behavior at small negative ϕ would be similar to that in the Coulomb potential, although the potential (13) features only a finite number ([λ]) of bound states at $E_n = -(\lambda^2 - n^2/2\lambda n)^2$.

V. LARGE q LIMIT

An alternate approach to the SYK models exploits the large-q approximation where the propagator is sought out in the form

$$G(\tau) = \frac{1}{2} sgn\tau (1 + \frac{2}{q}g(\tau) + \dots)$$
 (14)

Higher order corrections in 1/q can also be evaluated, albeit at the increasingly prohibitive $costs^{58-79}$.

In action in the path integral over the field g then takes the form

$$S(g) = \frac{N}{q^2} \int \int d\tau_1 d\tau_2 \left(\frac{1}{2} \frac{dg}{d\tau_1} \frac{dg}{d\tau_2} + W(g)\right)$$
(15)

with the corresponding equation of motion

$$\partial_{\tau}^2 g = -\frac{\partial W(g)}{\partial g} \tag{16}$$

Formally solving (16) one obtains the classical trajectory

$$\tau = \int_{g_0}^0 \frac{dg}{\sqrt{W(g_0) - W(g)}}$$
(17)

with the use of which thermodynamics of the system can be studied by putting $\tau = \beta/2^{52-54}$. In particular, the turning point $g_0 < 0$ of the potential can be directly related to the mean-field energy⁵⁴

$$E = \frac{N}{4q^2} (\beta W(g_0) - 2^{3/2} \int_{g_0}^0 dg \sqrt{W_0 - W(g)}) \qquad (18)$$

As already mentioned, one possible generalization of the bi-quadratic (Schwarzian plus tunneling) q = 4 action to the larger values of q is provided by the $SYK_q - SYK_{q/2}$ functional

$$F[G] = \frac{2^q J^2}{q^2} G^q(\tau_1, \tau_2) + \frac{2^{q/2} \Gamma^2}{q^2} G^{q/2}(\tau_1, \tau_2) \quad (19)$$

The corresponding effective potential

$$W(g) = J^2 e^{2g} + \Gamma^2 e^g$$
 (20)

allows for the explicit saddle point solution 52-54

$$g(\tau) = -\ln(1 + \sqrt{J^2 + 4\Gamma^2}\tau + \Gamma^2\tau^2)$$
(21)

that gives rise to the mean-field propagator

$$G_0(\tau) = \frac{1}{2} \frac{sgn\tau}{(1 + \sqrt{J^2 + 4\Gamma^2}\tau + \Gamma^2\tau^2)^{2/q}}$$
(22)

For future reference, the final-temperature counterpart of (21) reads $g(\tau) = -\ln[(\sqrt{v^2 J^2/\beta^2} + \Gamma^4 \cos(2v\tau/\beta - v) + \Gamma^2)(\beta^2/2v^2)]$ where the parameter v is to be determined from the relation $2v^2 = \Gamma^2\beta^2 + \cos v\sqrt{J^2v^2\beta^2} + \Gamma^4\beta^4$ and becomes $v = 1 - O(1/\beta J)$ for $\Gamma = 0^{52-54}$.

It is worth noting that, in the look-alike equations (10) and (20) the field variables ϕ and g depend on the 'center-of-mass' (cf. Eq.(8)) and relative times, respectively. Also, unlike the approximate conformal propagator G_0 , the expression (22) is UV-finite and naturally regularized at $\tau \sim min[1/J, 1/\Gamma]$. Hence, by contrast with the latter, the saddle-point solution (22) remains applicable at all γ , both large and small. Therefore, the fluctuations of $g(\tau)$ describe pseudo-Goldstone excitations about the fixed 'valley' in the space of field configurations which no longer needs to be adjusted.

VI. QUADRATIC FLUCTUATIONS

Small fluctuations about the mean-field solution (22) are described by the Gaussian action

$$S_2 = \frac{N}{2q^2} \int \int d\tau_1 d\tau_2 \delta g(\tau_1) \frac{\partial^2 S}{\partial g^2} |_{g_0} \delta g(\tau_2)$$
(23)

For a potential $W(g) = \sum_{n} c_n e^{ng}$ these fluctuations δg would then be governed by a functionally similar kernel $\partial^2 W/\partial g^2 = \sum_{n} c_n n(n-1)e^{ng}$. Albeit similar in its appearance to the previously discussed $S(\phi)$, this action is bi-local and can not be readily used for deriving the Hamiltonian and quantizing it by means of the substitution $g' \rightarrow -i\partial/\partial g$.

In contrast to the Schwarzian action (4) the δg fluctuations are scale-invariant and their strength is independent of energy or temperature, being instead controlled by the numerical parameter N/q^2 . For a finite q the strength of such fluctuations decreases with increasing N, yet it remains fixed in the double-scaling limit, $N \to \infty$ and $N/q^2 = const$.

Inverting the Hessian operator evaluated at the saddle point (21) requires one to find the Green function of the retarded kernel $D(T,\tau) = \langle \delta g(T+\frac{\tau}{2}) \delta g(T-\frac{\tau}{2}) \rangle = \langle 12|K^{-1}|12 \rangle$ which satisfies the equation

$$\int \int d\tau_5 d\tau_6 (-\partial_1 \partial_2 \delta_{15} \delta_{26} + < 12 |\frac{\partial^2 W}{\partial g^2}|_{56} >)$$

$$< 56 |K^{-1}|_{34} >= (\delta_{13} \delta_{24} - \delta_{14} \delta_{23}) \qquad (24)$$

Upon Fourier transforming with respect to the 'centerof-mass' time variable T one can use the spectral decomposition

$$D(T,\tau) = \sum_{n} \int \frac{d\omega}{2\pi} e^{-i\omega T} \frac{\psi_n(\frac{\tau}{2})\psi_n^*(-\frac{\tau}{2})}{\omega^2 - \omega_n^2 + i0}$$
$$= \frac{i}{2} \sum_n \frac{e^{i\omega_n T}}{\omega_n} \psi_n(\frac{\tau}{2})\psi_n^*(-\frac{\tau}{2}) \qquad (25)$$

in terms of the eigenfunctions of the equation

$$(-\partial_{\tau}^2 + \frac{\partial^2 W}{\partial g^2}|_{g_0})\psi_n(\tau) = \omega_n^2\psi_n(\tau)$$
(26)

By analogy with the aforementioned averaging over the ϕ -fluctuations the Gaussian average over δg in the vicinity of the saddle point (22) produces the 'Debye-Waller' factor

$$\frac{\langle G^{p}(\tau) \rangle}{G_{0}^{p}(\tau)} = \langle e^{2p\Delta\delta g(\tau)} \rangle = \exp(2p^{2}\Delta^{2}(D(0,\tau) - D(0,0)))$$
(27)

Notably, this averaging is to be performed over the entire function δg , thereby making no distinction between the 'angular' and 'radial' modes.

This might be somewhat similar to, e.g., the standard weak-coupling analysis of the two-dimensional nonlinear $O(N) \sigma$ -model which seems to emphasize a distinction between the longitudinal and transverse fluctuations of the order parameter (one gapped and N-1 Goldstone modes, respectively). By contrast, the exact solution demonstrates no such difference as the true O(N)symmetric spectrum consists of the N identical gapped modes.

Evaluating (20) on the classical trajectory (21) at zero temperature one finds the effective potential that asymptotically decays at large τ as $\sim 1/\tau^2$ in both cases of large and small γ . The one-dimensional Green function of the resulting eigenvalue equation

$$(-\partial_{\tau}^2 + \frac{\kappa}{\tau^2} - \omega^2)\psi = 0 \tag{28}$$

with $\kappa > -1/4$ can be found in the closed form

$$D_{\omega}(\tau, \tau') = \frac{\pi}{2i} \sqrt{\tau \tau'} \frac{H_{\nu}^{(1)}(\omega \tau_{>})}{H_{\nu}^{(1)}(\omega a)}$$
$$(H_{\nu}^{(1)}(\omega \tau) J_{i\nu}(\omega \tau_{<}) - H_{\nu}^{(1)}(\omega \tau_{<}) J_{\nu}(\omega a))$$
(29)

where $\tau_{>}$ and $\tau_{<}$ stand for the larger/smaller of τ and τ' , respectively, $\nu = \sqrt{1/4 + \kappa}$, and *a* is the UV cutoff.

For $\omega = 0$ (29) amounts to the previously derived expression whose finite-temperature version reads^{58–79}

$$D_0(x, x') = \frac{1}{V\pi} (1 + \tan x_< (\frac{V\pi}{2} + x_<)) (1 - \tan x_> (\frac{V\pi}{2} - x_>))$$
(30)

where $x = \pi \tau / \beta$ and $V = v + \frac{2}{\pi} \cot \pi v / 2$.

Expanding the Bessel functions one once again finds only a mild effect of the Gaussian fluctuations (this time around, of δg), as the ensuing reduction of the amplitude $\langle G^p(\tau) \rangle / G_0^p(\tau) = \exp(O(1)\Delta^2 p^2)$, does not alter the mean-field exponent of the power-law decay.

VII. QUADRATIC FLUCTUATIONS IN g-SPACE

As an alternative to (26) one can formulate the eigenvalue equation in terms of the g-variable⁵⁴

$$(-2\sqrt{W_0 - W}\partial_g\sqrt{W_0 - W}\partial_g + \frac{\partial^2 W}{\partial g^2}|_{g_0})\psi_n = \omega_n^2\psi_n$$
(31)

where $W_0 = W(g_0)$ without the need to explicitly solve for the classical trajectory $g(\tau)$.

However, a generally non-trivial derivative $\partial_{\tau}g$ precludes an immediate use of the known solutions such as (11) in the case of, e.g., the Morse potential W(g). Then treating (31) as a generic second-order equation

$$p(x)\partial_x^2\psi + q(x)\partial_x\psi + (E - V)\psi = 0$$
(32)

and eliminating the linear derivative term one can convert the equation (31) into the standard Schroedinger equation with the potential $V' = V + \frac{\partial_x^2 Q}{Q}$ in terms of the wavefunction $\chi = \psi Q$ with $Q(x) = \exp(\int dx q/2p)$.

Using this equation in the classically accessible domain $g_0 < g < 0$ one can study the system's thermodynamics. For example, in the case of the Hulten potential (13) one obtains non-trivial temperature dependences of energy $E = E_0 - O(J^{4/3}\beta^{1/3})$ and entropy $S = S_0 - O((J\beta)^{4/3})$ which suggest rather peculiar thermodynamic relations.

VIII. LADDER EIGENFUNCTIONS AND CHAOS EXPONENTS

A chaotic behavior may develop in the complementary (classically prohibited) regime $g < g_0$. One popular quantifier of chaos is provided by the out-of-timeorder correlator (OTOC) given by the averaged amplitude $\langle G_f(\tau_1, \tau_3)G_f(\tau_2, \tau_4) \rangle$ analytically continued from the domain $\tau_4 < \tau_2 < \tau_3 < \tau_1$ to the complex times $\tau_1 = \beta/4 - it/2, \tau_2 = -\beta/4 - it/2, \tau_3 = it/2,$ $\tau_4 = -\beta/2 + it/2.$

On top of a non-exponential regular part of zeroth order in 1/N the OTOC function demonstrates an exponentially growing first order correction. In the case of the $SYK_q - SYK_{q/2}$ model it reads

$$OTOC(t) = O(\frac{1}{\beta J}) - \frac{f(\gamma)}{N} e^{\lambda_L t}$$
(33)

where f(0) = O(1). It is controlled by the Lyapunov chaos exponent determined by the ladder eigenstate equation constructed out of the Wightman correlators $G_{lr}(t) = G(\tau = it + \beta/2)^{58-79}$. Taking ω_n in (26) to imaginary values $\omega_n \to i\lambda_L$ yields the exponentially growing ansatz $D(T, t) \sim e^{\lambda_L T} \psi(t)$ where the real-time eigenfunction $\psi(t)$ solves the equation⁵⁴

$$(-\partial_x^2 - \frac{\cos\theta}{\cosh x + \cos\theta} - \frac{2\sin^2\theta}{(\cosh x + \cos\theta)^2})\psi = -(\frac{\lambda_L\beta}{2\pi\nu})^2\psi$$
(34)

with $\theta = \tan^{-1}(v/J\beta\gamma^2)$. Importantly, the potential in (34) is monotonic and its sign is opposite of that in (26).

For $\gamma = 0$ this potential is the original SYK's one, $V_0(x) = -2/\cosh^2 x$, which supports no bound states other than the ground one, $\psi_0(x) \sim 1/\cosh x$, with the eigenvalue $E_0 = -(\lambda_L \beta/2\pi v)^2 = -1^{58-79}$. As has been repeatedly pointed out in the literature, this value of the chaotic operator growth is (almost) maximally possible, its reduction at strong coupling $(J\beta \gg 1)$ being solely due to the temperature-dependent factor v

$$\lambda_L = \frac{2\pi}{\beta} \left(1 - O(\frac{1}{\beta J})\right) \tag{35}$$

In the complementary weak coupling regime $(J\beta \ll 1)$ the chaotic exponent is $\lambda_L \sim J$.

In principle, the rest of the spectrum in (34) could provide for some slower growing terms. However, for $\gamma = 0$ no such terms appear as the next (single-node, hence first excited) state would be given by the function $\psi_1 \sim g'$ with the eigenvalue $E_1 = 0^{54}$.

Also, at longer times $t \gg \beta$ the behavior of the OTOC function is determined by the 2-particle density of states, resulting in another universal power-law, $OTOC(t) \propto 1/t^{681,82}$.

In Ref.⁹⁸ a weak SYK_2 term in (20) was found not to drastically alter the strong-coupling behavior, except for a reduction of the amplitude by a factor $O(1/J\beta\gamma^2) < 1$ in the entire interval $1/N \leq \gamma \leq 1/N^{1/2}$. At such parameter values the Schwarzian fluctuations were found to be suppressed, thus extending the validity of the SYK_4 mean-field solution beyond the energy scale J/N all the way down to $J\gamma^2$ at which the Fl behavior finally sets in. In Ref.⁵⁴, the chaotic exponent of the large-q biquadratic model was computed with the use of perturbation theory about the state ψ_0 for small γ , thereby finding

$$\lambda_L = \frac{2\pi}{\beta} (1 - O(\min[J\beta\gamma^2, \frac{1}{J\beta\gamma^2}]))$$
(36)

For comparison, Refs.^{99,100} found the exponent $\lambda_L = \frac{2\pi}{\beta}(1 - O(\beta^2 \Gamma^2))$ in the $SYK_q - SYK_2$ model, suggesting the possibility of a finite-temperature transition for arbitrarily small Γ .

The latter should, however, be contrasted against the result of Ref.¹⁰¹ which reported $\lambda_L \sim 1/J\beta^2\gamma^3$ for $\gamma \gg max[1, 1/J\beta]$. Such a behavior conforms to the generic quadratic temperature dependence of λ_L in disordered FL and could indicate the absence of a genuine finite-temperature phase transition for a sufficiently large Γ .

Adding to the list of possibilities, in Ref.⁵⁴ some nonmaximal (temperature-independent and growing with the increasing integer parameter n) values of λ_L were reported on the basis of a numerical solution of some other ('variable scaling') model with $W(g) \propto 1/(-g)^n$.

As far as more general potentials W(g) are concerned, the Hulten potential (13), for one, falls somewhere in between the 'super-symmetric' ($\gamma = 1$) point of the $SYK_q - SYK_{q/2}$ model and the 'variable scaling' one. The corresponding eigenvalue equation now reads

$$\left(-\partial_x^2 - \frac{2}{\delta}\left(\frac{1}{\cosh x} - \frac{1}{\cosh x + \delta}\right)\right)\psi = -\left(\frac{\lambda_L\beta}{2\pi v}\right)^2\psi \quad (37)$$

where $\delta = \sqrt{1 + 4\gamma^2}/J\beta\gamma^2$. At the super-symmetric point where $\delta = 5^{1/2}/J\beta$ and for low temperatures $(\delta \ll 1)$ the potential in the equation (37) approaches the original SYK's $V_0(x)$ and the maximally chaotic behavior $(\lambda_L \rightarrow \frac{2\pi}{\beta v})$ is once again restored. In the opposite limit of $\delta \gg 1$ the potential flattens out and the Lyapunov exponent decreases monotonically all the way to zero. It does not vanish at any finite temperature, though, thus calling for a closer look at any scenario of a finite-temperature phase transition - or a zero-temperature one predicted to occur at a critical γ_c vanishing as a power of 1/N.

IX. SUMMARY

This note discussed various generalizations of the SYK model that lead to the one-dimensional Liouvillean quantum mechanics. Of a particular interest are crossovers between the different conformal fixed points where all pertinent coupling constants are likely to be of the same order. Such 'SYK transits' are not directly amenable to perturbation theory in the vicinity of the fixed points in question but can still be explored in the large-q limit. To that end, one can utilize the already available -and seek out new - non-perturbative mean-field solutions akin to (22) which interpolate between the distinct conformal regimes. This way one could advance the previous studies of the bi-quadratic model (20) and its further extensions within a broader class of the effective potentials W(g).

In particular, this preliminary analysis finds that the Lyapunov exponent at the 'super-symmetric' point of the model (20) remains non-zero down to the lowest temper-

- ¹ S. Sachdev and Jinwu Ye, Gapless spin-fluid ground state in a random quantum Heisenberg magnet, Phys. Rev. Lett. **70**, 3339 (1993), arXiv:cond-mat/9212030
- ² S. Sachdev, Holographic metals and the fractionalized Fermi liquid, Phys. Rev. Lett. **105**, 151602 (2010)
- ³ S. Sachdev, Bekenstein-Hawking entropy and strange metals, Phys. Rev. X 5, 041025 (2015)
- ⁴ A. Kitaev, *KITP Seminars, 2015*, http://online.kitp.ucsb.edu/
- 5 A. Kitaev, Notes on $\tilde{SL}(2,R)$ representations, arXiv:1711.08169
- ⁶ A. Kitaev and S.J. Suh, The soft mode in the Sachdev-Ye-Kitaev model and its gravity dual, JHEP **2018**, 183 (2018), arXiv:1711.08467
- ⁷ A. Kitaev and S.J. Suh, Statistical mechanics of a two-dimensional black hole, JHEP **2019**, 198 (2019), arXiv:1808.07032
- ⁸ S. Sachdev, Statistical mechanics of strange metals and black holes, arXiv:2205.02285
- ⁹ G. Penington, Entanglement wedge reconstruction and the information paradox, arXiv:1905.08255
- ¹⁰ A. Almheiri, N. Engelhardt, D. Marolf, and H. Maxfield, The entropy of bulk quantum fields and the entanglement wedge of an evaporating black hole. JHEP **2019**, 63 (2019), arXiv:1905.08762
- ¹¹ S. Banerjee and E. Altman, Solvable model for a dynamical quantum phase transition from fast to slow scrambling, Phys. Rev. B **95**, 134302 (2017)
- ¹² Zhen Bi, Chao-Ming Jian, Yi-Zhuang You, K.A. Pawlak, and Cenke Xu, Instability of the non-Fermi-liquid state of the Sachdev-Ye-Kitaev model, Phys. Rev. B **95**, 205105 (2017)
- ¹³ Shao-Kai Jian and Hong Yao, Solvable Sachdev-Ye-Kitaev models in higher dimensions: From diffusion to manybody localization, Phys. Rev. Lett. **119**, 206602 (2017)
- ¹⁴ A. Haldar, S. Banerjee, and V.B. Shenoy, Higherdimensional Sachdev-Ye-Kitaev non-Fermi liquids at Lifshitz transitions, Phys. Rev. B 97, 241106 (2018)
- ¹⁵ Chao-Ming Jian, Zhen Bi, and Cenke Xu, Model for continuous thermal metal to insulator transition, Phys. Rev. B 96, 115122 (2017)
- ¹⁶ Xue-Yang Song, Chao-Ming Jian, and L. Balents, Strongly correlated metal built from Sachdev-Ye-Kitaev models, Phys. Rev. Lett. **119**, 216601 (2017)
- ¹⁷ Xin Chen, Ruihua Fan, Yiming Chen, Hui Zhai, and Pengfei Zhang, Competition between chaotic and nonchaotic phases in a quadratically coupled Sachdev-Ye-Kitaev model, Phys. Rev. Lett. **119**, 207603 (2017)
- ¹⁸ Pengfei Zhang, Dispersive Sachdev-Ye-Kitaev model: Band structure and quantum chaos, Phys. Rev. B 96, 205138 (2017)

atures. This observation may call for inspection of the earlier conclusions about the onset of the non-chaotic FL phase at a critical coupling γ_c which could be as weak as $O(1/N^{1/2})$ or even $O(1/N)^{11-26,98-100}$.

Also, further generalizations of the standard Liovillean action related to the various analytically solvable quantum mechanical Hamiltonians might be of interest well above and beyond the original SYK context.

- ¹⁹ Wenhe Cai, Xian-Hui Ge, and Guo-Hong Yang, Diffusion in higher dimensional SYK model with complex fermions, JHEP **2018**(01), 76 (2018)
- ²⁰ Zhong Yin, Periodic Anderson model meets Sachdev-Ye-Kitaev interaction: a solvable playground for heavy fermion physics, J. Phys. Commun. 2, 095014 (2018)
- ²¹ Xin Dai, Shao-Kai Jian, Hong Yao, Global phase diagram of the one-dimensional Sachdev-Ye-Kitaev model at finite N, Phys. Rev. B 100, 235144 (2019), arXiv:1802.10029
- ²² Pengfei Zhang and Hui Zhai, Topological Sachdev-Ye-Kitaev model, Phys. Rev. B **97**, 201112(R) (2018)
- ²³ Xiaochuan Wu, Xiao Chen, Chao-Ming Jian, Yi-Zhuang You, and Cenke Xu, Candidate theory for the strange metal phase at a finite-energy window, Phys. Rev. B 98, 165117 (2018)
- ²⁴ D. Ben-Zion and J. McGreevy, Strange metal from local quantum chaos, Phys. Rev. B **97**, 155117 (2018)
- ²⁵ A.A. Patel, J. McGreevy, D.P. Arovas, and S. Sachdev, Magnetotransport in a model of a disordered strange metal, Phys. Rev. X 8, 021049 (2018)
- ²⁶ D. Chowdhury, Y. Werman, E. Berg, and T. Senthil, Translationally invariant non-Fermi-liquid metals with critical Fermi surfaces: Solvable models, Phys. Rev. X 8, 031024 (2018)
- ²⁷ A.A. Patel and S. Sachdev, Theory of a Planckian metal, Phys. Rev. Lett. **123**, 066601 (2019)
- ²⁸ A.A. Patel and S. Sachdev, Critical strange metal from fluctuating gauge fields in a solvable random model, Phys. Rev. B **98**, 125134 (2018)
- ²⁹ D. Miserev, J. Klinovaja, and D. Loss, Fermi surface resonance and quantum criticality in strongly interacting Fermi gases, Phys. Rev. B **103**, 075104 (2021)
- ³⁰ D. Chowdhury, A. Georges, O. Parcollet, and S. Sachdev, Sachdev-Ye-Kitaev models and beyond: A window into non-Fermi liquids, arXiv:2109.05037
- ³¹ I. Esterlis, Haoyu Guo, A.A. Patel, and S. Sachdev, Large-N theory of critical Fermi surfaces, Phys. Rev. B 103, 235129 (2021)
- ³² D. Chowdhury and E. Berg, Intrinsic superconducting instabilities of a solvable model for an incoherent metal, Phys. Rev. Research 2, 013301 (2020)
- ³³ P. Cha, A.A. Patel, E. Gull, and Eun-Ah Kim, Slope invariant *T*-linear resistivity from local self-energy, Phys. Rev. Research 2, 033434 (2020)
- ³⁴ Haoyu Guo, Yingfei Gu, and S. Sachdev, Transport and chaos in lattice Sachdev-Ye-Kitaev models, Phys. Rev. B 100, 045140 (2019)
- ³⁵ I. Esterlis and J. Schmalian, Cooper pairing of incoherent electrons: An electron-phonon version of the Sachdev-Ye-Kitaev model, Phys. Rev. B **100**, 115132 (2019)
- ³⁶ Yuxuan Wang and A. V. Chubukov, Quantum phase tran-

sition in the Yukawa-SYK model, Phys. Rev. Research 2, 033084 (2020)

- ³⁷ B. Douçot, A. Mukhopadhyay, G. Policastro, and S. Samanta, Linear-in-*T* resistivity from semiholographic non-Fermi liquid models, Phys. Rev. D **104**, L081901 (2021)
- ³⁸ L. Classen and A. Chubukov, Superconductivity of incoherent electrons in the Yukawa Sachdev-Ye-Kitaev model, Phys. Rev. B **104**, 125120 (2021)
- ³⁹ Peter Cha, A.A. Patel, and Eun-Ah Kim, Strange metals from melting correlated insulators in twisted bilayer graphene, Phys. Rev. Lett. **127**, 266601 (2021)
- ⁴⁰ F. Salvati and A. Tagliacozzo, Superconducting critical temperature in the extended diffusive Sachdev-Ye-Kitaev model, Phys. Rev. Research **3**, 033117 (2021)
- ⁴¹ Gaopei Pan, Wei Wang, A. Davis, Yuxuan Wang, and Zi Yang Meng, Yukawa-SYK model and self-tuned quantum criticality, Phys. Rev. Research 3, 013250 (2021)
- ⁴² G. Jose, Kangjun Seo, and B. Uchoa, Non-Fermi liquid behavior in the Sachdev-Ye-Kitaev model for a onedimensional incoherent semimetal, Phys. Rev. Research 4, 013145 (2022)
- ⁴³ A.A. Patel, Haoyu Guo, I. Esterlis, and S. Sachdev, Universal, low temperature, *T*-linear resistivity in twodimensional quantum-critical metals from spatially random interactions, arXiv:2203.04990
- ⁴⁴ S.A. Hartnoll, Lectures on holographic methods for condensed matter physics, Class. Quant. Grav. 26, 224002 (2009)
- ⁴⁵ C.P. Herzog, Lectures on holographic superfluidity and superconductivity, J. Phys. A **42**, 343001 (2009)
- ⁴⁶ J. McGreevy, Holographic duality with a view toward many-body physics, Adv. High Energy Phys. **2010**, 723105 (2010)
- ⁴⁷ S. Sachdev, What can gauge-gravity duality teach us about condensed matter physics?, Annu. Rev. Cond. Matt. Phys. 3, 9 (2012)
- ⁴⁸ J. Zaanen, Y. Liu, Y.-W. Sun, and K. Schalm, *Holographic Duality in Condensed Matter Physics* (Cambridge University Press, 2015)
- ⁴⁹ M. Ammon and J. Erdmenger, *Gauge / Gravity Duality* (Cambridge University Press, 2015)
- ⁵⁰ S.A. Hartnoll, A. Lucas, and S. Sachdev, *Holographic Quantum Matter* (MIT Press, 2018)
- ⁵¹ D.V. Khveshchenko, Phase space holography with no strings attached, Lith. J. Phys. **61**, 233 (2021), arXiv:2102.01617
- ⁵² Chao-Ming Jian, Zhen Bi, and Cenke Xu, Model for continuous thermal metal to insulator transition, Phys. Rev. B 96, 115122 (2017), arXiv:1703.07793
- ⁵³ D. Anninos and D.A. Galante, Constructing AdS₂ flow geometries, JHEP **2021**, 45 (2021), arXiv:2011.01944
- ⁵⁴ Jiaqi Jiang and Zhenbin Yang, Thermodynamics and many body chaos for generalized large q SYK models, JHEP **2019** 19 (2019), arXiv:1905.00811
- ⁵⁵ A.V. Lunkin, K.S. Tikhonov, and M.V. Feigel'man, Sachdev-Ye-Kitaev model with quadratic perturbations: The route to a non-Fermi liquid, Phys. Rev. Lett. **121**, 236601 (2018), arXiv:1806.11211
- ⁵⁶ D.V. Khveshchenko, Thickening and sickening the SYK model, SciPost Phys. 5, 012 (2018), arXiv:1705.03956
- ⁵⁷ D.V. Khveshchenko, Seeking to develop global SYK-ness, Condens. Matter **2018**, 3(4), 40 (2018), arXiv:1805.00870
- ⁵⁸ J. Maldacena, S.H. Shenker, and D. Stanford, A bound

on chaos, JHEP 2016(08), 106 (2016), arXiv:1503.01409

- ⁵⁹ J. Maldacena and D. Stanford, Remarks on the Sachdev-Ye-Kitaev model, Phys. Rev. D **94**, 106002 (2016), arXiv:1604.07818
- ⁶⁰ J. Maldacena, D. Stanford, and Zhebin Yang, Conformal symmetry and its breaking in two dimensional nearly Anti-de-Sitter space, PTEP **2016**(12), 12C104 (2016), arXiv:1606.01857
- ⁶¹ D. Stanford and E. Witten, Fermionic localization of the schwarzian theory, JHEP **2017**(10), 8 (2017), arXiv:1703.04612
- ⁶² J. Polchinski and V. Rosenhaus, The spectrum in the Sachdev-Ye-Kitaev model, JHEP **2016**(04), 1 (2016), arXiv:1601.06768
- ⁶³ D.J. Gross and V. Rosenhaus, The bulk dual of SYK: cubic couplings, JHEP **2017**(05), 92 (2017), arXiv:1702.08016
- ⁶⁴ D.J. Gross and V. Rosenhaus, All point correlation functions in SYK, JHEP **2017**(12), 148 (2017), arXiv:1710.08113
- ⁶⁵ G. Sárosi, AdS₂ holography and the SYK model, in: XIII Modave Summer School in Mathematical Physics (Modave2017), Proc. Sci. vol. 323, arXiv:1711.08482, arXiv:1711.08482
- ⁶⁶ Henry W. Lin, J. Maldacena, and Ying Zhao, Symmetries near the horizon, JHEP **2019**, 49 (2019), arXiv:1904.12820
- ⁶⁷ E. Witten, An SYK-Like model without disorder, arXiv:1610.09758
- ⁶⁸ R. Gurau, The complete 1/N expansion of a SYKlike tensor model, Nucl. Phys. B **916**, 386–401 (2017), arXiv:1611.04032
- 69 R. Gurau, The ϵ prescription in the SYK model, arXiv:1705.08581
- ⁷⁰ I.R. Klebanov and G. Tarnopolsky, Uncolored random tensors, melon diagrams, and the Sachdev-Ye-Kitaev models, Phys. Rev. D **95**, 046004 (2017)
- ⁷¹ S. Giombi, I.R. Klebanov, and G. Tarnopolsky, Bosonic tensor models at large N and small ε , Phys. Rev. D **96**, 106014 (2017)
- ⁷² Yingfei Gu, Xiao-Liang Qi, and D. Stanford, Local criticality, diffusion and chaos in generalized Sachdev-Ye-Kitaev models, JHEP **2017**, 125 (2017)
- ⁷³ Yingfei Gu, A. Lucas, and Xiao-Liang Qi, Energy diffusion and the butterfly effect in inhomogeneous Sachdev-Ye-Kitaev chains, SciPost Phys. 2, 018 (2017)
- ⁷⁴ Yingfei Gu, A. Lucas, and Xiao-Liang Qi, Spread of entanglement in a Sachdev-Ye-Kitaev chain, JHEP **2017**, 120 (2017)
- ⁷⁵ Yingfei Gu and A. Kitaev, On the relation between the magnitude and exponent of OTOCs, JHEP **2019**(02), 75 (2019)
- ⁷⁶ Zhenbin Yang, The quantum gravity dynamics of near extremal black holes, JHEP **2019**(05), 205 (2019), arXiv:1809.08647
- ⁷⁷ A. Blommaert, T.G. Mertens, and H. Verschelde, The Schwarzian theory — a Wilson line perspective, JHEP **2018**(12), 22 (2018), arXiv:1806.07765
- ⁷⁸ A. Blommaert, T.G. Mertens, and H. Verschelde, Fine structure of Jackiw-Teitelboim quantum gravity, JHEP **2019**(9), 66 (2019), arXiv:1812.00918
- ⁷⁹ G. Tarnopolsky, Large q expansion in the Sachdev-Ye-Kitaev model, Phys. Rev. D **99**, 026010 (2019), arXiv:1801.06871

- ⁸⁰ A. Milekhin, Coupled Sachdev-Ye-Kitaev models without Schwartzian dominance, arXiv:2102.06651
- ⁸¹ D. Bagrets, A. Altland, and A. Kamenev, Sachdev-Ye-Kitaev model as Liouville quantum mechanics, Nucl. Phys. B **911**, 191–205 (2016)
- ⁸² D. Bagrets, A. Altland, and A. Kamenev, Power-law out of time order correlation functions in the SYK model, Nucl. Phys. B **921**, 727 (2017), arXiv:1702.08902
- ⁸³ N.V. Gnezdilov, J.A. Hutasoit, and C.W.J. Beenakker, Low-high voltage duality in tunneling spectroscopy of the Sachdev-Ye-Kitaev model, Phys. Rev. B **98**, 081413
- ⁸⁴ O. Can, E.M. Nica, and M. Franz, Charge transport in graphene-based mesoscopic realizations of Sachdev-Ye-Kitaev models, Phys. Rev. B **99**, 045419 (2019)
- ⁸⁵ A. Altland, D. Bagrets, and A. Kamenev, Sachdev-Ye-Kitaev non-Fermi-liquid correlations in nanoscopic quantum transport, Phys. Rev. Lett. **123**, 226801 (2019)
- ⁸⁶ A. Kruchkov, A.A. Patel, P. Kim, and S. Sachdev, Thermoelectric power of Sachdev-Ye-Kitaev islands: Probing Bekenstein-Hawking entropy in quantum matter experiments, Phys. Rev. B **101**, 205148 (2020), arXiv:1912.02835
- ⁸⁷ D.I.Pikulin and M.Franz, Black hole on a chip: Proposal for a physical realization of the Sachdev-Ye-Kitaev model in a solid-state system, Phys. Rev. X 7, 031006 (2017)
- ⁸⁸ A. Chew, A. Essin, and J. Alicea, Approximating the Sachdev-Ye-Kitaev model with Majorana wires, Phys. Rev. B **96**, 121119 (2017)
- ⁸⁹ A.Chen et al, Quantum holography in a graphene flake with an irregular boundary, Phys. Rev. Lett. **121**, 036403 (2018)
- ⁹⁰ E. Lantagne-Hurtubise, Chengshu Li, and M. Franz, Fam-

ily of Sachdev-Ye-Kitaev models motivated by experimental considerations, Phys. Rev. B **97**, 235124 (2018)

- ⁹¹ M. Franz and M. Rozali, Mimicking black hole event horizons in atomic and solid-state systems, arXiv:1808.00541
- ⁹² D.V. Khveshchenko, Connecting the SYK dots, Condens. Matter 5(2), 37 (2020), arXiv:2004.06646
- ⁹³ D.V. Khveshchenko, One SYK single electron transistor, Lith. J. Phys. **60**, 185 (2020), arXiv:1912.05691
- ⁹⁴ J. Maldacena and Xiao-Liang Qi, Eternal traversable wormhole, arXiv:1804.00491
- ⁹⁵ T.G. Mertens, G.J. Turiaci, and H.L. Verlinde, Solving the Schwarzian via the conformal bootstrap, JHEP 08, 136 (2017), arXiv:1705.08408
- ⁹⁶ T.G. Mertens, The Schwarzian theory origins, JHEP 2018(05), 36 (2018), arXiv:1801.09605
- ⁹⁷ D.V. Khveshchenko, On a (pseudo)holographic nature of the SYK-like models, Lith. J. Phys. **59**, 104 (2019), arXiv:1905.04381
- ⁹⁸ A.V. Lunkin, A.Yu. Kitaev, and M.V. Feigel'man, Perturbed Sachdev-Ye-Kitaev Model: A polaron in the hyperbolic plane, Phys. Rev. Lett. **125**, 196602 (2020), arXiv:2006.14535
- ⁹⁹ A.M. García-García, B. Loureiro, A. Romero-Bermúdez, and M. Tezuka, Chaotic-integrable transition in the Sachdev-Ye-Kitaev model, Phys. Rev. Lett. **120** 241603, (2018), arXiv:1707.02197
- ¹⁰⁰ A.M. García-García, B. Loureiro, A. Romero-Bermúdez, and M. Tezuka, *García-García et al.* Reply:, Phys. Rev. Lett. **126**, 109102 (2021)
- ¹⁰¹ Jaewon Kim and Xiangyu Cao, Comment on "Chaoticintegrable transition in the Sachdev-Ye-Kitaev model", Phys. Rev. Lett. **126**, 109101 (2021), arXiv:2004.05313