

---

# Taming the sign problem of explicitly antisymmetrized neural networks via rough activation functions

---

**Nilin Abrahamsen**

University of California, Berkeley  
nilin@berkeley.edu

**Lin Lin**

University of California, Berkeley & Lawrence Berkeley National Laboratory  
linlin@math.berkeley.edu

## Abstract

Explicit antisymmetrization of a two-layer neural network is a potential candidate for a universal function approximator for generic antisymmetric functions, which are ubiquitous in quantum physics. However, this strategy suffers from a sign problem, namely, due to near-exact cancellation of positive and negative contributions, the magnitude of the antisymmetrized function may be significantly smaller than before antisymmetrization. We prove that the severity of the sign problem is directly related to the smoothness of the activation function. For smooth activation functions (e.g., tanh), the sign problem of the explicitly antisymmetrized two-layer neural network deteriorates super-polynomially with respect to the system size. On the other hand, for rough activation functions (e.g., ReLU), the deterioration rate of the sign problem can be tamed to be at most polynomial with respect to the system size. Finally, the cost of a direct implementation of antisymmetrized two-layer neural network scales factorially with respect to the system size. We describe an efficient algorithm for approximate evaluation of such a network, of which the cost scales polynomially with respect to the system size and inverse precision.

## 1 Introduction

Accurate modeling of fermionic systems is one of the most challenging and interesting problems in science. For example, the solution of the Schrödinger equation underlies all chemical properties of a given atomic system. Due to the Pauli exclusion principle, the fermionic wavefunction is antisymmetric with respect to particle exchange. When the number of fermions grows, effective parametrization of such wavefunctions can become increasingly difficult for many systems of interest. The antisymmetry condition also results in near-exact cancellation between positive and negative contributions when computing observables. This leads to the so-called fermionic sign problem (FSP), which was originally discovered in quantum Monte Carlo (QMC) simulations [15, 10, 2].

In the past decade there has been an explosive growth of techniques using neural networks (NN) as universal function approximators. The practical applicability of NNs is brought about by new software tools, hardware optimizations, as well as improved algorithms. NN approximators also significantly broaden the parameterization class for antisymmetric functions in quantum physics [12, 5, 6, 13, 16, 9]. The combination of NN with QMC methods provides a new path towards a low-scaling, systematically improvable method to approach the exact solution.

Despite recent progresses it is unclear how to construct a universal NN representation of antisymmetric functions that does not obviously suffer from the curse of the dimensionality [12, 13, 8, 4]. It is worth noting that in the absence of symmetry constraints, very simple NN structures such as a two-layer NN is already a universal function approximator [3, 7, 1]. Therefore in principle, explicitly antisymmetrizing a two-layer NN can parameterize universal antisymmetric functions. One obvious drawback of this strategy is that the computational cost of the antisymmetrization step still increases factorially with respect to the system size. Nonetheless, such an explicitly antisymmetrized NN structure has been recently studied in QMC calculations, which can yield effectively the exact ground state energy for small atoms and molecules [9].

In this paper we demonstrate that the sign problem manifests itself in a different way in these explicitly antisymmetrized NN structures. The components with positive and negative signs in the antisymmetrization process can cancel with each other nearly exactly. As a result, the magnitude of the antisymmetrized NN function can be vanishingly small, even if the magnitude of the NN before antisymmetrization is large. When the ratio of the magnitudes exceeds the limit set by the decimal digits of precision, the computational result becomes meaningless. This is a severe problem for practical NN applications, and the problem is exacerbated by the wide usage of single-precision floating point arithmetic operations on GPU architectures.

The main results of this paper is to show that for a two-layer NN, the sign problem as described above can depend critically on the choice of activation functions, and specifically, the smoothness of the activation function characterized by the decay rate in the Fourier space. The sign problem is more severe for smooth activation functions (such as  $\tanh$ ) than for rough activation functions (such as ReLU). If the tail of the activation function decays super-polynomially in the Fourier space, then the sign problem also deteriorates super-polynomially with respect to the system size (Theorem 5). On the other hand, if the tail of the Fourier-transformed activation function decays inverse-polynomially, then under suitable conditions we prove the converse statement, namely the sign problem deteriorates at most polynomially with respect to the system size (Theorem 6). Therefore as a practical strategy to tame the sign problem, we should choose rough activation functions. Though our proof is constructed for two-layer NNs, numerical results indicate that the connection between the roughness of the activation function and the sign problem also holds for multi-layer NNs.

Among all activation functions, the exponential activation function (real or complex) plays a special role in our analysis. This is because antisymmetrizing a two-layer NN with an exponential activation function gives rise to a determinant (called a Slater determinant), which can be evaluated in polynomial time. By exploring the Fourier representation of a (rough) activation function, we can approximately express the explicitly antisymmetrized two-layer NN as a linear combination of polynomially (with respect to the system size and inverse precision) many Slater determinants. This overcomes the factorial scaling barrier, and gives rise to a polynomial-time algorithm for approximate evaluation of antisymmetrized two-layer neural networks (Theorem 7).

## 2 Background

The wave function  $\psi(x_1, \dots, x_n)$  of a system of  $n$  *indistinguishable particles* in a  $d$ -dimensional space ( $d = 1, 2, 3$ ) satisfies permutation symmetry of  $|\psi|$  under interchange of the  $n$  inputs  $x_i \in \mathbb{R}^d$ . *Fermions* are indistinguishable particles which satisfy the Pauli exclusion principle and correspond to an *antisymmetric* wave function  $\psi$ . For a permutation  $\pi \in S_n$  with sign  $(-1)^\pi$ , we have  $\pi(\psi) = (-1)^\pi \psi$  where we have defined  $\pi(\psi) : \mathbb{R}^{nd} \mapsto \mathbb{C}$  by  $\pi(\psi)(x) := \psi(x_{\pi(1)}, \dots, x_{\pi(n)})$  for  $x \in \mathbb{R}^{nd}$ . For any  $f : \mathbb{R}^{nd} \rightarrow \mathbb{C}$  we can define its explicit antisymmetrization

$$\mathcal{A}f = \frac{1}{\sqrt{n!}} \sum_{\pi \in S_n} (-1)^\pi \pi(f). \quad (1)$$

As will be shown later, the prefactor  $1/\sqrt{n!}$  is the natural scaling in the antisymmetrization process.

A function defined on  $\mathbb{R}^d$  is called a single-particle function. Let  $\rho$  be a fast-decaying probability density on  $\mathbb{R}^d$  and let  $\rho_n = \rho^{\otimes n}$  be a product of single-particle densities. For simplicity of analytic computation, we may take  $\rho$  to be the density of the standard Gaussian  $\mathcal{N}(0, I_d)$  (called a Gaussian envelope). We represent an  $n$ -particle fermionic wave function as

$$\psi = \sqrt{\rho_n} \odot \mathcal{A}f = \mathcal{A}(\sqrt{\rho_n} \odot f), \quad (2)$$




Figure 1: Log-plot of  $\mathbb{E}[\|\mathcal{A}f\|_\rho^2]$  as a function of  $n$  for different activation functions  $\tau$ : exp (magenta), tanh (red), ReLU (blue), and Heaviside step function (green). The weights are sampled from the Xavier initialization (Definition 4) with  $d = 3$ ,  $m = nd$ . Shaded areas represent 90% confidence regions. Values for  $n \leq 12$  are computed by direct antisymmetrization, and dotted lines on the right are computed from (13).

where  $\odot$  denotes multiplication of function values. The wave function should be normalized as  $\|\psi\|^2 = 1$ , where  $\|\cdot\|$  is the  $L^2$ -norm on  $\mathbb{R}^{nd}$ . Eq. (2) implies  $\|\psi\| = \|\mathcal{A}f\|_\rho$  where  $\|\cdot\|_\rho$  is the norm induced by the inner product  $\langle f|g \rangle_\rho = \int \bar{f}(x)g(x)d\rho_n(x)$ .

If  $f = f_1 \otimes \dots \otimes f_n$  is a product of single-particle functions, then so is  $\phi = \phi_1 \otimes \dots \otimes \phi_n$  where  $\phi_i = \sqrt{\rho} \odot f_i$ . In this case  $\psi(x) = (\mathcal{A}\phi)(x)$  is a determinant (called the Slater determinant) denoted by  $\phi_1 \wedge \dots \wedge \phi_n$  and defined by  $(\phi_1 \wedge \dots \wedge \phi_n)(x_1, \dots, x_n) = \frac{1}{\sqrt{n!}} \det[(\phi_i(x_j))_{i,j}]$ . The normalization in Eq. (1) is such that if  $\phi_i$  are orthonormal functions on  $L^2(\mathbb{R}^d)$  then  $\psi$  is normalized by Pythagoras' theorem,  $\|\psi\| = \|\mathcal{A}f\|_\rho = 1$ .

By letting  $f$  range over a universal class of functions on  $\mathbb{R}^{nd}$  we obtain a universal class of antisymmetric functions  $\psi = \sqrt{\rho_n} \odot \mathcal{A}f$  which are not in general normalized. However, this approach has two important drawbacks *a priori*:

1. The procedure of normalizing  $\psi$  becomes numerically unstable if cancellations in (1) cause the magnitudes of  $\mathcal{A}f$  to be too small compared to  $f$ . This can be viewed as a manifestation of the fermionic sign problem in this setting.
2. The sum (1) has  $n!$  terms, making it in general intractable to evaluate the sum for all but small values of  $n$ .

We consider the case when  $f$  is given by a two-layer NN

$$f_{W,a,b}(x) = \sum_{k=1}^m a_k \tau(w^{(k)} \cdot x + b_k), \quad (3)$$

where  $\tau : \mathbb{R} \rightarrow \mathbb{R}$  is some activation function,  $w^{(k)} \in \mathbb{R}^{nd}$ , and  $b_k, a_k \in \mathbb{R}$  for each  $k = 1, \dots, m$ .

To illustrate the sign problem, Section 2 shows the magnitude  $\|\mathcal{A}f\|_\rho^2$  for four activation functions exp, tanh, ReLU ( $\tau_{\text{ReLU}}(y) = \max\{0, y\}$ ), and the Heaviside step function ( $\tau_{\text{HS}}(y) = 1_{y>0}$ ). As the system size  $n$  increases, the norm decreases with respect to  $n$  for all activation functions. However, the decay rate depends on the smoothness of the activation function. The deterioration of the sign problem is much more severe for smooth activation functions (exp, tanh) than for rough activation

<sup>1</sup>Generating *complex* amplitudes can be done in a number of ways, for example by having two output nodes (two vectors  $a_{\text{real}}, a_{\text{imag}} \in \mathbb{R}^m$ ), which is equivalent with letting  $a \in \mathbb{C}^m$ . It suffices to consider the case of a real output. Our analysis includes complex activation functions for reasons not directly related to the fact that quantum wave functions can have complex amplitudes.

functions (ReLU, Heaviside). We aim to quantify this effect and investigate more precisely how the magnitude of  $\mathcal{A}_{\tau_w}$  depends on  $\tau$ .

### 3 Main results

We state our results in terms of the Fourier transform  $\widehat{\tau}$  of the activation function. A typical activation function does not have finite integral over  $\mathbb{R}$ , so its Fourier transform is not defined as a convergent integral but rather in the sense of *tempered distributions* [14]. We will not need the precise definition of  $\widehat{\tau}$  but only that it satisfies the *Fourier inversion formula* in the sense that for  $0 < \epsilon < 1$ ,

$$\tau(y) = \frac{1}{\sqrt{2\pi}} \int_{|\theta| > \epsilon} \widehat{\tau}(\theta) e^{i\theta y} d\theta + p(y) + C_\epsilon + O_{\epsilon \rightarrow 0}(\epsilon g(y)), \quad (4)$$

where  $p$  is a polynomial of bounded degree and  $g$  is bounded by a polynomial. In particular we will be able to take  $p, C_\epsilon \equiv 0$ ,  $g(y) = |y|$  for  $\tau = \tanh$  and  $p(y) = y/2$ ,  $C_\epsilon = \frac{1}{\pi\epsilon}$ ,  $g(y) = y^2$  for  $\tau = \text{ReLU}$ . The integral in (4) converges for these activation functions since  $\int_{|\theta| > \epsilon} |\widehat{\tau}(\theta)| d\theta < \infty$ :

Table 1: Fourier transforms of different activation functions.

$\tau(y)$	$\widehat{\tau}(\theta)$	Fourier tail decay $K$
ReLU( $y$ )	$\frac{-1}{\sqrt{2\pi}\cdot\theta^2} + \sqrt{\frac{\pi}{2}} i\delta'(\theta)$	3 (rough)
tanh( $y$ )	$\frac{-i\sqrt{\pi/2}}{\sinh(\pi\theta/2)}$	$\infty$ (smooth)

Consider the decomposition the an activation function into low- and high-frequency parts as follows:

**Definition 1.** For  $\tau : \mathbb{R} \rightarrow \mathbb{C}$  define its high-pass  $\tau^{\text{HP}(t)}$  at threshold  $t > 0$  by

$$\tau^{\text{HP}(t)}(y) = \frac{1}{\sqrt{2\pi}} \int_{|\theta| > t} \widehat{\tau}(\theta) e^{i\theta y} d\theta. \quad (5)$$

Define its low-pass as the remainder  $\tau^{\text{LP}(t)} = \tau - \tau^{\text{HP}(t)}$ .

Eq. (4) says that  $\tau^{\text{LP}(\epsilon)} = p + C_\epsilon + O(\epsilon g)$  as  $\epsilon \rightarrow 0$ .

**Definition 2.** For an activation function  $\tau$  define its frequency tail  $\mathfrak{T}_\tau : (0, \infty) \rightarrow [0, \infty)$  by

$$\mathfrak{T}_\tau(t) = \int_{|\theta| \geq t} |\widehat{\tau}(\theta)|^2 d\theta. \quad (6)$$

We define the tail decay  $K \geq 0$  of  $\widehat{\tau}$  as the largest  $K$  such that  $\mathfrak{T}_\tau(t) = O(t^{-K})$  as  $t \rightarrow \infty$ . More precisely,  $K = \limsup_{t \rightarrow \infty} \frac{-\log \mathfrak{T}_\tau(t)}{\log t}$ .

**Definition 3** (Smooth and rough activation functions).

1. A function  $\tau : \mathbb{R} \rightarrow \mathbb{C}$  is smooth if its Fourier transform  $\widehat{\tau}$  has tail decay  $\infty$ , i.e., if  $\mathfrak{T}_\tau(t) = t^{-\omega(1)}$  decays faster than polynomially as  $t \rightarrow \infty$ . In particular, any activation function with  $\widehat{\tau}(\theta) = \theta^{-\omega(1)}$  is smooth.
2.  $\tau$  is rough if there exists  $k > 1$  and non-zero constants  $z_+, z_- \in \mathbb{C}$  such that  $\widehat{\tau}(\theta) = z_+ \Theta(\theta^{-k})^2$  as  $\theta \rightarrow \infty$  and  $\widehat{\tau}(\theta) = z_- \Theta(|\theta|^{-k})$  as  $\theta \rightarrow -\infty$ . In this case  $\tau$  has Fourier tail decay  $K = 2k - 1$ .

<sup>2</sup>Here  $\Theta(b_t)$  is to be understood as non-negative by definition. More generally we use the standard  $O(\cdot)$ -notation: Write  $a_t = O(b_t)$  and  $b_t = \Omega(|a_t|)$  if  $|a_t| \leq Cb_t$  for all  $t$  and some constant  $C > 0$ . Write  $\tilde{b}_t = \Theta(b_t)$  if  $\tilde{b}_t = O(b_t)$  and  $\tilde{b}_t = \Omega(b_t)$ . Write  $a_t = \tilde{O}(b_t)$  if  $a_t = O(b_t |\log t|^{O(1)})$ . Write  $a_t = o(b_t)$  and  $b_t = \omega(|a_t|)$  if  $|a_t| = \epsilon_t b_t$  for some  $\epsilon_t$  which converges to 0.

Our results below for rough activation functions hold for a more general definition of roughness which allows the Fourier transform to take the value and have varying phase. We give this definition of *generalized rough* activation functions in Appendix F.

We consider the two-layer network (3) with randomly initialized weights using two standard initializations. It is typical to initialize the biases to zero.

**Definition 4.** We say that  $f_{W,a}(x) = \sum_{k=1}^m a_k \tau(w^{(k)} \cdot x)$  is chosen with the Xavier initialization or He initialization if the  $mn + m$  weights  $W = (w_{ij}^{(k)})$ ,  $a = (a_i)$  are chosen independently from  $a_k \sim \mathcal{N}(0, c_{\text{XH}}/m)$  for each  $k = 1, \dots, m$ , and  $w_{ij}^{(k)} \sim \mathcal{N}(0, c_{\text{XH}}/(nd))$  for  $k = 1, \dots, m$ ,  $i = 1, \dots, n$ , and  $j = 1, \dots, d$ . Here  $c_{\text{XH}} = 1$  corresponds to the Xavier initialization and  $c_{\text{XH}} = 2$  to the He initialization.

**Theorem 5** (Upper bound, sign problem deteriorates super-polynomially for smooth activation functions). Let  $f_{W,a} : \mathbb{R}^{nd} \rightarrow \mathbb{C}$  be given by a two-layer neural network (3) with activation function  $\tau$  and let  $\rho = \mathcal{N}(0, I_{nd})$  be the Gaussian envelope function. If  $\tau$  is smooth and  $f_{W,a}$  is sampled from the Xavier or He initializations, then with probability  $1 - o(1)$  over  $W$ ,  $\mathbb{E}_{a|W}[\|\mathcal{A}f_{W,a}\|_{\mathcal{N}}^2] = n^{-\omega(1)}$ .

Given an activation function  $\tau : \mathbb{R} \rightarrow \mathbb{C}$  and weight vector  $w \in \mathbb{R}^{nd}$  define  $\tau_w : \mathbb{R}^{nd} \rightarrow \mathbb{C}$  by  $\tau_w(x) = \tau(w \cdot x)$ . Let  $\ell_w = \tau_w^{\text{LP}(t)}$  be the low-passed part of the activation function. The main step of the proof of the upper bound of  $\|\mathcal{A}f\|_{\mathcal{N}}^2$  is to show that the norm of the antisymmetrized function  $\|\mathcal{A}\ell_w\|_{\mathcal{N}}^2$  decreases exponentially with respect to  $n$  when  $t = O(\sqrt{n}/\log n)$  (Lemma 17). The high-passed part can be directly controlled by the tail decay of the activation function.

**Theorem 6** (Lower bound, sign problem deteriorates at most polynomially for rough activation functions). Let  $f_{W,a} : \mathbb{R}^{nd} \rightarrow \mathbb{C}$  be given by a two-layer neural network (3) with activation function  $\tau$  and weights sampled from the Xavier or He initialization, and let  $\rho = \mathcal{N}(0, I_{nd})$  be the Gaussian envelope. If  $\tau$  is rough or generalized rough with tail decay  $K$ , then with probability  $1 - o(1)$  over  $W$ ,  $\mathbb{E}_{a|W}[\|\mathcal{A}f_{W,a}\|_{\mathcal{N}}^2] = \tilde{\Omega}(n^{-(1+2/d)K})$  where  $\tilde{\Omega}$  denotes a lower bound up to log-factors. In particular the magnitude of the antisymmetrized NN decays no faster than polynomially in  $n$ .

To prove Theorem 6 we show that for the high-passed part  $h = \tau^{\text{HP}(T)}$  at sufficiently large threshold, the norm of the antisymmetrized function  $\|\mathcal{A}h_w\|_{\mathcal{N}}^2$  can be approximated by the norm  $\|h_w\|_{\mathcal{N}}^2$  before antisymmetrization (Lemma 15). The lower bound with polynomial scaling in  $n$  can also be viewed as evidence that the prefactor  $1/\sqrt{n!}$  in Eq. (1) is the appropriate scaling.

We then show that when  $f$  is given by a two-layer NN with a rough activation function,  $\mathcal{A}f$  can be computed efficiently to any inverse-polynomial precision relative to  $\sqrt{\mathbb{E}[\|\mathcal{A}f\|_{\mathcal{N}}^2]}$ :

**Theorem 7** (Deterministic polynomial time algorithm for approximate evaluation of explicitly antisymmetrized two-layer neural network). Let  $f_{W,a} : \mathbb{R}^{nd} \rightarrow \mathbb{C}$  be given by a two-layer neural network as in Theorem 6 with a rough or generalized rough activation function, and let  $\epsilon = n^{-O(1)}$ . There exists a deterministic polynomial-time algorithm  $(W, a, b, x) \mapsto S_{W,a,b}(x)$  whose output is exactly antisymmetric in  $x$  and such that with probability  $1 - o(1)$  over  $W$ ,

$$\mathbb{E}_{a|W}[\|S_{W,a,0} - \mathcal{A}f_{W,a}\|_{\mathcal{N}}^2] \leq \epsilon \mathbb{E}_{a|W}[\|\mathcal{A}f_{W,a}\|_{\mathcal{N}}^2].$$

We consider the examples of activation functions in Table 1.  $\tanh$  is smooth, so  $\|\mathcal{A}f\|_{\mathcal{N}}^2$  decays super-polynomially with  $n$  with the  $\tanh$  or sigmoid activation function. The ReLU activation is rough with tail decay 3. By Theorem 6,  $\|\mathcal{A}f\|_{\mathcal{N}}^2$  is of order  $\tilde{\Omega}(n^{-(3+6/d)})$  with the ReLU activation function. By Theorem 7 there exists an efficient algorithm to compute the outputs of  $f$  with inverse polynomial relative error when the activation function is chosen to be ReLU.

## 4 Proof sketch

We present an outline of the proofs of the main theorems, and the full proofs are given in the appendix. We begin with the observation that low-degree polynomials vanish upon antisymmetrization.

**Lemma 8.** If  $f : \mathbb{R}^{nd} \rightarrow \mathbb{C}$  is a polynomial of degree  $\deg f \leq n - 2$ , then  $\mathcal{A}f \equiv 0$ . In particular  $\mathcal{A}\tau_w \equiv 0$  if  $\tau$  is an activation function which is a polynomial of degree  $\deg \tau \leq n - 2$ .

*Proof.* By linearity it suffices to prove the claim when  $f$  is a monomial  $f(x) = \prod_{i=1}^n x_i^{r_i}$  where  $r_i \in \mathbb{N}_0$ . Since  $\deg f = \sum_i r_i \leq n - 2$  there exists a pair  $i \neq j$  such that  $r_i, r_j = 0$ . Let  $\sigma_{ij}$  be the permutation which swaps  $i$  and  $j$ . Then  $f(\sigma_{ij}(x)) = f(x)$  because  $f$  does not depend on  $x_i, x_j$ . But we also have  $f(\sigma_{ij}(x)) = -f(x)$  by antisymmetry, so  $f(x) = 0$ .  $\square$

We reduce estimates of the magnitude of  $\mathcal{A}f$  to estimates on the magnitude of  $\mathcal{A}\tau_w$ .

**Lemma 9.** For  $f_{W,a}(x)$  given by the network (3) with the He initialization (Definition 4),

$$\mathbb{E}[\|\mathcal{A}f_{W,a}\|_\rho^2 | W] = \frac{c_{\text{XH}}}{m} \sum_{k=1}^m \|\mathcal{A}\tau_{w^{(k)}}\|_\rho^2. \quad (7)$$

In particular,  $\mathbb{E}[\|\mathcal{A}f_{W,a}\|_\rho^2] = c_{\text{XH}} \mathbb{E}[\|\mathcal{A}\tau_w\|_\rho^2]$ ,  $w \sim \mathcal{N}(0, \frac{c_{\text{XH}}}{nd} I_{nd})$  does not depend on  $m$ .

The next definition characterizes the weights of the first layer with high probability under the Xavier/He initializations.

**Definition 10.** Fix a constant  $C > 1$ . We say that  $w \in \mathbb{R}^{nd}$  is typical if  $c_{\text{XH}}/2 \leq \|w\|^2 \leq 2c_{\text{XH}}$  and  $\|w\|_\infty := \max_{ij} |w_{ij}| \leq C \sqrt{\frac{\log(nd)}{nd}}$ .

In particular a typical  $w$  has  $\|w\| = \Theta(1)$ . We view  $d$  as a constant, so  $\|w\|_\infty = O(\sqrt{(\log n)/n})$  for typical  $w$ . For lower bounds we need an additional property of  $w = w^{(k)}$  sampled from the Xavier/He initializations, namely that the  $w_i \in \mathbb{R}^d$  are sufficiently separated. Indeed, if  $w_i = w_j$  for  $i \neq j$  then  $\mathcal{A}\tau_w \equiv 0$ .

**Definition 11.** For  $w \in \mathbb{R}^{nd}$  write  $\delta_w = \frac{1}{2} \min_{1 \leq i < j \leq n} \|w_i \pm w_j\|$  where the minimum is over both choices of sign  $\pm$ .

#### 4.1 Antisymmetrizing an exponential function

Consider the case when the activation function is a complex exponential function. Let “expression( $[\mathbf{x}]$ )” denote the function  $x \mapsto \text{expression}(x)$ . Then  $e^{iw \cdot [\mathbf{x}]} = \otimes_{i=1}^n e^{iw_i \cdot [\mathbf{x}]_i}$  is a product of single-particle functions, so antisymmetrizing it yields a Slater determinant,

$$\mathcal{A}(e^{iw \cdot [\mathbf{x}]}) = \mathcal{A}(\otimes_{i=1}^n e^{iw_i \cdot [\mathbf{x}]_i}) = \wedge_{i=1}^n e^{iw_i \cdot [\mathbf{x}]_i}, \quad (8)$$

where the RHS is defined as  $\frac{1}{\sqrt{n!}} \det((e^{iw_i \cdot [\mathbf{x}]_j})_{ij})$ . The overlap between two Slater determinants is the determinant of the *overlap matrix* [11], meaning that

$$\langle \mathcal{A}e^{iv \cdot [\mathbf{x}]} | \mathcal{A}e^{iw \cdot [\mathbf{x}]} \rangle_\rho = \langle \wedge_i e^{iv_i \cdot [\mathbf{x}]_i} | \wedge_i e^{iw_i \cdot [\mathbf{x}]_i} \rangle_\rho = \det B^{(v,w)}, \quad (9)$$

where  $B^{(v,w)} \in \mathbb{R}^{n \times n}$  is given by  $B_{ij}^{(v,w)} = \langle e^{iv_i \cdot [\mathbf{x}]_i} | e^{iw_j \cdot [\mathbf{x}]_j} \rangle_\rho$ . We can evaluate this as

$$B_{ij}^{(v,w)} = \mathbb{E}_{X \sim \rho} [e^{-i(v_i - w_j) \cdot X}] = \mathfrak{F}\rho(v_i - w_j). \quad (10)$$

Here we have defined the *un-normalized* Fourier transform  $\mathfrak{F}$  (also denoted by  $\mathfrak{F}_d$ ) on  $\mathbb{R}^d$  by  $\mathfrak{F}\rho(v) = \int_{\mathbb{R}^d} e^{-iv \cdot x} d\rho(x)$ . In particular  $\mathfrak{F}\rho(0) = 1$ , and  $\mathfrak{F}\rho = (2\pi)^{d/2} \widehat{\rho}$ .

#### 4.2 Fourier decomposition of the activation function

By the Fourier inversion formula (4) we have the identity of functions on  $\mathbb{R}^{nd}$

$$\tau_w(x) = \frac{1}{\sqrt{2\pi}} \int_{|\theta| > \epsilon} \widehat{\tau}(\theta) e^{i\theta w \cdot x} d\theta + p(w \cdot x) + C_\epsilon + O(\epsilon g(w \cdot x)), \quad (11)$$

We antisymmetrize (11) and apply Lemma 8 which yields that for  $n \geq \deg p + 2$ ,

$$\mathcal{A}\tau_w = \lim_{\epsilon \rightarrow 0} \mathcal{A}\tau^{\text{HP}(\epsilon)} = \lim_{\epsilon \rightarrow 0} \frac{1}{\sqrt{2\pi}} \int_{|\theta| > \epsilon} \widehat{\tau}(\theta) \mathcal{A}(e^{i\theta w \cdot [\mathbf{x}]}) d\theta, \quad (12)$$

where convergence is in the  $\mathcal{L}^2(\mathbb{R}^{nd}; \rho_n)$ -norm.

**Definition 12** (Complex-exponential overlap kernel). Define  $D_\rho : \mathbb{R}^{nd} \times \mathbb{R}^{nd} \rightarrow \mathbb{C}$  by  $D_\rho(v, w) = \det B^{(v, w)}$ , where  $B^{(v, w)} \in \mathbb{C}^{n \times n}$  is given by (10). Given a vector of weights  $w \in \mathbb{R}^{nd}$  define  $D_\rho^{(w)} : \mathbb{R}^2 \rightarrow \mathbb{C}$  by  $D_\rho^{(w)}(\theta, \tilde{\theta}) = D_\rho(\theta w, \tilde{\theta} w)$ .

Note that  $D_\rho^{(w)}$  depends on the envelope function. Eq. (9) shows that  $D_\rho^{(w)}(\theta, \tilde{\theta}) = \langle \mathcal{A}e^{i\theta w \cdot [\mathbf{x}]} | \mathcal{A}e^{i\tilde{\theta} w \cdot [\mathbf{x}]} \rangle_\rho$ . The expansion (12) then yields Lemma 13.

**Lemma 13.** If  $\iint_{\mathbb{R}^2} |\hat{\tau}(\theta)\hat{\tau}(\tilde{\theta})D_\rho^{(w)}(\theta, \tilde{\theta})| d\theta d\tilde{\theta} < \infty$ , then

$$\|\mathcal{A}\tau_w\|_\rho^2 = \frac{1}{2\pi} \iint_{\mathbb{R}^2} \overline{\hat{\tau}(\theta)}\hat{\tau}(\tilde{\theta})D_\rho^{(w)}(\theta, \tilde{\theta}) d\theta d\tilde{\theta}. \quad (13)$$

### 4.3 Properties of the complex-exponential overlap kernel

On the diagonal  $v = w$  we have  $0 \leq D_\rho(w, w) \leq 1$ . Indeed,  $B_{ij}^{(w, w)}$  is the Gram matrix of vectors  $\mathcal{A}e^{iw_i \cdot [\mathbf{x}]} \in \mathcal{L}^2(\mathbb{R}^d; \rho)$  so it satisfies  $\det(B^{(w, w)}) \leq \prod_i B_{ii}^{(w, w)} = (\mathfrak{F}_d \rho(0))^n = 1$  by Hadamard's theorem for positive semidefinite matrices.  $D_\rho(v, w)$  is a positive semidefinite kernel because it is a (infinite) Gram matrix of vectors  $\mathcal{A}e^{iw \cdot [\mathbf{x}]} \in \mathcal{L}^2(\mathbb{R}^{nd}; \rho_n)$ . In particular,  $|D_\rho(v, w)|^2 \leq D_\rho(v, v)D_\rho(w, w) \leq 1$  for all  $v, w \in \mathbb{R}^{nd}$ . Specializing these observations to  $D_\rho^{(w)}$  yields  $0 \leq D_\rho^{(w)}(\theta, \theta) \leq 1$  on the diagonal and  $|D_\rho^{(w)}(\theta, \tilde{\theta})|^2 \leq 1$  everywhere.

To approximate the behavior of  $D_\rho^{(w)}$  at large  $\theta$  and  $\tilde{\theta}$  we define a probability distribution  $\rho_w$  on  $\mathbb{R}$ .

**Definition 14.** For any  $w \in \mathbb{R}^{nd}$ , let  $\rho_w$  on  $\mathbb{R}$  be the distribution of  $w \cdot X$  where  $X \sim \rho_n$ .

We expect  $\mathcal{A}e^{i\theta w \cdot [\mathbf{x}]}$  to be roughly orthogonal to  $\mathcal{A}e^{i\tilde{\theta} w \cdot [\mathbf{x}]}$  when the frequencies  $\theta$  and  $\tilde{\theta}$  are sufficiently different. We therefore expect  $D_\rho^{(w)}(\theta, \tilde{\theta}) = \langle \mathcal{A}e^{i\theta w \cdot [\mathbf{x}]} | \mathcal{A}e^{i\tilde{\theta} w \cdot [\mathbf{x}]} \rangle_\rho$  to vanish away from the diagonal  $\tilde{\theta} = \theta$ . Moreover, when  $\theta$  is large we expect the  $n!$  terms in (1) to be roughly orthogonal, so by Pythagoras' theorem and considering the normalization factor  $1/\sqrt{n!}$  we expect  $D_\rho^{(w)}(\theta, \theta) = \|\mathcal{A}e^{i\theta w \cdot [\mathbf{x}]} \|_\rho^2 \approx \|e^{i\theta w \cdot [\mathbf{x}]} \|_\rho^2 = 1$  for large  $\theta$ . We formalize this idea by approximating  $D_\rho^{(w)}(\theta, \tilde{\theta})$  with a convolution kernel  $\mathfrak{F}\rho_w(\theta - \tilde{\theta})$  when  $\max\{|\theta|, |\tilde{\theta}|\} \geq T$  for sufficiently large  $T > 0$ . We then apply the convolution theorem to obtain

**Lemma 15.** Let  $h = \tau^{\text{HP}(T)}$  be the high-passed activation function at threshold  $T > 1$ . Then,

$$\|\mathcal{A}h_w\|_\rho^2 = \|h_w\|_\rho^2 + \varepsilon(T), \quad (14)$$

with the error term  $|\varepsilon(T)| = O(n! \delta_w^{-2} \int_{\delta_w T}^\infty \epsilon_\rho(t) t dt)$ ,  $\epsilon_\rho(\theta) = \sup\{|\mathfrak{F}\rho(y)| : y \in \mathbb{R}^d, \|y\| \geq \theta\}$ .

Lemma 15 states that at sufficiently large threshold  $T$  the magnitude of the antisymmetrization of the high-passed activation function is approximately the magnitude before antisymmetrization.

The  $\delta_w$  in the error term was defined in Definition 11. The fast decay of  $\epsilon_\rho(\theta)$  compensates for the growth of the factor  $n!$  in (14), making the approximation meaningful at polynomially large  $T$ .

Lemma 15 explains the behavior of  $D_\rho^{(w)}(\theta, \tilde{\theta})$  at large  $\theta$ . We now show that it vanishes for small  $\theta$ .

### 4.4 Gaussian envelopes and proof of upper bound

When the envelope is the standard Gaussian  $\rho = \mathcal{N}$ , the overlap kernel takes the following form:

$$D_{\mathcal{N}}(v, w) = e^{-\frac{\|v\|^2 + \|w\|^2}{2}} \det((e^{v_i \cdot w_j})_{ij}). \quad (15)$$

To obtain an upper bound on  $D_{\mathcal{N}}$  we decompose  $(e^{v_i \cdot w_j})_{ij} = \sum_{k=0}^\infty Q_k$ , bounding the ranks and operator norms of the terms  $Q_k$ . For  $L = \sum_{k=1}^{p-1} \text{rank } Q_k$  we can then bound the  $L$ -th eigenvalue as the tail sum  $\sum_{k=p}^\infty \|Q_k\|$ . Taking the product of the eigenvalues yields (Appendix E)

**Proposition 16.** Let  $\nu = 2\sqrt{d\|v\|_\infty\|w\|_\infty}$ . Then,  $\det((e^{v_i \cdot w_j})_{ij}) \leq (\nu/2)^{pn}$  for  $\nu \leq 1$ , where  $p$  is any integer such that  $\binom{p+d-1}{d} \leq n/2$  and  $p! \geq 4n^2$ .




Figure 2: Left: Heatmap of  $\mathbb{E}_w D_\rho^{(w)}(\theta, \tilde{\theta})$  where  $w \sim \mathcal{N}(0, \frac{2}{nd} I_{nd})$  for  $d = 3$  and  $n = 2$ . Positive values are blue and negative values red. Right: Diagonal cross-section  $\mathbb{E}_w D_{\mathcal{N}}^{(w)}(\theta, \theta)$  for  $n = 100$ .

Apply the bound to  $D_{\mathcal{N}}(\theta w, \theta w)$  with  $p = \Theta(n^{1/d})$  to get that for  $\theta \leq t := (2\sqrt{d}\|w\|_\infty)^{-1}$ ,

$$D_{\mathcal{N}}^{(w)}(\theta, \theta) = (\frac{1}{2}\theta/t)^{\Omega(n^{1+1/d})}. \quad (16)$$

Eq. (12) implies the *triangle inequality*  $\|\mathcal{A}\tau_w\|_\rho \leq \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} |\hat{\tau}(\theta)| \sqrt{D_{\mathcal{N}}^{(w)}(\theta, \theta)} d\theta$  because  $\sqrt{D_\rho^{(w)}(\theta, \theta)} = \|\mathcal{A}e^{i\theta w \cdot [x]}\|_\rho$  by definition. Apply this triangle inequality to the low-pass part and bound the integrand using (16) to cancel the pole of  $|\hat{\tau}|$  at 0. We then obtain,

**Lemma 17.** *Let  $\ell = \tau^{\text{LP}(t)}$  be the low-pass at threshold  $t = (2\sqrt{d}\|w\|_\infty)^{-1}$ . If  $w$  is typical then  $t = \Omega(\sqrt{n/\log n})$  and  $\|\mathcal{A}\ell_w\|_{\mathcal{N}} = O(2^{-\Omega(n^{1+1/d})})$ .*

The high-pass part can be estimated using only the triangle inequality and the  $\ell^1$ -decay of  $\hat{\tau}$ , but for a sharper bound in terms of the  $\ell^2$ -decay we use (15) to obtain the relation

$$D_{\mathcal{N}}^{(w)}(\theta, \tilde{\theta}) = e^{-\frac{\|w\|^2}{2}(|\theta| - |\tilde{\theta}|)^2} D_{\mathcal{N}}^{(w)}(\theta_{\text{g.m.}}, \pm \theta_{\text{g.m.}}), \quad (17)$$

where the geometric mean  $\theta_{\text{g.m.}} = |\theta\tilde{\theta}|^{1/2}$  and  $\pm$  is the sign of  $\theta\tilde{\theta}$ . In particular  $D_{\mathcal{N}}^{(w)}(\theta, \tilde{\theta}) \leq e^{-\frac{1}{2}\|w\|^2(|\theta| - |\tilde{\theta}|)^2}$ . We then apply (13) and Cauchy-Schwarz to obtain

**Lemma 18.** *Let  $h = \tau^{\text{HP}(t)}$  be the high-passed activation function at  $t > 1$ . Then,  $\|\mathcal{A}h_w\|_{\mathcal{N}}^2 \leq \frac{4}{\sqrt{2\pi}\|w\|} \mathfrak{T}\hat{\tau}(t)$ . So  $\|\mathcal{A}h_w\|_{\mathcal{N}}^2 = O(t^{-K})$  for typical  $w$ .*

For typical  $w$  choose threshold  $t = \Omega(\sqrt{n/\log n})$  as in Lemma 17 and decompose  $\mathcal{A}\tau_w = \mathcal{A}\ell_w + \mathcal{A}h_w$ . Apply Lemma 17 and Lemma 18 to obtain

**Corollary 19.** *If  $\tau$  has frequency tail decay  $K < \infty$  then  $\|\mathcal{A}\tau_w\|_{\mathcal{N}}^2 = \tilde{O}(n^{-K/2})$  for typical  $w$ . If  $\tau$  is smooth ( $K = \infty$ ), then  $\|\mathcal{A}\tau_w\|_{\mathcal{N}}^2 = n^{-\omega(1)}$  for typical  $w$ .*

#### 4.5 Proof of lower bound

For the Gaussian envelope function the error term in Lemma 15 becomes  $O(\delta_w^{-2} e^{-\delta_w^2 T^2/2 + n \log n})$ . To prove the lower bound we pick  $T = \frac{2}{\delta_w}(\sqrt{n \log n} + \log(1/\delta_w))$  and decompose  $\tau$  into  $\ell = \tau^{\text{LP}(T)}$  and  $h = \tau^{\text{HP}(T)}$ . We then expand  $\|\mathcal{A}\ell + \mathcal{A}h\|_{\mathcal{N}}^2$  and use that  $\|\mathcal{A}\ell\|_{\mathcal{N}}^2 \geq 0$  to get  $\|\mathcal{A}\tau_w\|_{\mathcal{N}}^2 \geq \|\mathcal{A}h_w\|_{\mathcal{N}}^2 - 2|\langle \mathcal{A}\ell_w, \mathcal{A}h_w \rangle_{\mathcal{N}}|$ . The result follows by bounding  $|\langle \mathcal{A}\ell_w, \mathcal{A}h_w \rangle_{\mathcal{N}}|$  and applying Lemma 15 to get  $\|\mathcal{A}h_w\|_{\mathcal{N}}^2 \geq \|h_w\|_{\mathcal{N}}^2 - O(e^{-n \log n})$ . We then use the convolution theorem to show that under our definition of roughness (or generalized roughness),  $\|h_w\|_{\mathcal{N}}^2 = \Omega(\mathfrak{T}\hat{\tau}(T)) = \Omega(T^{-K})$ . There is a gap between the upper bound  $O(t^{-K}) = \tilde{O}(n^{-K/2})$  and this lower bound  $\Omega(T^{-K}) = \tilde{\Omega}(n^{-(1+2/d)K})$  due to the factor  $1/\delta_w$  in the threshold  $T$  compared to  $t$ .



## 4.6 Efficient algorithm

Recall (8) which gives the antisymmetrization with an exponential activation function as a determinant,  $\mathcal{A}(e^{i\mathbf{w} \cdot \mathbf{x}}) = \frac{1}{\sqrt{n!}} \det((e^{i\mathbf{w}_i \cdot \mathbf{x}})_{ij})$ . Let  $\ell^{(t)} = \tau^{\text{LP}(t)}$  and  $h^{(T)} = \tau^{\text{HP}(T)}$  be the low-pass and high-pass of  $\tau$  at thresholds  $t < T$ . We approximate  $\mathcal{A}\tau_w$  by removing the low-passed and high-passed components: Apply (12) to  $\tau^{\text{HP}(t)} - \tau^{\text{HP}(T)}$  to obtain  $\mathcal{A}\tau_w(x) = \alpha_w + \mathcal{A}\ell_w^{(t)} + \mathcal{A}h_w^{(T)}$ , where

$$\alpha_w(x) = \frac{1}{\sqrt{2\pi n!}} \int_{[-T, T] \setminus [-t, t]} \hat{\tau}(\theta) \det((e^{i\theta \mathbf{w}_i \cdot \mathbf{x}_j})_{ij}) d\theta. \quad (18)$$

The integrand can be computed at a single  $\theta$  in time  $O(n^3)$ . Apply the Lemma 17 and Lemma 18 (with a different choice of threshold for the high-pass) to bound the truncation error.

**Lemma 20** (Truncation error bound). *Suppose  $\hat{\tau}$  has tail decay  $K$ . Let  $t = \max\{\frac{1}{2\sqrt{d}\|w\|_\infty}, 1\}$  and, given  $\epsilon > 0$ , let  $T = \epsilon^{-1/K}$ . Then  $\|\mathcal{A}\tau_w - \alpha_w\|_{\mathcal{N}}^2 = O(2^{-n^{1+1/d}} + \epsilon)$  for typical  $w$ .*

We approximate  $\alpha_w(x)$  by a sum  $S_w(x) = \frac{1}{\sqrt{2\pi n!}} \sum_{p=\pm 1, \dots, \pm N} c_p \det((e^{i\theta_p \mathbf{w}_i \cdot \mathbf{x}_j})_{ij})$ , where  $\theta_1, \dots, \theta_N$  are a discretization of  $[t, T]$ . For evenly spaced  $\theta_p$  and appropriate coefficients  $c_p$ , and define  $\|\partial_\theta D_{\mathcal{N}}^{(w)}\|_\infty := \sup_{\theta, \tilde{\theta}} |\frac{\partial}{\partial \theta} D_{\mathcal{N}}^{(w)}(\theta, \tilde{\theta})|$ , we have

**Lemma 21.**  $\|S_w - \alpha_w\|_{\mathcal{N}}^2 \leq \frac{2}{\pi} \frac{T}{N} \|\partial_\theta D_{\mathcal{N}}^{(w)}\|_\infty (\int_{t \leq |\theta| \leq T} |\hat{\tau}(\theta)|)^2 = O(\frac{T}{N} \|\partial_\theta D_{\mathcal{N}}^{(w)}\|_\infty)$ .

In the presence of bias terms  $b_k$  in (3) we can efficiently compute an approximation  $S_{w,b}$  to the function  $\mathcal{A}\tau(w^{(k)} \cdot \mathbf{x} + b_k)$  with the same error bound. Indeed, the bias term results in a shift of the activation function which corresponds to multiplying the Fourier transform by an oscillating phase. Since the upper bounds do not depend on the phase of the Fourier transform, the same truncation error bound applies. Theorem 5 holds for arbitrary bias terms for the same reason.

*Proof of Theorem 7.* Because  $\tau$  is rough we have  $\mathbb{E}[\|\mathcal{A}f\|_{\mathcal{N}}|W] = n^{-O(1)}$  with probability  $1 - o(1)$ . Given target relative error  $\epsilon = n^{-O(1)}$  it suffices to achieve absolute error  $\epsilon' = \epsilon \cdot \mathbb{E}[\|\mathcal{A}f\|_{\mathcal{N}}|W] = n^{-O(1)}$ .  $\|S_w - \mathcal{A}\tau_w\|_{\mathcal{N}}^2 = O(\epsilon' + n^{O(1)} \|\partial_\theta D_{\mathcal{N}}^{(w)}\|_\infty / N)$  by Lemma 20 and Lemma 21. In Corollary 32 of Appendix H we show that  $\|\partial_\theta D_{\mathcal{N}}^{(w)}\|_\infty = n^{O(1)}$ , so  $\|S_w - \mathcal{A}\tau_w\|_{\mathcal{N}}^2 = O(\epsilon' + n^{O(1)}/N)$ . It then suffices to pick  $N = n^{O(1)}$ . Let  $S_{W,a,b} = \sum_{k=1}^m a_k S_{w^{(k)}, b_k}$  and apply Lemma 9 to the difference  $f_{W,a,b} - S_{W,a,b}$  to extend the termwise error bound to the sum (3). The computational cost of evaluating  $S_w(x)$  is  $O(n^3 N)$ .  $\square$

## 5 Empirical generalization to multi-layer networks

It is natural to ask whether the advantage of rough activation functions against cancellations remains as the depth of the neural network grows. We consider networks of depth  $L = 3, 4, 5$  and compare  $\|\mathcal{A}f\|_{\mathcal{N}}^2$  between two choices of activation functions: The smooth  $\tanh$  and the rough *normalized double ReLU* (DReLU)  $\tau_\kappa(y) = \kappa \max\{-1, \min\{1, y\}\}$  where  $\kappa \approx 0.875$  is chosen such that  $\mathbb{E}[\tau_\kappa(Z)^2] = \mathbb{E}[\tanh(Z)^2]$  for standard-Gaussian  $Z \sim \mathcal{N}(0, 1)$ .




Figure 3: Tanh and the normalized DReLU.

We take  $d = 3$ , let all layers have width  $m = 3n$ , and instantiate NNs from the Xavier initialization (independent Gaussian weights with variance  $1/m$  where  $m$  is the number of neurons in the preceding layer). Fig. 4 shows that the rough activation function maintains its advantage for networks with more layers.




Figure 4: Log-scale comparison between  $\mathbb{E}[\|\mathcal{A}f\|_{\mathcal{N}}^2]$  for smooth and rough activation functions (tanh vs normalized DReLU) for antisymmetrized neural networks of different depths. Shaded areas show 90% confidence regions.

## 6 Conclusion

Using the Fourier representation of the activation function, we observe that a rough activation function is necessary to tame the near-exact cancellations when antisymmetrizing two-layer NNs. The Fourier perspective also provides a polynomial-time algorithm for approximately evaluating explicitly antisymmetrized two-layer NNs. It may be possible that explicitly antisymmetrized two-layer NNs provides a path towards universal approximation of a class of antisymmetric functions without suffering from curse of dimensionality. Our work also raises intriguing open questions about how the cancellations and efficient algorithms generalize to antisymmetrized multi-layer NNs.

## Acknowledgment

This work was partially supported by the NSF Quantum Leap Challenge Institute (QLCI) program through grant number OMA-2016245 (N.A.), the U.S. Department of Energy under the CAMERA program and the SciDAC program (L.L.). L.L. is a Simons Investigator. The authors thank Jeffmin Lin and Lexing Ying for helpful discussions.

## References

- [1] Andrew R Barron. Universal approximation bounds for superpositions of a sigmoidal function. *IEEE Trans. Inform. Theory*, 39:930–945, 1993.
- [2] D. M. Ceperley. Fermion nodes. *J. Stat. Phys.*, 63(5-6):1237–1267, 1991.
- [3] George Cybenko. Approximation by superpositions of a sigmoidal function. *Mathematics of Control, Signals, and Systems*, 2(4):303–314, 1989.
- [4] Jiequn Han, Yingzhou Li, Lin Lin, Jianfeng Lu, Jiefu Zhang, and Linfeng Zhang. Universal approximation of symmetric and anti-symmetric functions. *arXiv:1912.01765*, 2019.
- [5] Jiequn Han, Linfeng Zhang, and Weinan E. Solving many-electron Schrödinger equation using deep neural networks. *J. Comput. Phys.*, 399:108929, 2019.
- [6] Jan Hermann, Zeno Schatzle, and Frank Noe. Deep neural network solution of the electronic schrödinger equation. *Nature Chem.*, 12:891–897, 2020.
- [7] Kurt Hornik, Maxwell Stinchcombe, and Halbert White. Multilayer feedforward networks are universal approximators. *Neural Networks*, 2(5):359–366, 1989.
- [8] Marcus Hutter. On representing (anti)symmetric functions. Technical Report arXiv:2007.15298, DeepMind, 2020.
- [9] Jeffmin Lin, Gil Goldshlager, and Lin Lin. Explicitly antisymmetrized neural network layers for variational Monte Carlo simulation. *arXiv:2112.03491*, 2021.
- [10] EY Loh Jr, JE Gubernatis, RT Scalettar, SR White, DJ Scalapino, and RL Sugar. Sign problem in the numerical simulation of many-electron systems. *Phys. Rev. B*, 41(13):9301, 1990.

- [11] Per-Olov Löwdin. Quantum theory of many-particle systems. i. physical interpretations by means of density matrices, natural spin-orbitals, and convergence problems in the method of configurational interaction. *Phys. Rev.*, 97:1474–1489, Mar 1955.
- [12] Di Luo and Bryan K. Clark. Backflow transformations via neural networks for quantum many-body wave functions. *Phys. Rev. Lett.*, 122:226401, 2019.
- [13] David Pfau, James S. Spencer, Alexander G. D. G. Matthews, and W. M. C. Foulkes. Ab initio solution of the many-electron schrödinger equation with deep neural networks. *Phys. Rev. Research*, 2:033429, 2020.
- [14] M. Reed and B. Simon. *I: Functional Analysis. Methods of Modern Mathematical Physics.* Elsevier Science, 1981.
- [15] Sandro Sorella, S Baroni, Roberto Car, and M Parrinello. A novel technique for the simulation of interacting fermion systems. *EPL*, 8(7):663, 1989.
- [16] James Stokes, Javier Robledo Moreno, Eftychios A. Pnevmatikakis, and Giuseppe Carleo. Phases of two-dimensional spinless lattice fermions with first-quantized deep neural-network quantum states. *Phys. Rev. B*, 102:205122, 2020.

## A The Fourier inversion formula holds for ReLU and tanh

We verify Eq. (4) which asserts that  $\tau^{\text{LP}(\epsilon)} = p + C_\epsilon + O(\epsilon g)$  where  $p$  is a low-degree polynomial and  $g$  is bounded by a polynomial. Here we have defined  $\tau^{\text{LP}(\epsilon)} = \tau - \tau^{\text{HP}(\epsilon)}$ .

1.  $\tau = \text{ReLU}$ : We first evaluate the high-pass part

$$\tau^{\text{HP}(\epsilon)}(y) = |y|/2 - \frac{\cos(\epsilon y)}{\pi \epsilon} - \frac{y \text{Si}(\epsilon y)}{\pi},$$

where  $\text{Si}(y) = \int_0^y \frac{\sin s}{s} ds$ . Since  $\text{ReLU}(y) = |y|/2 + y/2$ ,

$$\tau^{\text{LP}(\epsilon)}(y) = y/2 + \frac{\cos(\epsilon y)}{\pi \epsilon} + \frac{y \text{Si}(\epsilon y)}{\pi}. \quad (19)$$

Write  $\tau^{\text{LP}(\epsilon)} = p + C_\epsilon + \varepsilon$  where

$$p(y) = y/2, \quad C_\epsilon = \frac{1}{\pi \epsilon}.$$

Then the remainder satisfies

$$|\varepsilon| \leq \left| \frac{\cos(\epsilon y) - 1}{\pi \epsilon} \right| + \left| \frac{y \text{Si}(\epsilon y)}{\pi} \right| \leq \frac{(\epsilon y)^2}{2\pi \epsilon} + \frac{y \cdot (\epsilon y)}{\pi} = \epsilon g(y), \quad g(y) := \frac{3}{2\pi} y^2. \quad (20)$$

2.  $\tau = \tanh$ : We can write the low-pass part as an absolutely convergent integral as

$$\tau^{\text{LP}(\epsilon)}(y) = \frac{1}{\sqrt{2\pi}} \int_{-\epsilon}^{\epsilon} \frac{-i\sqrt{\pi/2}}{\sinh(\pi\theta/2)} (e^{i\theta y} - 1) d\theta.$$

Let  $p, C_\epsilon \equiv 0$  and bound

$$|\tau^{\text{LP}(\epsilon)}(y)| \leq \frac{1}{\sqrt{2\pi}} \int_{-\epsilon}^{\epsilon} \frac{\sqrt{\pi/2}}{|\pi\theta/2|} |\theta y| d\theta = \epsilon g(y), \quad g(y) := \frac{2}{\pi} |y|. \quad (21)$$

## B Typical weights and reduction to single-neuron case

**Lemma 9.** For  $f_{W,a}(x)$  given by the network (3) with the He initialization (Definition 4),

$$\mathbb{E}[\|\mathcal{A}f_{W,a}\|_\rho^2 | W] = \frac{c_{\text{XH}}}{m} \sum_{k=1}^m \|\mathcal{A}\tau_{w^{(k)}}\|_\rho^2. \quad (7)$$

In particular,  $\mathbb{E}[\|\mathcal{A}f_{W,a}\|_\rho^2] = c_{\text{XH}} \mathbb{E}[\|\mathcal{A}\tau_w\|_\rho^2]$ ,  $w \sim \mathcal{N}(0, \frac{c_{\text{XH}}}{nd} I_{nd})$  does not depend on  $m$ .

*Proof.* Expand  $\|\mathcal{A}f_{W,a}\|_\rho^2 = \sum_{k,l=1}^m a_k a_l \langle \mathcal{A}\tau_{w^{(k)}} | \mathcal{A}\tau_{w^{(l)}} \rangle_\rho$ . The  $a_k \sim \mathcal{N}(0, c_{\text{XH}}/m)$ 's are independent so  $\mathbb{E}[a_k a_l] = \delta_{kl} c_{\text{XH}}/m$ , and

$$\mathbb{E}[\|\mathcal{A}f_{W,a}\|_\rho^2 | W] = \frac{c_{\text{XH}}}{m} \sum_{k,l=1}^m \delta_{kl} \mathbb{E}[\langle \mathcal{A}\tau_{w^{(k)}} | \mathcal{A}\tau_{w^{(l)}} \rangle_\rho] = \frac{c_{\text{XH}}}{m} \sum_{k=1}^m \|\mathcal{A}\tau_{w^{(k)}}\|_\rho^2. \quad (22)$$

Taking the expectation over  $W$  yields  $\mathbb{E}_{W,a}[\|\mathcal{A}f_{W,a}\|_\rho^2] = c_{\text{XH}} \mathbb{E}[\|\mathcal{A}\tau_v\|_\rho^2]$ .  $\square$

**Lemma 22.** For  $w \sim \mathcal{N}(0, \frac{2}{nd} I_{nd})$  sampled from the He initialization,

$$P(\delta_w < \delta) \leq 2 \binom{n}{2} \left( \frac{2nd\delta^2}{c_{\text{XH}}\pi} \right)^{d/2} |B_d|, \quad (23)$$

where  $|B_d|$  is the volume of the unit ball in  $\mathbb{R}^d$ . For constant  $d$  this is  $O(n^{2+d/2}\delta^d)$ .

*Proof.* If  $w_i$  are sampled independently from a distribution  $\mathcal{W}$  on  $\mathbb{R}^d$ , then

$$\begin{aligned} P(\delta_w < \delta) &\leq \sum_{i<j} P\left(\frac{1}{2}\|w_i - w_j\| < \delta\right) + P\left(\frac{1}{2}\|w_i + w_j\| < \delta\right) \\ &\leq \sum_{i<j} \max_{w' \in \mathbb{R}^d} P(\|w_i - w_j\| < 2\delta \mid w_j = w') + \max_{w' \in \mathbb{R}^d} P(\|w_i + w_j\| < 2\delta \mid w_j = w') \\ &\leq 2 \binom{n}{2} (2\delta)^d |B_d| \|\mathcal{W}\|_\infty, \end{aligned} \quad (24)$$

where  $\|\mathcal{W}\|_\infty$  is the supremum of the density. For the He initialization  $w_i \sim \mathcal{N}(0, \frac{2}{nd} I_d)$  we have  $\|\mathcal{W}\|_\infty = (2\pi \cdot \frac{c_{\text{XH}}}{nd})^{-d/2} = (\frac{2c_{\text{XH}}\pi}{nd})^{-d/2}$ .  $\square$

For fixed  $d$ ,  $v = w^{(k)}$  satisfies that  $P(\delta_v < \delta) = O(n^{2+d/2}\delta^d)$ . Then with probability  $1 - o(1)$ ,  $v$  has typical separation as defined below.

**Definition 23.** Fix a function  $\delta(n) = o(n^{-(1/2+2/d)})$  as  $n \rightarrow \infty$  (for concreteness we let  $\delta(n) = n^{-(1/2+2/d)}/\sqrt{\log n}$ ). We say that  $v$  has typical separation if  $\delta_v \geq \delta(n)$ . We say that  $W \in \mathbb{R}^{m \times nd}$  is typical if each  $w^{(k)}$  is typical and at least half the  $w^{(k)}$  have typical separation.

**Lemma 24.** Let  $d$  be constant, let  $m = O(n^{C'})$  for some constant  $C'$ , and let  $f$  be sampled as in Definition 4. Then  $W$  is typical with probability  $1 - o(1)$  for some constant  $C$  in Definition 10 depending on  $C'$ . For such  $W$ ,

$$\frac{c_{\text{XH}}}{2} \inf_{w \in S'} \|\mathcal{A}\tau_w\|_\rho^2 \leq \mathbb{E}_{a|W}[\|\mathcal{A}f\|_\rho^2] \leq c_{\text{XH}} \sup_{w \in S} \|\mathcal{A}\tau_w\|_\rho^2, \quad (25)$$

where  $S$  is the set of typical  $w \in \mathbb{R}^{nd}$  and  $S' \subset S$  is the set of typical  $w$  which have typical separation.

*Proof.* Recall that by definition,  $W$  is typical if each  $w^{(k)}$  is typical and at least half the  $w^{(k)}$  have typical separation. The distribution of  $\frac{nd}{c_{\text{XH}}} \|w^{(k)}\|^2 \sim \chi^2(nd)$  implies that for each  $k = 1, \dots, m$ ,  $P(\|w^{(k)}\|^2 < c_{\text{XH}}/2) \leq (\sqrt{e}/2)^{nd/2}$  and  $P(\|w^{(k)}\|^2 > 2c_{\text{XH}}) \leq (2/e)^{nd/2}$  since  $\mathbb{E}[\|w^{(k)}\|^2] = c_{\text{XH}}$ . By a union bound we have  $c_{\text{XH}}/2 \leq \|w^{(k)}\|^2 \leq 2c_{\text{XH}}$  for all  $k = 1, \dots, m$  with probability  $1 - 2m2^{-\Omega(nd)}$ . Furthermore,  $w_{ij}^{(k)} \sim \mathcal{N}(0, \frac{c_{\text{XH}}}{nd})$  and a union bound imply that

$$P(|w_{ij}^{(k)}| \geq t \text{ for some } i, j, k) \leq 2mnd e^{-\frac{ndt^2}{2c_{\text{XH}}}} = O\left(e^{-\frac{ndt^2}{2c_{\text{XH}}} + (C'+1)\log n + \log d}\right).$$

Given any  $C'' > 0$  we may let  $t = 2\sqrt{C' + C'' + 1} \sqrt{\frac{\log(nd)}{nd}}$  and obtain that  $\|w^{(k)}\|_\infty \leq t$  for all  $k$  with probability at least  $1 - n^{-C''}$ . This shows that each  $w$  is typical with probability  $1 - 1/n$  for appropriate  $C$  in Definition 10.

If  $P(\delta_{w^{(k)}} \geq \delta) \rightarrow 1$  for each fixed  $k$  then the probability that  $\delta(w^{(k)}) \geq \delta$  for at least half the  $k = 1, \dots, m$  also converges to 1. Eq. (25) follows from Lemma 9.  $\square$

## C Properties of complex-exponential overlap kernel

**Lemma 25.** Let  $\delta_w = \frac{1}{2} \min_{i \neq j} \|w_i \pm w_j\|$  and let  $\rho_w$  be as in Definition 14. Then,

$$|D_\rho^{(w)}(\theta, \tilde{\theta}) - \mathfrak{F}\rho_w(\theta - \tilde{\theta})| \leq n! \epsilon_\rho(\delta_w \theta_{\max}), \quad (26)$$

where  $\theta_{\max} = \max\{|\theta|, |\tilde{\theta}|\}$  and  $\epsilon_\rho(\theta) = \sup\{|\mathfrak{F}_d \rho(y)| : y \in \mathbb{R}^d, \|y\| \geq \theta\}$ .  $\mathfrak{F}\rho_w$  can be expressed in terms of the  $nd$ -dimensional Fourier transform as  $\mathfrak{F}\rho_w(\theta) = \mathfrak{F}_{nd} \rho_n(\theta w)$ .

*Proof.* By definition we have

$$D_\rho^{(w)}(\theta, \tilde{\theta}) = \det(\mathfrak{F}_d \rho(\theta w_i - \tilde{\theta} w_j)_{ij}) = \sum_{\pi \in S_n} (-1)^\pi \mathfrak{F}_{nd} \rho_n(\theta w - \tilde{\theta} \pi(w)).$$

The  $\pi = 1$  term equals

$$\mathfrak{F}_{nd} \rho_n((\theta - \tilde{\theta})w) = \mathfrak{F}\rho_w(\theta - \tilde{\theta}). \quad (27)$$

To verify the identity (27) and the claim at the end of the lemma, write

$$\mathfrak{F}_{nd} \rho_n(\theta w) = \mathbb{E}_{X \sim \rho_n} [e^{-i(\theta w) \cdot X}] = \mathbb{E}_{X \sim \rho_n} [e^{-i\theta(w \cdot X)}] = \mathbb{E}_{y \sim \rho_w} [e^{-i\theta y}] = \mathfrak{F}\rho_w(\theta).$$

The difference on the LHS of (26) is exactly  $\varepsilon = \sum_{\pi \neq 1} (-1)^\pi \mathfrak{F}_{nd} \rho_n(\theta w - \tilde{\theta} \pi(w))$ . Apply the triangle inequality and use the fact that  $\|\epsilon_\rho\|_\infty = \|\mathfrak{F}_{nd} \rho_n\|_\infty \leq 1$  because  $\rho$  is a probability distribution to obtain

$$\begin{aligned} |\varepsilon| &\leq n! \max_{\pi \neq 1} |\mathfrak{F}_{nd} \rho_n(\theta w - \tilde{\theta} \pi(w))| \\ &\leq n! \max_{\pi \neq 1} \prod_{i=1}^n \epsilon_\rho(\|\theta w_i - \tilde{\theta} \pi(w)_i\|) \\ &\leq n! \max_{\pi \neq 1} \min_i \epsilon_\rho(\|\theta w_i - \tilde{\theta} \pi(w)_i\|) \leq n! \epsilon_\rho(|\theta| \tilde{\delta}(w)), \end{aligned} \quad (28)$$

where

$$\tilde{\delta}(v) = \min_{\tilde{\theta}} \min_{\pi \neq 1} \max_i \|v_i - \tilde{\theta} \pi(v)_i\|. \quad (29)$$

We show that  $\tilde{\delta}_w \geq \delta_w$  in Lemma 26. The result follows from symmetry in  $\theta$  and  $\tilde{\theta}$ .  $\square$

**Lemma 26.** Let  $\tilde{\delta}_w := \min_{\tilde{\theta} \in \mathbb{R}} \min_{\pi \neq 1} \max_i \|w_i - \tilde{\theta} \pi(w)_i\|$ . Then  $\tilde{\delta}_w \geq \delta_w$ .

*Proof.* Let  $\tilde{\theta}$  and  $\sigma = \pi^{-1} \neq 1$  be arbitrary. We need to show that  $\|w_i - \tilde{\theta} w_{\sigma(i)}\| \geq \delta_w$  for some  $i$ .

Since  $\sigma \neq 1$  there exists  $i$  such that  $\sigma(i) \neq i$ , therefore  $\|w_i - s w_{\sigma(i)}\| \geq 2\delta_w$  for  $s = \pm 1$ . If  $\|w_i - \tilde{\theta} w_{\sigma(i)}\| \geq \delta_w$  then we have found  $i$  and we are done. Otherwise, let  $s = \text{sign } \tilde{\theta}$ . Then,

$$\begin{aligned} 2\delta_w \leq \|w_i - s w_{\sigma(i)}\| &\leq \|w_i - \tilde{\theta} w_{\sigma(i)}\| + \|\tilde{\theta} w_{\sigma(i)} - s w_{\sigma(i)}\| \\ &= \|w_i - \tilde{\theta} w_{\sigma(i)}\| + |\tilde{\theta} - s| \cdot \|w_{\sigma(i)}\| \\ &< \delta_w + |\tilde{\theta} - s| \max_j \|w_j\|. \end{aligned} \quad (30)$$

Rearranging (30) we yields

$$|\tilde{\theta} - s|R > \delta_w, \quad R = \max_j \|w_j\|. \quad (31)$$

Pick  $j$  such that  $\|w_j\| = R$ . If  $|\tilde{\theta}| \geq 1$  then (31) says  $(|\tilde{\theta}| - 1)R > \delta_w$ . Let  $i = \pi(j)$  so that  $\sigma(i) = j$  and  $\|w_{\sigma(i)}\| = R$ . Then,

$$\|w_i - \tilde{\theta} w_{\sigma(i)}\| \geq \|\tilde{\theta} w_{\sigma(i)}\| - \|w_i\| = |\tilde{\theta}|R - \|w_i\| \geq (|\tilde{\theta}| - 1)R \geq \delta_w.$$

If instead  $|\tilde{\theta}| < 1$  then (31) says  $(1 - |\tilde{\theta}|)R > \delta_w$ . We then use

$$\|w_j - \tilde{\theta} w_{\sigma(j)}\| \geq \|w_j\| - \|\tilde{\theta} w_{\sigma(j)}\| = R - \|\tilde{\theta} w_{\sigma(j)}\| \geq (1 - |\tilde{\theta}|)R \geq \delta_w. \quad \square$$

**Lemma 15.** Let  $h = \tau^{\text{HP}(T)}$  be the high-passed activation function at threshold  $T > 1$ . Then,

$$\|\mathcal{A}h_w\|_\rho^2 = \|h_w\|_\rho^2 + \varepsilon(T), \quad (14)$$

with the error term  $|\varepsilon(T)| = O(n! \delta_w^{-2} \int_{\delta_w T}^\infty \varepsilon_\rho(t) t dt)$ ,  $\varepsilon_\rho(\theta) = \sup\{|\mathfrak{F}\rho(y)| : y \in \mathbb{R}^d, \|y\| \geq \theta\}$ .

*Proof.* By (13) we have  $\|\mathcal{A}h_w\|_\rho^2 = \frac{1}{2\pi} \iint_{|\theta|, |\tilde{\theta}| \geq T} D_\rho^{(w)}(\theta, \tilde{\theta}) \overline{\widehat{\tau}(\theta)} \widehat{\tau}(\tilde{\theta}) d\theta d\tilde{\theta}$ . Apply Lemma 25 to approximate  $D_\rho^{(w)}(\theta, \tilde{\theta})$  by  $\mathfrak{F}\rho_w(\theta - \tilde{\theta})$ . Recall that  $\widehat{h} = \widehat{\tau} \odot 1_{\mathbb{R} \setminus [-T, T]}$ . The resulting approximation is, by the convolution theorem and Plancherel's identity,

$$\begin{aligned} \frac{1}{2\pi} \iint \mathfrak{F}\rho_w(\theta - \tilde{\theta}) \overline{\widehat{h}(\theta)} \widehat{h}(\tilde{\theta}) d\theta d\tilde{\theta} &= \frac{1}{2\pi} \int \overline{\widehat{h}(\theta)} (\mathfrak{F}\rho_w * \widehat{h})(\theta) d\theta \\ &= \int |h(y)|^2 \rho_w(y) dy = \mathbb{E}_{X \sim \rho_n} [|h(w \cdot X)|^2] = \|h_w\|_\rho^2. \end{aligned} \quad (32)$$

$\widehat{\tau}$  is bounded on  $\mathbb{R} \setminus [-T, T]$  by assumption. The error of the approximation is then bounded by a constant times

$$\iint_{|\theta|, |\tilde{\theta}| \geq T} |D_\rho^{(w)}(\theta, \tilde{\theta}) - \mathfrak{F}\rho_w(\theta - \tilde{\theta})| d\theta d\tilde{\theta} \leq 8 \int_T^\infty \int_T^\theta n! \varepsilon_\rho(\delta_w \theta) d\tilde{\theta} d\theta \leq 8n! \int_T^\infty \varepsilon_\rho(\delta_w \theta) \theta d\theta.$$

The error bound follows by substituting  $t = \delta_w \theta$ .  $\square$

## D Properties of complex-exponential overlap kernel for the Gaussian envelope

For the Gaussian envelope  $\rho = \mathcal{N}(0, I_d)$  we have  $\mathfrak{F}_d \rho(v) = e^{-\|v\|^2/2}$ ,  $\mathfrak{F}\rho_w(\theta) = e^{-\|w\|^2 \theta^2/2}$ , and  $\varepsilon_\rho(\theta) = e^{-\theta^2/2}$ . In Lemma 15 the integral in the error term becomes  $\int_{\delta_w T}^\infty e^{-t^2/2} t dt = e^{-(\delta_w T)^2/2}$ , so we get the approximation

$$\|\mathcal{A}h_w\|_{\mathcal{N}}^2 = \|h_w\|_{\mathcal{N}}^2 + O(\delta_w^{-2} e^{-\delta_w^2 T^2/2 + n \log n}). \quad (33)$$

We have the two following expressions for  $D_{\mathcal{N}}^{(w)}$ .

$$D_{\mathcal{N}}^{(w)}(\theta, \tilde{\theta}) = \det[(e^{-\frac{1}{2}\|\theta w_i - \tilde{\theta} w_j\|^2})_{ij}] = e^{-\frac{\theta^2 + \tilde{\theta}^2}{2}\|w\|^2} \det[(e^{\theta \tilde{\theta} w_i \cdot w_j})_{ij}]. \quad (34)$$

It follows from (34) that  $D_{\mathcal{N}}^{(w)}(\theta, \tilde{\theta})$  can be determined from its values on the diagonal and anti-diagonal  $\tilde{\theta} = \pm\theta$ . More precisely we have (17),

$$D_{\mathcal{N}}^{(w)}(\theta, \tilde{\theta}) = e^{-\frac{\|w\|^2}{2}(|\theta| - |\tilde{\theta}|)^2} D_{\mathcal{N}}^{(w)}(\theta_{\text{g.m.}}, \pm\theta_{\text{g.m.}}),$$

where the geometric mean  $\theta_{\text{g.m.}} = |\theta \tilde{\theta}|^{1/2}$  and  $\pm$  is the sign of  $\theta \tilde{\theta}$ . To show (17) apply the rightmost expression of (34) on both sides and note that the determinants are equal. In particular we have that  $D_{\mathcal{N}}^{(w)}$  decays away from the diagonal and anti-diagonal,

$$|D_{\mathcal{N}}^{(w)}(\theta, \tilde{\theta})| \leq e^{-\frac{\|w\|^2}{2}(|\theta| - |\tilde{\theta}|)^2}. \quad (35)$$

The bound (35) gives concentration around the diagonal everywhere and not only for large  $\theta, \tilde{\theta}$  as Lemma 25. Integrating (35) yields

**Lemma 18.** Let  $h = \tau^{\text{HP}(t)}$  be the high-passed activation function at  $t > 1$ . Then,  $\|\mathcal{A}h_w\|_{\mathcal{N}}^2 \leq \frac{4}{\sqrt{2\pi}\|w\|} \mathfrak{F}\widehat{\tau}(t)$ . So  $\|\mathcal{A}h_w\|_{\mathcal{N}}^2 = O(t^{-K})$  for typical  $w$ .

*Proof.* Write the LHS as a double integral over  $|\theta|, |\tilde{\theta}| \geq t$  as in (13). By (35),

$$\frac{1}{2\pi} \iint_{\tilde{\theta} > \theta > t} D_N^{(w)}(\theta, \tilde{\theta}) |\hat{\tau}(\theta) \hat{\tau}(\tilde{\theta})| d\theta d\tilde{\theta} = \frac{1}{2\pi} \int_0^\infty e^{-\frac{\|w\|^2 s^2}{2}} \int_t^\infty |\hat{\tau}(\theta) \hat{\tau}(s + \theta)| d\theta ds \quad (36)$$

$$\leq \frac{1}{2\pi} \int_0^\infty e^{-\frac{\|w\|^2 s^2}{2}} ds \int_t^\infty |\hat{\tau}(\theta)|^2 d\theta \quad (37)$$

$$= \frac{1}{2\sqrt{2\pi}\|w\|} \int_t^\infty |\hat{\tau}(\theta)|^2 d\theta, \quad (38)$$

where (37) follows from Cauchy-Schwartz. The same bound holds for each of 8 regions in the  $\theta, \tilde{\theta}$ -plane.  $\square$

## E Bound on complex-exponential overlap kernel for the Gaussian envelope

**Lemma 27.** *Let  $v = (v_1, \dots, v_n)^T \in \mathbb{R}^{n \times d}$  and  $w = (w_1, \dots, w_n)^T \in \mathbb{R}^{n \times d}$ . Then  $(e^{v_i \cdot w_j})_{ij} = \sum_{k=0}^\infty Q_k$  where*

$$\text{rank } Q_k \leq \binom{k+d-1}{d-1}, \quad \|Q_k\| \leq \frac{n(\|v\|_\infty \|w\|_\infty d)^k}{k!}. \quad (39)$$

*Proof.* Let  $(c_1, \dots, c_d)$  and  $(\tilde{c}_1, \dots, \tilde{c}_d)$  be the columns of  $v$  and  $w$  and let  $\odot$  denote elementwise operations.

$$(e^{v_i \cdot w_j})_{ij} = e^{\odot \sum_{i=1}^d c_i \tilde{c}_i^T} = \odot_{i=1}^d e^{\odot c_i \tilde{c}_i^T}. \quad (40)$$

We first consider each factor  $e^{\odot c_i \tilde{c}_i^T}$  separately. Elementwise multiplication of rank-one matrices given as outer products corresponds to elementwise multiplication of the vectors,  $ab^T \odot \tilde{a}\tilde{b}^T = (a \odot \tilde{a})(b \odot \tilde{b})^T$ . Therefore, applying the Taylor expansion entrywise,

$$e^{\odot c \tilde{c}^T} = \sum_{k=0}^\infty \frac{(c \tilde{c}^T)^{\odot k}}{k!} = \sum_{k=0}^\infty \frac{(c^{\odot k})(\tilde{c}^{\odot k})^T}{k!}, \quad (41)$$

where  $c = c_i, \tilde{c} = \tilde{c}_i$  are column vectors. Apply (41) to each factor of (40),

$$\odot_{i=1}^d e^{\odot c_i \tilde{c}_i^T} = \sum_{k_1, \dots, k_d=0}^\infty \frac{(\odot_{i=1}^d c_i^{\odot k_i})(\odot_{i=1}^d \tilde{c}_i^{\odot k_i})^T}{\prod_{i=1}^d k_i!}. \quad (42)$$

$$= \sum_{k=0}^\infty \sum_{k_1 + \dots + k_d = k} \frac{1}{k!} \binom{k}{k_1, \dots, k_d} (\odot_{i=1}^d c_i^{\odot k_i})(\odot_{i=1}^d \tilde{c}_i^{\odot k_i})^T \quad (43)$$

Let  $Q_k$  be the innermost sum of (43). We estimate the maximum over the entries,

$$\|Q_k\|_{\max} \leq \frac{\|v\|_\infty^k \|w\|_\infty^k}{k!} \sum_{k_1 + \dots + k_d = k} \binom{k}{k_1, \dots, k_d} = \frac{\|v\|_\infty^k \|w\|_\infty^k d^k}{k!}$$

and apply the inequality  $\|Q_k\| \leq n \|Q_k\|_{\max}$ .  $\square$

**Lemma 28.** *Let  $\lambda_0 \geq \lambda_1 \geq \dots$  be the absolute values of the eigenvalues of  $(e^{v_i \cdot w_j})_{ij}$  and let  $\mu = \|v\|_\infty \|w\|_\infty d$ . Then  $\lambda_0 \leq ne^\mu$ , and for  $\mu \leq 1/2$ ,*

$$\lambda_L \leq \frac{2n}{p!} \mu^p, \quad L = \binom{p+d-1}{d}, \quad (44)$$

where the case  $p = 0$  of (44) holds with the interpretation  $L = \binom{d-1}{d} = 0$ ,  $\lambda_0 \leq ne^{1/2} \leq 2n$ .

*Proof.* From the identity

$$\binom{p+d-1}{d} = 1 + d + \binom{d+1}{d-1} + \dots + \binom{p+d-2}{d-1},$$

there are

$$1 + d + \binom{d+1}{d-1} + \cdots + \binom{p+d-2}{d-1} \geq \text{rank } Q_0 + \cdots + \text{rank } Q_{p-1}$$

eigenvalues in front of  $\lambda_L$  where  $L = \binom{p+d-1}{d}$ , and we have used Lemma 27. By the min-max principle,

$$\lambda_L \leq \left\| \sum_{k=p}^{\infty} Q_k \right\| \leq n \sum_{k=p}^{\infty} \frac{\mu^k}{k!} = \frac{n}{p!} \sum_{k=p}^{\infty} \mu^k = \frac{n}{p!} \frac{\mu^p}{1-\mu} \leq \frac{2n}{p!} \mu^p.$$

□

**Proposition 16.** *Let  $\nu = 2\sqrt{d\|v\|_{\infty}\|w\|_{\infty}}$ . Then,  $\det((e^{v_i \cdot w_j})_{ij}) \leq (\nu/2)^{pn}$  for  $\nu \leq 1$ , where  $p$  is any integer such that  $\binom{p+d-1}{d} \leq n/2$  and  $p! \geq 4n^2$ .*

*Proof.* By Lemma 28 and the assumptions on  $p$  we have  $\lambda_{\lfloor n/2 \rfloor} \leq \frac{2n}{p!} \mu^p \leq \frac{\mu^p}{2n}$  and  $\lambda_0 \leq 2n$  where  $\mu = d\|v\|_{\infty}\|w\|_{\infty}$ , so it follows that  $|\det((e^{v_i \cdot w_j})_{ij})| \leq \lambda_0^{n/2} \lambda_{n/2}^{n/2} \leq (\mu^p)^{n/2} = (\nu/2)^{pn}$ . □

**Lemma 17.** *Let  $\ell = \tau^{\text{LP}(t)}$  be the low-pass at threshold  $t = (2\sqrt{d}\|w\|_{\infty})^{-1}$ . If  $w$  is typical then  $t = \Omega(\sqrt{n/\log n})$  and  $\|\mathcal{A}\ell_w\|_{\mathcal{N}} = O(2^{-\Omega(n^{1+1/d})})$ .*

*Proof.* The lower bound on  $t$  follows directly from its definition and the definition of  $w$  being typical.

Bound  $D_{\mathcal{N}}^{(w)}$  as in (16) and write  $|\widehat{\tau}(\theta)| = O(|\theta|^{-r} + 1)$ . The triangle inequality yields  $\|\tau^{\text{LP}(t)}\|_{\mathcal{N}} = O(2^{-pn/2} \int_{-t}^t (|\theta|/t)^{pn/2} (|\theta|^{-r} + 1) d\theta) = O(2^{-pn/2} t(t^{-r} + 1))$  where  $p = \Omega(n^{1/d})$ . Here we have cancelled the pole  $|\theta|^{-r}$  by writing  $(|\theta|/t)^{pn/2} |\theta|^{-r} \leq (|\theta|/t)^r |\theta|^{-r} = t^{-r}$  so that the integrand is bounded by  $t^{-r} + 1$ . □

## F Generalized definition of roughness

**Definition 29** (Generalized rough activation functions).  $\tau : \mathbb{R} \rightarrow \mathbb{C}$  is generalized rough if its Fourier transform  $\widehat{\tau}$  has tail decay  $K < \infty$ , i.e., if  $\mathfrak{I}\widehat{\tau}(t) = t^{-O(1)}$ , and if additionally,

1.  $|\widehat{\tau}(\theta)|^2 / \mathfrak{I}\widehat{\tau}(\theta) \rightarrow 0$  as  $\theta \rightarrow \infty$ .
2. There exists  $\gamma > 0$  such that for  $1/2 \leq \sigma \leq 2$  and all  $\theta \in \mathbb{R} \setminus [-1, 1]$  where  $\widehat{\tau}(\theta) \neq 0$ ,

$$\text{Re} \left( \frac{\mathbb{E}[\widehat{\tau}(\theta \pm |Z|)]}{\widehat{\tau}(\theta)} \right) \geq \gamma, \quad (45)$$

where  $Z \sim \mathcal{N}(0, \sigma^2)$ . Here, “ $\pm$ ” is taken the sign of  $\theta$ , and  $\text{Re}$  is the real part of a complex number.

The assumptions of Item 1 and Item 2 prevent activation functions with excessive negative correlations between nearby frequencies. The constraint on the standard deviation  $1/2 \leq \sigma \leq 2$  is related to the Xavier / He initialization in Definition 4.

## G Proof of Theorem 6 (lower bound for Gaussian envelope)

**Lemma 30.** *For activation functions satisfying Item 1 of Definition 29 we have that*

$$\|\mathcal{A}\tau_w\|_{\mathcal{N}}^2 \geq \|h_w^{(T)}\|_{\mathcal{N}}^2 - O(e^{-n \log n}) + o(\mathfrak{I}\widehat{\tau}(T)). \quad (46)$$

for  $T = \frac{2}{\delta_v} (\sqrt{n \log n + \log(1/\delta_v)})$  and typical  $w$ .



*Proof.* Let  $\ell^{(T)} = \tau^{\text{LP}(T)}$  and  $h^{(T)} = \tau^{\text{HP}(T)}$ . As discussed in Section 4.5 we have the lower bound

$$\|\mathcal{A}\tau_w\|_{\mathcal{N}}^2 \geq \|\mathcal{A}h_w^{(T)}\|_{\mathcal{N}}^2 - 2|\langle \mathcal{A}\ell_w^{(T)} | \mathcal{A}h_w^{(T)} \rangle_{\mathcal{N}}| \geq \|h_w^{(T)}\|_{\mathcal{N}}^2 - e^{-n \log n} - 2|\langle \mathcal{A}\ell_w^{(T)} | \mathcal{A}h_w^{(T)} \rangle_{\mathcal{N}}|, \quad (47)$$

We further divide the low-frequency part into  $|\theta|$  in  $[0, 1]$ ,  $[1, T/2]$ , and  $[T/2, T]$ . We write  $\alpha^{(t, T)} = h^{(t)} - h^{(T)}$ . Then,

$$|\langle \mathcal{A}\ell_w^{(T)} | \mathcal{A}h_w^{(T)} \rangle_{\mathcal{N}}| \leq \|\mathcal{A}\ell_w^{(1)}\|_{\mathcal{N}} \|\mathcal{A}h_w^{(T)}\|_{\mathcal{N}} + |\langle \mathcal{A}\alpha_w^{(1, T/2)} | \mathcal{A}h_w^{(T)} \rangle_{\mathcal{N}}| + \langle \mathcal{A}\alpha_w^{(T/2, T)} | \mathcal{A}h_w^{(T)} \rangle_{\mathcal{N}}|. \quad (48)$$

By applying the polarization identity to (13) we obtain the overlap between the antisymmetrization with different activation functions, so by the bound (35) on  $D_{\mathcal{N}}^{(w)}(\theta, \tilde{\theta})$ ,

$$|\langle \mathcal{A}\alpha_w^{(T/2, T)} | \mathcal{A}h_w^{(T)} \rangle_{\mathcal{N}}| \leq \frac{(\sup_{|\theta| \geq T/2} |\widehat{\tau}(\theta)|)^2}{2\pi} \int_{T/2 < |\theta| < T} \int_{T < |\tilde{\theta}|} e^{-\frac{\|w\|^2}{2}(|\tilde{\theta}| - |\theta|)^2} d\tilde{\theta} d\theta \quad (49)$$

$$\leq \frac{4(\sup_{|\theta| \geq T/2} |\widehat{\tau}(\theta)|)^2}{2\pi} \int_{T/2 < \theta < T} \int_{T-\theta}^{\infty} e^{-\frac{\|w\|^2}{2}t^2} dt d\theta \quad (50)$$

$$= \frac{4(\sup_{|\theta| \geq T/2} |\widehat{\tau}(\theta)|)^2}{2\pi} \int_0^{\infty} \min\{t, T/2\} e^{-\frac{\|w\|^2}{2}t^2} dt \quad (51)$$

$$\leq \frac{4 \sup_{|\theta| \geq T/2} |\widehat{\tau}(\theta)|^2}{2\pi} \int_0^{\infty} t e^{-\frac{\|w\|^2}{2}t^2} dt \quad (52)$$

$$= \frac{2 \sup_{|\theta| \geq T/2} |\widehat{\tau}(\theta)|^2}{\pi \|w\|^2}, \quad (53)$$

where in (52) the factor 4 comes from the choice of signs of  $\theta, \tilde{\theta}$ , and we have substituted  $t = \tilde{\theta} - \theta$ . We apply Item 1 of Definition 29 to obtain that

$$|\langle \mathcal{A}\alpha_w^{(T/2, T)} | \mathcal{A}h_w^{(T)} \rangle_{\mathcal{N}}| = o(\mathfrak{F}_{\widehat{\tau}}(T)/\|w\|^2). \quad (54)$$

For the first term on the RHS of (48) we have for  $T$  sufficiently large (so that  $|\widehat{\tau}(\theta)\widehat{\tau}(\tilde{\theta})| \leq 2\pi$  for  $|\theta| > 1$  and  $|\tilde{\theta}| > T$ ),

$$\begin{aligned} |\langle \mathcal{A}\alpha_w^{(1, T/2)} | \mathcal{A}h_w^{(T)} \rangle_{\mathcal{N}}| &\leq \int_{1 < |\theta| < T/2} \int_{T < |\tilde{\theta}|} e^{-\frac{\|w\|^2}{2}(|\tilde{\theta}| - |\theta|)^2} d\tilde{\theta} d\theta \\ &\leq \frac{T}{2} \int_{T/2}^{\infty} e^{-\frac{\|w\|^2}{2}t^2} dt \\ &\leq \frac{e^{-\frac{\|w\|^2 T^2}{8}}}{\|w\|^2}. \end{aligned} \quad (55)$$

By (54) and (55),  $|\langle \alpha_w^{(1, T/2)} | h_w^{(T)} \rangle_{\mathcal{N}}| + |\langle \alpha_w^{(T/2, T)} | h_w^{(T)} \rangle_{\mathcal{N}}| = O(e^{-\Omega(T^2)}) + o(\mathfrak{F}_{\widehat{\tau}}(T))$  for typical  $w$ . Finally, for typical  $w$  we have  $t > 1$  in Lemma 17 and  $\|h_w^{(T)}\|_{\mathcal{N}} = O(t^{-K}) = O(1)$  by Lemma 18, so  $\|\ell_w^{(1)}\|_{\mathcal{N}} \|\mathcal{A}h_w^{(T)}\|_{\mathcal{N}} = O(\|\ell_w^{(t)}\|_{\mathcal{N}}) = 2^{-\Omega(n^{1+1/d})}$ . The claim follows by substituting back into (47) and (48).  $\square$

*Proof of Theorem 6.* By Lemma 24 it suffices to show the claim for  $f = \tau_w$  where  $w$  is typical and has typical separation.

By the convolution theorem (32) and the expression  $\mathfrak{F}\rho_v(\theta) = e^{-\|v\|^2\theta^2/2}$  for the Gaussian case we have

$$\|h_w^{(T)}\|_{\mathcal{N}}^2 = \frac{1}{2\pi} \iint_{|\theta|, |\tilde{\theta}| > T} e^{-\|w\|^2(\theta - \tilde{\theta})^2/2} \overline{\widehat{\tau}(\theta)} \widehat{\tau}(\tilde{\theta}) d\theta d\tilde{\theta}. \quad (56)$$

We bound the contribution of the first quadrant to (56) from below. The same argument holds for the third quadrant ( $\theta, \tilde{\theta} < -T$ ). Let  $\sigma = 1/\|w\|$ . Then,

$$\begin{aligned} \iint_{\theta, \tilde{\theta} > T} e^{-\|w\|^2(\theta-\tilde{\theta})^2/2} \overline{\hat{\tau}(\theta)} \hat{\tau}(\tilde{\theta}) d\theta d\tilde{\theta} &= 2 \operatorname{Re} \int_T^\infty \overline{\hat{\tau}(\theta)} \int_\theta^\infty e^{-\|w\|^2(\tilde{\theta}-\theta)^2/2} \hat{\tau}(\tilde{\theta}) d\tilde{\theta} d\theta \\ &= 2 \operatorname{Re} \int_T^\infty \overline{\hat{\tau}(\theta)} \int_\theta^\infty e^{-\frac{(\tilde{\theta}-\theta)^2}{2\sigma^2}} \hat{\tau}(\tilde{\theta}) d\tilde{\theta} d\theta \\ &= \sqrt{2\pi}\sigma \int_T^\infty \operatorname{Re} \left( \overline{\hat{\tau}(\theta)} \mathbb{E}[\hat{\tau}(\theta + |Y|)] \right) d\theta, \end{aligned} \quad (57)$$

where  $Y \sim \mathcal{N}(0, \sigma^2)$ . To obtain a lower bound on the integrand we write

$$\overline{\hat{\tau}(\theta)} \cdot \mathbb{E}[\hat{\tau}(\theta + |Y|)] = \|\hat{\tau}(\theta)\|^2 \cdot \frac{\mathbb{E}[\hat{\tau}(\theta + |Y|)]}{\hat{\tau}(\theta)}. \quad (58)$$

So from Item 2 of Definition 29 we then have a lower bound

$$\operatorname{Re} \left( \overline{\hat{\tau}(\theta)} \cdot \mathbb{E}[\hat{\tau}(\theta + |Y|)] \right) = \|\hat{\tau}(\theta)\|^2 \operatorname{Re} \left( \frac{\mathbb{E}[\hat{\tau}(\theta + |Y|)]}{\hat{\tau}(\theta)} \right) = \Omega(|\hat{\tau}(\theta)|^2), \quad (59)$$

when  $1/2 \leq \sigma \leq 2$ , i.e., when  $1/4 \leq \|w\|^2 \leq 4$ . This holds because  $w$  is typical (and because  $1/2 \leq c_{\text{XH}} \leq 2$ ). For typical  $w$  (56) and (59) then show that

$$\iint_{\theta, \tilde{\theta} > T} e^{-\|w\|^2(\theta-\tilde{\theta})^2/2} \overline{\hat{\tau}(\theta)} \hat{\tau}(\tilde{\theta}) d\theta d\tilde{\theta} = \Omega \left( \int_T^\infty |\hat{\tau}(\theta)|^2 d\theta \right) = \Omega(\mathfrak{T}_{\hat{\tau}}(T)), \quad (60)$$

and the same lower bound holds for the integral over  $\theta, \tilde{\theta} < T$ . Finally we bound the second quadrant  $\theta < -T, \tilde{\theta} > T$  (the fourth quadrant  $\theta > T, \tilde{\theta} < -T$  is analogous) by writing

$$\begin{aligned} \left| \iint_{\theta < -T, \tilde{\theta} > T} e^{-\|w\|^2(\theta-\tilde{\theta})^2/2} \overline{\hat{\tau}(\theta)} \hat{\tau}(\tilde{\theta}) d\theta d\tilde{\theta} \right| &= o \left( \iint_{\theta < -T, \tilde{\theta} > T} e^{-\|w\|^2(\tilde{\theta}-\theta)^2/2} d\theta d\tilde{\theta} \right) \\ &= o \left( \int_{2T}^\infty t e^{-\|w\|^2 t^2/2} dt \right) \\ &= o(e^{-2\|w\|^2 T^2} / \|w\|^2) = o(e^{-T^2}), \end{aligned} \quad (61)$$

where the last expression is for typical  $w$ . The identity (56), the lower bound for the diagonal part (60), and the bound on the magnitude of the off-diagonal part (61) show that  $\|h_w^{(T)}\|_{\mathcal{N}}^2 = \Omega(\mathfrak{T}_{\hat{\tau}}(T)) = \Omega(T^{-K})$ . For typical  $w$  with typical separation we have that  $\delta_w \geq n^{-(1/2+2/d)}/\sqrt{\log n}$  and  $T \sim 2n^{1+2/d} \log n$ , so  $\|h_w^{(T)}\|_{\mathcal{N}}^2 = \tilde{\Omega}(n^{-(1+2/d)})$ .  $\square$

## H Analysis of discretization error in the efficient algorithm

Let  $t = \Omega(1)$  and let  $t = t_0 < t_1 < \dots < t_N = T$  be evenly spaced and for each  $p = 1, \dots, N$  write

$$I_{-p} = [-t_p, -t_{p-1}], \quad I_p = [t_{p-1}, t_p].$$

For  $p = \pm 1, \dots, \pm N$ , let  $\theta_p \in I_p$  and define

$$c_p := \int_{I_p} \hat{\tau}(\theta) d\theta.$$

**Lemma 21.**  $\|S_w - \alpha_w\|_{\mathcal{N}}^2 \leq \frac{2}{\pi} \frac{T}{N} \|\partial_\theta D_{\mathcal{N}}^{(w)}\|_\infty \left( \int_{t \leq |\theta| \leq T} |\hat{\tau}(\theta)| \right)^2 = O\left(\frac{T}{N} \|\partial_\theta D_{\mathcal{N}}^{(w)}\|_\infty\right)$ .

*Proof.* is exactly the antisymmetrization  $\mathcal{A}_{S_w}$  where

$$s(t) = \frac{1}{\sqrt{2\pi}} \sum_{\pm q=1}^N c_q e^{i\theta_q t}.$$

Passing from the truncated activation to its discretization incurs the error

$$\|S_w - \tilde{\alpha}_w\|_{\mathcal{N}}^2 = \|\mathcal{A}(s - \tau_{\text{trc}})_w\|_{\mathcal{N}}^2 = \frac{1}{2\pi} \iint_{\mathbb{R}^2} D_{\mathcal{N}}^{(w)}(\theta, \tilde{\theta}) d\bar{\mu}(\theta) d\mu(\tilde{\theta}), \quad (62)$$

where  $\tau_{\text{trc}} = \tau - \tau^{\text{LP}(1)} - \tau^{\text{HP}(T)}$ , and  $\mu$  is the complex-valued measure

$$\mu(\theta) = \sum_q c_q \delta(\theta - \theta_q) - (\widehat{\tau}(\theta) d\theta).$$

For each square  $S_{pq} = I_p \times I_q$ ,

$$\begin{aligned} & \iint_{S_{pq}} D_{\mathcal{N}}^{(w)}(\theta, \tilde{\theta}) d\bar{\mu}(\theta) d\mu(\tilde{\theta}) \\ &= \bar{c}_p c_q D_{\mathcal{N}}^{(w)}(\theta_p, \theta_q) + \iint_{S_{pq}} D_{\mathcal{N}}^{(w)}(\theta, \tilde{\theta}) \widehat{\tau}(\theta) \overline{\widehat{\tau}(\tilde{\theta})} d\theta d\tilde{\theta} \\ & \quad - \bar{c}_p \int_{I_q} D_{\mathcal{N}}^{(w)}(\theta_p, \tilde{\theta}) \widehat{\tau}(\tilde{\theta}) d\tilde{\theta} - c_q \int_{I_p} D_{\mathcal{N}}^{(w)}(\theta, \theta_q) \overline{\widehat{\tau}(\theta)} d\theta \\ &= \iint_{S_{pq}} \left[ D_{\mathcal{N}}^{(w)}(\theta_p, \theta_q) + D_{\mathcal{N}}^{(w)}(\theta, \tilde{\theta}) - D_{\mathcal{N}}^{(w)}(\theta_p, \tilde{\theta}) - D_{\mathcal{N}}^{(w)}(\theta, \theta_q) \right] \overline{\widehat{\tau}(\theta)} \widehat{\tau}(\tilde{\theta}) d\theta d\tilde{\theta} \\ &\leq \iint_{S_{pq}} \left[ 2 \max D_{\mathcal{N}}^{(w)}(I_p, I_q) - 2 \min D_{\mathcal{N}}^{(w)}(I_p, I_q) \right] |\widehat{\tau}(\theta) \widehat{\tau}(\tilde{\theta})| d\theta d\tilde{\theta} \\ &= 2(\text{diam } D_{\mathcal{N}}^{(w)}(I_p, I_q)) \int_{I_p} |\widehat{\tau}(\theta)| d\theta \int_{I_q} |\widehat{\tau}(\tilde{\theta})| d\tilde{\theta}, \end{aligned} \quad (63)$$

where  $\text{diam}(D_{\mathcal{N}}^{(w)}(I_p, I_q))$  is the diameter of the set  $D(I_p, I_q) = \{D_{\mathcal{N}}^{(w)}(\theta, \tilde{\theta}) \mid (\theta, \tilde{\theta}) \in I_p \times I_q\}$ . Let  $\|\partial D_{\mathcal{N}}^{(w)}\|_{\infty} = \sup_{\theta, \tilde{\theta}} \max\{|\frac{\partial}{\partial \theta} D_{\mathcal{N}}^{(w)}(\theta, \tilde{\theta})|, |\frac{\partial}{\partial \tilde{\theta}} D_{\mathcal{N}}^{(w)}(\theta, \tilde{\theta})|\}$  and let  $t_0 = 1, \dots, t_N = T$  be evenly spaced so that  $|I_p| \leq T/N$ . Then by (63),

$$\iint_{S_{pq}} D_{\mathcal{N}}^{(w)}(\theta, \tilde{\theta}) d\mu(\theta) d\mu(\tilde{\theta}) \leq \frac{4T}{N} \|\partial D_{\mathcal{N}}^{(w)}\|_{\infty} \int_{I_p} |\widehat{\tau}(\theta)| d\theta \int_{I_q} |\widehat{\tau}(\tilde{\theta})| d\tilde{\theta}. \quad (64)$$

Summing over  $p, q$  and applying (62) we get

$$\|S_w - \tilde{\alpha}_w\|_{\mathcal{N}}^2 \leq \frac{2}{\pi} \frac{T}{N} \|\partial D_{\mathcal{N}}^{(w)}\|_{\infty} \left( \int_{t \leq |\theta| \leq T} |\widehat{\tau}(\theta)| \right)^2.$$

□

**Lemma 31.** Let  $B_{ij}^{(v,w)} = \mathfrak{F} \rho(v_i - w_j)$  as in Definition 12. Then,  $|\frac{\partial}{\partial B_{ij}} \det B| \leq 1$  at any  $B = B^{(v,w)}$ .

*Proof.*  $\frac{\partial}{\partial B_{ij}} \det B = (-1)^{i+j} m_{i,j}^B$  where  $m_{i,j}^B$  is the  $i, j$ th minor of  $B$ . But  $m_{i,j}^B = \det B^{(\tilde{v}, \tilde{w})}$  where  $\tilde{v} = (v_{i'})_{i' \neq i}$  and  $\tilde{w} = (w_{j'})_{j' \neq j}$ , and  $|\det B^{(\tilde{v}, \tilde{w})}| = |D_{\mathcal{N}}(\tilde{v}, \tilde{w})| \leq 1$  by the properties mentioned in Appendix C. □

**Corollary 32.**  $|\frac{\partial}{\partial \theta} D_{\mathcal{N}}^{(w)}(\theta, \tilde{\theta})| \leq n^{3/2} \|w\| / \sqrt{e}$ .

*Proof.* By the chain rule,

$$\frac{\partial}{\partial \theta} D_{\mathcal{N}}^{(w)}(\theta, \tilde{\theta}) = \sum_{ij} \frac{\partial B_{ij}(\theta w, \tilde{\theta} w)}{\partial \theta} \frac{\partial \det B}{\partial B_{ij}},$$

where

$$\left| \frac{\partial B_{ij}(\theta w, \tilde{\theta} w)}{\partial \theta} \right| = |w_i \cdot (\nabla \mathfrak{F}_d \rho)(\theta w_i - \tilde{\theta} w_j)| \leq \|w_i\| \cdot \|\nabla \mathfrak{F}_d \rho\|_{\infty}$$

Lemma 31 and the triangle inequality then imply that

$$\left| \frac{\partial}{\partial \theta} D_{\mathcal{N}}^{(w)}(\theta, \tilde{\theta}) \right| \leq n \|\nabla \mathfrak{F}_d \rho\|_{\infty} \sum_i \|w_i\| \leq n^{3/2} \|\nabla \mathfrak{F}_d \rho\|_{\infty} \|w\|,$$

where the last inequality is by Cauchy-Schwarz. For  $\rho = \mathcal{N}(0, I_d)$  we have  $\mathfrak{F}_d \mathcal{N}(w) = e^{-\|w\|^2/2}$  and  $\|\nabla \mathfrak{F}_d \mathcal{N}\|_{\infty} = 1/\sqrt{e}$ .  $\square$