The complexity of blocking (semi)total dominating sets with edge contractions

Esther Galby¹

¹CISPA Helmholtz Center for Information Security, Saarbrücken, Germany

Abstract

We consider the problem of reducing the (semi)total domination number of graph by one by contracting edges. It is known that this can always be done with at most three edge contractions and that deciding whether one edge contraction suffices is an NP-hard problem. We show that for every fixed $k \in \{2, 3\}$, deciding whether exactly k edge contractions are necessary is NP-hard and further provide for k = 2 complete complexity dichotomies on monogenic graph classes.

1 Introduction

A blocker problem asks whether given a graph G, a graph a parameter π , a set \mathcal{O} of one or more graph operations and an integer $k \geq 1$, G can be transformed into a graph H such that $\pi(H) \leq \pi(G) - d$ for some threshold $d \geq 1$, by using at most k operations from \mathcal{O} . These problems are so called because the set of vertices or edges involved can be seen as "blocking" the parameter π . Identifying such sets may provide important information on the structure of the input graph: for instance, if $\pi = \alpha$, k = d = 1 and $\mathcal{O} = \{\text{vertex deletion}\}$, then the problem is equivalent to testing whether the input graph contains a vertex which belongs to every maximum independent set [20]. While the set \mathcal{O} generally consists of a single operation (namely vertex deletion, edge deletion, edge addition or edge contraction), a variety of parameters have been considered in the literature including the chromatic number [2, 5, 6, 19, 21], the stability number [1, 20], the clique number [16, 17], the matching number [3, 25], domination-like parameters [7, 9, 10, 11, 12, 18] and others [4, 13, 14, 22, 24]. In this paper, we focus on two well-known variants of the domination number, namely the total domination number and the semitotal domination number, let \mathcal{O} consists of an edge contraction and set the threshold d to one.

Formally, let G be a graph. The contraction of an edge $xy \in E(G)$ removes vertices x and y from G and replaces them with a new vertex that is made adjacent to precisely those vertices which were adjacent to x or y (without introducing self-loops nor multiple edges). For a parameter π , we denote by $ct_{\pi}(G)$ the minimum number of edge contractions required to modify G into a graph H such that $\pi(H) = \pi(G) - 1$. A set $D \subseteq V(G)$ is a total dominating set of G if every vertex in V(G) has a neighbour in D, and the total domination number $\gamma_t(G)$ of G is the size of a minimum total dominating set of G. A set $D \subseteq V(G)$ is a semitotal dominating set of G if every vertex in $V(G) \setminus D$ has a neighbour in D and every vertex in D is at distance at most two from another vertex in D. The semitotal domination number $\gamma_{t2}(G)$ of G is the size of a minimum semitotal dominating set of G. We are interested in the following problem for $\pi \in \{\gamma_t, \gamma_{t2}\}$.

It is known [10, 12] that, contrary to other parameters such as the chromatic number, the stability number or the clique number¹, ct_{π} is bounded for $\pi \in \{\gamma_t, \gamma_{t2}\}$, namely by three in

CONTRACTION NUMBER (π, k) Instance: A graph G. Question: Is $ct_{\pi}(G) = k$?

¹To see that the minimum number of edge contractions required to decrease the value of the parameter may be arbitrarily large for (1) the chromatic number, consider e.g. stars, (2) the stability number, consider e.g., graphs obtained by identifying one vertex in two otherwise disjoint cliques, (3) the clique number, consider e.g. paths.

both cases. It follows that for $\pi \in \{\gamma_t, \gamma_{t2}\}$ and k > 3, every instance of the above problem is always negative. Similarly, for $\pi \in \{\gamma_t, \gamma_{t2}\}$ and $k \geq 3$, deciding $ct_{\pi}(G) \leq k$ (the so-called *k*-EDGE CONTRACTION(π) problem [9]) is trivial. In contrast, it was shown [10, 11] that for $\pi \in \{\gamma_t, \gamma_{t2}\}$, 1-EDGE CONTRACTION(π) (or, equivalently, CONTRACTION NUMBER($\pi, 1$)) is NPhard; the complexity status for k = 2, 3 remained open. In this paper, we settle these questions and show that for $\pi \in \{\gamma_t, \gamma_{t2}\}$ and k = 2, 3, CONTRACTION NUMBER(π, k) is NP-hard. Thus, combined with [10, 11], the following dichotomy holds.

Theorem 1.1. For $\pi \in \{\gamma_t, \gamma_{t2}\}$, CONTRACTION NUMBER (π, k) is NP-hard if and only if $k \leq 3$.

Let us note that since for $\pi \in \{\gamma_t, \gamma_{t2}\}$, a graph G is a NO-instance for CONTRACTION NUMBER $(\pi, 3)$ if and only if G is a YES-instance for 2-EDGE CONTRACTION (π) , the above implies the following.

Theorem 1.2. For $\pi \in \{\gamma_t, \gamma_{t2}\}$, k-EDGE CONTRACTION (π) is (co)NP-hard if and only if $k \leq 2$.

It is, however, not difficult to find cases which are easy to solve: for instance, in any graph class closed under edge contraction and where π can be efficiently computed, the CONTRACTION NUMBER (π, k) problem can efficiently be solved through a simple brute force approach. Based on this observation, a natural question is whether there are other efficiently solvable instances for which computing π is in fact hard. Motivated by such questions, we consider these problems on monogenic graph classes (that is, graph classes excluding one graph as an induced subgraph) for which the complexity status of TOTAL DOMINATION and SEMITOTAL DOMINATION [8] is known: both problems are polynomial-time solvable on *H*-free graph if $H \subseteq_i P_4 + tK_1$ for some $t \ge 0$, and NP-hard otherwise. This investigation led us to establish complete complexity dichotomies for $\pi \in \{\gamma_t, \gamma_{t2}\}$ and k = 2 on monogenic graph classes, which read as follows.

Theorem 1.3. CONTRACTION NUMBER $(\gamma_t, 2)$ is polynomial-time solvable on *H*-free graphs if $H \subseteq_i P_5 + tK_1$ for some $t \ge 0$ or $H \subseteq_i P_4 + tP_3$ for some $t \ge 0$, and (co)NP-hard otherwise.

Theorem 1.4. CONTRACTION NUMBER $(\gamma_{t2}, 2)$ is polynomial-time solvable on *H*-free graphs if $H \subseteq_i P_5 + tK_1$ for some $t \ge 0$ or $H \subseteq_i P_3 + tP_2$ for some $t \ge 0$, and (co)NP-hard otherwise.

Related work. The study of the blocker problem for $\pi = \gamma_t$ and $\mathcal{O} = \{\text{edge contraction}\}\$ was initiated by Huang and Xu [12] who characterised for every $k \in [3]$, the graphs for which $ct_{\gamma_t}(G) = k$ in terms of the structure of their total dominating sets. More specifically, they proved the following theorem (see Section 2 for missing definitions).

Theorem 1.5 ([12]). For any graph G, the following holds.

- (i) $ct_{\gamma_t}(G) = 1$ if and only if there exists a minimum total dominating set of G that contains a (not necessarily induced) P_3 .
- (ii) $ct_{\gamma_t}(G) = 2$ if and only if no minimum total dominating set of G contains a P_3 and there exists a total dominating set of G of size $\gamma_t(G) + 1$ that contains a (not necessarily induced) P_4 , $K_{1,3}$ or $2P_3$.

On the other hand, the algorithmic study of CONTRACTION NUMBER $(\gamma_t, 1)$ was initiated in [11] where the following dichotomy theorem was established.

Theorem 1.6 ([11]). CONTRACTION NUMBER $(\gamma_t, 1)$ is polynomial-time solvable on *H*-free graphs if $H \subseteq_i P_5 + tK_1$ for some $t \ge 0$ or $H \subseteq_i P_4 + tP_3$ for some $t \ge 0$, and (co)NP-hard otherwise.

Similarly to the total domination parameter, Galby et al. [10] later characterised for every $k \in [3]$ the graphs for which $ct_{\gamma_{t2}}(G) = k$ in terms of the structure of their semitotal dominating sets, as follows (see Section 2 for missing definitions).

Theorem 1.7 ([10]). For any graph G, the following holds.

- (i) $ct_{\gamma_{t2}}(G) = 1$ if and only if there exists a minimum semitotal dominating set of G that contains a friendly triple.
- (ii) $ct_{\gamma_{t2}}(G) = 2$ if and only if no minimum semitotal dominating set of G contains a friendly triple and there exists a semitotal dominating set of G of size $\gamma_{t2}(G) + 1$ that contains a (not necessarily induced) ST-configuration.




Figure 1: The ST-configurations (the dashed lines indicate that the corresponding vertices are at distance two and the serpentine line indicates that the corresponding vertices can be identified).

In the same paper [10], the authors further established the following dichotomy theorem on monogenic graph classes.

Theorem 1.8. CONTRACTION NUMBER $(\gamma_{t2}, 1)$ is polynomial-time solvable for *H*-free graphs if $H \subseteq_i P_5 + tK_1$ for some $t \ge 0$ or $H \subseteq_i P_3 + tP_2$ for some $t \ge 0$, and (co)NP-hard otherwise.

The paper is organised as follows. In Section 2, we present the necessary definitions and notations, as well as some preliminary results. In Section 3, we study the complexity of CONTRACTION NUMBER (γ_t, k) for k = 2, 3, and in Section 4, that of CONTRACTION NUMBER (γ_{t2}, k) for k = 2, 3. The proof of Theorem 1.3 can be found in Section 3.3 and that of Theorem 1.4, in Section 4.3.

2 Preliminaries

2.1 Definitions and notations

Throughout this paper, we only consider finite and simple graphs. Furthermore, since for any parameter π and any $k \geq 0$, a non-connected graph G is a YES-instance for CONTRACTION NUMBER (π, k) if and only if at least one connected component of G is a YES-instance for CONTRACTION NUMBER (π, k) , we restrict our study to connected graphs.

For a graph G, we denote its vertex set by V(G) and its edge set by E(G). For any edge $e = \{x, y\} \in E(G)$, we call x and y the *endvertices* of e. We may also denote the edge e by xy instead of $\{x, y\}$. Given a set $S \subseteq V(G)$, we let G[S] denote the graph *induced* by S, that is, the graph with vertex set S and edge set $\{xy \in E(G) \mid x, y \in S\}$. A graph H is an *induced subgraph* of G, which we denote by $H \subseteq_i G$, if there exists a set $S \subseteq V(G)$ such that H is isomorphic to G[S]. The neighbourhood N(x) of a vertex $x \in V(G)$ is the set $\{y \in V(G) \mid xy \in E(G)\}$ and the closed neighbourhood N[x] of x is the set $N(x) \cup \{x\}$. Similarly, for a set $X \subseteq V(G)$, the neighbourhood N(X) of X is the set $\{y \in V(G) \mid \exists x \in X \text{ s.t. } xy \in E(G)\} \setminus X$ and the closed neighbourhood N[X] of X is the set $N(X) \cup X$. The degree d(x) of a vertex $x \in V(G)$ is the size of its neighbourhood. For any two vertices $x, y \in V(G)$, the distance between x and y is the number of edges in a shortest path from x to y and is denoted d(x, y). Similarly, for any two sets $X, Y \subseteq V(G)$, the distance d(X, Y) from X to Y is the number of edges in a shortest path from X to Y, that is, $d(X,Y) = \min_{x \in X, y \in Y} d(x,y)$. If $X = \{x\}$, we may write d(x,Y) instead of $d({x}, Y)$. For any two sets $X, Y \subseteq V(G), X$ is complete (anticomplete, respectively) to Y, which we denote by X - Y ($X \cdots Y$, respectively) if every vertex in X is adjacent (nonadjacent, respectively) to every vertex in Y.

For $n \ge 1$, we denote by P_n the path on n vertices and by $K_{1,n}$ the complete bipartite graph with partitions of size one and n. If H is a graph and $k \in \mathbb{N}$, we denote by kH the graph consisting of k disjoint copies of H. The paw is the graph depicted in Figure 2a and unless specified otherwise, the vertices of a paw P will be denoted by $P(1), \ldots, P(4)$ (as indicated in Figure 2a) throughout the paper. The long paw is the graph depicted in Figure 2b and unless specified otherwise, the vertices of a long paw P will be denoted by $P(1), \ldots, P(5)$ (as indicated in Figure 2b) throughout the paper.

Let G be a graph with at least two vertices. A set $D \subseteq V(G)$ is dominating if every vertex in $V(G) \setminus D$ has a neighbour in D. A set $D \subseteq V(G)$ is a total dominating set (TD set for short) of G if every vertex in V(G) has a neighbour in D. The total domination number $\gamma_t(G)$ of G is the size of a minimum total dominating set of G. Note that $\gamma_t(G) \ge 2$ by definition. For every $x \in V(G)$ and every neighbour $y \in N(x) \cap D$, we say that y dominates x or that x is dominated by



Figure 2: Some special graphs.

y. A set $D \subseteq V(G)$ is a semitotal dominating set (SD set for short) of G if D is dominating and every vertex in D is at distance at most two from another vertex in D. The semitotal domination number $\gamma_{t2}(G)$ is the size of a minimum semitotal dominating set of G. Note that $\gamma_{t2}(G) \geq 2$ by definition. For every $x \in V(G)$ and every $y \in N[x] \cap D$, we say that y dominates x or that x is dominated by y (note that contrary to a total dominating set, x can dominate itself). If $x, y \in D$ and $d(x, y) \leq 2$ then y is called a witness for x. The set of witnesses of a vertex $x \in D$ is denoted by $w_D(x)$. Note that with this terminology, a semitotal dominating set D is equivalently a dominating set where every vertex in D has a witness. For every $x \in D$, we denote by $PN_D(x) = \{y \in V(G) \mid N(y) \cap D = \{x\}\}$ the private neighbourhood of x with respect to D. A friendly triple is a subset of three vertices x, y and z such that $xy \in E(G)$ and $d_G(y, z) \leq 2$.

Finally, let us define the POSITIVE EXACTLY 3-BOUNDED 1-IN-3 3-SAT problem which will be used in several of our hardness reductions: it is an NP-hard [15] variant of the 3-SAT problem where given a formula Φ in which all literals are positive, every clause contains exactly three literals and every variable appears in exactly three clauses, the problem is to determine whether there exists a truth assignment for Φ such that each clause contains exactly one true literal.

2.2 Preliminary results

In this section, we present some useful technical results.

Lemma 2.1. Let G be a graph such that $\gamma_t(G) \geq 3$. Then for any minimum TD set D of G, if there exist $x, y, z \in D$ such that $xy \in E(G)$ and $d(z, \{x, y\}) \leq 2$, then $ct_{\gamma_t}(G) \leq 2$.

Proof. Suppose that there exist three such vertices $x, y, z \in D$. If $d(z, \{x, y\}) = 1$, say z is adjacent to y without loss of generality, then D contains the P_3 xyz and thus, $ct_{\gamma_t}(G) = 1$ by Theorem 1.5. Suppose next that $d(z, \{x, y\}) = 2$, say $d(z, \{x, y\}) = d(z, y)$ without loss of generality, and let t be a common neighbour of y and z. If t belongs to D as well, then D contains the P_3 ytz and we conclude as previously. Otherwise, $t \notin D$ in which case $D \cup \{t\}$ contains the P_4 xytz and thus, $ct_{\gamma_t}(G) \leq 2$ by Theorem 1.5.

Lemma 2.2. Let G be a graph such that $\gamma_{t2}(G) \geq 3$. If G has a minimum SD set D such that one of the following holds:

(i) D contains an edge, or

(ii) there exists $x \in D$ such that $|w_D(x)| \ge 2$, then $ct_{\gamma_{t2}}(G) \le 2$.

Proof. Assume first that G has a minimum SD set D containing an edge uv. Let $w \in D \setminus \{u, v\}$ be a closest vertex from the edge uv, that is, $d(w, \{u, v\}) = \min_{x \in D \setminus \{u, v\}} d(x, \{u, v\})$. Then $d(w, \{u, v\}) \leq 3$: indeed, if $d(w, \{u, v\}) > 3$ then the vertex at distance two from $\{u, v\}$ on a shortest path from $\{u, v\}$ to w is not dominated, by the choice of w. Now if $d(w, \{u, v\}) \leq 2$ then u, v, w is a friendly triple of D and thus, $ct_{\gamma_{t2}}(G) = 1$ by Theorem 1.7. Otherwise, $d(w, \{u, v\}) = 3$ in which case, denoting by x the vertex at distance one from $\{u, v\}$ on a shortest path from $\{u, v\}$ to $w, D \cup \{x\}$ contains the O_4 u, v, x, w and thus, $ct_{\gamma_{t2}}(G) \leq 2$ by Theorem 1.7 which proves item (i).

To prove item (ii), assume that G has a minimum SD set D such that there exists $x \in D$ where $|w_D(x)| \ge 2$, say $y, z \in w_D(x)$. If $d(x, \{y, z\}) = 1$ then we conclude as previously that $ct_{\gamma_{t2}}(G) \le 2$. Otherwise, d(x, y) = d(x, z) = 2 in which case, denoting by t a common neighbour of x and $y, D \cup \{t\}$ contains the O_4 y, t, x, z and thus, $ct_{\gamma_{t2}}(G) \le 2$ by Theorem 1.7. **Lemma 2.3.** Let G be a graph such that $ct_{\gamma_{t2}}(G) \leq 2$ and for every minimum SD set D of G the following hold.

(i) D contains no edge.

(ii) For every $x \in D$, $|w_D(x)| = 1$.

Then every SD set D of G of size $\gamma_{t2}(G) + 1$ containing an ST-configuration is either a minimal SD set of G or contains an O_6 .

Proof. Let D be an SD set of size $\gamma_{t2}(G) + 1$ containing an ST-configuration and suppose that D is not minimal. Then there exists $x \in D$ such that $D_x = D \setminus \{x\}$ is a minimum SD set of G; in particular, D_x contains no edge by item (i). It follows that D contains no O_1, O_2, O_3 and O_7 : indeed, since these ST-configurations contain two vertex-disjoint edges, if D contains one of them, then D_x must contain at least one edge. Now if D contains an O_4 on vertices u, v, w, t where $uv, vw \in E(G)$ and d(w, t) = 2, necessarily x = v (D_x would otherwise contain an edge); but then, $u, t \in w_{D_x}(w)$, a contradiction to item (ii). Similarly, if D contains an O_5 on vertices u, v, w, t where t is the vertex of degree three, necessarily x = t (D_x would otherwise contain an edge); but then, $u, v \in w_{D_x}(w)$, a contradiction to item (ii). Thus, D contains an O_6 .

Let us finally remark the following.

Observation 1. Let $\pi \in {\gamma_t, \gamma_{t2}}$ and let \mathcal{G} be a graph class where CONTRACTION NUMBER $(\pi,3)$ is polynomial-time solvable. Then CONTRACTION NUMBER $(\pi,1)$ is NP-hard on \mathcal{G} if and only if CONTRACTION NUMBER $(\pi,2)$ is coNP-hard on \mathcal{G} .

3 Total Domination

In this section, we consider the CONTRACTION NUMBER (γ_t, k) problem for $k \in \{2, 3\}$. In Section 3.1, we cover polynomial-time solvable cases and in Section 3.2, we examine hard cases. The proof of Theorem 1.3 can be found in Section 3.3.

3.1 Polynomial-time solvable cases

The algorithms for CONTRACTION NUMBER $(\gamma_t, 2)$ and CONTRACTION NUMBER $(\gamma_t, 3)$ outlined thereafter will rely on the following key result.

Lemma 3.1. For every $k \ge 0$, 2-EDGE CONTRACTION (γ_t) is polynomial-time solvable on $(P_6 + kP_3)$ -free graphs.

Proof. We prove the result by induction on k.

Base Case. k = 0. Let G be a P_6 -free graph such that $\gamma_t(G) \geq 3$. We show that $ct_{\gamma_t}(G) \leq 2$, that is, G is a YES-instance for 2-EDGE CONTRACTION (γ_t) . To this end, let D be a minimum TD set of G. Consider two adjacent vertices $u, v \in D$ and let $w \in D \setminus \{u, v\}$ be a closest vertex from $\{u, v\}$, that is, $d(w, \{u, v\}) = \min_{x \in D \setminus \{u, v\}} d(x, \{u, v\})$. Then $d(w, \{u, v\}) \leq 3$: indeed, if $d(w, \{u, v\}) > 3$ then the vertex at distance two from $\{u, v\}$ on a shortest path from $\{u, v\}$ to w is not dominated. Now if $d(w, \{u, v\}) \leq 2$ then $ct_{\gamma_t}(G) \leq 2$ by Lemma 2.1; thus, we assume henceforth that $d(w, \{u, v\}) = 3$, say $d(w, \{u, v\}) = d(w, u)$ without loss of generality.

Let P = uxyw be a shortest path from u to w and let $t \in D$ be a neighbour of w. Suppose first that t is nonadjacent to y (note that t is nonadjacent to x as $d(w, \{u, v\}) = \min_{b \in D \setminus \{u, v\}} d(b, \{u, v\})$ = 3). Then any neighbour a of t must be adjacent to w, y or x, for atwyxv would otherwise induce a P_6 (note indeed that a is nonadjacent to v as $d(w, \{u, v\}) = \min_{b \in D \setminus \{u, v\}} d(b, \{u, v\}) = 3$). But then, $(D \setminus \{t\}) \cup \{x, y\}$ is a TD set of G of size $\gamma_t(G) + 1$ containing the P_4 vxyw and thus, $ct_{\gamma_t}(G) \leq 2$ by Theorem 1.5. Second, suppose that t is adjacent to y. If every neighbour of vis adjacent to x or y, then $(D \setminus \{v\}) \cup \{x, y\}$ is a TD set of G of size $\gamma_t(G) + 1$ containing the P_4 xywt and so, $ct_{\gamma_t}(G) \leq 2$ by Theorem 1.5. Thus, assume that v has a neighbour z which is nonadjacent to both x and y. If z = u then, by symmetry, we conclude as in the previous case. Suppose therefore that $z \neq u$. Then every neighbour a of w is adjacent to z, x or y, for awyxvzwould otherwise induce a P_6 ; and we conclude symmetrically that every neighbour of t is adjacent to z, x or y. It then follows that $(D \setminus \{w,t\}) \cup \{x, y, z\}$ is a TD set of G of size $\gamma_t(G) + 1$ containing the P_4 yxvz and thus, $ct_{\gamma_t}(G) \leq 2$ by Theorem 1.5.

Inductive step. Let G be a $(P_6 + kP_3)$ -free graph. We aim to show that if G contains an induced $P_6 + (k-1)P_3$ and G is a NO-instance for 2-EDGE CONTRACTION (γ_t) then $\gamma_t(G)$ is bounded by some function f of k (and k only). Assuming for now that this claim is correct, the following algorithm solves the 2-EDGE CONTRACTION (γ_t) problem on $(P_6 + kP_3)$ -free graphs.

- 1. If G contains no induced $P_6 + (k-1)P_3$ then use the algorithm for $(P_6 + (k-1)P_3)$ -free graphs to determine whether G is a YES-instance for 2-EDGE CONTRACTION (γ_t) or not.
- 2. Check whether there exists a TD set of G of size at most f(k).
 - 2.1 If the answer is no then output YES.
 - 2.2 Check whether there exists a minimum TD set of G containing a P_3 , or a TD set of G of size $\gamma_t(G) + 1$ containing a $P_4, K_{1,3}$ or $2P_3$ (see Theorem 1.5) and output the answer accordingly.

Now observe that checking whether G is $(P_6 + (k-1)P_3)$ -free can be done in time $|V(G)|^{O(k)}$ and that step 2 can be done in time $|V(G)|^{O(f(k))}$ by simple brute force. Since 2-EDGE CONTRAC-TION (γ_t) is polynomial-time solvable on $(P_6 + (k-1)P_3)$ -free graphs by the induction hypothesis, the above algorithm indeed runs in polynomial time (for fixed k). The remainder of this proof is devoted to showing that $f(k) = k^4 + 4k^2 + 21k + 19$.

Assume henceforth that G contains an induced $P_6 + (k-1)P_3$. Let $A \subseteq V(G)$ be a set of vertices such that G[A] is isomorphic to $P_6 + (k-1)P_3$, let $B \subseteq V(G) \setminus A$ be the set of vertices at distance one from A and let $C = V(G) \setminus (A \cup B)$. Note that since G is $(P_6 + kP_3)$ -free, Cis a disjoint union of cliques; we denote by \mathcal{K} the set of maximal cliques in C. Now let D be a minimum TD set of G such that $|D \cap B|$ is maximum amongst all minimum TD set of G. Denote by $\mathcal{K}_D \subseteq \mathcal{K}$ the set of cliques $K \in \mathcal{K}$ such that $D \cap (N(K) \cap B) \neq \emptyset$, and set $\overline{\mathcal{K}_D} = \mathcal{K} \setminus \mathcal{K}_D$. We aim to upperbound $|D \cap N[\mathcal{K}_D]|$ and $|D \cap N[\overline{\mathcal{K}_D}]|$ when G is a No-instance for 2-EDGE CONTRACTION (γ_t) . To this end, we first prove the following.

Claim 1. If G is a NO-instance for 2-EDGE CONTRACTION (γ_t) then for every $K \in \mathcal{K}$, exactly one of the following holds.

(i) $D \cap V(K) \neq \emptyset$ and $D \cap N[K] = \{x, y\}$ for some edge $xy \in E(G)$. (ii) $N[K] \cap D \subseteq B$.

Proof. Assume that G is a NO-instance for 2-EDGE CONTRACTION(γ_t) and consider a clique $K \in \mathcal{K}$. Then $D \cap N[K] \neq \emptyset$ as the vertices of K must be dominated. Now if $D \cap V(K) \neq \emptyset$, say $x \in D \cap V(K)$, then x has a neighbour $y \in D$ as x must be dominated; but then, $D \cap N[K] = \{x, y\}$ for otherwise $ct_{\gamma_t}(G) \leq 2$ by Lemma 2.1 (note indeed that every vertex in N[K] is within distance at most two from x). Otherwise, $D \cap V(K) = \emptyset$ in which case $D \cap (N(K) \cap B) \neq \emptyset$.

Claim 2. If G is a NO-instance for 2-EDGE CONTRACTION (γ_t) then $|D \cap N[\mathcal{K}_D]| \leq 4|A|$.

Proof. Assume that G is a No-instance for 2-EDGE CONTRACTION(γ_t) and suppose for a contradiction that $|D \cap N[\mathcal{K}_D]| > 4|A|$. We contend that at least half of those vertices belong to B, that is, $|D \cap (N(\mathcal{K}_D) \cap B)| > 2|A|$. Indeed, for every $x \in D \cap (N(\mathcal{K}_D) \cap B)$, denote by \mathcal{K}_D^x the set of cliques $K \in \mathcal{K}_D$ such that $x \in N(K)$. Let us first show that for every $x \in D \cap (N(\mathcal{K}_D) \cap B)$, there exists at most one clique $K \in \mathcal{K}_D^x$ such that $D \cap V(K) \neq \emptyset$, and that furthermore, if such a clique exists then it in fact contains only one element of D. Consider $x \in D \cap (N(\mathcal{K}_D) \cap B)$. If $|\mathcal{K}_D^x| = 1$ then the result readily follows from Claim 1. Suppose therefore that $|\mathcal{K}_D^x| \ge 2$. Then for all but at most one of the cliques in \mathcal{K}_D^x , item (ii) of Claim 1 must hold: indeed, if there exist $K_1, K_2 \in \mathcal{K}_D^x$ such that K_1 and K_2 both satisfy item (i) of Claim 1, then D contains the $P_3 y_1 x y_2$ where $y_1 \in V(K_1)$ and $y_2 \in V(K_2)$, a contradiction to Theorem 1.5.

Now observe that for any $x, y \in D \cap (N(\mathcal{K}_D) \cap B)$, if there exist $K_x \in \mathcal{K}_D^x$ and $K_y \in \mathcal{K}_D^y$ such that $D \cap V(K_x) \neq \emptyset$ and $D \cap V(K_y) \neq \emptyset$, then surely $K_x \neq K_y$ as otherwise $|D \cap N[K_x]| \geq 3$, a contradiction to Claim 1(i). Since for every $K \in \mathcal{K}_D$, there exists $x \in D \cap (N(\mathcal{K}_D) \cap B)$ such that

 $K \in \mathcal{K}_D^x$, by definition of \mathcal{K}_D , it follows that

$$\begin{split} |D \cap (N(\mathcal{K}_D) \cap B)| &\geq \sum_{x \in D \cap (N(\mathcal{K}_D) \cap B)} |D \cap \bigcup_{K \in \mathcal{K}_D^x} V(K)| \\ &\geq |D \cap \bigcup_{x \in D \cap (N(\mathcal{K}_D) \cap B)} \bigcup_{K \in \mathcal{K}_D^x} V(K)| \\ &\geq |D \cap \bigcup_{K \in \mathcal{K}_D} V(K)| \end{split}$$

But

$$|D \cap N[\mathcal{K}_D]| = |D \cap (N(\mathcal{K}_D) \cap B)| + |D \cap \bigcup_{K \in \mathcal{K}_D} V(K)|$$

and so, we conclude that

$$2|D \cap (N(\mathcal{K}_D) \cap B)| \ge |D \cap N[\mathcal{K}_D]| > 4|A|$$

as claimed. It follows that there must exist at least three vertices $x, y, z \in D \cap (N(\mathcal{K}_D) \cap B)$ such that x, y and z have a common neighbour in A: indeed, if no such three vertices exist then every vertex in A has at most two neighbours in $D \cap (N(\mathcal{K}_D) \cap B)$ and so, $|D \cap (N(\mathcal{K}_D) \cap B)| \leq 2|A|$, a contradiction to the above. This implies, in particular, that for every $u, v \in \{x, y, z\}, d(u, v) \leq 2$; but x must have a neighbour in D (possibly y or z) and so, $ct_{\gamma_t}(G) \leq 2$ by Lemma 2.1, a contradiction to our assumption. Therefore, $|D \cap N[\mathcal{K}_D]| \leq 4|A|$.

Now note that since for every $K \in \overline{\mathcal{K}_D}$, $D \cap (N(K) \cap B) = \emptyset$ by definition, the following ensues from Claim 1.

Observation 2. If G is a NO-instance for 2-EDGE CONTRACTION (γ_t) then for every $K \in \overline{\mathcal{K}_D}$, $|D \cap N[K]| = |D \cap V(K)| = 2$.

For every $K \in \overline{\mathcal{K}_D}$, let us denote by $x_K y_K \in E(G)$ the edge contained in $V(K) \cap D$ when G is a NO-instance for 2-EDGE CONTRACTION (γ_t) . Observe that by Lemma 2.1, the following holds.

Observation 3. If G is a NO-instance for 2-EDGE CONTRACTION (γ_t) then for every clique $K \in \overline{\mathcal{K}_D}, d(\{x_K, y_K\}, D \setminus \{x_K, y_K\}) > 2.$

Claim 3. If G is a NO-instance for 2-EDGE CONTRACTION (γ_t) then for every $K \in \overline{\mathcal{K}_D}$, the following hold.

(i) For every $u \in \{x_K, y_K\}$, $PN_D(u) \neq \emptyset$.

(ii) For $u \in \{x_K, y_K\}$ and $v \in \{x_K, y_K\} \setminus \{u\}$, no vertex of $PN_D(u)$ is complete to $PN_D(v)$.

Proof. Assume that *G* is a No-instance for 2-EDGE CONTRACTION(γ_t) and consider a clique $K \in \overline{\mathcal{K}_D}$. Observe first the $PN_D(x_K) \cup PN_D(y_K) \subseteq B$ as *K* is a clique. Furthermore, we may assume that $(N(x_K) \cup N(y_K)) \cap B \neq \emptyset$ for it otherwise suffices to consider, in place of *D*, the TD set $(D \setminus \{x_K\}) \cup \{x\}$, where $x \in V(K)$ has at least one neighbour in *B*. Now suppose that $PN_D(u) = \emptyset$ for some $u \in \{x_K, y_K\}$, say $PN_D(x_K) = \emptyset$ without loss of generality. Then $N(y_K) \cap B \neq \emptyset$: if not, then $N(x_K) \cap B \neq \emptyset$ by the above, and since no vertex in $N(x_K) \cap B$ can be adjacent to another vertex in $D \setminus \{x_K, y_K\}$ by Observation 3, necessarily $N(x_K) \cap B \subseteq PN_D(x_K)$, a contradiction to our assumption. But then, letting $y \in B$ be a neighbour of y_K , the set $(D \setminus \{x_K\}) \cup \{y\}$ is a minimum TD set of *G* containing strictly more vertices from *B* than *D*, a contradiction to the choice of *D*. Thus, $PN_D(u) \neq \emptyset$ for every $u \in \{x_K, y_K\}$. Now if for some $u \in \{x_K, y_K\}$, there exists $y \in PN_D(u)$ such that y is complete to $PN_D(v)$ where $v \in \{x_K, y_K\} \setminus \{u\}$, then the set $(D \setminus \{v\}) \cup \{y\}$ is a minimum TD set of *G* containing strictly more vertices from *B* than *D*, a contradiction to the choice of *D*.

Claim 4. If G is a NO-instance for 2-EDGE CONTRACTION (γ_t) and there exists a set $S \subseteq \overline{\mathcal{K}_D}$ such that for every $K, K' \in S$, $PN_D(x_K) \cup PN_D(y_K) \cdots PN_D(x_{K'}) \cup PN_D(y_{K'})$ then $|S| \leq |A|$.

Proof. Assume that G is a No-instance for 2-EDGE CONTRACTION(γ_t) and there exists such a set $S \subseteq \overline{\mathcal{K}_D}$. Suppose for a contradiction that |S| > |A|. Then there must exist $u \in PN_D(x_K) \cup PN_D(y_K)$ and $v \in PN_D(x_{K'}) \cup PN_D(y_{K'})$ for two distinct cliques $K, K' \in S$, such that u and v have a common neighbour $w \in A$. Then for all but at most k - 1 cliques $K'' \in S$, w is complete to $PN_D(x_{K''}) \cup PN_D(y_{K''})$: indeed, if there exist at least k cliques $K_1, \ldots, K_k \in S$ such that for every $i \in [k]$, w is nonadjacent to a vertex $u_i \in PN_D(x_{K_i}) \cup PN_D(y_{K_i})$, then $\{x_K, y_K, u, w, v, x_{K'}, y_{K'}\} \cup \bigcup_{i \in [k]} \{u_i, x_{K_i}, y_{K_i}\}$ induces a $P_7 + kP_3$, a contradiction. Since |S| > |A| = 6 + 3(k - 1), it follows that there are at least two cliques $K_1, K_2 \in S \setminus \{K, K'\}$ such that w is complete to $PN_D(x_{K_1}) \cup PN_D(y_{K_1})$ and to $PN_D(x_{K_2}) \cup PN_D(y_{K_2})$. However, by Claim 3(i), $PN_D(y_{K_1}) \neq \emptyset$ and $PN_D(y_{K_2}) \neq \emptyset$ and so, the set $(D \setminus \{x_{K_1}, x_{K_2}\}) \cup \{w, u_1, u_2\}$ where $u_1 \in PN_D(y_{K_1})$ and $u_2 \in PN_D(y_{K_2})$, is a TD set of G of size $\gamma_t(G) + 1$ containing the P_4 $y_{K_1}u_1wu_2$, a contradiction to Theorem 1.5.

Claim 5. Let $K, K' \in \overline{\mathcal{K}_D}$. If there exist $u \in PN_D(x_K) \cup PN_D(y_K)$ and $v \in PN_D(x_{K'}) \cup PN_D(y_{K'})$ such that $uv \in E(G)$, then there exist a set $S_{uv} \subseteq \overline{\mathcal{K}_D} \setminus \{K, K'\}$ and a set $T_{uv} \subseteq \bigcup_{L \in S_{uv}} PN_D(x_L) \cup PN_D(y_L)$ such that the following hold. (i) For every $L \in S_{uv}$, every vertex in $PN_D(x_L) \cup PN_D(y_L)$ is adjacent to at least one vertex

(i) For every $L \in S_{uv}$, every vertex in $PN_D(x_L) \cup PN_D(y_L)$ is adjacent to at least one vertex of $\underline{T}_{uv} \cup \{u, v\}$.

(*ii*)
$$|\overline{\mathcal{K}_D} \setminus S_{uv}| \le 2 + k(k-1)/2.$$

Proof. Assume that there exist $u \in PN_D(x_K) \cup PN_D(y_K)$ and $v \in PN_D(x_{K'}) \cup PN_D(y_{K'})$ such that $uv \in E(G)$. Let $T \subseteq \bigcup_{L \in \overline{\mathcal{K}_D} \setminus \{K, K'\}} PN_D(x_L) \cup PN_D(y_L)$ be a maximum size independent set such that $T \cdots \{u, v\}$ and for every $L \in \overline{\mathcal{K}_D}$, $|T \cap (PN_D(x_L) \cup PN_D(y_L))| \leq 1$. Further denote by $S = \{L \in \overline{\mathcal{K}_D} \mid T \cap (PN_D(x_L) \cup PN_D(y_L)) \neq \emptyset\} \cup \{K, K'\}$. Observe that since $\{u, v\} \cup T \cup \bigcup_{L \in S} \{x_L, y_L\}$ induces $P_6 + |T|P_3$, necessarily $|T| \leq k - 1$ and thus, $|S| \leq k + 1$. Now consider the sequence of sets constructed according to the following procedure.

- 1. Initialise $i = 1, T_1 = T$ and $R_1 = S_1 = S$.
- 2. Increase i by one.
- 3. Let $T_i \subseteq \bigcup_{L \in \overline{\mathcal{K}_D} \setminus S_{i-1}} PN_D(x_L) \cup PN_D(y_L)$ be a maximum size independent set such that $T_i \cdots \{u, v\}$ and for every $L \in \overline{\mathcal{K}_D} \setminus S_{i-1}, |T_i \cap (PN_D(x_L) \cup PN_D(y_L))| \leq 1$. Set $R_i = \{L \in \overline{\mathcal{K}_D} \mid T_i \cap (PN_D(x_L) \cup PN_D(y_L)) \neq \emptyset\}$ and $S_i = S_{i-1} \cup R_i$.
- 4. If $|T_i| = |T_{i-1}|$ then stop the procedure.
- 5. Return to step 2.

Consider the value of i at the end of the procedure (note that $i \ge 2$). Let us show that we may take

$$T_{uv} = T_{i-1} \cup T_i$$

and

$$S_{uv} = \begin{cases} \overline{\mathcal{K}_D} \setminus S_{i-2} & \text{if } i > 2\\ \overline{\mathcal{K}_D} \setminus \{K, K'\} & \text{otherwise.} \end{cases}$$

Observe first that by construction, for every $L \in \overline{\mathcal{K}_D} \setminus S_i$ and every vertex $x \in PN_D(x_L) \cup PN_D(y_L)$, x is adjacent to at least one vertex in $T_i \cup \{u, v\}$ for otherwise, the procedure would have output $T_i \cup \{x\}$ in place of T_i . Similarly, for every $L \in R_i$ and every vertex $x \in PN_D(x_L) \cup PN_D(y_L)$, x is adjacent to at least one vertex in $T_{i-1} \cup \{u, v\}$ for otherwise, the procedure would have output $T_{i-1} \cup \{x\}$ in place of T_{i-1} ; and for $L \in R_{i-1}$ and every vertex $x \in PN_D(x_L) \cup PN_D(y_L)$, x is adjacent to at least one vertex in $T_i \cup \{u, v\}$ for otherwise, the procedure would have output $T_i \cup \{x\}$ in place of T_{i-1} ; and for $L \in R_{i-1}$ and every vertex $x \in PN_D(x_L) \cup PN_D(y_L)$, x is adjacent to at least one vertex in $T_i \cup \{u, v\}$ for otherwise, the procedure would have output $T_i \cup \{x\}$ in place of T_{i-1} (recall that by construction $|T_{i-1}| = |T_i|$). In particular, every vertex in T_i is adjacent to at least one vertex in T_{i-1} and thus, item (i) holds true. Now for every $1 \le p < q \le i-1$, $|T_p| < |T_p|$ by construction; and since $|T_1| \le k-1$, it follows that $i \le k+1$ and for every $1 \le p \le i-1$, $|T_p| \le k-p$. Thus, if i > 2 then

$$|S_{i-2}| = 2 + \sum_{1 \le p \le i-2} |T_p| \le 2 + \sum_{1 \le p \le i-2} k - p \le 2 + \frac{k(k-1)}{2}$$

and so, item (ii) holds true.

In the following, for any $K, K' \in \overline{\mathcal{K}_D}$, $u \in PN_D(x_K) \cup PN_D(y_K)$ and $v \in PN_D(x_{K'}) \cup PN_D(y_{K'})$ such that $uv \in E(G)$, we denote by $S_{uv} \subseteq \overline{\mathcal{K}_D}$ and $T_{uv} \subseteq B$ the two sets given by Claim 5.

Claim 6. If there exist $K_1, K'_1, K_2, K'_2 \in \overline{\mathcal{K}_D}$ such that

- (i) $K_2, K'_2 \in S_{u_1u'_1}$ for some $u_1 \in PN_D(x_{K_1}) \cup PN_D(y_{K_1})$ and $u'_1 \in PN_D(x_{K'_1}) \cup PN_D(y_{K'_1})$ where $u_1u'_1 \in E(G)$, and
- (ii) one of K_1 and K'_1 belongs to $S_{u_2u'_2}$ for some $u_2 \in PN_D(x_{K_2}) \cup PN_D(y_{K_2})$ and $u'_2 \in PN_D(x_{K'_2}) \cup PN_D(y_{K'_2})$ where $u_2u'_2 \in E(G)$,

then $ct_{\gamma_t}(G) \leq 2$.

Proof. Assume that four such cliques $K_1, K'_1, K_2, K'_2 \in \overline{\mathcal{K}_D}$ exist and let u_1, u'_1, u_2, u'_2 be the vertices given by items (i) and (ii). Then for every $z \in T_{u_1u'_1}$, there exists, by construction, a clique $L_z \in S_{u_1u'_1}$ such that z is a private neighbour of x_{L_z} or y_{L_z} : let us denote by $w_z \in \{x_{L_z}, y_{L_z}\}$ the vertex such that z is not a private neighbour of w_z . Similarly, for every $z \in T_{u_2u'_2}$, there exists a clique $L_z \in S_{u_2u'_2}$ such that z is not a private neighbour of w_z . Now assume without loss of generality that for every $i \in [2], u_i \in PN_D(y_{K_i})$ and $u'_i \in PN_D(y_{K'_i})$, and that furthermore, $K'_1 \in S_{u_2u'_2}$. We contend that the set $D' = (D \setminus (\{w_z \mid z \in T_{u_1u'_1} \cup T_{u_2u'_2}\} \cup \{x_{K'_1}, x_{K_2}, x_{K'_2}\}) \cup T_{u_1u'_1} \cup T_{u_2u'_2} \cup \{u_1, u'_1, u_2, u'_2\}$ is a TD set of G. Indeed, by Claim 5(i), every vertex in $PN_D(w_z)$ for $z \in T_{u_1u'_1}$, is adjacent to at least one vertex in $T_{u_1u'_1} \cup \{u_1, u'_1\}$; and similarly, every vertex in $PN_D(w_z)$ for $z \in T_{u_2u'_2}$, here $K_2, K'_2 \in S_{u_1u'_1}$ and $K'_1 \in S_{u_2u'_2}$, every vertex in $PN_D(x_{K_2}) \cup PN_D(x_{K'_2})$ is adjacent to at least one vertex in $T_{u_2u'_2} \cup \{u_2, u'_2\}$. Furthermore, by Claim 5(i), since $K_2, K'_2 \in S_{u_1u'_1}$ and $K'_1 \in S_{u_2u'_2}$, every vertex in $PN_D(x_{K'_1})$ is adjacent to at least one vertex in $T_{u_2u'_2} \cup \{u_2, u'_2\}$. It follows that D' is indeed a TD set of G; and since $|D'| = \gamma_t(G) + 1$ and D' contains the $P_4 y_{K_1}u_1u'_1y'_{K'_1}$, we conclude by Theorem 1.5 that $ct_{\gamma_t}(G) \leq 2$.

Claim 7. If G is a NO-instance for 2-EDGE CONTRACTION (γ_t) then $|\overline{\mathcal{K}_D}| \leq k^2(k(k-1)/2 + 2) + |A| - 1$.

Proof. Assume that G is a NO-instance for 2-EDGE CONTRACTION(γ_t) and suppose to the contrary that $|\overline{\mathcal{K}_D}| > k^2(k(k-1)/2+2) + |A| - 1$. Let us show that there then exist four cliques in $\overline{\mathcal{K}_D}$ satisfying items (i) and (ii) of Claim 6, which would contradict the fact that $ct_{\gamma_t}(G) > 2$. Let $K_1, K'_1 \in \overline{\mathcal{K}_D}$ be two cliques for which there exist $u_1 \in PN_D(x_{K_1}) \cup PN_D(y_{K_1})$ and $u'_1 \in PN_D(x_{K'_1}) \cup PN_D(y_{K'_1})$ such that $u_1u'_1 \in E(G)$ (the existence of such cliques is guaranteed by Claim 4). We claim that the algorithm below always outputs four cliques of $\overline{\mathcal{K}_D}$ satisfying items (i) and (ii) of Claim 6.

- 1. Initialise i = 1 and $C_1 = S_{u_1u'_1}$.
- 2. Increase i by one.
- 3. If there exist $K, K' \in C_{i-1}$ with $u \in PN_D(x_K) \cup PN_D(y_K)$ and $u' \in PN_D(x_{K'}) \cup PN_D(y_{K'})$ such that
 - $\cdot uu' \in E(G)$ and
 - $\cdot K_{i-1} \in S_{uu'} \text{ or } K'_{i-1} \in S_{uu'},$
 - then output K_{i-1}, K'_{i-1}, K, K' .
- 4. If $|C_{i-1}| \ge |A|$ then let $K_i, K'_i \in C_{i-1}$ be two cliques for which there exist $u_i \in PN_D(x_{K_i}) \cup PN_D(y_{K_i})$ and $u'_i \in PN_D(x_{K'_i}) \cup PN_D(y_{K'_i})$ such that $u_i u'_i \in E(G)$; and set $C_i = C_{i-1} \cap S_{u_i u'_i}$.
- 5. If $|C_{i-1}| < |A|$ then set $C_i = C_{i-1}$.
- 6. Return to step 2.

Before showing correctness of the above algorithm, let us first note that for every $j \ge 1$ satisfying the condition in step 4 (that is, $|C_j| \ge |A|$), the existence of the cliques K_j and K'_j is guaranteed by Claim 4. Now let us show that if the above algorithm terminates then its output is indeed as claimed. Assume that the algorithm terminates when the counter *i* equals some value $j \ge 1$ and let K_{j-1}, K'_{j-1}, K, K' be the output. Observe first that since the algorithm terminates, the condition in step 5 is never satisfied during its run (the algorithm would have otherwise looped indefinitely); in particular $C_{j-1} = \bigcap_{1 \le \ell \le j-1} S_{u_\ell u'_\ell}$. Since by construction, K and K' both belong to C_{j-1} , they belong in particular to $S_{u_{j-1}u'_{j-1}}$ and so, item (i) of Claim 6 indeed holds. Furthermore, by construction, there exist $u \in PN_D(x_K) \cup PN_D(y_K)$ and $u' \in PN_D(x_{K'}) \cup PN_D(y_{K'})$ with $uu' \in E(G)$, such that $K_{j-1} \in S_{uu'}$ or $K'_{j-1} \in S_{uu'}$, and thus, item (ii) of Claim 6 holds true as well.

There remains to show that the algorithm indeed terminates. To this end, let us first show by induction that for every $j \in [k^2+1]$, the condition in step 4 is always satisfied, that is, $|C_{j-1}| \ge |A|$. More specifically, we show that for every $j \in [k^2]$, $|C_j| \ge (k^2 - j)(k(k-1)/2 + 2) + |A|$. For j = 1, the result readily follows from the fact that $|\overline{\mathcal{K}_D}| = |C_1| + |\overline{\mathcal{K}_D} \setminus C_1| \ge k^2(k(k-1)/2 + 2) + |A|$ by assumption and $|\overline{\mathcal{K}_D} \setminus C_1| \le k(k-1)/2 + 2$ by Claim 5(ii). For j > 1, $|C_j| = |S_{u_ju'_j} \cap C_{j-1}|$ by definition and so,

$$\begin{aligned} |C_j| &= |\overline{\mathcal{K}_D}| - |\overline{\mathcal{K}_D} \setminus C_j| \\ &= |\overline{\mathcal{K}_D}| - |(\overline{\mathcal{K}_D} \setminus S_{u_j u'_j}) \cup (\overline{\mathcal{K}_D} \setminus C_{j-1})| \\ &\geq |\overline{\mathcal{K}_D}| - (|\overline{\mathcal{K}_D} \setminus S_{u_j u'_j}| + |\overline{\mathcal{K}_D} \setminus C_{j-1}|) \\ &= |C_{j-1}| - |\overline{\mathcal{K}_D} \setminus S_{u_j u'_j}|. \end{aligned}$$

But $|C_{j-1}| \ge (k^2 - (j-1))(k(k-1)/2 + 2) + |A|$ by the induction hypothesis and $|\overline{\mathcal{K}_D} \setminus S_{u_j u'_j}| \ge k(k-1)/2 + 2$ by Claim 5(ii) and thus, $|C_j| \ge (k^2 - j)(k(k-1)/2 + 2) + |A|$ as claimed.

Now suppose to the contrary that the algorithm has not yet terminated by the time the counter i reaches the value t = k(k-1)/4 + 3. Then for every j < t, $K_j, K'_j \notin S_{u_t u'_t}$: indeed, if there exist indices $j \in [t-1]$ such that one of K_j and K'_j belongs to $S_{u_t u'_t}$ then, letting ℓ be the smallest such index, we have $C_{t-1} \subseteq C_{\ell}$ and $K_t, K'_t \in C_{t-1}$; but then, the algorithm would have terminated after setting the counter i to $\ell + 1$ and output $K_{\ell}, K'_{\ell}, K = K_t, K' = K'_t$. It follows that $\{K_j, K'_j \mid j \in [t-1]\} \subseteq \overline{\mathcal{K}_D} \setminus S_{u_t u'_t}$; but $|\{K_j, K'_j \mid j \in [t-1]\}| = 2(t-1) = k(k-1)/2 + 4$, a contradiction to Claim 5(ii).

To conclude, assume that G is a NO-instance for 2-EDGE CONTRACTION(γ_t). Then by Observation 2 and Claim 7,

$$|D \cap N[\overline{\mathcal{K}_D}]| = 2|\overline{\mathcal{K}_D}| \le k^2(k(k-1)+4) + 2(|A|-1)$$

and thus, combined with Claim 2, we conclude that

$$|D \cap N[C]| = |D \cap N[\mathcal{K}_D]| + |D \cap N[\overline{\mathcal{K}_D}| \le k^2(k(k-1)+4) + 6|A| - 2.$$

Now observe that $D' = D \setminus N[C]$ has size at most |A|: indeed, since D' dominates (only) vertices from $A \cup B$, if |D'| > |A| then $(D \setminus D') \cup A$ is a TD set of G of size strictly less than that of D, a contradiction. It follows that

$$\gamma_t(G) = |D \cap N[C]| + |D \setminus N[C]| \le k^2(k(k-1)+4) + 7|A| - 2$$

and so, we may take $f(k) = k^4 + 4k^2 + 21k + 19$ as claimed.

Since for any graph G, G is a YES-instance for 2-EDGE CONTRACTION(γ_t) if and only if G is a NO-instance for CONTRACTION NUMBER(γ_t , 3), the following ensues from Lemma 3.1.

Corollary 3.2. For every $k \ge 0$, CONTRACTION NUMBER $(\gamma_t, 3)$ is polynomial-time solvable on $(P_6 + kP_3)$ -free graphs.

Lemma 3.3. CONTRACTION NUMBER $(\gamma_t, 2)$ is polynomial-time solvable on H-free graphs if $H \subseteq_i P_5 + tK_1$ for some $t \ge 0$, or $H \subseteq_i P_4 + tP_3$ for some $t \ge 0$.

Proof. Assume that $H \subseteq_i P_5 + tK_1$ for some $t \ge 0$ (the case where $H \subseteq_i P_4 + tP_3$ for some $t \ge 0$ is handled similarly) and let G be an H-free graph. Since H is a fortiori an induced subgraph of $P_6 + tP_3$, we may use the polynomial-time algorithm for $(P_6 + tP_3)$ -free graphs given by Corollary 3.2 to determine whether G is a YES-instance for CONTRACTION NUMBER $(\gamma_t, 3)$ or not. If the answer is yes then we output NO; otherwise, we use the polynomial-time algorithm for $(P_5 + tK_1)$ -free graphs given by Theorem 1.6 to determine whether G is a YES-instance for CONTRACTION NUMBER $(\gamma_t, 1)$ or not, and output the answer accordingly.




Figure 3: The variable gadget G_x for a variable x contained in clauses c, c' and c'' (a rectangle indicates that the corresponding set of vertices induces a clique).




Figure 4: The clause gadget G_c for a clause $c = x \lor y \lor z$ is the disjoint union of G_c^T and G_c^F (a rectangle indicates that the corresponding set of vertices is a clique).

3.2 Hardness results

In this section, we show that CONTRACTION NUMBER $(\gamma_t, 2)$ and CONTRACTION NUMBER $(\gamma_t, 3)$ are NP-hard on a number of monogenic graph classes. We first consider the case k = 2.

Lemma 3.4. CONTRACTION NUMBER $(\gamma_t, 2)$ is NP-hard on $K_{1,3}$ -free graphs.

Proof. We use the same construction as in [11, Theorem 6]. More precisely, we reduce from POSITIVE EXACTLY 3-BOUNDED 1-IN-3 3-SAT (see Section 2 for a precise definition of this problem): given an instance Φ of this problem, with variable set X and clause set C, we construct an instance G of CONTRACTION NUMBER $(\gamma_t, 2)$ as follows. For every variable $x \in X$ appearing in clauses c, c' and c'', we introduce the gadget G_x depicted in Figure 3 (where a rectangle indicates that the corresponding set of vertices is a clique). For every clause $c \in C$ containing variables x, yand z, we introduce the gadget G_c which is the disjoint union of the graph G_c^T and the graph G_c^F depicted in Figure 4 (where a rectangle indicates that the corresponding set of vertices is a clique) and further add for every $\ell \in \{x, y, z\}$, an edge between t_{ℓ}^c and t_c^ℓ , and an edge between f_{ℓ}^c and f_c^r . We let G be the resulting graph. Let us show that Φ is satisfiable if and only if $ct_{\gamma_t}(G) = 2$. To do so, we will rely on the following key result shown in [11].

Claim 8 ([11]). The following statements are equivalent.

(i) Φ is satisfiable.

(*ii*) $\gamma_t(G) = 14|X| + 8|C|$.

(iii) $ct_{\gamma_t}(G) > 1.$

Now assume that Φ is satisfiable and consider a truth assignment satisfying Φ . We construct a minimum TD set D of G as follows. For every variable $x \in X$ appearing in clauses c, c' and c'', if x is true then we include $\{u_x, T_x\} \cup \{d_x^p, t_x^p, h_x^p, j_x^p \mid p \in \{c, c', c''\}\}$ in D; otherwise, we include $\{u_x, F_x\} \cup \{b_x^p, d_x^p, j_x^p, f_x^p \mid p \in \{c, c', c''\}\}$ in D. For every clause $c \in C$ containing variables x, yand z, exactly one variable is true, say x without loss of generality, in which case we include $\{v_c, g_c^x\} \cup \{c_c^x, a_c^x\} \cup \{c_c^t, d_c^t \mid t \in \{y, z\}\}$ in D. It is not difficult to see that the constructed set D is indeed a TD set and since |D| = 14|X| + 8|C|, we conclude by Claim 8 that D is minimum. Now consider a clause $c \in C$ containing variables x, y and z and assume without loss of generality that x is true (and thus y and z are false). Then $D \cup \{a_c^y\}$ contains the P_4 $c_c^x a_c^x a_c^y c_c^y$ and thus, $ct_{\gamma_t}(G) \leq 2$ by Theorem 1.5; but $ct_{\gamma_t}(G) > 1$ by Claim 8 and so, $ct_{\gamma_t}(G) = 2$. Conversely, if $ct_{\gamma_t}(G) = 2$ then Φ is satisfiable by Claim 8. Since G is $K_{1,3}$ -free, the lemma follows.

Lemma 3.5. CONTRACTION NUMBER $(\gamma_t, 2)$ is coNP-hard on P_6 -free graphs.

Proof. If G is a P_6 -free graph then $ct_{\gamma_t}(G) \leq 2$ as shown in the proof of Lemma 3.1; and since CONTRACTION NUMBER $(\gamma_t, 1)$ is NP-hard on P_6 -free graphs by Theorem 1.6, the lemma follows from Observation 1.

Lemma 3.6. CONTRACTION NUMBER $(\gamma_t, 2)$ is NP-hard on $2P_4$ -free graphs.

Proof. We use the same reduction as in [11, Theorem 5]. More precisely, we reduce from 3-SAT: given an instance Φ of this problem, with variable set X and clause set C, we construct an instance G of CONTRACTION NUMBER $(\gamma_t, 2)$ as follows. For every variable $x \in X$, we introduce the gadget G_x consisting of a triangle on vertex set $\{x, \overline{x}, u_x\}$ and an additional vertex v_x adjacent to u_x (that is, G_x is a paw); the vertices x and \overline{x} are referred to as *literal vertices*. For every clause $c \in C$, we introduce a *clause vertex*, denoted by c, and add an edge between c and every literal vertex whose corresponding literal appears in the clause c. Finally, we add an edge between every two clause vertices so that the set of clause vertices induces a clique. We let G be the resulting graph. We next show that Φ is satisfiable if and only if $ct_{\gamma_t}(G) = 2$. To do so, we will rely on the following key results proved in [11].

Claim 9 ([11]). The following statements are equivalent.

(i) Φ is satisfiable.

(*ii*) $\gamma_t(G) = 2|X|$.

(iii) $ct_{\gamma_t}(G) > 1.$

Now assume that Φ is satisfiable and consider a truth assignment satisfying Φ . We construct a minimum TD set D of G as follows. For every variable $x \in X$, if x is true then we include $\{u_x, x\}$ in D; otherwise, we include $\{u_x, \overline{x}\}$ in D. It is not difficult to see that the constructed set D is indeed a TD set of G and since |D| = 2|X|, we conclude by Claim 9 that D is minimum. Now consider a clause $c \in C$ containing variables x, y and z, and assume without loss of generality that x and y both appear positive in c. Then the set $(D \setminus (V(G_x) \cup V(G_y))) \cup \{x, u_x, y, u_y, c\}$ is a TD set of G of size $\gamma_t(G) + 1$ containing the $P_4 \ u_x, x, c, u_y$ and thus, $ct_{\gamma_t}(G) \leq 2$ by Theorem 1.5; but $ct_{\gamma_t}(G) > 1$ by Claim 9 and so, $ct_{\gamma_t}(G) = 2$. Conversely, if $ct_{\gamma_t}(G) = 2$ then Φ is satisfiable by Claim 9. Since G is readily seen to be $2P_4$ -free, the lemma follows.

Lemma 3.7. CONTRACTION NUMBER $(\gamma_t, 2)$ is coNP-hard on $(P_5 + P_2)$ -free graphs.

Proof. Since $P_5 + P_2 \subseteq_i P_6 + P_3$, CONTRACTION NUMBER $(\gamma_t, 3)$ is polynomial-time solvable on $(P_5 + P_2)$ -free graphs by Corollary 3.2; but CONTRACTION NUMBER $(\gamma_t, 1)$ is NP-hard on $(P_5 + P_2)$ -free graphs by Theorem 1.6 and thus, the lemma follows from Observation 1.

Lemma 3.8. Let G be a graph such that $\gamma_t(G) \ge 3$ and let H be the graph obtained by 4-subdividing every edge of G. Then $ct_{\gamma_t}(G) = ct_{\gamma_t}(H)$.

Proof. It was shown in [11, Lemma 7] that $\gamma_t(H) = \gamma_t(G) + 2|E(G)|$. Let us show how to construct from a TD set D of G a TD set T(D) of H of size |D| + 2|E(G)| (see Figure 5). For every edge $e = uv \in E(G)$, we will denote by $ue_1e_2e_3e_4v$ the P_6 in H replacing the edge e. Firstly, we include in T(D) every vertex of D. Then for every edge $e = uv \in E(G)$, if $D \cap \{u, v\} = \emptyset$, we further include $\{e_2, e_3\}$ in T(D); if $|D \cap \{u, v\}| = 1$, say $u \in D$ without loss of generality, we further include $\{e_3, e_4\}$ in T(D); and if $u, v \in D$, we further include $\{e_1, e_4\}$ in T(D). It is not difficult to see that the constructed set T(D) is indeed a TD set of H and that |T(D)| = |D| + 2|E(G)|.

Conversely, given a TD set D of H, let us show how to construct from D a TD set $T^{-1}(D)$ of G of size at most |D| - 2|E(G)|. To this end, we show how, given a graph F, a graph F' obtained from F by 4-subdividing one edge $uv \in E(F)$ and a TD set $D_{F'}$ of F', we can construct from




Figure 5: Constructing a total dominating set of H from a total dominating set of G (vertices in red belong to the corresponding total dominating set).

 $D_{F'}$ a TD set D_F of F of size at most $|D_{F'}| - 2$. Then by iterating this procedure on H, we will obtain the desired TD set $T^{-1}(D)$ of G.

Let $ue_1e_2e_3e_4v$ be the path in F' corresponding to the 4-subdivision of the edge $uv \in E(F)$. If $e_1 \in D_{F'}$ and $u \notin D_{F'}$ then necessarily $e_2 \in D_{F'}$, for e_1 would otherwise not be dominated. Similarly, if $e_4 \in D_{F'}$ and $v \notin D_{F'}$ then necessarily $e_3 \in D_{F'}$, for e_4 would otherwise not be dominated. Thus, if $e_1, e_4 \in D_{F'}$, we let $D_F = (D_{F'} \setminus \{e_1, e_2, e_3, e_4\}) \cup \{u, v\}$. Suppose next that $e_4 \notin D_{F'}$. Then $e_2 \in D_{F'}$, for e_3 would otherwise not be dominated; and if $v \notin D_{F'}$ then $e_3 \in D_{F'}$, for e_4 would otherwise not be dominated. Thus, if $e_1 \in D_{F'}$ and $e_4 \notin D_{F'}$, then either $v \in D_{F'}$ in which case we let $D_F = D_{F'} \setminus \{e_1, e_2, e_3, e_4\}$, or $v \in D_{F'}$ in which case we let $D_F = (D_{F'} \setminus \{e_1, e_2, e_3, e_4\}) \cup \{v\}$. We proceed symmetrically if $e_1 \notin D_{F'}$ and $e_4 \in D_{F'}$. Finally, if $e_1, e_4 \notin D_{F'}$ then $e_2, e_3 \in D_{F'}$ and so, we may take $D_F = D_{F'} \setminus \{e_1, e_2, e_3, e_4\}$. Now let us observe that the TD set $T^{-1}(D)$ constructed as such satisfies the following property.

Observation 4. For every edge $e = uv \in E(G)$, if $e_1 \in D$ ($e_4 \in D$, respectively) then $v \in T^{-1}(D)$ ($u \in T^{-1}(D)$, respectively).

We next show that $ct_{\gamma_t}(G) = ct_{\gamma_t}(H)$. As the following was shown in [11, Lemma 7], it in fact suffices to show that $ct_{\gamma_t}(G) = 2$ if and only if $ct_{\gamma_t}(H) = 2$.

Claim 10 ([11]). $ct_{\gamma_t}(G) = 1$ if and only if $ct_{\gamma_t}(H) = 1$.

Assume first that $ct_{\gamma_t}(G) = 2$ and let D be a TD set of G of size $\gamma_t(G) + 1$ containing a P_4 , a $2P_3$ or a $K_{1,3}$ (see Theorem 1.5). Suppose first that D contains a P_4 with edge set $e^1 = uv$, $e^2 = vw$ and $e^3 = wt$. Then by construction, the TD set T(D) of G contains the $2P_3 e_4^1 v e_1^2, e_4^2 w e_1^3$. Second, suppose that D contains a $2P_3$ with edge set $e^1 = uv$, $e^2 = vw$, $e^3 = xy$ and $e^4 = yz$. Then by construction, the TD set T(D) of G contains the $2P_3 e_4^1 v e_1^2, e_4^3 w e_1^4$. Suppose finally that Dcontains a $K_{1,3}$ on edge set $e^1 = uv$, $e^2 = wv$ and $e^3 = tv$. Then by construction, the TD set T(D)contains the $K_{1,3} v, e_4^1, e_4^2, e_4^3$. In all three cases, we conclude by Theorem 1.5 that $ct_{\gamma_t}(H) \leq 2$ and by Claim 10 that in fact $ct_{\gamma_t}(H) = 2$.

Conversely assume that $ct_{\gamma_t}(H) = 2$ and denote by \mathcal{D} the set of TD sets of H of size $\gamma_t(H) + 1$ containing a P_4 , a $2P_3$ or a $K_{1,3}$ (note that $\mathcal{D} \neq \emptyset$ by Theorem 1.5). Observe that if there exists $D \in \mathcal{D}$ such that $T^{-1}(D)$ contains a P_4 , a $2P_3$ or a $K_{1,3}$, then since $ct_{\gamma_t}(G) > 1$ by Claim 10, it must be that $|T^{-1}(D)| = \gamma_t(G) + 1$ and thus, $ct_{\gamma_t}(G) = 2$ by Theorem 1.5. Now if there exists $D \in \mathcal{D}$ such that D contains a $K_{1,3}$ u, e_1^1, e_1^2, e_1^3 where u is the vertex of degree three and $e^i = ux_i$ for every $i \in [3]$, then by Observation 4, the TD set $T^{-1}(D)$ contains the $K_{1,3}$ u, x_1, x_2, x_3 and so, $ct_{\gamma_t}(G) = 2$ by the above.

Assume henceforth that no TD set in \mathcal{D} contains a $K_{1,3}$. We contend that there then exists a TD set in \mathcal{D} containing a $2P_3$. To prove this claim, let us first show that there exists $D \in \mathcal{D}$ such that for every edge $e \in E(G)$, $\{e_1, e_2, e_3, e_4\} \not\subseteq D$. In the following, given a TD set $D \in \mathcal{D}$ and an edge $e \in E(G)$, if $\{e_1, e_2, e_3, e_4\} \subseteq D$ then D is said to accommodate the edge uv. Consider a TD set $D \in \mathcal{D}$ accommodating the minimum number of edges amongst every TD set in \mathcal{D} , and suppose to the contrary that D accommodates an edge $e = uv \in E(G)$. Let us show that $\mathcal{D} \setminus \{D\}$ contains a TD set accommodating fewer edges than D, which would contradict the choice of D. First note that one of u and v does not belong to D: indeed, if $\{u, v\} \subseteq D$ then $D \setminus \{e_3\}$ is a minimum TD set of H containing the P_3 ue_1e_2 , a contradiction to Theorem 1.5. Furthermore, since $\gamma_t(G) \geq 3$, at least one of u and v has degree at least two in G.

Suppose first that there exists $x \in \{u, v\} \setminus D$ such that $d_G(x) \geq 2$, say x = v without loss of generality, and let f = vw be an edge of G. Note that since $v \notin D$ by assumption, necessarily $f_2 \in D$, for f_1 would otherwise not be dominated. Now if $f_1 \notin D$ then $f_3 \in D$ as f_2 should be dominated; but then, the TD set $(D \setminus \{e_4\}) \cup \{f_1\}$ belongs to \mathcal{D} (note indeed that it contains $e_1e_2e_3$ and $f_1f_2f_3$) and accommodates fewer edges than D, a contradiction to the choice of D. Thus, it must be that $f_1 \in D$; but then, the TD set $(D \setminus \{e_3\}) \cup \{v\}$ belongs to \mathcal{D} (note indeed that it contains the $P_4 \ e_4vf_1f_2$) and accommodates fewer edges than D, a contradiction to the choice of D.

Suppose second that for every $x \in \{u, v\} \setminus D$, $d_G(x) = 1$. By the above, $\{u, v\} \setminus D \neq \emptyset$ and at least one of u and v has degree at least two in G: let us assume without loss of generality that $u \notin D$ and $d_G(v) \ge 2$ (note that then $d_G(u) = 1$ and $v \in D$). Observe that since $D \setminus \{e_3\}$ is a minimum TD set of H, it cannot contain a P_3 by Theorem 1.5. Now among the neighbours of v, there must be one of degree at least two, for G is otherwise a star thereby contradicting the fact that $\gamma_t(G) \ge 3$. Thus, let f = vw be an edge of G such that $d_G(w) \ge 2$. Then since $f_1 \notin D$ $(D \setminus \{e_3\}$ would otherwise contain the $P_3 e_4vf_1$), necessarily $f_3 \in D$ as f_2 should be dominated; and since f_3 should be dominated but $D \setminus \{e_3\}$ cannot contain a P_3 , necessarily $|D \cap \{f_2, f_4\}| = 1$. Now if $f_2 \in D$ then the TD set $(D \setminus \{e_4\}) \cup \{f_1\}$ belongs to \mathcal{D} (note indeed that it contains the P_4 $vf_1f_2f_3$) and accommodates fewer edges than D (recall that in this case $f_4 \notin D$), a contradiction to the choice of D. Now if $f_4 \in D$ then $w \notin D$, for $D \setminus \{e_3\}$ would otherwise contain the $P_3 f_3f_4w$; but then, by considering the TD set $(D \setminus \{e_3, e_4\}) \cup \{f_1, f_2\}$ in place of D, we may argue as in the previous case (recall indeed that $w \notin D$ and $d_G(w) \ge 2$) and conclude similarly to a contradiction. Therefore, D accommodates no edge.

Let us next show that among those TD sets of \mathcal{D} accommodating no edge, there exists a TD set D such that D contains a $2P_3$. Indeed, let $D \in \mathcal{D}$ be a TD set which accommodates no edge and suppose that D contains no $2P_3$. Let us show how obtain from D a TD set with a $2P_3$. Note that since D contains no $2P_3$, D contains a P_4 by assumption. Now suppose first that D contains a P_4 of the form $ue_1e_2e_3$, where $e = uv \in E(G)$. Since D does not accommodate the edge uv, $e_4 \notin D$ and so, there must exist $f = vw \in E(G)$ such that $f_1 \in D$ (v would otherwise not be dominated). But then, either $v \notin D$ in which case $f_2 \in D$ (f_1 would otherwise not be dominated) and thus, the TD set ($D \setminus \{e_3\} \cup \{v\}$ contains the $2P_3$ ue_1e_2, vf_1f_2 ; or $v \in D$ in which case the TD set ($D \setminus \{e_3\} \cup \{e_4\}$ contains the $2P_3$ ue_1e_2, e_4vf_1 . Second, suppose that D contains a P_4 of the form $e_4vf_1f_2$, where $e = uv, f = vw \in E(G)$. Then we may assume that $f_3 \notin D$ (we revert to the previous case otherwise) which implies that $w \in D$ for otherwise, f_4 would not be dominated. But then, either $f_4 \in D$ in which case the TD set ($D \setminus \{f_2\} \cup \{f_3\}$ contains the $2P_3$ e_4vf_1, f_3f_4w ; or $f_4 \notin D$ in which case the TD set ($D \setminus \{f_2\} \cup \{f_4\}$ contains the $2P_3$ e_4vf_1, f_4wt where $t \in N_H(w) \cap D$.

Finally, let us show that among those TD sets in \mathcal{D} accommodating no edge and containing a 2P₃, there exists a TD set D such that (1) no P₃ in D is of the form $e_1e_2e_3$ where $e \in E(G)$, and (2) if D contains a P_3 of the form ue_1e_2 , where $e = uv \in E(G)$, then v has a neighbour $f_1 \in D \setminus \{e_4\}$ where $f = vw \in E(G)$. To this end, let $D \in \mathcal{D}$ be a TD set accommodating no edge and containing a $2P_3$. Suppose that D contains a P_3 of the form $e_1e_2e_3$ where $e = uv \in E(G)$. Since D does not accommodate the edge $uv, e_4 \notin D$, which implies that v must have a neighbour $f_1 \in D$ where $f = vw \in E(G)$. But then, either $w \in D$ in which case we may consider the TD set $(D \setminus \{e_3\}) \cup \{e_4\}$ in place of D; or $w \notin D$ in which case $f_2 \in D$ $(f_1$ would otherwise not be dominated) and we may consider the TD set $(D \setminus \{e_3\}) \cup \{e_4\}$ in place of D. By iterating this process, we eventually reach a TD set $D \in \mathcal{D}$ accommodating no edge, containing a $2P_3$ and satisfying (1). Now suppose that this TD set D contains a P_3 of the form ue_1e_2 , where $e = uv \in E(G)$, and suppose that $N_H(v) \cap D = \{e_4\}$. Then since D does not accommodate the edge uv, necessarily $e_3 \notin D$ and so, $v \in D$ as e_4 would otherwise not be dominated. Thus, if there exists an edge $f = vw \in E(G)$, then we may consider the TD set $(D \setminus \{e_4\}) \cup \{f_1\}$ in place of D. Assume therefore that no such edge exists, that is, $d_G(v) = 1$. Then surely, there exists an edge $f = uw \in E(G)$ since $\gamma_t(G) \geq 3$. But then, either $f_1 \in D$ in which case we may consider the TD set $D' = (D \setminus \{e_2\}) \cup \{e_3\}$ in place of D (note indeed that the path ve_4e_3 in D' then satisfies (2) as $f_1 \in D'$; or $f_1 \notin D$ in which case we may consider the TD set $(D \setminus \{e_2\}) \cup \{f_1\}$ (note that we here simply discard the path ue_1e_2). By iterating this process, we eventually reach a TD set in \mathcal{D} accommodating no edge, containing a $2P_3$ and satisfying both (1) and (2).

To conclude, let $D \in \mathcal{D}$ be a TD set accommodating no edge, containing a $2P_3$ and satisfying




Figure 6: The variable gadget G_x for a variable x contained in clause c, c' and c'' (a rectangle indicates that the corresponding set of vertices induces a clique).

both (1) and (2). Then by (1), D contains only P_3 s of the form f_4ve_1 where $f = uv, e = vw \in E(G)$, or the form ue_1e_2 where $e = uv \in E(G)$. Now if D contains a P_3 of the form f_4ve_1 where $f = uv, e = vw \in E(G)$, then uvw is a P_3 of $T^{-1}(D)$ by Observation 4. Similarly, if D contains a P_3 of the form ue_1e_2 where $e = uv \in E(G)$, then by (2), v has a neighbour $f_1 \in D \setminus \{e_4\}$ where f = vw, and thus, uvw is a P_3 of $T^{-1}(D)$ by Observation 4. Since any two P_{3s} uvw and u'v'w' in $T^{-1}(D)$ corresponding to two distinct P_{3s} in D have at most two common vertices (it may be indeed that $\{u, w\} \cap \{u', w'\} \neq \emptyset$), we conclude that $T^{-1}(D)$ contains a $2P_3$ if $\{u, w\} \cap \{u', w'\} = \emptyset$, and a P_4 otherwise. Therefore, $ct_{\gamma_t}(G) \leq 2$ by Theorem 1.5 and since $ct_{\gamma_t}(G) > 1$ by Claim 10, in fact $ct_{\gamma_t}(G) = 2$.

By 4-subdividing an instance of CONTRACTION NUMBER (γ_t, k) sufficiently many times, the following ensues from Lemma 3.8.

Lemma 3.9. For every $k \in \{2,3\}$ and $\ell \geq 3$, CONTRACTION NUMBER (γ_t, k) is NP-hard on $\{C_3, \ldots, C_\ell\}$ -free graphs.

The last result of this section concerns CONTRACTION NUMBER(γ_t ,3). Combining Lemma 3.8 with the following result, we obtain in particular that if CONTRACTION NUMBER(γ_t ,3) is polynomial-time solvable on *H*-free graphs, then *H* must be a linear forest.

Lemma 3.10. CONTRACTION NUMBER $(\gamma_t, 3)$ is NP-hard on $K_{1,3}$ -free graphs.

Proof. We reduce from POSITIVE EXACTLY 3-BOUNDED 1-IN-3 3-SAT (see Section 2 for a precise definition of this problem): given an instance Φ of this problem, with variable set X and clause set C, we construct an instance G of CONTRACTION NUMBER(γ_t ,3) as follows. For every variable $x \in X$ appearing in clauses c, c' and c'', we introduce the gadget G_x depicted in Figure 6 (where a rectangle indicates that the corresponding set of vertices is a clique). In the following, we denote by P_x the paw induced by $\{T_x, F_x, v_x, u_x\}$ and we may refer to the vertices of P_x as $P_x(1), \ldots, P_x(4)$ where $P_x(1) = T_x$ and $P_x(2) = F_x$. For every clause $c \in C$ containing variables x, y and z, we introduce the gadget G_c which is the disjoint union of the graphs G_c^T and G_c^F depicted un Figure 7 (where a rectangle indicates that the corresponding set of vertices is a clique) and further add for every $\ell \in \{x, y, x\}$, an edge between $P_{\ell,T}^c(1)$ t_{ℓ}^c , and an edge between $P_{x,F}^c(2)$ and f_{ℓ}^c . In the following, we denote by K_c^T the clique induced by p_c^x, p_c^y and p_c^z , and by K_c^F the clique induced by q_c^x, q_x^y and q_c^z . We let G be the resulting graph. We next show that $ct_{\gamma_t}(G) = 3$ if and only if Φ is satisfiable through a series of claims.

Claim 11. For every TD set D of G, the following hold.

(i) For every clause $c \in C$ and for every $R \in \{T, F\}$, $|D \cap V(G_c^R)| \ge 2$. Furthermore, $v_c \in D$. (ii) For every variable $x \in X$ and every paw P of G_x , $|D \cap V(P)| \ge 2$ and $P(3) \in D$.

Proof. To prove (i), let $c \in C$ be a clause containing variables x, y and z. Then since u_c should be dominated, necessarily $v_c \in D$. Furthermore, $D \cap \{u_c, q_c^x, q_c^y, q_c^z\} \neq \emptyset$ as v_c should be dominated. Thus, $|D \cap V(G_c^F)| \geq 2$. Now since every vertex of K_c^T must be dominated, either $D \cap V(K_c^T) = \emptyset$



Figure 7: The clause gadget G_c for a clause $c = x \lor y \lor z$ is the disjoint union of G_c^T and G_c^F (a rectangle indicates that the corresponding set of vertices is a clique).

in which case $\{t_c^x, t_c^y, t_c^z\} \subseteq D$; or $D \cap V(K_c^T) \neq \emptyset$, say $p_c^x \in D$ without loss of generality, in which case $D \cap \{t_c^x, p_c^y, p_c^z\} \neq \emptyset$ as p_c^x should be dominated as well. In both cases, $|D \cap V(G_c^T)| \ge 2$.

To prove (ii), let $x \in X$ be a variable and let P be a paw in G_x . Then since P(4) should be dominated, necessarily $P(3) \in D$; and since P(3) should be dominated, necessarily $D \cap \{P(1), P(2), P(4)\} \neq \emptyset$. Thus, $|D \cap V(P)| \ge 2$.

The following is an immediate consequence of Claim 11(ii).

Observation 5. For every TD set D of G and every variable $x \in X$ contained in clauses c, c'and c'', if $|D \cap V(G_x)| = 14$ then $D \cap \{q_x^\ell, p_x^\ell \mid \ell \in \{c, c', c''\}\} = \emptyset$.

The following is an immediate consequence of Claim 11(i).

Observation 6. Let D be a TD set of G. If $|D \cap V(G_c^F)| = 2$ for some clause $c \in C$ containing variables x, y and z, then $D \cap \{f_c^x, f_c^y, f_c^z\} = \emptyset$.

Claim 12. Let D be a TD set of G and let $x \in X$ be a variable appearing in clauses c, c' and c''. If $|D \cap V(G_x)| = 14$ then the following hold.

(i) If there exists $\ell \in \{c, c', c''\}$ such that $P_{x,T}^{\ell}(1) \in D$ then $T_x \in D$.

(ii) It there exists $\ell \in \{c, c', c''\}$ such that $P_{x,F}^{\ell}(2) \in D$ then $F_x \in D$.

Proof. Assume that $|D \cap V(G_x)| = 14$. If there exists $\ell \in \{c, c', c''\}$ such that $P_{x,T}^{\ell}(1) \in D$, then $P_{x,T}^{\ell}(2) \notin D$ by Claim 11(i); and since $D \cap \{p_x^c, p_x^{c'}, p_x^{c''}\} = \emptyset$ by Observation 5, necessarily $T_x \in D$ as p_x^{ℓ} should be dominated. Item (ii) follows by symmetry.

Claim 13. Φ is satisfiable if and only if $\gamma_t(G) = 14|X| + 4|C|$.

Proof. Assume first that Φ is satisfiable and consider a truth assignment satisfying Φ . We construct a TD set D of G as follows. For every variable $x \in X$ appearing in clauses c, c' and c'', if x is true then we include $\{v_x, T_x\} \cup \{P_{x,R}^{\ell}(1), P_{x,R}^{\ell}(3) \mid \ell \in \{c, c', c''\}$ and $R \in \{T, F\}\}$ in D; otherwise, we include $\{v_x, F_x\} \cup \{P_{x,R}^{\ell}(2), P_{x,R}^{\ell}(3) \mid \ell \in \{c, c', c''\}$ and $R \in \{T, F\}\}$ in D. For every clause $c \in C$ containing variables x, y and z, exactly one variable is true, say x without loss of generality, in which case we include $\{v_c, q_c^x\} \cup \{p_c^y, p_c^z\}$ in D. It is easy to see that the constructed set D is indeed a TD set of G and since |D| = 14|X| + 4|C|, we conclude by Claim 11 that D is minimum.

Conversely, assume that $\gamma_t(G) = 14|X| + 4|C|$ and let D be a minimum TD set of G. Consider a clause $c \in C$ containing variables x, y and z. Since $|D \cap V(G_c^T)| = 2$ by Claim 11(i), at least one of t_c^x, t_c^y and t_c^z is not dominated by a vertex in $V(G_c^T)$, say $N(t_c^x) \cap D \subseteq V(G_x)$ without loss of generality. Then since $P_{x,T}^c(1) \in D$, it follows from Claim 12(i) that $T_x \in D$ and so, $F_x \notin D$ by Claim 11(ii). But then, $P_{x,F}^c(2) \notin D$ by Claim 12(ii) and since $f_c^x \notin D$ by Observation 6, necessarily $q_c^x \in D$ as f_c^x should be dominated. It then follows from Claim 11(i) that $D \cap \{q_c^y, q_c^z, f_c^y, f_c^z\} = \emptyset$, which implies that $P_{y,F}^c(2), P_{z,F}^c(2) \in D$ (one of f_c^y and f_c^z would otherwise not be dominated) and so, $F_y, F_z \in D$ by Claim 12(ii). Thus, the truth assignment obtained by setting a variable x to true if $T_x \in D$ and to false otherwise, satisfies Φ .

Claim 14. Let D be a TD set of G. If there exists a clause $c \in C$ containing variables x, y and z such that $p_c^{\ell}, q_c^{\ell} \notin D$ for some $\ell \in \{x, y, z\}$, then $|D \cap V(G_\ell)| > 14$.

Proof. Let $c \in C$ be a clause containing variables x, y and z, and suppose that $p_c^{\ell}, q_c^{\ell} \notin D$ for some $\ell \in \{x, y, z\}$. Then $P_{\ell,T}^c(1), P_{\ell,F}^c(2) \in D$ since t_c^{ℓ} and f_c^{ℓ} should be dominated. Thus, if $|D \cap V(G_\ell)| = 14$ then by Claim 12(i) and (ii), $T_\ell, F_\ell \in D$, a contradiction to Claim 11(ii).

Claim 15. Let *D* be a *TD* set of *G*. If there exists a clause $c \in C$ such that $|D \cap V(G_c^T)| = |D \cap V(G_c^F)| = 2$ and $t_c^{\ell} \in D$ for some variable ℓ contained in *c*, then there exists a variable $v \neq \ell$ contained in *c* such that $|D \cap V(G_v)| > 14$.

Proof. Let $c \in C$ be a clause containing variables x, y and z, and assume that $|D \cap V(G_c^T)| = |D \cap V(G_c^T)| = 2$ and $t_c^{\ell} \in D$ for some variable $\ell \in \{x, y, z\}$. Then $D \cap V(G_c^T) = \{t_c^{\ell}, p_c^{\ell}\}$: indeed, $D \cap V(K_c^T) \neq \emptyset$ for one of p_c^x, p_c^y and p_c^z would otherwise not be dominated; and if $p_c^v \in D$ for some $v \neq \ell$, then p_c^v is not dominated. Now by Claim 11(i), $|D \cap V(K_c^F)| \leq 1$ and so, there exists a variable $v \in \{x, y, z\} \setminus \{\ell\}$ such that $q_c^v \notin D$; and since $P_c^v \notin D$ by the above, the result then follows from Claim 14.

Claim 16. $ct_{\gamma_t}(G) = 3$ if and only if $\gamma_t(G) = 14|X| + 4|C|$.

Proof. Assume first that $ct_{\gamma_t}(G) = 3$ and let D be a minimum TD set of G. Let us show that |D| = 14|X| + 4|C|. To this end, consider first a variable $x \in X$ contained in clauses c, c' and c". Since D contains no P_3 by Theorem 1.5(i), it is clear that $|D \cap V(P)| \leq 2$ for every paw P contained in G_x ; and we conclude by Claim 11 that, in fact, equality holds. Now if there exists $\ell \in \{c,c',c''\} \text{ such that } p_x^\ell \in D \text{ then } (D \setminus V(P_{x,T}^\ell)) \cup \{P_{x,T}^\ell(1), P_{x,T}^\ell(2), P_{x,T}^\ell(3)\} \text{ is a TD set of } G \in \mathcal{F}_{x,T}^\ell(2)$ of size $|D| + 1 = \gamma_t(G) + 1$ containing the $P_4 P_{x,T}^{\ell}(1) P_{x,T}^{\ell}(3) P_{x,T}^{\ell}(2) p_x^{\ell}$, a contradiction to Theorem 1.5(ii). Thus, $D \cap \{p_x^c, p_x^{c'}, p_x^{c''}\} = \emptyset$ and by symmetry, we conclude that $D \cap \{q_x^c, q_x^{c'}, q_x^{c''}\} = \emptyset$ as well. Therefore, $|D \cap V(G_x)| = 14$. Second, consider a clause $c \in C$ containing variables x, yand z. Suppose to the contrary that there exists $\ell \in \{x, y, z\}$ such that $t_c^{\ell} \in D$. If $P_{\ell,T}^c(1) \in D$ as well, then D contains the $P_3 t_c^{\ell} P_{\ell,T}^c(1) P_{\ell,T}^c(3)$ by Claim 11(ii), a contradiction to Theorem 1.5(i). Thus, $P_{\ell,T}^c(1) \notin D$; but then, by Claim 11(ii), $D \cup \{P_{\ell,T}^c(1)\}$ contains the $P_4 t_c^\ell P_{x,T}^c(1) P_{x,T}^c(3) w$ where $w \in D \cap \{P_{x,T}^c(2), P_{x,T}^c(4)\}$, a contradiction to Theorem 1.5. Thus, $D \cap \{t_c^x, t_c^y, t_c^z\} = \emptyset$. Now if there exists $\ell \in \{x, y, z\}$ such that $f_c^{\ell} \in D$, then either $P_{\ell,F}^c(2) \in D$ in which case D contains the $P_3 f_c^{\ell} P_{x,F}^c(2) P_{x,F}^c(3)$ by Claim 11(ii); or $P_{\ell,F}^c(2) \notin D$ in which case, by Claim 11(ii), $D \cup P_{\ell,F}^c(2)$ contains the $P_4 f_c^\ell P_{\ell,F}^c(2) P_{\ell,F}^c(3) w$ where $t \in D \cap \{P_{\ell,F}^c(1), P_{\ell,F}^c(4)\}$, a contradiction in both cases to Theorem 1.5. Thus, $D \cap \{f_c^x, f_c^y, f_c^z\} = \emptyset$. Finally, since D contains no P_3 by Theorem 1.5(i), it is clear that $|D \cap V(K_c^T)| \leq 2$ and $|D \cap \{u_c, v_c, q_c^x, q_c^y, q_c^x\}| \leq 2$, and so, by Claim 11(i), $|D \cap V(G_c^T)| = |D \cap V(G_c^F)| = 2$. Therefore, $\gamma_t(G) = |D| = 14|X| + 4|C|$.

Conversely, assume that $\gamma_t(G) = 14|X| + 4|C|$ and consider a minimum TD set D of G. Let us show that D contains no P_3 , which by Theorem 1.5(i), would imply that $ct_{\gamma_t}(G) > 1$. Since for every clause $c \in C$, $|D \cap V(G_c^T)| = |D \cap V(G_c^F)| = 2$ by Claim 11(i), clearly $D \cap V(G_c)$ contains no P_3 . Similarly, for every variable $x \in X$ appearing in clauses c, c' and $c'', D \cap V(G_x)$ contains no P_3 : indeed, by Claim 11(ii), $|D \cap V(P)| = 2$ for every paw P of G_x and $D \cap \{q_x^\ell, p_x^\ell \mid \ell \in \{c, c', c''\}\} = \emptyset$ by Observation 5. Thus, if D contains a P_3 , then there must exist a clause $c \in C$ such that (1) $f_c^\ell \in D$ for some variable ℓ contained in c, or (2) $t_c^\ell \in D$ for some variable ℓ contained in c. However, by Observation 6, (1) cannot hold; and by Claim 15, (2) cannot hold. Thus, D contains no P_3 .

Now suppose for a contradiction that G has a TD set D of size $\gamma_t(G)+1$ containing a P_4 , a $K_{1,3}$ or a $2P_3$ (see Theorem 1.5(ii)). Then by Claim 11, there exists either a variable $x \in X$ such that $|D \cap V(G_x)| = 15$ or a clause $c \in C$ such that $|D \cap V(G_c)| = 5$. We next distinguish these two cases.

Case 1. There exists a variable $x \in X$ such that $|D \cap V(G_x)| = 15$. Let $c, c', c'' \in C$ be the three clauses in which x is contained. First note that, since by Claim 11(ii), for every variable $v \in X \setminus \{x\}, |D \cap V(P)| = 2$ for every paw P in G_x and $q_v^\ell, p_v^\ell \notin D$ for every clause ℓ containing v, clearly $D \cap V(G_v)$ contains no $P_4, K_{1,3}$ or $2P_3$. Similarly, for every clause $\ell \in C, |D \cap V(G_\ell^T)| = |D \cap V(G_\ell^F)| = 2$ by Claim 11(i), and so, $D \cap V(G_\ell)$ contains no $P_4, K_{1,3}$ or $2P_3$. Since for every clause $\ell \in C, |D \cap V(G_\ell^T)| = |D \cap V(G_\ell^F)| = 2$ by Claim 11(i), and so, $D \cap V(G_\ell)$ contains no $P_4, K_{1,3}$ or $2P_3$. Since for every clause $\ell \in C \setminus \{c, c', c''\}$ and every variable $v \in X$ appearing in $\ell, f_\ell^v, t_\ell^v \notin D$ by Observation 6 and Claim 15, it follows that $D \cap (V(G_\ell) \cup V(G_v))$ contains no $P_4, K_{1,3}$ or $2P_3$. Now suppose that there exists a clause $\ell \in \{c, c', c''\}$ containing, apart from x, variables y and z, such that $t_\ell^v \in D$

for some $v \in \{y, z\}$ (note that by Claim 15, $t_{\ell}^x \notin D$). We contend that t_{ℓ}^v cannot be part of a P_4 , a $K_{1,3}$ or a $2P_3$. To prove this claim, let us show that $P_{v,T}^{\ell}(1) \notin D$ (since $|D \cap V(G_{\ell}^T)| = 2$, this would indeed imply our claim). Observe first that $D \cap V(G_{\ell}^T) = \{t_{\ell}^v, p_{\ell}^v\}$: indeed, $D \cap V(K_{\ell}^T) \neq \emptyset$ for one of p_{ℓ}^x, p_{ℓ}^y and p_{ℓ}^z would otherwise not be dominated; and if $p_{\ell}^u \in D$ for some $u \neq v$, then p_{ℓ}^u is not dominated, a contradiction. By Claim 14, it must then be that $q_{\ell}^u \in D$ for $u \in \{y, z\} \setminus \{v\}$, as $p_{\ell}^u \notin D$ and $|D \cap V(G_u)| = 14$. It then follows from Claim 11(i) that $q_{\ell}^v \notin D$, which implies that $P_{v,F}^{\ell}(2) \in D$, as f_{ℓ}^v should be dominated. But then, $F_{\ell} \in D$ by Claim 12(i), and since then, $T_{\ell} \notin D$ by Claim 11(ii), $P_{v,T}^{\ell}(1) \notin D$ by Claim 12(i), as claimed. Since for every $\ell \in \{c, c', c''\}$, $f_{\ell}^x \notin D$ by Observation 6 and $t_{\ell}^x \notin D$ by Claim 15, it follows that any P_4 , $K_{1,3}$ or $2P_3$ of D is in fact contained in $D \cap V(G_x)$.

Now suppose that $|D \cap V(P)| = 3$ for some paw P of G_x . Then by Claim 11(i), $D \cap \{p_x^\ell, q_x^\ell \mid \ell \in \{c, c', c''\}\} = \emptyset$ and $|D \cap V(P')| = 2$ for any paw $P' \neq P$ of G_x . Thus, $D \cap V(G_x)$ contains no P_4 , $K_{1,3}$ or $2P_3$. Suppose next that $D \cap \{p_x^c, p_x^{c'}, p_x^{c''}\} \neq \emptyset$ (the case where $D \cap \{q_x^c, q_x^c, q_x^{c''}, q_x^{c''}\} \neq \emptyset$ is symmetric). Then by Claim 11(ii), in fact $|D \cap \{p_x^c, p_x^c, p_x^{c''}\}| = 1$; furthermore, $D \cap \{q_x^c, q_x^c, q_x^{c''}, q_x^{c'''}\} = \emptyset$ and $|D \cap V(P)| = 2$ for every paw P of G_x . Now assume without loss of generality that $p_x^c \in D$. If $T_x \notin D$, then it is easy to that D at most one P_3 (namely $p_x^c P_{x,T}^c(2) P_{x,T}^c(3)$ if $P_{x,T}^c(2) \in D$). Similarly, if $P_{x,T}^c(2) \notin D$ then D contains at most one P_3 (namely $p_x^c T_x v_x$ if $T_x \in D$). Thus, it must be that $T_x, P_{x,T}^c(2) \in D$ which by Claim 11(ii), implies that $F_x, P_{x,T}^c(1) \notin D$. Since t_x^c should be dominated, it follows that $p_x^c \in D$. Furthermore, $P_{x,F}^c(2) \notin D$ by Claim 12(ii) and so, $q_c^c \in D$, as f_c^c should be dominated. Since by Claim 11(i), $|D \cap V(G_c^T)| = |D \cap V(G_c^c)| = 2$ and $v_c \in D$, it follows that there exists a variable $v \neq x$ contained in c such that $q_c^v, p_c^v \notin D$; but $|D \cap V(G_v)| = 14$, a contradiction to Claim 14.

Case 2. There exists a clause $c \in C$ such that $|D \cap V(G_c)| = 5$. Let x, y and z be the variables contained in c. First note that, since by Claim 11(ii), for every variable $v \in X$, $|D \cap V(P)| = 2$ for every paw P of G_v and $q_v^{\ell}, p_v^{\ell} \notin D$ for every clause ℓ containing v, clearly $D \cap V(G_v)$ contains no P_4 , $K_{1,3}$ or $2P_3$. Similarly, for every clause $\ell \in C \setminus \{c\}, |D \cap V(G_\ell^T)| = |D \cap V(G_\ell)^F|| = 2$ by Claim 11(i) and so, $D \cap V(G_\ell)$ contains no P_4 , $K_{1,3}$ or $2P_3$. Since for every clause $\ell \in C \setminus \{c\}$ and every variable $v \in X$ appearing in ℓ , $f_\ell^v, t_\ell^v \notin D$ by Observation 6 and Claim 15, it follows that $D \setminus (V(G_c) \cup V(G_x) \cup V(G_y) \cup V(G_z)$ contains no P_4 , $K_{1,3}$ or $2P_3$.

Suppose first that $|D \cap V(G_c^F)| = 3$ (note that then $|D \cap V(G_c^T)| = 2$). Suppose further that $D \cap \{t_c^x, t_c^y, t_c^z\} \neq \emptyset$, say $t_c^x \in D$ without loss of generality. We contend that t_c^x cannot be part of a $P_4, K_{1,3}$ or $2P_3$ in D. To prove this claim, let us show that $P_{x,T}^{(1)} \notin D$ (since $|D \cap V(G_c^T)| = 2$, this would indeed prove our claim). Observe first that $D \cap V(G_c^T) = \{p_c^x, t_c^x\}$, for if $p_c^x \notin D$, then one of p_c^y and p_c^z is not dominated. Since $|D \cap V(K_c^F)| \le 2$ by Claim 11(i), it follows that either (1) $q_c^v \notin D$ for some $v \in \{y, z\}$, or (2) $q_c^x \notin D$. If (1) holds then $p_c^v, q_c^v \notin D$ by the above; but $|D \cap V(G_v)| = 14$, a contradiction to Claim 14. Thus, (2) holds; in particular, $D \cap V(K_c^F) = \{q_c^y, q_c^z\}$. Since $v_c \in D$ by Claim 11(i), it follows that $P_{x,F}^c(2) \in D$ as f_c^x should be dominated. Then by Claim 12(ii), $F_x \in D$ which, by Claim 11(ii), implies that $T_x \notin D$; and so, by Claim 12(i), $P_{x,T}^c(1) \notin D$, as claimed. It follows that any P_4 , $K_{1,3}$ or $2P_3$ in D is in fact contained in $V(G_c^F) \cup V(G_x) \cup V(G_y) \cup V(G_z)$. Since for any $\ell \in \{x, y, z\}$, $D \cap V(G_{\ell})$ contains no P_3 by Claim 11(ii) and Observation 5, it follows that $D \cap \{f_c^x, f_c^y, f_c^z\} \neq \emptyset$; and since $v_c \in D$ by Claim 11(i) and v_c should be dominated, in fact $|D \cap \{f_c^x, f_c^y, f_c^z\}| = 1$. Assume without loss of generality that $f_c^x \in D$. Then $q_c^x \in D$: indeed, if $q_c^x \notin D$ then $D \cap V(G_c^F)$ contains no P_3 ; and since for every $\ell \in \{x, y, z\}, D \cap V(G_\ell)$ contains no P_3 by Claim 11(ii) and Observation 5, clearly $D \cap (V(G_c^F) \cup V(G_x) \cup V(G_y) \cup V(G_z))$ then contains no P_4 , $K_{1,3}$ or $2P_3$. Similarly, $P_{x,F}^c(2) \in D$: indeed, if $P_{x,F}^c(2) \notin D$ then since for any $\ell \in \{x, y, z\}, D \cap V(G_{\ell})$ contains no P_3 , clearly $D \cap (V(G_c^F) \cup V(G_x) \cup V(G_y) \cup V(G_z))$ then contains no P_4 , $K_{1,3}$ or $2P_3$. It then follows from Claim 12(ii) that $F_x \notin D$ which, by Claim 11(ii), implies that $T_x \notin D$ and so, $P_{x,T}^c(1) \notin D$ by Claim 12(i). Since t_c^x should be dominated, it follows that $p_c^x \in D$ which implies that $|D \cap \{p_c^y, p_c^z\}| \leq 1$, as $|D \cap V(G_c^T)| = 2$. Since $D \cap \{q_c^y, q_c^z\} = \emptyset$ by the above, there then exists $v \in \{y, z\}$ such that $p_c^v, q_c^v \notin D$; but $|D \cap V(G_v)| = 14$, a contradiction to Claim 14. Thus, $|D \cap V(G_c^F)| < 3$.

Second, suppose that $|D \cap V(G_c^T)| = 3$. Since then, $|D \cap V(G_c^F)| = 2$, it follows from Observation 6 that $D \cap \{f_c^x, f_c^y, f_c^z\} = \emptyset$ and so, any P_4 , $K_{1,3}$ or $2P_3$ in D is in fact contained in $D \cap \{V(G_c^T) \cup V(G_x) \cup V(G_y) \cup V(G_z)$. But then $D \cap \{t_c^x, t_c^y, t_c^z\} \neq \emptyset$: indeed, if $D \cap \{t_c^x, t_c^y, t_c^z\} = \emptyset$, since for any $\ell \in \{x, y, z\}$, $D \cap V(G_\ell)$ contains no P_3 by Claim 11(ii) and Observation 5, clearly

 $\begin{array}{l} D\cap (V(G_c^T)\cup V(G_x)\cup V(G_y)\cup V(G_z) \text{ then contains no } P_4, K_{1,3} \text{ or } 2P_3. \text{ Assume without loss of generality that } t_c^x\in D. \text{ Then } p_c^x\in D: \text{ indeed, if } p_c^x\notin D \text{ then } D\cap V(G_c^T) \text{ contains no } P_3 \text{ and since for any } \ell\in\{x,y,z\}, \ D\cap V(G_\ell) \text{ contains no } P_3, \text{ clearly } D\cap (V(G_c^T)\cup V(G_x)\cup V(G_y)\cup V(G_z)) \text{ contains no } P_4, K_{1,3} \text{ or } 2P_3. \text{ Similarly, } P_{x,T}^c(1)\in D: \text{ if not, then since for any } \ell\in\{x,y,z\}, \ D\cap V(G_\ell) \text{ contains no } P_3, \text{ clearly } D\cap (V(G_c^T)\cup V(G_y)\cup V(G_z)) \text{ contains no } P_4, K_{1,3} \text{ or } 2P_3. \text{ Similarly, } P_{x,T}^c(1)\in D: \text{ if not, then since for any } \ell\in\{x,y,z\}, \ D\cap V(G_\ell) \text{ contains no } P_3, \text{ clearly } D\cap (V(G_c^T)\cup V(G_x)\cup V(G_y)) \text{ or } V(G_z) \text{ contains no } P_4, K_{1,3} \text{ or } 2P_3. \text{ If then follows from Claim 12(i) that } T_x\in D \text{ which, by Claim 11(ii), implies that } F_x\notin D \text{ and so, } P_{x,F}^c(2)\notin D \text{ by Claim 12(ii). Since } f_c^x \text{ should be dominated, necessarily } q_c^x\in D \text{ and so, } D\cap \{q_c^y,q_c^z\}=\varnothing \text{ by Claim 11(i). Since } |D\cap \{p_c^y,p_c^z\}| \leq 1 \text{ by the above, it follows that there exists } v\in\{y,z\} \text{ such that } p_c^v,q_c^v\notin D; \text{ but } |D\cap V(G_v)| = 14, \text{ a contradiction to Claim 14 which concludes the proof.} \end{array}$

Now by Claims 13 and 16, Φ is satisfiable if and only if $ct_{\gamma_t}(G) = 3$; and since G is easily seen to be $K_{1,3}$ -free, the lemma follows.

3.3 Proof of Theorem 1.3

Let H be a graph. If H contains a cycle then CONTRACTION NUMBER $(\gamma_t, 2)$ is NP-hard by Lemma 3.9. Assume henceforth that H is a forest. If H contains a vertex of degree at least three then CONTRACTION NUMBER $(\gamma_t, 2)$ is NP-hard on H-free graphs by Lemma 3.4. Suppose therefore that H is a linear forest. If H contains a connected component on at least six vertices then CONTRACTION NUMBER $(\gamma_t, 2)$ is coNP-hard on H-free graphs by Lemma 3.5. Thus we may assume that every connected component of H has size at most five. If H contains at least two connected component on at least four vertices then CONTRACTION NUMBER $(\gamma_t, 2)$ is NPhard on H-free graphs by Lemma 3.6. Assume therefore that H contains at most one connected component of size at least four and at most five. Now if H contains a P_5 and at least one other connected component on at least two vertices then CONTRACTION NUMBER $(\gamma_t, 2)$ is coNP-hard on H-free graphs by Lemma 3.7; and if H contains a P_5 and every other connected component has size one then CONTRACTION NUMBER $(\gamma_t, 2)$ is polynomial-time solvable on H-free graphs by ??. Otherwise H contains at most one connected component of size at most four while every other connected component has size at most three in which case CONTRACTION NUMBER $(\gamma_t, 2)$ is polynomial-time solvable on H-free graphs by Lemma 3.3.

4 Semitotal Domination

In this section, we consider the CONTRACTION NUMBER (γ_{t2}, k) problem for k = 2, 3. In Section 4.1, we cover those polynomial-time solvable cases and in Section 4.2, we examine hard cases. The proof of Theorem 1.4 can be found in Section 4.3.

4.1 Polynomial-time solvable cases

The algorithms for CONTRACTION NUMBER(γ_{t2} ,2) and CONTRACTION NUMBER(γ_{t2} ,3) outlined thereafter will rely on the following key result.

Lemma 4.1. Let H be a graph. If 2-EDGE CONTRACTION (γ_{t2}) is polynomial-time solvable on H-free graphs then it is polynomial-time solvable on $(H + P_3)$ -free graphs.

Proof. Let G be an $(H + P_3)$ -free graph. We aim to show that if G contains an induced H and G is a NO-instance for 2-EDGE CONTRACTION (γ_{t2}) , then $\gamma_{t2}(G)$ is bounded by some function f of |V(H)| (and |V(H)| only). Assuming for now that this claim is correct, the following algorithm solves the 2-EDGE CONTRACTION (γ_{t2}) problem on $(H + P_3)$ -free graphs.

- 1. If G contains no induced H then use the algorithm for H-free graphs to determine whether G is a YES-instance for 2-EDGE CONTRACTION(γ_{t2}) or not.
- 2. Check whether there exists an SD set of G of size at most f(|V(H)|).
 - 2.1 If the answer is no then output YES.
 - 2.2 Check whether there exists a minimum SD set of G containing a friendly triple, or an SD set of G of size $\gamma_{t2}(G) + 1$ containing an ST-configuration (see Theorem 1.7) and output the answer accordingly.

Now observe that checking whether G is H-free can be done in time $|V(G)|^{O(|V(H)|)}$ and that step 2 can be done in time $|V(G)|^{O(f(|V(H)|)}$ by simple brute force. Since 2-EDGE CONTRACTION(γ_{t2}) is polynomial-time solvable on H-free graphs by assumption, the above algorithm indeed runs in polynomial time (for fixed H). We next show that f(|V(H)|) = 8|V(H)|.

Assume henceforth that G contains an induced H. Let $A \subseteq V(G)$ be a set of vertices such that G[A] is isomorphic to H, let $B \subseteq V(G) \setminus A$ be the set of vertices at distance one from A and let $C = V(G) \setminus (A \cup B)$. Note that since G is $(H + P_3)$ -free, C is a disjoint union of cliques. In the following, we denote by \mathcal{K} the set of maximal cliques in C.

Now observe that if G is a No-instance for 2-EDGE CONTRACTION(γ_{t2}), then no minimum SD set of G contains an edge by Lemma 2.2 and thus, given a minimum SD set D of G, \mathcal{K} can be partitioned into two sets: $\mathcal{K}_1^D = \{K \in \mathcal{K} \mid |D \cap V(K)| = 1\}$ and $\mathcal{K}_0^D = \{K \in \mathcal{K} \mid D \cap V(K) = \emptyset\}$.

Claim 17. If G is a NO-instance for 2-EDGE CONTRACTION(γ_{t2}) and D is a minimum SD set of G then

$$|(D \cap N[\mathcal{K}_1^D]) \cup w_D(D \cap N[\mathcal{K}_1^D])| \le 4|A|.$$

Proof. Assume that G is a NO-instance for 2-EDGE CONTRACTION(γ_{t2}) and let D be a minimum SD set of G. For every $K \in \mathcal{K}_1^D$, denote by $x_K \in V(K) \cap D$ the unique vertex in K belonging to D. Observe that for every $K \in \mathcal{K}_1^D$, $N(x_K) \cap B \neq \emptyset$: indeed, if $N(x_K) \cap B = \emptyset$ for some $K \in \mathcal{K}_1^D$, then there must exist $y \in V(K)$ such that y is adjacent to the unique witness z for x_K (recall that by Lemma 2.2, $|w_D(x_K)| = 1$ and x_K is at distance exactly two from its witness); but then $(D \setminus \{x_K\}) \cup \{y\}$ is a minimum SD set of G containing the edge yz, a contradiction to Lemma 2.2. We now contend that there exist no two cliques $K, K' \in \mathcal{K}_1^D$ such that $d(x_K, x_{K'}) = 2$. Indeed, suppose to the contrary that there exist two such cliques $K, K' \in \mathcal{K}_1^D$. Let $z \in B$ be a common neighbour of x_K and $x_{K'}$, and let $t \in A$ be a neighbour of z. Then $z \notin D(x_K, z, x_{K'})$ would otherwise be a friendly triple) which implies, in particular, that t is not dominated by z. Since t should be dominated nonetheless, it follows that either $t \in D$ in which case $D \cup \{z\}$ contains the $O_6 x_K, z, x_{K'}, u$, a contradiction in both cases to Theorem 1.7.

It follows that for any clique $K \in \mathcal{K}_1^D$, any witness for x_K can only belong to $A \cup B$: let us denote by $W_A = A \cap w_D(\{x_K \mid K \in \mathcal{K}_1^D\})$ and by $W_B = B \cap w_D(\{x_K \mid K \in \mathcal{K}_1^D\})$. Further note that since by Lemma 2.2, $|w_D(x)| = 1$ for any $x \in D$,

$$|W_B| + |W_A| = |w_D(\{x_K \mid K \in \mathcal{K}_1^D\})| = |\mathcal{K}_1^D|.$$

Now if there exist two vertices $u, v \in W_B$ such that u and v have a common neighbour in A, then $d(u, v) \leq 2$, a contradiction to Lemma 2.2 as, by construction, u and v both have a witness in $\{x_K \mid K \in \mathcal{K}_1^D\}$. Thus, no two vertices in W_B have a common neighbour in A, which implies that $|W_B| \leq |A|$. Since $|W_A| \leq |A|$, it follows that $|\{w_K \mid K \in \mathcal{K}_1^D\}| \leq 2|A|$. Now note that $|(D \cap N[\mathcal{K}_1^D]) \cup w_D(D \cap N[\mathcal{K}_1^D])| = |\mathcal{K}_1^D| + |w_D(\{x_K \mid K \in \mathcal{K}_1^D\})|$: indeed, for every $K \in \mathcal{K}_1^D$, any vertex in $D \cap N[K]$ is at distance at most two from x_K , and by Lemma 2.2, $|w_D(x)| = 1$ for every $x \in D$. Thus, the lemma follows.

Claim 18. If G is a NO-instance for 2-EDGE CONTRACTION(γ_{t2}) and D is a minimum SD set of G, then

$$|(D \cap N[\mathcal{K}_0^D]) \cup w_D(D \cap N[\mathcal{K}_0^D])| \le 2|A|.$$

Proof. Assume that G is a NO-instance for 2-EDGE CONTRACTION (γ_{t2}) and let D be a minimum SD set of G. Since for every $K \in \mathcal{K}_0^D$, $D \cap V(K) = \emptyset$ by definition, necessarily $D \cap N[\mathcal{K}_0^D] = D \cap N(\mathcal{K}_0^D) \subseteq B$; in particular, every vertex in $D \cap N(\mathcal{K}_0^D)$ has at least one neighbour in A. Since no two vertices $x, y \in D \cap N(\mathcal{K}_0^D)$ such that $y \notin w_D(x)$, have a common neighbour in A, and for every $x \in D$, $|w_D(x)| = 1$ by Lemma 2.2, denoting by

$$D_1 = \{x \mid x \in D \cap N(\mathcal{K}_0^D) \text{ and } w_D(x) \cap (D \cap N(\mathcal{K}_0^D)) = \emptyset\}$$

and by

$$D_2 = \{\{x, y\} \mid x, y \in D \cap N(\mathcal{K}_0^D) \text{ and } y \in w_D(x)\}$$

it follows that $|D_1| + |D_2| \le |A|$. But

$$|(D \cap N(\mathcal{K}_0^D)) \cup w_D(D \cap N(\mathcal{K}_0^D))| = 2|D_1| + 2|D_2|$$

and thus, the upperbound follows.

To conclude, assume that G is a NO-instance for 2-EDGE CONTRACTION(γ_{t2}) and let D be a minimum SD set of G. Then $D' = D \setminus ((D \cap N[\mathcal{K}]) \cup (w_D(D \cap N[\mathcal{K}])))$ has size at most 2|A|: indeed, since D' dominates only vertices in $A \cup B$ and no vertex in $D \setminus D'$ is witnessed by a vertex D' by Lemma 2.2, if |D'| > 2|A| then $(D \setminus D') \cup (A \cup N_A)$ where N_A contains one neighbour for each vertex in A, is an SD set of G of size strictly smaller than that of D, a contradiction to the minimality of D. Since

$$|(D \cap N[\mathcal{K}]) \cup (w_D(D \cap N[\mathcal{K}]))| \le \sum_{i \in \{0,1\}} |(D \cap N[\mathcal{K}_i^D]) \cup (w_D(D \cap N[\mathcal{K}_i^D]))| \le 6|A|$$

by Claims 17 and 18, we conclude that $\gamma_{t2}(G) = |D| \le 8|A| = 8|V(H)|$, as claimed.

٦

Lemma 4.2. If G is a P_8 -free graph then $ct_{\gamma_{t2}}(G) \leq 2$.

Proof. Let G be a P_8 -free graph and let D be a minimum SD set of G. If D contains an edge or there exists $x \in D$ such that $|w_D(x)| \ge 2$, then $ct_{\gamma_{t2}}(G) \le 2$ by Lemma 2.2. Thus, assume that D is an independent set and for every $x \in D$, $|w_D(x)| = 1$. Let $u, v \in D$ be two vertices at distance two and further let $w \in D \setminus \{u, v\}$ be a closest vertex from $\{u, v\}$, that is, $d(w, \{u, v\}) =$ $\min_{x \in D \setminus \{u,v\}} d(x, \{u,v\})$. Then $d(w, \{u,v\}) > 2$ by assumption, and since $d(w, \{u,v\}) \leq 3$ as shown in the proof of Lemma 2.2, in fact $d(w, \{u, v\}) = 3$. Now assume without loss of generality that $d(w, \{u, v\}) = d(w, v)$, and let $t \in D$ be the witness for w. Note that if there exists $a \in A$ $N(u) \cap N(v)$ and $b \in N(w)$ such that $at \in E(G)$, then $D \cup \{a\}$ contains the $O_6(u, a, v, w)$ and thus, $ct_{\gamma_{12}}(G) \leq 2$. Assume henceforth that no common neighbour of u and v is adjacent to a neighbour of w (or t), and that, symmetrically, no common neighbour of w and t is adjacent to a neighbour of u (or v). Let P = vxyw be a shortest path from v to w. Then by the aforementioned assumption, u is nonadjacent to x, and t is nonadjacent to y (also note that u is nonadjacent to y, and t is nonadjacent to x, by minimality of the distance between v and w). Let a be a common neighbour of u and v, and let b be a common neighbour of w and t. Then both a and x are nonadjacent to b by assumption; and similarly, y is nonadjacent to a. It follows that $ax \in E(G)$ or $by \in E(G)$ for otherwise, uavxywbt would induce a P_8 ; we next distinguish two cases depending on these adjacencies.

Suppose first that $ax, by \in E(G)$. Observe that if every private neighbour of t is adjacent to b or y, then $(D \setminus \{t\}) \cup \{b, y\}$ is an SD set of G containing the O_4 b, w, y, v and thus, $ct_{\gamma_{t2}}(G) \leq 2$ by Lemma 2.2. Assume therefore that t has a private neighbour c which is nonadjacent to both b and y, and that, symmetrically, u has a private neighbour d which is nonadjacent to both a and x. Let us show that then G contains an induced P_8 . Indeed, since by assumption d is nonadjacent to b, and c is nonadjacent to a, it must be that $dc \in E(G)$, $dy \in E(G)$ or $cx \in E(G)$ for otherwise, dyaxybtc would induce a P_8 . However, if $dc \in E(G)$ then vaudctbw induces a P_8 . Thus, $dy \in E(G)$ or $cx \in E(G)$, say the latter holds without loss of generality, in which case wbtcxaud induces a P_8 .

Second, suppose that either a is nonadjacent to x, or b is nonadjacent to y, say $ax \in E(G)$ and $by \notin E(G)$ (the other case is symmetric). As previously, we show that if t has a private neighbour c which is nonadjacent to both b and y, and that u has a private neighbour d which is nonadjacent to both a and x, then G contains an induced P_8 (observe that if this does not hold, then $ct_{\gamma_{t2}}(G) \leq 2$ as mentioned above). Since c is nonadjacent to a by assumption, c must be adjacent to x for otherwise, ctbwyxau would induce a P_8 . But then, c must be adjacent to d for otherwise, wbtcxaud induces a P_8 ; however, if $dc \in E(G)$ then wbtcduav induces a P_8 , a contradiction which concludes the proof.

Lemma 4.3. If G is a 2P₄-free graph then $ct_{\gamma_{t2}}(G) \leq 2$.

Proof. The proof is similar to that of Lemma 4.2. Let G be a $2P_4$ -free graph and let D be a minimum SD set of G. If D contains an edge or there exists $x \in D$ such that $|w_D(x)| \ge 2$, then $ct_{\gamma_{t2}}(G) \le 2$ by Lemma 2.2. Thus, assume that D is an independent set and for every $x \in D$,

 $|w_D(x)| = 1$. Let $u, v \in D$ be two vertices at distance two and further let $w \in D \setminus \{u, v\}$ be a closest vertex from $\{u, v\}$, that is, $d(w, \{u, v\}) = \min_{x \in D \setminus \{u, v\}} d(x, \{u, v\})$. Then $d(w, \{u, v\}) > 2$ by assumption, and since $d(w, \{u, v\}) \leq 3$ as shown in the proof of Lemma 2.2, in fact $d(w, \{u, v\}) = 3$. Now assume without loss of generality that $d(w, \{u, v\}) = d(w, v)$, and let $t \in D$ be the witness for w. Note that if there exists $a \in N(u) \cap N(v)$ and $b \in N(w)$ such that $at \in E(G)$, then $D \cup \{a\}$ contains the O_6 u, a, v, w and thus, $ct_{\gamma_{t2}}(G) \leq 2$ by Theorem 1.7. Assume henceforth that no common neighbour of u and v is adjacent to a neighbour of w (or t), and that, symmetrically, no common neighbour of w and t is adjacent to a neighbour of u (or v). Let P = vxyw be a shortest path from v to w. Then by the aforementioned assumption, u is nonadjacent to x, and t is nonadjacent to y (also note that u is nonadjacent to y, and t is nonadjacent to x, by minimality of the distance between v and w). Let a be a common neighbour of u and v, and let b be a common neighbour of w and t. Then a and x are nonadjacent to b by assumption; and similarly, y is nonadjacent to a. Now if b is nonadjacent to y, then every private neighbour p of u must be adjacent to a or y for otherwise, puav and ywbt would induce a $2P_4$ (note that p is nonadjacent to b by assumption); but then, $(D \setminus \{u\}) \cup \{a, y\}$ is an SD set of G containing the $O_7 a, v, y, w$ and thus, $ct_{\gamma_{t2}}(G) \leq 2$ by Theorem 1.7. Assume therefore that b is adjacent to y. If every private neighbour of u is adjacent to a, then $(D \setminus \{u\}) \cup \{a\}$ is a minimum SD set of G containing an edge and thus, $ct_{\gamma_{t2}}(G) \leq 2$ by Lemma 2.2. Thus, we may assume that u has a private neighbour p nonadjacent to a. Then every private neighbour p' of t must be adjacent to b or p for otherwise, puav and wbtp' would induce a $2P_4$ (note that p is nonadjacent to b, and p' is nonadjacent to a by assumption). By symmetry, we conclude that every private neighbour of w is adjacent to por b. Now if, in fact, no private neighbour of w is adjacent to p, then $D' = (D \setminus \{w\}) \cup \{b\}$ is a minimum SD set of G, as every private neighbour of w w.r.t. D is then adjacent to b; but D' contains an edge and thus, $ct_{\gamma_{t2}}(G) \leq 2$ by Lemma 2.2. Thus, we may assume that at least one private neighbour of w is adjacent to p; in particular, d(w,p) = 2. But then, $(D \setminus \{t\}) \cup \{p,b\}$ is an SD of G containing the O_7 b, w, p, u and thus, $ct_{\gamma_{t2}}(G) \leq 2$ by Theorem 1.7.

Since for any graph G, G is a YES-instance for 2-EDGE CONTRACTION(γ_{t2}) if and only if G is a NO-instance for CONTRACTION NUMBER(γ_{t2} ,3), the following ensues from Lemmas 4.1, 4.2 and 4.3.

Corollary 4.4. CONTRACTION NUMBER $(\gamma_{t2},3)$ is polynomial-time solvable on *H*-free graphs if $H \subseteq_i P_8 + kP_3$ for some $k \ge 0$, or $H \subseteq 2P_4 + kP_3$ for some $k \ge 0$.

Lemma 4.5. CONTRACTION NUMBER $(\gamma_{t2}, 2)$ is polynomial-time solvable on *H*-free graphs if $H \subseteq_i P_5 + tK_1$ for some $t \ge 0$, or $H \subseteq_i P_3 + tP_2$ for some $t \ge 0$.

Proof. Assume that $H \subseteq_i P_3 + tP_2$ for some $t \ge 0$ (the case where $H \subseteq_i P_5 + tK_1$ for some $t \ge 0$ is handled similarly) and let G be an H-free graph. Since H is a fortiori an induced subgraph of $P_8 + tP_3$, we may use the polynomial-time algorithm for $(P_8 + tP_3)$ -free graphs given by Corollary 4.4 to determine whether G is a YES-instance for CONTRACTION NUMBER $(\gamma_{t2},3)$ or not. If the answer is yes then we output NO; otherwise, we use the polynomial-time algorithm for $(P_3 + tP_2)$ -free graphs given by Theorem 1.8 to determine whether G is a YES-instance for CONTRACTION NUMBER $(\gamma_{t2},1)$ or not, and output the answer accordingly.

4.2 Hardness results

In this section, we show that CONTRACTION NUMBER $(\gamma_{t2}, 2)$ and CONTRACTION NUMBER $(\gamma_{t2}, 3)$ are NP-hard on a number of monogenic graph classes. We first consider the case k = 2.

Firstly, since for any graph G such that $ct_{\gamma_{t2}}(G) \leq 2$, G is a YES-instance for CONTRACTION NUMBER $(\gamma_{t2}, 1)$ if and only if G is a NO-instance for CONTRACTION NUMBER $(\gamma_{t2}, 2)$, the following ensues from Theorem 1.8 and Lemma 4.2.

Lemma 4.6. CONTRACTION NUMBER $(\gamma_{t2}, 2)$ is (co)NP-hard on P_6 -free graphs, $2P_3$ -free graphs and $(P_4 + P_2)$ -free graphs.

Lemma 4.7. CONTRACTION NUMBER $(\gamma_{t2}, 2)$ is NP-hard in $K_{1,3}$ -free graphs.

Proof. We use the same construction as in [10, Theorem 7]. More precisely, we reduce from POSITIVE EXACTLY 3-BOUNDED 1-IN-3 3-SAT (see Section 2 for a precise definition of this




Figure 8: The gadget G_x for a variable $x \in X$ contained in clauses c, c' and c'' (a rectangle indicates that the corresponding set of vertices induces a clique).




Figure 9: The gadget G_c for a clause $c = x \vee y \vee z$ is the disjoint union of G_c^T and G_c^F .

problem): given an instance Φ of this problem, with variable set X and clause set C, we construct an instance G of CONTRACTION NUMBER $(\gamma_{t2}, 2)$ as follows. For every variable $x \in X$ contained in clauses c, c' and c'', we introduce the gadget G_x depicted in Figure 8 (where a rectangle indicates that the corresponding set of vertices is a clique). For every clause $c \in C$ containing variables x, yand z, we introduce the gadget G_c , which is the disjoint union of the graph G_c^T and the graph G_c^F depicted in Figure 9, and further add the following edges.

- For every $p \in \{x, y, z\}$, we connect $P_{p,T}^c(2)$ to f_c^{ab} if and only if $p \in \{a, b\}$. For every $p \in \{x, y, z\}$, we connect $P_{p,T}^c(1)$ to t_c^p and further connect $P_{p,T}^c(1)$ to w_c^{ab} if and only if $p \in \{a, b\}$.

We let G be the resulting graph. Let us show that Φ is satisfiable if and only if $ct_{\gamma_{t2}}(G) = 2$. To do so, we will rely on the following key result shown in [10]

Claim 19 ([10]). The following statements are equivalent.

- (i) Φ is satisfiable.
- (*ii*) $\gamma_{t2}(G) = 14|X| + |C|.$
- (*iii*) $ct_{\gamma_{t2}}(G) > 1$.

Observe first that if $ct_{\gamma_{t2}}(G) = 2$ then Φ is satisfiable by Claim 19. Assume, conversely, that Φ is satisfiable and consider a truth assignment satisfying Φ . Let us show how to construct a minimum SD set D of G given this assignment. For every variable $x \in X$ appearing in clauses $c, c' \text{ and } c'', \text{ if } x \text{ is true, we include } \{v_x, T_x\} \cup \{P_{x,R}^p(1), P_{x,R}^p(4) \mid R \in \{T, F\} \text{ and } p \in \{c, c', c''\}\}$ in D; otherwise, we include $\{v_x, F_x\} \cup \{P_{x,R}^p(2), P_{x,R}^p(4) \mid R \in \{T, F\} \text{ and } p \in \{c, c', c''\}\}$ in D. For every clause $c \in C$ containing variables x, y and z, exactly one variable is set to true, say xwithout loss of generality, in which case we add t_c^y to D (note that w_c^{xz}, t_c^x and w_c^{xy} are already dominated by $P_{x,T}^{c}(1)$, which also serves as a witness for t_{c}^{y}). It is not difficult to see that the constructed set D is indeed an SD set of G and since |D| = 14|X| + |C|, we conclude by Claim 19 that D is minimum. Now consider a clause $c \in C$ containing variables x, y and z, and assume without loss of generality that x is true and $t_c^y \in D$. Then $D \cup \{t_c^x\}$ is an SD set of G of size $\gamma_{t2}(G) + 1$ containing the $O_4 t_c^y, t_c^x, P_{x,T}^c(1), P_{x,T}^c(4)$ and thus, $ct_{\gamma_{t2}}(G) \le 2$ by Theorem 1.7. But $ct_{\gamma_{t2}}(G) > 1$ by Claim 19 and so, $ct_{\gamma_{t2}}(G) = 2$. Since G is $K_{1,3}$ -free, the lemma follows.

Lemma 4.8. CONTRACTION NUMBER $(\gamma_{t2}, 2)$ is NP-hard on $\{C_k \mid k \geq 5\}$ -free graphs.

Proof. The reduction is similar to that of Lemma 3.6. More precisely, we reduce from 3-SAT: given an instance Φ of this problem, with variable set X and clause set C, we construct an instance G of CONTRACTION NUMBER $(\gamma_{t2}, 2)$ as follows. For every variable $x \in X$, we introduce a long paw G_x on vertex set $\{G_x(1) = x, G_x(2) = \overline{x}, G_x(3) = u_x, G_x(4) = v_x, G_x(5) = w_x\}$ and refer to the vertices x and \overline{x} as *literal vertices*. For every clause $c \in C$, we introduce a *clause vertex*, denoted by c, and add an edge between c and every literal vertex whose corresponding literal appears in the clause c. Finally, we add an edge between every two clauses vertices so that the set of clause vertices induces a clique, denoted by K in the following. We let G be the resulting graph. We next show that G is a YES-instance for CONTRACTION NUMBER $(\gamma_{t2}, 2)$ if and only if Φ is satisfiable through a series of claims. Observe first that since for any variable $x \in X$, the vertex w_x should be dominated and any witness for a vertex dominating w_x can only belong to G_x , the following holds.

Observation 7. For any SD set D of G and any variable $x \in X$, $|D \cap V(G_x)| \ge 2$.

Claim 20. Φ is satisfiable if and only if $\gamma_{t2}(G) = 2|X|$.

Proof. Assume first that Φ is satisfiable and consider a truth assignment satisfying Φ . We construct an SD set D of G as follows. For every variable $x \in X$, if x is true then we include v_x and x to D; otherwise, we include v_x and \overline{x} to D. It is easy to see that the constructed set D is indeed an SD set of G and since |D| = 2|X|, we conclude by Observation 7 that D is minimum.

Conversely assume that $\gamma_{t2}(G) = 2|X|$ and let D be a minimum SD set of G. Then by Observation 7, $D \cap V(K) = \emptyset$ and thus, every clause vertex must have a neighbouring literal vertex in D. Therefore, the truth assignment obtained by setting a variable x to true when $x \in D$ and to false otherwise, satisfies Φ .

Claim 21. $ct_{\gamma_{t2}}(G) = 2$ if and only if $\gamma_{t2}(G) = 2|X|$.

Proof. Assume first that $ct_{\gamma_{t2}}(G) = 2$ and consider a minimum SD set D of G. If there exists a variable $x \in X$ such that $|D \cap V(G_x)| \geq 3$, then $D \cap V(G_x)$ contains a friendly triple, a contradiction to Theorem 1.7. Thus, $|D \cap V(G_x)| \leq 2$ for every variable $x \in X$, and we conclude by Observation 7 that, in fact, equality holds. Now suppose that $D \cap V(K) \neq \emptyset$, say $c \in D \cap V(K)$, and let $x \in X$ be a variable occurring in c, say x appears positive in c without loss of generality. Then the set $(D \setminus V(G_x)) \cup \{x, v_x\}$ is a minimum SD set of G containing the friendly triple c, x, v_x , a contradiction to Theorem 1.7. Thus, $D \cap V(K) = \emptyset$ and so, $\gamma_{t2}(G) = |D| = 2|X|$.

Conversely, assume that $\gamma_{t2}(G) = 2|X|$ and let D be a minimum SD set of G. Then by Observation 7, $D \cap V(K) = \emptyset$ and $|D \cap V(G_x)| = 2$ for every variable $x \in X$. Since for every variable $x \in X$, w_x should be dominated, it follows that if for some variable $x \in X$, $D \cap V(G_x)$ contains an edge e, then $D \cap \{x, \overline{x}\} = \emptyset$; but then, $d(e, D \setminus e) \ge 3$ and thus, e cannot be part of a friendly triple. Therefore, D contains no friendly triple. Consider now a clause $c \in C$ and let $x, y \in X$ be two variables occurring in c, say x and y both appear positive in c without loss of generality. Then the set $(D \setminus (V(G_x) \cup V(G_y))) \cup \{c, x, v_x, y, v_y\}$ is an SD set of G of size $\gamma_{t2}(G) + 1$ containing the O_4 x, c, y, v_y and thus, $ct_{\gamma_{t2}}(G) = 2$ by Theorem 1.7.

Now Claims 20 and 21, Φ is satisfiable if and only if $ct_{\gamma_{t2}}(G) = 2$ if and only if Φ is satisfiable. Since G is easily seen to contain no induced cycle of length at least five, the lemma follows.

Lemma 4.9. For every $\ell \in \{3, 4\}$, CONTRACTION NUMBER $(\gamma_{t2}, 2)$ is NP-hard on C_{ℓ} -free graphs.

Proof. We reduce from POSITIVE NOT-ALL-EQUAL 3-SAT which is a NP-complete variant [23] of the 3-SAT problem where given a formula in which all literals are positive, the problem is to determine whether there exists a truth assignment such that in no clause, all three literals have the same truth value. Given an instance Φ of this problem, with variable set X and clause set C, we construct two instances G_3 and G_4 of CONTRACTION NUMBER($\gamma_{t2}, 2$), one C_3 -free and the other C_4 -free, respectively. For both graphs, we start as follows. For every variable $x \in X$, we introduce the gadget G_x depicted in Figure 10, which has two distinguished truth vertices T_x and F_x . For every clause $c \in C$ containing variables x, y and z, we introduce two clause vertices c and \overline{c} , and add for every $v \in \{x, y, z\}$, an edge between c and T_v , and an edge between \overline{c} and F_v . We




Figure 10: The variable gadget G_x .

denote by $V_C = \{c \mid c \in C\}$ the set of *positive* clause vertices, and by $V_{\overline{C}} = \{\overline{c} \mid c \in C\}$ the set of *negated* clause vertices. This concludes the construction of G_3 . For the graph G_4 , we further add edges so that V_C is a clique and $V_{\overline{C}}$ is a clique.

We next show that for every $\ell \in \{3,4\}$, $ct_{\gamma_{t2}}(G_\ell) = 2$ if and only if Φ is satisfiable through a series of claims. Observe first that since for every variable $x \in X$, the vertex r_x should be dominated and any witness for a vertex dominating r_x can only belong to G_x , the following holds.

Observation 8. Let $\ell \in \{3,4\}$. Then for every SD set D of G_{ℓ} and every variable $x \in X$, $|D \cap V(G_x)| \ge 2$.

Claim 22. Let $\ell \in \{3,4\}$. Then Φ is satisfiable if and only if $\gamma_{t2}(G_{\ell}) = 2|X|$.

Proof. Assume first that Φ is satisfiable and consider a truth assignment satisfying Φ . We construct an SD set D for G_{ℓ} . For every variable $x \in X$, if x is true, then we include w_x and T_x in D; otherwise, we include w_x and F_x in D. It is not difficult to see that the constructed set D is indeed an SD set for both G_{ℓ} and since |D| = 2|X|, we conclude by Observation 8 that D is minimum.

Conversely, assume that $\gamma_{t2}(G_{\ell}) = 2|X|$ and let D be a minimum SD set of G_{ℓ} . Then by Observation 8, $D \cap V_C = D \cap V_{\overline{C}} = \emptyset$ and thus, every (positive or negated) clause vertex must have a neighbouring truth vertex in D. Therefore, the truth assignment obtained by setting a variable x to true when $T_x \in D$ and to false otherwise, satisfies Φ .

Claim 23. Let $\ell \in \{3, 4\}$. Then $ct_{\gamma_{t2}}(G_{\ell}) = 2$ if and only if $\gamma_{t2}(G_{\ell}) = 2|X|$.


Proof. Assume that $ct_{\gamma_{t2}}(G_{\ell}) = 2$ and let D be a minimum SD set of G_{ℓ} . If there exists a variable $x \in X$ such that $|D \cap V(G_x)| \geq 3$, then $D \cap V(G_x)$ contains a friendly triple, a contradiction to Theorem 1.7. Thus, $|D \cap V(G_x)| \leq 2$ for every variable $x \in X$ and we conclude by Observation 8 that, in fact, equality holds. Furthermore, we may assume that for every variable $x \in X$, $D \cap \{F_x, T_x\} \neq \emptyset$ and $w_x \in D$, for it otherwise suffices to consider e.g. $(D \setminus V(G_x)) \cup \{F_x, w_x\}$ in place of D. Now suppose to the contrary that $D \cap V_C \neq \emptyset$. We distinguish cases depending on the value of ℓ .


Case 1. $\ell = 3$. Let $c \in V_C$ be a vertex in D and let $x, y, z \in X$ be the three variables occurring in c. If there exists $v \in \{x, y, z\}$ such that $T_v \in D$, then c, T_v, w_v is a friendly triple, a contradiction to Theorem 1.7. Thus, for every $v \in \{x, y, z\}$, $T_v \notin D$ and so, $F_v \in D$ by the aforementioned assumption. But then, $(D \setminus \{c\}) \cup \{T_x\}$ is a minimum SD of G_3 (recall that for $\ell = 3$, V_C is an independent set) containing the friendly triple F_x, T_x, w_x , a contradiction to Theorem 1.7.

Case 2. $\ell = 4$. Then $|D \cap V_C| = 1$: indeed, if there exist $c, c' \in D \cap V_C$ then $d(\{c, c'\}, D \setminus \{c, c'\}) \leq 2$ (recall that by assumption, $D \cap \{t_v, F_v\} \neq \emptyset$ for every variable $v \in X$) and so, D contains a friendly triple, a contradiction to Theorem 1.7. Now if there exists a variable $x \in X$ such that $T_x \in D$ then letting $c \in C$ be a clause containing x, the set $(D \setminus V_C) \cup \{c\}$ is a minimum SD set of G_4 containing the friendly triple c, T_x, w_x , a contradiction to Theorem 1.7. It follows that for every variable $x \in X$, $F_x \in D$; but then, it suffices to consider, in place of D, the SD set $(D \setminus \{F_x\}) \cup \{T_x\}$ for some variable $x \in X$, and argue as previously.

Thus, in both cases, $D \cap V_C = \emptyset$; and by symmetry, we conclude that $D \cap V_{\overline{C}} = \emptyset$ as well. Therefore, $\gamma_{t2}(G_{\ell}) = |D| = 2|X|$.

Conversely, assume that $\gamma_{t2}(G_{\ell}) = 2|X|$ and let D be a minimum SD set of G_{ℓ} . Then by Observation 8, $D \cap V_C = D \cap V_{\overline{C}} = \emptyset$ and $|D \cap V(G_x)| = 2$ for every variable $x \in X$. Since for every variable $x \in X$, r_x should be dominated, it follows that if for some variable $x \in X$, $D \cap V(G_x)$ contains an edge e, then $D \cap \{F_x, T_x\} = \emptyset$; but then, $d(e, D \setminus e) \ge 3$ and thus, ecannot be part of a friendly triple. Therefore, D contains no friendly triple. Now let us assume, for the following, that for every variable $x \in X$, $D \cap \{F_x, T_x\} \ne \emptyset$ and $w_x \in D$ (if this does





(b) The graph G_c^F (the rectangle indicates that the corresponding set of vertices is a clique).

Figure 11: The clause gadget G_c for a clause $c = x \lor y \lor z$ is the disjoint union of G_c^T and G_c^F .

not hold for a variable $x \in X$, we may consider e.g. $(D \setminus V(G_x)) \cup \{F_x, w_x\}$ in place of D). Consider a clause $c \in C$ and let $x, y, z \in X$ be the three variables occurring in c. By assumption, either $|D \cap \{T_x, T_y, T_z\}| \ge 2$ or $|D \cap \{F_x, F_y, F_z\}| \ge 2$, say $F_x, F_y \in D$ without loss of generality. Then the set $D \cup \{\overline{c}\}$ is an SD set of G_ℓ of size $\gamma_{t2}(G) + 1$ containing the O_4 $F_x, \overline{c}, F_y, w_y$ and so, $ct_{\gamma_{t2}}(G) = 2$ by Theorem 1.7.

Now by Claims 22 and 23, we conclude that for every $\ell \in \{3,4\}$, $ct_{\gamma_{t2}}(G_\ell) = 2$ if and only if Φ is satisfiable. Since for every $\ell \in \{3.4\}$, G_ℓ is easily seen to contain no induced C_ℓ , the lemma follows.

The following now ensues from Lemmas 4.8 and 4.9.

Corollary 4.10. For every $\ell \geq 3$, CONTRACTION NUMBER $(\gamma_{t2}, 2)$ is NP-hard on C_{ℓ} -free graphs.

We next investigate the complexity of the CONTRACTION NUMBER(γ_{t2} ,3) problem on several monogenic graph classes.

Lemma 4.11. CONTRACTION NUMBER $(\gamma_{t2},3)$ is NP-hard on $K_{1,3}$ -free graphs.

Proof. We reduce from POSITIVE EXACTLY 3-BOUNDED 1-IN-3 3-SAT (see Section 2 for a precise definition of this problem): given an instance Φ of this problem, with variable set X and clause set C, we construct an instance G of CONTRACTION NUMBER $(\gamma_{t2},3)$ as follows. For every variable $x \in X$, we introduce the gadget G_x depicted in Figure 8 (it is the same gadget as the one used in the proof of Lemma 4.7). In the following, we denote by P_x the long paw induced by $\{T_x, F_x, u_x, v_x, w_x\}$ and we may refer to the vertices of P_x as $P_x(1), \ldots, P_x(5)$ where $P_x(1) = t_x$ and $P_x(2) = F_x$. For every clause $c \in C$ containing variables x, y and z, we introduce the gadget G_c which is the disjoint union of the graph G_c^T and the graph G_c^F depicted in Figure 11 (where a rectangle indicates that the corresponding set of vertices is a clique), and further add for every $v \in \{x, y, z\}$ an edge between t_c^v and $P_{v,T}^c(1)$, and an edge between f_c^v and $P_{v,F}^c(2)$. Note that $\{u_c^x, u_c^y, u_c^z, a_c\}$ and $\{v_c^x, v_c^y, v_c^z, b_c\}$ induce cliques; in particular, G_c^T is $K_{1,3}$ -free. We let G be the resulting graph. We next show that $ct_{\gamma_{t2}}(G) = 3$ if and only if Φ is satisfiable through a series of claims.

Claim 24. For any SD set D of G, the following hold.

- (i) For every variable $x \in X$ and every long paw P contained in G_x , $|D \cap V(P)| \ge 2$ and $D \cap \{P(4), P(5)\} \neq \emptyset$. In particular, $|D \cap V(G_x)| \ge 14$.
- (ii) For every clause $c \in C$ containing variables x, y and $z, |D \cap \{u_c, v_c, w_c, q_c^x, q_c^y, q_c^z\}| \ge 2$ and $\min(|D \cap \{u_c^x, u_c^y, u_c^z, a_c\}|, |D \cap \{v_c^x, v_c^y, v_c^z, b_c\}|) \ge 1$. In particular, $|D \cap V(G_c)| \ge 4$.

Proof. Let D be an SD set of G. Since in a long paw P, the vertex P(5) should be dominated and any witness for a vertex dominating P(5) can only belong to P, item (i) follows. To prove (ii), consider a clause $c \in C$ containing variables x, y and z. Since w_c must be dominated and any witness for a vertex dominating w_c can only belong to $\{u_c, v_c, w_c, q_c^x, q_c^y, q_c^z\}$, the first part of item (ii) follows. Now since a_c should be dominated, $D \cap \{u_c^x, u_c^y, u_c^z, a_c\} \neq \emptyset$; and similarly, since b_c should be dominated, $D \cap \{v_c^x, v_c^y, v_c^z, b_c\} \neq \emptyset$ which proves the second part of item (ii).

The following is an immediate consequence of Claim 24(i).

Observation 9. For every SD set D of G and every variable $x \in X$ contained in clauses c, c' and c'', if $|D \cap V(G_x)| = 14$ then $D \cap \{q_x^\ell, p_x^\ell \mid \ell \in \{c, c', c''\}\} = \emptyset$.

Claim 25. Let D be an SD set of G and let $c \in C$ be a clause containing variables x, y and z. If $|D \cap V(G_c^F)| = 2$ then $D \cap \{f_c^x, f_c^y, f_c^z\} = \emptyset$. Similarly, if $|D \cap V(G_c^T)| = 2$ then $D \cap \{t_c^x, t_c^y, t_c^z\} = \emptyset$.

 $\begin{array}{l} \textit{Proof. If } |D \cap V(G_c^F)| = 2 \text{ then by Claim 24(ii), } |D \cap V(G_c^F)| = |D \cap \{u_c, v_c, w_c, q_c^x, q_c^y, q_c^z\}| = 2 \text{ and thus, } D \cap \{f_c^x, f_c^y, f_c^z\} = \varnothing. \text{ Now if } |D \cap V(G_c^T)| = 2 \text{ then by Claim 24(ii), } |D \cap \{u_c^x, u_c^y, u_c^z, a_c\}| = |D \cap \{v_c^x, v_c^y, v_c^z, b_c\}| = 1 \text{ and thus, } D \cap \{t_c^x, t_c^y, t_c^z\} = \varnothing. \end{array}$

Claim 26. Let D be an SD set of G. If for some variable $x \in X$ contained in clauses c, c' and $c'', |D \cap V(G_x)| = 14$ then the following hold.

(i) If there exists $\ell \in \{c, c', c''\}$ such that $P_{x,T}^{\ell}(1) \in D$ then $T_x \in D$.

(ii) If there exists $\ell \in \{c, c', c''\}$ such that $P_{x,F}^{\ell}(2) \in D$ then $F_x \in D$. In particular, if $P_{x,T}^{\ell}(1) \in D$ for some $\ell \in \{c, c', c''\}$, then $D \cap \{P_{x,F}^{c}(2), P_{x,F}^{c'}(2), P_{x,F}^{c''}(2)\} = \emptyset$. Similarly, if $P_{x,F}^{\ell}(2) \in D$ for some $\ell \in \{c, c', c''\}$, then $D \cap \{P_{x,T}^{c}(1), P_{x,T}^{c''}(1), P_{x,T}^{c''}(1)\} = \emptyset$.

Proof. Let $x \in X$ be a variable such that $|D \cap V(G_x)| = 14$ and let $c, c', c'' \in C$ be the clauses in which x appears. Then by Observation 9, $D \cap \{p_x^\ell, q_x^\ell \mid \ell \in \{c, c', c''\}\} = \emptyset$ and by Claim 24(i), $|D \cap \{P(1), P(2)\}| \leq 1$ for every long paw P of G_x . Thus if $P_{x,T}^\ell(1) \in D$ for some $\ell \in \{c, c', c''\}$, then $T_x \in D$ as p_x^ℓ should be dominated; but then, $F_x \notin D$ which implies that $\{P_{x,F}^c(1), P_{x,F}^{c''}(1)\} \subseteq D$ (one of $q_x^c, q_x^{c'}$ and $q_x^{c''}$ would otherwise not be dominated) and so, $D \cap \{P_{x,F}^c(2), P_{x,F}^{c''}(2)\} = \emptyset$. Item (ii) follows by symmetry.

Claim 27. Φ is satisfiable if and only if $\gamma_{t2}(G) = 14|X| + 4|C|$.

Proof. Assume first that Φ is satisfiable and consider a truth assignment satisfying Φ . We construct an SD set D of G as follows. For every variable $x \in X$ contained in clauses c, c' and c'', if x is true then we include $\{v_x, T_x\} \cup \{P_{x,R}^{\ell}(1), P_{x,R}^{\ell}(4) \mid R \in \{T, F\} \text{ and } \ell \in \{c, c', c''\}\}$ in D; otherwise we include $\{v_x, F_x\} \cup \{P_{x,R}^{\ell}(2), P_{x,R}^{\ell}(4) \mid R \in \{T, F\} \text{ and } \ell \in \{c, c', c''\}\}$ in D. For every clause $c \in C$ containing variables x, y and z, exactly one variable is true, say x without loss of generality, in which case we include $\{w_c, q_c^x\} \cup \{v_c^y, u_c^z\}$ in D. Now it is not difficult to see that the constructed set D is indeed an SD set of G and since |D| = 14|X| + 4|C|, we conclude by Claim 24 that D is minimum.

Conversely, assume that $\gamma_{t2}(G) = 14|X| + 4|C|$ and let D be a minimum SD set of G. Consider a clause $c \in C$ and let $x, y, z \in X$ be the variables contained in c. Then $|D \cap \{q_c^x, q_c^y, q_c^z\}| \leq 1$: indeed, $D \cap \{w_c, v_c\} \neq \emptyset$ as w_c should be dominated, and $|D \cap \{u_c, v_c, w_c, q_c^x, q_c^y, q_c^z\}| = 2$ by Claim 24(i). Since $D \cap \{f_c^x, f_c^y, f_c^z\} = \emptyset$ by Claim 25, it follows that at least two vertices amongst f_c^x, f_c^y and f_c^z are not dominated by a vertex in G_c^F , say f_c^y and f_c^z without loss of generality. Then $P_{y,F}^c(2), P_{z,F}^c(2) \in D$ as f_c^y and f_c^z should be dominated, and so $F_y, F_z \in D$ by Claim 26(ii). It then follows from Claim 24(i) that $T_y, T_z \notin D$ which, by Claim 26(i), implies that $P_{y,T}^c(1), P_{z,T}^c(1) \notin D$; in particular, t_c^y and t_c^z must be dominated by vertices in G_c^T . But then, $D \cap \{t_c^x, v_c^x, u_c^x\} = \emptyset$: indeed, since $|D \cap V(G_c^T)| = 2$ by assumption, if $D \cap \{t_c^x, v_c^x, u_c^x\} \neq \emptyset$ then one of t_c^y and t_c^z is not dominated. Since t_c^x should be dominated nonetheless, it follows that $P_{x,T}^c(1) \in D$ and thus, $T_x \in D$ by Claim 26(i). Therefore, the truth assignment obtained by setting a variable x to true when $T_x \in D$ and to false otherwise, satisfies Φ .

Claim 28. Let D be an SD set of G and let $c \in C$ be a clause containing variables x, y and z. If $|D \cap V(G_c)| = 4$ and $|D \cap V(G_\ell)| = 14$ for every $\ell \in \{x, y, z\}$ then $D \cap V(G_c^T)$ is an independent set.

Proof. Assume that $|D \cap V(G_c)| = 4$ and $|D \cap V(G_\ell)| = 14$ for every $\ell \in \{x, y, z\}$ (note that then $|D \cap V(G_c^T)| = |D \cap V(G_c^F)| = 2$ by Claim 24(ii)). Suppose to the contrary that $D \cap V(G_c^T)$ contains an edge uv. Then $\{u, v\} = \{u_c^{\ell}, v_c^{\ell}\}$ for some $\ell \in \{x, y, z\}$: indeed, $\{u, v\} \cap \{u_c^x, u_c^y, u_c^z, a_c\} \neq \emptyset$ and $\{u, v\} \cap \{v_c^x, v_c^y, v_c^z, b_c\} \neq \emptyset$ by Claim 24(ii), and any edge between $\{u_c^x, u_c^y, u_c^z, a_c\}$ and $\{v_c^x, v_c^y, v_c^z, b_c\}$ connect vertices with the same superscript. Assume without loss of generality that $\{u, v\} = \{u_c^x, v_c^x\}$. Then $P_{y,T}^c(1), P_{z,T}^c(1) \in D$ as t_c^y and t_c^z should be dominated, which, by Claim 26, implies that $P_{y,F}^c(2), P_{z,F}^c(2) \notin D$. By Claim 25, it must then be that $q_c^y, q_c^z \in D$ for

one of f_c^y and f_c^z would otherwise not be dominated; but $|D \cap V(G_c^F)| = 2$ and thus, w_c is not dominated.

Claim 29. Let D be an SD set of G and let $c \in C$ be a clause containing variables $x, y, z \in X$. If $|D \cap V(G_c)| = 4$ and $|D \cap V(G_\ell)| = 14$ for every $\ell \in \{x, y, z\}$ then $D \cap V(G_c^F)$ is an independent set.

Proof. Assume that $|D \cap V(G_c)| = 4$ and $|D \cap V(G_\ell)| = 14$ for every $\ell \in \{x, y, z\}$ (note that then $|D \cap V(G_c^T)| = |D \cap V(G_c^F)| = 2$ by Claim 24(ii)). Suppose for a contradiction that $D \cap V(G_c^F)$ contains an edge uv. Then, since w_c should be dominated, $\{u, v\} \subseteq \{u_c, v_c, w_c\}$; in particular $D \cap \{q_c^\ell, f_c^\ell \mid \ell \in \{x, y, z\}\} = \emptyset$ by Claims 24(ii) and 25. But then, $\{P_{\ell,F}^c(2) \mid \ell \in \{x, y, z\}\} \subseteq D$ as f_c^x, f_c^y and f_c^z should be dominated, which, by Claim 26, implies that $\{P_{\ell,T}^c(1) \mid \ell \in \{x, y, z\}\} \subseteq D$. But $|D \cap V(G_c^T)| = 2$ and so, one of t_c^x, t_c^y and t_c^z is not dominated.

Claim 30. Let D be an SD set of G and let $x \in X$ be a variable. If $|D \cap V(G_x)| = 14$ and $D \cap V(G_x)$ contains an edge uv then $\{u, v\} = \{P(3), P(4)\}$ for some long paw P of G_x .

Proof. Assume that $|D \cap V(G_x)| = 14$ and $D \cap V(G_x)$ contains an edge uv. Then by Observation 9, $u, v \in V(P)$ for some long paw P of G_x . But $|D \cap V(P)| = 2$ and $D \cap \{P(4), P(5)\} \neq \emptyset$ by Claim 24(i) and so, $D \cap V(P) = \{u, v\} \subseteq \{P(3), P(4), P(5)\}$. Now suppose for a contradiction that $P(3) \notin D$. If $P = P_x$ then both T_x and F_x are not dominated since by Observation 9, $q_x^\ell, p_x^\ell \notin D$ for every clause ℓ containing x. If $P = P_{x,T}^\ell$ for some clause ℓ containing x, then $P_{x,T}^{\ell}(2)$ is not dominated since $p_x^{\ell} \notin D$ by Observation 9. Finally if $P = P_{x,F}^{\ell}$ for some clause ℓ containing x, then $P_{x,F}^{\ell}(1)$ is not dominated since $q_x^{\ell} \notin D$ by Observation 9. Thus, $P(3) \in D$ and so, $\{u, v\} = \{P(3), P(4)\}.$ ٦

Claim 31. Let D be an SD set of G and let $c \in C$ be a clause containing variables x, y and z. If $|D \cap V(G_c)| = 4$ and $|D \cap V(G_\ell)| = 14$ for every $\ell \in \{x, y, z\}$, then for every $\ell \in \{x, y, z\}$, the following hold.

(i) $D \cap \{u_c^{\ell}, v_c^{\ell}\} = \emptyset$ if and only if $P_{\ell,T}^c(1) \in D$.

- (ii) $q_c^{\ell} \notin D$ if and only if $P_{\ell,F}^c(2) \in D$. (iii) $P_{\ell,F}^c(2) \in D$ if and only if $P_{\ell,T}^c(1) \notin D$

Proof. Assume that $|D \cap V(G_c)| = 4$ and $|D \cap V(G_\ell)| = 14$ for every $\ell \in \{x, y, z\}$. Now consider $\ell \in \{x, y, z\}$ and let us first prove (i). If $D \cap \{u_c^\ell, v_c^\ell\} = \emptyset$ then since $t_c^\ell \notin D$ by Claim 25, necessarily $P_{\ell,T}^c(1) \in D$ as t_c^ℓ should be dominated. Conversely, assume that $P_{\ell,T}^c(1) \in D$ and suppose for a contradiction that $D \cap \{u_c^{\ell}, v_c^{\ell}\} \neq \emptyset$. Then since $|D \cap V(G_c^T)| = 2$ by Claim 24(ii), it must be that $D \cap \{u_c^p, v_c^p\} = \emptyset$ for some $p \neq \ell$; and since $t_c^p \notin D$ by Claim 25, necessarily $P_{p,T}^c(1) \in D$ as t_c^p should be dominated. Then $P_{\ell,F}^c(2), P_{p,F}^c(2) \notin D$ by Claim 26 and since $f_c^{\ell}, f_c^{p} \notin D$ by Claim 25, it follows that $q_c^\ell, q_c^p \in D$ as f_c^ℓ and f_c^p should be dominated. But $|D \cap V(G_c^F)| = 2$ by Claim 24 and thus, w_c is not dominated.

To prove (ii), observe that if $q_c^{\ell} \notin D$ then since $f_c^{\ell} \notin D$ by Claim 25, necessarily $P_{\ell,F}^c(2) \in D$ for f_c^{ℓ} would otherwise not be dominated. Conversely, assume that $P_{\ell,F}^c(2) \in D$ for some $\ell \in \{x, y, z\}$, and suppose for a contradiction that $q_c^{\ell} \in D$. Then since $|D \cap V(G_c^F)| = 2$ by Claim 24(ii) and w_c must be dominated, $q_c^p \notin D$ for every $p \neq \ell$; and since by Claim 25, $f_c^p \notin D$ for every $p \neq \ell$, necessarily $P_{p,F}^{c}(2) \in D$ for every $p \neq \ell$ (f_{c}^{p} would otherwise not be dominated). It then follows from Claim 26 that $D \cap \{P_{x,T}^{c}(1), P_{y,T}^{c}(1), P_{z,T}^{c}(1)\} = \emptyset$; but $|D \cap V(G_{c}^{T})| = 2$ by Claim 24(ii) and so, one of t_c^x, t_c^y and t_c^z is not dominated.

Let us finally prove (iii). If $P_{\ell,F}^c(2) \in D$ then by Claim 26, $P_{\ell,T}^c(1) \notin D$. Conversely, assume that $P_{\ell,T}^c(1) \notin D$. Then by item (i), $D \cap \{u_c^\ell, v_c^\ell\} \neq \emptyset$ which implies that $D \cap \{u_c^p, v_c^p\} = \emptyset$ for some $p \neq \ell$, since $|D \cap V(G_c^T)| = 2$ by Claim 24(ii). It then follows from item (i) that $P_{p,T}^c(1) \in D$ and so, $P_{p,F}^c(2) \notin D$ by Claim 26. Thus, by item (ii), $q_c^p \in D$; but $|D \cap V(G_c^F)| = 2$ by Claim 24(ii) and w_c must be dominated, and so $q_c^{\ell} \notin D$ which, by item (ii), implies that $P_{\ell F}^c(2) \in D$.

Claim 32. Let D be an SD set of G and let $x \in X$ be a variable contained in clauses c, c' and c''. If $|D \cap V(G_{\ell})| = 4$ for every $\ell \in \{c, c''c''\}$ and $|D \cap V(G_{\nu})| = 14$ for every variable ν appearing in c, c' or c'', then $D \cap V(G_x)$ is an independent set.

Proof. Assume that $|D \cap V(G_{\ell})| = 4$ for every $\ell \in \{c, c''c''\}$ and $|D \cap V(G_v)| = 14$ for every variable v appearing in c, c' or c''. Suppose for a contradiction that $D \cap V(G_x)$ contains an edge uv. Then by Claim 30, $\{u, v\} = \{P(3), P(4)\}$ for some long paw P of G_x . If $P = P_x$ then since $D \cap \{q_x^\ell, p_x^\ell \mid \ell \in \{c, c', c''\}\} = \emptyset$ by Observation 9, necessarily $P_{x,T}^c(2), P_{x,F}^c(1) \in D$ (one of p_x^c and q_x^c would otherwise not be dominated); but then by Claim 24(i), $P_{x,T}^c(1), P_{x,F}^c(2) \notin D$, a contradiction to Claim 31(iii). Thus $P = P_{x,R}^{\ell}$ for some $\ell \in \{c, c', c''\}$ and $R \in \{T, F\}$. Assume that R = T (the case where R = F is symmetric). Then since $D \cap \{q_x^\ell, p_x^\ell \mid \ell \in \{c, c', c''\}\} = \emptyset$ by Observation 9, necessarily $T_x \in D$ $(p_x^{\ell}$ would otherwise not be dominated) and so, $F_x \notin D$ by Claim 24(i). But then by Claim 26(ii), $P_{x,F}^{\ell}(2) \notin D$, a contradiction to Claim 31(ii) as $P_{x,T}^{\ell}(1) \notin D.$

Claim 33. If $\gamma_{t2}(G) = 14|X| + 4|C|$ then for every minimum SD set D of G, the following hold.

(i) D is an independent set.

(ii) For every $u \in D$, $|w_D(u)| = 1$.

Proof. Assume that $\gamma_{t2}(G) = 14|X| + 4|C|$ and let D be a minimum SD set of G. Then by Claims 25, 28, 29 and 32, D contains no edge. Let us next show that for every $u \in D$, $|w_D(u)| = 1$. Consider first a clause $c \in C$ containing variables $x, y, z \in X$ and let $u \in D \cap V(G_c)$. Note that since $|D \cap V(G_c^T)| = |D \cap V(G_c^F)| = 2$ by Claim 24(ii), if $|w_D(u)| \ge 2$ then $d(u, D \setminus V(G_c)) \le 2$. Now if $u \in \{u_c^{\ell}, v_c^{\ell}\}$ for some $\ell \in \{x, y, z\}$, then by Claim 31(i), $P_{\ell,T}^c(1) \notin D$ and thus, $d(u, D \setminus V(G_c)) > 0$ 2. Similarly, if $u = q_c^{\ell}$ for some $\ell \in \{x, y, z\}$, then by Claim 31(ii), $P_{\ell,F}^c(2) \notin D$ and thus, $d(u, D \setminus V(G_c)) > 2$. Since $u \notin \{f_c^\ell, t_c^\ell \mid \ell \in \{x, y, z\}\}$ by Claim 25 and $d(w, D \setminus V(G_c)) \ge 3$ for every $w \in \{a_c, b_c\}$, it follows that $|w_D(u)| = 1$. Consider next a variable $x \in X$ contained in clauses $c, c', c'' \in C$, and suppose to the contrary that there exists a vertex $u \in D \cap V(G_x)$ such that $|w_D(u)| \ge 2$. Since $|D \cap V(P)| = 2$ for ever long paw P in G_x and $D \cap \{q_x^\ell, p_x^\ell \mid \ell \in \{c, c', c''\}, \ell \in \{c, c', c''\}$ necessarily $u \in \{P(1), P(2)\}$ for some long paw P of G_x . Suppose first that $u = T_x$ (the case where $u = F_x$ is symmetric). Then by Observation 9, there must exist $\ell \in \{c, c', c''\}$ such that $P_{x,T}^{\ell}(2) \in \mathbb{C}$ D; in particular, $P_{x,T}^{\ell}(1) \notin D$ by Claim 24(i). But $F_x \notin D$ by Claim 24(i) and so, $P_{x,F}^{\ell}(2) \notin D$ by Claim 26(ii), a contradiction to Claim 31(iii). Suppose next that $u \in \{P_{x,R}^{\ell}(1), P_{x,R}^{\ell}(2)\}$ for some $\ell \in \{c, c', c'\}$ and $R \in \{T, F\}$. Assume that R = F (the case where R = T is symmetric). Then u must have a witness in $D \cap V(G_{\ell}^F)$: indeed, if u had two witnesses in $D \cap V(G_x)$ then one of them would be F_x and so, $|w_D(F_x)| \ge 2$ which is excluded by the previous case. However, $f_\ell^x \notin D$ by Claim 25, and if $q_\ell^x \in D$ then $P_{x,F}^\ell(2) \notin D$ by Claim 31(ii); thus $d(u, D \cap V(G_\ell^F)) > 2$, a contradiction. Therefore, every vertex in $D \cap V(G_x)$ has a unique witness.

Claim 34. Let D be an SD set of G and let $x \in X$ be a variable. If $|D \cap V(G_x)| = 14$ then for every clause $c \in C$ containing x, the following hold.

(i) If $D \cap \{P_{x,T}^c(1), P_{x,T}^c(3)\} \neq \emptyset$ then $P_{x,F}^c(2) \notin D$. (ii) If $D \cap \{P_{x,F}^c(2), P_{x,F}^c(3)\} \neq \emptyset$ then $P_{x,T}^c(1) \notin D$.

Proof. Assume that $|D \cap V(G_x)| = 14$ and let $c \in C$ be a clause containing x. Suppose that $D \cap \{P_{x,T}^c(1), P_{x,T}^c(3)\} \neq \emptyset$. Then by Claim 24(i), $P_{x,T}^c(2) \notin D$ which, by Observation 9, implies that $T_x \in D$ (p_x^c) would otherwise not be dominated); but then $F_x \notin D$ by Claim 24(i) and so, $P_{x,F}^{c}(2) \notin D$ by Claim 26(ii). Item (ii) follows by symmetry. ٦

Claim 35. $ct_{\gamma_{t2}}(G) = 3$ if and only if $\gamma_{t2}(G) = 14|X| + 4|C|$.

Proof. Assume first that $ct_{\gamma_{t2}}(G) = 3$ and let D be a minimum SD set of G. Consider a variable $x \in X$ contained in clauses c, c' and c''. If $|D \cap V(P)| \geq 3$ for some long paw P of G_x then $D \cap V(P)$ contains a friendly triple, a contradiction to Theorem 1.7. Thus $|D \cap V(P)| \leq 2$ for every long paw P of G_x and we conclude by Claim 24(i) that, in fact, equality holds. Let us assume, in the following, that for every long paw P of G_x , $D \cap \{P(1), P(2)\} \neq \emptyset$ and $P(4) \in D$ (if this does not hold for a long paw P then it suffices to consider, e.g., $(D \setminus V(P)) \cup \{P(1), P(4)\}$ in place of D). Now suppose that $D \cap \{p_x^c, p_x^{c'}, p_x^{c''}\} \neq \emptyset$, say $p_x^c \in D$ without loss of generality. Then $T_x \notin D$ (D would otherwise contain the friendly triple p_x^c, T_x, v_x) and so, $(D \setminus \{p_x^c\}) \cup \{T_x\}$ is a minimum SD set of G containing the friendly triple T_x, F_x, w_x , a contradiction to Theorem 1.7. Thus $D \cap \{p_x^c, p_x^{c'}, p_x^{c''}\} = \emptyset$ and by symmetry, we conclude that $D \cap \{q_x^c, q_x^{c'}, q_x^{c''}\} = \emptyset$ as well. Therefore, $|D \cap V(G_x)| = 14$.

Next, consider a clause $c \in C$ and let $x, y, z \in X$ be the variables contained in c. Then $|D \cap$ $\{u_c, v_c, w_c, q_c^x, q_c^y, q_c^z\} \le 2$ for $D \cap \{u_c, v_c, w_c, q_c^x, q_c^y, q_c^z\}$ would otherwise contain a friendly triple; and we conclude by Claim 24 that, in fact, equality holds. Now suppose that $D \cap \{f_c^x, f_c^y, f_c^z\} \neq \emptyset$, say $f_c^x \in D$ without loss of generality. Then $q_c^x \notin D$: indeed, D would otherwise contain the friendly triple f_c^x, q_c^x, t , where $t \in D \cap \{u_c, v_c, w_c, q_c^y, q_c^z\}$, a contradiction to Theorem 1.7. But by assumption, $P_{x,F}^c(2)$ is not a private neighbour of f_c^x and the vertex in $D \cap \{P_{x,F}^c(1), P_{x,F}^c(2)\}$ is witnessed by $P_{x,F}^c(4)$, which implies that $D' = (D \setminus \{f_c^x\}) \cup \{q_c^x\}$ is a minimum SD set of G where $|D' \cap \{u_c, v_c, w_c, q_c^x, q_c^y, q_c^z\}| \geq 3$, a contradiction to the above. Thus, $D \cap \{f_c^x, f_c^y, f_c^z\} = \emptyset$ and so, $|D \cap V(G_c^F)| = 2$. Suppose next that $|D \cap V(G_c^T)| \ge 3$. If $D \cap \{t_c^x, t_c^y, t_c^z\} = \emptyset$, say $t_c^x \in D$ without loss of generality, then surely one of u_c^x and v_c^x does not belong to D (D would otherwise contain the friendly triple t_c^x, u_c^x, v_c^x), say $u_c^x \notin D$ without loss of generality. Since by assumption, $P_{x,T}^c(1)$ is not a private neighbour of t_c^x and the vertex in $D \cap \{P_{x,T}^c(1), P_{x,T}^c(2)\}$ is witnessed by $P_{x,T}^c(4)$, the set $D' = (D \setminus \{t_c^x\} \cup \{u_c^x\}$ is a minimum SD of G (note that u_c^x is within distance two of every vertex in $D \cap V(G_c^T)$). By repeating this argument if necessary, we obtain a minimum SD set D'' of G such that $D'' \cap \{t_c^x, t_c^y, t_c^z\} = \emptyset$ and $|D'' \cap V(G_c^T)| \ge 3$; but then, it is easy to see that $D'' \cap (\{a_c, b_c\} \cup \{u_c^{\ell}, v_c^{\ell} \mid \ell \in \{x, y, z\}\}$ contains a friendly triple, a contradiction to Theorem 1.7. Thus, $|D \cap V(G_c^T)| \leq 2$ and we conclude by Claim 24(ii) that, in fact, equality holds. Therefore, $|D \cap V(G_c)| = 4$ and thus, $\gamma_{t2}(G) = |D| = 14|X| + 4|C|$.

Conversely, assume that $\gamma_{t2}(G) = 14|X| + 4|C|$. Observe first that since by Claim 33, every minimum SD set of G is an independent set, no minimum SD set of G contains a friendly triple; thus by Theorem 1.7(i), $ct_{\gamma_{t2}}(G) > 1$. Now suppose to the contrary that $ct_{\gamma_{t2}}(G) = 2$ and let D be an SD set of size $\gamma_{t2}(G) + 1$ containing an ST configuration (see Theorem 1.7(ii)). Then by Claim 24, there exists either a clause $c \in C$ such that $|D \cap V(G_c)| = 5$ or a variable $x \in X$ such that $|D \cap V(G_x)| = 15$. We next distinguish these two cases.

Case 1. There exists a clause $c \in C$ such that $|D \cap V(G_c)| = 5$. Let $x, y, z \in X$ be the three variables appearing in c. Then for any clause $\ell \in C \setminus \{c\}, D \cap V(G_{\ell})$ contains no edges by Claims 28 and 29; and for any variable $v \in X \setminus \{x, y, z\}, D \cap V(G_v)$ contains no edge by Claim 32. Thus by Claim 25, $D \setminus (V(G_c) \cup V(G_x) \cup V(G_y) \cup V(G_z))$ contains no edge and a fortiori, no ST-configurations. Further observe if for some $\ell \in \{x, y, z\}, D \cap V(G_{\ell})$ contains an edge uv then by Claim 30, $\{u, v\} = \{P(3), P(4)\}$ for some long paw P in G_{ℓ} . Now if $P \neq P_{\ell,T}^c, P_{\ell,F}^c$ then by Observation 9 and Claim 25, $d(\{u,v\}, D \setminus \{u,v\}) \ge 3$ and so uv cannot be part of an ST-configuration in D; and if $P = P_{\ell,T}^c$ (resp. $P = P_{\ell,F}^c$) then $d(\{u,v\}, D \setminus \{u,v\}) \ge 3$ unless $t_c^\ell \in D$ (resp. $f_c^{\ell} \in D$). Thus any edge uv of an ST-configuration in D satisfies one of the following.

- (aT) $\{u, v\} = \{P_{\ell,T}^c(3), P_{\ell,T}^c(4)\}$ for some $\ell \in \{x, y, z\}$ and $t_c^\ell \in D$.
- (aF) $\{u, v\} = \{P_{\ell, F}^{c}(3), P_{\ell, F}^{c}(4)\}$ for some $\ell \in \{x, y, z\}$ and $f_{c}^{\ell} \in D$.
- (bT) $\{u, v\} = \{P_{\ell,T}^{(1)}, t_c^{\ell}\}$ for some $\ell \in \{x, y, z\}$.
- (bF) $\{u, v\} = \{P_{\ell, F}^{(\ell)}2), f_c^{\ell}\}$ for some $\ell \in \{x, y, z\}.$
- (cT) $u, v \in V(G_c^T)$. (cF) $u, v \in V(G_c^F)$.

Note that since for any $\ell \in \{x, y, z\}$ and any $R \in \{T, F\}, |D \cap \{P_{\ell,R}^c(1), P_{\ell,R}^c(2), P_{\ell,R}^c(3)\}| \leq 1$ by Claim 24(i), if there is an edge satisfying (aR) and an edge satisfying (bR), then these two edges are vertex-disjoint. We now distinguish cases depending on whether $|D \cap V(G_c^T)| = 3$ or $|D \cap V(G_c^F)| = 3.$

Case 1.a. $|D \cap V(G_c^T)| = 3$. Then $D \cap \{f_c^x, f_c^y, f_c^z\} = \emptyset$ by Claim 25, which implies that no edge of an ST-configuration in D satisfies (aF) or (bF). Now if $D \cap V(G_c^F)$ contains an edge uvthen $\{u, v\} \subseteq \{w_c, v_c, u_c\}$ as w_c should be dominated; but then $d(\{u, v\}, D \setminus \{u, v\}) \ge 3$ and so, uvcannot be part of an ST-configuration in D. Thus no edge of an ST-configuration in D satisfies (cF). Further note that since $|D \cap \{t_c^x, t_c^y, t_c^z\}| \leq 1$ by Claim 24(ii), there is at most one edge of an ST-configuration in D satisfying (aT) or (bT).

Suppose first that $D \cap V(G_c^T)$ is an independent set and let uv be an edge of an ST-configuration in D. If uv satisfies (aT) for some $\ell \in \{x, y, z\}$, then $D \cap \{P_{\ell,T}^{(1)}(1), P_{\ell,T}^{c}(2)\} = \emptyset$ by Claim 24(i) and so, uv cannot be part of a P_3 . Similarly, if uv satisfies (bT) for some $\ell \in \{x, y, z\}$, then

 $D \cap \{P_{\ell,T}^c(2), P_{\ell,T}^c(3)\} = \emptyset$ by Claim 24(i) and so, uv cannot be part of a P_3 . It follows that D contains no P_3 , which implies that D must contain an O_3 or an O_7 ; in particular, D contains two vertex-disjoint edges. However, any such edge would satisfy (aT) or (bT), a contradiction to the above.

Suppose next that $D \cap V(G_c^T)$ contains an edge. We claim that there exists at most one variable $\ell \in \{x, y, z\}$ such that $D \cap \{P_{\ell,T}^c(1), P_{\ell,T}^c(3)\} \neq \emptyset$. Indeed, suppose for a contradiction that there are two such variables, say x and y without loss of generality. Then $P_{x,F}^c(2), P_{y,F}^c(2) \notin D$ by Claim 34, which implies that $|D \cap \{q_c^x, q_c^y, f_c^x, f_c^y\}| \ge 2$ as f_c^x and f_c^y should be dominated; but $|D \cap V(G_c^F)| = 2$ and so, w_c is not dominated, a contradiction. Thus assume, without loss of generality, that $D \cap \{P_{x,T}^c(1), P_{x,T}^c(3)\} = D \cap \{P_{y,T}^c(1), P_{y,T}^c(3)\} = \emptyset$ (note that $D \cap (\{t_c^x\} \cup V(P_{x,T}^c))$) and $D \cap (\{t_c^y\} \cup V(P_{y,T}^c))$ can then contain no edge). Since t_c^x and t_c^y should be dominated, necessarily $D \cap \{t_c^x, u_c^x, v_c^x\} \neq \emptyset$ and $D \cap \{t_c^y, u_c^y, v_c^y\} \neq \emptyset$. Now if $D \cap \{t_c^z, u_c^z, v_c^z\} \neq \emptyset$ then since $D \cap V(G_c^T)$ contains an edge by assumption, it must be that $u_c^a, u_c^b \in D$ or $v_c^a, v_c^b \in D$ for two variables $a, b \in \{x, y, z\}$. In the first case, since $D \cap \{v_c^x, v_c^y, v_c^z\} \neq \emptyset$ by Claim 24(ii), necessarily $D \cap \{t_c^x, t_c^y, t_c^z\} = \emptyset$; and in the second case, since $D \cap \{u_c^x, u_c^y, u_c^z\} \neq \emptyset$ by Claim 24(ii), necessarily $D \cap \{t_c^x, t_c^y, t_c^z\} = \emptyset$. Thus, in both cases, we conclude that D contains no edge satisfying (aT) or (bT); but then D contains only one edge, a contradiction. Thus $D \cap \{t_c^z, v_c^z, v_c^z\} =$ \varnothing which implies, in particular, $D \cap (\{t_c^z\} \cup V(P_{z,T}^c))$ contains no edge. Since by assumption $\{t_c^x\} \cup V(P_{x,T}^c)$ and $\{t_c^y\} \cup V(P_{y,T}^c)$ contain no edge as well, it follows that $D \cap V(G_c^T)$ must contain a P_3 (D would otherwise contain only one edge). However, since $D \cap \{t_c^x, u_c^x, v_c^x\} \neq \emptyset$ and $D \cap \{t_c^y, u_c^y, v_c^y\} \neq \emptyset$, any such P_3 must then be contained in $\{u_c^x, v_c^x, u_c^y, v_c^y\}$ by Claim 24(ii); but $d(\{u_c^x, v_c^x, u_c^y, v_c^y\}, D \setminus V(G_c^T)) \geq 3, \text{ a contradiction}.$

Case 1.b. $|D \cap V(G_c^F)| = 3$. Then $D \cap \{t_c^x, t_c^y, t_c^z\} = \emptyset$ by Claim 25, which implies that no edge of an ST-configuration satisfies (aT) or (bT). Let us next show that no edge of an STconfiguration in D satisfies (cT). Suppose that $D \cap V(G_c^T)$ contains an edge uv. Then by ??(ii), $\{u,v\} = \{u_c^\ell, v_c^\ell\}$ for some $\ell \in \{x, y, z\}$, say x without loss of generality. If $P_{x,T}^c(1) \notin D$ then $d(\{u,v\}, D \setminus \{u,v\} \ge 3 \text{ and so, } uv \text{ cannot be part of an ST-configuration in } D$. Thus suppose that $P_{x,T}^c(1) \in D$. Since t_c^y and t_c^z should be dominated and $D \cap \{t_c^v, u_c^v, v_c^v, t_c^z, u_c^z, v_c^z\} = \emptyset$, necessarily $P_{y,T}^c(1), P_{z,T}^c(1) \in D$. Then $D \cap \{P_{x,F}^c(2), P_{y,F}^c(2), P_{z,F}^c(2)\} = \emptyset$ by Claim 34 and so, $|D \cap \{f_c^\ell, q_c^\ell \mid \ell \in \{x, y, z\}\}| \ge 3$ as f_c^x, f_c^y and f_c^z should be dominated. But $|D \cap V(G_c^F)| = 3$ and so, w_c is not dominated, a contradiction. Thus no edge of an ST-configuration satisfies (cT). Now observe that $|D \cap \{f_c^x, f_c^y, f_c^z\}| \le 1$: indeed, if, say, $t_c^x, t_c^y \in D$ then $D \cap \{u_c, q_c^x, q_c^y, q_c^z, f_c^z\} \neq \emptyset$ as q_c^z should be dominated; but then $D \cap \{w_c, v_c\} = \emptyset$ and so, w_c is not dominated, a contradiction. This implies, in particular, that there is at most one edge of an ST-configuration in D satisfying (aF) or (bF).

Now suppose that $D \cap V(G_c^F)$ is an independent set and let uv be an edge of an ST-configuration in D. If uv satisfies (aF) for some $\ell \in \{x, y, z\}$, then $D \cap \{P_{\ell,F}^{(1)}, P_{\ell,F}^{c}(2)\} = \emptyset$ by Claim 24(i) and so, uv cannot be part of a P_3 . Similarly, if uv satisfies (bF) for some $\ell \in \{x, y, z\}$, then $D \cap \{P_{\ell,T}^c(1), P_{\ell,T}^c(3)\} = \emptyset$ by Claim 24(i) and so, uv cannot be part of a P_3 . It follows that Dcontains no P_3 , which implies that D must contain an O_3 or an O_7 ; in particular, D contains two vertex-disjoint edges. However, any such edge would satisfy (aF) or (bF), a contradiction to the above.

Suppose next that $D \cap V(G_c^F)$ contains an edge. We claim that there exists at least one variable $\ell \in \{x, y, z\}$ such that $D \cap \{P_{\ell,F}^c(2), P_{\ell,F}^c(3)\} = \emptyset$. Indeed, if there is no such variable then by Claim 34, $D \cap \{P_{x,T}^c(1), P_{y,T}^c(1), P_{z,T}^c(1)\} = \emptyset$; but $|D \cap V(G_c^T)| = 2$ and so, one of t_c^x, t_c^y and t_c^z is not dominated. Thus assume, without loss of generality, that $D \cap \{P_{x,F}^c(2), P_{x,F}^c(3)\} = \emptyset$ (note that $D \cap (\{f_c^x\} \cup V(P_{x,F}^c)\})$ can then contain no edge). Since f_c^x should be dominated, necessarily $D \cap \{q_c^x, f_c^x\} \neq \emptyset$. Now if $q_c^x, f_c^x \in D$ then since $|D \cap \{w_c, v_c\}| = 1$, $D \cap V(G_c^F)$ contains only one edge and so, D contains no ST-configuration, a contradiction. Suppose next that $f_c^x \in D$. Since $D \cap \{w_c, v_c\} \neq \emptyset$ and $D \cap V(G_c^F)$ contains only one edge and so, D contains no ST-configuration, a contradiction. Suppose next that $p_c^x \in D$. Since $D \cap \{w_c, v_c\} \neq \emptyset$ and $D \cap V(G_c^F)$ contains only one edge and so, D contains no ST-configuration, a contradiction. Suppose next that $p_c^x \in D$. Since $D \cap \{w_c, v_c\} \neq \emptyset$ and $D \cap V(G_c^F)$ contains only one edge and so, D contains no ST-configuration. Suppose finally that $q_c^x \in D$. Since $D \cap \{w_c, v_c\} \neq \emptyset$ and $D \cap V(G_c^F)$ contains only one edge and so, D contains no ST-configuration. Suppose finally that $q_c^x \in D$. Since $D \cap \{w_c, v_c\} \neq \emptyset$ and $D \cap V(G_c^F)$ contains only one edge and so, D contains no ST-configuration. Suppose finally that $q_c^x \in D$. Since $D \cap \{w_c, v_c\} \neq \emptyset$ in which case $D \cap V(G_c^F)$ contains only one edge and so, D contains no ST-configuration. Thus it must be that $|D \cap \{w_c, v_c, u_c\}| = 2$; but then $d(D \cap V(G_c^F), D \setminus V(G_c^F)) \geq 3$ and so, D contains no ST-configuration.

ST-configuration.

Case 2. There exists a variable $x \in X$ such that $|D \cap V(G_x)| = 15$. Let $c, c', c'' \in C$ be the three clauses in which x appears. If for some variable $\ell \neq x$ appearing in c, c' or $c'', D \cap V(G_\ell)$ contains an edge uv then by Claim 30, $\{u, v\} = \{P(3), P(4)\}$ for some long paw P of G_ℓ . But then $d(\{u, v\}, D \setminus \{u, v\}) \geq 3$ by Claim 25 and so, uv cannot be part of an ST-configuration in D. Now for any clause $\ell \in C \setminus \{c, c', c''\}, D \cap V(G_\ell)$ contains no edges by Claims 28 and 29; and for any variable $\ell \in X$ not appearing in c, c' or $c'', D \cap V(G_v)$ contains no edge by Claim 32. Thus by Claim 25, any ST-configuration in D is in fact contained in $D \cap (V(G_x) \cup V(G_c) \cup V(G_{c'}) \cup V(G_{c''}))$.

Now suppose that for some clause $\ell \in \{c, c', c''\}$, $D \cap V(G_\ell)$ contains an edge uv. If $u, v \in V(G_\ell^F)$ then, since w_ℓ must be dominated, necessarily $\{u, v\} \subseteq \{w_c, v_c, u_c\}$. But then $d(\{u, v\}, D \setminus \{u, v\}) \ge 3$ and so, uv cannot be part of an ST-configuration in D. Thus assume that $u, v \in V(G_\ell^T)$ and let $y, z \in X$ be the other two variables appearing in ℓ . Then by Claim 24(ii), $\{u, v\} = \{u_\ell^a, v_\ell^a\}$ for some variable $a \in \{x, y, z\}$. Now if $P_{a,T}^\ell(1) \notin D$ then $d(\{u, v\}, D \setminus \{u, v\}) \ge 3$ and so, uv cannot be part of an ST-configuration in D. Thus suppose that $P_{a,T}^\ell(1) \in D$. Since for every $b \neq a, t_\ell^b$ should be dominated, it follows that $P_{b,T}^\ell(1) \in D$ as well. Then by Claim 34, $P_{y,F}^\ell(2), P_{z,F}^\ell(2) \notin D$ and so $|D \cap \{q_\ell^y, f_\ell^y, q_\ell^z, f_\ell^z\}| \ge 2$ since f_ℓ^y and f_ℓ^z should be dominated, But $|D \cap V(G_\ell^F)| = 2$ and so, w_c is not dominated, a contradiction. Thus no edge of an ST-configuration in D is contained in $D \cap (V(G_c) \cup V(G_{c'}) \cup V(G_{c''})$. By Claim 25, it follows that every edge of an ST-configuration in D is contained in G_x . We next distinguish cases depending on whether $D \cap \{q_x^\ell, p_x^\ell \mid \ell \in \{c, c', c''\}\} \neq \emptyset$ or $|D \cap V(P)| = 3$ for some long paw P of G_x .

Case 2.a. $D \cap \{q_x^\ell, p_x^\ell \mid \ell \in \{c, c', c''\}\} \neq \emptyset$. Then by Claim 24(i), $|D \cap \{q_x^\ell, p_{x'}^\ell \mid \ell \in \{c, c', c''\}\}| = 1$. Now suppose that $D \cap \{q_x^c, q_x^{c'}, q_x^{c''}\} \neq \emptyset$ (the case where $D \cap \{p_x^c, p_x^c, p_x^c, p_x^c\} \neq \emptyset$ is symmetric), say $q_x^c \in D$ without loss of generality. Suppose first that $F_x \in D$. Since $f_x^c \notin D$ by ?? and $P_{x,F}^c(2)$ must be dominated, $D \cap \{P_{x,F}^c(1), P_{x,F}^c(2), P_{x,F}^c(3)\} \neq \emptyset$ and so, by Claim 24(i), $D' = D \setminus \{q_x^c\}$ is an SD set of G. Now since D' is in fact minimum, it follows from Claim 33 that every vertex in D' has a unique witness; in particular, $D' \cap \{P_{x,F}^c(1), P_{x,F}^{c'}(1), P_{x,F}^{c''}(1)\} = \emptyset$ as $F_x \in D$. On the other hand, since D is not minimal, D must contain an O_6 by Lemma 2.3 and Claim 33; in particular, $D \cap V(G_x)$ contains a $P_3 Q$. Now since $|D \cap V(P)| = 2$ for every long paw P of G_x and $D \cap \{q_x^{c'}, q_x^{c''}, p_x^{c}, p_x^{c''}, p_x^{c''}\} = \emptyset$, necessarily $q_x^c \in V(Q)$. However $N(q_x^c) \cap D = \{F_x\}$; and since $D \cap \{v_x, w_x\} \neq \emptyset$ by Claim 24(i), we also have that $N(F_x) \cap D = \{q_x^c\}$, a contradiction. Second, suppose that $F_x \notin D$. Then $D \cap V(G_c)$ contains no P_3 : indeed, by Claim 24(i),

Second, suppose that $F_x \notin D$. Then $D \cap V(G_c)$ contains no P_3 : indeed, by Claim 24(i), $D \cap \{F_x, q_x^{c'}, q_x^{c''}, p_x^{c}, p_x^{c''}\} = \emptyset$ and for every long paw P of G_x , $|D \cap \{P(4), P(5)\}| \ge 1$ and $|D \cap \{P(1), P(2), P(3)\}| \le 1$. It follows that D contains an O_3 or an O_7 ; in particular, $D \cap V(G_x)$ contains two vertex-disjoint edges. Now suppose that some long paw P of G_x contains an edge uv. Then since $D \cap \{P(4), P(5)\} \neq \emptyset$ by Claim 24(i), necessarily $\{u, v\} \subseteq$ $\{P(3), P(4), P(5)\}$. Thus, if $P \neq P_x, P_{x,F}^c$ then $d(\{u, v\}, D \setminus \{u, v\}) \ge 3$ by Claim 25 and so, uvcannot be part of an ST-configuration in D. It follows that $D \cap V(P_x)$ must contain an edge, as $D \cap (\{q_x^c\} \cup V(P_{x,F}^c))$ can contain at most one edge of an ST-configuration in D; in particular, $D \cap V(P_x) = \{v_x, u_x\}$. Note that since p_x^c should be dominated and $T_x \notin D$, $P_{x,T}^c(2) \in D$ and so, $P_{x,T}^c(1) \notin D$. Now, similarly, $D \cap (\{q_x^c\} \cup V(P_{x,F}^c))$ must contain an edge uv; and either $q_x^c \in \{u, v\}$ in which case $D \cap \{P_{x,F}^c(1), P_{x,F}^c(2), P_{x,F}^(3)\} = \{P_{x,F}^c(1)\}$, or $u, v \in V(P_{x,F}^c)$ in which case $D \cap V(P_x) = \{P_{x,F}^c(3), P_{x,F}^c(4)\}$. In both cases, we conclude that $P_{x,F}^c(2) \notin D$. Since, as shown above, $P_{x,T}^c(1) \notin D$ as well, it follows from Claim 25 that $q_x^c \in D$ and $D \cap \{u_x^c, v_x^c\} \neq \emptyset$ as f_x^c and t_x^c should be dominated, respectively. Now $|D \cap V(G_c^T)| = |D \cap V(G_c^F)| = 2$ and so, by Claim 24(ii), there must exist a variable $\ell \neq x$ contained in c such that $D \cap \{q_c^c, u_c^\ell, v_c^\ell\} = \emptyset$. But f_c^ℓ and t_c^ℓ should be dominated and so, by Claim 25, $P_{\ell,F}^c(2), P_{\ell,T}^c(1) \in D$, a contradiction to Claim 26.

Case 2.b. $|D \cap V(P)| = 3$ for some long paw P of G_x . Suppose first that $P = P_x$. If $D \cap V(P) = \{P(3), P(4), P(5)\}$ then $D \setminus \{P(3)\}$ is an SD set of G and so, by Lemma 2.3 and Claim 33, D must contain an O_6 ; in particular, $D \cap V(G_x)$ contains a $P_3 Q$. Since $D \cap \{q_x^\ell, p_x^\ell \mid \ell \in \{c, c', c''\}\} = \emptyset$ and for any long paw $P' \neq P$, $|D \cap V(P')| = 2$, necessarily Q = P(3), P(4), P(5); but then, by Claim 25 and because $D \cap \{q_x^\ell, p_x^\ell \mid \ell \in \{c, c', c''\}\} = \emptyset$, $d(P(4), D \setminus \{P(3), P(4), P(5)\}) \geq 3$, a contradiction. Suppose next that $|D \cap \{P(3), P(4), P(5)\}| = 2$. If $P(4), P(5) \in D$ then $D \setminus \{P(5)\}$ is an SD set of G and so, by Lemma 2.3 and Claim 33, D




Figure 12: The graph H_c (a rectangle indicates that the corresponding set of vertices is a clique).

must contain an O_6 ; but D contains no P_3 in this case, a contradiction. Now suppose that $D \cap \{P(3), P(4), P(5)\} = \{P(3), P(4)\}$ and assume that $P(1) \in D$ (the case where $P(2) \in P(3)$) D is symmetric). Then $D' = D \setminus \{P(3)\}$ is a minimum SD set of G and so, D must contain an O_6 by Lemma 2.3 and Claim 33; in particular, $D \cap V(G_c)$ contains a P_3 Q. Since $D \cap \{q_x^\ell, p_x^\ell \mid \ell \in \{c, c', c''\}\} = \emptyset$ and for any long paw $P' \neq P, |D \cap V(P')| = 2$, necessarily Q = P(1), P(3), P(4). However, by Claim 33(ii), $w_{D'}(P(1)) = \{P(4)\}$ and so $d(P(3), D \setminus D)$ $\{P(1), P(3), P(4)\} \geq d(P(1), D' \setminus \{P(1), P(4)\}) \geq 3$, a contradiction. Suppose finally that $D \cap \{P(3), P(4), P(5)\} = \{P(3), P(5)\}$ and assume that $P(1) \in D$ (the case where $P(2) \in D$ is symmetric). Then, since $D \cap \{q_x^{\ell}, p_x^{\ell} \mid \ell \in \{c, c', c''\}\} = \emptyset$ and for every long paw $P' \neq P$, $|D \cap V(P')| = 2, D \cap V(G_x)$ contains no P_3 . It follows that D contains an O_3 or O_6 ; in particular, $D \cap V(G_x)$ must contain an edge uv distinct from P(1)P(3). Now uv can only be contained in some long paw $P' \neq P$; and since $|D \cap \{P'(4), P'(5)\}| \geq 1$ and $|D \cap \{P'(1), P'(2), P'(3)\}| \leq 1$ by Claim 24(i), in fact $\{u, v\} \subseteq \{P'(3), P'(4), P'(5)\}$. But then, by Claim 25 and because $D \cap \{q_x^\ell, p_x^\ell \mid \ell \in \{c, c', c''\}\} = \emptyset, d(\{u, v\}, D \setminus \{u, v\}) \ge 3$, a contradiction. Thus, it must be that $|D \cap \{P(3), P(4), P(5)\}| = 1$; and since P(5) must be dominated and any vertex dominating P(5)must have a witness, in fact $D \cap \{P(3), P(4), P(5)\} = P(4)$. Since $D \cap \{q_x^{\ell}, p_x^{\ell} \mid \ell \in \{c, c', c''\}\} = \emptyset$ and for every long paw $P' \neq P$, $|D \cap V(P')| = 2$, it follows that D contains no P_3 and so, D contains an O_3 or an O_6 . In particular, D must contain an edge uv distinct from P(1)P(2); but then, we conclude, as in the previous case, that $\{u, v\} \subseteq \{P'(3), P'(4), P'(5)\}$ for some long paw $P' \neq P$ and $d(\{u, v\}, D \setminus \{u, v\}) \geq 3$, a contradiction which concludes the proof.

Now by Claims 27 and 35, Φ is satisfiable if and only if $ct_{\gamma_{t2}}(G) = 3$; and since G is $K_{1,3}$ -free, the lemma follows.

Lemma 4.12. CONTRACTION NUMBER $(\gamma_{t2},3)$ is NP-hard on $3P_4$ -free graphs.

Proof. We reduce from POSITIVE EXACTLY 3-BOUNDED 1-IN-3 3-SAT (see Section 2 for a precise definition of this problem): given an instance Φ of this problem, with variable set X and clause set C, we construct an instance G of CONTRACTION NUMBER $(\gamma_{t2},3)$ as follows. For every clause $c \in C$, we introduce the gadget G_c which is constructed as follows. Let $x, y, z \in X$ be the variables appearing in c. For every $\ell \in \{x, y, z\}$, denote by a_{ℓ} and b_{ℓ} the two other clauses in C in which ℓ appears. To construct the gadget G_c for c, we first construct the auxiliary graph H_c depicted in Figure 12 (where a rectangle indicates that the corresponding set is a clique). More precisely, H_c consists of two cliques $V_c = \{p_c, q_c, v_c^x, v_c^y, v_c^z\}$ and $K_c = \{u_c^{\ell, a_\ell}, u_c^{\ell, b_\ell} \mid \ell \in \mathcal{C}\}$ $\{x, y, z\}\} \cup \{t_c^{x, y}, t_c^{x, z}, t_c^{y, z}\},$ and the following edges.

- For every $\ell \in \{x, y, z\}$, u_c^{ℓ} is complete to $\{u_c^{\ell, a_{\ell}}, u_c^{\ell, b_{\ell}}\}$. For every $\ell \in \{x, y, z\}$, u_c^{ℓ} is adjacent to $t_c^{a, b}$ if and only if $\ell \in \{a, b\}$.

The gadget G_c then consists of two disjoint copies of H_c , denoted by $H_{c,1}$ and $H_{c,2}$, with the following additional edges. For $i \in [2]$, let us denote by $\{p_{c,i}, q_{c,i}, v_{c,i}^x, v_{c,c}^y, v_{c,i}^z\} \cup \{u_{c,i}^{\ell,a_\ell}, u_{c,i}^{\ell,b_\ell} \mid \ell \in \{x, y, z\}\} \cup \{t_{c,i}^{x,y}, t_{c,i}^{x,z}, t_{c,i}^{y,z}\}$ the vertex set of $H_{c,i}$, and by $K_{c,i}$ (resp. $V_{c,i}$) the copy of K_c (resp. V_c) in $H_{c,i}$.

- (1) For every $\ell \in \{x, y, z\}$, $v_{c,1}^{\ell}$ is adjacent to $t_{c,2}^{a,b}$ if and only if $\ell \notin \{a, b\}$. (2) For every $\ell \in \{x, y, z\}$, $v_{c,2}^{\ell}$ is adjacent to $t_{c,1}^{a,b}$ if and only if $\ell \notin \{a, b\}$.
- (3) $p_{c,1}$ is complete to $V_{c,2} \setminus \{p_{c,2}\}.$
- (4) $p_{c,2}$ is complete to $V_{c,1} \setminus \{p_{c,1}\}$.




Figure 13: The graph G_c (a rectangle indicates that the corresponding set of vertices is a clique, and thick edges between a vertex and a rectangle indicates that the vertex is complete to the corresponding set).

This completes the construction of G_c (see Figure 13 for an illustration - the edges of (1) and (2) have been omitted for clarity).

To complete the construction of G, we proceed as follows. For every two clauses $a, b \in C$ containing a same variable $u \in X$, we further add edges between G_a and G_b as described below.

- $v_{a,1}^u$ is adjacent to $u_{b,2}^{v,a}$, for every variable $v \neq u$ appearing in b. $v_{a,2}^u$ is adjacent to $u_{b,1}^{v,a}$, for every variable $v \neq u$ appearing in b.
- $v_{b,1}^u$ is adjacent to $u_{a,2}^{v,b}$, for every variable $v \neq u$ appearing in a.
- $v_{b,2}^{u}$ is adjacent to $u_{a,1}^{v,b}$, for every variable $v \neq u$ appearing in a.

Finally, we add edges so that $K_1 = \bigcup_{c \in C} K_{c,1}$ is a clique and $K_2 = \bigcup_{c \in C} K_{c,2}$ is a clique. We let G be the resulting graph. We next show that Φ is satisfiable if and only if $ct_{\gamma_{t2}}(G) = 3$.

Before turning to the proof, let us briefly explain the idea behind the construction. In a clause gadget G_c , the vertices $v_{c,1}^{\ell}$ and $v_{c,2}^{\ell}$ should be seen as representing the variable ℓ . As we will show, for any minimum SD set D of G, $D \cap V(G_c)$ contains only one vertex from $V_{c,1} \setminus \{p_{c,1}, q_{c,1}\}$ and one vertex from $V_{c,2} \setminus \{p_{c,2}, q_{c,2}\}$, both corresponding to the same variable ℓ . This variable ℓ can then be set to true to satisfy c. To ensure that the choice in every clause gadget is consistent (that is, if ℓ is chosen in G_c then ℓ should be chosen in G_a , for every clause a containing ℓ), we make use of the vertices $u_{c,i}^{v,a}$, which should be seen as representing the clause a in G_c . By construction, $u_{c,i}^{v,a}$ is adjacent to only $v_{c,i}^v$ (representing v) and $v_{a,(i+1) \mod 2}^u$ (representing u), for $u \neq v$. Thus, if $v_{c,i}^v$ is not in the SD set then $v_{a,i}^v$ is not in the SD set either, as $u_{c,i}^{v,a}$ would otherwise not be dominated.

Claim 36. For any SD D of G and any clause $c \in C$, $|D \cap (V_{c,1} \cup V_{c,2})| \ge 2$.

Proof. Let D be an SD set of G and let $c \in C$ be a clause. Since $q_{c,1}$ should be dominated, $D \cap (V_{c,1} \cup \{p_{c,2}\} \neq \emptyset;$ and similarly, since $q_{c,2}$ should be dominated, $D \cap (V_{c,2} \cup \{p_{c,1}\}) \neq \emptyset$. Suppose first that $p_{c,1} \in D$. Then $D \cap (V_{c,2} \cup V_{c,1} \setminus \{p_{c,1}\}) \neq \emptyset$, as $p_{c,2}$ should be dominated, and so $|D \cap (V_{c,1} \cup V_{c,2})| \ge 2$. We conclude symmetrically if $p_{c,2} \in D$. Otherwise, $D \cap \{p_{c,1}, p_{c,2}\} = \emptyset$ and so, $D \cap (V_{c,i} \setminus \{p_{c,i}\}) \neq \emptyset$ for every $i \in [2]$. _

Claim 37. If $\gamma_{t2}(G) = 2|C|$ then for every minimum SD set D of G and every clause $c \in C$, $D \cap V(G_c) = \{v_{c,1}^{\ell}, v_{c,2}^{\ell}\}$ for some variable ℓ appearing in c.

Proof. Assume that $\gamma_{t2}(G) = 2|C|$ and let D be a minimum SD set of G. Consider a clause $c \in C$ containing variables x, y and z. We contend that $D \cap \{p_{c,1}, q_{c,1}, p_{c,2}, q_{c,2}\} = \emptyset$. Indeed, observe first that by Claim 36, $|D \cap \{p_{c,1}, q_{c,1}, p_{c,2}, q_{c,2}\}| \le 2$. Now if $|D \cap \{p_{c,1}, q_{c,1}, p_{c,2}, q_{c,2}\}| = 2$ then the vertices in $\{t_{c,i}^{x,y}, t_{c,i}^{x,z}, t_{c,i}^{y,z} \mid i \in [2]\}$ are not dominated, as $D \cap (K_1 \cup K_2) = \emptyset$ by Claim 36. Similarly, if $|D \cap \{p_{c,1}, q_{c,1}, p_{c,2}, q_{c,2}\}| = 1$ then by Claim 36, $|D \cap \{v_{c,i}^x, v_{c,i}^y, v_{c,i}^z| i \in [2]\}| = 1$, say $v_{c,i}^\ell \in D$, and so the vertex $t_{c,i}^{a,b}$ with $\ell \notin \{a,b\}$, is not dominated. Thus, $D \cap \{p_{c,1}, q_{c,1}, p_{c,2}, q_{c,2}\} = \emptyset$ which by Claim 36, implies that $|D \cap \{v_{c,i}^x, v_{c,i}^y, v_{c,i}^z| i \in [2]\}| = 2$; and since $q_{c,1}$ and $q_{c,2}$ should be dominated, in fact $|D \cap \{v_{c,1}^x, v_{c,1}^y, v_{c,1}^z\}| = |D \cap \{v_{c,2}^x, v_{c,2}^y, v_{c,2}^z\}| = 1$. Now if $v_{c,1}^u, v_{c,2}^v \in D$ for two distinct variables $u, v \in \{x, y, z\}$, then the vertex $t_{c,1}^{v,w}$ where $w \in \{x, y, z\} \setminus \{u, v\}$, is not dominated. Thus, $D \cap V(G_c) = \{v_{c,1}^{\ell}, v_{c,2}^{\ell}\}$ for some variable $\ell \in \{x, y, z\}$.

Claim 38. Φ is satisfiable if and only if $\gamma_{t2}(G) = 2|C|$.

Proof. Assume first that Φ is satisfiable and consider a truth assignment satisfying Φ . We construct an SD set D of G as follows: for every clause $c \in C$, exactly one variable in c is true, say ℓ , in which case we add $v_{c,1}^{\ell}$ and $v_{c,2}^{\ell}$ to D. Let us show that the constructed set D is indeed an SD set of G. Consider a clause $c \in C$ containing variables x, y and z, and assume without loss of generality that x is true. For every $\ell \in \{x, y, z\}$, let a_{ℓ} and b_{ℓ} be the two other clauses in which ℓ appears. Note that since by construction, $D \cap V(G_c) = \{v_{c,1}^x, v_{c,2}^x\}$ and $d(v_{c,1}^x, v_{c,2}^z) = 2, v_{c,1}^x$ and $v_{c,2}^x$ witness each other. Now it is clear that every vertex in $V_{c,1} \cup V_{c,2} \cup \{u_{c,i}^{x,a_x}, u_{c,i}^{x,b_x} \mid i \in [2]\} \cup \{t_{c,i}^{x,y}, t_{c,i}^{x,z} \mid i \in [2]\}$ is dominated. Furthermore, by construction, $v_{c,1}^x$ is adjacent to $t_{c,2}^{y,z}$, and $v_{c,2}^x$ is adjacent to $t_{c,1}^{y,z}$, and thus, these two vertices are dominated as well. There remains to show that for every $i \in [2]$ and every $\ell \neq x$, the vertices $u_{c,i}^{\ell,a_{\ell}}$ and $u_{c,i}^{\ell,b_{\ell}}$ are dominated. But this readily holds true: for every $\ell \in \{y, z\}$ and every $c_{\ell} \in \{a_{\ell}, b_{\ell}\}$, since ℓ is false, $D \cap V(G_{c_{\ell}}) = \{v_{c_{\ell},1}^v, v_{c_{\ell},2}^v\}$ for some variable $v \neq \ell$ appearing in c_{ℓ} ; and by construction, $v_{c_{\ell+1}}^v$ is adjacent to $u_{c,2}^{\ell,c_{\ell}}$ and $v_{c_{\ell+2}}^v$ is adjacent to $u_{c,1}^{\ell,c_{\ell}}$. Thus, D is an SD set of G and since |D| = 2|C|, we conclude by Claim 36 that D is minimum.

Conversely, assume that $\gamma_{t2}(G) = 2|C|$ and let D be a minimum SD set of G. Then by Claim 37, $D \cup (K_1 \cup K_2) = \emptyset$ and for every clause $c \in C$, $D \cap V(G_c) = \{v_{c,1}^{\ell}, v_{c,2}^{\ell}\}$ for some variable ℓ contained in c. We claim that if $a, b \in C$ are two clauses containing a same variable ℓ and $D \cap V(G_a) = \{v_{a,1}^{\ell}, v_{a,2}^{\ell}\}$, then $D \cap V(G_b) = \{v_{b,1}^{\ell}, v_{b,2}^{\ell}\}$. Indeed, suppose for a contradiction that $D \cap V(G_b) = \{v_{b,1}^{\ell}, v_{b,2}^{\ell}\}$ for some variable $v \neq \ell$, and let $p, q \in X$ be the other two variables appearing in a. Then $u_{b,1}^{\ell,a}$ and $u_{b,2}^{\ell,a}$ are not dominated: indeed, $N(u_{b,1}^{\ell,a}) \setminus (K_1 \cup K_2) = \{v_{b,1}^{\ell}, v_{a,2}^{p}, v_{a,2}^{q}\}$ and $N(u_{b,2}^{\ell,a}) \setminus (K_1 \cup K_2) = \{v_{b,2}^{\ell}, v_{a,1}^{p}, v_{a,1}^{q}\}$, and so $D \cap N(u_{b,1}^{\ell,a}) = D \cap N(u_{b,2}^{\ell,a}) = \emptyset$. Thus, $V(G_b) = \{v_{b,1}^{\ell}, v_{a,2}^{\ell}\}$ as claimed. It follows that the truth assignment obtained by setting a variable ℓ to true if $v_{c,1}^{\ell} \in D$ for some clause c containing ℓ , and to false otherwise, satisfies Φ .

Claim 39. $ct_{\gamma_{t2}}(G) = 3$ if and only if $\gamma_{t2}(G) = 2|C|$.

Proof. Assume first that $ct_{\gamma_{t2}}(G) = 3$ and let D be a minimum SD set of G. If there exists a clause $c \in C$ such that $|D \cap (V_{c,1} \cup V_{c,2})| \geq 3$ then $D \cap (V_{c,1} \cup V_{c,2})$ contains a friendly triple, a contradiction to Theorem 1.7. Thus, $|D \cap (V_{c,1} \cup V_{c,2})| \leq 2$ for every clause $c \in C$, and we conclude by Claim 36 that, in fact, equality holds. Now suppose that $D \cap K \neq \emptyset$, say $u \in D \cap (K_{c,1} \cup K_{c,2})$ for some clause $c \in C$. If u has a neighbour $v \in D \cap (V_{c,1} \cup V_{c,2})$, then u, v, w where $w \in D \cap (V_{c,1} \cup V_{c,2} \setminus \{v\})$, is a friendly triple, a contradiction to ??. Thus, $N(u) \cap (D \cup (V_{c,1} \cup V_{c,2})) = \emptyset$. Now let $v \in V_{c,1} \in V_{c,2}$ be a neighbour of u, and let $w_1, w_2 \in D \cap (V_{c,1} \cup V_{c,2})$. Then the vertices in $\{v, w_1, w_2\}$ are pairwise at distance at most two, and at least two of them must be adjacent. However, if $w_1w_2 \in E(G)$ then $D \cup \{v\}$ contains the O_7 u, v, w_1, w_2 ; and if $vw_i \in E(G)$ for some $i \in [2]$, then $D \cup \{v\}$ contains the O_7 u, v, w_1, w_2 ; and if $vw_i \in E(G)$ for some $i \in [2]$, then $D \cup \{v\}$ contains the O_7 u, v, w_1, w_2 and if $vw_i \in E(G)$ for some $i \in [2]$.

Conversely, assume that $\gamma_{t2}(G) = 2|C|$ and let D be a minimum SD set of G. Then by Claim 37, $D \cap (K_1 \cup K_2) = \emptyset$ and for every clause $c \in C$, $D \cap V(G_c) = \{v_{c,1}^{\ell}, v_{c,2}^{\ell}\}$ for some variable ℓ contained in c. Since for every clause $c \in C$ and every variable $v \in X$ contained in c, $d(v_{c,1}^v, v_{c,2}^v) = 2$, it follows that D is an independent set and thus, D contains no friendly triple. Furthermore, as shown in the proof of Claim 38, if two clauses $a, b \in C$ contain a same variable ℓ and $D \cap V(G_a) = \{v_{a,1}^\ell, v_{a,2}^\ell\}$, then $D \cap V(G_b) = \{v_{b,1}^\ell, v_{b,2}^\ell\}$. It follows that for any two clauses $a, b \in C$ containing a same variable, $d(D \cap V(G_a), D \cap V(G_b)) \ge 3$; and since for any two clauses $a, b \in C$ with no common variable, $d(V_{a,1} \cup V_{a,2}, V_{b,1} \cup V_{b,2}) \ge 3$, every vertex in D has a unique witness. Thus, the following hold.

Observation 10. If $\gamma_{t2}(G) = 2|C|$ then for every minimum SS set D of G, the following hold. (i) D is an independent set.

(ii) For every $x \in D$, $|w_D(x)| = 1$.

Now suppose for a contradiction that G has an SD set D of size $\gamma_{t2}(G) + 1$ containing an ST-configuration (see Theorem 1.7). Then by Claim 36, either there exists a clause $c \in C$ such that $|D \cap (V_{c,1} \cup V_{c,2})| = 3$, or $|D \cap (K_1 \cup K_2)| = 1$. We next distinguish these two cases.

Case 1. There exists a clause $c \in C$ such that $|D \cap (V_{c,1} \cup V_{c,2})| = 3$. Let us first show that for every clause $a \in C \setminus \{c\}$, $D \cap (V_{a,1} \cup V_{a,2})$ contains no edge. Suppose for a contradiction that there exists a clause $a \in C \setminus \{a\}$ such that $D \cap (V_{a,1} \cup V_{a,2})$ contains an edge, and let $x, y, z \in X$ be the three variables appearing in a. Since by Claim 36, $|D \cap (V_{a,1} \cup V_{a,2})| = 2$, necessarily $D \cap \{q_{a,i}, v_{a,i}^x, v_{a,i}^y, v_{a,i}^z\} = \emptyset$ for some $i \in [2]$. But $D \cap K_i = \emptyset$ and $|D \cap \{v_{a,j}^x, v_{a,j}^y, v_{a,j}^z\}| \le 2$ for $j \neq i$ and so, one at least of $t_{a,i}^{x,y}, t_{a,i}^{x,z}$ and $t_{a,i}^{y,z}$ is not dominated, a contradiction.

 $D \cap \{q_{a,i}, v_{a,i}^x, v_{a,i}^y, v_{a,i}^z\} = \emptyset \text{ for some } i \in [2]. \text{ But } D \cap K_i = \emptyset \text{ and } |D \cap \{v_{a,i}^x, v_{a,j}^y, v_{a,j}^z\}| \le 2 \text{ for } j \neq i \text{ and so, one at least of } t_{a,i}^{x,y}, t_{a,i}^{x,z} \text{ and } t_{a,i}^{y,z} \text{ is not dominated, a contradiction.} \\ \text{Now let } x, y, z \in X \text{ be the three variables appearing in } c. We contend that for every } i \in [2], \\ D \cap \{v_{c,i}^x, v_{c,i}^y, v_{c,i}^z\} \neq \emptyset. \text{ Indeed, suppose for a contradiction that } D \cap \{v_{c,i}^x, v_{c,i}^y, v_{c,i}^z\} = \emptyset \text{ for some } i \in [2], \text{ say } i = 1 \text{ without loss of generality. Then since every vertex in } \{t_{c,1}^{x,y}, t_{c,1}^{x,z}, t_{c,1}^{y,z}\} \text{ must be dominated and } D \cap K_1 = \emptyset, \text{ necessarily } v_{c,2}^\ell \in D \text{ for every } \ell \in \{x, y, z\}. \text{ But then, } D \cap \{v_{c,1} \cup V_{c,2}\} \neq \emptyset: \text{ indeed, if } p_{c,1}, p_{c,2} \notin D \text{ then } D \cap \{v_{c,1} \cup V_{c,2}\} \text{ contains only one edge; and since } D \setminus \{v_{c,1} \cup V_{c,2}\} \text{ contains no edge, as shown above, } D \text{ contains no ST-configuration, a contradiction to our assumption. Now assume without loss of generality that <math>p_{c,1} \in D$, and for $i \in [2], \text{ let } w_i \in D \cap \{v_{c,i}^x, v_{c,i}^y, v_{c,i}^z\}. \text{ Then since } D \setminus \{p_{c,1}\} \text{ is an SD set of } G, \text{ it follows from Lemma 2.3 and Observation 10 that } D \text{ contains an } O_6; \text{ but } w_1p_{c,1}w_2 \text{ is the only } P_3 \text{ contained in } D \text{ and } d(p_{c,1}, D \setminus \{w_1, w_2, p_{c,1}\}) \geq 3, \text{ a contradiction.}$

Case 2. $|D \cap (K_1 \cup K_2)| = 1$. Assume without loss of generality that $D \cap K_1 \neq \emptyset$. Let us first show that for every clause $c \in C$, $|D \cap (V_{c,1} \setminus \{p_{c,1}\})| = D \cap (V_{c,2} \setminus \{p_{c,2}\}) = 1$. Suppose for a contradiction that there exists a clause $c \in C$ such that $D \cap (V_{c,i} \setminus \{p_{c,i}\}) = \emptyset$ for some $i \in [2]$, and let $x, y, z \in X$ be the three variables appearing in c. Then $i \neq 2$: indeed, since $D \cap K_2 = \emptyset$ and $|D \cap \{v_{c,1}^i v_{c,1}^y, v_{c,1}^z\}| \leq 2$, if i = 2 then one of $t_{c,2}^{x,y}, t_{c,2}^{x,z}, t_{c,2}^{y,z}$ is not dominated. Thus, it must be that i = 1; but then, $D \cap \{p_{c,1}, p_{c,2}\} \neq \emptyset$ as $q_{c,1}$ should be dominated, which implies that $|D \cap \{q_{c,2}, v_{c,2}^x, v_{c,2}^y, v_{c,2}^z\}| \leq 1$, and so, one of $t_{c,2}^{x,y}, t_{c,2}^{x,z}$ and $t_{c,2}^{y,z}$ is not dominated, a contradiction. Thus, $|D \cap (V_{c,1} \setminus \{p_{c,1}\})| = D \cap (V_{c,2} \setminus \{p_{c,2}\})| = 1$ for every clause $c \in C$; in particular, $D \cap (V_{c,1} \cup V_{c,2})$ contains no edge.

Now let $u \in D \cap K_1$, say $u \in K_{c,1}$ for some clause $c \in C$. Then $|N(u) \cap D| > 1$: if not then u is the endvertex of at most one edge in D; and since $D \setminus \{u\}$ contains no edge, as shown above, D then contains no ST-configuration. Suppose first that $u \in \{t_{c,1}^{x,y}, t_{c,1}^{x,z}, t_{c,1}^{y,z}\}$, say $u = t_{c,1}^{x,y}$ without loss of generality. Then since $|D \cap \{v_{c,1}^{,}v_{c,1}^{,}, v_{c,1}^{,z}\}| \leq 1$ as shown above, it must be that $D \cap \{v_{c,1}^{,}, v_{c,1}^{,y}\} \neq \emptyset$ and $v_{c,2}^{z} \in D$ (u would otherwise has at most one neighbour in D); but then, $t_{c,2}^{x,y}$ is not dominated, a contradiction. Second, suppose that $u \in K_{c,1} \setminus \{t_{c,1}^{x,y}, t_{c,1}^{x,z}, t_{c,1}^{y,z}\}$, say $u = u_{c,1}^{x,a}$ where $a \neq c$ is a clause containing x. Then since $N(u_{c,1}^{x,a} \subseteq V_{c,1} \cup V_{a,2}$ and $|D \cap V_{c,1}| = |D \cap V_{a,2}| = 1$ as shown above, it must be that $D \cap V_{c,1} = \{v_{c,1}^x\}$ and $D \cap V_{a,2} = \{v_{a,2}^\ell\}$ for some variable $\ell \neq x$ contained in a; but then, $u_{a,2}^{x,c}$ is not dominated, a contradiction which concludes the proof.

Now by Claims 38 and 39, Φ is satisfiable if and only if $ct_{\gamma_{t_2}}(G) = 3$. There remains to show that G is $3P_4$ -free. To see this, observe that for any clause $c \in C$, $G[V_{c,1} \cup V_{c,2}]$ is P_4 -free. Thus, if G contains a P_4 P, then $V(P) \cap (K_1 \cup K_2) \neq \emptyset$; but K_1 are K_2 are both cliques and so, G contains no induced $3P_4$.

4.3 Proof of Theorem 1.4

Let *H* be a graph. If *H* contains a cycle then CONTRACTION NUMBER($\gamma_{t2}, 2$) is NP-hard on *H*-free graphs by Corollary 4.10. Assume henceforth that *H* is a forest. If *H* contains a vertex of degree at least three then CONTRACTION NUMBER($\gamma_{t2}, 2$) is NP-hard on *H*-free graphs by Lemma 4.7. Suppose therefore that *H* is a linear forest. If *H* has a connected component on at least six vertices then CONTRACTION NUMBER($\gamma_{t2}, 2$) is (co)NP-hard on *H*-free graphs by Lemma 4.6. Thus we may assume that every connected component of *H* has size at most five. Now suppose that *H* has a connected component on at least four vertices, Then if every other connected component of *H* has size one, CONTRACTION NUMBER($\gamma_{t2}, 2$) is polynomial-time solvable on *H*-free graphs by Lemma 4.5; otherwise CONTRACTION NUMBER($\gamma_{t2}, 2$) is (co)NP-hard on *H*-free graphs by Lemma 4.6. Assume finally that every connected component of H has at most three vertices. If H has at least two connected component of size three then CONTRACTION NUMBER($\gamma_{t2}, 2$) is (co)NP-hard on H-free graphs by Lemma 4.6. Otherwise H has at most one connected component of size three in which case CONTRACTION NUMBER($\gamma_{t2}, 2$) is polynomial-time solvable on H-free graphs by Lemma 4.5 which concludes the proof.

References

- C. Bazgan, S. Toubaline, and Z. Tuza. The most vital nodes with respect to independent set and vertex cover. *Discrete Applied Mathematics*, 159(17):1933–1946, 2011.
- [2] C. Bazgan, C. Bentz, C. Picouleau, and B. Ries. Blockers for the stability number and the chromatic number. *Graphs and Combinatorics*, 31(1):73–90, 2015.
- [3] C. Bentz, M.-C. Costa, D. de Werra, C. Picouleau, and B. Ries. Blockers and transversals in some subclasses of bipartite graphs: when caterpillars are dancing on a grid. *Discrete Mathematics*, 310(1):132 – 146, 2010.
- [4] G. Chen, J. Ehrenmüller, C. G. Fernandes, C. G. Heise, S. Shan, P. Yang, and A. N. Yates. Nonempty intersection of longest paths in series-parallel graphs. *Discrete Mathematics*, 340 (3):287–304, 2017. doi: https://doi.org/10.1016/j.disc.2016.07.023.
- [5] O. Y. Diner, D. Paulusma, C. Picouleau, and B. Ries. Contraction blockers for graphs with forbidden induced paths. In Algorithms and Complexity - 9th International Conference (CIAC 2015), volume 9079 of Lecture Notes in Computer Science, pages 194–207, 2015.
- [6] O. Y. Diner, D. Paulusma, C. Picouleau, and B. Ries. Contraction and deletion blockers for perfect graphs and H-free graphs. *Theoretical Computer Science*, 746:49–72, 2018.
- [7] E. Galby, P. T. Lima, and B. Ries. Blocking dominating sets for H-Free graphs via edge contractions. In 30th International Symposium on Algorithms and Computation (ISAAC 2019), volume 149 of Leibniz International Proceedings in Informatics (LIPIcs), pages 21:1– 21:14, 2019.
- [8] E. Galby, A. Munaro, and B. Ries. Semitotal domination: New hardness results and a polynomial-time algorithm for graphs of bounded mim-width. *Theoretical Computer Science*, 814, 2020.
- [9] E. Galby, P. T. Lima, and B. Ries. Reducing the domination number of graphs via edge contractions and vertex deletions. *Discrete Mathematics*, 344(1):112–169, 2021.
- [10] E. Galby, F. Mann, Lima. P.T., and B. Ries. Using edge contractions to reduce the semitotal domination number. *manuscript*, 2021.
- [11] E. Galby, F. Mann, and B. Ries. Blocking total dominating sets via edge contractions. *Theoretical Computer Science*, 877:18–35, 2021.
- [12] J. Huang and J.-M. Xu. Domination and total domination contraction numbers of graphs. Ars Combinatoria, 94:431–443, 2010.
- [13] C. Keller and M. A. Perles. Blockers for simple hamiltonian paths in convex geometric graphs of even order. *Discrete & Computational Geometry*, 60(1):1–8, 2018.
- [14] P. T. Lima, V. F. dos Santos, I. Sau, and U. S. Souza. Reducing graph transversals via edge contractions. In 45th International Symposium on Mathematical Foundations of Computer Science (MFCS 2020), volume 170 of Leibniz International Proceedings in Informatics (LIPIcs), pages 64:1–64:15, 2020.
- [15] C. Moore and J. M. Robson. Hard tiling problems with simple tiles. Discrete & Computational Geometry, 26(4):573–590, 2001.

- [16] F. Nasirian, F. M. Pajouh, and J. Namayanja. Exact algorithms for the minimum cost vertex blocker clique problem. *Computers & Operations Research*, 103:296–309, 2019.
- [17] F. M. Pajouh, V. Boginski, and E. Pasiliao. Minimum vertex blocker clique problem. Networks, 64:48–64, 2014.
- [18] F. M. Pajouh, J. L. Walteros, V. Boginski, and E. L. Pasiliao. Minimum edge blocker dominating set problem. *European Journal of Operational Research*, 247(1):16–26, 2015.
- [19] D. Paulusma, C. Picouleau, and B. Ries. Reducing the clique and chromatic number via edge contractions and vertex deletions. In 4th International Symposium on Combinatorial Optimisation (ISCO 2016), volume 9849 of Lecture Notes in Computer Science, pages 38–49, 2016.
- [20] D. Paulusma, C. Picouleau, and B. Ries. Blocking independent sets for H-free graphs via edge contractions and vertex deletions. In 14th International Conference in Theory and Applications of Models of Computation (TAMC 2017), volume 10185 of Lecture Notes in Computer Science, pages 470–483, 2017.
- [21] D. Paulusma, C. Picouleau, and B. Ries. Critical vertices and edges in H-free graphs. Discrete Applied Mathematics, 257:361 – 367, 2019.
- [22] D. Rautenbach and J.-S. Sereni. Transversals of longest paths and cycles. SIAM Journal on Discrete Mathematics, 28(1):335–341, 2014.
- [23] T. J. Schaefer. The complexity of satisfiability problems. In Proceedings of the Tenth Annual ACM Symposium on Theory of Computing, STOC '78, pages 216–226, 1978.
- [24] R. Zenklusen. Network flow interdiction on planar graphs. Discrete Applied Mathematics, 158(13):1441–1455, 2010.
- [25] R. Zenklusen. Matching interdiction. Discrete Applied Mathematics, 158(15):1676–1690, 2010.