

# COMPARISON OF LIMIT SHAPES FOR BERNOULLI FIRST-PASSAGE PERCOLATION

NAOKI KUBOTA AND MASATO TAKEI

ABSTRACT. We consider Bernoulli first-passage percolation on the  $d$ -dimensional hypercubic lattice with  $d \geq 2$ . The passage time of edge  $e$  is 0 with probability  $p$  and 1 with probability  $1-p$ , independently of each other. Let  $p_c$  be the critical probability for percolation of edges with passage time 0. When  $0 \leq p < p_c$ , there exists a nonrandom, nonempty compact convex set  $\mathcal{B}_p$  such that the set of vertices to which the first-passage time from the origin is within  $t$  is well-approximated by  $t\mathcal{B}_p$  for all large  $t$ , with probability one. The aim of this paper is to prove that for  $0 \leq p < q < p_c$ , the Hausdorff distance between  $\mathcal{B}_p$  and  $\mathcal{B}_q$  grows linearly in  $q-p$ . Moreover, we mention that the approach taken in the paper provides a lower bound for the expected size of the intersection of geodesics, that gives a nontrivial consequence for the *critical case*.

## 1. INTRODUCTION

First-passage percolation was first introduced by Hammersley and Welsh [9]. In this paper, let  $d \geq 2$  and consider Bernoulli first-passage percolation on the  $d$ -dimensional hypercubic lattice  $\mathbb{L}^d = (\mathbb{Z}^d, \mathbb{E}^d)$ : The family  $\{t(e)\}_{e \in \mathbb{E}^d}$  of independent random variables satisfies that

$$P_p(t(e) = 0) = p \quad \text{and} \quad P_p(t(e) = 1) = 1 - p \quad \text{for each } e \in \mathbb{E}^d.$$

The expectation of a random variable  $X$  with respect to  $P_p$  is denoted by  $E_p[X]$ . For a self-avoiding path  $\gamma$  from  $\mathbf{x} \in \mathbb{Z}^d$  to  $\mathbf{y} \in \mathbb{Z}^d$ , we define

$$t(\gamma) := \sum_{e \in \gamma} t(e).$$

For each  $\mathbf{x} \in \mathbb{R}^d$ , we can find a unique  $\mathbf{x}' \in \mathbb{Z}^d$  with  $\mathbf{x} \in \mathbf{x}' + [0, 1)^d$ . The first-passage time from  $\mathbf{x} \in \mathbb{R}^d$  to  $\mathbf{y} \in \mathbb{R}^d$  is defined by

$$T(\mathbf{x}, \mathbf{y}) := \inf_{\gamma: \mathbf{x}' \rightarrow \mathbf{y}'} t(\gamma).$$

The wet region at time  $t \geq 0$  is defined by

$$B(t) := \{\mathbf{y} \in \mathbb{R}^d : T(\mathbf{O}, \mathbf{y}) \leq t\},$$

where  $\mathbf{O} := (0, 0, \dots, 0)$ . Let

$$p_c = p_c(d) := \inf \left\{ p \in [0, 1] : \begin{array}{l} \text{there exists an infinite self-avoiding path} \\ \text{of edges } e \text{ with } t(e) = 0 \end{array} \right\}.$$

---

N.K. was supported by JSPS KAKENHI Grant Number JP20K14332. M.T. is partially supported by JSPS KAKENHI Grant Numbers JP19H01793, JP19K03514 and JP22K03333.

It is known that when  $0 \leq p < p_c$ , there exists a nonrandom, nonempty compact convex set  $\mathcal{B}_p$  such that for any  $\varepsilon > 0$ ,

$$P_p \left( (1 - \varepsilon)\mathcal{B}_p \subset \frac{1}{t}B(t) \subset (1 + \varepsilon)\mathcal{B}_p \text{ for all large } t \right) = 1. \quad (1.1)$$

This is called the shape theorem, a kind of law of large numbers for the wet region. For fundamental results on first-passage percolation (including the shape theorem above), refer to Kesten [13] and Auffinger, Damron and Hanson [1].

In this paper, we compare limit shapes  $\mathcal{B}_p$ ,  $0 \leq p < p_c$ , and prove that a certain distance between  $\mathcal{B}_p$  and  $\mathcal{B}_q$  is of order  $|q - p|$ . Moreover, the method used to compare limit shapes also gives a nontrivial property for the size of the intersection of paths realizing the first-passage time.

## 2. RESULTS

For  $p \in [0, 1]$  and  $\mathbf{x} \in \mathbb{Z}^d$ , the following limit exists (see Chapter 2 in [13]):

$$\mu_p(\mathbf{x}) := \lim_{n \rightarrow \infty} \frac{E_p[T(\mathbf{O}, n\mathbf{x})]}{n} = \inf_{n \geq 1} \frac{E_p[T(\mathbf{O}, n\mathbf{x})]}{n}. \quad (2.1)$$

It is known that  $\mu_p(\mathbf{x})$  can be extended to a seminorm on  $\mathbb{R}^d$ , and  $\mu_p(\mathbf{x})$  is called the time constant. In particular, for  $0 \leq p < p_c$ ,  $\mu_p(\mathbf{x}) > 0$  for any  $\mathbf{x} \in \mathbb{R}^d \setminus \{\mathbf{O}\}$ , and the limit shape  $\mathcal{B}_p$  in (1.1) is the unit ball for the norm  $\mu_p(\cdot)$ :

$$\mathcal{B}_p = \{\mathbf{x} \in \mathbb{R}^d : \mu_p(\mathbf{x}) \leq 1\}.$$

Note that if  $0 \leq p < q \leq 1$ , then  $\mu_p(\mathbf{x}) \geq \mu_q(\mathbf{x})$  for any  $\mathbf{x} \in \mathbb{R}^d$ . For  $p \geq p_c$ , it is known that  $\mu_p(\mathbf{x}) = 0$  for any  $\mathbf{x} \in \mathbb{R}^d$ , and the limit shape  $\mathcal{B}_p = \mathbb{R}^d$ , in that for any compact  $K \subset \mathbb{R}^d$ ,  $K \subset t^{-1}B(t)$  for all large  $t$  with probability one. We can see that  $\mathcal{B}_p \subset \mathcal{B}_q$  if  $0 \leq p < q \leq 1$ .

For  $\mathbf{e}_1 := (1, 0, \dots, 0)$ , the general theory developed in [5, 6, 13] tells us that  $\mu_p(\mathbf{e}_1)$  is continuous in  $p$ . Moreover, from another general result by van den Berg and Kesten [2],  $\mu_p(\mathbf{e}_1)$  is strictly decreasing in  $p \in [0, p_c]$ . Furthermore, Wu and Feng [15] derived the following concrete lower bound for the decrease: If  $0 \leq p < q < p_c$ , then

$$\mu_p(\mathbf{e}_1) - \mu_q(\mathbf{e}_1) \geq \frac{\mu_q(\mathbf{e}_1)}{1 - q} \cdot (q - p). \quad (2.2)$$

Our first theorem extends (2.2) to all directions.

**Theorem 2.1.** If  $0 \leq p < q < p_c$ , then for each  $\mathbf{x} \in \mathbb{R}^d$ ,

$$\mu_p(\mathbf{x}) - \mu_q(\mathbf{x}) \geq \frac{\mu_q(\mathbf{x})}{1 - q} \cdot (q - p). \quad (2.3)$$

The next theorem gives a modulus of continuity of  $\mu_p(\mathbf{x})$  in  $p$ .

**Theorem 2.2.** Assume that  $0 < q_0 < p_c$ . There exists a constant  $K = K(q_0) > 0$  such that for any  $0 < p < q \leq q_0$  and any  $\mathbf{x} \in \mathbb{R}^d$ ,

$$\mu_p(\mathbf{x}) - \mu_q(\mathbf{x}) \leq \frac{K\mu_q(\mathbf{x})}{q} \cdot (q - p). \quad (2.4)$$

In particular,  $\mu_p(\mathbf{x})$  is Lipschitz continuous over any closed subinterval  $[p_0, q_0]$  of  $(0, p_c)$ .

For  $A \subset \mathbb{R}^d$  and  $\delta > 0$ , set

$$N_\delta(A) := \{\mathbf{y} \in \mathbb{R}^d : \|\mathbf{y} - \mathbf{x}\|_2 < \delta \text{ for some } \mathbf{x} \in A\}.$$

For  $A, B \subset \mathbb{R}^d$ , the Hausdorff distance between  $A$  and  $B$  is defined by

$$d_{\mathcal{H}}(A, B) := \inf\{\delta > 0 : A \subset N_\delta(B) \text{ and } B \subset N_\delta(A)\}.$$

As a corollary to Theorems 2.1 and 2.2, we can derive linear bounds for  $d_{\mathcal{H}}(\mathcal{B}_p, \mathcal{B}_q)$  in  $|q - p|$ .

**Corollary 2.3.** Assume that  $0 < p_0 < q_0 < p_c$ . We can find a constant  $C > 0$  such that for  $p_0 \leq p < q \leq q_0$ ,

$$q - p \leq d_{\mathcal{H}}(\mathcal{B}_p, \mathcal{B}_q) \leq C(q - p).$$

We review literatures closely related to our main results. In bond percolation on  $\mathbb{L}^d$  with density  $\rho$  (i.e. each edge is open with probability  $\rho$ ), if  $\rho > p_c$ , then there exists a unique infinite open cluster  $\mathcal{C}_\infty$  with probability one. The first-passage percolation on  $\mathcal{C}_\infty$  is studied by Garet, Marchand, Procaccia, and Th  ret [11]. They show that the continuity of the time constant and the limit shape with respect to the probability law of  $t(e)$ . This model is equivalent to allow  $t(e)$  to be  $+\infty$  in first-passage percolation on  $\mathbb{L}^d$ . In the special case where

$$P_\rho(t(e) = 1) = \rho \quad \text{and} \quad P_\rho(t(e) = +\infty) = 1 - \rho,$$

the regularity of the time constant and the limit shape with respect to  $\rho \in (p_c, 1]$  is explored by Dembin [8] and Cerf and Dembin [3]. It is shown in [3] that for  $p_0 > p_c$ , the time constant is Lipschitz continuous on  $\rho \in [p_0, 1]$ . Note that although this result looks similar to our Theorem 2.2, the results in [3] do not imply it.

The proofs of our main results will be given in Section 3, where geometric properties of geodesics play important roles. As an application of ideas in the proof of Theorem 2.1, we can obtain a new result on the expected size of intersection of geodesics for the *critical* case. It will be discussed in Section 4.

### 3. PROOFS

**3.1. Proof of Theorem 2.1.** Fix  $n \in \mathbb{N}$ ,  $\mathbf{x} \in \mathbb{Z}^d$ , and  $N \in \mathbb{N}$  with  $N > \|\mathbf{n}\mathbf{x}\|_\infty$ . The first-passage time over paths from  $\mathbf{O}$  to  $\mathbf{n}\mathbf{x}$  contained in  $S(N) := [-N, N]^d$  is denoted by  $T^N(\mathbf{O}, \mathbf{n}\mathbf{x})$ . A path  $\gamma$  from  $\mathbf{O}$  to  $\mathbf{n}\mathbf{x}$  with  $t(\gamma) = T^N(\mathbf{O}, \mathbf{n}\mathbf{x})$  is called a geodesic for  $T^N(\mathbf{O}, \mathbf{n}\mathbf{x})$ . The edge intersection of geodesics for  $T^N(\mathbf{O}, \mathbf{n}\mathbf{x})$  is denoted by  $\overline{\text{GEO}}^N(\mathbf{O}, \mathbf{n}\mathbf{x})$ .

**Lemma 3.1.** For  $p \in (0, 1)$ ,

$$-\frac{d}{dp} E_p[T^N(\mathbf{O}, \mathbf{n}\mathbf{x})] = \frac{1}{1-p} E_p[\#\{e \in \overline{\text{GEO}}^N(\mathbf{O}, \mathbf{n}\mathbf{x}) : t(e) = 1\}]. \quad (3.1)$$

*Proof.* Let  $A_{n,k} := \{T^N(\mathbf{O}, n\mathbf{x}) \geq k\}$ . We have

$$E_p[T^N(\mathbf{O}, n\mathbf{x})] = \sum_{k=1}^{n\|\mathbf{x}\|_1} P_p(A_{n,k}). \quad (3.2)$$

The event  $A_{n,k}$  depends on the edges in  $S(N)$ , and decreasing in the configuration of passage times. Define

$$\mathcal{N}(A_{n,k}) := \#\{e \in S(N) : e \text{ is pivotal for } A_{n,k}\}.$$

By Russo's formula (see e.g. Theorem 2.25 in [12]),

$$\begin{aligned} -\frac{d}{dp}P_p(A_{n,k}) &= E_p[\mathcal{N}(A_{n,k})] \\ &= \sum_{e \in S(N)} P_p(e \text{ is pivotal for } A_{n,k}) \\ &= \frac{1}{1-p} \sum_{e \in S(N)} P_p(\{e \text{ is pivotal for } A_{n,k}\} \cap A_{n,k}) \\ &= \frac{1}{1-p} E_p[\mathcal{N}(A_{n,k}) : A_{n,k}]. \end{aligned}$$

Noting that

$$\mathcal{N}(A_{n,k}) = \begin{cases} \#\{e \in \overline{\text{GEO}}^N(\mathbf{O}, n\mathbf{x}) : t(e) = 1\} & \text{if } T^N(\mathbf{O}, n\mathbf{x}) = k, \text{ and} \\ 0 & \text{if } T^N(\mathbf{O}, n\mathbf{x}) > k, \end{cases}$$

we have

$$\begin{aligned} -\frac{d}{dp}P_p(A_{n,k}) &= \frac{1}{1-p} E_p[\#\{e \in \overline{\text{GEO}}^N(\mathbf{O}, n\mathbf{x}) : t(e) = 1\} : T^N(\mathbf{O}, n\mathbf{x}) = k]. \quad (3.3) \end{aligned}$$

Now (3.1) follows from (3.2) and (3.3).  $\square$

Since

$$\#\{e \in \overline{\text{GEO}}^N(\mathbf{O}, n\mathbf{x}) : t(e) = 1\} \geq t(\gamma) = T^N(\mathbf{O}, n\mathbf{x}) \quad (3.4)$$

for any geodesic  $\gamma$  for  $T^N(\mathbf{O}, n\mathbf{x})$ , we have

$$-\frac{d}{dp}E_p[T^N(\mathbf{O}, n\mathbf{x})] \geq \frac{1}{1-p} \cdot E_p[T^N(\mathbf{O}, n\mathbf{x})],$$

which implies that

$$\frac{d}{dp} \left( \frac{E_p[T^N(\mathbf{O}, n\mathbf{x})]}{1-p} \right) \leq 0 \quad \text{for } p \in (0, 1).$$

Thus we obtain

$$\frac{E_p[T^N(\mathbf{O}, n\mathbf{x})]}{1-p} \geq \frac{E_q[T^N(\mathbf{O}, n\mathbf{x})]}{1-q} \quad \text{for } 0 \leq p < q < 1.$$

Letting  $N \rightarrow \infty$ , dividing both sides by  $n$ , and then letting  $n \rightarrow \infty$ , we have

$$\frac{\mu_p(\mathbf{x})}{1-p} \geq \frac{\mu_q(\mathbf{x})}{1-q},$$

which is equivalent to (2.3).  $\square$

**3.2. Proof of Theorem 2.2.** Our proof is inspired by that of Lemma 3 in Zhang [17]. We begin with a lemma. The geodesic for  $T(\mathbf{O}, \mathbf{x})$  with maximal number of edges is denoted by  $\text{GEO}(\mathbf{O}, \mathbf{x})$ .

**Lemma 3.2.** Let  $q_0 \in (0, p_c)$ . There exists a constant  $K = K(q_0) > 0$  such that for any  $q \in (0, q_0]$  and any  $\mathbf{x} \in \mathbb{Z}^d$ ,

$$E_q[\#\text{GEO}(\mathbf{O}, \mathbf{x})] \leq K \cdot E_q[T(\mathbf{O}, \mathbf{x})]. \quad (3.5)$$

*Proof.* We closely follows that of Proposition 4.7 (1) in Auffinger, Damron and Hanson [1]. There exist constants  $a, C_1 > 0$  depending only on  $q_0$  such that for any  $q \in (0, q_0]$  and any  $m \in \mathbb{N}$ ,

$$P_q \left( \begin{array}{l} \text{there is a self-avoiding path } \gamma \text{ starting from } \mathbf{O} \\ \text{with } \#\gamma \geq m \text{ but } t(\gamma) \leq am \end{array} \right) \leq e^{-C_1 m}. \quad (3.6)$$

To prove this, we have only to choose the constant  $M$  in the proof of Lemma 4.5 in [1] as

$$\sum_{\mathbf{y} \in \mathbb{Z}^d: \|\mathbf{y}\|_\infty = M} P_{q_0}(T(\mathbf{O}, \mathbf{y}) = 0) \leq \frac{1}{2}.$$

We estimate

$$E_q[\#\text{GEO}(\mathbf{O}, \mathbf{x})] \leq \frac{1}{a} E_q[T(\mathbf{O}, \mathbf{x})] + E_q[Y_{\mathbf{x}}],$$

where

$$Y_{\mathbf{x}} := \#\text{GEO}(\mathbf{O}, \mathbf{x}) \cdot 1_{\{T(\mathbf{O}, \mathbf{x}) < a\#\text{GEO}(\mathbf{O}, \mathbf{x})\}}.$$

Following the proof of Lemma 4.6 in [1] together with (3.6), we can find a constant  $C_2 = C_2(q_0) > 0$  such that

$$P_q(Y_{\mathbf{x}} \geq n) \leq C_2 e^{-C_1 n} \quad \text{for } n \in \mathbb{N},$$

which implies that

$$E_q[Y_{\mathbf{x}}] \leq C_3$$

for some  $C_3 = C_3(q_0) > 0$ . Since  $E_{q_0}[T(\mathbf{O}, \mathbf{x})] \rightarrow \infty$  as  $\|\mathbf{x}\|_1 \rightarrow \infty$ , we can take a constant  $C_4 = C_4(q_0)$  such that

$$C_3 \leq C_4 \cdot E_{q_0}[T(\mathbf{O}, \mathbf{x})] \quad \text{for all } \mathbf{x} \in \mathbb{Z}^d.$$

Noting that  $E_q[T(\mathbf{O}, \mathbf{x})] \geq E_{q_0}[T(\mathbf{O}, \mathbf{x})]$ , (3.5) holds true with  $K := \frac{1}{a} + C_4$ .  $\square$

Let  $\{U(e)\}_{e \in \mathbb{E}^d}$  be independent uniform random variables on  $[0, 1]$ , and define

$$t_p(e) := \begin{cases} 0 & (U(e) \leq p), \\ 1 & (U(e) > p) \end{cases} \quad \text{for each } e \in \mathbb{E}^d.$$

Suppose that  $0 < p < q < p_c$  and  $\mathbf{x} \in \mathbb{Z}^d$ . The first-passage time from  $\mathbf{O}$  to  $n\mathbf{x}$  in the configuration  $\{t_p(e)\}_{e \in \mathbb{E}^d}$  is denoted by  $T_p(\mathbf{O}, n\mathbf{x})$ . A geodesic

from  $\mathbf{O}$  to  $n\mathbf{x}$  in the configuration  $\{t_q(e)\}_{e \in \mathbb{E}^d}$  with maximal number is denoted by  $\text{GEO}_q(\mathbf{O}, n\mathbf{x})$ . Then we have

$$\begin{aligned} T_p(\mathbf{O}, n\mathbf{x}) &\leq \sum_{b \in \text{GEO}_q(\mathbf{O}, n\mathbf{x})} t_p(b) \\ &= \sum_{b \in \text{GEO}_q(\mathbf{O}, n\mathbf{x})} t_q(b) + \sum_{b \in \text{GEO}_q(\mathbf{O}, n\mathbf{x})} \{t_p(b) - t_q(b)\} \\ &= T_q(\mathbf{O}, n\mathbf{x}) + \sum_{b \in \text{GEO}_q(\mathbf{O}, n\mathbf{x})} \{t_p(b) - t_q(b)\}. \end{aligned} \quad (3.7)$$

We prove

$$E \left[ \sum_{b \in \text{GEO}_q(\mathbf{O}, n\mathbf{x})} \{t_p(b) - t_q(b)\} \right] \leq \frac{q-p}{q} \cdot E_q[\#\text{GEO}(\mathbf{O}, n\mathbf{x})]. \quad (3.8)$$

The left hand side of (3.8) is

$$\sum_{\gamma: \mathbf{O} \rightarrow n\mathbf{x}} \sum_{b \in \gamma} P(t_p(b) = 1, t_q(b) = 0, \text{GEO}_q(\mathbf{O}, n\mathbf{x}) = \gamma). \quad (3.9)$$

We define a new configuration  $\{t_q^*(e)\}_{e \in \mathbb{E}^d}$  by

$$t_q^*(e) := \begin{cases} 0 & (e = b), \\ t_q(e) & (e \neq b) \end{cases} \quad \text{for each } e \in \mathbb{E}^d.$$

A geodesic from  $\mathbf{O}$  to  $n\mathbf{x}$  in the configuration  $\{t_q^*(e)\}_{e \in \mathbb{E}^d}$  with maximal number is denoted by  $\text{GEO}_q^*(\mathbf{O}, n\mathbf{x})$ . Using independence,

$$\begin{aligned} &P(t_p(b) = 1 \mid t_q(b) = 0, \text{GEO}_q(\mathbf{O}, n\mathbf{x}) = \gamma) \\ &= P(t_p(b) = 1 \mid t_q(b) = 0, \text{GEO}_q^*(\mathbf{O}, n\mathbf{x}) = \gamma) \\ &= P(t_p(b) = 1 \mid t_q(b) = 0) = \frac{q-p}{q}. \end{aligned}$$

As

$$P(t_q(b) = 0, \text{GEO}_q(\mathbf{O}, n\mathbf{x}) = \gamma) \leq P_q(\text{GEO}(\mathbf{O}, n\mathbf{x}) = \gamma),$$

we can see that (3.9) is not more than

$$\begin{aligned} &\sum_{\gamma: \mathbf{O} \rightarrow n\mathbf{x}} \sum_{b \in \gamma} \frac{q-p}{q} \cdot P_q(\text{GEO}(\mathbf{O}, n\mathbf{x}) = \gamma) \\ &= \frac{q-p}{q} \cdot \sum_{\gamma: \mathbf{O} \rightarrow n\mathbf{x}} (\#\gamma) \cdot P_q(\text{GEO}(\mathbf{O}, n\mathbf{x}) = \gamma) \\ &= \frac{q-p}{q} \cdot E_q[\#\text{GEO}(\mathbf{O}, n\mathbf{x})]. \end{aligned}$$

This completes the proof of (3.8). By (3.7) and (3.8), we obtain

$$E_p[T(\mathbf{O}, n\mathbf{x})] \leq E_q[T(\mathbf{O}, n\mathbf{x})] + \frac{q-p}{q} \cdot E_q[\#\text{GEO}(\mathbf{O}, n\mathbf{x})].$$

(2.4) follows from Lemma 3.2 and (2.1).  $\square$

*Proof of Corollary 2.3.* Assume that  $0 \leq p < q < p_c$ . Since  $\mathcal{B}_p \subset \mathcal{B}_q$ ,

$$d_{\mathcal{H}}(\mathcal{B}_p, \mathcal{B}_q) = \inf\{\delta > 0 : \mathcal{B}_q \subset N_{\delta}(\mathcal{B}_p)\}.$$

Define

$$\mathbb{S}^{d-1} := \{\mathbf{u} \in \mathbb{R}^d : \|\mathbf{u}\|_2 = 1\}.$$

We can see that

$$\inf_{\mathbf{u} \in \mathbb{S}^{d-1}} \left\| \frac{\mathbf{u}}{\mu_q(\mathbf{u})} - \frac{\mathbf{u}}{\mu_p(\mathbf{u})} \right\|_2 \leq d_{\mathcal{H}}(\mathcal{B}_p, \mathcal{B}_q) \leq \sup_{\mathbf{u} \in \mathbb{S}^{d-1}} \left\| \frac{\mathbf{u}}{\mu_q(\mathbf{u})} - \frac{\mathbf{u}}{\mu_p(\mathbf{u})} \right\|_2,$$

and

$$\left\| \frac{\mathbf{u}}{\mu_q(\mathbf{u})} - \frac{\mathbf{u}}{\mu_p(\mathbf{u})} \right\|_2 = \left| \frac{1}{\mu_q(\mathbf{u})} - \frac{1}{\mu_p(\mathbf{u})} \right| \cdot \|\mathbf{u}\|_2 = \frac{\mu_p(\mathbf{u}) - \mu_q(\mathbf{u})}{\mu_p(\mathbf{u})\mu_q(\mathbf{u})}.$$

By Theorem 2.1,

$$\begin{aligned} d_{\mathcal{H}}(\mathcal{B}_p, \mathcal{B}_q) &\geq \frac{q-p}{1-q} \cdot \inf_{\mathbf{u} \in \mathbb{S}^{d-1}} \frac{1}{\mu_p(\mathbf{u})} \\ &\geq (q-p) \cdot \inf_{\mathbf{u} \in \mathbb{S}^{d-1}} \frac{1}{\mu_0(\mathbf{u})} = q-p, \end{aligned}$$

where we used  $\mu_0(\mathbf{u}) = \|\mathbf{u}\|_1$ . On the other hand, by Theorem 2.2,

$$d_{\mathcal{H}}(\mathcal{B}_p, \mathcal{B}_q) \leq \frac{K(q-p)}{q} \cdot \sup_{\mathbf{u} \in \mathbb{S}^{d-1}} \frac{1}{\mu_p(\mathbf{u})} \leq \frac{K(q-p)}{p_0} \cdot \sup_{\mathbf{u} \in \mathbb{S}^{d-1}} \frac{1}{\mu_{q_0}(\mathbf{u})}.$$

□

#### 4. A REMARK ON THE INTERSECTION OF GEODESICS

Let  $\mathbf{x} \in \mathbb{Z}^d$ . The proof of (4.10) in Auffinger, Damron and Hanson [1] contains the following inequality:

$$E_p[\#\underline{\text{GEO}}(\mathbf{O}, \mathbf{x})] \geq \frac{p}{1-p} \cdot E_p[\#\{e \in \overline{\text{GEO}}(\mathbf{O}, \mathbf{x}) : t(e) = 1\}] \quad (4.1)$$

holds for *any*  $p \in (0, 1)$  (set  $I = 0$  and  $\delta = 1/4$  in that proof). Similarly to (3.4), we can deduce that

$$E_p[\#\underline{\text{GEO}}(\mathbf{O}, \mathbf{x})] \geq \frac{p}{1-p} \cdot E_p[T(\mathbf{O}, \mathbf{x})]. \quad (4.2)$$

For  $p \in [0, p_c)$ , it follows from (4.2), (3.5) and (2.1) that there exists a constant  $C = C(p) > 0$  such that

$$\frac{p}{1-p} \cdot \mu_p(\mathbf{x})n \leq E_p[\#\underline{\text{GEO}}(\mathbf{O}, n\mathbf{x})] \leq Cn\|\mathbf{x}\|_1 \quad (4.3)$$

for any  $\mathbf{x} \in \mathbb{Z}^d$  and  $n \in \mathbb{N}$ . Nakajima [14] proves that  $E_p[\#\underline{\text{GEO}}(\mathbf{O}, n\mathbf{e}_1)]$  is of order  $n$  if  $t(e)$  is square-integrable and satisfies an additional condition called “useful”. Our proof of (4.3) based on (4.2) gives a simple alternative proof for the Bernoulli case. In view of (4.1), we can see that  $E_p[\#\{e \in \overline{\text{GEO}}(\mathbf{O}, n\mathbf{x}) : t(e) = 1\}]$  is also of order  $n$ , which is a key for proving that  $E_p[\overline{\text{GEO}}(\mathbf{O}, n\mathbf{x})]$  is of order  $n$  (See Theorem 2 in Zhang [17], Theorem 4.12 in Auffinger, Damron and Hanson [1], and Theorem 3 in Nakajima [14]).

We emphasize that (4.2) gives an interesting information about the critical case. Let  $\partial S(n) := \{\mathbf{x} \in \mathbb{Z}^d : \|\mathbf{x}\|_\infty = n\}$ , and the first-passage time from  $\mathbf{O}$  to  $\partial S(n)$  is denoted by  $T(\mathbf{O}, \partial S(n))$ . Under the hypothesis

$$P_{p_c} \left( \begin{array}{c} \text{there exists an infinite self-avoiding path} \\ \text{of edges } e \text{ with } t(e) = 0 \end{array} \right) = 0, \quad (4.4)$$

we have

$$P_{p_c} \left( \lim_{n \rightarrow \infty} T(\mathbf{O}, \partial S(n)) = +\infty \right) = 1,$$

and (4.2) implies that

$$\lim_{n \rightarrow \infty} E_{p_c} [\#\underline{\text{GEO}}(\mathbf{O}, n\mathbf{x})] = +\infty.$$

This suggests that the size of the intersection of geodesics can be large even if  $p = p_c$ . The verification of the hypothesis (4.4) for all  $d \geq 2$  is a long-standing problem in the percolation theory. At the moment (4.4) is shown to be true for  $d = 2$  and for sufficiently large  $d$  (see [10] and [12]). Chayes, Chayes, and Durrett [4] prove that there exist constants  $C_1, C_2 > 0$  such that

$$C_1 \log n \leq E_{p_c}[T(\mathbf{O}, \partial S(n))] \leq C_2 \log n \quad \text{for } n \geq 3.$$

On the other hand, Zhang [16] shows that for sufficiently large  $d$ , there exists a constant  $C > 0$  such that

$$E_{p_c}[T(\mathbf{O}, \partial S(n))] \geq C \log \log n \quad \text{for } n \geq 3.$$

Those results give lower bounds of  $E_{p_c} [\#\underline{\text{GEO}}(\mathbf{O}, n\mathbf{x})]$ . One of important future problems is to give the precise order for  $E_{p_c} [\#\underline{\text{GEO}}(\mathbf{O}, n\mathbf{x})]$  and  $E_{p_c} [\#\overline{\text{GEO}}(\mathbf{O}, n\mathbf{x})]$ . It is proved in Damron and Tang [7] that when  $d = 2$ , there exist constants  $c > 0$  and  $s > 1$  such that for any  $\mathbf{x} \in \mathbb{Z}^2 \setminus \{\mathbf{O}\}$ ,

$$P_{p_c}(N_{\mathbf{O}, \mathbf{x}} \leq \|\mathbf{x}\|_1^s) \leq \frac{1}{c} e^{-\|\mathbf{x}\|_1^c},$$

where  $N_{\mathbf{O}, \mathbf{x}}$  is the minimal number of edges in any geodesic from  $\mathbf{O}$  to  $\mathbf{x}$ .

## REFERENCES

- [1] Auffinger, A., Damron, M., and Hanson, J. (2017). 50 years of first-passage percolation, *University Lecture Series*, **68**, American Mathematical Society.
- [2] van den Berg, J. and Kesten, H. (1993). Inequalities for the time constant in first-passage percolation, *Ann. Appl. Probab.*, **3**, 56–80.
- [3] Cerf, R. and Dembin, B. (2021). The time constant for Bernoulli percolation is Lipschitz continuous strictly above  $p_c$ , arXiv:2101.11858, to appear in *Ann. Probab.*
- [4] Chayes, J.T., Chayes, L., and Durrett, R. (1986). Critical behavior of two-dimensional first-passage times, *J. Stat. Phys.*, **45**, 933–951.
- [5] Cox, J.T. (1980). The time constant of first-passage percolation on the square lattice, *Adv. in Appl. Probab.*, **12**, 864–879.
- [6] Cox, J.T. and Kesten, H. (1981). On the continuity of the time constant of first-passage percolation, *J. Appl. Probab.*, **18**, 809–819.
- [7] Damron, M. and Tang, P. (2019). Superlinearity of geodesic length in 2D critical first-passage percolation, *Sojourns in probability theory and statistical physics. II. Brownian web and percolation, a festschrift for Charles M. Newman*, *Proc. Math. Stat.*, **299**, 101–122, Springer.
- [8] Dembin, B. (2021). Regularity of the time constant for a supercritical Bernoulli percolation, *ESAIM, Probab. Stat.*, **25**, 109–132.

- [9] Hammersley, J.M. and Welsh, D.J.A. (1965). First-passage percolation, subadditive processes, stochastic networks, and generalized renewal theory, *Bernoulli, Bayes, Laplace Anniversary volume*, 61–110, Springer.
- [10] Heydenreich, M. and van der Hofstad, R. (2017). Progress in high-dimensional percolation and random graphs, *CRM Short Courses*, Springer.
- [11] Garet, O., Marchand, R., Procaccia, E.B., and Th  ret, M. (2017). Continuity of the time and isoperimetric constants in supercritical percolation, *Electron. J. Probab.*, **22**, Paper No. 78, 35 p.
- [12] Grimmett, G. (1999). Percolation, Second edition, *Grundlehren math. Wissenschaften*, **321**, Springer.
- [13] Kesten, H. (1986). Aspects of first passage percolation, *  cole d'  t   de probabilit  s de Saint-Flour, XIV – 1984, Lecture Notes in Math.*, **1180**, 125–264, Springer.
- [14] Nakajima, S. (2019). On properties of optimal paths in first-passage percolation, *J. Stat. Phys.*, **174**, 259–275.
- [15] Wu, X.-Y. and Feng, P. (2009). A lower bound for the time constant of first-passage percolation, *Acta Math. Sinica (Chin. Ser.)*, **52**, 495–500; (2008). English version, arXiv:0807.0839
- [16] Zhang, Y. (1999). Double behavior of critical first-passage percolation, *Perplexing problems in probability, Progr. Probab.*, **44**, 143–158, Birkh  user.
- [17] Zhang, Y. (2006). The divergence of fluctuations for the shape of first passage percolation, *Probab. Theory Relat. Fields*, **136**, 298–320.

COLLEGE OF SCIENCE AND TECHNOLOGY, NIHON UNIVERSITY

*Email address:* `kubota.naoki08@nihon-u.ac.jp`

DEPARTMENT OF APPLIED MATHEMATICS, FACULTY OF ENGINEERING, YOKOHAMA NATIONAL UNIVERSITY, YOKOHAMA, JAPAN

*Email address:* `takei-masato-fx@ynu.ac.jp`