# DEPARTMENT OF MATHEMATICAL SCIENCES TECHNICAL REPORT SERIES

The Generalized Law of Total Covariance

by

Charles W. Champ and Andrew V. Sills Department of Mathematical Sciences Georgia Southern University, Statesboro, GA 30460-8093

> Number 2022-001 Submitted: May 31, 2022 © 2022

#### 1 Introduction

The covariance  $Cov(Y_1, Y_2)$  between the random measurements  $Y_1$  and  $Y_2$  to be taken on each individual in the population is the average value of the random variable  $[Y_1 - E(Y_1)][Y_2 - E(Y_2)]$  over all individuals in the population. That is,

$$Cov (Y_1, Y_2) = Cov_{Y_1, Y_2} (Y_1, Y_2) = E_{Y_1, Y_2} ([Y_1 - E(Y_1)] [Y_2 - E(Y_2)])$$
$$= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} [y_1 - E(Y_1)] [y_2 - E(Y_2)] f_{Y_1, Y_2} (y_1, y_2) dy_1 dy_2$$

if both  $Y_1$  and  $Y_2$  are continuous random variables. The subscript " $Y_1, Y_2$ " is a more explicit way of indicating that the covariance is a parameter of the joint distribution of  $Y_1$  and  $Y_2$ . If  $Y_1$  and/or  $Y_2$  is a discrete random variable, integration is replaced with summation. For convenience, we let  $\mathbf{Y} = [Y_1, Y_2]^{\mathbf{T}}$ . We will also express the  $Cov(Y_1, Y_2)$  by  $Cov(\mathbf{Y})$  and  $Cov_{\mathbf{Y}}(\mathbf{Y})$ . It is not difficult to show that

$$Cov_{\mathbf{Y}}(\mathbf{Y}) = E_{\mathbf{Y}}(Y_1Y_2) - E_{\mathbf{Y}}(Y_1)E_{\mathbf{Y}}(Y_2).$$

Supposing that the random vector  $\mathbf{Y}$  and the random variable  $X_1$  have a joint distribution, the Total Law of Covariance states that

$$Cov_{\mathbf{Y}}(\mathbf{Y}) = E_{X_1}\left[Cov_{\mathbf{Y}|X_1}(Y_1, Y_2)\right] + Cov_{X_1}\left(E_{\mathbf{Y}|X_1}(Y_1), E_{\mathbf{Y}|X_1}(Y_2)\right).$$

The total law of covariance is also referred to as covariance decomposition formula or conditional covariance formula.

For the case in which  $Y_1 = Y_2 = Y$ , then Cov(Y, Y) = V(Y). For this case it follows that

$$V(Y) = E_{X_1} \left[ V_{Y|X_1} (Y_1, Y_2) \right] + V_{X_1} \left[ E_{Y|X_1} (Y) \right].$$
(1)

This special case is known as the total law of variance. The law of total variance is also referred to as the *variance decomposition formula*, the *conditional variance formulas*, or *Law of Iterated Variances* – known as *Eve's Law* (see Blitzstein and Hwang (2014)). Bain and Engelhardt (1992), Ross (2010), Wackerly et al. (2008). The preceding references provide proofs of (1). Applications of (1) in actuarial science can be found in Mahler and Dean (2001).

It is our interest to examine the unconditional mean  $E_{\mathbf{Y}}[(\mathbf{Y})]$  and the unconditional variance  $Cov_{\mathbf{Y}}(\mathbf{Y})$ of the distribution of  $\mathbf{Y}$  with respect to the joint distribution of  $\mathbf{Y}$  and the k random variables  $X_1, \ldots, X_k$ . In the next section, we will present results obtained by Champ and Sills (2019) for the case in which  $Y_1 = Y_2 = Y$ . They refer to their results as the generalized law of total variance. In Section 3, we present a generalization of the law of total covariance.

Key Words: Law of total variance, conditional variance and covariance, conditional expectation, Eve's law, variance decomposition, covariance decomposition

#### 2 The Generalized Law of Total Expectation and Variance

Champ and Sills (2019) present and prove the following theorems which we include for completeness in this artile. The first theorem is about expectation. This is a generalization of the what is known as the law of total expectation, the law of iterated expectations, the tower rule, Adam's law, and the smoothing theorem. Champ and Sills (2019) refer to their results as the generalized law of total expection.

**Theorem 1.** Assume that Y and the k random variables  $X_1, \ldots, X_k$  have a joint distribution. Then

$$E(Y) = E_{X_1} \left[ E_{X_2 | \mathbf{X}_1} \cdots \left[ E_{X_k | \mathbf{X}_{k-1}} \left[ E_{Y | \mathbf{X}_k} \left( Y \right) \right] \right] \cdots \right]$$

where  $\mathbf{X}_{i} = [X_{1}, ..., X_{i}]^{T}$  for i = 1, ..., k.

Champ and Sills (2019) refer to the next theorem as the Generalized Law of Total Variance.

**Theorem 2.** Assume that Y and the k random variables  $X_1, \ldots, X_k$  have a joint distribution. Then

$$V(Y) = \sum_{i=1}^{k+1} Q_{X_1}^{\langle i,1\rangle} \left[ Q_{X_2|X_1}^{\langle i,2\rangle} \left[ \cdots \left[ Q_{Y|X_1,\dots,X_k}^{\langle i,k+1\rangle} \left(Y\right) \right] \cdots \right] \right],$$
(2)

where the operator  $Q^{\langle i,j \rangle}$  is the variance operator V if i = j and the expectation operator E if  $i \neq j$ , for i, j = 1, ..., k + 1.

For k = 1, we have

$$V(Y) = E_{X_1} \left[ V_{Y|\mathbf{X}_1} (Y) \right] + V_{X_1} \left[ E_{Y|\mathbf{X}_1} (Y) \right]$$

This is the law of total variance. For the case in which k = 2, the variance of Y can be expressed as

$$V(Y) = E_{X_1} \left[ E_{X_2 | \mathbf{X}_1} \left[ V_{Y | \mathbf{X}_2} (Y) \right] \right] + E_{X_1} \left[ V_{X_2 | \mathbf{X}_1} \left[ E_{Y | \mathbf{X}_2} (Y) \right] \right] + V_{X_1} \left[ E_{X_2 | \mathbf{X}_1} \left[ E_{Y | \mathbf{X}_2} (Y) \right] \right].$$

An equivalent result is given in Geweke and Amisano (2014). Klugman (Panjer) provide an example applying the law of total variance. Also, Maheshwari (2020) provides an example of that applies the law of total variance.

# 3 The Generalized Law of Total Covariance

**Theorem 3.** If  $p \times 1$  random vector  $\mathbf{Y}$  and the k random variables  $X_1, \ldots, X_k$  have a joint distribution, then the expection of  $u(\mathbf{Y})$ , a real valued function of  $\mathbf{Y}$ , can be expressed as

$$E\left[u\left(\mathbf{Y}\right)\right] = E_{X_1}\left[E_{X_2|\mathbf{X}_1}\cdots\left[E_{X_k|\mathbf{X}_{k-1}}\left[E_{\mathbf{Y}|\mathbf{X}_k}\left[u\left(\mathbf{Y}\right)\right]\right]\right]\cdots\right],$$

where  $\mathbf{X}_{i} = [X_{1}, \dots, X_{i}]^{\mathbf{T}}$  for  $i = 1, \dots, k$ .

Proof: We see for k = 1 that

$$E\left[u\left(\mathbf{Y}\right)\right] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} u\left(\mathbf{y}\right) f_{\mathbf{Y}}\left(\mathbf{y}\right) d\mathbf{y} = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} u\left(\mathbf{y}\right) \left[\int_{-\infty}^{\infty} f_{\mathbf{Y}}\left(\mathbf{y}, x_{1}\right) dx_{1}\right] d\mathbf{y}$$
$$= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} u\left(\mathbf{y}\right) \left[\int_{-\infty}^{\infty} f_{\mathbf{Y}|X_{1}}\left(\mathbf{y}|x_{1}\right) f_{X_{1}}\left(x_{1}\right) dx_{1}\right] d\mathbf{y}$$
$$= \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} u\left(\mathbf{y}\right) f_{\mathbf{Y}|X_{1}}\left(\mathbf{y}|x_{1}\right) d\mathbf{y}\right] f_{X_{1}}\left(x_{1}\right) dx_{1}$$
$$= \int_{-\infty}^{\infty} E_{\mathbf{Y}|x_{1}}\left[u\left(\mathbf{Y}\right)\right] f_{X_{1}}\left(x_{1}\right) dx_{1}$$
$$= E_{X_{1}}\left[E_{\mathbf{Y}|\mathbf{X}_{1}}\left[u\left(\mathbf{Y}\right)\right]\right].$$

Thus, the theorem is true for k = 1.

Now assume the theorem is true for k > 1. Further, assume that **Y** and the k + 1 random variables  $X_1, \ldots, X_{k+1}$  have a joint distribution. Observe that

$$\begin{split} E_{\mathbf{Y}|\mathbf{X}_{k}}\left[u\left(\mathbf{Y}\right)\right] &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} u\left(\mathbf{y}\right) f_{\mathbf{Y}|\mathbf{X}_{k}}\left(\mathbf{y}\left|\mathbf{x}_{k}\right) d\mathbf{y} = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} u\left(\mathbf{y}\right) \left[\frac{f_{\mathbf{Y},\mathbf{X}_{k}}\left(\mathbf{y},\mathbf{x}_{k}\right)}{f_{\mathbf{X}_{k}}\left(\mathbf{x}_{k}\right)}\right] d\mathbf{y} \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} u\left(\mathbf{y}\right) \left[\frac{\int_{-\infty}^{\infty} f_{\mathbf{Y},\mathbf{X}_{k},X_{k+1}}\left(\mathbf{y}\left|\mathbf{x}_{k},x_{k+1}\right)f_{X_{k+1}|\mathbf{X}_{k}}\left(x_{k+1}\left|\mathbf{x}_{k}\right)f_{\mathbf{X}_{k}}\left(\mathbf{x}_{k}\right)dx_{k+1}\right]}{f_{\mathbf{X}_{k}}\left(\mathbf{x}_{k}\right)}\right] d\mathbf{y} \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} u\left(\mathbf{y}\right) \left[\frac{\int_{-\infty}^{\infty} f_{\mathbf{Y}|\mathbf{X}_{k},X_{k+1}}\left(\mathbf{y}\left|\mathbf{x}_{k},x_{k+1}\right)f_{X_{k+1}|\mathbf{X}_{k}}\left(x_{k+1}\left|\mathbf{x}_{k}\right)dx_{k+1}\right]}{f_{\mathbf{X}_{k}}\left(\mathbf{x}_{k}\right)}\right] d\mathbf{y} \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} u\left(\mathbf{y}\right) \left[\int_{-\infty}^{\infty} f_{\mathbf{Y}|\mathbf{X}_{k},X_{k+1}}\left(\mathbf{y}\left|\mathbf{x}_{k},x_{k+1}\right)f_{X_{k+1}|\mathbf{X}_{k}}\left(x_{k+1}\left|\mathbf{x}_{k}\right)dx_{k+1}\right]d\mathbf{y} \right] \\ &= \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} u\left(\mathbf{y}\right)f_{\mathbf{Y}|\mathbf{X}_{k},X_{k+1}}\left(\mathbf{y}\left|\mathbf{x}_{k},x_{k+1}\right)d\mathbf{y}\right]f_{X_{k+1}|\mathbf{X}_{k}}\left(x_{k+1}\left|\mathbf{x}_{k}\right)dx_{k+1}\right] \\ &= \int_{-\infty}^{\infty} E_{\mathbf{Y}|\mathbf{X}_{k},x_{k+1}}\left[u\left(\mathbf{Y}\right)\right]f_{X_{k+1}|\mathbf{X}_{k}}\left(x_{k+1}\left|\mathbf{x}_{k}\right)dx_{k+1} \\ &= E_{X_{k+1}|\mathbf{X}_{k}}\left[E_{\mathbf{Y}|\mathbf{X}_{k},X_{k+1}}\left[u\left(\mathbf{Y}\right)\right]\right] \\ &= E_{X_{k+1}|\mathbf{X}_{k}}\left[E_{\mathbf{Y}|\mathbf{X}_{k+1}}\left[u\left(\mathbf{Y}\right)\right]\right]. \end{split}$$

Substituting  $E_{\mathbf{Y}|\mathbf{X}_{k}}\left[u\left(\mathbf{Y}\right)\right]$  with  $E_{X_{k+1}|\mathbf{X}_{k}}\left[E_{\mathbf{Y}|\mathbf{X}_{k+1}}\left[u\left(\mathbf{Y}\right)\right]\right]$  in the theorem results in

$$E\left[u\left(\mathbf{Y}\right)\right] = E_{X_1}\left[E_{X_2|\mathbf{X}_1}\cdots\left[E_{X_k|\mathbf{X}_{k-1}}\left[E_{X_{k+1}|\mathbf{X}_k}\left[E_{\mathbf{Y}|\mathbf{X}_{k+1}}\left[u\left(\mathbf{Y}\right)\right]\right]\right]\right]\cdots\right].$$

Hence, the theorem is true for k + 1. Therefore, by the Axiom of Induction, the theorem is true for all positive integers k.

**Theorem 4.** (Generalized Law of Total Covariance) Assume that  $\mathbf{Y} = [Y_1, Y_2]^{\mathbf{T}}$  and the k random variables  $X_1, \ldots, X_k$  have a joint distribution. Then

$$Cov (\mathbf{Y}) = Cov (Y_1, Y_2)$$

$$= Cov_{X_1} \left( E_{X_2 | \mathbf{X}_1} \left[ \cdots \left[ E_{\mathbf{Y} | \mathbf{X}_k} (Y_1) \right] \cdots \right], E_{X_2 | \mathbf{X}_1} \left[ \cdots \left[ E_{\mathbf{Y} | \mathbf{X}_k} (Y_2) \right] \cdots \right] \right)$$

$$+ E_{X_1} \left[ Cov_{X_2 | \mathbf{X}_1} \left( E_{X_3 | \mathbf{X}_2} \left[ \cdots \left[ E_{\mathbf{Y} | \mathbf{X}_k} (Y_1) \right] \cdots \right], E_{X_3 | \mathbf{X}_2} \left[ \cdots \left[ E_{\mathbf{Y} | \mathbf{X}_k} (Y_2) \right] \cdots \right] \right) \right]$$

$$\vdots$$

$$+ E_{X_1} \left[ E_{X_2 | \mathbf{X}_1} \left[ E_{X_3 | \mathbf{X}_2} \left[ \cdots \left[ Cov_{\mathbf{Y} | \mathbf{X}_k} (Y_1, Y_2) \right] \cdots \right] \right] \right],$$

where  $\mathbf{X}_{i} = [X_{1}, \dots, X_{i}]^{\mathbf{T}}$  for  $i = 1, \dots, k$ .

Proof: As previously stated,

$$Cov_{\mathbf{Y}}(\mathbf{Y}) = E_{\mathbf{Y}}(Y_1Y_2) - E_{\mathbf{Y}}(Y_1)E_{\mathbf{Y}}(Y_2).$$

For k = 1, using theorem 3 with  $u(\mathbf{Y}) = Y_1 Y_2$ , we can write

$$E_{\mathbf{Y}}(Y_1Y_2) = E_{X_1}\left[\left[E_{\mathbf{Y}|X_1}(Y_1Y_2)\right]\right]$$

and with  $u(\mathbf{Y}) = Y_i$  for i = 1, 2,

$$E_{\mathbf{Y}}(Y_1) E_{\mathbf{Y}}(Y_2) = E_{X_1} \left[ \left[ E_{\mathbf{Y}|X_1}(Y_1) \right] \right] E_{X_1} \left[ \left[ E_{\mathbf{Y}|X_1}(Y_2) \right] \right].$$

Thus,

$$Cov_{\mathbf{Y}}(\mathbf{Y}) = E_{X_{1}}\left[\left[E_{\mathbf{Y}|X_{1}}(Y_{1}Y_{2})\right]\right] - E_{X_{1}}\left[\left[E_{\mathbf{Y}|X_{1}}(Y_{1})\right]\right] E_{X_{1}}\left[\left[E_{\mathbf{Y}|X_{1}}(Y_{2})\right]\right]$$
$$= E_{X_{1}}\left[\left[E_{\mathbf{Y}|X_{1}}(Y_{1}Y_{2})\right] - \left[E_{\mathbf{Y}|X_{1}}(Y_{1})\right]\left[E_{\mathbf{Y}|X_{1}}(Y_{2})\right]\right]$$
$$+ E_{X_{1}}\left[\left[E_{\mathbf{Y}|X_{1}}(Y_{1})\right]\left[E_{\mathbf{Y}|X_{1}}(Y_{2})\right]\right] - E_{X_{1}}\left[\left[E_{\mathbf{Y}|X_{1}}(Y_{1})\right]\right] E_{X_{1}}\left[\left[E_{\mathbf{Y}|X_{1}}(Y_{2})\right]\right]$$
$$= E_{X_{1}}\left[Cov\left(E_{\mathbf{Y}|X_{1}}(Y_{1}), E_{\mathbf{Y}|X_{1}}(Y_{2})\right)\right] + Cov_{X_{1}}\left(E_{\mathbf{Y}|X_{1}}(Y_{1}), E_{\mathbf{Y}|X_{1}}(Y_{2})\right).$$

Hence, the theorem is true forrem is true for k > 1. Further, assume that **Y** and the k + 1 random variables  $X_1, \ldots, X_k, X_{k+1}$  have a joint distribution. Observe that

$$E_{\mathbf{Y}|\mathbf{X}_{k}}\left(Y_{i}\right) = E_{X_{k+1}|\mathbf{X}_{k}}\left[E_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(Y_{i}\right)\right]$$

for i = 1, 2 which follows from the proof of theorem 3. Again using the results of theorem 3, we have that

$$\begin{aligned} Cov_{\mathbf{Y}|\mathbf{X}_{k}} (Y_{1}, Y_{2}) &= E_{\mathbf{Y}|\mathbf{X}_{k}} (Y_{1}Y_{2}) - E_{\mathbf{Y}|\mathbf{X}_{k}} (Y_{1}) E_{\mathbf{Y}|\mathbf{X}_{k}} (Y_{2}) \\ &= E_{X_{k+1}|\mathbf{X}_{k}} \left[ E_{\mathbf{Y}|\mathbf{X}_{k+1}} (Y_{1}Y_{2}) - E_{\mathbf{Y}|\mathbf{X}_{k+1}} (Y_{1}) E_{\mathbf{Y}|\mathbf{X}_{k+1}} (Y_{2}) \right] \\ &+ E_{X_{k+1}|\mathbf{X}_{k}} \left[ E_{\mathbf{Y}|\mathbf{X}_{k+1}} (Y_{1}) E_{\mathbf{Y}|\mathbf{X}_{k+1}} (Y_{2}) \right] \\ &- E_{X_{k+1}|\mathbf{X}_{k}} \left[ E_{\mathbf{Y}|\mathbf{X}_{k+1}} (Y_{1}) \right] E_{X_{k+1}|\mathbf{X}_{k}} \left[ E_{\mathbf{Y}|\mathbf{X}_{k+1}} (Y_{2}) \right] \\ &= E_{X_{k+1}|\mathbf{X}_{k}} \left[ Cov_{\mathbf{Y}|\mathbf{X}_{k+1}} (Y_{1}, Y_{2}) \right] \\ &+ E_{X_{k+1}|\mathbf{X}_{k}} \left[ Cov_{\mathbf{Y}|\mathbf{X}_{k+1}} \left( E_{\mathbf{Y}|\mathbf{X}_{k+1}} (Y_{1}), E_{\mathbf{Y}|\mathbf{X}_{k+1}} (Y_{2}) \right) \right] \\ &= E_{X_{k+1}|\mathbf{X}_{k}} \left[ Cov_{\mathbf{Y}|\mathbf{X}_{k+1}} \left( E_{\mathbf{Y}|\mathbf{X}_{k+1}} (Y_{1}), E_{\mathbf{Y}|\mathbf{X}_{k+1}} (Y_{2}) \right) \right] \\ &+ E_{X_{k+1}|\mathbf{X}_{k}} \left[ Cov_{\mathbf{Y}|\mathbf{X}_{k+1}} \left( E_{\mathbf{Y}|\mathbf{X}_{k+1}} (Y_{1}), E_{\mathbf{Y}|\mathbf{X}_{k+1}} (Y_{2}) \right) \right] \\ &+ E_{X_{k+1}|\mathbf{X}_{k}} \left[ Cov_{\mathbf{Y}|\mathbf{X}_{k+1}} \left( Y_{1}, Y_{2} \right) \right]. \end{aligned}$$

Substituting  $E_{\mathbf{Y}|\mathbf{X}_{k}}(Y_{i})$  with  $E_{X_{k+1}|\mathbf{X}_{k}}[E_{\mathbf{Y}|\mathbf{X}_{k+1}}(Y_{i})]$  for i = 1, 2 and  $Cov_{\mathbf{Y}|\mathbf{X}_{k}}(Y_{1}, Y_{2})$  with

 $E_{X_{k+1}|\mathbf{X}_{k}}\left[Cov_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(E_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(Y_{1}\right),E_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(Y_{2}\right)\right)\right]+E_{X_{k+1}|\mathbf{X}_{k}}\left[Cov_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(Y_{1},Y_{2}\right)\right]$ 

yields

$$\begin{split} &Cov\left(\mathbf{Y}\right) \\ &= Cov_{X_{1}}\left(E_{X_{2}|\mathbf{X}_{1}}\left[\cdots\left[E_{X_{k+1}|\mathbf{X}_{k}}\left[E_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(Y_{1}\right)\right]\right]\cdots\right], E_{X_{2}|\mathbf{X}_{1}}\left[\cdots\left[E_{X_{k+1}|\mathbf{X}_{k}}\left[E_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(Y_{2}\right)\right]\right]\cdots\right]\right) \\ &+ E_{X_{1}}\left[Cov_{X_{2}|\mathbf{X}_{1}}\left(\frac{E_{X_{3}|\mathbf{X}_{2}}\left[\cdots\left[E_{X_{k+1}|\mathbf{X}_{k}}\left[E_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(Y_{1}\right)\right]\right]\cdots\right]\right)\right] \\ &\vdots \\ &+ E_{X_{1}}\left[E_{X_{2}|\mathbf{X}_{1}}\left[E_{X_{3}|\mathbf{X}_{2}}\left[\cdots\left[\frac{E_{X_{k+1}|\mathbf{X}_{k}}\left[Cov_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(\frac{E_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(Y_{1}\right)\right)}{E_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(Y_{2}\right)}\right)\right]\right]\cdots\right]\right]\right] \\ &= Cov_{X_{1}}\left(\frac{E_{X_{2}|\mathbf{X}_{1}}\left[\cdots\left[E_{X_{k+1}|\mathbf{X}_{k}}\left[E_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(Y_{1}\right)\right]\right]\cdots\right]}{E_{X_{3}|\mathbf{X}_{2}}\left[\cdots\left[E_{X_{k+1}|\mathbf{X}_{k}}\left[E_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(Y_{1}\right)\right]\right]\cdots\right]\right) \\ &+ E_{X_{1}}\left[Cov_{X_{2}|\mathbf{X}_{1}}\left(\frac{E_{X_{3}|\mathbf{X}_{2}}\left[\cdots\left[E_{X_{k+1}|\mathbf{X}_{k}}\left[E_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(Y_{1}\right)\right]\right]\cdots\right]\right)\right] \\ &\vdots \\ &+ E_{X_{1}}\left[E_{X_{3}|\mathbf{X}_{1}}\left(\frac{E_{X_{3}|\mathbf{X}_{2}}\left[\cdots\left[E_{X_{k+1}|\mathbf{X}_{k}}\left[E_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(Y_{1}\right)\right]\right]\cdots\right]\right)\right] \\ &\vdots \\ &+ E_{X_{1}}\left[E_{X_{3}|\mathbf{X}_{1}}\left[E_{X_{3}|\mathbf{X}_{2}}\left[\cdots\left[E_{X_{k+1}|\mathbf{X}_{k}}\left[Cov_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(\frac{E_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(Y_{1}\right)\right)\right]\cdots\right]\right)\right] \\ &\vdots \\ &+ E_{X_{1}}\left[E_{X_{2}|\mathbf{X}_{1}}\left[E_{X_{3}|\mathbf{X}_{2}}\left[\cdots\left[E_{X_{k+1}|\mathbf{X}_{k}}\left[Cov_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(\frac{E_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(Y_{1}\right)\right)\right]\right]\cdots\right]\right]\right] \\ &\vdots \\ &+ E_{X_{1}}\left[E_{X_{2}|\mathbf{X}_{1}}\left[E_{X_{3}|\mathbf{X}_{2}}\left[\cdots\left[E_{X_{k+1}|\mathbf{X}_{k}}\left[Cov_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(\frac{E_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(Y_{1}\right)\right)\right]\cdots\right]\right]\right] \\ &\vdots \\ &+ E_{X_{1}}\left[E_{X_{2}|\mathbf{X}_{1}}\left[E_{X_{3}|\mathbf{X}_{2}}\left[\cdots\left[E_{X_{k+1}|\mathbf{X}_{k}}\left[Cov_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(\frac{E_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(Y_{1}\right)\right)\right]\cdots\right]\right]\right] \\ &\vdots \\ &+ E_{X_{1}}\left[E_{X_{2}|\mathbf{X}_{1}}\left[E_{X_{3}|\mathbf{X}_{2}}\left[\cdots\left[E_{X_{k+1}|\mathbf{X}_{k}}\left[Cov_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(Y_{1}\right)\left(Y_{1}\right)\right]\cdots\right]\right]\right] \\ &\vdots \\ &+ E_{X_{1}}\left[E_{X_{2}|\mathbf{X}_{1}}\left[E_{X_{3}|\mathbf{X}_{2}}\left[\cdots\left[E_{X_{k+1}|\mathbf{X}_{k}}\left[Cov_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(Y_{1}\right)\left(Y_{1}\right)\right]\right]\cdots\right]\right]\right] \\ &\vdots \\ &= E_{X_{1}}\left[E_{X_{2}|\mathbf{X}_{1}}\left[E_{X_{2}|\mathbf{X}_{2}}\left[\cdots\left[E_{X_{k+1}|\mathbf{X}_{k}}\left[E_{\mathbf{Y}|\mathbf{X}_{k+1}}\left(Y_{1}\right)\right]\right]\cdots\right]\right] \\ &= E_{X_{1}}\left[E_{X_{2}|\mathbf{X}_{1}}\left[E_{X_{2}|\mathbf{X}_{2}}\left[\cdots\left[E_{X_{k+1}|\mathbf{X}_{k}}\left[E_{\mathbf{Y}|\mathbf{X}$$

Hence, the theorem is true for k + 1. Therefore, by the Axiom of Induction, the theorem is true for all positive integers k.

Klugman, Panjer, and Willmot (2019) and Maheshwari (2020) provide examples applying the law of total covariance.

Theorem 5. The generalized law of total variance is a special case of the generalized law of total covariance.

Proof: Suppose that  $Y_1 = Y_2 = Y$ . We have

$$\begin{split} V(Y) &= Cov\left(Y,Y\right) \\ &= Cov_{X_{1}}\left(E_{X_{2}|\mathbf{X}_{1}}\left[\cdots\left[E_{\mathbf{Y}|\mathbf{X}_{k}}\left(Y\right)\right]\cdots\right], E_{X_{2}|\mathbf{X}_{1}}\left[\cdots\left[E_{\mathbf{Y}|\mathbf{X}_{k}}\left(Y\right)\right]\cdots\right]\right) \\ &+ E_{X_{1}}\left[Cov_{X_{2}|\mathbf{X}_{1}}\left(E_{X_{3}|\mathbf{X}_{2}}\left[\cdots\left[E_{\mathbf{Y}|\mathbf{X}_{k}}\left(Y\right)\right]\cdots\right], E_{X_{3}|\mathbf{X}_{2}}\left[\cdots\left[E_{\mathbf{Y}|\mathbf{X}_{k}}\left(Y\right)\right]\cdots\right]\right)\right] \\ &\vdots \\ &+ E_{X_{1}}\left[E_{X_{2}|\mathbf{X}_{1}}\left[E_{X_{3}|\mathbf{X}_{2}}\left[\cdots\left[Cov_{\mathbf{Y}|\mathbf{X}_{k}}\left(Y,Y\right)\right]\cdots\right]\right]\right] \\ &= V_{X_{1}}\left(E_{X_{2}|\mathbf{X}_{1}}\left[\cdots\left[E_{\mathbf{Y}|\mathbf{X}_{k}}\left(Y\right)\right]\cdots\right], E_{X_{2}|\mathbf{X}_{1}}\left[\cdots\left[E_{\mathbf{Y}|\mathbf{X}_{k}}\left(Y\right)\right]\cdots\right]\right) \\ &+ E_{X_{1}}\left[V_{X_{2}|\mathbf{X}_{1}}\left(E_{X_{3}|\mathbf{X}_{2}}\left[\cdots\left[E_{\mathbf{Y}|\mathbf{X}_{k}}\left(Y\right)\right]\cdots\right], E_{X_{3}|\mathbf{X}_{2}}\left[\cdots\left[E_{\mathbf{Y}|\mathbf{X}_{k}}\left(Y\right)\right]\cdots\right]\right)\right] \\ &\vdots \\ &+ E_{X_{1}}\left[E_{X_{2}|\mathbf{X}_{1}}\left[E_{X_{3}|\mathbf{X}_{2}}\left[\cdots\left[V_{\mathbf{Y}|\mathbf{X}_{k}}\left(Y,Y\right)\right]\cdots\right]\right]\right]. \end{split}$$

This is the generalized law of total variance.■

### 4 Conclusion

A generalization of the law of total covariance was given. It was proven that the generalized law of total variance is a special case of the generalized law of the total covariance. Examples were referenced.

# References

- Anderson, T.W. (2003), An Introduction to Multivariate Statistical Analysis, Third Edition, New York: John Wiley and Sons, Inc.
- Bain, L.J. and Engelhardt, M. (1992), Introduction to Probability and Mathematical Statistics, Boston: PWS-KENT Publishing Company.

Billingsley, P. (1995), Probability and Measure, New York: John Wiley & Sons, Inc.

Blitzstein, J.K. and Hwang, J. (2014), Introduction to Probability, CRC Press.

- Bowsher, C.G. and P.S. Swain, (2012), "Identifying Sources of Variation and the Flow of Information in Biochemical Networks," *Proceeding of the National Academy Sciences USA* 109, 1320-1329.
- Champ, C.W. and Sills, A.V. (2019), "The Generalized Law of Total Variance," Department of Mathematical Sciences Technical Report Series, Georgia Southern University.
- Geweke, J. and Amisano, G. (2014), "Analysis of Variance for Bayesian Inference," *Econometric Reviews* **33**(1-4), 270-288.
- Klugman, S.A., Panjer, H.H., and Willmot, G.E. (2019), Loss Models: From Data to Decisions, 5th Edition, Wiley and Sons, Inc.: New York.
- Maheshwari, S. (2020), "Solving Conditional Probability Problems with the Laws of Total Expectation, Variance, and Covariance," Blog: https://towardsdatascience.com/solving-conditional-probability-problemswith-the-laws-of-total-expectation-variance-and-c38c07cfebfa.
- Mahler, H.C. and Dean, C.G. (2001). "Chapter 8: Credibility," in Casualty Actuarial Society. Foundations of Casualty Actuarial Science, Fourth Edition, Casualty Actuarial Society.
- Ross, S.M. (2010), A First Course in Probability, Eighth Edition, Upper Saddle River, NJ: Prentice Hall.
- Wackerly, D.D., Mendenhall III, W., and Scheaffer, R.L. (2008), Mathematical Statistics with Applications, Seventh Edition, Belmont, CA: Duxbury, Thomson Higher Education.
- Weiss, N.A. (2005), A Course in Probability, Addison-Wesley.