# Symmetries and conservation laws for the generalized n-dimensional Ermakov system

Andronikos Paliathanasis<sup>\*†</sup>

Institute of Systems Science, Durban University of Technology PO Box 1334, Durban 4000, Republic of South Africa Instituto de Ciencias Físicas y Matemáticas, Universidad Austral de Chile, Valdivia, Chile,

Genly Leon

Departamento de Matemáticas, Universidad Católica del Norte, Avda. Angamos 0610, Casilla 1280 Antofagasta, Chile Institute of Systems Science, Durban University of Technology PO Box 1334, Durban 4000, Republic of South Africa

P.G.L. Leach

Institute of Systems Science, Durban University of Technology PO Box 1334, Durban 4000, Republic of South Africa

May 31, 2022

#### Abstract

We revise recent results on the classification of the generalized three-dimensional Hamiltonian Ermakov system. We show that a statement published recently is incorrect, while the solution for the classification problem was incomplete. We present the correct classification for the three-dimensional system by using results which related the background space with the dynamics. Finally, we extend our results for the generalized *n*-dimensional Hamiltonian Ermakov system.

<sup>\*</sup>Email: anpaliat@phys.uoa.gr

<sup>&</sup>lt;sup>†</sup>Coresponding Author

Keywords: Lie symmetries; Noether symmetries; Ermakov system; Integrability

## 1 Introduction

At the end of the 19th century, Ermakov [1] solved the problem for the derivation of first integral for the time-dependent linear equation

$$\ddot{x} + \omega^2(t)x = 0,\tag{1}$$

where an overdot means total derivative with respect to the time-variable, t, that is,  $\dot{x} = \frac{dx}{dt}$ .

By introducing the new variable  $\rho(t)$  which satisfies the second-order differential equation

$$\ddot{\rho} + \omega^2(t)\rho = \rho^{-3} \tag{2}$$

the time-dependent term in (1) can be eliminated and the following function  $J(\rho, \dot{\rho}, x, \dot{x})$  can be constructed

$$J(\rho, \dot{\rho}, x, \dot{x}) = \frac{1}{2} \left[ (\rho \dot{x} - \dot{\rho} x)^2 + (x/\rho)^2 \right],$$
(3)

which is a conservation law, i.e.  $\dot{J} = 0$ . The second-order ordinary differential equation (2) was solved by Pinney in [2]. The analytic solution is

$$\rho^{2}(t) = c_{1}(\rho_{1}(t))^{2} + c_{2}(\rho_{2}(t))^{2} + 2c_{3}\rho_{1}(t)\rho_{2}(t), \qquad (4)$$

where  $\rho_1(t)$  and  $\rho_2(t)$  are solutions of equation (1) and  $c_1$ ,  $c_2$  and  $c_3$  are constants, of which only two are independent. Hence, equation (2) is known as the Ermakov-Pinney equation. Furthermore, in [3], Lewis investigated the integral properties for the time-dependent harmonic oscillator, of equation (1) in classical and quantum systems. Lewis rediscovered the the Ermakov invariant (3) and function  $J(\rho, \dot{\rho}, x, \dot{x})$  nowadays is called Ermakov-Lewis invariant. The oscillator appears in many physical problems and other areas of Applied Mathematics. Consequently, the Ermakov-Lewis invariant and the Ermakov-Pinney equation have appeared in many studies, from quantum mechanics [3–5], to gravitational theory [6,7] and others [8–11]. Recently, in [12] the physical interpretation of the Ermakov-Lewis invariant was discussed.

The application of the Lie theory on the Ermakov-Pinney equation provides that equation (2) admits three Lie point symmetries which form the SL(2, R) algebra [13]. The representation of the Lie algebra depends upon the term  $\omega^2(t)$ . Thus, there exists a point transformation in which the term  $\omega^2(t)$  can be eliminated and equation (2) to be reduced in the simple form

$$\ddot{\rho} = \rho^{-3}.\tag{5}$$

There are many extensions and generalization of the Ermakov-Pinney equation in the literature. The twodimensional system,

$$\ddot{x} + \omega^2(t) x = \frac{f\left(\frac{y}{x}\right)}{x^3} \tag{6}$$

$$\ddot{y} + \omega^2(t) y = \frac{g\left(\frac{y}{x}\right)}{y^3},\tag{7}$$

is known as an Ermakov system [14]. System (6), (7) admits the conservation law [15]

$$I(x, \dot{x}, y, \dot{x}) = \frac{1}{2} (x\dot{y} - \dot{x}y)^2 + \int^{y/x} \left( uf(u) - u^{-3}g(u) \right) du$$
(8)

which is a generalization of the Ermakov-Lewis invariant. We remark that there exist a point transformation such that the system (6), (7) can be written in an equivalent form without the  $\omega^2(t)$  term. Moreover, the two-dimensional Ermakov system admits three point symmetries which form the the SL(2, R) algebra. In the special case for which  $f\left(\frac{y}{x}\right) = V(x, y)_{,x}$  and  $g\left(\frac{y}{x}\right) = V(x, y)_{y}$ , the dynamical system follows from a variational principle and the Lie symmetries are also Noether symmetries.

By using the latter property for the Ermakov system, that is, the admitted Lie symmetries for a given dynamical system to form the SL(2, R) algebra, there have been various generalizations of the Ermakov system in higher-dimensions [13], in curved geometries [16] or many others see for instance [18–21] and references therein. The three-dimensional Ermakov system introduced in [13], for which it was found that in the case that the Ermakov system is Hamiltonian the Ermakov-Lewis invariant is a function of the variational conservation laws which are determined by the applications of Noether's second theorem for the elements of the SL(2, R)algebra, see also the discussion in [17]. However, such a property was found also in the case of curved geometries. Moreover, for the Hamiltonian Ermakov system, the Ermakov-Lewis invariant can be constructed by Noether's second theorem for contact symmetries. Indeed, there was found to be an exact relation for the existence of the Killing tensor which provides the Ermakov-Lewis invariant and of the main generator vector for the background space of the SL(2, R) Lie algebra [16]. The SL(2, R) Lie algebra plays an important role in the study of the integrability of various Hamiltonian systems, we refer the reader in some recent studies [22–28]

A complete classification of variational symmetries for the Hamiltonian three-dimensional Ermakov [13] system performed in [29]. The authors determined the functional form for the unknown potential functions for the generalized Ermakov system to admit additional variational symmetries which produce conservation laws except from the elements of the SL(2, R) algebra. The authors determined eight different potential functions of which they claim that four of these functions are new in the literature and new exact solutions were determined. However, this claim is false. As we shall see with a simple construction method the potential functions the authors determined in [29] are well known in the literature. Nevertheless, inspired by the problem which the authors of [29] study, we extend our analysis and we find a general result for the generalized Hamiltonian Ermakov system in n-dimensional Euclidian geometry. The plan of the paper is as follow.

In Section 2 we present basic properties and definitions of the symmetry analysis. In Section 3 we discuss the two-dimensional Ermakov system and the Lewis invariant. The analysis of the three-dimensional generalized Ermakov system is presented in Section 4. Finally, in Section 5 we draw our conclusions.

### 2 Preliminaries

In this Section we present the basic properties and definitions for the theory of symmetries of differential equations. We assume a system of second-order ordinary differential equations of the form

$$\ddot{x}^i = \Omega^i \left( t, x^j, \dot{x}^j \right) \ , \tag{9}$$

in which  $x^i$  are the dependent variables and t is the independent variable. Furthermore, consider the oneparameter point transformation  $\Phi(t, x; \varepsilon)$  with infinitesimal generator  $\xi(t, x^j) \partial_t + \eta^i(t, x^j) \partial_i$  defined in the augmented space  $\{t, x^i\}$ . Thus, we say that the vector field X is a Lie point symmetry for the system (9) if the set of differential equations remains invariant under the action of the one-parameter point transformation  $\Phi(t, x; \varepsilon)$ , equivalently when the following condition is true [30]

$$X^{[2]}\left(\ddot{x}^{i} - \Omega^{i}\left(t, x^{j}, \dot{x}^{j}\right)\right) = 0 , \qquad (10)$$

where  $X^{[2]}$  is the second prolongation of X defined by the formula  $X^{[2]} = X + \left(\dot{\eta}^i - \dot{x}^i \dot{\xi}\right) \partial_{\dot{x}^i} + \left(\ddot{\eta}^i - \dot{x}^i \ddot{\xi} - 2\ddot{x}^i \dot{\xi}\right) \partial_{\ddot{x}^i}$ . An equivalent way to express the symmetry condition (10) is [30]

$$\left[X^{[1]}, A\right] = \lambda\left(x^k\right) A , \qquad (11)$$

in which now  $X^{[1]}$  is the first prolongation of X and A is the Hamiltonian vector field  $A = \partial_t + \dot{x}\partial_x + \Omega^i(t, x^j, \dot{x}^j) \partial_{\dot{x}^i}$ .

If the system of differential equations results from a variational principle, that is, there exists a Lagrangian function  $L = L(t, x^j, \dot{x}^j)$ , then the infinitesimal generator X for the one-parameter point transformation is characterized as variational symmetry or as Noether symmetry for the given Lagrangian function, if there exists a function G such that the condition of Noether's first theorem is satisfied [31]

$$X^{[1]}L + L\frac{d\xi}{dt} = \frac{dG}{dt}$$
(12)

Function G is boundary term introduced to allow for the infinitesimal changes in the value of the Action Integral produced by the infinitesimal change in the boundary of the domain caused by the infinitesimal transformation of the variables in the Action Integral.

The importancy of the determination of Noether symmetries for a given system is that there exist a unique relation between Noether symmetries and conservation laws. Indeed, if X is a Noether symmetry vector with corresponding boundary term G, then according to Noether's second theorem the following function

$$I(t, x^{j}, \dot{x}^{j}) = \xi \left( \dot{x}^{i} \frac{\partial L}{\partial \dot{x}^{i}} - L \right) - \frac{\partial L}{\partial \dot{x}^{i}} \eta^{i} + G$$
(13)

is a conservation law for the dynamical system, that is  $\dot{I}(t, x^j, \dot{x}^j) = 0$ . For a recent review on Noether's theorems we refer the reader to [32].

#### 3 The Ermakov system

In the following we consider the Ermakov system and without loss of generality we can omit the linear term with the time-dependent component  $\omega^2(t)$ .

As we mentioned before the Ermakov-Pinney equation (5) admits three Lie point symmetries, the vector fields

$$X^{1} = \partial_{t} , \ X^{2} = 2t\partial_{t} + \rho\partial_{\rho} , \ X^{3} = t^{2}\partial_{t} + t\rho\partial_{\rho} .$$

$$\tag{14}$$

Equation (5) admits the Lagrangian function

$$L(\rho, \dot{\rho}) = \frac{1}{2} \left( \dot{\rho}^2 - \rho^{-2} \right).$$
(15)

Thus, by replacing the vector fields  $\{X^1, X^2, X^3\}$  in Noether's condition we find that the three vector fields are Noether symmetries with corresponding boundary terms  $G(X^1) = G_0$ ,  $G(X^2) = 0$  and  $G(X^3) = \frac{1}{2}\rho^2$  respectively.

Hence, from expression (13) we define the conservation laws

$$I_1(X^1) = \frac{1}{2}(\dot{\rho}^2 + \rho^{-2}) \equiv \mathcal{H},$$
(16)

$$I_2(X^2) = 2t\mathcal{H} - \rho\dot{\rho} , \qquad (17)$$

$$I_3(X^3) = t^2 \mathcal{H} - t\rho \dot{\rho} + \frac{1}{2}\rho^2 , \qquad (18)$$

where  $\mathcal{H}$  is the Hamiltonian function.

Consider now the two-dimensional Hamiltonian Ermakov system

$$\ddot{x} = \frac{\left(V\left(\frac{y}{x}\right)\right)_x}{x^3}, \ \ddot{y} = \frac{\left(V\left(\frac{y}{x}\right)\right)_y}{y^3},\tag{19}$$

or in polar coordinates  $(x, y) = (r \cos \theta, r \sin \theta),$ 

$$\ddot{\rho} - \rho \dot{\theta}^2 = \frac{V(\theta)}{\rho^3} , \qquad (20)$$

$$\ddot{\theta} + 2\rho \dot{\theta} = \frac{V(\theta)_{,\theta}}{2\rho^4}.$$
(21)

with Lagrangian function

$$L\left(\rho,\dot{\rho},\theta,\dot{\theta}\right) = \frac{1}{2}\left(\dot{\rho}^2 + \rho^2\dot{\theta}^2 - \frac{V\left(\theta\right)}{\rho^2}\right).$$
(22)

For arbitrary potential function  $V(\theta)$  the dynamical system (20), (21) admits as Lie point symmetries the vector fields  $\{X^1, X^2, X^3\}$  which are also Noether symmetries for the Lagrangian function (22). The corresponding conservation laws are the  $I_1(X^1)$ ,  $I_2(X^2)$  and  $I_3(X^3)$  where now

$$\mathcal{H} = \frac{1}{2} \left( \dot{\rho}^2 + \rho^2 \dot{\theta}^2 + \frac{V\left(\theta\right)}{\rho^2} \right).$$
(23)

By using these three functions we can define the following expression [13]

$$J(\rho,\dot{\rho}) = 4I_3(X^1)\mathcal{H} - (I_2(X^2))^2$$
(24)

which is nothing else than the Ermakov-Lewis invariant, that is

$$J(\rho, \dot{\rho}) = \frac{1}{2} \left( \dot{\rho}^2 + \rho^{-2} \right).$$
(25)

However, for specific forms of  $V(\theta)$  additional Noether symmetries can exist. Indeed when  $V(\theta) = V_0$ , equation (21) reads  $(\mathcal{L})^{\cdot} = 0$ , where  $\mathcal{L} = \rho^2 \dot{\theta}$  is the angular momentum. The additional Noether symmetry in

this case is the vector field  $X^4 = \partial_{\phi}$ . The functional forms of  $V(\theta)$  for which additional variational symmetries exist are presented in [33]. We omit the presentation of the symmetry vector and we present the potential functions.

The potentials are  $V_A(\theta) = V_0$ ,  $V_B(\theta) = V_0(\cos\theta)^{-2}$ ,  $V_C(\theta) = V_0(\sin\theta)^{-2}$ ,  $V_D(\theta) = V_1(\cos\theta)^{-2} + V_2(\sin\theta)^{-2}$  and  $V_E(\theta) = V_0(V_1\sin\theta - \cos\theta)^{-2}$ . We remark that potentials  $V_B(\theta)$  and  $V_C(\theta)$  are the same potential function, because  $\sin\theta = \cos\left(\theta - \frac{\pi}{2}\right)$ . Thus there are only four potentials. As we discussed, potential  $V_A(\theta)$  provides the conservation law of angular momentum.

For potential  $V_B(\theta)$ , if we go into Cartesian coordinates the Lagrangian function for the Ermakov system (19) is

$$L_B(x, \dot{x}, y, \dot{y}) = \frac{1}{2} \left( \dot{x}^2 + \dot{y}^2 - \frac{V_0}{x^2} \right) , \qquad (26)$$

which provides the second-order differential equations  $\ddot{x} = \frac{V_0}{x^3}$ ,  $\ddot{y} = 0$ . The time-independent conservation laws are  $J(x, \dot{x}) = \frac{1}{2} (\dot{x}^2 + V_0 x^{-2})$  and  $I^2 = \dot{y}^2$ . The potential  $V_C(\theta)$  gives the same conservation laws with  $(x, y) \to (y, x)$ .

For potential  $V_D(\theta)$  in Cartesian coordinates the Lagrangian function becomes

$$L_D(x, \dot{x}, y, \dot{y}) = \frac{1}{2} \left( \dot{x}^2 + \dot{y}^2 - \frac{V_1}{x^2} - \frac{V_2}{y^2} \right) , \qquad (27)$$

with equations of motion  $\ddot{x} = \frac{V_1}{x^3}$  and  $\ddot{y} = \frac{2}{y^3}$ , for which the conservation laws are  $J_1(x, \dot{x}) = \frac{1}{2} \left( \dot{x}^2 + V_1 x^{-2} \right)$  and  $J_2(y, \dot{y}) = \frac{1}{2} \left( \dot{y}^2 + V_1 y^{-2} \right)$ .

Finally, for the potential function  $V_{E}(\theta)$  we have the Lagrangian

$$L_E(x, \dot{x}, y, \dot{y}) = \frac{1}{2} \left( \dot{x}^2 + \dot{y}^2 - \frac{V_0}{(\alpha y - x)^2} \right).$$
(28)

By doing the second change of variables  $x = X + \alpha y$  it follows that  $L_E\left(X, \dot{X}, y, \dot{y}\right) = \frac{1}{2}\left(\dot{X}^2 + 2\alpha\dot{X}\dot{Y} + (1+\alpha^2)\dot{y}^2 - \frac{V_0}{X^2}\right)$ for which the equations of motion are  $\ddot{X} = \frac{V_0(1+\alpha^2)}{X^3}, \ \ddot{y} = -\frac{\alpha V_0}{x^3}$ . The time-independent conservation laws are  $J\left(X, \dot{X}\right) = \frac{1}{2}\left(\dot{X}^2 - V_0 X^{-2}\right)$  and  $I\left(X, \dot{X}, \dot{Y}\right) = \left(2\alpha\dot{X}\dot{Y} + (1+\alpha^2)\dot{y}^2\right)$ .

We have seen that in all cases for which additional point symmetries exist the Hamiltonian Ermakov system, when it is written in Cartesian coordinates, the potential function  $\frac{V(\theta)}{\rho^2}$  becomes  $V(x, y) = \frac{V_1}{x^2} + \frac{V_2}{y^2}$  and  $V(x, y) = \frac{V_0}{(\alpha y - x)^2}$ . In these cases the additional point symmetries follow from the the gradient Killing symmetries of the Euclidian space.

## 4 The 3D generalized Ermakov system

Consider now the three-dimensional Ermakov system with Lagrangian function

$$L\left(\rho,\dot{\rho},\theta,\dot{\theta},\phi,\phi\right) = \frac{1}{2}\left(\dot{\rho}^2 + \rho^2\dot{\theta} + \rho^2\sin^2\theta \ \dot{\phi}^2 - \frac{V\left(\theta,\phi\right)}{\rho^2}\right)$$
(29)

and equations of motion

$$\ddot{\rho} - \rho \dot{\phi}^2 - \rho \sin^2 \theta \, \dot{\phi}^2 - \frac{V(\theta, \phi)}{R^3} = 0, \qquad (30)$$

$$\ddot{\theta} + \frac{2}{\rho}\dot{\rho}\dot{\theta} - \sin\theta\cos\theta \ \dot{\phi}^2 + \frac{V(\theta,\phi)_{,\theta}}{2R^4} = 0$$
(31)

$$\ddot{\phi} + \frac{2}{\rho}\dot{\rho}\dot{\phi} + \cot\theta \,\dot{\theta}\dot{\phi} + \frac{V(\theta,\phi)_{,\phi}}{2R^4\sin^2\theta} = 0.$$
(32)

At this point we discuss the analysis presented in [29]. The authors investigated the functional forms of  $V(\theta, \phi)$  for the dynamical system to admit additional conservation laws based upon point symmetries and contact symmetries. The authors determined the following functional forms for the potential  $V(\theta, \phi)$ ,

$$V_A(\theta,\phi) = V_0 , \ V_B(\theta,\phi) = V_0(\cos\theta)^{-2} , \ V_C(\theta,\phi) = V_0(\sin\theta)^{-2} ,$$
(33)

$$V_D(\theta,\phi) = V_0(\cos\theta\sin\phi)^{-2} , V_E(\theta,\phi) = V_0(\cos\theta\cos\phi)^{-2} , \qquad (34)$$

$$V_F(\theta,\phi) = \frac{\delta}{\cos\left(2\phi - 2\theta\right) + \cos\left(2\phi + 2\theta\right) + 2\cos\left(2\theta\right) - 2\cos\left(2\phi\right) + 6},$$
(35)

$$V_G(\theta,\phi) = \frac{8}{2\cos(2\phi) + 2\cos(2\theta) - \cos(2\phi - 2\theta) - \cos(2\phi + 2\theta) + 6}$$
(36)

If we simplify potential functions  $V_F$  and  $V_G$  by using trigonometric identities we find

$$V_F(\theta,\phi) = \left(\cos^2\phi\sin^2\theta - 1\right)^{-1},\tag{37}$$

$$V_G(\theta,\phi) = \left(1 - \sin^2 \phi \sin^2 \theta\right)^{-1}.$$
(38)

As we discussed above the potentials are related, for instance  $V_B$  and  $V_C$  are the same potential function, as are also potentials  $V_D$ ,  $V_E$  and  $V_F$ ,  $V_G$ . In order to understand that we should write the dynamical systems in Cartesians coordinates. we assume the transformation

$$(x, y, z) = (r \sin \theta, r \cos \theta \sin \phi, r \cos \theta \cos \phi).$$
(39)

Then

$$\rho^{-2}V_C(\theta,\phi) = x^{-2} , \ \rho^{-2}V_D(\theta,\phi) = y^{-2}$$
(40)

and

$$\rho^{-2} V_F(\theta, \phi) = \left(x^2 + z^2\right)^{-1} \tag{41}$$

while in a similar way the remaining potentials reduce to similar potentials in Cartesians coordinates, by doing a change on the variables  $\{x, y, z\}$ . These potentials are not new and have been found before and widely studied in [34]. In addition the classification scheme in [29] is incomplete, for which see below.

We want to omit any calculation and use results from the previous section as also important properties of the symmetries for the Euclidian space  $\mathbb{R}^3$ . There is a direct relation of the Noether symmetries with the elements of the Homothetic algebra of the Euclidian space, for more details we refer the reader in [33]. The proper homothetic vector of the  $\mathbb{R}^3$  is the generator of the two vector fields of the SL(2, R) algebra. However, the  $\mathbb{R}^3$  space admits six isometries. Three rotations which form the O(3) Lie algebra and three translations which form the T(3) Lie algebra. From the nature of the O(3), if an element of O(3) is a symmetry vector then we can change the coordinates  $(\theta, \phi) \to (\bar{\theta}, \bar{\phi})$ , the symmetry vector to be always the  $\partial_{\bar{\phi}}$  which provides the conservation law for the angular momentum  $\mathcal{L} = \rho^2 \sin^2 \theta \ \dot{\phi}^2$ . If we assume that two elements of O(3) to be Noether symmetries, then it follows that  $V(\theta, \phi) = V_0$ .

Therefore, when  $V(\theta, \phi) = V_A^1(\theta)$  or  $V(\theta, \phi) = V_A^{20}$ , the additional Noether symmetries follow from the elements of the O(3) Lie algebra. On the other hand, if we assume that the symmetry is generated by an element of the T(3) Lie algebra, it follows that in the Cartesians coordinates the system is similar to that found above for the two-dimensional system.

Indeed, we perform the symmetry classification

$$V_A(x, y, z) = \frac{V_1}{x^2} + \frac{V_2}{y^2} + \frac{V_3}{z^2} , \qquad (42)$$

$$V_B(x, y, z) = \frac{V_0}{x^2 + y^2} + \frac{V_1}{z^2},$$
(43)

$$V_E = \frac{V_0}{(\alpha x - \beta y - \gamma z)^2} , \quad V_E = \frac{V_0}{(\alpha x - \beta y)^2} + \frac{V_1}{z^2}, \quad (44)$$

where  $V(x, y, z) = \rho^{-2}V(\theta, \phi)$ .

It is obvious that we have found new potentials which have not been presented before in [29]. Furthermore, if we apply the Killing tensors to determine new potential functions, from the algorithm described in [35]; it is clear that there are no additional potential functions.

### 5 Conclusions

In this work we revised previous results on the classification problem of the three-dimensional Hamiltonian Ermakov system by using point and contact symmetries. We show that the classification problem solved in [29] is not correct and incomplete. We were able to show that with a constructive approach by using previous results which relate properties of the background space with the dynamical system.

We show that there is a natural extension on the classification scheme from the two-dimensional Ermakov system to the three-dimensional problem. Hence we can use the analysis to elevate the solution in higher dimensions. We summarize our analysis in the following proposition.

**Proposition:** Consider the n-dimensional Hamiltonian Ermakov system generated by the Lagrangian function

$$L\left(\rho,\dot{\rho},\theta_{1},\dot{\theta}_{1},...,\theta_{n-1},\dot{\theta}_{n-1}\right) = \frac{1}{2}\left(\dot{\rho}^{2} + \rho^{2}\left(\dot{\theta}_{1}^{2} + \sin^{2}\theta_{1}\left(\dot{\theta}_{2}^{2} + \sin^{2}\left(...\right)\right)\right) - \frac{V\left(\theta_{1},\theta_{2},...,\theta_{n-1}\right)}{\rho^{2}}\right),\tag{45}$$

which admits the SL(2, R) as Noether symmetries. The functional forms for the potential  $V(\theta_1, \theta_2, ..., \theta_{n-1})$ in which additional conservation laws linear or quadratic in the momentum are:

$$V\left(\theta_1, \theta_2, \dots, \theta_{n-1}\right) = V_0 \tag{46}$$

$$V(x_1, x_2, ..., x_n) = \sum_{i} V_i (x^i)^{-2}$$
(47)

$$V(x_1, x_2, ..., x_n) = \sum_{i} \frac{V_i}{\sum_{\kappa} x^{\kappa} x_{\kappa}} + \sum_{i} \frac{V_i}{\left(x^{\lambda}\right)^2} , \ \kappa \neq \lambda,$$
(48)

$$V(x_1, x_2, ..., x_n) = \sum_{i} \frac{V_i}{\sum_{\kappa} (\alpha^{\kappa} x_{\kappa})^2} + \sum_{i} \frac{\alpha_{\lambda} V_i}{(x^{\lambda})^2} , \ \kappa \neq \lambda,$$
(49)

where  $(x_1, x_2, ..., x_n)$  are the Cartesians coordinates for the n-dimensional Euclidian space and  $\rho^{-2}V(\theta_1, \theta_2, ..., \theta_{n-1}) = V(x_1, x_2, ..., x_n).$ 

In a future work we plan to extend this procedure to other attempts on the generalization of the Ermakov system.

#### Acknowledgements

GL was funded by Comisión Nacional de Investigación Científica y Tecnológica (CONICYT) through FONDE-CYT Iniciación 11180126. GL thanks the Department of Mathematics and the Vicerrectoría de Investigación y Desarrollo Tecnológico at Universidad Católica del Norte for financial support. This work does not have any conflicts of interest.

#### References

- [1] V. Ermakov, Univ. Isz. Kiev Series III 9, (1880) (translated by Harin A.O.)
- [2] E.Pinney, Proc. Am. Math. Soc. 1 681 (1950)
- [3] H.R. Lewis, Phys. Rev. Lett. 18 510 (1967)
- [4] H.R. Lewis and W.B. Riesenfeld, J. Math. Phys. 10, 1458 (1969)
- [5] E.R. Floyd, Phys. Lett. A 214, 259 (1996)
- [6] M. Tsamparlis and A. Paliathanasis, Symmetry 10, 233 (2018)
- [7] A. Paliathanasis and P.G.L. Leach, Phys. Lett. A 381, 1277 (2017)
- [8] S.C. Mancas and H.C. Rosu, Phys. Lett. A 378, 2113 (2014)
- [9] A. Sen and Z. Silagadze, Int. J. Theor. Phys. 59, 2187 (2020)
- [10] C. Rogers, C. Hoenselaers and J. R. Ray, J. Phys. A: Math. and Gen. 26 2625 (1993)
- [11] W.K. Schief, C. Rogers and A.P Bassom, J. Phys. A: Math. and Gen. 29 903 (1996)
- [12] T. Padmanabhan, MPLA 33, 1830005 (2018)
- [13] S. Moyo and P.G.L. Leach, J. Math. Phys. A: Math. Gen. 35, 5333 (2002)

- [14] J.R. Ray and J.L Reid, Phys. Lett. A 71, 317 (1979)
- [15] P.G.L. Leach, Phys. Lett. A 158, 102 (1991)
- [16] M. Tsamparlis and A. Paliathanasis, J. Phys. A: Math. Theor. 45, 275202 (2012)
- [17] F. Haas and J. Goedert, Phys. Lett. A 279 281 (2001)
- [18] J. Goedert and F. Haas, Phys. Lett. A 239, 348 (1998)
- [19] C. Athorne, J. Phys. A: Math and Gen 24, L1385 (1991)
- [20] K.S. Govinder and P.G.L. Leach, Phys. Lett. A 186 391 (1974)
- [21] A. Mitsopoulos and M. Tsamparlis, Eur. Phys. J. Plus 136, 933 (2021)
- [22] P. Morando, C. Muriel and A. Ruiz, Mathematics 8, 2167 (2020)
- [23] R. Campoamor-Stursberg, The Lie algebra SL(2,R) and Noether point symmetries of Lagrangian systems, Proceedings of the twentieth international conference on Geometry, Integrability and Quantization, pages 99-114, edited by I.M. Mladenov et al. (2019)
- [24] M.C. Nucci and R. Campoamor-Stursberg, J. Math. Phys. 62, 012702 (2021)
- [25] A. Ballesteros and A. Blasco, Annals of Physics 325, 2787 (2010)
- [26] J. Roberts, S. Shkoller and T.C. Sideris, Comm. Math. Phys. 375, 1003 (2020)
- [27] X. Gu, W.X. Ma and W.Y. Zhang, J. Math. Phys. 58, 053512 (2017)
- [28] M. de la Cruz, N. Gaspar, L. Jimenez-Lara and R. Linares, Annals of Physics 379, 112 (2017)
- [29] R. Naz and F.M. Mahomed, Math. Meth. Appl. Sci. 44, 4467 (2021)
- [30] H. Stephani, Differential Equations: Their Solutions using Symmetry, Campridge: Cambridge University Press, (1989)
- [31] E. Noether, Nachr. v.d. Ges. d. Wiss. zu Gottingen 235, (1918)
- [32] A. Halder, A. Paliathanasis and P.G.L. Leach, Symmetry 10, 744 (2018)
- [33] M. Tsamparlis and A. Paliathanasis, J. Phys. A: Math. Theor. 44, 175202 (2011)
- [34] M. Tsamparlis, A. Paliathanasis and L. Karpathopoulos, J. Phys. A: Math. Theor. 45, 275201 (2012)
- [35] L. Karpathopoulos, M. Tsamparlis and A. Paliathanasis, J. Geom. Phys. 133, 279 (2018)