Recurrence and transience of Rademacher series

Satyaki Bhattacharya and Stanislav Volkov*

October 14, 2022

Abstract

We introduce the notion of **a**-walk $S(n) = a_1X_1 + \cdots + a_nX_n$, based on a sequence of positive numbers $\mathbf{a} = (a_1, a_2, \ldots)$ and a Rademacher sequence X_1, X_2, \ldots . We study recurrence/transience (properly defined) of such walks for various sequences of **a**. In particular, we establish the classification in the cases where $a_k = \lfloor k^\beta \rfloor$, $\beta > 0$, as well as in the case $a_k = \lceil \log_{\gamma} k \rceil$ or $a_k = \log_{\gamma} k$ for $\gamma > 1$.

Keywords: recurrence, transience, Rademacher distribution, non-homogeneous Markov chains. **AMS subject classification**: 60G50, 60J10.

1 Introduction

We will say that a random variable X has a Rademacher distribution and write $X \sim \mathcal{R}ademacher$, if $\mathbb{P}(X = +1) = \mathbb{P}(X = -1) = \frac{1}{2}$.

Let $X_i \sim \mathcal{R}ademacher$, i = 1, 2, ..., be i.i.d., and $\mathcal{F}_n = \sigma(X_1, X_2, ..., X_n)$ be the sigma-algebra generated by the first *n* members of this sequence. Let $\mathbf{a} = (a_1, a_2, ...)$ be a non-random sequence of positive numbers. Define the **a**-walk as

$$S(n) = a_1 X_1 + a_2 X_2 + \dots + a_n X_n = \sum_{k=1}^n a_k X_k$$

with the convention S(0) = 0.

Definition 1. Let $C \ge 0$. We call the **a**-walk S defined above C-recurrent, if the event $\{|S(n)| \le C\}$ occurs for infinitely many n. (In case when C = 0, this is equivalent to the usual recurrence, i.e., S(n) = 0 for infinitely many n, so we will call the walk just recurrent.)

We call the **a**-walk transient, if it is not C-recurrent for any $C \ge 0$.

^{*}Centre for Mathematical Sciences, Lund University, Box 118 SE-22100, Lund, Sweden

Our aim is to determine the probability that the **a**-walk for given **a** and *C* is recurrent; in principle, this probability may be different¹ from 0 and 1. A simplest example of an **a**-walk is when all $a_i \equiv a \in \mathbb{R}_+$. Such a random walk is obviously a.s. recurrent since it is equivalent to the one-dimensional simple random walk.

The question of recurrence is naturally related to the Littlewood-Offord problem which deals with the maximization of probability $\mathbb{P}(S(n) = v)$ over all v, subject to various hypotheses on **a**. In particular, in [7] the authors develop an inverse Littlewood-Offord theory, using which they show that this probability is large only when the elements of **a** are contained in a generalized arithmetic progression; see also [4].

The study of **a**-walk is also somewhat relevant to the conjecture by Boguslaw Tomaszewski (1986), which says that $\mathbb{P}\left(|S(n)| \leq \sqrt{a_1^2 + \cdots + a_n^2}\right) \geq \frac{1}{2}$ for all sequences **a** and all *n*. The conjecture was recently proved by Nathan Keller and Ohad Klein in [2].

Let us first start with some general statements. First, we show that the choice of C > 0 is sometimes unimportant for the definition of C-recurrence.

Theorem 1. Suppose that $a_n \to \infty$ and at the same time $|a_n - a_{n-1}| \to 0$ as $n \to \infty$. Then if an **a**-walk is *C*-recurrent with a positive probability for some C > 0 then it is \tilde{C} -recurrent with a positive probability for all $\tilde{C} > 0$.

Proof. Since the notion of *C*-recurrence is monotone in *C*, i.e. if an **a**-walk is C_1 -recurrent for $C_1 > 0$ then it is C_2 -recurrent for all $C_2 \ge C_1$, it suffices to prove that *C*-recurrence implies $\frac{2C}{3}$ -recurrence.

Indeed, suppose the \mathbf{a} -walk is C-recurrent; formally, if we define the events

$$E = \{S(n) \in [-C, C] \text{ for infinitely many } n\},$$

$$\tilde{E} = \{S(n) \in [-2C/3, 2C/3] \text{ for infinitely many } n\}$$

then $\mathbb{P}(E) > 0$. We want to show that $\mathbb{P}(E) > 0$ as well.

Let n_1 be so large that $|a_i - a_{i-1}| < C/6$ for all $i \ge n_1$. Define the sequence $n_k, k \ge 2$, by setting

$$n_k = \min\{i \ge n_{k-1} + 1 : a_i \ge a_{n_{k-1}} + C/6\}$$

(which is well-defined since $a_i \to \infty$), then trivially

$$\frac{C}{6} \le a_{n_{k+1}} - a_{n_k} \le \frac{C}{3} \qquad \text{for each } k = 1, 2, \dots$$

$$\tag{1}$$

¹For example, if $\mathbf{a} = (1, 1, 3, 3, 3, 3, ...)$ then the **a**-walk is recurrent with probability 1/2.

Fix a positive integer K and for $y = (y_1, y_2, \dots, y_K) \in \Omega_K := \{-1, +1\}^K$ define

$$\bar{X}_K = \{X_1, X_2, \dots, X_K\};$$

 $s_y = a_1 y_1 + a_2 y_2 + \dots a_k y_K.$

Let $y \in \Omega_K$ be such that $\mathbb{P}(\{\bar{X}_K = y\} \cap E) > 0$. Observe that

$$\{\bar{X}_K = y\} \cap E = \{\bar{X}_K = y\} \cap B_K(s_y)$$

where

$$B_{K}^{+}(u) = \{ \text{there exist } m_{1} < m_{2} < \dots \text{ such that } u + \sum_{i=K+1}^{m_{j}} a_{i}X_{i} \in [0, C] \};$$

$$B_{K}^{-}(u) = \{ \text{there exist } m_{1}' < m_{2}' < \dots \text{ such that } u + \sum_{i=K+1}^{m_{j}'} a_{i}X_{i} \in [-C, 0] \};$$

$$B_{K}(u) = B_{K}(u)^{+} \cup B_{K}(u)^{-}.$$

Since $\{\bar{X}_K = y\}$ and $B_K(u)$ are independent, we have

$$\mathbb{P}(\{\bar{X}_K = y\} \cap B_K(s_y)) = \mathbb{P}(\bar{X}_K = y) \mathbb{P}(B_K(s_y)).$$

Consequently, $\mathbb{P}(B_K(s_y)) > 0$, and as a result, $\mathbb{P}(B_K^+(s_y)) > 0$ or $\mathbb{P}(B_K^-(s_y)) > 0$ (or both).

Let $\Omega_K^* \subseteq \Omega_K$ contain those y_s for which there is an index k such that $n_{k+2} \leq K$ and $y_{n_k} = -1$, $y_{n_{k+1}} = +1$, $y_{n_{k+2}} = -1$; let k be the smallest such index. For $y \in \Omega_K^*$ define the mappings $\sigma^+, \sigma^- : \Omega_K^* \to \Omega_K$ by

$$\sigma^{+}(y) = \begin{cases} -y_i, & \text{if } i = n_k \text{ or } i = n_{k+1}; \\ y_i, & \text{otherwise}; \end{cases}$$
$$\sigma^{-}(y) = \begin{cases} -y_i, & \text{if } i = n_{k+1} \text{ or } i = n_{k+2}; \\ y_i, & \text{otherwise.} \end{cases}$$

Then for $y \in \Omega_K^*$

$$s_{\sigma^+(y)} = s_y + 2a_{n_k} - 2a_{n_k+1} \in [s_y - 2C/3, s_y - C/3],$$

$$s_{\sigma^-(y)} = s_y - 2a_{n_k} + 2a_{n_k+1} \in [s_y + C/3, s_y + 2C/3].$$

As a result, it is not hard to see that

$$\{\bar{X}_K = \sigma^+(y)\} \cap B_K^+(s_y) \subseteq \left\{\sum_{i=1}^m a_i X_i \in [-2C/3, 2C/3] \text{ for infinitely many } ms\right\} = \tilde{E},$$
$$\{\bar{X}_K = \sigma^-(y)\} \cap B_K^-(s_y) \subseteq \left\{\sum_{i=1}^m a_i X_i \in [-2C/3, 2C/3] \text{ for infinitely many } ms\right\} = \tilde{E}.$$

Since at least one of $B_K^+(s_y)$ and $B_K^-(s_y)$ has a positive probability, $\mathbb{P}\left(\bar{X}_K = \sigma^{\pm}(y)\right) = 2^{-K}$ and the events on the LHS are independent, we conclude that $\mathbb{P}(\tilde{E}) > 0$.

Now it only remains to show that there exists $y \in \Omega_K^*$ such that $\mathbb{P}(\{\bar{X}_K = y\} \cap E) > 0$. Let $\kappa := \kappa(K) = \max\{k \in \mathbb{Z}_+ : n_k \leq K\}$; obviously, $\kappa(K) \to \infty$ as $K \to \infty$. If we choose y from Ω_K uniformly, we can trivially bound the probability that $y \notin \Omega_K^*$ by²

$$\left(1-\frac{1}{8}\right)^{\lfloor\kappa/3\rfloor} \to 0 \quad \text{as } \kappa \to \infty$$

by grouping together triples $(X_{n_1}, X_{n_2}, X_{n_3}), (X_{n_4}, X_{n_5}, X_{n_6})$, etc.; in each such a triple

$$\mathbb{P}\left((X_{n_k}, X_{n_{k+1}}, X_{n_{k+2}}) = (-1, +1, -1)\right) = 1/8.$$

Hence

$$\mathbb{P}(E) = \sum_{y \in \Omega_K} \mathbb{P}(\{\bar{X}_K = y\} \cap E) = \sum_{y \in \Omega_K^*} \mathbb{P}(\{\bar{X}_K = y\} \cap E) + \mathbb{P}\left(\{\bar{X}_K \in \Omega_K \setminus \Omega_K^*\} \cap E\right).$$
(2)

Since $\mathbb{P}\left(\left\{\bar{X}_{K} \in \Omega_{K} \setminus \Omega_{K}^{*}\right\} \cap E\right) \leq \mathbb{P}\left(\bar{X}_{K} \in \Omega_{K} \setminus \Omega_{K}^{*}\right)$, by making K sufficiently large, we can ensure that the second term on the RHS of (2) is less than $\mathbb{P}(E)$, implying that there exist some $y \in \Omega_{K}^{*}$ such that $\mathbb{P}(\left\{\bar{X}_{K} = y\right\} \cap E) > 0$, as required.

Our next result shows that if the sequence \mathbf{a} is non-decreasing, then the walk will always "jump" over 0 infinitely many times, even if the walk is not *C*-recurrent.

Theorem 2. Suppose that a_i is a non-decreasing positive sequence. Then the event $\{S(n) > 0\}$ holds for infinitely many n a.s. The same is true for the event $\{S(n) < 0\}$.

The theorem immediately follows from symmetry and the more general

Proposition 1. Suppose that a_i is a non-decreasing sequence, m is an integer such that $a_{m+1} > 0$, and S(m) = A > 0. Define

$$\tau = \inf\{k \ge 0 : S(m+k) \le 0\}.$$

Let $Y_j \sim \mathcal{R}$ ademacher be i.i.d., and

$$\tilde{\tau} = \inf\{k \ge 0: Y_1 + Y_2 + \dots + Y_k \le -r\}$$

where $r = \lceil A/a_{m+1} \rceil$; note that $\tilde{\tau} < \infty$ a.s. and that, in fact, $Y_1 + \cdots + Y_{\tilde{\tau}} = -r$. Then τ is stochastically smaller than $\tilde{\tau}$, that is,

$$\mathbb{P}(\tau > m) \le \mathbb{P}(\tilde{\tau} > m), \quad m = 0, 1, 2, \dots$$

²exact: see the sequence A005251 in the online encyclopedia of integer sequences (https://oeis.org/A005251), $\mathbb{P}\left(\bar{X}_{K} \notin \Omega_{K}^{*}\right) \approx \lambda^{\kappa}, \ \lambda = \frac{\sqrt[3]{100+12\sqrt{69}}}{6} + \frac{2}{3\sqrt[3]{100+12\sqrt{69}}} + \frac{2}{3} = 0.877..., \ \kappa = \kappa(K)$

Proof. We will use coupling. Indeed, we can write

$$S(m+j) = A + a_{m+1}Y_1 + a_{m+2}Y_2 + \dots + a_{m+j}Y_j, \quad j = 1, 2, \dots$$

Suppose that $\tilde{\tau} = k$, that is

$$Y_1 > -r, Y_1 + Y_2 > -r, \dots, Y_1 + Y_2 + \dots + Y_{k-1} > -r;$$

 $Y_1 + Y_2 + \dots + Y_{k-1} + Y_k = -r.$

Then, recalling that a_i is a non-decreasing sequence,

$$S(m+k) = A + a_{m+1}Y_1 + \dots + a_{m+k-1}Y_{k-1} + a_{m+k}Y_k$$

$$\leq A + a_{m+1}Y_1 + \dots + a_{m+k-2}Y_{k-2} + a_{m+k-1}Y_{k-1} + a_{m+k-1}Y_k \quad \text{(since } Y_k = -1\text{)}$$

$$= A + a_{m+1}Y_1 + \dots + a_{m+k-2}Y_{k-2} + a_{m+k-1}[Y_{k-1} + Y_k]$$

$$\leq A + a_{m+1}Y_1 + \dots + a_{m+k-2}Y_{k-2} + a_{m+k-2}[Y_{k-1} + Y_k]$$

$$= A + a_{m+1}Y_1 + \dots + a_{m+k-2}[Y_{k-2} + Y_{k-1} + Y_k]$$

$$\leq \dots \leq A + a_{m+1}[Y_1 + \dots + Y_k] = A - ra_{m+1} \leq 0,$$

since Y_k , $Y_{k-1} + Y_k$, $Y_{k-2} + Y_{k-1} + Y_k$, ..., $Y_1 + \cdots + Y_k$ are all negative. Therefore, $\tau \leq \tilde{\tau}$.

Throughout the paper we will use a version of the Azuma-Hoeffding inequality; compare with the results of [3].

Lemma 1. Suppose that b_1, b_2, \ldots, b_m is a sequence of non-negative numbers and $S = b_1Y_1 + b_2Y_2 + \cdots + b_mY_m$, where $Y_j \sim \mathcal{R}$ ademacher are *i.i.d.* Then

$$\mathbb{P}\left(|\mathcal{S}| \ge A\right) \le 2\exp\left(-\frac{A^2}{2(b_1^2 + \dots + b_m^2)}\right) \text{ for all } A > 0.$$
(3)

We also state the following fairly standard result.

Lemma 2. Let $T_i = Y_1 + \cdots + Y_i$ be a simple random walk. Suppose that L_k and y_k , $k = 1, 2, \ldots$, are two sequences such that $L_k \to \infty$, $y_k \to \infty$ and $y_k/\sqrt{L_k} \to r > 0$. Then

$$\lim_{k \to \infty} \mathbb{P}\left(\max_{1 \le i \le L_k} T_i \ge y_k\right) = 2 \mathbb{P}(\eta \ge r) = 2 - 2 \Phi(r)$$

where $\eta \sim \mathcal{N}(0, 1)$ and $\Phi(\cdot)$ is its CDF.

Proof. Let $\tilde{y}_k = \lceil y_k \rceil \in \mathbb{Z}_+$. By the reflection principle,

$$\mathbb{P}\left(\max_{1\leq i\leq L_{k}}T_{i}\geq y_{k}\right) = \mathbb{P}\left(\max_{1\leq i\leq L_{k}}T_{i}\geq \tilde{y}_{k}\right) = 2\mathbb{P}(T_{L_{k}}\geq \tilde{y}_{k}) - \mathbb{P}(T_{L_{k}}=\tilde{y}_{k})$$
$$= 2\mathbb{P}\left(\frac{T_{L_{k}}}{\sqrt{L_{k}}}\geq \frac{\tilde{y}_{k}}{\sqrt{L_{k}}}\right) + O\left(\frac{1}{\sqrt{L_{k}}}\right) \to 2\mathbb{P}(\eta\geq r)$$

by the Central Limit Theorem, using also the fact that $\tilde{y}_k/y_k \to 1$.

5

2 Integer-valued a-walks

Suppose that the sequence **a** contains only integers.

Proposition 2. Let $z \in \mathbb{Z}$. Suppose that the sequence

$$\int_0^{\pi} \cos(tz) \prod_{k=1}^n \cos(ta_k) dt, \qquad n = 1, 2, \dots$$

is summable. Then the events $\{S(n) = z\}$ occur for finitely many n a.s.

Proof. The result follows from standard Fourier analysis. Indeed,

$$\mathbb{E}e^{itS(n)} = \sum_{k \in \mathbb{Z}} e^{itk} \mathbb{P}(S(n) = k)$$

where the sum above goes, in fact, effectively over a finite number of ks (as $|S(n)| \le a_1 + \cdots + a_n$). At the same time,

$$\int_{-\pi}^{\pi} e^{it(k-z)} \mathrm{d}t = \begin{cases} 2\pi, & \text{if } k = z; \\ 0, & \text{if } k \in \mathbb{Z} \setminus \{z\}. \end{cases}$$

By changing the order of summation and integration, we obtain

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} \mathbb{E}e^{it(S(n)-z)} dt = \frac{1}{2\pi} \sum_{k \in \mathbb{Z}} \int_{-\pi}^{\pi} e^{it(k-z)} \mathbb{P}(S(n) = k) dt = \mathbb{P}(S(n) = z).$$

On the other hand,

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} \mathbb{E}e^{it(S(n)-z)} dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-itz} \prod_{k=1}^{n} \mathbb{E}e^{ita_k X_k} dt = \frac{1}{\pi} \int_{0}^{\pi} \cos(tz) \prod_{k=1}^{n} \cos(ta_k) dt$$

by the symmetry of cos and the fact that the imaginary part must equal zero. Now the result follows from the Borel-Cantelli lemma, since $\sum_{n} \mathbb{P}(S(n) = z) < \infty$.

Corollary 1. Suppose that the sequence

$$\int_0^{\pi} \left| \prod_{k=1}^n \cos(ta_k) \right| \mathrm{d}t, \qquad n = 1, 2, \dots$$

is summable. Then the \mathbf{a} -walk is transient a.s.

Proof. From Proposition 2 we know that for each $z \in \mathbb{Z}$

$$\pi \mathbb{P}(S(n) = z) = \int_0^\pi \cos(tz) \prod_{k=1}^n \cos(ta_k) dt \le \int_0^\pi \left| \cos(tz) \prod_{k=1}^n \cos(ta_k) \right| dt \le \int_0^\pi \left| \prod_{k=1}^n \cos(ta_k) \right| dt.$$

Hence the event $\{S(n) = z\}$ occurs finitely often a.s. for each z. Since for each C > 0 there are only finitely many integers in [-C, C] we conclude that the walk is not C-recurrent a.s. for every C.

An interesting and quite natural example is when $\mathbf{a} = (1, 2, 3, ...)$, i.e., $a_i = i$. It was previously published in the IMS Bulletin [8], in the Student Puzzle Corner no. 37.

Theorem 3. The **a**-walk with $\mathbf{a} = (1, 2, 3, ...)$ is a.s. transient.

This statement follows from a much stronger Theorem 4, but for the sake of completeness, we present its self-contained proof.

Proof of Theorem 3. Let $A_n = \{S(n) = 0\} = \{X_1 + 2X_2 + \dots + nX_n = 0\}$. Then $\mathbb{P}(A_n) = Q_n/2^n$, where

 Q_n = number of ways to put \pm in the sequence $*1 * 2 * 3 * \cdots * n$ such that the sum equals 0.

For example, $Q_1 = Q_2 = 0$, $Q_3 = Q_4 = 2$, $Q_5 = Q_6 = 0$, $Q_7 = 8$, $Q_8 = 14$, etc. It was essentially shown in [6] that

$$Q_n \sim \sqrt{\frac{6}{\pi}} \frac{2^n}{n^{3/2}} \qquad \text{when } n \mod 4 \in \{0, 3\}$$

(and zero otherwise) as $n \to \infty$, meaning that the ratio of the RHS and the LHS converges to one. Consequently, $\sum_{n} \mathbb{P}(A_n) \sim \sum_{n} \frac{\text{const}}{n^{3/2}} < \infty$ and the events A_n occur a.s. finitely often by the Borel-Cantelli lemma. Hence the walk is a.s. not recurrent.

Moreover, since for any $m \in \mathbb{Z}$

$$\mathbb{P}(S(n+2|m|) = S(n) - m \mid \mathcal{F}_n) \ge \frac{1}{2^{2m}}$$

(by making the signs of $X_{n+1}, X_{n+2}, \ldots, X_{n+2|m|}$ alternate), we conclude that if the event $\{S(n) = m\}$ occurs infinitely often, then A_n shall also occur infinitely often a.s., leading to contradiction. As a result, $\mathbb{P}(\{S(n) = m\} \text{ i.o.}) = 0$ for all integer ms, and thus the walk is a.s. not C-recurrent for any non-negative C.

Remark 1. Though the (1, 2, 3, ...)-walk is transient, it still can jump over zero infinitely many times, as it was shown by Theorem 2.

In fact, Theorem 3 can be generalized greatly, using the result from [5], or even a weaker result of [1], which provide the estimates for the maximum number of solutions of the equation $\sum_{i=1}^{n} \varepsilon_i a_i = t \text{ where } \epsilon_i \in \{0, 1\} \text{ while } a_i \text{'s and } t \text{ are all integers.}$

Theorem 4. Let \mathbf{a} be such that all a_i 's are distinct integers. Then \mathbf{a} -walk is a.s. transient.

Proof. The main result of [5] implies that for any $\epsilon > 0$

$$\operatorname{card}(\{(x_1, x_2, \dots, x_n) : \text{ all } x_i = \pm 1, \ a_1 x_1 + \dots + a_n x_n = m\}) \le \frac{(1+\epsilon)2^{n+3}}{n^{3/2}\sqrt{\pi}}$$

for all $n \ge n_0(\epsilon)$ and all m. Setting $\epsilon = 1$, and fixing $m \in \mathbb{Z}$, we obtain that

$$\sum_{n=n_0(1)}^{\infty} \mathbb{P}(S(n)=m) \le \sum_{n=n_0(1)}^{\infty} \frac{2 \cdot 2^{n+3}}{n^{3/2} \sqrt{\pi}} \times \frac{1}{2^n} = \frac{16}{\sqrt{\pi}} \sum_{n=n_0(1)}^{\infty} \frac{1}{n^{3/2}} < \infty.$$

Therefore, by the Borel-Cantelli lemma, only finitely many events $\{S(n) = m\}$ occur a.s. Since S(n) takes only integer values, this implies that $\{|S(n)| \leq C\}$ happens finitely often a.s. for any C > 0.

Remark 2.

(a) It is not difficult to see that under the condition of Theorem 4 it suffices that all a_k 's are distinct only starting from some $k_0 \ge 1$.

(b) If $a_k = \lfloor k^{\beta} \rfloor$ with $\beta \ge 1$, then we immediately have a.s. transience by Theorem 4.

3 A non-trivial recurrent example

We assume here that $\mathbf{a} = (B_1, B_2, B_3, ...)$ where each B_k is a consecutive block of k's of length precisely $L_k \ge 1$. Denote also by $i_k = 1 + L_1 + L_2 + \cdots + L_{k-1}$ the index of the first element of the k-th block. For example, if $L_k = 2^k$, then $i_k = 2^k - 1$ and

$$\mathbf{a} = (1, 1, \underbrace{2, 2, 2, 2}_{L_2 \text{ times}}, \underbrace{3, 3, 3, 3, 3, 3, 3, 3, 3}_{L_3 \text{ times}}, \underbrace{4, \dots, 4}_{L_4 \text{ times}}, \dots),$$

one can also notice that $a_i = \lfloor \log_2(i+1) \rfloor = \lceil \log_2(i+2) \rceil - 1.$

Theorem 5. Suppose that for some $\varepsilon > 0$, r > 0, and k_0 we have

$$\frac{\frac{L_k}{L_1 + L_2 + \dots + L_{k'}} \ge (2 + \varepsilon) \ln k;}{\frac{L_k}{L_{k'+1} + L_{k'+2} + \dots + L_{k-1}} \ge 2r;} \qquad (4)$$

$$\frac{L_k}{L_k \ge k^4,}$$

whenever $k - k' \geq \frac{k}{\ln k} - 2$ and $k, k' \geq k_0$. Then the **a**-walk described above is a.s. recurrent.

Remark 3. One can easily check that the conditions of the theorem are satisfied if $a_k = \lfloor (\log_{\gamma} k)^{\beta} \rfloor$, where $\gamma > 1$ and $\beta \in (0, 1]$.

Proof of Theorem 5. We will proceed in FIVE steps.

Step 1: Preliminaries

First, we need the following lemma, which is probably known.

Lemma 3. Let $m \in \mathbb{Z}_+$ and T_m be a simple symmetric random walk on \mathbb{Z}^1 , that is, $T_m = Y_1 +$ $\cdots + Y_m$, where $Y_i \sim \mathcal{R}$ advector are *i.i.d.* There exists a universal constant $c_1 > 0$ such that for all integers z such that $|z| \leq 2\sqrt{m}$, assuming that m is sufficiently large and m + z is even,

$$\mathbb{P}(T_m = z) \ge \frac{c_1}{\sqrt{m}}.$$

Proof. W.l.o.g. assume $z \ge 0$. We have

$$\mathbb{P}(T_m = z) = \mathbb{P}\left(\frac{T_m + m}{2} = \frac{z + m}{2}\right) = \mathbb{P}(\tilde{T} = w)$$

where $\tilde{T} \sim \text{Bin}(m, 1/2)$ and $w = \frac{z+m}{2} \in \mathbb{Z}_+$. Note that $\tilde{m} \leq w \leq \tilde{m} + \sqrt{m}$ where $\tilde{m} = m/2$. So

$$\mathbb{P}(\tilde{T} = w) = \binom{m}{w} 2^{-m} = \binom{2\tilde{m}}{\tilde{m}} \frac{1}{2^{2\tilde{m}}} \frac{\tilde{m}! \tilde{m}!}{w!(m-w)!} = \frac{1+o(1)}{\sqrt{\pi\tilde{m}}} \frac{(2\tilde{m}-w+1)(2\tilde{m}-w+2)\dots\tilde{m}}{(\tilde{m}+1)(\tilde{m}+2)\dots w} \\
= \frac{1+o(1)}{\sqrt{\pi\tilde{m}}} \left(1 - \frac{w-\tilde{m}}{\tilde{m}+1}\right) \left(1 - \frac{w-\tilde{m}}{\tilde{m}+2}\right) \dots \left(1 - \frac{w-\tilde{m}}{w}\right) \\
\ge \frac{1+o(1)}{\sqrt{\pi\tilde{m}}} \left(1 - \frac{\sqrt{m}}{\tilde{m}+1}\right)^{w-\tilde{m}} \ge \frac{1+o(1)}{\sqrt{\pi\tilde{m}}} \left(1 - \frac{\sqrt{2}+o(1)}{\sqrt{\tilde{m}}}\right)^{\sqrt{2\tilde{m}}} = \frac{e^{-2}+o(1)}{\sqrt{\pi m/2}} \ge \frac{0.1}{\sqrt{m}} \\
\text{or large enough } m.$$

for large enough m.

Corollary 2. Let T_m , m = 0, 1, 2, ..., be as simple symmetric random walk as in Lemma 3. Assume that m and k are positive integers such that $k^2 \leq m$. Let $u \in \mathbb{Z}$, and either k is odd, or both k and m-u are even. Then for large ks

$$\mathbb{P}(T_m - u \mod k = 0) \ge \frac{c_1}{2k}$$

where c_1 is the constant from Lemma 3.

Proof of Corollary 2. First, assume that m, and hence u, are both even. Since $(T_m - u) \mod k =$ $0 \iff T_m = \tilde{u} \mod k$, where $\tilde{u} = (u \mod k) \in \{0, 1, 2, \dots, k-1\}$, it suffices to show the statement for \tilde{u} .

Let $M = \lfloor 2\sqrt{m} \rfloor \in (2\sqrt{m} - 1, 2\sqrt{m}]$ and define

$$\mathbb{I} = [-M, -M+1, \dots, -1, 0, 1, \dots, M] = \mathbb{I}_0 \cup \mathbb{I}_1;$$
$$\mathbb{I}_0 = \{z \in \mathbb{I} : z \text{ is even}\}; \quad \mathbb{I}_1 = \{z \in \mathbb{I} : z \text{ is odd}\}.$$

There are at least M elements in each \mathbb{I}_0 and \mathbb{I}_1 .

If k is odd, then each of these two sets contains at least $\lfloor \frac{M}{k} \rfloor$ elements z such that $z = \tilde{u} \mod k$. If m is even (odd, resp.) for all z either in \mathbb{I}_0 (in \mathbb{I}_1 , resp.) by Lemma 3 for large ks (and hence large m) we have $\mathbb{P}(T_m = z) \ge c_1/\sqrt{m}$. Consequently,

$$\mathbb{P}(T_m = \tilde{u} \mod k) \ge \sum_{z \in \mathbb{I}, \ z = \tilde{u} \mod k} \mathbb{P}(T_m = z) \ge \left\lfloor \frac{M}{k} \right\rfloor \times \frac{c_1}{\sqrt{m}} \ge \left(\frac{M}{k} - 1\right) \times \frac{c_1}{\sqrt{m}}$$
$$\ge \left(\frac{2\sqrt{m} - 1}{k} - 1\right) \times \frac{c_1}{\sqrt{m}} \ge \left(1 - \frac{1}{k}\right) \times \frac{c_1}{\sqrt{m}} = \frac{c_1}{k} - O(k^{-2})$$

since $m \ge k^2$.

If k is even, then if m is even (and thus a is also even) then \mathbb{I}_0 contains at least $\lfloor \frac{M}{k} \rfloor$ elements z such that $z = \tilde{u} \mod k$ and at the same time Lemma 3 is applicable for $z \in \mathbb{I}_0$. On the other hand, if m (and so u) is odd then \mathbb{I}_1 contains at least $\lfloor \frac{M}{k} \rfloor$ elements z such that $z = \tilde{u} \mod k$ and Lemma 3 is applicable for $z \in \mathbb{I}_1$. The rest of the proof is the same as for the case when k is odd.

Step 2: Splitting S(n)

Recall that that i_k denotes the first index of block k and note that the sum of all the steps within block k can be represented as

$$S(i_{k+1}-1) - S(i_k-1) = k \cdot T_k, \qquad T_k = X_1^{(k)} + \dots + X_{L_k}^{(k)}$$

where $X_i^{(k)}$'s are i.i.d. *Rademacher* random variables.

For $m = 2, \ldots$, let

$$k_m = \begin{cases} \lfloor m \ln m \rfloor & \text{if } \lfloor m \ln m \rfloor \text{ is odd;} \\ \lfloor m \ln m \rfloor + 1 & \text{if } \lfloor m \ln m \rfloor \text{ is even.} \end{cases}$$
(5)

Thus k_m is always odd; k_m , m = 2, 3, ... equal 1, 3, 5, 7, 9, 13, 15, 19, 23, etc. Define also

$$A_m = \{ S(j) = 0 \text{ for some } i_{k_m} \le j < i_{k_m+1} \},\$$

the event that S(j) hits zero for the steps within block B_{k_m} , and the sequence of sigma-algebras

$$\mathcal{G}_m = \mathcal{F}_{i_{k_{m+1}}-1} = \sigma\left(\bigcup_{\ell=1}^{k_m} \sigma\left(X_1^{(\ell)}, X_2^{(\ell)}, \dots, X_{L_\ell}^{(\ell)}\right)\right).$$

Intuitively, \mathcal{G}_m contains all the information about the walk during its steps corresponding to the first k_m blocks.

To simplify notations, let us now write $k = k_m$ and $k' = k_{m-1}$, and observe that

$$k - k' = k_m - k_{m-1} \ge m \ln m - (m-1)\ln(m-1) - 2 = \ln m - 1 + O\left(\frac{1}{m}\right) \ge \ln m - 2 \qquad (6)$$

for large m.

Let us split S(j) where $j \in [i_k, i_{k+1})$, as follows:

$$S(j) = S(i_{k'}) + \sum_{n=k'+1}^{k-1} \left(X_1^{(n)} + \dots + X_{L_n}^{(n)} \right) + k \cdot \left(X_1^{(k)} + \dots + X_{j-i_k}^{(k)} \right)$$
$$= S(i_{k'}) + \left[\sum_{n=k'+1}^{k-2} \left(X_1^{(n)} + \dots + X_{L_n}^{(n)} \right) + (k-1) \sum_{\ell=1}^{i_k - 2k^2 - 1} X_{\ell}^{(k-1)} \right]$$
$$+ (k-1) \cdot \Sigma_3 + k \cdot \left(X_1^{(k)} + \dots + X_{j-i_k}^{(k)} \right)$$
$$= \Sigma_1 + \Sigma_2 + (k-1) \cdot \Sigma_3 + k \cdot \Sigma_4$$

where $\Sigma_1 = S(i_{k'})$ and

$$\Sigma_{2} = \sum_{n=k'+1}^{k-2} nT_{n} + (k-1)T'_{k-1}, \quad T'_{k-1} = \sum_{\ell=i_{k-1}}^{i_{k}-1-2k^{2}} X_{\ell}^{(k-1)};$$

$$\Sigma_{3} = X_{i_{k}-2k^{2}}^{(k-1)} + X_{i_{k}-2k^{2}+1}^{(k-1)} + \dots + X_{i_{k}-2}^{(k-1)} + X_{i_{k}-1}^{(k-1)};$$

$$\Sigma_{4} = X_{1}^{(k)} + X_{2}^{(k)} + \dots + X_{j-i_{k}}^{(k)}.$$

Note that Σ_i , i = 1, 2, 3, 4 are independent, and Σ_3 has precisely $2k^2$ terms.

Step 3: Estimating Σ_1

Recall that $k = k_m, k' = k_{m-1}$ and let

$$E_{m-1} = \left\{ |\Sigma_1| < k\sqrt{L_k} \right\} \in \mathcal{G}_{m-1}.$$

By Lemma 1 and (4), assuming k is large,

$$\mathbb{P}(E_{m-1}^{c}) \leq \mathbb{P}(|S(k')| \geq k'\sqrt{L_{k}}) \leq 2\exp\left(-\frac{k'^{2} \cdot L_{k}}{2(L_{1}+2^{2} \cdot L_{2}+3^{2} \cdot L_{3}+\dots+k'^{2} \cdot L_{k'})}\right)$$

$$\leq 2\exp\left(-\frac{L_{k}}{2(L_{1}+L_{2}+L_{3}+\dots+L_{k'})}\right) \leq 2\exp\left(-(1+\varepsilon/2)\ln k\right) = \frac{2}{k^{1+\varepsilon/2}} =:\varepsilon_{m}.$$
(7)

Step 4: Estimating Σ_2

Again, by Lemma 1 and (4), assuming that k is sufficiently large,

$$\mathbb{P}\left(|\Sigma_2| \ge k\sqrt{\frac{L_k}{r}}\right) \le 2\exp\left(-\frac{k^2r^{-1}L_k}{2[(k'+1)^2L_{k'+1} + \dots + (k-2)^2L_{k-2} + (k-1)^2(L_{k-1} - 2k^2)]}\right) \\
\le 2\exp\left(-\frac{r^{-1}L_k}{2[L_{k'+1} + \dots + L_{k-1} - 2k^2]}\right) \le 2\exp\left(-1\right) = 0.7357588824\dots$$

Consequently,

$$\mathbb{P}\left(|\Sigma_2| < k\sqrt{L_k/r}\right) \ge 0.2 \quad \text{for large } k.$$
(8)

Step 5: Finishing the proof

We have a trivial lower bound

$$\mathbb{P}(A_m \mid E_{m-1}, \mathcal{G}_{m-1}) \ge \mathbb{P}\left(A_m \mid |\Sigma_2| < k\sqrt{\frac{L_k}{r}}, E_{m-1}, \mathcal{G}_{m-1}\right) \times \mathbb{P}\left(|\Sigma_2| < k\sqrt{\frac{L_k}{r}} \mid E_{m-1}, \mathcal{G}_{m-1}\right)$$
$$=: (*) \times 0.2 \quad \text{for large } k \tag{9}$$

by (8), since the second multiplier equals $\mathbb{P}\left(|\Sigma_2| < k\sqrt{L_k/r}\right)$ by independence. Let

$$\mathsf{Div}_{k} = \{\Sigma_{1} + \Sigma_{2} + (k-1)\Sigma_{3} = 0 \mod k\} = \{\Sigma_{1} + \Sigma_{2} - \Sigma_{3} = 0 \mod k\}$$

Since only on the event Div_k , it is possible that S(j) = 0 for some j (since the step sizes are $\pm k$ in the block B_k), we conclude that for large k

$$(*) = \mathbb{P}\left(A_m \mid \mathsf{Div}_k, |\Sigma_2| < k\sqrt{\frac{L_k}{r}}, E_{m-1}, \mathcal{G}_{m-1}\right) \times \mathbb{P}\left(\mathsf{Div}_k \mid |\Sigma_2| < k\sqrt{\frac{L_k}{r}}, E_{m-1}, \mathcal{G}_{m-1}\right)$$
$$\geq \mathbb{P}\left(A_m \mid \mathsf{Div}_k, |\Sigma_2| < k\sqrt{\frac{L_k}{r}}, E_{m-1}, \mathcal{G}_{m-1}\right) \times \frac{c_1}{2k} \tag{10}$$

due to the fact that by Corollary 2, $\mathbb{P}(\text{Div}_k \mid \mathcal{F}_{i_k-2k^2-1}) \geq c_1/(2k)$. On the other hand,

$$\mathbb{P}\left(A_m \mid \mathsf{Div}_k, |\Sigma_2| < k\sqrt{\frac{L_k}{r}}, E_{m-1}, \mathcal{G}_{m-1}\right) \ge \min_{z \in Z_k} \mathbb{P}(z + T_m = 0 \text{ for some } m \in [0, L_k])$$
$$\ge \beta := 1 - \Phi\left(r^{-1/2} + 3\right) > 0 \tag{11}$$

where $z + T_m$ is a simple random walk starting at z (see Lemma 3), and

$$Z_k = \left\{ z \in \mathbb{Z} : |z| \le (r^{-1/2} + 3) \sqrt{L_k} \right\}$$

Indeed, using the last part of (4), and the conditions we imposed, we have

$$|\Sigma_1 + \Sigma_2 + (k-1)\Sigma_3| \le k\sqrt{L_k} + k\sqrt{L_k/r} + 2(k-1)k^2 < (1+r^{-1/2}+2)k\sqrt{L_k}$$

for large $k, S(j) = [\Sigma_1 + \Sigma_2 + (k-1)\Sigma_3] + k \cdot \Sigma_4$, and by Lemma 2

$$\liminf_{k \to \infty} \min_{z \in Z_k} \mathbb{P}(z + T_m = 0 \text{ for some } m \in [0, L_k]) \ge 2 \mathbb{P}(\eta > r^{-1/2} + 3) = 2\beta,$$

so the minimum in (11) is $\geq \beta$ for all sufficiently large k.

Finally, from (9), (10), and (11) we get that

$$\sum_{m} \mathbb{P}\left(A_{m} \mid E_{m-1}, \mathcal{G}_{m-1}\right) \geq \sum_{m} \frac{0.2 c_{1}\beta}{2m \log m} = +\infty$$

and $\mathbb{P}(E_m^c)$ is summable by (7), so we can apply Lemma 5 of Appendix 1 to conclude that events A_m occur infinitely often and thus our **a**-walk is recurrent.

4 Continuous example

The example of **a**-walk described in Theorem 5 roughly corresponds to the case $a_k = \lceil \log_{\gamma} k \rceil$, $k = 1, 2, \ldots$ But what if a_k 's take non-integer values, but, for example, equal

$$a_k = \log_\gamma k \equiv c \ln k, \qquad k = 1, 2, \dots,$$

where $\gamma = e^{1/c} > 1$? In this Section we will study this example. It is unreasonable to assume that such **a**-walk is recurrent, because of the irrationality of the step sizes, however, we might want to investigate if this walk is *C*-recurrent for *some* C > 0. Our main result is the following

Theorem 6. Let c > 0 and $a_k = c \ln k$. Then the **a**-walk is a.s. C-recurrent for every C > 0.

To prove this theorem, it is sufficient to show that whatever the value c > 0 is, $\{|S(n)| \leq 3\}$ happens i.o. almost surely. Indeed, take any C > 0. Then the statement that **a'**-walk with $a'_k = \frac{3c}{C} \ln k$, $k = 1, 2, \ldots$, is 3-recurrent is equivalent to the statement that **a**-walk with $a_k = c \ln k$ is *C*-recurrent.

The proof will proceed similarly to that of Theorem 5. Let us define k_m slightly differently from (5); namely, let

$$k_m = \begin{cases} \lfloor m \ln m \rfloor & \text{if } \lfloor m \ln m \rfloor \text{ is even}; \\ \lfloor m \ln m \rfloor - 1 & \text{if } \lfloor m \ln m \rfloor \text{ is odd.} \end{cases}$$

Thus now k_m are always even. As before, set $k = k_m$, and $k' = k_{m-1}$, and define

$$i_k = \lceil \gamma^k \rceil = \max\{i \ge 1 : a_i < k\} + 1 = \min\{i \ge 1 : a_i \ge k\} \in [\gamma^k, \gamma^k + 1),$$

i.e., the first index when a_i starts exceeding k. For $i \in J_k := [i_k, i_{k+1})$ write

$$S(i) = S(i_{k'} - 1) + [S(i_k - 1) - S(i_{k'} - 1) - \Sigma_3] + \Sigma_3 + [S(i) - S(i_k - 1)]$$

= $\Sigma_1 + \Sigma_2 + \Sigma_3 + \Sigma_4(i)$ (12)

where

$$\Sigma_3 = \left[S(i_{k-1} + k^2 - 1) - S(i_{k-1} - 1)\right] + \left[S(i_k) - S(i_k - k^2)\right].$$

Note that Σ_i , i = 1, 2, 3, 4 are independent, and Σ_3 has $2 \cdot k^2$ terms, and contains the first k^2 and the last k^2 steps of the walk, when the step sizes lie in [k, k + 1).

Let

$$E_{m-1} = \left\{ |\Sigma_1| < k\sqrt{i_k} \right\} = \left\{ S(i_{k_{m-1}}) < k_m \sqrt{i_{k_m}} \right\}$$
(13)

By Lemma 1, since $a_i < k' < k$ for $i < i_{k'}$,

$$\mathbb{P}(E_{m-1}^{c}) = \mathbb{P}\left(|\Sigma_{1}| \ge k\sqrt{i_{k}}\right) \le 2\exp\left(-\frac{i_{k}k^{2}}{2\sum_{j=1}^{i_{k'}}a_{j}^{2}}\right) \le 2\exp\left(-\frac{i_{k}}{2i_{k'}}\right) \le 2\exp\left(-\frac{\gamma^{k}-1}{2\gamma^{k'}}\right)$$
$$= 2\exp\left(-\frac{\gamma^{k-k'}(1+o(1))}{2}\right) = 2\exp\left(-\frac{\gamma^{\ln m-2}}{2+o(1)}\right) = 2\exp\left(-\frac{m^{\ln \gamma}}{2\gamma^{2}+o(1)}\right) =:\varepsilon_{m-1}$$
(14)

using (6) for k sufficiently large³. Observe that ε_m is summable.

Similarly, by Lemma 1

$$\mathbb{P}\left(|\Sigma_2| \ge 2k\sqrt{i_k}\right) \le 2\exp\left(-\frac{4k^2i_k}{2k^2\left(i_k - i_{k'} - 2k^2\right)}\right) < 2e^{-2} = 0.27\dots$$

Hence,

$$\mathbb{P}(F_m) \ge 0.72 \quad \text{where } F_m = \left\{ |\Sigma_2| < 2k\sqrt{i_k} \right\}.$$
(15)

Lemma 4. Let $n = k^2$ where k is an even positive integer, and assume also that k is sufficiently large. Suppose that X_i , Y_i , i = 1, 2, ..., n, are *i.i.d.* Rademacher. Let

$$T = (k-1)(X_1 + X_2 + \dots + X_n) + k(Y_1 + Y_2 + \dots + Y_n).$$
(16)

Then

$$\mathbb{P}(T=j) \ge \frac{c_1^2}{4n} \quad for \ each \ j=0, \pm 2, \pm 4, \dots, \pm n$$

where c_1 is the constant from Lemma 3.

³Note that (6) was stated for k_m defined slightly differently, however, it holds here as well.

Proof. It follows from Corollary 2 that

$$\mathbb{P}(X_1 + \dots + X_n = \ell) \ge \frac{c_1}{2k}, \quad \mathbb{P}(Y_1 + \dots + Y_n = \ell) \ge \frac{c_1}{2k} \quad \text{for all even } \ell \text{ such that } |\ell| \le 2k.$$
(17)

Let $j = 2\tilde{j} \in \{0, 2, 4, \dots, n-2, n\}$. Consider the sequence of k-1 numbers

$$\tilde{j}, \ \tilde{j}-k, \ \tilde{j}-2k, \ \tilde{j}-3k, \dots, \tilde{j}-(k-2)k;$$

they all give different remainders when divided by k-1. Hence there must be an $m \in \{0, 1, \ldots, k-2\}$ such that $\tilde{j} - mk = b(k-1)$ and b is an integer; moreover, since $0 \leq \tilde{j} \leq n/2$, we have $b \in \left[-\frac{k(k-2)}{k-1}, \frac{n}{2(k-1)}\right]$. For such m and b we have $j = 2\tilde{j} = (2m)k + (2b)(k-1)$, and, since both |2m| and $|2b| \leq 2k$,

$$\mathbb{P}(T=j) \ge \mathbb{P}(X_1 + \dots + X_n = 2b) \cdot \mathbb{P}(Y_1 + \dots + Y_n = 2m) \ge \left(\frac{c_1}{2k}\right)^2 = \frac{c_1^2}{4n}$$

by (17). The result for negative j follows by symmetry.

Corollary 3. Let $\varepsilon = \frac{2ck^4}{\gamma^{k-1}}$. Then for large even k

$$\mathbb{P}\left(\Sigma_3 \in [j-\varepsilon, j+\varepsilon]\right) \ge \frac{c_1^2}{4k^2} \quad for \ each \ j=0, \pm 2, \pm 4, \dots, \pm k^2.$$

Proof. Σ_3 has the same distribution as

$$\sum_{\ell=1}^{k^2} c \ln(i_{k-1} - 1 + \ell) X_{\ell} + \sum_{\ell=1}^{k^2} c \ln(i_k - \ell) Y_{\ell}$$

for some i.i.d. $X_{\ell}, Y_{\ell} \sim \mathcal{R}ademacher$. At the same time, for $\ell \geq 1$,

$$\begin{aligned} |c\ln(i_{k-1} - 1 + \ell) - (k - 1)| &= |c\ln(\lceil \gamma^{k-1} \rceil + \ell - 1) - (k - 1)| \\ &\leq |c\ln(\gamma^{k-1} + \ell) - (k - 1)| = c\ln\left(1 + \frac{\ell}{\gamma^{k-1}}\right) \leq \frac{c\ell}{\gamma^{k-1}} \end{aligned}$$

Similarly,

$$k - c\ln(i_k - \ell) = k - c\ln(\lceil \gamma^k \rceil - \ell) = k - c\ln(\gamma^k - \ell') = -c\ln\left(1 - \frac{\ell'}{\gamma^k}\right) \in \left[0, \frac{c\ell}{\gamma^{k-1}}\right]$$

for some $\ell' \in [\ell - 1, \ell]$, assuming $\ell = o(\gamma^k)$. As a result, for T defined by (16),

$$|\Sigma_3 - T| \le \sum_{\ell=1}^{k^2} \frac{2c\ell}{\gamma^{k-1}} = \frac{ck^2(k^2 + 1)}{\gamma^{k-1}} \le \frac{2ck^4}{\gamma^{k-1}}.$$

Now the result follows from Lemma 4.

Proof of Theorem 6. Recall that $J_k = [i_k, i_{k+1})$ and define

$$A_m = \{S(i) = 0 \text{ for some } i \in J_{k_m}\}.$$

Then

$$\mathbb{P}(A_m \mid E_{m-1}, \mathcal{G}_{m-1}) \ge 0.72 \times \mathbb{P}(A_m \mid F_m, E_{m-1}, \mathcal{G}_{m-1})$$
(18)

(please see the definition of F_m in (15)). Recall formula (12) and write

$$\tilde{S}(i) = S(i) - \Sigma_3 = \Sigma_1 + \Sigma_2 + \Sigma_4(i).$$

From now on assume that $|\Sigma_1| < k\sqrt{i_k}$ and $|\Sigma_2| < 2k\sqrt{i_k}$, that is, E_{m-1} and F_m occur. Also assume w.l.o.g. that $\Sigma_1 + \Sigma_2 \ge 0$. Let

$$L_k = i_{k+1} - i_k - k^2 = (\gamma - 1)\gamma^k + o(\gamma^k).$$

Consider a simple random walk with steps $Y_i \sim \mathcal{R}ademacher$ during its first L_k steps. The probability that its minimum will be equal to or below the level $-3\sqrt{i_k} = -\frac{3+o(1)}{\sqrt{\gamma-1}}\sqrt{L_k}$ converges by Lemma 2 to

$$2\mathbb{P}\left(\eta > \frac{3}{\sqrt{\gamma - 1}}\right) = 2 - 2\Phi\left(\frac{3}{\sqrt{\gamma - 1}}\right) =: 2c_2 \in (0, 1)$$

as $k \to \infty$ (recall that $\eta \sim \mathcal{N}(0,1)$). As a result, by Proposition 1, the probability that for some $j_0 \in \{i_k, i_k + 1, i_k + 2, \dots, i_k + L_k - 1\}$ we have the downcrossing, that is,

 $\tilde{S}(j_0 - 1) \ge 0 > \tilde{S}(j_0)$

is bounded below by c_2 for k sufficiently large. Formally, let

$$j_0 = \inf\{j > i_k : \tilde{S}(j) < 0\};$$

$$\mathcal{C}_0 = \{i_k \le j_0 \le i_k + L_k - 1\},$$

so we have showed that on $E_{m-1} \cap F_m \cap \{\Sigma_1 + \Sigma_2 > 0\}$ we have $\mathbb{P}(\mathcal{C}_0) > c_2$ for large k.

Now assume that event \mathcal{C}_0 occurred and define additionally

$$b_0 = \tilde{S}(j_0) \in (-k - 1, 0];$$

$$\mathcal{C} = \left\{ \max_{0 \le h \le k^2} \sum_{g=1}^h X_{j_0 + g} \ge k \right\}$$

Again, from Lemma 2,

$$\mathbb{P}(\mathcal{C}) = 2 \mathbb{P}(X_{j_0+1} + X_{j_0+2} + \dots + X_{j_0+k^2} \ge k) \to 2(1 - \Phi(1)) = 0.3173\dots \text{ as } k \to \infty.$$

From now on assume that k is so large that $\mathbb{P}(\mathcal{C}) > 0.2$. On the event \mathcal{C} there exists an increasing sequence j_1, j_2, \ldots, j_k such that

$$j_0 < j_1 < j_2 < \dots < j_k \le j_0 + k^2 < i_{k+1}$$

such that $X_{j_0+1} + X_{j_0+2} + \cdots + X_{j_\ell} = \ell$ for each $\ell = 1, 2, \ldots, k$, since the random walk must pass through each integer in $\{1, 2, \ldots, k\}$ in order to reach level k.

For $\ell = 1, 2, \ldots, k$, define

$$b_{\ell} := \tilde{S}(j_{\ell}) = b_0 + \sum_{h=j_0+1}^{j_{\ell}} a_h X_h = b_0 + a_{j_0} \left[\sum_{h=j_0+1}^{j_{\ell}} X_h \right] + \sum_{h=j_0+1}^{j_{\ell}} (a_h - a_{j_0}) X_h$$
$$= b_0 + a_{j_0}\ell + O\left(\frac{k^4}{\gamma^k}\right)$$

since for $h \in [j_0, j_0 + k^2] \subseteq [i_k, i_{k+1})$ we have

$$|a_h - a_{j_0}| = c \left| \ln \frac{a_h}{a_{j_0}} \right| \le c \left| \ln \frac{i_k + k^2}{j_k} \right| = O\left(\frac{k^2}{\gamma^k}\right).$$

As a result,

$$-(k+1) < b_0 < b_1 < b_2 < \dots < b_{k-1} < (k-1)(k+1) < k^2$$

and moreover the distance between consecutive b_g 's is at least two (provided k is large). For $\ell = 0, 1, ..., k - 1$ define

$$\tilde{b}_{\ell} = \sup\left\{x \in 2\mathbb{Z} : x + b_{\ell} \in \left[\frac{1}{2}, 3 - \frac{1}{2}\right)\right\} \equiv -2\left\lfloor\frac{b_{\ell}}{2} - \frac{1}{4}\right\rfloor.$$

Then \tilde{b}_{ℓ} 's are all distinct even integers satisfying $|\tilde{b}_{\ell}| \leq k^2$.

As a result,

$$\mathbb{P}(A_m \mid F_m, E_{m-1}, \mathcal{G}_{m-1}) \geq c_2 \times 0.2 \times \mathbb{P}(S(i) \in [0, 3] \text{ for some } i \in J_k \mid \mathcal{C}, \mathcal{C}_0)$$

$$\geq c_2 \times 0.2 \times \mathbb{P}(\tilde{S}(i_\ell) + \Sigma_3 \in [0, 3] \text{ for some } \ell = 0, 1, \dots, k-1 \mid \mathcal{C}, \mathcal{C}_0)$$

$$= c_2 \times 0.2 \times \mathbb{P}(b_\ell + \Sigma_3 \in [0, 3] \text{ for some } \ell = 0, 1, \dots, k-1 \mid \mathcal{C}, \mathcal{C}_0)$$

$$\geq c_2 \times 0.2 \times \mathbb{P}(\tilde{\Sigma}_3 \in [\tilde{b}_\ell - \varepsilon, \tilde{b}_\ell + \varepsilon] \text{ for some } \ell = 0, 1, \dots, k-1 \mid \mathcal{C}, \mathcal{C}_0)$$

$$= c_2 \times 0.2 \times \sum_{\ell=0}^{k-1} \mathbb{P}(\Sigma_3 \in [\tilde{b}_\ell - \varepsilon, \tilde{b}_\ell + \varepsilon] \mid \mathcal{C}, \mathcal{C}_0) \geq c_2 \times 0.2 \times k \times \frac{c_1^2}{4k^2} = \frac{c_1^2 c_2}{20k}$$

assuming that ε in Corollary 3 is sufficiently small. Finally,

$$\mathbb{P}(A_m \mid E_{m-1}, \mathcal{G}_{m-1}) \ge 0.72 \times \mathbb{P}(A_m \mid F_m, E_{m-1}, \mathcal{G}_{m-1}) \ge \frac{0.72 c_1^2 c_2}{20k_m} \ge \frac{0.036 c_1^2 c_2}{m \ln m}$$

the sum of which diverges. Hence, recalling (14), we can again apply Lemma 5.

5 Sublinear growth of step sizes

Throughout this Section we assume

$$a_k = \lfloor k^\beta \rfloor, \qquad 0 < \beta < 1$$

Proposition 3. Let $S(n) = a_1 X_1 + \dots + a_n X_n$ where $a_k = \lfloor k^{\beta} \rfloor$, $0 < \beta < 1$. Then

$$\mathbb{P}(|S(n)| = z) \le \frac{\nu}{n^{1/2+\beta}}$$
 for all large n ,

for some $\nu > 0$.

Proof. Let $F_n(t) = \prod_{k=1}^n |\cos(ta_k)|$. For all $z \in \mathbb{Z}$ we have

$$\mathbb{P}(S(n) = z) = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-itz} \mathbb{E}e^{itS(n)} dt \le \frac{1}{2\pi} \int_{-\pi}^{\pi} \left| \mathbb{E}e^{itS(n)} \right| dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} F_n(t) dt \le \frac{\nu}{n^{1/2+\beta}}$$

by Lemma 6 for some $\nu > 0$, for all large n.

Theorem 7. Suppose that $a_k = \lfloor k^{\beta} \rfloor$, $0 < \beta < 1$. Then the **a**-walk is a.s. transient.

Proof. In the case $\beta > 1/2$ the a.s. transience follows immediately from Borel-Cantelli lemma and Proposition 3, as $\sum_{n} \mathbb{P}(|S(n)| \leq C) < \infty$ for each $C \geq 1$.

Assume from now on that $0 < \beta \leq 1/2$. Define k_m , Δ_m , m_n as in Case 3 of the proof of Lemma 6. Fix a positive integer M and consider now only those n for which $m_n = M$. Let $I_M = \{k_M, k_M + 1, \ldots, k_{M+1} - 1\}$. Note that the elements of I_M are precisely those n for which $a_n = M$, and that the caridnality of I_M is of order $M^{1/\beta-1}$. Next, fix some $z \in \mathbb{Z}$ and define

$$E_M = E_M(z) = \{S(n) = z \text{ for some } n \in I_M\}\}.$$

For each z we will show that $\sum_{M} \mathbb{P}(E_M) < \infty$, and so by the Borel-Cantelli lemma a.s. only finitely many events E_M occur. Since S(n) takes only integer values, this will imply that the walk is not *C*-recurrent for any $C \ge 0$.

So, fix z from now on, write $S(n) = S(k_M) + R(n)$ where

$$R(n) = \sum_{i=k_M}^n a_i X_i = M \sum_{i=k_M}^n X_i,$$

Observe also that $S(k_M)$ and R(n) are independent. In order S(n) = z for some $n \in I_M$, we need that $S(k_M) = z \mod M$. Let $Q = M^{\frac{1}{2\beta}+1-\varepsilon}$ for an $\varepsilon \in (0, 1/2)$. Assuming M is so large that $Q \ge 2|z|$,

$$\mathbb{P}(|R(n)| \ge Q - |z|) \le 2 \exp\left(-\frac{(Q/2)^2}{2M^2 \cdot (k_{M+1} - k_M)}\right) = 2 \exp\left(-\frac{\beta Q^2}{(8 + o(1))M^2 \cdot M^{\frac{1}{\beta} - 1}}\right)$$
$$= 2 \exp\left(-\frac{\beta M^{1 - 2\varepsilon}}{8 + o(1)}\right) \quad \text{for all } n \in I_M$$

by Lemma 1; hence

$$\mathbb{P}(\max_{n \in I_M} |R(n)| \ge Q - |z|) \le |I_M| \times 2 \exp\left(-\frac{\beta M^{1-2\varepsilon}}{8 + o(1)}\right) =: \alpha_M = O\left(M^{\frac{1}{\beta} - 1} e^{-\frac{\beta M^{1-2\varepsilon}}{8 + o(1)}}\right),$$

which is summable in M. So,

$$\mathbb{P}(E_M) \le \mathbb{P}\left(E_M, \max_{n \in I_M} |R(n)| < Q - |z|\right) + \mathbb{P}\left(\max_{n \in I_M} |R(n)| \ge Q - |z|\right)$$
$$= \mathbb{P}\left(E_M, S(k_M) = z \mod M, \max_{n \in I_M} |R(n)| < Q - |z|\right) + \alpha_M = (*) + \alpha_M$$

where the term α_M is summable since $1 - 2\varepsilon > 0$. Since E_M implies implies $-S(k_M) = R(n) - z$ for some $n \in I_M$,

$$(*) \leq \mathbb{P}(|S(k_M)| < Q, \ S(k_M) = z \ \text{mod} \ M) = \sum_{j: \ |j| < Q, \ j = z \ \text{mod} \ M} \mathbb{P}(|S(k_M)| = j)$$

$$\leq \frac{\nu}{k_M^{1/2+\beta}} \times |\{j: \ |j| < Q, \ j = z \ \text{mod} \ M\}| \leq \frac{\nu + o(1)}{M^{1+\frac{1}{2\beta}}} \times \left[\frac{2Q+1}{M} + 1\right] = \frac{(\nu + o(1))}{\pi M^{1+\varepsilon}}$$

by Proposition 3. The RHS is summable in M, which concludes the proof.

Remark 4. By setting $\varepsilon = 1/2 - \delta/2$, where $\delta > 0$ is very close to zero, we can ensure that

$$\mathbb{P}(|S(n)| < M^{1/2-\delta} \text{ for some } n \in I_M) \le \sum_{\substack{z:|z| < M^{1/2-\delta}}} \mathbb{P}(E_M(z)) \le \left[\frac{\text{const}}{M^{1+\varepsilon}} + \alpha_M\right] \times 2M^{1/2-\delta}$$
$$= \frac{2 \text{ const}}{M^{1+\delta/2}} + 2M^{1/2-\delta}\alpha_M$$

is summable. Hence, a.s. eventually |S(n)| will be larger that $n^{\beta/2-\delta}$ for any $\delta > 0$.

Appendix 1: Modified conditional Borel-Cantelli lemma

Lemma 5. Suppose that we have an increasing sequence of sigma-algebras \mathcal{G}_m and a sequence of \mathcal{G}_m -measurable events A_m and E_m such that

$$\mathbb{P}(A_m \mid E_{m-1}, \mathcal{G}_{m-1}) \ge \alpha_m, \quad \mathbb{P}(E_m^c) \le \varepsilon_m \quad a.s.$$

where the non-negative α_n and ε_n satisfy

$$\sum_{m} \alpha_m = \infty, \quad \sum_{m} \varepsilon_m < \infty.$$
(19)

Then $\mathbb{P}(A_m \ i.o.) = 1.$

Proof. Let $m > \ell \ge 1$ and $B_{\ell,m} = \bigcap_{i=\ell}^m A_i^c$. We need to show that for any $\ell \ge 1$

$$\mathbb{P}(B_{\ell,\infty}) = \mathbb{P}\left(A_{\ell}^c \cap A_{\ell+1}^c \cap A_{\ell+2}^c \cap \dots\right) = 0.$$
(20)

We have for $m \ge \ell + 1$

$$\mathbb{P}(B_{\ell,m}) = \mathbb{P}\left(A_m^c \cap B_{\ell,m-1}\right) \leq \mathbb{P}\left(A_m^c \cap B_{\ell,m-1} \cap E_{m-1}\right) + \mathbb{P}\left(E_{m-1}^c\right) \\
= \mathbb{P}\left(A_m^c \mid B_{\ell,m-1} \cap E_{m-1}\right) \mathbb{P}\left(B_{\ell,n-1} \cap E_{m-1}\right) + \mathbb{P}\left(E_{m-1}^c\right) \\
\leq \mathbb{P}\left(A_m^c \mid B_{\ell,m-1} \cap E_{m-1}\right) \mathbb{P}\left(B_{\ell,m-1}\right) + \varepsilon_{m-1} \leq (1 - \alpha_m) \mathbb{P}\left(B_{\ell,m-1}\right) + \varepsilon_{m-1}.$$
(21)

By induction over $m, m-1, m-2, \ldots, \ell+1$ in (21), we get that

$$\mathbb{P}(B_{\ell,m}) \leq \varepsilon_{m-1} + (1 - \alpha_m) \left[(1 - \alpha_{m-1}) \mathbb{P}(B_{\ell,m-2} \mid \mathcal{G}_m) + \varepsilon_{m-2} \right] \leq \dots$$

$$\leq \varepsilon_{m-1} + (1 - \alpha_m) \varepsilon_{m-2} + (1 - \alpha_m) (1 - \alpha_{m-1}) \varepsilon_{m-3} + \dots$$

$$+ (1 - \alpha_m) (1 - \alpha_{m-1}) \dots (1 - \alpha_{\ell+2}) \varepsilon_{\ell} + (1 - \alpha_m) (1 - \alpha_{m-1}) \dots (1 - \alpha_{\ell+1}).$$

Hence, for any integer $M \in (\ell, m)$

$$\mathbb{P}(B_{\ell,m}) \leq \varepsilon_{m-1} + \varepsilon_{m-2} + \dots + \varepsilon_M$$

$$+ (1 - \alpha_m)(1 - \alpha_{m-1}) \cdots (1 - \alpha_{M+1})\varepsilon_{M-1} + \dots + (1 - a_m)(1 - \alpha_{m-1}) \dots (1 - \alpha_{\ell+2})\varepsilon_\ell$$

$$+ (1 - \alpha_m)(1 - \alpha_{m-1}) \dots (1 - \alpha_{\ell+1})$$

$$\leq [\varepsilon_{m-1} + \varepsilon_{m-2} + \dots + \varepsilon_M]$$

$$+ (1 - \alpha_m)(1 - \alpha_{m-1}) \cdots (1 - \alpha_{M+1})[\varepsilon_{M-1} + \varepsilon_{M-2} + \dots + \varepsilon_\ell + 1]$$

Fix any $\delta > 0$. By (19) we can find an M be so large that $\sum_{i=M}^{\infty} \varepsilon_i < \delta/2$. Then, again by (19), there exists an $m_0 > M$ such that $\prod_{i=M+1}^{m_0} (1 - \alpha_i) < \frac{\delta}{2\left(1 + \sum_{i=\ell}^{M-1} \varepsilon_i\right)}$. Hence, for all $m \ge m_0$ we have $\mathbb{P}(B_{\ell,m}) \le \delta/2 + \delta/2 = \delta$. Since $\delta > 0$ is arbitrary, and $B_{\ell,m}$ is a decreasing sequence of events in m, we conclude that $\mathbb{P}(B_{\ell,\infty}) = 0$, as required.

Appendix 2: Generalization of Blair Sullivan's results

Let $a_k = \lfloor k^{\beta} \rfloor$, where $0 < \beta < 1$. Lemma 6. Let $F_n(t) = \prod_{k=1}^n |\cos(ta_k)|$. Then $\int_{0}^{\pi} F_n(t) \ln \sqrt{8\pi(1+2\beta)} + o(1)$

$$\int_{-\pi}^{\pi} F_n(t) dt = \frac{\sqrt{8\pi (1+2\beta) + o(1)}}{n^{\beta+1/2}} \quad as \ n \to \infty$$

Remark 5. Note that for $\beta = 1$ we would have obtained the same result as in [6].

Proof. We will proceed in the spirit of [6]. Note that by symmetry

$$\int_{-\pi}^{\pi} F_n(t) dt = 2 \int_0^{\pi} F_n(t) dt = 2 \int_0^{\pi/2} F_n(t) dt + 2 \int_0^{\pi/2} F_n(\pi - t) dt = 4 \int_0^{\pi/2} F_n(t) dt$$

since $|\cos((\pi - t)a_k)| = |\cos(\pi a_k - ta_k)| = |\cos(ta_k)|$ as a_k is an integer. Let $\varepsilon > 0$ be very small and define

$$I_0 = \left[0, \frac{1}{n^{\beta+1/2-\varepsilon}}\right], \quad I_1 = \left[\frac{1}{n^{\beta+1/2-\varepsilon}}, \frac{1}{n^{\beta}}\right], \quad I_2 = \left[\frac{1}{n^{\beta}}, \frac{c_1}{n^{\beta}}\right], \quad I_3 = \left[\frac{c_1}{n^{\beta}}, \frac{\pi}{2}\right]$$

for some $c_1 > 1$ to be determined later. Then

$$\int_0^{\pi/2} F_n(t) dt = \int_{I_0} F_n(t) dt + \int_{I_1} F_n(t) dt + \int_{I_2} F_n(t) dt + \int_{I_3} F_n(t) dt.$$

We will show that the contribution of all the integrals, except the first one, is negligible, and estimate the value of the first one.

First, observe that when $0 \le ta_k \le \pi/2$ for all $k \le n$, by the elementary inequality $|\cos u| \le e^{-u^2/2}$ valid for $|u| \le \pi/2$, we have

$$F_n(t) \le \prod_{k=1}^n \exp\left(-\frac{t^2 a_k^2}{2}\right) = \exp\left(-\frac{t^2}{2} \sum_{k=1}^n a_k^2\right) = \exp\left(-\frac{t^2 n^{2\beta+1}(1+o(1))}{2(1+2\beta)}\right)$$
(22)

since $a_k^2 = k^{2\beta} (1 + o(1)).$

<u>Case 0</u>: $t \in I_0$

Here $ta_k \leq \frac{1}{n^{1/2-\varepsilon}} \ll 1$, hence for *n* large enough

$$\frac{(ta_k)^2}{2} \le -\ln\cos(ta_k) = \frac{(ta_k)^2}{2} + O\left((ta_k)^4\right) \le (1+o(1))\frac{(ta_k)^2}{2}$$

yielding $F_n(t) = \exp\left(-\frac{t^2 n^{2\beta+1}\rho_{n,t}}{2(1+2\beta)}\right)$ where $\rho_{n,t} = 1 + o(1)$ for large *n* (compare with (22)). Since for any r > 0 we have

$$\int_{0}^{n^{-\beta-1/2+\varepsilon}} \exp\left(-\frac{t^2 n^{2\beta+1} r}{2(1+2\beta)}\right) dt = \frac{1}{n^{1/2+\beta}} \int_{0}^{n^{\varepsilon}} \exp\left(-\frac{s^2 r}{2(1+2\beta)}\right) ds$$
$$= \frac{1}{n^{1/2+\beta}} \left[\sqrt{\frac{\pi(1+2\beta)}{2r}} + o(1)\right]$$

where the main term is monotone in r, by substituting $r = \rho_{n,t} = 1 + o(1)$ we conclude that

$$\int_{I_0} F_n(t) dt = \frac{1}{n^{1/2+\beta}} \left[\sqrt{\pi(1/2+\beta)} + o(1) \right].$$

<u>Case 1</u>: $t \in I_1$

Since $ta_k \leq 1 < \pi/2$, by (22) for some $C_2 > 0$ we have $F_n(t) \leq \exp\left(-\frac{n^{2\varepsilon}(1+o(1))}{2(1+2\beta)}\right) \leq e^{-C_2 n^{2\varepsilon}}$, so $\int_{I_1} F_n(t) dt \leq e^{-C_2 n^{2\varepsilon}}$, which decays faster than polynomially.

<u>Case 2</u>: $t \in I_2$

As in Case 2 in [6], we will use monotonicity of $F_n(t)$ in n. Let $r = c_1^{-1/\beta} \in (0,1)$ then $\lfloor rn \rfloor^{\beta} \leq (rn)^{\beta} = \frac{n^{\beta}}{c_1}$, consequently by (22), since $t \leq \frac{c_1}{n^{\beta}}$,

$$F_{\lfloor rn \rfloor}(t) \le \exp\left(-\frac{t^2 (rn)^{2\beta+1} (1+o(1))}{2(1+2\beta)}\right)$$

and since $F_n(t) \leq F_{\lfloor rn \rfloor}(t)$, we get a similar bound as in Case 1.

Case 3:
$$t \in I_3$$

Let

$$k_{m} = \inf\{k \in \mathbb{Z}_{+} : k^{\beta} \ge m\} = \lceil m^{1/\beta} \rceil, \quad m = 1, 2, \dots;$$

$$\Delta_{m} = k_{m+1} - k_{m} = \beta^{-1} m^{\gamma} + O(m^{1/\beta-2}) + \rho_{0}, \qquad \gamma := \frac{1-\beta}{\beta},$$

where $|\rho_0| \leq 1$. Then

 $a_k = m$ if and only if $k \in \{k_m, k_m + 1, \dots, k_m + \Delta_m - 1 (\equiv k_{m+1} - 1)\}.$

For $n \in \mathbb{Z}_+$ let

$$m_n = \max\{m: k_m \le n\} = n^{\beta} (1 + o(1)), \quad n \in [k_{m_n}, k_{m_n} + \Delta_{m_n} - 1].$$

By the inequality between the mean geometric and the mean arithmetic,

$$F_n(t) = \sqrt{\prod_{k=1}^n \cos^2(ta_k)} = \left(\sqrt[n]{\prod_{k=1}^n \cos^2(ta_k)}\right)^{n/2} \le \left(\frac{\sum_{k=1}^n \cos^2(ta_k)}{n}\right)^{n/2} \\ = \left(\frac{1}{2} + \frac{U_n(t)}{2n}\right)^{n/2} \quad \text{where } U_n(t) = \sum_{k=1}^n \cos(2ta_k).$$

We will show that if t is not too small, then for some $0 \le c < 1$ we have $U_n(t) \le cn$ and hence $F_n(t) \le \left(\frac{1+c}{2}\right)^{n/2}$. In order to do that, first note that

$$U_n(t) \le \sum_{k=1}^{k_{m_n}} \cos(2ta_k) + (n - k_{m_n}) = \sum_{m=1}^{m_n} \Delta_m \cos(2tm) + (n - k_{m_n}).$$

Let $r \in (0, 1)$ and assume w.l.o.g. that rm_n is an integer. For $m \in [rm_n + 1, m_n]$ we have

$$A \leq \Delta_m \leq \overline{A}$$
 where $A = \beta^{-1} (rm_n)^{\gamma} + O(1), \quad \overline{A} = \beta^{-1} m_n^{\gamma} + O(1).$

Consequently,

$$\sum_{m=rm_{n}+1}^{m_{n}} \Delta_{m} \cos(2tm) \leq \sum_{m=rm_{n}+1}^{m_{n}} \left[\bar{A} \cdot \mathbf{1}_{\cos(2tm) \geq 0} + A \cdot \mathbf{1}_{\cos(2tm) < 0} \right] \cos(2tm)$$

$$= \sum_{m=rm_{n}+1}^{m_{n}} \left[\bar{A} - A \right] \cos(2tm) \mathbf{1}_{\cos(2tm) \geq 0} + \sum_{m=rm_{n}+1}^{m_{n}} A \cos(2tm)$$

$$\leq (1 - r)m_{n} \left(\bar{A} - A \right) + A \sum_{m=rm_{n}+1}^{m_{n}} \cos(2tm) = (1 - r)m_{n} \left(\bar{A} - A \right)$$

$$+ A \left(\cos^{2} \left(rtm_{n} + t \right) - \cos^{2} (tm_{n} + t) + \frac{\cos t}{2 \sin t} \left[\sin(2t(m_{n} + 1)) - \sin(2t(rm_{n} + 1)) \right] \right)$$

$$\leq \frac{m_{n}^{1/\beta}}{\beta} (1 - r) \left[1 - r^{\gamma} + O(m_{n}^{-\gamma}) \right] + m_{n}^{\gamma} \left(\frac{r^{\gamma}}{\beta} + O(m_{n}^{-\gamma}) \right) \left(1 + \frac{1}{|\sin t|} \right)$$

Hence, since $m_n^{1/\beta} = n + o(n)$,

$$\begin{aligned} U_n(t) &\leq \sum_{m=1}^{rm_n} \Delta_m + \sum_{m=rm_n+1}^{m_n} \Delta_m \cos(2tm) + (n-k_{m_n}) = k_{rm_n+1} + \sum_{m=rm_n+1}^{m_n} \Delta_m \cos(2tm) + O(\Delta_{m_n}) \\ &\leq r^{1/\beta}n + \frac{m_n^{1/\beta}}{\beta} (1-r) \left[1 - r^{\gamma} + O(m_n^{-\gamma}) \right] + m_n^{\gamma} \left(\frac{r^{\gamma}}{\beta} + O(m_n^{-\gamma}) \right) \left(1 + \frac{1}{|\sin t|} \right) + O(m_n^{\gamma\beta}) \\ &\leq n \left(r^{1/\beta} + \frac{(1-r)(1-r^{\gamma})}{\beta} \right) + \frac{4n^{1-\beta}\beta^{-1}r^{\gamma}}{|\sin t|} + o(n) \end{aligned}$$

Consider now the the function

$$h(r,\beta) := r^{1/\beta} + \frac{(1-r)(1-r^{\gamma})}{\beta} = r^{1/\beta} + \frac{(1-r)\left(1-r^{1/\beta-1}\right)}{\beta}$$

and note that

$$h(1-\beta,\beta) = (1-\beta)^{1/\beta} + 1 - (1-\beta)^{1/\beta-1} = 1 - \beta(1-\beta)^{1/\beta-1} \le 1 - e^{-1}\beta < 1 - \beta/3$$

since $\sup_{\beta \in (0,1)} (1-\beta)^{1/\beta-1} = e^{-1}$ by elementary calculus. So if we set $r = 1 - \beta \in (0,1)$, by noting $t \leq 2 \sin t$ for $t \in [0, \pi/2]$, we conclude that $U_n(t) \leq (1 - \frac{\beta}{4}) n$ provided that $t \geq \frac{c_1}{n^{\beta}}$ for some $c_1 > 0$. Consequently, $\int_{I_3} F_n(t) dt \leq (1 - \frac{\beta}{8})^{n/2}$ for large n, which converges to zero exponentially fast.

Acknowledgment

We thank Edward Crane and Andrew Wade for providing us with useful references. The research is partially supported by Swedish Science Foundation grants VR 2019-04173 and Crafoord foundation grant no. 20190667.

References

- Erdős, P. Extremal problems in number theory. 1965 Proc. Sympos. Pure Math., Vol. VIII pp. 181–189 Amer. Math. Soc., Providence, R.I.
- [2] Keller, Nathan and Klein, Ohad. (2021). Proof of Tomaszewski's Conjecture on Randomly Signed Sums. https://arxiv.org/abs/2006.16834
- [3] Montgomery-Smith, S. J. The distribution of Rademacher sums. Proc. Amer. Math. Soc. 109 (1990), no. 2, 517–522.
- [4] Nguyen, Hoi H. A new approach to an old problem of Erdős and Moser. J. Combin. Theory Ser. A 119 (2012), no. 5, 977–993.
- [5] Sárközi, A.; Szemerédi, E. Uber ein Problem von Erdös und Moser. (German) Acta Arith. 11 (1965), 205–208.
- [6] Sullivan, Blair D. On a conjecture of Andrica and Tomescu. J. Integer Seq. (2013), 16, no. 3, Article 13.3.1, 6 pp.
- [7] Tao, Terence and Vu, Van H. Inverse Littlewood-Offord theorems and the condition number of random discrete matrices. Ann. of Math. 169 (2009), no. 2, 595–632.
- [8] IMS Bulletin, Volume 51, Issue 2: March 2022.