Edge coloring of graphs of signed class 1 and 2

Robert Janczewski¹

Department of Algorithms and System Modeling, Faculty of Electronics, Telecommunication and Informatics, Gdańsk University of Technology, ul. Narutowicza 11/12, Gdańsk, Poland

Krzysztof Turowski^{2,*}

Theoretical Computer Science Department, Jagiellonian University, Kraków, 30–348, Poland

Bartłomiej Wróblewski^{3,**}

Department of Algorithms and System Modeling, Faculty of Electronics, Telecommunication and Informatics, Gdańsk University of Technology, ul. Narutowicza 11/12, Gdańsk, Poland

Abstract

Recently, Behr [1] introduced a notion of the chromatic index of signed graphs and proved that for every signed graph (G, σ) it holds that

$$\Delta(G) \le \chi'(G, \, \sigma) \le \Delta(G) + 1,$$

where $\Delta(G)$ is the maximum degree of G and χ' denotes its chromatic index.

In general, the chromatic index of (G, σ) depends on both the underlying graph G and the signature σ . In the paper we study graphs G for which $\chi'(G, \sigma)$ does not depend on σ . To this aim we introduce two new classes of graphs, namely 1^{\pm} and 2^{\pm} , such that graph G is of class 1^{\pm} (respectively, 2^{\pm}) if and only if $\chi'(G, \sigma) = \Delta(G)$ (respectively, $\chi'(G, \sigma) = \Delta(G) + 1$) for all possible signatures σ .

We prove that all wheels, necklaces, complete bipartite graphs $K_{r,t}$ with $r \neq t$ and almost all cacti graphs are of class 1^{\pm} . Moreover, we give sufficient and necessary conditions for a graph to be of class 2^{\pm} , i.e. we show that these graphs must have odd maximum degree and give examples of such graphs with arbitrary odd maximum degree bigger than 1.

 $^{^{*}}$ Krzysztof Turowski's research was funded in whole by Polish National Science Center 2020/39/D/ST6/00419 grant. For the purpose of Open Access, the author has applied a CC-BY public copyright licence to any Author Accepted Manuscript (AAM) version arising from this submission.

^{**}Corresponding author

¹E-mail: skalar@eti.pg.gda.pl

²E-mail: krzysztof.szymon.turowski@gmail.com

³E-mail: bart.wroblew@gmail.com

Keywords: signed graphs, edge coloring signed graphs, complete bipartite graphs, cacti

1. Introduction

In this paper we only consider simple, finite and undirected graphs. Graph G has a set of vertices V(G) and a set of edges E(G) with n(G), m(G) denoting their cardinalities, respectively. By $\deg_G(v)$ we denote degree of a vertex v in a graph G and by $\Delta(G)$ the maximum degree of all vertices of G. Incidence is a pair (v, e), where v is a vertex, e is an edge and v is one of the endpoints of e. Incidence (v, e) will be shortly denoted by v: e and the set of all graph's incidences will be denoted by I(G). All other definitions and symbols are consistent with those defined in Diestel [2].

Signed graphs were introduced in 1950s by Harary [3] as a generalization of simple graphs. Their main purpose was to better model social relations of disliking, indifference and liking. A signed graph is a pair (G, σ) where G is a graph and $\sigma: E(G) \to \{\pm 1\}$ is a function. G and σ are called the underlying graph of (G, σ) and the signature of (G, σ) , respectively. Edge $e \in E(G)$ will be called positive (respectively, negative) if and only if $\sigma(e) = 1$ (respectively, $\sigma(e) = -1$). Cycle in (G, σ) is called positive (respectively, negative) if product of signs of its edges is positive (respectively, negative). A signed graph with only positive cycles is balanced otherwise it is unbalanced.

Switching of a set $V' \subseteq V(G)$ of a signed graph (G, σ) is an operation resulting in a signed graph (G, σ') such that for every edge $uv \in E(G)$ we have

$$\sigma'(uv) = \begin{cases} -\sigma(uv), & \text{if exactly one of } u, v \text{ belongs to } V', \\ \sigma(uv), & \text{otherwise.} \end{cases}$$

For example, switching a vertex, i.e. switching a subset of vertices of cardinality one, negates the signs of its incident edges. If a signed graph (G, σ') can be obtained by switching some of the vertices of (G, σ) , we say that (G, σ') and (G, σ) are *switching equivalent*. It is well-known that switching equivalence is an equivalence relation in the set of all signed graphs with a fixed underlying graph.

In 2020, Behr [1] introduced a concept of edge coloring of signed graphs as a generalization of ordinary graph edge coloring. Let n be a positive integer and

$$M_n = \begin{cases} \{0, \pm 1, \dots, \pm k\}, & \text{if } n = 2k+1, \\ \{\pm 1, \dots, \pm k\}, & \text{if } n = 2k. \end{cases}$$

An *n*-edge-coloring of a signed graph (G, σ) is a function $f: I(G) \to M_n$ such that $f(u: uv) = -\sigma(uv)f(v: uv)$ for each edge $uv \in E(G)$ and $f(u: e_1) \neq f(u: e_2)$ for all edges $e_1 \neq e_2$ such that $u: e_1, u: e_2 \in I(G)$. By $\chi'(G, \sigma)$ we denote the chromatic index of a signed graph (G, σ) , i.e. the smallest *n* for which (G, σ) has an *n*-edge-coloring.

Behr [1] proved that every signed path can be colored using 2 colors and a signed cycle can be colored with 2 colors if and only if it's balanced. The main result of Behr's article [1] is the generalized Vizing's theorem. We will call it the Behr's theorem to emphasize the importance of Behr's achievement.

Theorem 1 (Behr [1]). $\Delta(G) \leq \chi'(G, \sigma) \leq \Delta(G) + 1$ for all signed graphs (G, σ).

In this paper we introduce two new classes of graphs, namely 1^{\pm} and 2^{\pm} , such that graph G is of class 1^{\pm} (respectively, 2^{\pm}) if and only if $\chi'(G, \sigma) = \Delta(G)$ (respectively, $\chi'(G, \sigma) = \Delta(G) + 1$) for all possible signatures σ . In Section 2 we relate these classes to previously known notions. We also show sufficient and necessary conditions for a regular signed graph (G, σ) to be $\Delta(G)$ -edgecolorable and use these results to obtain sufficient and necessary conditions for a regular signed graph to be of class 2^{\pm} . Next, we show that graphs of class 2^{\pm} must be of odd maximum degree and there are such graphs for any greater than 1 odd value of the degree. In Section 3 we prove that all cacti except cycles are of class 1^{\pm} . Section 4 deals with wheels—they are all of class 1^{\pm} . In Section 5 we show that all necklaces except cycles are of class 1^{\pm} . In Section 6 we deal with complete bipartite graphs by showing that graphs $K_{r,t}$ are of class 1^{\pm} for any $r \neq t$. The paper ends with some open problems and conjectures.

2. 1^{\pm} and 2^{\pm}

Behr [4] defined class ratio $\mathcal{C}(G)$ of graph G to be the number of possible signatures $\sigma: E(G) \to \{\pm 1\}$ such that the signed graph (G, σ) is $\Delta(G)$ -colorable, divided by the number of all possible signatures defined on E(G), i.e. $2^{m(G)}$. The class ratio is a rational number satisfying $0 \leq \mathcal{C}(G) \leq 1$ and, by previous definitions, G is of class 1^{\pm} (respectively, 2^{\pm}) if and only if $\mathcal{C}(G) = 1$ (respectively, $\mathcal{C}(G) = 0$). Clearly, there are graphs that are neither of class 1^{\pm} nor 2^{\pm} . The obvious example is a cycle—all signed cycles are 2-colorable if are balanced and require 3 colors otherwise. It is also easy to see that if G is of class 1^{\pm} (2^{\pm}), then it is of class 1 (2), where class 1 (2) is defined as all graphs G whose ordinary chromatic index equals $\Delta(G)$ ($\Delta(G) + 1$). This relation provides justification for our notation: class X^{\pm} is both a signed counterpart of class X and a subclass of X.

Recall that graph G is r-regular if and only if $\deg_G(v) = r$ for all $v \in V(G)$. Recall also that $M \subseteq E(G)$ is a *(perfect) matching* in G if every vertex of G is incident to at most (exactly) one edge of M.

Lemma 2. Let G be a k-regular graph and $\sigma: E(G) \to \{\pm 1\}$ be a signature. $\chi'(G, \sigma) = \Delta(G)$ if and only if one of the following conditions holds:

- 1. k = 2r where r is a positive integer and (G, σ) admits a decomposition into exactly r spanning edge disjoint 2-regular balanced subgraphs.
- 2. k = 2r + 1 where r is a positive integer and (G, σ) admits a decomposition into a perfect matching and exactly r spanning edge disjoint 2-regular balanced subgraphs.

Proof. We prove the equivalence by proving both implications.

(\Leftarrow) In the k = 2r case, each of the r spanning edge disjoint 2-regular balanced subgraphs can be colored using exactly 2 colors, so in total 2r colors are sufficient to color the whole graph. In the k = 2r + 1 case, the matching can be colored using color 0, each of the r spanning edge disjoint 2-regular balanced subgraphs can be colored using 2 colors, so in total 2r + 1 colors are also sufficient. Since $\Delta(G) = 2r$ in the first case and $\Delta(G) = 2r + 1$ in the second, we get $\chi'(G, \sigma) = \Delta(G)$.

 (\Rightarrow) We consider the two cases separately:

- G is a 2r-regular graph, so m(G) = r ⋅ n(G). Let c be an arbitrary Δ(G)-edge-coloring of (G, σ). For every pair of opposite colors used by c, edges colored with them form a subgraph with maximum degree not exceeding 2. Clearly, all such subgraphs are edge disjoint. Each of the subgraphs has at most n(G) edges. There are exactly r such subgraphs in G and G has r ⋅ n(G) edges, so every subgraph contains exactly n(G) edges. If a subgraph with maximum degree not exceeding 2 contains n(G) edges, it clearly is spanning and 2-regular. If it can be colored with two colors, it has to be balanced, too.
- (2) G is a (2r + 1)-regular graph, so $m(G) = r \cdot n(G) + \frac{1}{2}n(G)$. Let c be an arbitrary $\Delta(G)$ -edge-coloring of (G, σ) . The set of edges colored in c with color 0 must form a matching in G. Matching in G can have maximum size of $\frac{1}{2}n(G)$, so there have to be at least $r \cdot n(G)$ edges colored by c with colors different than 0. Such edges form r subgraphs with maximum degree not exceeding 2, each of them is colored with $\{-k, k\}$ for some $k \neq 0$. Clearly, they are edge disjoint subgraphs with degree not exceeding 2, so they have in total at most $r \cdot n(G)$ edges. Combining these observations, we conclude that these subgraphs have exactly $r \cdot n(G)$ edges. This implies that they are 2-regular, spanning and balanced since otherwise they could not be colored using only colors $\{-k, k\}$. This also implies that our matching has $\frac{1}{2}n(G)$ edges, which means it is perfect.

Let $\mathbf{1}_X$ denote the function $X \ni x \mapsto 1 \in \{\pm 1\}$.

Corollary 3. If G is a graph with even $\Delta(G)$, then G is not of class 2^{\pm} .

Proof. Every graph has a regular supergraph with the same maximum degree. Let H be a $\Delta(G)$ -regular supergraph of G. It follows from Petersen's 2-factor theorem [5] that H admits a decomposition into r edge disjoint 2-regular, spanning subgraphs. Since all edges in a signed graph $(H, \mathbf{1}_{E(H)})$ are positive, these subgraphs are balanced. Lemma 2 implies that $\chi'(H, \mathbf{1}_{E(H)}) = \Delta(H)$. Hence $\chi'(G, \mathbf{1}_{E(G)}) \leq \chi'(H, \mathbf{1}_{E(H)}) = \Delta(H) = \Delta(G)$, which shows that G is not of class 2^{\pm} .

Theorem 4. Let G be a graph. G is of class 2^{\pm} if and only if $\Delta(G)$ is odd and there is no matching M in G such that $\Delta(G \setminus M) < \Delta(G)$.

Proof. If follows from Corollary 3 that if graph is of class 2^{\pm} then its maximum degree must be odd.

(⇒) Let G be a graph with odd maximum degree. Suppose that there is a matching $M \subseteq E(G)$ such that $\Delta(G \setminus M) < \Delta(G)$. We will show that G is not of class 2^{\pm} . Clearly, $\Delta(G \setminus M)$ is even since $\Delta(G \setminus M) = \Delta(G) - 1$. The proof of Corollary 3 shows that a signed graph $(G \setminus M, \mathbf{1}_{E(G) \setminus M})$ is $\Delta(G \setminus M)$ -colorable. Since $\Delta(G \setminus M)$ is even, $\Delta(G \setminus M)$ -colorings of $G \setminus M$ does not use color 0. We can extend any of these colorings to a coloring of G by using color 0 on M, so $(G, \mathbf{1}_{E(G)})$ is $\Delta(G)$ -edge-colorable. This completes the proof.

 (\Leftarrow) Let us assume $\Delta(G)$ is odd, there is no matching $M \subseteq E(G)$ such that $\Delta(G \setminus M) < \Delta(G)$ and G is not of class 2^{\pm} . We show that this leads to a contradiction. There must exist a signature σ such that (G, σ) is $\Delta(G)$ -edge-colorable. Let c be an arbitrary $\Delta(G)$ -edge-coloring of (G, σ) . Edges colored in c with color 0 must form a matching M. Clearly, $\Delta(G \setminus M) = \Delta(G)$, so graph $(G \setminus M, \sigma|_{E(G) \setminus M})$ requires at least $\Delta(G)$ colors different than 0. It follows that (G, σ) cannot be colored using $\Delta(G)$ colors, so G is of class 2^{\pm} .

It's easy to observe that signed graphs with $\Delta = 1$ can be colored using color 0 regardless of their signature, so graphs with $\Delta = 1$ are of class 1^{\pm} .

Theorem 5. For every $k \ge 1$ there is a graph G of class 2^{\pm} such that $\Delta(G) = 2k + 1$.

Proof. We will describe the procedure of construction of a graph G such that $\Delta(G) = 2k + 1$ and G is of class 2^{\pm} . We start by constructing a complete bipartite graph H with parts $\{u_1, \ldots, u_{2k}\}$ and $\{v_1, \ldots, v_{2k}\}$. Next we create a graph H' by adding edges $u_{2i-1}u_{2i}$ for $1 \le i \le k$ to H, a new vertex v and edges $v_i v$ for $1 \le i \le 2k$. It is easy to see that

$$\deg_{H'}(w) = \begin{cases} 2k, & \text{if } v = w, \\ 2k+1, & \text{otherwise.} \end{cases}$$

Now we construct the graph G by creating two disjoint copies of H' and connecting their lowest degree vertices with path P_3 in such a way that these vertices are the endpoints of the path (see Figure 1 for an example). By v'_1 , v_c , v'_2 we denote vertices of that path and by v'_1v_c , $v_cv'_2$ —its edges. Let's observe that $\deg_G(v_c) = 2$ and other vertices of G have degree 2k + 1.

To complete the proof it suffices to show that G is of class 2^{\pm} . Clearly, $\Delta(G)$ is odd, so Theorem 4 tells us that we need to show that there is no matching $M \subseteq E(G)$ such that $\Delta(G \setminus M) < \Delta(G)$. Let's assume there is such a matching M in G. We show that it leads to a contradiction. Clearly, M must cover all the vertices of G other than v_c —it must cover all the vertices of both copies of H'. Let's observe that H' has an odd number of vertices, so there is no perfect matching in it. It means that in order to cover vertices of both copies of H', M must contain edges v'_1v_c and $v_cv'_2$. Then M contains two edges incident with v_c , so M is not a matching, a contradiction that completes the proof.




Figure 1: Example graph of class 2^{\pm} with $\Delta = 5$.

3. Cacti

A connected graph G is called *cactus* if and only if every edge of G belongs to at most one cycle. Clearly every tree is a cactus as it doesn't have cycles at all. It is a well known fact [6] that graph G is a cactus if and only if there exists a sequence of graphs G_1, \ldots, G_k called a *decomposition of* G, such that:

- 1. G_i is a cycle or a path P_2 for $1 \le i \le k$;
- 2. G_i has exactly one vertex in common with $G_1 \cup \ldots \cup G_{i-1}$ for $2 \le i \le k$;
- 3. $G = G_1 \cup \ldots \cup G_k$.

We prove that almost all signed cacti are of class 1^{\pm} .

Theorem 6. Let G be a cactus. If G is not a cycle, then G is of class 1^{\pm} .

Proof. Let $\sigma: E(G) \to \{\pm 1\}$ be a function. If $\Delta(G) \leq 2$, then G is a path, so our claim is obvious. Therefore we assume that $\Delta(G) \geq 3$ in the remainder of the proof.

Let G_1, \ldots, G_k be an arbitrary decomposition of G meeting the three above conditions. By G'_l we denote $G_1 \cup \ldots \cup G_l$. Clearly, $G'_k = G$. We will use induction on l to prove that $(G'_l, \sigma|_{E(G'_l)})$ is $\Delta(G)$ -edge-colorable. Let us observe that G'_1 is a cycle or path, so $(G'_1, \sigma|_{E(G'_1)})$ can be colored using $\Delta(G) \geq 3$ colors. Let's assume that $(G'_l, \sigma|_{E(G'_l)})$ is $\Delta(G)$ -edge-colorable for any l < x. We will show that $(G'_x, \sigma|_{E(G'_x)})$ can be colored using at most $\Delta(G)$ colors. Let us note that $G'_x = G'_{x-1} \cup G_x$ and $(G'_{x-1}, \sigma|_{E(G'_{x-1})})$ is $\Delta(G)$ -edge-colorable. By c we denote an arbitrary $\Delta(G)$ -edge-coloring of $(G'_{x-1}, \sigma|_{E(G'_{x-1})})$. It follows from the definition of the decomposition of G that G_x is either a path P_2 or a cycle and G_x has exactly one vertex in common with G'_{x-1} . Let us consider the two cases separately.

- 1. G_x is a path P_2 . By u, v we denote vertices of G_x and without loss of generality we assume $u \in V(G'_{x-1})$. Clearly, there are at most $\Delta(G) 1$ edges incident to u in G'_{x-1} , so there must be a color $\alpha \in M_{\Delta(G)}$ not used on any of these edges by c. Let's color u: uv with α and v: uv with $-\sigma(uv)\alpha$. That way $(G'_x, \sigma|_{E(G'_x)})$ can be colored using $\Delta(G)$ colors.
- 2. G_x is a cycle. By $u \in V(G_x)$ we denote a vertex such that $u \in V(G'_{x-1})$. By uv_1 , uv_2 we denote two different edges of G_x . Clearly, there are at




Figure 2: Three possible edge colorings of $(G_x, \sigma|_{E(G'_x)})$ considered in the case 2 of the proof of Theorem 6.

most $\Delta(G) - 2$ edges incident to u in G'_{x-1} , so there must be two different colors $\alpha, \beta \in M_{\Delta(G)}$ not used on any of these edges by c. Let us consider three possible subcases.

- (a) $\alpha = -\beta$ and $\Delta(G) = 3$. Since $\Delta(G) = 3$, $0 \in M_{\Delta(G)}$ and $\alpha \neq 0$, $\beta \neq 0$. By w we denote vertex in G_x such that w is adjacent to v_1 and $w \neq u$. Incidences $v_1 : v_1 w$ and $w : v_1 w$ of G'_x can be colored with color 0. Let us observe that other edges of G_x form a path, so their incidences can be colored with colors $\pm \alpha$ in the coloring of $(G'_x, \sigma|_{E(G'_x)})$.
- (b) $\alpha = -\beta$ and $\Delta(G) > 3$. Since $\Delta(G) > 3$, there must be colors γ , $-\gamma \in M_{\Delta(G)}$ such that $\gamma \neq 0, \gamma \neq \alpha, \gamma \neq \beta$. Let us observe that edges $v_1 u, uv_2$ form a path in G_x , so their incidences can be colored with colors $\pm \alpha$ in the coloring of $(G'_x, \sigma|_{E(G'_x)})$. Other edges of G_x also form a path—can be colored with colors $\pm \gamma$.
- (c) $\alpha \neq -\beta$. Without loss of generality let us assume $\beta \neq 0$. Incidence $u: uv_1$ can be colored with α in the coloring of $(G'_x, \sigma|_{E(G'_x)})$ and $v_1: uv_1$ —with $-\sigma(uv_1)\alpha$. Edges of G_x other than uv_1 form a path and can be colored with $\pm\beta$ in such a way that $u: uv_2$ gets color β .

The decomposition of an *n*-vertex cactus can be done in O(n) time. This implies that the coloring procedure being a part of the above proof is also linear.

4. Wheels

Wheel W_n is a graph on *n* vertices consisting of a cycle C_{n-1} , a vertex $v \notin V(C_{n-1})$ and edges between *v* and all the vertices of a cycle C_{n-1} .

Theorem 7. All wheels are of class 1^{\pm} .

Proof. We consider three cases separately:

1. n = 4. Wheel W_4 is a complete graph K_4 . It's easy to observe that W_4 has a decomposition into three perfect matchings and edges of any two of




Figure 3: The decomposition of W_7 into three paths. Distinct paths are marked with solid, dashed and dotted lines.

them create a spanning cycle. Let us consider an arbitrary signed graph (W_4, σ) and it's decomposition D into three perfect matchings. If there is an odd number of negative edges in (W_4, σ) , there must be a matching in D with an odd number of negative edges. Remaining edges create a balanced spanning cycle. In the opposite case—when there is an even number of negative edges. Remaining edges create a balanced spanning cycle. In both cases, (W_4, σ) , there must be a matching in D with an even number of negative edges. Remaining edges create a balanced spanning cycle. In both cases, (W_4, σ) has a decomposition into a perfect matching and a balanced spanning cycle. Obviously, W_4 is 3-regular, so it follows from Lemma 2 that (W_4, σ) is Δ -edge-colorable.

2. $n = 2k + 1, k \ge 2$. Let $V(W_n) = \{u, v_0, \ldots, v_{n-2}\}$ and $E(G) = \{uv_0, \ldots, uv_{n-2}\} \cup \{v_0v_1, v_1v_2, \ldots, v_{n-2}v_0\}$. For $0 \le i \le k-2$, let G_i be a subgraph of W_n , such that $V(G_i) = \{v_i, v_{i+1}, u, v_{i+k}, v_{i+k+1}\}$ and $E(G_i) = \{v_{i+1}v_i, v_iu, uv_{i+k}, v_{i+k}v_{i+k+1}\}$. Clearly, G_i is a path and for any different i_1, i_2 paths G_{i_1}, G_{i_2} are edge disjoint.

Let P be a subgraph of W_n , such that $V(P) = \{v_{k-1}, v_k, u, v_{n-2}, v_0\}$ and $E(P) = \{v_k v_{k-1}, v_{k-1}u, uv_{n-2}, v_{n-2}v_0\}$. Clearly, P is a path. We will show that paths P, G_i are edge disjoint for any $0 \le i \le k-2$. It's sufficient to show that $\{v_i u, uv_{i+k}\} \cap \{v_{k-1}u, uv_{n-2}\} = \emptyset$ and $\{v_{i+1}v_i, v_{i+k}v_{i+k+1}\} \cap \{v_k v_{k-1}, v_{n-2}v_0\} = \emptyset$. The first equality follows from the fact that i < k-1 < i+k < n-2. Let's assume that the second equality is false, so one of the following cases must hold:

- (a) $v_{i+1}v_i = v_k v_{k-1}$. Impossible because i < k-1 < k.
- (b) $v_{i+1}v_i = v_{n-2}v_0$. Impossible because i < i + 1 < n 2.
- (c) $v_{i+k}v_{i+k+1} = v_k v_{k-1}$. Impossible because k-1 < i+k < i+k+1.
- (d) $v_{i+k}v_{i+k+1} = v_{n-2}v_0$. Impossible because $i+k \neq 0$ and $i+k+1 \neq 0$.

Let's observe that $m(\bigcup_{i=0}^{k-2} G_i) + m(P) = 4(k-1) + 4 = 4k = m(W_n)$, so k-2

 $\bigcup_{i=0}^{k-2} G_i \cup P = W_n \text{ and } W_n \text{ has a decomposition into exactly } k = \Delta(W_n)/2$ paths. It follows that any signed graph with W_n as an underlying graph

can be colored using $\Delta(W_n)$ colors because edges of each path can be colored with exactly two colors.

3. $n = 2k, k \ge 3$. Let us consider wheel W_{n-1} . By u we denote the center vertex of W_{n-1} —the one with degree $\Delta(W_{n-1})$ and by v_1, v_2 we denote




Figure 4: The decomposition of W_8 into three paths and a single edge. Edge is marked with a bold line and distinct paths are marked with solid, dashed and dotted lines.

arbitrary vertices adjacent to u such that $v_1v_2 \in E(W_{n-1})$. It follows from the previous case that W_{n-1} admits a decomposition D into k-1paths. We construct graph W'_{n-1} from W_{n-1} by adding new vertex v and replacing edge v_1v_2 by path v_1, v, v_2 . It's easy to observe that W'_{n-1} also admits some decomposition D' into k-1 paths. It can be constructed directly from D—path containing edge v_1v_2 in D contains edges v_1v, vv_2 in D'. Clearly, graph W_n can be obtained from W'_{n-1} by adding edge vu. It follows that W_n admits a decomposition into k-1 paths and a single edge, so every signed graph with W_n as an underlying graph can be colored using $2k - 1 = \Delta(W_n)$ colors.

5. Necklaces

Necklace is a connected graph having a decomposition into $k \ge 2$ paths, in which all vertices except selected two—u, v—are different. Vertices u, v are starting and ending vertices of all paths. If k = 2, a necklace is a cycle, so it's possible that a signed graph on such necklace is negative and requires $\Delta + 1$ colors to be properly colored. We prove necklaces are Δ -colorable in all other cases.

Theorem 8. Let G be a necklace. If G is not a cycle, then G is of class 1^{\pm} .

Proof. G is not a cycle, so $\Delta(G) > 3$. By $D = G_1, \ldots, G_{\Delta(G)}$ we denote the decomposition of G into paths. By u, v we denote the starting and ending vertices of all the paths from D. Without loss of generality, we assume that $m(G_i) \leq m(G_{i+1})$ for $1 \leq i < \Delta(G)$. It follows that if there is a path of length 1 in D, it must be G_1 . By u_i^j we denote a vertex belonging to path G_i such that the distance between u_i^j and u in G_i is j. The definition of v_i^j is analogous. By $S = (G, \sigma)$ we denote an arbitrary signed graph with G being its underlying graph. We consider two cases separately:

1. $\Delta(G) = 2k+1, k \ge 1$. We will construct necklace S starting with necklace S_0 that contains three paths from $D-G_1, G_2, G_3$ and extending it with




Figure 5: Necklace S_0 considered in case 1 of the proof of Theorem 8. Edges colored with color 0 are marked with bold lines. Remaining edges span a path.

consecutive pairs of paths. By S_p we denote the necklace constructed from S_{p-1} by adding paths G_{2p+2} , G_{2p+3} . Clearly, $\Delta(S_0) = 3$. We color edges uu_2^1 , vv_3^1 with color 0. Let us observe that the remaining edges of S_0 span a path that contains following vertices: $u_2^1, \ldots, v_2^1, v, v_1^1, \ldots, u_1^1, u, u_3^1, \ldots, v_3^1$. It can be colored with colors ± 1 .

Necklace S_p is constructed from S_{p-1} by adding paths G_{2p+2} , G_{2p+3} . Clearly, $\Delta(S_p) = \Delta(S_{p-1}) + 2$, so there are two new colors $\pm \alpha$ available for coloring of S_p . Let us assume there is an edge uu_l^1 in S_{p-1} colored with color 0. We can color edge uu_{2p+2}^1 with color 0 and path u_l^1 , u, u_{2p+3}^1 , \ldots , v_{2p+3}^1 , v, v_{2p+2}^1 , \ldots , u_{2p+2}^1 with colors $\pm \alpha$. That way edge uu_l^1 , previously colored with 0, was recolored. Let us observe that this gives us a $\Delta(S_p)$ edge-coloring of S_p . Let us also observe that there is still an edge incident to u such that it's colored with color 0 in the coloring of S_p , so such an edge must exist in the colorings of all S_0, \ldots, S_p .

2. $\Delta(G) = 2k, k \geq 2$. We will construct necklace S starting with necklace S_0 that contains four paths from $D-G_1, \ldots, G_4$ and extending it with consecutive pairs of paths. By S_p we denote the necklace constructed from S_{p-1} by adding paths G_{2p+3}, G_{2p+4} . Clearly, $\Delta(S_0) = 4$. Let us observe that vertices $u_3^1, u, u_1^1, \ldots, v_1^1, v, v_2^1$ span a path in S_0 and its edges can be colored with colors ± 1 . We can assume that incidence $u: uu_3^1$ gets color 1. Let us observe that the remaining edges also span a path, end its edges can be colored with colors ± 2 .

Necklace S_p is constructed from S_{p-1} by adding paths G_{2p+3} , G_{2p+4} . Clearly, $\Delta(S_p) = \Delta(S_{p-1}) + 2$, so there are two new colors $\pm \alpha$ available for coloring of S_p . Let us assume there is an incidence $u: uu_l^1$ in S_{p-1} colored with color 1. We can color incidence $u: uu_{2p+3}^1$ with color 1 and $u_{2p+3}^1: uu_{2p+3}^2$ —with color $-\sigma(uu_{2p+3}^1)$. Path u_l^1 , $u, u_{2p+4}^1, \ldots, v_{2p+4}^1$, $v, v_{2p+3}^1, \ldots, u_{2p+3}^1$ can be colored with colors $\pm \alpha$. That way incidence $u: uu_l^1$, previously colored with 1, was recolored. Let us observe that this gives us a $\Delta(S_p)$ -edge-coloring of S_p . Let us also observe that there is




Figure 6: Necklace S_0 considered in case 2 of the proof of Theorem 8. Edges colored with colors ± 1 are marked with bold lines. Remaining edges are colored with colors ± 2 .

still an incidence incident to u such that it's colored with color 1 in the coloring of S_p , so such an incidence must exist in the colorings of all S_0 , ..., S_p .

6. Complete bipartite graphs

A graph is bipartite if and only if its vertices can be divided into two sets V_1 , V_2 such that each edge has endpoints in both sets. These sets are usually called *parts*. Graph is a complete bipartite graph if and only if it's bipartite and for every $u \in V_1$, $v \in V_2$ there is an edge uv.

Theorem 9. Let r, t be positive integers and G be a complete bipartite graph $K_{r,t}$. If $r \neq t$, then G is of class 1^{\pm} .

Proof. Without loss of generality, we assume that r < t. Let t = 2s + k, $s \in \mathbb{N}$ and $k \in \{0, 1\}$. Let $V(G) = \{u_0, \ldots, u_{r-1}\} \cup \{v_0, \ldots, v_{t-1}\}$ and $E(G) = \{u_i v_j : 0 \le i \le r-1, 0 \le j \le t-1\}.$

Let $0 \leq j \leq s - 1$. Let G_j be a subgraph of G such that $V(G_j) = \{u_0, \ldots, u_{r-1}\} \cup \{v_{2j}, \ldots, v_{(2j+r) \mod t}\}$ and $E(G_j) = \{u_i v_{(i+2j) \mod t}: 0 \leq i \leq r-1\} \cup \{u_i v_{(i+2j+1) \mod t}: 0 \leq i \leq r-1\}$. Let us observe that:

- $\deg_{G_i}(u_i) = 2$ for $i = 0, \ldots, r 1;$
- $\deg_{G_i}(v_{2j}) = \deg_{G_i}(v_{(2j+r) \mod t}) = 1;$
- $\deg_{G_i}(v_{(2j+i) \mod t}) = 2$ for $i = 1, \ldots, r-1$.




Figure 7: Paths G_0 (solid lines) and G_1 (dashed lines) created in the procedure of coloring of complete bipartite graph $K_{3,4}$ described in the proof of Theorem 9.

It follows from the definition of G_j and above observations that G_j is a path. We will show that any two distinct paths G_{j_1} , G_{j_2} are edge disjoint. Let us assume it's not true. Then one of the following cases must hold for some i_1 , $i_2 \in \{0, \ldots, r-1\}$.

- 1. $u_{i_1}v_{(i_1+2j_1) \mod t} = u_{i_2}v_{(i_2+2j_2) \mod t}$. Clearly, $u_{i_1} = u_{i_2}$, so $i_1 = i_2$. Moreover, $v_{(i_1+2j_1) \mod t} = v_{(i_2+2j_2) \mod t}$, so $2j_1 \equiv 2j_2 \pmod{t}$. It follows that $0 \equiv 2(j_1 - j_2) \pmod{t}$. Let us consider two cases separately:
 - (a) t = 2s + 1. Then $t \equiv 2s + 1 \pmod{t}$, so $0 \equiv 2s + 1 \pmod{t}$ and $2s \equiv -1 \pmod{t}$. We observe that if $0 \equiv 2(j_1 j_2) \pmod{t}$, then $0 \equiv 2s(j_1 j_2) \pmod{t}$, so $0 \equiv j_2 j_1 \pmod{t}$ and $j_1 \equiv j_2 \pmod{t}$. It follows that $j_1 = j_2$, a contradiction.
 - (b) t = 2s. If $0 \equiv 2(j_1 j_2) \pmod{t}$, then $0 \equiv 2(j_1 j_2) \pmod{2s}$ and $0 \equiv j_1 j_2 \pmod{s}$, so $j_1 \equiv j_2 \pmod{s}$. It follows that $j_1 = j_2$, a contradiction.
- 2. $u_{i_1}v_{(i_1+2j_1+1) \mod t} = u_{i_2}v_{(i_2+2j_2+1) \mod t}$. We observe that $i_1 = i_2$, so $2j_1 + 1 \equiv 2j_2 + 1 \pmod{t}$ and then $2(j_1 j_2) \equiv 0 \pmod{t}$. The remaining part of that case is identical to the previous case and leads to a contradiction.
- 3. Either $u_{i_1}v_{(i_1+2j_1) \mod t} = u_{i_2}v_{(i_2+2j_2+1) \mod t}$ or $u_{i_1}v_{(i_1+2j_1+1) \mod t} = u_{i_2}v_{(i_2+2j_2) \mod t}$. Without loss of generality, we assume that the first of two holds. Clearly, $i_1 = i_2$, so $2j_1 \equiv 2j_2+1 \pmod{t}$ and then $2(j_1-j_2) \equiv 1 \pmod{t}$. We consider two cases separately:
 - (a) t = 2s+1. Then $2s \equiv -1 \pmod{t}$. Let us observe that $2s(j_1-j_2) \equiv s \pmod{t}$, so $j_2 j_1 \equiv s \pmod{t}$. It's a contradiction because j_1 , $j_2 \in \{0, \ldots, s-1\}$.
 - (b) t = 2s. We observe that $2(j_1 j_2) 1 \equiv 0 \pmod{2s}$, so the odd number $2(j_1 j_2) 1$ is divisible by an even number 2s, a contradiction.

All of the cases lead to a contradiction, so paths G_{j_1} , G_{j_2} are edge disjoint. Clearly, $\bigcup_{i=0}^{s-1} G_j \subseteq G$. We will continue the proof separately for two cases:

1. t = 2s. Since G is a complete bipartite graph, m(G) = rt = 2rs. We observe that $m(G_j) = 2r$ for $0 \le j \le s - 1$. It follows that $m(\bigcup_{j=0}^{s-1} G_j) =$

$$\sum_{j=0}^{s-1} m(G_j) = 2rs, \text{ so } m(G) = m(\bigcup_{j=0}^{s-1} G_j). \text{ Since paths } G_j \text{ are edge disjoint,}$$

 $G = \bigcup_{j=0}^{s-1} G_j$, so G admits a decomposition into exactly s paths. Let us observe that an arbitrary signed graph with G being its underlying graph

can be colored with $2s = \Delta(G)$ colors, since each path can be colored with just two colors. It completes a proof for this case.

2. t = 2s + 1. It's clear that m(G) = rt = r(2s + 1) and $m(\bigcup_{j=0}^{s-1} G_j) = 2rs$. Let M be a subgraph of G such that $V(M) = \{u_0, \ldots, u_{r-1}\} \cup \{v_{(i-1) \mod t} : 0 \le i \le r-1\}$ and $E(M) = \{u_i v_{(i-1) \mod t} : 0 \le i \le r-1\}$. It's easy to observe that all the vertices of M have degree equal to 1, so M is a matching.

We will show that $E(M) \cap E(\bigcup_{j=0}^{s-1} G_j) = \emptyset$. We assume that it's not true. Then for some j and $i_1, i_2 \in \{0, \ldots, r-1\}$ one of the following cases must hold:

- (a) $u_{i_1}v_{(i_1-1) \mod t} = u_{i_2}v_{(i_2+2j) \mod t}$. Clearly, $i_1 = i_2$. We observe that $i_1 1 \equiv i_2 + 2j \pmod{t}$, so $2j + 1 \equiv 0 \pmod{t}$ and then $2js + s \equiv 0 \pmod{t}$. Since t = 2s + 1, $2s \equiv -1 \pmod{t}$, so $j \equiv s \pmod{t}$. It's a contradiction because $j \in \{0, \ldots, s-1\}$.
- (b) $u_{i_1}v_{(i_1-1) \mod t} = u_{i_2}v_{(i_2+2j+1) \mod t}$. It holds that $i_1 1 \equiv i_2 + 2j + 1 \pmod{t}$, so $2j \equiv -2 \pmod{t}$ and then $2js \equiv -2s \pmod{t}$. Since $t = 2s + 1, j + 1 \equiv 0 \pmod{t}$. It's a contradiction because t = 2s + 1 and $j \in \{0, \ldots, s 1\}$.

All of the cases lead to a contradiction, so $E(M) \cap E(\bigcup_{j=0}^{s-1} G_j) = \emptyset$.

Since m(M) = r, $G = E(M) \cup E(\bigcup_{j=0}^{s-1} G_j)$, G admits a decomposition into exactly s paths and one matching. Let us observe that an arbitrary signed graph with G being its underlying graph can be colored with $2s+1 = \Delta(G)$ colors, since each path can be colored with two non-zero colors and a matching can be colored with color 0.

We briefly consider complete bipartite graphs with equal parts. Complete bipartite graph $K_{r,r}$ is an r-regular graph, so it follows from Lemma 2 that if r is

even and $K_{r,r}$ has an odd number of negative edges, it cannot be colored using Δ colors. We hypothesize that for all other cases, complete bipartite graphs with equal parts are Δ -colorable. If the conjecture is true, graphs with odd r must have a perfect matching M such that $K_{r,r} \setminus M$ have an even number of negative edges and have a decomposition into (r-1)/2 spanning 2-regular subgraphs with positive cycles being their components.

Conjecture 10. Let $r \in \mathbf{N}$ and S be a signed complete bipartite graph $K_{r,r}$. $\chi'(S) = \Delta(S)$ if one of the following conditions holds:

- 1. r is odd;
- 2. r is even and there is an even number of negative edges in S.

References

- R. Behr, Edge coloring signed graphs, Discrete Mathematics 343 (2) (2020) 111654.
- [2] R. Diestel, Graph Theory, Springer-Verlag Berlin Heidelberg, 2017.
- [3] F. Harary, On the notion of balance of a signed graph, Michigan Mathematical Journal 2 (1953) 143–146.
- [4] R. Behr, Edge coloring signed graphs (2018).
 URL https://arxiv.org/abs/1807.11465
- [5] H. M. Mulder, Julius Petersen's theory of regular graphs, Discrete Mathematics 100 (1-3) (1992) 157–175.
- [6] R. Janczewski, K. Turowski, An O(n log n) algorithm for finding edge span of cacti, Journal of Combinatorial Optimization 31 (2016) 1373–1382.