

ALMA Confirmation of an Obscured Hyperluminous Radio-Loud AGN at $z = 6.853$ Associated with a Dusty Starburst in the 1.5 deg^2 COSMOS Field

Ryan Endsley^{1*}, Daniel P. Stark¹, Jianwei Lyu¹, Feige Wang¹, Jinyi Yang^{1,2}, Xiaohui Fan¹, Renske Smit³, Rychard Bouwens⁴, Kevin Hainline¹, Sander Schouws⁴

¹*Steward Observatory, University of Arizona, 933 N Cherry Ave, Tucson, AZ 85721 USA*

²*Strittmatter Fellow*

³*Astrophysics Research Institute, Liverpool John Moores University, 146 Brownlow Hill, Liverpool L3 5RF, United Kingdom*

⁴*Leiden Observatory, Leiden University, NL-2300 RA Leiden, Netherlands*

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

We present band 6 ALMA observations of a heavily-obscured radio-loud ($L_{1.4 \text{ GHz}} = 10^{25.4} \text{ W Hz}^{-1}$) AGN candidate at $z_{\text{phot}} = 6.83 \pm 0.06$ found in the 1.5 deg^2 COSMOS field. The ALMA data reveal detections of exceptionally strong [CII] $158\mu\text{m}$ ($z_{\text{[CII]}} = 6.8532$) and underlying dust continuum emission from this object (COS-87259), where the [CII] line luminosity, line width, and $158\mu\text{m}$ continuum luminosity are comparable to that seen from $z \sim 7$ sub-mm galaxies and quasar hosts. The $158\mu\text{m}$ continuum detection suggests a total infrared luminosity of $9 \times 10^{12} L_{\odot}$ with corresponding very large obscured star formation rate ($1300 M_{\odot}/\text{yr}$) and dust mass ($2 \times 10^9 M_{\odot}$). The strong break seen between the VIRCcam and IRAC photometry perhaps suggests that COS-87259 is an extremely massive reionization-era galaxy with $M_{*} \approx 1.7 \times 10^{11} M_{\odot}$. Moreover, the MIPS, PACS, and SPIRE detections imply that this object harbors an AGN that is heavily obscured ($\tau_{9.7\mu\text{m}} = 2.3$) with a bolometric luminosity of approximately $5 \times 10^{13} L_{\odot}$. Such a very high AGN luminosity suggests this object is powered by an $\approx 1.6 \times 10^9 M_{\odot}$ black hole if accreting near the Eddington limit, and is effectively a highly-obscured version of an extremely UV-luminous ($M_{1450} \approx -27.3$) $z \sim 7$ quasar. Notably, these $z \sim 7$ quasars are an exceedingly rare population ($\sim 0.001 \text{ deg}^{-2}$) while COS-87259 was identified over a relatively small field. Future very wide-area surveys with, e.g., *Roman* and *Euclid* have the potential to identify many more extremely red yet UV-bright $z \gtrsim 7$ objects similar to COS-87259, providing richer insight into the occurrence of intense obscured star formation and supermassive black hole growth among this population.

Key words: galaxies: high-redshift – quasars: supermassive black holes – submillimetre: galaxies – dark ages, reionization, first stars – galaxies: evolution

1 INTRODUCTION

For much of the past two decades, deep imaging surveys with the Hubble Space Telescope (*HST*) have served as key windows on galaxies present in the first billion years of cosmic history (e.g. [Giavalisco et al. 2004](#); [Bouwens et al. 2011](#); [Grogin et al. 2011](#); [Koekemoer et al. 2011](#); [Trenti et al. 2011](#); [Ellis et al. 2013](#); [Illingworth et al. 2013](#); [Lotz et al. 2017](#)). To date, these *HST* surveys have enabled the identification of over 1000 Lyman-break galaxies at $z \gtrsim 6$ (e.g. [McLure et al. 2013](#); [Finkelstein et al. 2015](#); [Oesch et al. 2018](#); [Ishigaki et al. 2018](#); [Bouwens et al. 2021](#)), the bulk of which are faint ($M_{\text{UV}} > -20$) star-forming systems thought largely responsible for cosmic reionization ([Bouwens et al. 2015](#); [Finkelstein et al. 2019](#); [Robertson 2022](#)). Overlapping deep rest-optical imaging with *Spitzer*/IRAC has shown that these galaxies tend to possess low stellar masses ($M_{*} < 10^9 M_{\odot}$) and strong rest-optical nebular line emission consistent with young ages (e.g. [Labbé et al. 2013](#); [Stark et al. 2013](#); [Smit et al. 2014, 2015](#); [Song et al. 2016](#); [De Barros et al. 2019](#); [Endsley et al. 2021a](#);

[Stefanon et al. 2021](#)). However, while these *HST* surveys can push to extremely deep depths ($J \sim 29$), they tend to cover relatively small areas ($\lesssim 0.01 \text{ deg}^2$) which do not statistically sample $z \gtrsim 6$ galaxies that are sufficiently bright for detailed characterization with existing facilities ($J \lesssim 25$). Consequently, our understanding of very early galaxies has long been primarily limited to their rest-UV+optical photometric properties.

Within the past few years, several deep wide-area ($> \text{deg}^2$) optical+near-infrared imaging surveys have reached (near) completion ([Lawrence et al. 2007](#); [McCracken et al. 2012](#); [Jarvis et al. 2013](#); [Inoue et al. 2020](#); [Aihara et al. 2022](#)), providing large samples of UV-bright ($-23 \lesssim M_{\text{UV}} \lesssim -21$) Lyman-break $z \gtrsim 7$ galaxies ([Bowler et al. 2017, 2020](#); [Stefanon et al. 2019](#); [Endsley et al. 2021a](#); [Bouwens et al. 2022](#); [Harikane et al. 2022](#); [Schouws et al. 2022a](#)). This has in turn led to substantial improvements in our understanding of the UV-luminous reionization-era galaxy population. Rest-UV spectroscopic follow-up has revealed significant Lyman-alpha emission from several of these systems ([Ono et al. 2012](#); [Furusawa et al. 2016](#); [Endsley et al. 2021b, 2022b](#); [Valentino et al. 2022](#)), while (in some cases) simultaneously uncovering strong large-scale over-

* E-mail: rendsley@email.arizona.edu

densities likely tracing very large ionized bubbles (Endsley & Stark 2022). ALMA follow-up has also been highly successful in detecting far-IR cooling lines ([CII]158 μm and [OIII]88 μm) from several UV-bright $z \gtrsim 7$ galaxies (Smit et al. 2018; Hashimoto et al. 2019; Bouwens et al. 2022; Bowler et al. 2022; Schouws et al. 2022a), with a substantial fraction additionally exhibiting strong far-IR continuum emission indicating substantial obscured star formation activity (Hashimoto et al. 2019; Inami et al. 2022; Schouws et al. 2022b).

Alongside efforts to study Lyman-break $z \gtrsim 7$ galaxies, there has also been a dedicated push to identify and subsequently characterize UV-luminous quasars as well as IR-luminous sub-millimeter galaxies in this epoch. The 2500 deg² South Pole Telescope Survey (Everett et al. 2020) has resulted in the discovery of a sub-mm system at $z = 6.900$ (SPT-0311; Strandet et al. 2017), the only such source currently known at $z > 6.5$. ALMA follow-up revealed that this object is actually a merger of two galaxies exhibiting intense dust-obscured star formation activity ($\approx 3500 M_{\odot}/\text{yr}$ combined) that likely traces one of the most massive halos in the reionization era ($\sim 3 \times 10^{12} M_{\odot}$; Marrone et al. 2018; Spilker et al. 2022). Thanks to a rich collection of $\gtrsim 1000$ deg² optical+near-IR surveys, there are now more than 40 known extremely UV-luminous ($-28 \leq M_{\text{UV}} \leq -25$) quasars at $z > 6.5$ (Mortlock et al. 2011; Venemans et al. 2013, 2015; Mazzucchelli et al. 2017; Reed et al. 2017; Wang et al. 2017, 2018, 2019, 2021; Bañados et al. 2018, 2021; Matsuoka et al. 2018, 2019, 2022; Reed et al. 2019; Yang et al. 2019a, 2020b, 2021). These unobscured active galactic nuclei (AGN) are powered by the earliest known supermassive black holes with masses of $M_{\text{BH}} \approx (0.3 - 3) \times 10^9 M_{\odot}$ accreting near the Eddington limit (e.g. Yang et al. 2021). Far-IR follow-up of these quasars has provided a window into their host galaxies, revealing that several of these systems show very bright [CII], CO, and dust continuum emission similar to sub-mm galaxies (e.g. Venemans et al. 2016, 2017a,b, 2020; Decarli et al. 2018, 2022; Yang et al. 2019b; Pensabene et al. 2021). One population that would help bridge the gap between known quasar and sub-mm galaxies at $z \gtrsim 7$ are heavily-obscured AGN. However, even though these obscured AGN are expected to be more abundant than their unobscured counterparts (i.e. quasars) in the early Universe (Vito et al. 2018; Davies et al. 2019; Trebitsch et al. 2019; Ni et al. 2020; Lupi et al. 2022), to date no such object has been confirmed at $z > 6.5$.

We previously reported the photometric identification of a heavily-obscured, hyperluminous AGN candidate at $z \approx 6.83$ located in the 1.5 deg² COSMOS field (Endsley et al. 2022a, hereafter E22a). This source, COS-87259, was originally identified as a Lyman-break $z \approx 6.6 - 6.9$ galaxy using deep optical+near-IR imaging (Endsley et al. 2021a), and was later found to exhibit a sharp spectral discontinuity between two intermediate/narrow-bands at 9450 Å and 9700 Å suggesting $z_{\text{phot}} = 6.83 \pm 0.06$ (E22a). While similar in UV luminosity to many other wide-area selected Lyman-break $z \sim 7$ galaxies ($J = 25.0$ AB; $M_{\text{UV}} = -21.7$), COS-87259 was found to show both an exceptionally red rest-UV slope ($\beta = -0.6$) and a prominent Balmer break in the *Spitzer*/IRAC bands, implying unusually strong dust attenuation as well as a relatively old, massive stellar population (Endsley et al. 2021a). Subsequent investigations revealed that this galaxy is also detected in *Spitzer*/MIPS, *Herschel*/PACS, *Herschel*/SPIRE, JCMT/SCUBA-2, VLA, LOFAR, and MeerKAT bands (E22a). These data imply that, if at $z \approx 6.8$, COS-87259 is undergoing rapid, highly-obscured star formation activity and harbors a heavily-obscured hyperluminous radio-loud ($L_{1.4 \text{ GHz}} = 10^{25.4} \text{ W Hz}^{-1}$) AGN (E22a).

Here, we report the ALMA spectroscopic confirmation of COS-87259 as well as detections of exceptionally strong [CII]158 μm and

underlying dust continuum emission from this system. We describe the ALMA observations in §2 and then report the [CII] and dust continuum properties of COS-87259 in §3 comparing to other known objects at $z > 6.5$. We then discuss the physical properties of COS-87259 in §4. Our conclusions are summarized in §5, followed by a brief outlook on the potential of future *Roman* and *Euclid* surveys to identify many more extremely red, UV-bright $z \gtrsim 7$ galaxies similar to COS-87259. Throughout this paper, we quote magnitudes in the AB system (Oke & Gunn 1983), adopt a flat Λ CDM cosmology with parameters $h = 0.7$, $\Omega_{\text{M}} = 0.3$, and $\Omega_{\Lambda} = 0.7$, and report 68% confidence interval uncertainties unless otherwise specified.

2 OBSERVATIONS

ALMA observations were conducted targeting the [CII]158 μm emission line from COS-87259 on January 6, 2022 (project 2021.1.01311.S). Given the precise photometric redshift estimate of COS-87259 ($z_{\text{phot}} = 6.83 \pm 0.06$; E22a), we used the band 6 receiver to target a frequency range of 239.42–244.79 GHz corresponding to $z_{[\text{CII}]} = 6.76 - 6.94$. During the ALMA observations, a total of 45 antennae were used in a configuration with baseline lengths spanning 15–977 meters. The total integration time on COS-87259 was 33 minutes with an average precipitable water vapor level of 1.1 mm. The source J0948+0022 was used for calibration. We use the reduced data supplied by the observatory which were processed using CASA (McMullin et al. 2007) v.6.2.1.7 with pipeline v.2021.2.0.128. The resulting naturally-weighted continuum-subtracted spectra have an rms sensitivity of $\approx 440 \mu\text{Jy}/\text{beam}$ in 9.7 km/s channels, with a beam size of 0.49 arcsec by 0.37 arcsec and a position angle of -76.6° . The output continuum image ($\nu = 242.14$ GHz) has very similar beam properties with a size of 0.49 arcsec by 0.35 arcsec and a position angle of -75.1° , and reaches an rms sensitivity of 22 $\mu\text{Jy}/\text{beam}$.

3 RESULTS

In this section, we begin by reporting the [CII]158 μm and underlying dust continuum data for COS-87259 (§3.1), summarizing the measured and derived properties in Table 1. We then compare the observed far-IR properties of COS-87259 to other known systems at $z > 6.5$ (§3.2). Finally, we quantify the relative positions of the far-IR, rest-UV, and radio emission, and subsequently investigate whether there is any evidence that source confusion is significantly impacting the mid-IR detections of COS-87259 (§3.3).

3.1 [CII]158 μm and Dust Continuum Emission

The primary goal of our ALMA observations is to test the high-redshift ($z \approx 6.83$) nature of COS-87259, as implied by the optical through sub-mm photometric SED (E22a). We visually inspect the spectral data cubes, finding a clear emission line detection coincident with the near-IR position of COS-87259. To generate a 1D spectrum of this emission line, we create a moment-0 map for the extraction. We determine the frequency range over which to make the moment-0 map by first extracting the (continuum-subtracted) spectra over a single beam centered on the rest-UV emission of COS-87259 and subsequently fitting a Gaussian profile to determine the line FWHM and central frequency. Following Schouws et al. (2022a), a moment-0 map is then generated by summing all channels that fall within $\pm \text{FWHM}$ determined above, and we next extract the spectrum within the 3σ contours and refit a Gaussian profile to determine the line




Figure 1. (a) ALMA [CII] $158\mu\text{m}$ spectrum of COS-87259 (after continuum subtraction) extracted within the 3σ contours of the moment-0 map. The spectrum is binned to half the original resolution for clarity and the red line shows the Gaussian fit. (b) ALMA [CII] (blue), $158\mu\text{m}$ dust continuum (red), and VLA 3 GHz (green) 2, 3, and 4σ contour maps of COS-87259 overlaid on the stacked rest-UV image. The ALMA and VLA beams are shown in the bottom corners with their respective colors. The 3 GHz detection peak (green cross) is clearly offset (≈ 6 kpc projected) from the [CII] and dust continuum centroids (blue and red crosses), as expected if the radio emission is powered by an extended relativistic jet from the AGN. The centroid of the rest-UV emission (black cross) is also slightly offset from that of the far-IR (≈ 1.5 kpc), consistent with strong dust opacity variations across the galaxy.

FWHM and central frequency. This process is performed iteratively until the central frequency and FWHM converge (within a few iterations) which we then use to create the final moment-0 map and 1D extracted spectrum.

From a Gaussian fit to the final 1D spectrum (Fig. 1 a), we measure a $\text{FWHM} = 652 \pm 17 \text{ km s}^{-1}$ and a central frequency of 242.009 ± 0.007 GHz. This central frequency corresponds to a [CII] redshift of $z_{\text{[CII]}} = 6.8532 \pm 0.0002$, in excellent agreement with the precise pho-

Table 1. Measured and Derived Properties of COS-87259. The reported [CII] and dust continuum sizes are the deconvolved values.

Rest-UV and radio properties (see E22a)	
UV right ascension	09:58:58.27
UV declination	+01:39:20.2
z_{phot}	6.83 ± 0.06
Rest-UV slope	-0.59 ± 0.27
M_{1450}	-21.7 ± 0.1
1.32–3 GHz radio slope	$-1.57^{+0.22}_{-0.21}$
0.144–1.32 GHz radio slope	$-0.86^{+0.22}_{-0.16}$
1.4 GHz luminosity	$(2.5 \pm 0.5) \times 10^{25} \text{ W Hz}^{-1}$
ALMA results	
$z_{\text{[CII]}}$	6.8532 ± 0.0002
[CII] FWHM	$652 \pm 17 \text{ km s}^{-1}$
[CII] size (core)	$(1.9 \pm 0.4 \text{ kpc}) \times (< 1.0 \text{ kpc}^a)$
[CII] size (extended)	$(9.6 \pm 1.1 \text{ kpc}) \times (4.9 \pm 0.6 \text{ kpc})$
$F_{\text{[CII]}}$ (core)	$1.97 \pm 0.16 \text{ Jy km s}^{-1}$
$F_{\text{[CII]}}$ (extended)	$5.76 \pm 0.64 \text{ Jy km s}^{-1}$
Total [CII] luminosity	$(8.82 \pm 0.75) \times 10^9 L_{\odot}$
Dust continuum size (core)	$(1.4 \pm 0.2 \text{ kpc}) \times (1.0 \pm 0.1 \text{ kpc})$
Dust continuum size (extended)	$(7.8 \pm 10 \text{ kpc}) \times (3.4 \pm 0.5 \text{ kpc})$
$F_{242 \text{ GHz}}$ (core)	$1.48 \pm 0.05 \text{ mJy}$
$F_{242 \text{ GHz}}$ (extended)	$1.23 \pm 0.15 \text{ mJy}$
Total $F_{242 \text{ GHz}}$	$2.71 \pm 0.16 \text{ mJy}$
$L_{\text{IR,SF}}^b$	$(8.7 \pm 0.5) \times 10^{12} L_{\odot}$
M_{dust}^b	$(1.9 \pm 0.1) \times 10^9 M_{\odot}$
SFR_{IR}^c	$1300 \pm 70 M_{\odot} \text{ yr}^{-1}$
$\text{SFR}_{\text{[CII]}}^d$	$2970 \pm 250 M_{\odot} \text{ yr}^{-1}$
SED fitting results	
$L_{\text{IR,AGN}}$	$(2.5 \pm 0.2) \times 10^{13} L_{\odot}$
$\tau_{9.7\mu\text{m}}$	2.3 ± 0.1
$L_{\text{bol,AGN}}$	$(5.1 \pm 0.5) \times 10^{13} L_{\odot}$
$M_{1450, \text{unobscured}}$	-27.3 ± 0.1
$M_{\text{BH}} \lambda_{\text{Edd}}$	$(1.6 \pm 0.2) \times 10^9 M_{\odot}$
$L_{\text{IR,SF}}$	$(9.1 \pm 1.1) \times 10^{12} L_{\odot}$
SFR	$1610 \pm 190 M_{\odot} \text{ yr}^{-1}$
M_*	$(1.7 \pm 0.7) \times 10^{11} M_{\odot}$

^a 2σ upper limit on minor axis size from IMFIT.

^b Computed from the $158\mu\text{m}$ dust continuum measurement adopting $T_{\text{dust}} = 47 \text{ K}$ and $\beta_{\text{dust}} = 1.6$.

^c Converted from $L_{\text{IR,SF}}$ using the relation of Murphy et al. (2011).

^d Converted from $L_{\text{[CII]}}$ using the relation of Herrera-Camus et al. (2015).

tometric redshift of $z_{\text{phot}} = 6.83 \pm 0.06$ set by the sharp spectral break seen between the Subaru/Hyper Suprime-Cam intermediate/narrowbands at 9450 \AA and 9700 \AA (E22a). We therefore conclude that the ALMA-detected emission feature seen from COS-87259 is the [CII] $158\mu\text{m}$ line.

The [CII] emission of COS-87259 is spatially resolved with our moderate-resolution (beam ≈ 0.4 arcsec) ALMA data (Fig. 1 b). To determine the flux and size of the [CII] emission, we use the CASA task IMFIT which fits one or more 2D Gaussians to the moment-0 map. We find that a single elliptical Gaussian poorly reproduces the observed [CII] surface brightness profile, while a two-component model provides a satisfactory match to the data (Fig. 2). This two-component fit suggests the presence of a compact (≈ 2 kpc) [CII]-emitting core in addition to a more extended (≈ 10 kpc) region of emission, consis-




Figure 2. Illustration of the 2D Gaussian fits to the [CII] moment 0 map (top) and the $158\mu\text{m}$ dust continuum map (bottom). The data are shown in the leftmost panels, with the best-fitting single elliptical Gaussian models shown in the second panels from left. These single Gaussian fits provide a poor match to the measured surface brightness profiles of both the [CII] and dust continuum data, as can be seen by the strong residuals in the middle panels. The data are much better fit as the superposition of two elliptical Gaussians as illustrated in the two rightmost panels. Each panel is 7.3 arcsec (39 kpc) on a side.

tent with the [CII] surface brightness profiles of $z > 6$ quasar hosts (Novak et al. 2020). The extended [CII] component of COS-87259 has a deconvolved size of $(1.82 \pm 0.21 \text{ arcsec}) \times (0.93 \pm 0.11 \text{ arcsec})$ and an integrated flux of $5.76 \pm 0.64 \text{ Jy km s}^{-1}$. At the [CII] redshift of COS-87259 ($z_{[\text{CII}]} = 6.853$), these values translate to a physical size and [CII] line luminosity of $(9.6 \pm 1.1 \text{ kpc}) \times (4.9 \pm 0.6 \text{ kpc})$ and $(6.57 \pm 0.73) \times 10^9 L_{\odot}$, respectively. The compact component is only resolved along the major axis with an estimated deconvolved size of $0.35 \pm 0.07 \text{ arcsec}$ ($1.9 \pm 0.4 \text{ kpc}$) and its integrated flux is found to be $1.97 \pm 0.16 \text{ Jy km s}^{-1}$, which corresponds to a luminosity of $(2.25 \pm 0.18) \times 10^9 L_{\odot}$. The two-component `IMFIT` output therefore implies a total [CII] luminosity of $(8.82 \pm 0.75) \times 10^9 L_{\odot}$, the bulk of which is originating from the extended component ($75 \pm 10\%$).

The [CII] emission line is frequently used as a valuable tracer of the ISM kinematics for reionization-era galaxies (e.g. Smit et al. 2018; Hashimoto et al. 2019; Matthee et al. 2019; Neeleman et al. 2019). We begin investigating the kinematics of COS-87259 by generating maps of the [CII] mean velocity and velocity dispersion fields using the code `QUBEFIT` (Neeleman et al. 2021). When running `QUBEFIT`, we apply a 1σ clipping as a compromise between mitigating the impact of noise on the resulting moment maps while preserving true signal (see discussion in Neeleman et al. 2021). We have verified that our conclusions do not significantly change if we instead adopt a clipping threshold in the range 0.5 – 2σ . A prominent gradient is found in the mean velocity field with a maximum difference of $\approx 360 \text{ km s}^{-1}$ between spaxels in the northeast and southwest corners (Fig. 3). The velocity dispersion map also shows that the outermost spaxels tend to have a larger fitted FWHM (~ 600 – 900 km s^{-1}) than those located near the peak of emission ($\approx 550 \text{ km s}^{-1}$; see Fig. 3). This perhaps suggests the presence of a broader [CII] wing in COS-87259 as reported in other $z > 6$ AGN hosts (Maiolino et al. 2012; Izumi et al. 2021) and stacked spectra of $z = 4$ – 6 galaxies (Ginolfi et al. 2020), implying significant outflowing gas. However, because the

[CII] FWHM only varies by a factor of ≈ 1.5 across COS-87259 (c.f. the factor of 4 in Izumi et al. 2021), it is much more challenging to determine if a broad wing is indeed present in this system with existing moderate-resolution data. Upon extracting the [CII] spectrum across the outermost spaxels in the moment-0 footprint, we find that the resulting profile has a slightly larger fitted FWHM ($690^{+60}_{-50} \text{ km/s}$), though consistent with the full profile shown in Fig. 1a within uncertainties. We also extract position-velocity information along the major axis (as determined by a single 2D Gaussian fit) and find no clear evidence of multiple distinct components separated in velocity space (see Fig. A1). Higher-resolution [CII] data will be required to better assess whether the kinematics of COS-87259 are more consistent with a rotating disk or a disturbed, possibly merging system.

Strong dust continuum emission is also detected from COS-87259 (Fig. 1b). Similar to the [CII] emission, we find that the dust continuum surface brightness profile is much better fit by the superposition of two elliptical Gaussians as opposed to a single Gaussian (Fig. 2). With `IMFIT`, we find a deconvolved angular size of $(1.47 \pm 0.19 \text{ arcsec}) \times (0.65 \pm 0.10 \text{ arcsec})$ and $(0.26 \pm 0.03 \text{ arcsec}) \times (0.18 \pm 0.02 \text{ arcsec})$ for the more extended and compact components, respectively. These scales correspond to physical sizes of $(7.8 \pm 1.0 \text{ kpc}) \times (3.4 \pm 0.5 \text{ kpc})$ and $(1.4 \pm 0.2 \text{ kpc}) \times (1.0 \pm 0.1 \text{ kpc})$, respectively, which are systematically ≈ 20 – 30% smaller than that found for the [CII] emission. The integrated flux densities of the more compact and extended components are nearly equal at $1.48 \pm 0.05 \text{ mJy}$ and $1.23 \pm 0.15 \text{ mJy}$, respectively, yielding a total $158\mu\text{m}$ dust continuum flux density of $2.71 \pm 0.16 \text{ mJy}$ for COS-87259.

Given the highly luminous nature of COS-87259, we investigate whether this system may be significantly magnified by nearby foreground objects. We consider all foreground objects within 10 arcsec of COS-87259 identified in the COSMOS2020 catalog (Weaver et al. 2022), assuming each is a singular isothermal sphere with photometric redshift and stellar mass taken from that same catalog. The




Figure 3. Maps of the [CII] mean velocity (top) and velocity dispersion (bottom) fields of COS-87259 derived using QUBEFIT. The footprint of each map is confined to pixels with >3 sigma [CII] flux and the [CII] beam (≈ 0.4 arcsec) is shown in the bottom left corner of each panel. A prominent gradient is seen in the mean velocity field and there is evidence of larger velocity dispersion towards the outer edges of the [CII] emission.

velocity dispersion of each neighboring source is computed in two independent ways. First, we assume that each galaxy occupies a virialized halo with $200\times$ the mean density of the Universe at its assumed redshift and mass $100\times$ that of the stellar mass. Our second approach follows that of Mason et al. (2015) where we adopt the relation between velocity dispersion and stellar mass from Auger et al. (2010). Both methods to estimate velocity dispersions suggest that COS-87259 is only weakly lensed by neighboring foreground sources, with a small combined magnification factor of $\mu_{\text{tot}} \approx 1.15 - 1.25$. Because this weak magnification does not significantly impact our main conclusions, we choose not to correct for it when reporting the measured and inferred properties of COS-87259.

3.2 Comparison of Far-IR Properties to Other $z > 6.5$ Sources

We compare the observed far-infrared properties ([CII] luminosity, [CII] FWHM, and monochromatic $158\mu\text{m}$ dust continuum luminosity) of COS-87259 to other known objects in the reionization-era. From the literature, we compile such measured properties from




Figure 4. Comparison of the observed far-infrared properties of COS-87259 with other known objects at $z > 6.5$. We compare the [CII] luminosity, [CII] FWHM, and monochromatic $158\mu\text{m}$ dust continuum luminosities in panels (a), (b), and (c), respectively. A full description of the objects shown in this plot is provided in §3.2. The two known red type 1 quasars at $z > 6.5$ are shown as blue diamonds with red edges. Serendipitously-detected companions of quasars and Lyman-break galaxies are shown as empty markers. We correct for strong gravitational lensing where appropriate.

known $z > 6.5$ sub-mm galaxies (Strandet et al. 2017; Marrone et al. 2018), extremely UV-luminous quasars ($M_{\text{UV}} \leq -25$; Venemans et al. 2012; Bañados et al. 2015; Venemans et al. 2016, 2017b; Mazzucchelli et al. 2017; Decarli et al. 2018; Yang et al. 2019b, 2020b; Wang et al. 2021; Yue et al. 2021), UV-bright Lyman-break galaxies ($M_{\text{UV}} \leq -21$; Pentericci et al. 2016; Smit et al. 2018; Hashimoto et al. 2019; Bouwens et al. 2022; Inami et al. 2022; Schouws et al. 2022b,a), reddened type 1 quasars (Izumi et al. 2021; Fujimoto et al. 2022), as well as companion galaxies serendipitously identified from ALMA observations (Decarli et al. 2017; Willott et al. 2017; Neeleman et al. 2019; Venemans et al. 2019, 2020; Fudamoto et al. 2021). We correct the properties of the sub-mm system SPT0311-58 and the quasar J0439+1634 for strong gravitational lensing given reported magnification factors (Marrone et al. 2018; Yue et al. 2021).

With a [CII] luminosity of $(8.82 \pm 0.75) \times 10^9 L_{\odot}$, COS-87259 is the




Figure 5. Figure showing the *Spitzer*/IRAC 3.6 μm (blue), 4.5 μm (red), as well as *Spitzer*/MIPS 24 μm (green) emission around COS-87259. We show 7 and 12 σ contours for IRAC, and 4, 6, and 8 σ contours for MIPS. The background image is the near-IR stack centered on the rest-UV position of COS-87259. The centroids of each *Spitzer* detection near COS-87259 are closely aligned with the rest-UV centroid (<0.5 arcsec) and there is no evidence of significant confusion impacting any of the *Spitzer* detections.

second most [CII] luminous object known at $z > 6.5$ (Fig. 4a), being outshined by only the sub-mm galaxy SPT0311-58W at $z = 6.900$ ($1.02 \times 10^{10} L_{\odot}$). The [CII] line of COS-87259 is nearly $5 \times$ brighter than typical extremely bright $z > 6.5$ quasars (median $L_{[\text{CII}]}$ = $2.0 \times 10^9 L_{\odot}$) and $16 \times$ brighter than typical UV-bright Lyman-break $z > 6.5$ galaxies (median $L_{[\text{CII}]}$ = $0.55 \times 10^9 L_{\odot}$). COS-87259 additionally shows a [CII] line that is $4.7\text{--}8.0 \times$ brighter than the two reddened type 1 quasars thus far identified at $z > 6.5$ (Izumi et al. 2021; Fujimoto et al. 2022).

The [CII] line width of COS-87259 ($\text{FWHM} = 652 \pm 17 \text{ km s}^{-1}$) is also exceptionally broad for known systems at $z > 6.5$ (Fig. 4b), only surpassed by SPT0311-58W ($\text{FWHM} = 840 \pm 10 \text{ km s}^{-1}$; Marrone et al. 2018) and the $z = 6.66$ quasar PSO J338+29 ($\text{FWHM} = 740^{+540}_{-310} \text{ km s}^{-1}$; Mazzucchelli et al. 2017). For reference, the typical [CII] line widths of extremely UV-luminous $z > 6.5$ quasars and Lyman-break galaxies are 2.0 and $2.7 \times$ narrower than that of COS-87259, respectively. Finally, we note that the monochromatic $158\mu\text{m}$ dust continuum luminosity of COS-87259 ($(9.3 \pm 0.5) \times 10^{11} L_{\odot}$) is among the brightest of any known object at $z > 6.5$ (Fig. 4c). This $158\mu\text{m}$ continuum luminosity is very similar to that of $z > 6.5$ quasars, with only SPT0311-58W substantially outshining COS-87259 in the continuum around [CII]. While the conversion between $158\mu\text{m}$ luminosity to total infrared luminosity depends on dust temperature and emissivity, it is nonetheless clear that COS-87259 is among the most extreme objects known in the reionization era in terms of its observed [CII] and dust continuum properties. This suggests that COS-87259 is likely one of the most massive galaxies at $z \sim 7$ with a very large star formation rate, as supported by the full near-infrared to millimeter SED of this system (see §4).

3.3 Relative Positions of the Multi-wavelength Emission

To further characterize COS-87259, we compare the relative positions of the [CII], dust continuum, rest-UV, and radio emission, all of which are imaged with $\text{FWHM} \approx 0.4\text{--}0.8$ arcsec ($\approx 2\text{--}4$ kpc) resolution. The centroids of the [CII] and dust continuum emission are consistent within uncertainties ($\Delta\theta = 0.04 \pm 0.03$ arcsec), indicating that the $158\mu\text{m}$ dust continuum is concentrated near the dynamical center of the galaxy. Here, we have computed 1σ positional uncertainties from the ALMA data as¹ (beam FWHM)/(peak S/N)/0.9, enforcing a minimum positional uncertainty of $\sigma_{\text{min}} = (\text{beam FWHM})/20/0.9$ due to limits imposed by atmospheric phase fluctuations. We also find that the 3 GHz radio emission peak from COS-87259 (beam = 0.75 arcsec; Smolčić et al. 2017) is significantly offset from the dust continuum centroid with a separation of 1.21 ± 0.17 arcsec (see Fig. 1b). This large (6.4 ± 0.9 kpc projected) offset is to be expected if the radio emission seen from COS-87259 is powered by an extended relativistic jet emerging from the AGN.

The rest-UV emission centroid from COS-87259 also appears slightly offset from both the [CII] and dust continuum centroids (Fig. 1b). We measure the rest-UV centroid from a stack of the HSCSSP PDR2 y and UltraVISTA DR4² Y , J , H , and K_s images, each of which have a seeing-limited $\text{FWHM} \approx 0.77$ arcsec and are calibrated to the Gaia astrometric frame with an absolute position uncertainty of ≤ 0.05 arcsec (McCracken et al. 2012; Aihara et al. 2019). The stacked rest-UV detection of COS-87259 has a $S/N = 15$ suggesting a relative positional uncertainty of $\text{FWHM}/(S/N) \approx 0.05$ arcsec, comparable to the absolute precision. To be conservative, we adopt a total position uncertainty of 0.10 arcsec for the rest-UV emission. This results in a total offset of 0.30 ± 0.10 arcsec between the rest-UV and [CII] emission centroids, as well as a very similar offset from the dust continuum. Such moderate ($\sim 1\text{--}2$ kpc) offsets between the dust continuum and rest-UV centroids have been found among other UV-bright Lyman-break $z \sim 7$ galaxies and are often coupled with strong spatial gradients in the rest-UV slope, indicating considerable dust opacity variations across the galaxy (e.g. Carniani et al. 2018; Bowler et al. 2022).

COS-87259 is also detected at fairly high S/N ($\sim 10\text{--}20\sigma$) in the *Spitzer*/IRAC 3.6 μm , 4.5 μm , and MIPS 24 μm bands (see Fig. 5; E22a). Given the relatively poor angular resolution of the *Spitzer* images ($\text{FWHM} \approx 1.8\text{--}6$ arcsec), we investigate whether there is any evidence that these detections are significantly impacted by confusion. After calibrating the *Spitzer* images to the Gaia astrometric frame (using the IRAF task CCMAP), we find that the centroids of all detections are closely aligned with the rest-UV centroid of COS-87259 ($\Delta\theta < 0.5$ arcsec). Moreover, none of the *Spitzer* contours show any sign of strong asymmetry (see Fig. 5). The next nearest MIPS 24 μm detection is 12 arcsec (i.e. $2 \times$ the FWHM) away from COS-87259, while the next nearest strong IRAC detection is from an object 5.4 arcsec away (i.e. $3 \times$ the FWHM) located at the bottom of Fig. 5. Overall, we find no evidence that nearby objects are contributing significantly to the *Spitzer* detections of COS-87259. Nonetheless, the IRAC, MIPS, PACS, and SPIRE photometry of COS-87259 used below were all computed after performing deconvolution using the relatively high-resolution imaging as a prior (see E22a for further details).

¹ <https://help.almascience.org/kb/articles/what-is-the-absolute-astrometric-accuracy-of-alma>

² https://ultravista.org/release4/dr4_release.pdf

4 THE PHYSICAL PROPERTIES OF COS-87259

Now equipped with measurements of the [CII] systemic redshift and $158\mu\text{m}$ dust continuum emission, we reassess the inferred physical properties of COS-87259 derived in E22a. To start, we fit the observed near-infrared to millimeter SED of COS-87259 using a custom Bayesian SED fitting package (Lyu et al. 2022) that utilizes dynamic nested sampling (DYNesty; Speagle 2020) and features (semi-)empirical templates of AGN and star-forming dust emission. The validity of this tool has been demonstrated by finding and characterizing obscured AGN within the GOODS-S field in Lyu et al. (2022), and we refer the interested reader to that paper as well as Lyu & Rieke (2022, their section 4.2) for details of this code. For the purposes of this work, we adopt the Lyu et al. (2017) warm-dust-deficient AGN SED (which is suitable for very luminous AGN like COS-87259; see below) and account for obscuration using an empirical AGN infrared attenuation curve (Lyu & Rieke 2022). The galaxy dust emission is represented by the infrared SED template of Haro 11 which successfully matches the observed infrared SEDs and expected physical properties of massive $5 < z < 7$ galaxies (De Rossi et al. 2018), including the hosts of $z > 5$ quasars (Lyu et al. 2016). Along with these AGN and galaxy dust emission components, we include stellar emission using the Flexible Stellar Population Synthesis code (FSPS; Conroy et al. 2009; Conroy & Gunn 2010) under the SED fitting framework provided by PROSPECTOR (Johnson et al. 2021). For this stellar emission, we adopt a delayed star formation history, a Calzetti et al. (2000) attenuation curve (as is likely appropriate for a dust and hence metal rich system like COS-87259; e.g. Shivaeei et al. 2020), and a Kroupa (2001) stellar initial mass function.

In the SED fitting procedure, we include all VIRCcam, WFC3, IRAC, MIPS, PACS, SPIRE, and SCUBA-2 photometry of COS-87259 (see table 1 of E22a) as well as the $158\mu\text{m}$ dust continuum measurement from our new ALMA data. The only difference from the photometry adopted in E22a is that we here use the deboosted SCUBA-2 $850\mu\text{m}$ flux density measurement of 4.60 ± 1.65 mJy (Simpson et al. 2019). The IRAC, MIPS, PACS, and SPIRE photometry were all computed using deconfusion algorithms (see E22a). We note that the reported uncertainties on inferred physical properties below are largely dominated by the statistical uncertainties from the photometric measurements and do not include systematic uncertainties arising from our adopted set of SED templates. Using the x-CIGALE SED-fitting code (Yang et al. 2020a) as described in E22a, we infer similar values for the physical parameters (within $\lesssim 0.3$ dex).

Before reporting the results from our SED fit, we briefly discuss the photometric properties of COS-87259 to provide context on what is driving the inferred physical parameters (see E22a for further details). From the VIRCcam photometry, COS-87259 shows an extremely red rest-UV slope ($\beta = -0.6$ where $f_\lambda \propto \lambda^\beta$) compared to typical UV-bright $z \sim 7$ Lyman-break galaxies ($\beta \approx -1.8$; Bowler et al. 2017; Endsley et al. 2021a), indicating unusually strong dust attenuation. At slightly redder wavelengths, COS-87259 shows a jump in flux density between the K_s ($0.67 \pm 0.09 \mu\text{Jy}$) and IRAC $3.6+4.5\mu\text{m}$ bands ($\approx 2.75 \mu\text{Jy}$) that is much stronger than expected from extrapolating the rest-UV slope, consistent with a prominent Balmer break. The flux density then steeply rises between the IRAC $8\mu\text{m}$ ($< 4.9 \mu\text{Jy}$ 2σ), MIPS $24\mu\text{m}$ ($180 \pm 6 \mu\text{Jy}$), and PACS $100\mu\text{m}$ ($5100 \pm 1400 \mu\text{Jy}$) bands, as would be expected from hot dust emission powered by a highly-obscured AGN. Finally, we note that COS-87259 shows a relatively flat and bright SED between the PACS, SPIRE, and SCUBA-2 $160\text{--}850\mu\text{m}$ bands ($\approx 5\text{--}10$ mJy in each filter), indicative of very luminous hot (AGN powered) and warm (star formation powered) dust emission,




Figure 6. Near-infrared through millimeter SED of COS-87259. The photometric data (see E22a) and ALMA band 6 continuum measurement are shown as red circles with the best-fitting SED and its corresponding model photometry shown via a thick, light blue line and blue squares, respectively. We also show the relative contributions of the stellar, AGN, and (star-forming) dust emission as green, blue, and gold lines, respectively. The translucent lines show 20 realizations of the fit randomly sampled from the output posterior distribution to illustrate the fitting uncertainties. COS-87259 is inferred to be an extremely massive reionization-era galaxy harboring a heavily-obscured luminous AGN and powering intense dust-obscured star formation.

the latter of which is now corroborated by the strong ALMA $158\mu\text{m}$ continuum detection. As discussed in E22a, the MIPS, PACS, SPIRE, and SCUBA-2 photometry of COS-87259 are consistent with the SEDs of hot dust-obscured galaxies identified at lower redshifts (e.g. Eisenhardt et al. 2012; Tsai et al. 2015; Fan et al. 2016).

From the SED fit (see Fig. 6), COS-87259 is inferred to harbor a very luminous AGN ($L_{\text{IR,AGN}} = (2.5 \pm 0.2) \times 10^{13} L_\odot$) that is heavily obscured with an optical depth at rest-frame $9.7\mu\text{m}$ of $\tau_{9.7\mu\text{m}} = 2.3 \pm 0.1$. Such inferred obscuration is consistent with Compton-thick absorption ($N_H \gtrsim 10^{24} \text{ cm}^{-2}$; Shi et al. 2006) and helps explain the weak X-ray luminosity of COS-87259, which remains undetected in the 160 ks *Chandra* imaging over COSMOS (Civano et al. 2016; see E22a for further details). Using the Runnoe et al. (2012) bolometric conversion from rest-frame 5100 \AA luminosity, the intrinsic (de-reddened) AGN SED implied by our fits suggests a bolometric AGN luminosity of $L_{\text{bol,AGN}} = (5.1 \pm 0.5) \times 10^{13} L_\odot$. Such a large bolometric AGN luminosity is comparable to that of $z \sim 1\text{--}4$ hot dust obscured galaxies (e.g. Eisenhardt et al. 2012; Tsai et al. 2015; Fan et al. 2016) as well as $z \sim 7$ quasars (e.g. Mazzucchelli et al. 2017; Yang et al. 2021), and implies a black hole mass of $M_{\text{BH}} = (1.6 \pm 0.2) \times 10^9 M_\odot / \lambda_{\text{Edd}}$ for COS-87259 where λ_{Edd} is the Eddington ratio. We come back to discuss potential implications of identifying such an early supermassive black hole over a relatively small area (1.5 deg^2) at the end of this section.

Due to the strong jump in flux density between the VIRCcam K_s and IRAC bands, the rest UV+optical emission of COS-87259 is inferred to be powered by star formation (with a prominent Balmer break) rather than by the AGN (Fig. 6). To further assess whether the rest UV+optical photometry of COS-87259 may plausibly be due to reddened AGN emission, we also compare these data to empirical spectra of lower-redshift quasars. Specifically, we fit the VIRCcam Y , J , H , and K_s as well as the IRAC $4.5\mu\text{m}$ photometry of COS-87259 to the composite $z \sim 1.5$ blue quasar spectrum from Selsing et al.

(2016) after redshifting to $z = 6.853$ and applying a range of V-band optical depths ($0 < A_V < 2$) using the SMC attenuation law (Pei 1992) motivated by previous studies of red quasars (e.g. Richards et al. 2003; Hopkins et al. 2004). In this fit, we do not include the $3.6\mu\text{m}$ band as it can be contaminated by strong [OIII]+H β emission at the redshift of COS-87259. This fitting procedure results in a best-fitting reduced $\chi^2 = 11.8$ with 3–6 σ offsets in the K_s and $4.5\mu\text{m}$ bands, indicating that the rest UV+optical photometry of COS-87259 is poorly explained by such AGN spectra. Ultimately, UV+optical spectra of COS-87259 (taken with e.g. JWST/NIRSpec) will be necessary to determine the physical origin of this emission.

Assuming that the UV+optical emission is dominated by star formation, the fiducial SED fit shown in Fig. 6 returns a total stellar mass of $M_* = (1.7 \pm 0.7) \times 10^{11} M_\odot$, implying that COS-87259 is an extremely massive $z \sim 7$ galaxy as perhaps expected given its extreme far-IR properties (Fig. 4). However, as discussed elsewhere (van Mierlo et al. 2022a; Whittler et al. 2023; see also E22a), alternative assumptions to the star formation history can yield significantly lower stellar mass estimates for COS-87259 ($M_* \sim (1-3) \times 10^{10} M_\odot$) wherein the strong jump in flux density between K_s and [3.6] is inferred to be strongly assisted by prominent [OIII]+H β emission. Nonetheless, the full range of estimated stellar masses for COS-87259 ($\sim 10^{10-11} M_\odot$) still falls substantially below that expected given local relations between the properties of supermassive black holes and their host galaxies. With an inferred black hole mass of $\sim 1.5 \times 10^9 M_\odot$, the relation of Häring & Rix (2004) predicts a stellar mass of $\sim 10^{12} M_\odot$ for COS-87259. Similar inferred offsets towards higher black hole mass at fixed galaxy mass have been reported for luminous high-redshift quasars (e.g. Trakhtenbrot et al. 2015; Venemans et al. 2016; Pensabene et al. 2020) as well as hot dust obscured galaxies (Assef et al. 2015; Tsai et al. 2015), potentially indicating that the growth of their supermassive black holes outpaced that of the host galaxies at early times.

The models also suggest that intense obscured star formation within COS-87259 is powering a total (8–1000 μm rest-frame) IR luminosity of $L_{\text{IR,SF}} = (9.1 \pm 1.1) \times 10^{12} L_\odot$, with a corresponding very large star formation rate of $1610 \pm 190 M_\odot/\text{yr}$. For comparison, we also derive the obscured star formation properties of COS-87259 adopting the standard approach for $z > 6$ quasars. In these objects, the dust emission is often assumed to follow a modified blackbody with temperature $T = 47$ K and emissivity index $\beta = 1.6$ (e.g. Beelen et al. 2006; Decarli et al. 2018; Venemans et al. 2020). After fixing the SED normalization to match the $158\mu\text{m}$ dust continuum measurement of COS-87259, this approach yields an IR luminosity and obscured star formation rate (Murphy et al. 2011) of $(8.7 \pm 0.5) \times 10^{12} L_\odot$ and $1300 \pm 70 M_\odot/\text{yr}$, respectively, similar to that obtained from the SED fit using the Haro 11 template. The modified blackbody fit also implies a very large dust mass of $M_{\text{dust}} = (1.9 \pm 0.1) \times 10^9 M_\odot$. Finally, we note that if we adopt the SFR- $L_{\text{[CII]}}$ conversion of (Herrera-Camus et al. 2015), we infer a total star formation rate of $2970 \pm 250 M_\odot/\text{yr}$. While this [CII]-based SFR is $2.3\times$ larger than that inferred from the dust SED, we note that there remain considerable uncertainties on both the dust temperature and emissivity index for COS-87259. Regardless, the very bright ALMA detections of COS-87259 make it clear that this object is undergoing vigorous obscured star formation activity comparable to $z \sim 7$ sub-mm galaxies and quasar hosts (Fig. 4). The range of stellar masses ($\sim 10^{10-11} M_\odot$) and SFRs (1300–3000 M_\odot/yr) inferred for COS-87259 suggest a specific SFR between 13 to 300 Gyr^{-1} , consistent with the range of sSFRs inferred among otherwise typical extremely UV-luminous ($M_{\text{UV}} \sim -22$) $z \sim 7-8$ galaxies (Topping et al. 2022). Additional

data will be required to determine whether the AGN within COS-87259 is significantly impacting its star formation efficiency.

With an inferred bolometric AGN luminosity of $\sim 5 \times 10^{13} L_\odot$, the supermassive black hole within COS-87259 is comparable to that powering the brightest quasars known at $z > 6$ (e.g. Yang et al. 2021) which are discovered using extremely wide-area ($\geq 1000 \text{ deg}^2$) surveys. The identification of COS-87259 within the relatively small 1.5 deg^2 COSMOS field is therefore very surprising. To better place COS-87259 into context of the UV-luminous quasar population, we extract its de-reddened rest-frame 1450 \AA AGN luminosity from the SED fit described in §4. This suggests that COS-87259 is a highly-obscured version of a $M_{1450} \approx -27.3$ quasar which, at $z \sim 7$, have been found to be powered by $\approx 1.5 \times 10^9 M_\odot$ black holes accreting near the Eddington limit (Yang et al. 2021). Such UV-luminous quasars ($M_{1450} \leq -27$) are an exceedingly rare population in the very early Universe with a surface density of ≈ 1 per 3000 deg^2 at $z = 6.6-6.9$ (Wang et al. 2019), the approximate redshift interval over which COS-87259 was selected. While COS-87259 is only one object and thus may have simply been a fortuitous discovery, we briefly discuss what the spectroscopic confirmation of this object may imply for the assembly of supermassive black holes at $z \geq 7$. We refer the interested reader to section 4.2 of E22a for a more extended discussion of what COS-87259 may imply for the statistical properties of UV-bright ($M_{\text{UV}} \lesssim -21.25$) massive ($M_* > 10^{10} M_\odot$) $z \sim 7$ galaxies, as well as the possible contribution of this population to the obscured star formation rate density at early times.

The identification of COS-87259 within the 1.5 deg^2 COSMOS field versus the very low surface density of similarly luminous $z \sim 6.8$ quasars ($\approx 1/3000 \text{ deg}^{-2}$; see above) raises the question of whether rest-UV selections may be missing a very large fraction ($\geq 1000\times$) of heavily-obscured black hole growth in the early Universe. Observations of both the local Universe and the $z > 6$ quasar population provide some insight. The known population of early UV-luminous quasars suggest that approximately 0.1% of the total black hole mass density inferred at present day had already formed by $z = 6$ (e.g. Marconi et al. 2004; Shen et al. 2020). Therefore, boosting the total black hole mass density at $z > 6$ by a factor of ≥ 1000 (to correct for an extremely dominant population of heavily-obscured AGN) would imply little cosmic black hole growth between $0 < z < 6$. Such a scenario is inconsistent with the census of AGN activity at these redshifts (Shen et al. 2020). This line of reasoning suggests it is unlikely that there are ≥ 3 orders of magnitude more buried AGN at $z > 6$ than expected based on results from rest-UV selected samples.

However, it remains unknown whether the most massive ($\sim 10^9 M_\odot$) black holes at $z \geq 7$ very frequently grew in heavily-obscured phases. Indirect evidence for such a scenario has emerged in recent years, with spectroscopic observations showing that a large fraction of UV-luminous $z > 6$ quasars possess much smaller Ly α proximity zones than expected given their black hole masses (Davies et al. 2019; Eilers et al. 2020, 2021). Even though these early supermassive black holes have presumably been growing for several hundred million years (Inayoshi et al. 2020), their proximity zones imply that they have only been ionizing their surrounding IGM for $\sim 10^6$ years on average (Morey et al. 2021). This apparent tension can be alleviated if the supermassive black holes powering $z > 6$ quasars were heavily obscured for a large fraction ($\geq 95\%$) of their lifetime (Eilers et al. 2018; Davies et al. 2019). However, it is not clear what physical mechanism might drive such a very dominant population of heavily-obscured luminous AGN at $z \geq 7$ given that X-ray observations imply an obscured ($N_H > 10^{23} \text{ cm}^{-2}$) fraction of $\sim 80\%$ among luminous ($L_{\text{bol}} \sim 10^{13} L_\odot$) AGN at $z \sim 3-6$ (Vito et al. 2018). One potential contributing factor could be the increasing gas and dust density within

massive galaxies towards higher redshifts (Circosta et al. 2019), as suggested by recent hydrodynamic simulations (Trebitsch et al. 2019; Ni et al. 2020; Lupi et al. 2022). Ultimately, wider-area searches for very luminous ($L_{\text{bol}} \gtrsim 10^{13} L_{\odot}$) heavily-obscured AGN at $z > 6$ will be required to better assess how common this population is relative to quasars, with subsequent detailed follow-up necessary to characterize the origin of their obscuration.

5 SUMMARY AND OUTLOOK

In this work, we have reported the results of ALMA follow-up observations targeting the [CII]158 μm line and underlying dust continuum from a previously-reported candidate heavily-obscured radio-loud ($L_{1.4\text{ GHz}} = 10^{25.4} \text{ W Hz}^{-1}$) AGN at $z \simeq 6.83$ identified in the 1.5 deg² COSMOS field (COS-87259; E22a). Our summary and conclusions are listed below:

(i) The ALMA data confidently reveal a luminous emission line with a central frequency of 242.009 ± 0.007 GHz coincident with the near-infrared position of COS-87259 (Fig. 1). We conclude that this emission feature is the [CII]158 μm cooling line at $z_{\text{[CII]}} = 6.8532 \pm 0.0002$ which is in excellent agreement with the precise photometric redshift of COS-87259 ($z_{\text{phot}} = 6.83 \pm 0.06$) set by the sharp Lyman-alpha break seen between two narrow/intermediate-bands at 9450 Å and 9700 Å (E22a).

(ii) The [CII]158 μm emission of COS-87259 is spatially resolved with our moderate-resolution (beam \approx 0.4 arcsec) ALMA data and its surface brightness profile suggests the presence of a compact (major axis = 1.9 ± 0.4 kpc) [CII]-emitting core in addition to a more extended (major axis = 9.6 ± 1.1 kpc) region of emission (Fig. 2), similar to that of $z > 6$ quasar hosts (Novak et al. 2020). Summing the two emission components, we measure a total [CII] line luminosity of $(8.82 \pm 0.75) \times 10^9 L_{\odot}$ from COS-87259, the bulk of which is originating from the more extended region (75 \pm 10%). The [CII] emission is extremely broad (FWHM = 652 ± 17 km s⁻¹) and a prominent gradient is seen in the mean velocity field with a maximum difference of ≈ 360 km s⁻¹ between spaxels in the northeast versus southwest corners of COS-87259 (Fig. 3).

(iii) Strong 158 μm dust continuum emission is also detected from COS-87259 with a surface brightness profile consistent with a compact (major axis = 1.4 ± 0.2 kpc) core and a more extended (major axis = 7.8 ± 1.0 kpc) region of emission (Fig. 2), similar to that of the [CII] line. The combined flux density of these two dust continuum components is measured to be 2.71 ± 0.16 mJy and is almost equally divided among the two components (55 \pm 4% from the compact core).

(iv) COS-87259 exhibits the second most luminous and third broadest [CII] line known at $z > 6.5$ and the 158 μm dust continuum monochromatic luminosity of this system is also very similar to that of the IR-brightest $z \sim 7$ quasar host galaxies (Fig. 4). The ALMA data therefore indicate that COS-87259 is among the most extreme objects known in the reionization-era in terms of its far-IR properties, suggesting it is likely one of the most massive galaxies at $z \sim 7$ with a very large star formation rate.

(v) The centroids of the [CII] and dust continuum emission are consistent within uncertainties, though we identify significant offsets between the ALMA centroids and the rest-UV as well as radio centroids. The ≈ 6 kpc projected separation between the 3 GHz detection peak and the [CII]+dust continuum centroids is to be expected if the radio emission is powered by an extended relativistic jet emerging from the AGN.

(vi) The MIPS, PACS, and SPIRE detections of COS-87259 in-

dicating that this source harbors a heavily-obscured AGN with a rest-frame 9.7 μm optical depth of $\tau_{9.7\mu\text{m}} = 2.3 \pm 0.1$, consistent with Compton-thick absorption. The very large bolometric AGN luminosity of $(5.1 \pm 0.5) \times 10^{13} L_{\odot}$ implies that COS-87259 is a highly-obscured version of a $M_{1450} \approx -27.3$ quasar at $z \sim 7$ and is powered by a $\gtrsim 1.5 \times 10^9 M_{\odot}$ black hole (assuming an Eddington ratio of ≤ 1). Comparably luminous quasars are an exceedingly rare population at $z \sim 7$ (~ 0.001 deg⁻²; Wang et al. 2019) while COS-87259 was identified over a relatively small (1.5 deg²) area. Wider-area searches for very luminous ($L_{\text{bol}} \gtrsim 10^{13} L_{\odot}$) heavily-obscured AGN at $z > 6$ will be required to better assess how common this population is relative to quasars.

(vii) The 158 μm dust continuum emission suggests that COS-87259 is forming stars at a rate of $\approx 1300 M_{\odot}/\text{yr}$, respectively, consistent with the estimate from the near-IR through millimeter SED fit ($\approx 1610 M_{\odot}/\text{yr}$). Regardless of the exact value, it is clear from the strong ALMA detections that COS-87259 is undergoing vigorous obscured star formation activity comparable to $z \sim 7$ sub-mm galaxies and quasar hosts. The VIRCAM and IRAC photometry of COS-87259 suggest that its rest UV+optical emission is powered by star formation rather than by the AGN. Under this assumption, we estimate an extremely large stellar mass of $(1.7 \pm 0.7) \times 10^{11} M_{\odot}$ for COS-87259, though different assumptions on star formation history can yield significantly lower mass estimates given existing broadband photometry (E22a; van Mierlo et al. 2022a; Whittler et al. 2023). Nonetheless, all stellar mass estimates of COS-87259 are at least an order of magnitude lower than that expected from local scaling relations between the properties of supermassive black holes and their host galaxies.

Future wide-area datasets will provide a greatly improved understanding of the incidence of heavily-obscured supermassive black hole growth and dust-enshrouded stellar mass buildup within the most massive reionization-era galaxies. Within the coming decade, surveys with *Roman* and *Euclid* are expected to cover hundreds to thousands of square degrees with optical+near-infrared (0.9–2 μm) depths of $m_{5\sigma} \sim 26 - 27$ (Spergel et al. 2015; van Mierlo et al. 2022b; Euclid Collaboration et al. 2022). These surveys will therefore enable the identification of extremely red ($\beta > -1$) yet UV-bright ($M_{\text{UV}} \lesssim -21.5$) $z \gtrsim 7$ galaxies like COS-87259 over a much wider area via the presence of very strong ($\gtrsim 3$ mag) Lyman-alpha breaks. Sensitive follow-up at mid/far-infrared, radio, and X-ray wavelengths will provide the opportunity to investigate both obscured star formation as well as heavily-obscured AGN activity within these sources, delivering a much clearer picture of the typical characteristics of the most massive reionization-era galaxies. Such studies will complement searches of red yet relatively unobscured quasars (e.g. Kato et al. 2020; Fujimoto et al. 2022), radio-selected AGN (e.g. Smith et al. 2016; Saxena et al. 2018; Broderick et al. 2022), and X-ray selected AGN (Nandra et al. 2013; Weisskopf et al. 2015) in the reionization epoch to build a more complete census of early supermassive black hole growth mechanisms.

ACKNOWLEDGEMENTS

The authors sincerely thank the anonymous referee for providing helpful, constructive comments that significantly improved the quality of this work. RE and DPS acknowledge funding from JWST/near-IRCam contract to the University of Arizona, NAS5-02015. DPS acknowledges support from the National Science Foundation through the grant AST-2109066. FW acknowledges the support provided by NASA through the NASA Hubble Fellowship grant #HST-HF2-51448.001-A awarded by the Space Telescope Science Institute,

which is operated by the Association of Universities for Research in Astronomy, Incorporated, under NASA contract NAS5-26555. JY and XF acknowledge support from the US NSF Grant AST 19-08284. RS acknowledges support from a STFC Ernest Rutherford Fellowship (ST/S004831/1). RJB acknowledges support from TOP grant TOP1.16.057 provided by the Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO).

This paper makes use of the following ALMA data: ADS/JAO.ALMA#2021.1.01311.S. ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada), MOST and ASIAA (Taiwan), and KASI (Republic of Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO and NAOJ. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

This work is based [in part] on observations made with the Spitzer Space Telescope, which was operated by the Jet Propulsion Laboratory, California Institute of Technology under a contract with NASA. Herschel is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA. Based on data products from observations made with ESO Telescopes at the La Silla Paranal Observatory under ESO programme ID 179.A-2005 and on data products produced by CALET and the Cambridge Astronomy Survey Unit on behalf of the UltraVISTA consortium. This research has made use of the NASA/IPAC Infrared Science Archive, which is funded by the National Aeronautics and Space Administration and operated by the California Institute of Technology.

This research made use of *ASTROPY*, a community-developed core PYTHON package for Astronomy (*Astropy Collaboration et al. 2013; Price-Whelan et al. 2018*); *MATPLOTLIB* (*Hunter 2007*); *NUMPY* (*Harris et al. 2020*); and *SCIPY* (*Virtanen et al. 2020*).

DATA AVAILABILITY

The ALMA data that is the focus of this article is publicly available on the ALMA science archive (2021.1.01311.S). The remaining data on COS-87259 are detailed in [Endsley et al. \(2022a\)](#).

REFERENCES

- Aihara H., et al., 2019, *PASJ*, 71, 114
Aihara H., et al., 2022, *PASJ*, 74, 247
Assef R. J., et al., 2015, *ApJ*, 804, 27
Astropy Collaboration et al., 2013, *A&A*, 558, A33
Auger M. W., Treu T., Bolton A. S., Gavazzi R., Koopmans L. V. E., Marshall P. J., Moustakas L. A., Burles S., 2010, *ApJ*, 724, 511
Bañados E., Decarli R., Walter F., Venemans B. P., Farina E. P., Fan X., 2015, *ApJ*, 805, L8
Bañados E., et al., 2018, *Nature*, 553, 473
Bañados E., et al., 2021, *ApJ*, 909, 80
Beelen A., Cox P., Benford D. J., Dowell C. D., Kovács A., Bertoldi F., Omont A., Carilli C. L., 2006, *ApJ*, 642, 694
Bouwens R. J., et al., 2011, *ApJ*, 737, 90
Bouwens R. J., Illingworth G. D., Oesch P. A., Caruana J., Holwerda B., Smit R., Wilkins S., 2015, *ApJ*, 811, 140
Bouwens R. J., et al., 2021, *AJ*, 162, 47
Bouwens R. J., et al., 2022, *ApJ*, 931, 160
Bowler R. A. A., Dunlop J. S., McLure R. J., McLeod D. J., 2017, *MNRAS*, 466, 3612
Bowler R. A. A., Jarvis M. J., Dunlop J. S., McLure R. J., McLeod D. J., Adams N. J., Milvang-Jensen B., McCracken H. J., 2020, *MNRAS*, 493, 2059
Bowler R. A. A., Cullen F., McLure R. J., Dunlop J. S., Avison A., 2022, *MNRAS*, 510, 5088
Broderick J. W., et al., 2022, *Publ. Astron. Soc. Australia*, 39, e061
Calzetti D., Armus L., Bohlin R. C., Kinney A. L., Koornneef J., Storchi-Bergmann T., 2000, *ApJ*, 533, 682
Carniani S., et al., 2018, *MNRAS*, 478, 1170
Circosta C., et al., 2019, *A&A*, 623, A172
Civano F., et al., 2016, *ApJ*, 819, 62
Conroy C., Gunn J. E., 2010, *ApJ*, 712, 833
Conroy C., Gunn J. E., White M., 2009, *ApJ*, 699, 486
Davies F. B., Hennawi J. F., Eilers A.-C., 2019, *ApJ*, 884, L19
De Barros S., Oesch P. A., Labbé I., Stefanon M., González V., Smit R., Bouwens R. J., Illingworth G. D., 2019, *MNRAS*, 489, 2355
De Rossi M. E., Rieke G. H., Shivaee I., Bromm V., Lyu J., 2018, *ApJ*, 869, 4
Decarli R., et al., 2017, *Nature*, 545, 457
Decarli R., et al., 2018, *ApJ*, 854, 97
Decarli R., et al., 2022, *A&A*, 662, A60
Eilers A.-C., Hennawi J. F., Davies F. B., 2018, *ApJ*, 867, 30
Eilers A.-C., et al., 2020, *ApJ*, 900, 37
Eilers A.-C., Hennawi J. F., Davies F. B., Simcoe R. A., 2021, *ApJ*, 917, 38
Eisenhardt P. R. M., et al., 2012, *ApJ*, 755, 173
Ellis R. S., et al., 2013, *ApJ*, 763, L7
Endsley R., Stark D. P., 2022, *MNRAS*, 511, 6042
Endsley R., Stark D. P., Chevallard J., Charlot S., 2021a, *MNRAS*, 500, 5229
Endsley R., Stark D. P., Charlot S., Chevallard J., Robertson B., Bouwens R. J., Stefanon M., 2021b, *MNRAS*, 502, 6044
Endsley R., et al., 2022a, *MNRAS*, 512, 4248
Endsley R., et al., 2022b, *MNRAS*, 517, 5642
Euclid Collaboration et al., 2022, *A&A*, 662, A112
Everett W. B., et al., 2020, *ApJ*, 900, 55
Fan L., Han Y., Nikutta R., Drouart G., Knudsen K. K., 2016, *ApJ*, 823, 107
Finkelstein S. L., et al., 2015, *ApJ*, 810, 71
Finkelstein S. L., et al., 2019, *ApJ*, 879, 36
Fudamoto Y., et al., 2021, *Nature*, 597, 489
Fujimoto S., et al., 2022, *Nature*, 604, 261
Furusawa H., et al., 2016, *ApJ*, 822, 46
Gialavisco M., et al., 2004, *ApJ*, 600, L93
Ginolfi M., et al., 2020, *A&A*, 633, A90
Grogan N. A., et al., 2011, *ApJS*, 197, 35
Harikane Y., et al., 2022, *ApJS*, 259, 20
Håring N., Rix H.-W., 2004, *ApJ*, 604, L89
Harris C. R., et al., 2020, *Nature*, 585, 357
Hashimoto T., et al., 2019, *PASJ*, 71, 71
Herrera-Camus R., et al., 2015, *ApJ*, 800, 1
Hopkins P. F., et al., 2004, *AJ*, 128, 1112
Hunter J. D., 2007, *Computing in Science & Engineering*, 9, 90
Illingworth G. D., et al., 2013, *ApJS*, 209, 6
Inami H., et al., 2022, *MNRAS*, 515, 3126
Inayoshi K., Visbal E., Haiman Z., 2020, *ARA&A*, 58, 27
Inoue A. K., et al., 2020, *PASJ*,
Ishigaki M., Kawamata R., Ouchi M., Oguri M., Shimasaku K., Ono Y., 2018, *ApJ*, 854, 73
Izumi T., et al., 2021, *ApJ*, 908, 235
Jarvis M. J., et al., 2013, *MNRAS*, 428, 1281
Johnson B. D., Leja J., Conroy C., Speagle J. S., 2021, *ApJS*, 254, 22
Kato N., et al., 2020, *PASJ*, 72, 84
Koekemoer A. M., et al., 2011, *ApJS*, 197, 36
Kroupa P., 2001, *MNRAS*, 322, 231
Labbé I., et al., 2013, *ApJ*, 777, L19
Lawrence A., et al., 2007, *MNRAS*, 379, 1599
Lotz J. M., et al., 2017, *ApJ*, 837, 97
Lupi A., Volonteri M., Decarli R., Bovino S., Silk J., 2022, *MNRAS*, 510, 5760
Lyu J., Rieke G., 2022, *Universe*, 8, 304
Lyu J., Rieke G. H., Albers S., 2016, *ApJ*, 816, 85

- Lyu J., Rieke G. H., Shi Y., 2017, *ApJ*, **835**, 257
- Lyu J., Alberts S., Rieke G. H., Rujopakarn W., 2022, *ApJ*, **941**, 191
- Maiolino R., et al., 2012, *MNRAS*, **425**, L66
- Marconi A., Risaliti G., Gilli R., Hunt L. K., Maiolino R., Salvati M., 2004, *MNRAS*, **351**, 169
- Marrone D. P., et al., 2018, *Nature*, **553**, 51
- Mason C. A., Trenti M., Treu T., 2015, *ApJ*, **813**, 21
- Matsuoka Y., et al., 2018, *ApJS*, **237**, 5
- Matsuoka Y., et al., 2019, *ApJ*, **883**, 183
- Matsuoka Y., et al., 2022, *ApJS*, **259**, 18
- Matthee J., et al., 2019, *ApJ*, **881**, 124
- Mazzucchelli C., et al., 2017, *ApJ*, **849**, 91
- McCracken H. J., et al., 2012, *A&A*, **544**, A156
- McLure R. J., et al., 2013, *MNRAS*, **432**, 2696
- McMullin J. P., Waters B., Schiebel D., Young W., Golap K., 2007, in Shaw R. A., Hill F., Bell D. J., eds, *Astronomical Society of the Pacific Conference Series Vol. 376, Astronomical Data Analysis Software and Systems XVI*. p. 127
- Morey K. A., Eilers A.-C., Davies F. B., Hennawi J. F., Simcoe R. A., 2021, *ApJ*, **921**, 88
- Mortlock D. J., et al., 2011, *Nature*, **474**, 616
- Murphy E. J., et al., 2011, *ApJ*, **737**, 67
- Nandra K., et al., 2013, arXiv e-prints, p. arXiv:1306.2307
- Neeleman M., et al., 2019, *ApJ*, **882**, 10
- Neeleman M., et al., 2021, *ApJ*, **911**, 141
- Ni Y., Di Matteo T., Gilli R., Croft R. A. C., Feng Y., Norman C., 2020, *MNRAS*, **495**, 2135
- Novak M., et al., 2020, *ApJ*, **904**, 131
- Oesch P. A., Bouwens R. J., Illingworth G. D., Labbé I., Stefanon M., 2018, *ApJ*, **855**, 105
- Oke J. B., Gunn J. E., 1983, *ApJ*, **266**, 713
- Ono Y., et al., 2012, *ApJ*, **744**, 83
- Pei Y. C., 1992, *ApJ*, **395**, 130
- Pensabene A., Carniani S., Perna M., Cresci G., Decarli R., Maiolino R., Marconi A., 2020, *A&A*, **637**, A84
- Pensabene A., et al., 2021, *A&A*, **652**, A66
- Pentericci L., et al., 2016, *ApJ*, **829**, L11
- Price-Whelan A. M., et al., 2018, *AJ*, **156**, 123
- Reed S. L., et al., 2017, *MNRAS*, **468**, 4702
- Reed S. L., et al., 2019, *MNRAS*, **487**, 1874
- Richards G. T., et al., 2003, *AJ*, **126**, 1131
- Robertson B. E., 2022, *ARA&A*, **60**, 121
- Runnoe J. C., Brotherton M. S., Shang Z., 2012, *MNRAS*, **422**, 478
- Saxena A., et al., 2018, *MNRAS*, **480**, 2733
- Schouws S., et al., 2022a, arXiv e-prints, p. arXiv:2202.04080
- Schouws S., et al., 2022b, *ApJ*, **928**, 31
- Selsing J., Fynbo J. P. U., Christensen L., Krogager J. K., 2016, *A&A*, **585**, A87
- Shen X., Hopkins P. F., Faucher-Giguère C.-A., Alexander D. M., Richards G. T., Ross N. P., Hickox R. C., 2020, *MNRAS*, **495**, 3252
- Shi Y., et al., 2006, *ApJ*, **653**, 127
- Shivaei I., et al., 2020, *ApJ*, **899**, 117
- Simpson J. M., et al., 2019, *ApJ*, **880**, 43
- Smit R., et al., 2014, *ApJ*, **784**, 58
- Smit R., et al., 2015, *ApJ*, **801**, 122
- Smit R., et al., 2018, *Nature*, **553**, 178
- Smith D. J. B., et al., 2016, in Reylé C., Richard J., Cambrésy L., Deleuil M., Pécontal E., Tresse L., Vauglin I., eds, *SF2A-2016: Proceedings of the Annual meeting of the French Society of Astronomy and Astrophysics*. pp 271–280 (arXiv:1611.02706)
- Smolčić V., et al., 2017, *A&A*, **602**, A1
- Song M., et al., 2016, *ApJ*, **825**, 5
- Speagle J. S., 2020, *MNRAS*, **493**, 3132
- Spergel D., et al., 2015, arXiv e-prints,
- Spilker J. S., et al., 2022, *ApJ*, **929**, L3
- Stark D. P., Schenker M. A., Ellis R., Robertson B., McLure R., Dunlop J., 2013, *ApJ*, **763**, 129
- Stefanon M., et al., 2019, *ApJ*, **883**, 99
- Stefanon M., Bouwens R. J., Labbé I., Illingworth G. D., Gonzalez V., Oesch P. A., 2021, *ApJ*, **922**, 29
- Strandet M. L., et al., 2017, *ApJ*, **842**, L15
- Topping M. W., et al., 2022, *MNRAS*, **516**, 975
- Trakhtenbrot B., et al., 2015, *Science*, **349**, 168
- Trebitsch M., Volonteri M., Dubois Y., 2019, *MNRAS*, **487**, 819
- Trenti M., et al., 2011, *ApJ*, **727**, L39
- Tsai C.-W., et al., 2015, *ApJ*, **805**, 90
- Valentino F., et al., 2022, *ApJ*, **929**, L9
- Venemans B. P., et al., 2012, *ApJ*, **751**, L25
- Venemans B. P., et al., 2013, *ApJ*, **779**, 24
- Venemans B. P., et al., 2015, *ApJ*, **801**, L11
- Venemans B. P., Walter F., Zschaechner L., Decarli R., De Rosa G., Findlay J. R., McMahon R. G., Sutherland W. J., 2016, *ApJ*, **816**, 37
- Venemans B. P., et al., 2017a, *ApJ*, **845**, 154
- Venemans B. P., et al., 2017b, *ApJ*, **851**, L8
- Venemans B. P., Neeleman M., Walter F., Novak M., Decarli R., Hennawi J. F., Rix H.-W., 2019, *ApJ*, **874**, L30
- Venemans B. P., et al., 2020, *ApJ*, **904**, 130
- Virtanen P., et al., 2020, *Nature Methods*, **17**, 261
- Vito F., et al., 2018, *MNRAS*, **473**, 2378
- Wang F., et al., 2017, *ApJ*, **839**, 27
- Wang F., et al., 2018, *ApJ*, **869**, L9
- Wang F., et al., 2019, *ApJ*, **884**, 30
- Wang F., et al., 2021, *ApJ*, **907**, L1
- Weaver J. R., et al., 2022, *ApJS*, **258**, 11
- Weisskopf M. C., Gaskin J., Tananbaum H., Vikhlinin A., 2015, in Hudec R., Pina L., eds, *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series Vol. 9510, EUV and X-ray Optics: Synergy between Laboratory and Space IV*. p. E2 (arXiv:1505.00814), doi:10.1117/12.2185084
- Whitler L., Stark D. P., Endsley R., Leja J., Charlot S., Chevallard J., 2023, *MNRAS*,
- Willott C. J., Bergeron J., Omont A., 2017, *ApJ*, **850**, 108
- Yang J., et al., 2019a, *AJ*, **157**, 236
- Yang J., et al., 2019b, *ApJ*, **880**, 153
- Yang G., et al., 2020a, *MNRAS*, **491**, 740
- Yang J., et al., 2020b, *ApJ*, **897**, L14
- Yang J., et al., 2021, *ApJ*, **923**, 262
- Yue M., et al., 2021, *ApJ*, **917**, 99
- van Mierlo S. E., Caputi K. I., Kokorev V., 2022a, arXiv e-prints, p. arXiv:2212.04511
- van Mierlo S. E., et al., 2022b, *A&A*, **666**, A200

APPENDIX A: PV DIAGRAM

We show the position-velocity diagram of [CII]158 μ m emission from COS-87259 in Fig. A1.

This paper has been typeset from a \LaTeX file prepared by the author.




Figure A1. The position-velocity diagram of [CII]158μm emission from COS-87259 extracted along the major axis. There is no clear evidence of multiple distinct components separated in velocity space.