A barycentric trigonometric Hermite interpolant via an iterative approach

Giacomo Elefante^a

^aDepartment of Mathematics "Tullio Levi-Civita", University of Padova, Via Trieste 63, Padova, Italy, giacomo.elefante@unipd.it

Abstract

In this work we extend and generalise an iterative approach, introduced by Cirillo and Hormann [9] for the Floater-Hormann family of interpolants, to construct an Hermite interpolant for general interpolant with basis functions which satisfy a Lagrange property. In particular, we apply this scheme to produce an effective barycentric rational trigonometric Hermite interpolant using the basis functions of the trigonometric interpolant introduced by Berrut [5]. Moreover, in order to give an easy construction of such an interpolant we compute analytically the differentation matrix and we conclude with various examples and a numerical study of the rate of convergence at equidistant nodes.

Keywords: barycentric interpolation, rational interpolation, trigonometric interpolation, Hermite interpolation 2020 MSC: 65D05, 65T40, 42A15.

1 Introduction

Hermite interpolation is a classical problem which has been widely explored. It has many applications, and therefore, it is still vivid and studied.

The problem consists into finding a good interpolant that satisfies the Hermite conditions

$$r_m^{(j)}(x_i) = f_i^{(j)} = f^{(j)}(x_i), \quad i = 0, \dots, n; \ j = 0, \dots, m,$$
(1)

which means that it has to interpolant not only a function on some nodes $x_i, i = 0, \ldots, n$ but also the values of its first *m* derivatives.

A good proposal for a solution to the problem could be given by interpolants based on rational interpolant since they may take advantage of the well-known stability of the interpolants for the classical problem of interpolation (see e.g. [20]). In fact, some recent works go in that direction, as the family of Hermite barycentric interpolants introduced in [16], the family of interpolants in [9] or the combination of rational functions and six-point Shepard basis functions introduced in [13]. Multivariate Hermite interpolants are as well of great interest and widely studied as it shows the recently introduced interpolant in [12].

A good stable interpolant that achieves a fast and accurate solution is preferable since it may be useful for the many applications where an Hermite interpolant is used, such as in Treecode algorithm [17] or constructing curves and surfaces for images [18] or approximating the reliability of a hammock network of arbitrary size [14].

In the present paper we focus on the method introduced by Cirillo and Hormann in [9] which allows to construct an effective Hermite interpolant in an iterative way starting from the basis $b_i(x)$ of the renowned Floater-Hormann family of interpolants, i.e., the basis functions that, given $d \in \mathbb{N}$ and x_0, \ldots, x_n interpolation nodes, are constructed as

$$b_i(x) = \frac{w_i}{x - x_i} \Big/ \sum_{j=0}^n \frac{w_j}{x - x_k}, i = 0, \dots, n,$$

with

$$w_i = (-1)^{i+d} \sum_{j=\max(0,i-d)}^{i,n-d} \prod_{k=j,k\neq i}^{j+d} \frac{1}{|x_i - x_k|}, i = 0, \dots, n$$

Therefore, they considered the functions

$$b_{i,j}(x) = \frac{1}{j!}(x - x_i)^j b_i(x)^{j+1}$$

and they used them to construct an interpolant that satisfies the Hermite conditions.

It has been investigated also the effectiveness of such construction for FHinterpolants by studying in [10] the Lebesgue constant at equidistant nodes and giving an estimate of the interpolation error in [11].

In more specific, we are going to generalise the iterative Hermite construction of [9] and use this generalisation to introduce a barycentric rational trigonometric Hermite interpolant.

The paper is construct as follows: in Section 2 is presented a generalisation of the iterative approach presented by Cirillo and Hormann. In Section 3 is computed analytically the differentiation matrix of the Hermite basis constructed starting by the basis functions of Berrut's trigonometric interpolant; this allows a fast construction and computation of the barycentric rational trigonometric Hermite interpolant. In Section 4 various numerical examples are presented and analysed. Finally, we conclude in Section 5.

2 A general iterative approach

Let us consider a set $\Omega \subset \mathbb{R}$, some nodes $x_i \in \Omega$, i = 0, ..., n and consider some basis functions b_i for an interpolant on Ω . We define, therefore, the following functions,

$$b_{i,j}(x) := \frac{1}{j!} d_i(x)^j b_i(x)^{j+1}$$

for a smooth function $d_i(x)$ that vanishes in x_i and not in x_j for $j \neq i$ and such that $d'_i(x_i) = 1$.

These functions will be the foundation to construct the Hermite interpolants, in fact we get that

Lemma 1. Let $\mathcal{X}_n = \{x_0, \ldots, x_n\} \subset \Omega \subset \mathbb{R}$ be some ordered nodes and let $d_i(x)$ be a function that vanishes in x_j only if j = i and such that $d'_i(x_i) = 1$. Let $b_i(x), i = 0, \ldots, n$ be a basis of some space \mathcal{V} that satisfy the Lagrange property at the nodes. Then, the functions $b_{i,j}$ defined as

$$b_{i,j}(x) := \frac{1}{j!} d_i(x)^j b_i(x)^{j+1}$$
(2)

satisfy

$$b_{i,j}^{(k)}(x_{\ell}) = \begin{cases} 0, & \text{if } k < j, \\ \delta_{i,\ell}, & \text{if } k = j. \end{cases}$$
(3)

Proof. For j = 0 the statement follows directly from the Lagrange property of $b_i(x)$. Let us now consider j > 0 and prove the statement by induction on j. Let

$$c_i(x) = d_i(x)b_i(x);$$

then, it is clear that

$$b_{i,j}(x) = \frac{1}{j}c_i(x)b_{i,j-1}(x).$$

Using the Leibniz rule, we get

$$b_{i,j}^{(k)}(x) = \frac{1}{j} \sum_{s=0}^{k} \binom{k}{s} c_i(x)^{(k-s)}(x) b_{i,j-1}^{(s)}(x).$$

Therefore, since $c_i(x_p) = 0$ for p = 0, ..., n and, furthermore, we have that $b_{i,j-1}^{(s)}(x_p) = 0$ for s < j-1 because of the induction hypothesis, we get

$$b_{i,j}^{(k)}(x_p) = \frac{1}{j} \sum_{s=j-1}^{k-1} \binom{k}{s} c_i^{(k-s)}(x_p) b_{i,j-1}^{(s)}(x_p).$$

Let us observe that the sum is empty in the case when k < j, whereas when k = j we obtain

$$b_{i,j}^{(j)}(x_p) = \frac{1}{j} \binom{j}{j-1} c'_i(x_p) b_{i,j-1}^{(j-1)}(x_p).$$

Finally, we notice that $b_{i,j}^{(j)}(x_p) = 0$ if $p \neq i$ and in the case when p = i we have that the product is one since $c'_i(x_i) = 1$ by construction.

Remark 1. Given a function $h_i(x)$ vanishing only in x_i and with a non-zero derivative in x_i , we remark that we can always construct a function that satisfies the assumption of Lemma 1 by setting

$$d_i(x) \coloneqq \frac{h_i(x)}{h'_i(x_i)}.$$

Once we have the functions $b_{i,j}$ we can construct the Hermite interpolant starting from

$$r_0(x) = \sum_{i=0}^{n} b_{i,0}(x) f_i^{(0)},$$

which corresponds to the classical interpolant since $b_{i,0} = b_i$. For j > 0, since we have that normally

$$r_{j-1}^{(j)}(x_\ell) \neq f_\ell^{(j)},$$

we introduce, at each step, a correction function q_j which we add to the previous interpolants, in order to construct a new one and that allows us to interpolate correctly the derivatives up to the *j*-th, i.e.,

$$r_j(x) = r_{j-1}(x) + q_j(x), \quad j = 1, \dots, m.$$
 (4)

We define the correction term as

$$q_j(x) = \sum_{i=0}^n b_{i,j}(x) \left(f_i^{(j)} - r_{j-1}^{(j)}(x_i) \right), \quad j = 1, \dots, m.$$
(5)

The resulting function r_m , which we may write shortly as

$$r_m(x) := \sum_{j=0}^m \sum_{i=0}^n b_{i,j}(x) g_i^{(j)}$$

for proper $g_i^{(j)}$, construct via the iterative approach described above, thanks to Lemma 1, will satisfy the Hermite conditions. In fact, we have the following theorem.

Theorem 1. Let $\mathcal{X}_n = \{x_0, \ldots, x_n\} \subset \Omega \subset \mathbb{R}$ be some ordered nodes and let $d_i(x)$ be a function that vanishes in x_j only if j = i and such that $d'_i(x_i) = 1$. Let $b_i(x)$, $i = 0, \ldots, n$ be a basis of some space \mathcal{V} that satisfy the Lagrange property at the nodes and let $b_{i,j}$ be defined as (2). Then, the function r_m , defined as

$$r_m(x) \coloneqq \sum_{j=0}^m \sum_{i=0}^n b_{i,j}(x) g_i^{(j)},$$
(6)

with

$$g_i^{(j)} = \begin{cases} f_i^{(0)} & \text{if } j = 0\\ f_i^{(j)} - r_{j-1}^{(j)}(x_i) & \text{if } j > 0, \end{cases}$$
(7)

satisfies the Hermite conditions (1)

Proof. It is straightforward that,

$$r_m^{(k)}(x_\ell) = \sum_{i=0}^n \sum_{j=0}^m b_{i,j}^{(k)}(x)_{|_{x=x_\ell}} g_{i,j},$$

which due to Lemma 1 is

$$\begin{aligned} r_m^{(k)}(x_\ell) &= \sum_{i=0}^n \sum_{j=0}^k b_{i,j}^{(k)}(x)_{|_{x=x_\ell}} g_{i,j} \\ &= g_{\ell,k} + \sum_{i=0}^n \sum_{j=0}^{k-1} b_{i,j}^{(k)}(x)_{|_{x=x_\ell}} g_{i,j} = g_{\ell,k} + r_{k-1}^{(k)}(x_\ell) \\ &= f_\ell^{(k)}. \end{aligned}$$

3 A barycentric rational trigonometric interpolant

Barycentric rational trigonometric interpolants are suitable for approximating periodic functions and many of them have been introduced in the last years.

The barycentric form of the classical interpolant at equidistant nodes has been introduced by Henrici [15], and, later, Berrut [5] proposed to use the same function as the Henrici's interpolant but for other ordered nodes.

More recently, Bandiziol and De Marchi [4] proposed a barycentric rational trigonometric interpolant constructed similarly as the well-known Floater-Hormann interpolant. Moreover, Baddoo [1] introduced a trigonometric equivalent of the renowned algorithm AAA, the AAAtrig, which construct a barycentric trigonometric rational approximant selecting the nodes progressively via a greedy algorithm.

In this section, we focus into Berrut's trigonometric interpolant [5]. Let us consider *n* ordered nodes $0 \leq \theta_0 < \cdots < \theta_{n-1} < 2\pi$, then, we define the interpolant as

$$T_n(\theta) = \frac{\sum_{i=0}^{n-1} (-1)^i \operatorname{cst}\left(\frac{\theta - \theta_i}{2}\right) f(\theta_i)}{\sum_{i=0}^{n-1} (-1)^i \operatorname{cst}\left(\frac{\theta - \theta_i}{2}\right)},\tag{8}$$

where the function cst is

$$\operatorname{cst}(\theta) = \begin{cases} \operatorname{csc}(\theta), & \text{if } n \text{ is odd,} \\ \operatorname{ctg}(\theta), & \text{if } n \text{ is even.} \end{cases}$$
(9)

It has no poles in $[0, 2\pi)$ and enjoys a logarithmic growth of its Lebesgue constant for a wide family of interpolation nodes [8]. This class of nodes includes also the images of equidistant nodes via a conformal map, such as those in [2, 7]. As said earlier, it corresponds to the classical trigonometric interpolant when the nodes are equidistant [15] and, therefore, it enjoys its properties, such as the exponential convergence. In addition, the interpolant converges exponentially also when the nodes are images of equidistant nodes via a conformal map [3] and it has been used effectively in [6] to interpolate functions on two-dimensional starlike domains.

In order to construct the Hermite interpolant and to retain the periodic behaviour, we may choose

$$d_i(\theta) = 2\sin\left(\frac{\theta - \theta_i}{2}\right),\tag{10}$$

which clearly satisfies the conditions we need to define the functions $b_{i,j}$.

In particular, we get

$$\begin{split} b_{i,j}(\theta) &\coloneqq \frac{2^j}{j!} \sin^j \left(\frac{\theta - \theta_i}{2}\right) (-1)^{ij+i} \frac{\operatorname{cst}^{j+1} \left(\frac{\theta - \theta_i}{2}\right)}{\left(\sum_{k=0}^{n-1} (-1)^k \operatorname{cst} \left(\frac{\theta - \theta_k}{2}\right)\right)^{j+1}} \\ &= \frac{2^j}{j!} (-1)^{ij+i} \frac{\operatorname{cst} \left(\frac{\theta - \theta_i}{2}\right)}{\left(\sum_{k=0}^{n-1} (-1)^k \operatorname{cst} \left(\frac{\theta - \theta_k}{2}\right)\right)^{j+1}} \cdot \begin{cases} 1, & \text{if } n \text{ is odd,} \\ \cos^j \left(\frac{x - x_i}{2}\right), & \text{if } n \text{ is even,} \end{cases} \\ &= \frac{2^j (-1)^{i(j+1)}}{j!} \operatorname{cst} \left(\frac{\theta - \theta_i}{2}\right) \begin{cases} \frac{1}{\left(\sum_k \frac{(-1)^k}{\sin((\theta - \theta_k)/2)}\right)^{j+1}} & \text{if } n \text{ is odd,} \end{cases} \\ &\frac{\cos^j \left(\frac{\theta - \theta_i}{2}\right)}{\left(\sum_k \frac{(-1)^k}{\sin((\theta - \theta_k)/2)}\right)^{j+1}} & \text{if } n \text{ is even.} \end{cases} \end{split}$$

This formulation will allow us to calculate analytically the differential matrix of the Hermite interpolant

$$t_m(\theta) = \sum_{i=0}^{n-1} \sum_{j=0}^m b_{i,j}(\theta) g_{i,j},$$
(12)

which will be useful for a fast computation of the interpolant.

Theorem 2. Let us consider the basis functions (11), then when s > j the differential matrix, whose elements are $(D_j^s)_{ik} = b_{k,j}^{(s)}(\theta_i)$, is defined as

$$(D_{j}^{s})_{ik} = \frac{1}{\sin\left(\frac{\theta_{i}-\theta_{k}}{2}\right)} \left((-1)^{(k-i)(j+1)} \sum_{q=0}^{\lfloor\frac{s-j-1}{2}\rfloor} {s \choose 2q+1} \frac{(-1)^{q}}{2^{2q+1}} (D_{j}^{s-2q-1})_{ii} - \sum_{q=j}^{s-1} {s \choose s-q} \left(\sin\left(\frac{\theta-\theta_{k}}{2}\right) \right)_{|_{\theta=\theta_{i}}}^{(s-q)} (D_{j}^{q})_{ik} \right)$$
(13)

when n is odd and

$$(D_{j}^{s})_{ik} = \frac{1}{\tan\left(\frac{\theta_{i}-\theta_{k}}{2}\right)} \left((-1)^{(k-i)(j+1)} \sum_{q=0}^{\lfloor \frac{s-j}{2} \rfloor} {s \choose 2q+1} 2^{-2q-1} \cdot \left(\sum_{j=0}^{q} a(2q+1,2j+1) \right) (D_{j}^{s-2q-1})_{ii} - \sum_{q=j}^{s-1} {s \choose s-q} \left(\tan\left(\frac{\theta-\theta_{k}}{2}\right) \right)_{|_{\theta=\theta_{i}}}^{(s-q)} (D_{j}^{q})_{ik} \right).$$
(14)

when n is even, with

$$a(p,q) = (-1)^{\frac{1}{2}(p-\frac{3+(-1)^{p}}{2})} 2 \sum_{j=0}^{\frac{p-q-1}{2}} (-1)^{\frac{p-q-1}{2}-j} {p+1 \choose j} \left(\frac{p-q-1}{2}-j+1\right)^{p}.$$

Proof. Therefore, we can write the basis as

$$b_{k,j}(\theta) := u_{k,j}\alpha(\theta)\operatorname{cst}\left(\frac{\theta - \theta_k}{2}\right) \tag{15}$$

where $u_{k,j} \coloneqq \frac{2^{j}(-1)^{k(j+1)}}{j!}$ and, if n is odd,

$$\alpha(\theta) := \frac{1}{\left(\sum_{i=0}^{n-1} (-1)^i \operatorname{cst}((\theta - \theta_i)/2)\right)^{j+1}}$$

and, if n is even,

$$\alpha(\theta) := \frac{\cos^j(\frac{\theta - \theta_k}{2})}{\left(\sum_{i=0}^{n-1} (-1)^i \operatorname{cst}\left((\theta - \theta_i)/2\right)\right)^{j+1}}$$

In this way we can compute the differentiation matrices as done for the classical barycentric trigonometric interpolant in [3], which will be useful to compute the values of the derivatives of the interpolant r_m at the nodes in the previous iterations.

If n is odd, considering equation (15), differentiating both side and evaluating at the points x_i , we get that

$$\begin{aligned} \left(\alpha(\theta)\right)_{|\theta=\theta_{k}}^{(s)} &= \frac{1}{u_{k,j}} \sum_{q=0}^{s} {\binom{s}{q}} \left(\sin\left(\frac{\theta-\theta_{k}}{2}\right)\right)^{(q)} |_{\theta=\theta_{k}} \left(b_{k,j}(\theta)\right)^{(s-q)} |_{\theta=\theta_{k}} \\ &= \frac{1}{u_{k,j}} \sum_{q=0}^{s} {\binom{s}{q}} \left(\sin\left(\frac{\theta-\theta_{k}}{2}\right)\right)^{(q)} |_{\theta=\theta_{k}} (D_{j}^{s-q})_{kk} \\ &= \frac{1}{u_{k,j}} \sum_{q=1}^{s} {\binom{s}{q}} \left(\sin\left(\frac{\theta-\theta_{k}}{2}\right)\right)^{(q)} |_{\theta=\theta_{k}} (D_{j}^{s-q})_{kk}, \end{aligned}$$
(16)

where we defined $(D_j^s)_{ik} = b_{k,j}^{(s)}(\theta_i)$. Moreover, by evaluating the derivative in a different nodes we have

$$\left(\alpha(\theta)\right)^{(s)}\Big|_{\theta=\theta_i} = \frac{1}{u_{k,j}} \sum_{q=0}^s \binom{s}{q} \left(\sin\left(\frac{\theta-\theta_k}{2}\right)\right)^{(q)}\Big|_{\theta=\theta_i} \left(b_{k,j}(\theta)\right)^{(s-q)}\Big|_{\theta=\theta_i}$$
$$= \frac{1}{u_{k,j}} \sum_{q=0}^s \binom{s}{q} \left(\sin\left(\frac{\theta-\theta_k}{2}\right)\right)^{(q)}\Big|_{\theta=\theta_i} (D_j^{s-q})_{ik}.$$
(17)

In (17) we can isolate the term q = 0 of the sum, that gives,

$$\left(\alpha(\theta)\right)^{(s)}\Big|_{\theta=\theta_i} = \frac{1}{u_{k,j}} \sum_{q=1}^s \binom{s}{q} \left(\sin\left(\frac{\theta-\theta_k}{2}\right)\right)^{(q)}\Big|_{\theta=\theta_i} (D_j^{s-q})_{ik} + \frac{1}{u_{k,j}} \left(\sin\left(\frac{\theta_i-\theta_k}{2}\right)\right) (D_j^s)_{ik},$$
(18)

which together with (16) gives us the following recursive formula for the differentiation matrix

$$(D_{j}^{s})_{ik} = \frac{1}{\sin\left(\frac{\theta_{i}-\theta_{k}}{2}\right)} \left(u_{k,j} \left(\alpha(\theta)\right)^{(s)} \Big|_{\theta=\theta_{i}} -\sum_{q=1}^{s} {s \choose q} \left(\sin\left(\frac{\theta-\theta_{k}}{2}\right) \right)^{(q)} \Big|_{\theta=\theta_{i}} (D_{j}^{s-q})_{ik} \right)$$
(19)
$$= \frac{1}{\sin\left(\frac{\theta_{i}-\theta_{k}}{2}\right)} \left(\frac{u_{k,j}}{u_{i,j}} \sum_{q=1}^{s} {s \choose q} \left(\sin\left(\frac{\theta-\theta_{i}}{2}\right) \right)^{(q)} \Big|_{\theta=\theta_{i}} (D_{j}^{s-q})_{ii} -\sum_{q=1}^{s} {s \choose q} \left(\sin\left(\frac{\theta-\theta_{k}}{2}\right) \right)^{(q)} \Big|_{\theta=\theta_{i}} (D_{j}^{s-q})_{ik} \right).$$
(20)

Notice that since the terms $b_{i,j}$ satisfy (3), D_j^j corresponds to the identity matrix and $D_j^s = 0$ when s < j. Then, the formula becomes

$$(D_j^s)_{ik} = \frac{1}{\sin\left(\frac{\theta_i - \theta_k}{2}\right)} \left(\frac{u_{k,j}}{u_{i,j}} \sum_{q=j}^{s-1} {s \choose s-q} \left(\sin\left(\frac{\theta - \theta_i}{2}\right)\right)^{(s-q)} \Big|_{\theta = \theta_i} (D_j^q)_{ii} - \sum_{q=j}^{s-1} {s \choose s-q} \left(\sin\left(\frac{\theta - \theta_k}{2}\right)\right)^{(s-q)} \Big|_{\theta = \theta_i} (D_j^q)_{ik}\right).$$
(21)

Similarly, when n is even, we have that

$$\left(\alpha(\theta)\right)^{(s)}\Big|_{\theta=\theta_k} = \frac{1}{u_{k,j}} \sum_{q=1}^s \binom{s}{q} \left(\tan\left(\frac{\theta-\theta_k}{2}\right)\right)^{(q)}\Big|_{\theta=\theta_k} (D_j^{s-q})_{kk}$$
(22)

$$\left(\alpha(\theta)\right)^{(s)}\Big|_{\theta=\theta_i} = \frac{1}{u_{k,j}} \sum_{q=0}^s \binom{s}{q} \left(\tan\left(\frac{\theta-\theta_k}{2}\right)\right)^{(q)}\Big|_{\theta=\theta_i} (D_j^{s-q})_{ik}.$$
 (23)

Hence,

$$(D_j^s)_{ik} = \frac{1}{\tan\left(\frac{\theta_i - \theta_k}{2}\right)} \left(\frac{u_{k,j}}{u_{i,j}} \sum_{q=j}^{s-1} \binom{s}{s-q} \left(\tan\left(\frac{\theta - \theta_i}{2}\right)\right)^{(s-q)} \Big|_{\theta = \theta_i} (D_j^q)_{ii} - \sum_{q=j}^{s-1} \binom{s}{s-q} \left(\tan\left(\frac{\theta - \theta_k}{2}\right)\right)^{(s-q)} \Big|_{\theta = \theta_i} (D_j^q)_{ik}\right).$$
(24)

Moreover, since

$$\left(\sin\left(\frac{\theta-\theta_k}{2}\right)\right)^{(q)} = \begin{cases} \frac{(-1)^{\frac{q}{2}}}{2^q} \sin\left(\frac{\theta-\theta_k}{2}\right), & \text{if } q \text{ even} \\ \frac{(-1)^{\frac{q-1}{2}}}{2^q} \cos\left(\frac{\theta-\theta_k}{2}\right), & \text{if } q \text{ odd} \end{cases}$$

and since the *j*-th derivative of $\tan((\theta - \theta_i)/2)$ computed in θ_i is a given by (see [19])

$$\left(\left(\tan\left(\frac{\theta - \theta_i}{2}\right) \right)^{(n)} \Big|_{\theta = \theta_i} = \begin{cases} 0, & \text{if } n = 2k, \\ 2^{-n} \sum_{j=0}^{k-1} a(2k-1, 2j+1), & \text{if } n = 2k-1, \end{cases} \right)$$

for some $k\in\mathbb{N}$ and with

$$a(p,q) = (-1)^{\frac{1}{2}(p-\frac{3+(-1)^p}{2})} 2 \sum_{j=0}^{\frac{p-q-1}{2}} (-1)^{\frac{p-q-1}{2}-j} {p+1 \choose j} \left(\frac{p-q-1}{2}-j+1\right)^p$$

we can simplify equation (21) to

$$(D_{j}^{s})_{ik} = \frac{1}{\sin\left(\frac{\theta_{i}-\theta_{k}}{2}\right)} \left((-1)^{(k-i)(j+1)} \sum_{q=0}^{\lfloor\frac{s-j-1}{2}\rfloor} \left(\frac{s}{2q+1}\right) \frac{(-1)^{q}}{2^{2q+1}} (D_{j}^{s-2q-1})_{ii} - \sum_{q=j}^{s-1} \left(\frac{s}{s-q}\right) \left(\sin\left(\frac{\theta-\theta_{k}}{2}\right)\right)_{|_{\theta=\theta_{i}}}^{(s-q)} (D_{j}^{q})_{ik} \right)$$
(25)

and equation (24) to

$$(D_{j}^{s})_{ik} = \frac{1}{\tan\left(\frac{\theta_{i}-\theta_{k}}{2}\right)} \left((-1)^{(k-i)(j+1)} \sum_{q=0}^{\lfloor \frac{s-j}{2} \rfloor} {\binom{s}{2q+1}} 2^{-2q-1} \cdot \left(\sum_{j=0}^{q} a(2q+1,2j+1) \right) (D_{j}^{s-2q-1})_{ii} - \sum_{q=j}^{s-1} {\binom{s}{s-q}} \left(\tan\left(\frac{\theta-\theta_{k}}{2}\right) \right)_{|_{\theta=\theta_{i}}}^{(s-q)} (D_{j}^{q})_{ik} \right).$$
(26)

We remark that, when we compute the matrix for s = j + 1, the elements simplify to the following

$$(D_j^{j+1})_{ik} = \frac{(-1)^{(j+1)(k-i)}(j+1)}{2} \operatorname{cst}\left(\frac{\theta_i - \theta_k}{2}\right), \qquad k \neq i,$$
(27)

which for j = 0 corresponds to the values in [3]; for the diagonal elements, we use the relation $\sum_{i=0}^{n-1} b_{i,j}^{(j)}(\theta) = 1$, since we want to reproduce the constants for the *j*-th derivate. This implies that

$$\sum_{i=0}^{n-1} b_{i,j}^{(j+1)}(\theta) = 0,$$

so that we obtain

$$(D_j^{j+1})_{ii} = -\sum_{\substack{k=0\\k\neq i}}^{n-1} (D_j^{j+1})_{ik}.$$

Moreover, since for the construction of the Hermite interpolant (12), we need to compute $r_{j-1}^{(s)}(x_i)$ and therefore D_j^s , in the next section, we decided to proceed by computing the differentiation matrices of higher order as

$$D_j^s = (D_j^{j+1})^{s-j} \qquad s \ge j+1.$$
(28)

as done for the polynomial case in [21].



Figure 1: The functions f_1 (left) and f_2 (right) in red dashed and in blue line the interpolant t_3 at 7 equidistant nodes (left) and the interpolant t_2 at 6 equidistant nodes (right). Interpolation nodes are labeled in green.

4 Numerical Experiments

Here we present some numerical tests, some Matlab demos to reproduce the experiments are available at

https://github.com/gelefant/TrigonometricHermite

We considered the following periodic test functions in $[0, 2\pi)$,

$$f_1(\theta) = e^{2\sin(\theta) + \cos(\theta)},$$

$$f_2(\theta) = \cos(3\theta) + \log(\cos(\theta) + 1.5),$$

and we interpolated them by means of the interpolant (12) with the basis presented in Section 3 with N equidistant nodes.

We tested the interpolants which satisfy the Hermite conditions from one to four derivatives. In Figure 1 we display both test functions together with two interpolants, one satisfying the Hermite conditions up to the second derivative and one satisfying the conditions up to the third derivative.

For a fixed m, we compute, the absolute error between the function and the interpolant with N nodes $\operatorname{err}_N = \max_{x \in [0,2\pi)} |f(x) - t_m(x)|$ and we can notice that the interpolant, in the case of conditions up to the second derivative retains the exponential convergence of the classical interpolant (8) at equidistant nodes (see Figures 2 and 3), on the other hand, when we have conditions on the third or the fourth derivatives, the convergence slows down (see Figures 4 and 5). We can numerically estimate the convergence as $\mathcal{O}(N^{-3})$ as we can see in Table 1 where we computed $-\log_2(\frac{\operatorname{err}_2N}{\operatorname{err}_N})$ for N = 5, 10, 20, 40, 80, 160. In order to test the interpolant with conformally shifted nodes and explore

In order to test the interpolant with conformally shifted nodes and explore the convergence to functions with fronts, we tested the interpolant moreover with the function

$$f_3(\theta) = \tanh(50\cos(x + \pi/3));$$

which has two fronts located in $\theta_1 = \pi/6$ and $\theta_2 = 7\pi/6$. For this reason we compared the performances of the interpolant with equidistant nodes and

| | Ν | 5 | 10 | 20 | 40 | 80 | 160 |
|-------|-------|------|------|------|------|------|------|
| t_3 | f_1 | 3.70 | 2.59 | 3.03 | 2.97 | 3.00 | 3.00 |
| | f_2 | 4.18 | 2.96 | 2.89 | 2.97 | 3.01 | 2.99 |
| t_4 | f_1 | 4.37 | 2.93 | 2.93 | 2.95 | 2.87 | 2.96 |
| | f_2 | 5.50 | 2.84 | 2.94 | 2.99 | 2.92 | 2.89 |

Table 1: Estimate of the rate of convergence for t_3 and t_4 with both test functions



Figure 2: Convergence of t_1 for f_1 (left) and f_2 (right) and N = 5, 10, 20, 40.



Figure 3: Convergence of t_2 for f_1 (left) and f_2 (right) and N = 5, 10, 20, 40.



Figure 4: Convergence of t_3 for f_1 (left) and f_2 (right) and N = 5, 10, 20, 40, 80, 160, 320.



Figure 5: Convergence of t_4 for f_1 (left) and f_2 (right) and N = 5, 10, 20, 40, 80, 160, 320.



Figure 6: Error of function f_3 with equidistant nodes (blue line) and conformally shifted nodes (red line) by using (a) t_1 , (b) t_2 , (c) t_3 , (d) t_4 , with $N = 5, 10, 20, \ldots, 1280$

conformally shifted nodes, via the conformal map introduced in [7] in the generalisation for two fronts and with the same density parameter for both fronts set as $\alpha = 0.85$.

As we can see in Figures 6, the advantage of clustering the nodes in the locations of the fronts becomes less important as we consider more derivatives. In fact, it speeds up a lot the convergence in the case of one derivative but, then, once we add the conditions on more derivatives, clustering the nodes decreases the speed up until it even slows the convergence when we have conditions up to the fourth derivative.

We present another numerical experiment which is the reconstruction of a closed simple curve, given some points, as might be a figure or a sketch. The two dimensions are reconstructed separately and each coordinates is consider as a periodic function. Therefore, in order to enjoy the fast exponential convergence, the data are considered as the values of a function at equidistant nodes. Then, we approximate the first derivatives by using the coordinates of the difference quotients of the points.

We did therefore, the reconstruction of the sketch in Figure 7a and starting by extracting 362 points (which are the red dots in 7b), we used the values of the abscissae and ordinates as datae of two periodic function with those values at equidistant nodes and which we intend to approximate by the Hermite interpolant t_1 , which is the interpolant with conditions on one derivative. Moreover, since the datae of the real tangent vectors are unknown, we approximate them with the difference quotient vector between the extracted points ordered in anticlockwise sense. The result is in Figure 7b.

In Figure 7(c)-(e) we used a subset of the initial set of nodes: half points, a third and a fifth, respectively.

As we can see by the results in Figures 7, the interpolant performs quite well even by using an approximation of the derivative and by decreasing the number of nodes.

In particular, even by using a fifth of the points initially extracted the reconstruction is still resembling quite a lot the sketch.

5 Conclusion

The present work introduced an iterative scheme to construct an Hermite interpolant starting from a general interpolation basis and in particular, we focused on Berrut's trigonometric interpolant. In order to construct the values of the derivative of the precedent iterative step, we compute analytically the differentation matrix which allows to obtain these values easily as a matrix product.

Our numerical experiments suggest that the produced interpolant retains the exponential convergence of the classical interpolant at equidistant nodes and conformally shifted nodes, once we interpolate the first two derivatives and slows down to a rate of $\mathcal{O}(N^{-3})$, where N is the number of nodes used, when we interpolate also the third or the fourth derivatives.

Moreover, the tests suggests that clustering nodes in the locations of fronts via a conformal map speeds up the convergence in case of one derivative but, instead, could be of negative impact when we need to satisfy conditions also for higher derivatives.

Future work will be aimed into analysing the convergence more deeply, also for other class of nodes, as the one introduced in [7], which includes conformally shifted nodes, and also into analysing the Lebesgue constant of the interpolant.

Acknowledgments. This research has been accomplished within Rete ITaliana di Approssimazione (RITA), the thematic group on "Teoria dell'Approssimazione e Applicazioni" (TAA) of the Italian Mathematical Union and partially supported by GNCS-IN δ AM "Finanziamenti Giovani Ricercatori 2022" (Young Researchers Fund 2022) for the project "An iterative approach for Hermite interpolants", INdAM - GNCS Project cod. CUP E55F22000270001.

References

- P. J. Baddoo. "The AAAtrig algorithm for rational approximation of periodic functions". In: SIAM J. Sci. Comput. 43.5 (2021), A3372–A3392.
- P. J. Baddoo and D. G. Crowdy. "Periodic Schwarz-Christoffel mappings with multiple boundaries per period". In: *Proc. A*. 475.2228 (2019), pp. 20190225, 20.



Figure 7: Reconstruction of the sketch (a), by using 362 nodes (b), 181 nodes (c), 121 nodes (d) and 73 nodes (e).

- [3] R. Baltensperger. "Some results on linear rational trigonometric interpolation". In: Comput. Math. Appl. 43.6-7 (2002), pp. 737–746.
- [4] C. Bandiziol and S. De Marchi. "On the Lebesgue constant of the trigonometric Floater-Hormann rational interpolant at equally spaced nodes". In: *Dolomites Res. Notes Approx.* 12 (2019), pp. 51–67.
- [5] J.-P. Berrut. "Rational functions for guaranteed and experimentally wellconditioned global interpolation". In: *Comput. Math. Appl.* 15.1 (1988), pp. 1–16.
- [6] J.-P. Berrut and G. Elefante. "A linear barycentric rational interpolant on starlike domains". In: *Electron. Trans. Numer. Anal.* 55 (2022), pp. 726– 743.
- [7] J.-P. Berrut and G. Elefante. "A periodic map for linear barycentric rational trigonometric interpolation". In: Appl. Math. Comput. 371 (2020), pp. 124924, 8.
- [8] J.-P. Berrut and G. Elefante. "Bounding the Lebesgue constant for a barycentric rational trigonometric interpolant at periodic well-spaced nodes". In: J. Comput. Appl. Math. 398 (2021), pp. 113664, 11.
- [9] E. Cirillo and K. Hormann. "An iterative approach to barycentric rational Hermite interpolation". In: *Numer. Math.* 140.4 (2018), pp. 939–962.
- [10] E. Cirillo and K. Hormann. "On the Lebesgue constant of barycentric rational Hermite interpolants at equidistant nodes". In: J. Comput. Appl. Math. 349 (2019), pp. 292–301.
- [11] E. Cirillo, K. Hormann, and J. Sidon. "Convergence rates of a Hermite generalization of Floater-Hormann interpolants". In: J. Comput. Appl. Math. 371 (2020), pp. 112624, 9.
- [12] F. Dell'Accio et al. "Fast and accurate scattered Hermite interpolation by triangular Shepard operators". In: J. Comput. Appl. Math. 382 (2021), Paper No. 113092, 15.
- [13] F. Dell'Accio et al. "Rational Hermite interpolation on six-tuples and scattered data". In: Appl. Math. Comput. 386 (2020), pp. 125452, 11.
- [14] L. Dăuş and M. Jianu. "Full Hermite interpolation of the reliability of a hammock network". In: Appl. Anal. Discrete Math. 14.1 (2020), pp. 198– 220.
- [15] P. Henrici. "Barycentric formulas for interpolating trigonometric polynomials and their conjugates". In: Numer. Math. 33.2 (1979), pp. 225–234.
- [16] K. Jing et al. "A convergent family of linear Hermite barycentric rational interpolants". In: J. Math. Res. Appl. 40.6 (2020), pp. 628–646.
- [17] R. Krasny and L. Wang. "A treecode based on barycentric Hermite interpolation for electrostatic particle interactions". In: Comput. Math. Biophys. 7 (2019), pp. 73–84.
- [18] F. Oumellal and A. Lamnii. "Curve and Surface Construction Using Hermite Trigonometric Interpolant". In: *Mathematical and Computational Applications* 26.1 (2021).
- [19] F. Qi. "Derivatives of tangent function and tangent numbers". In: Appl. Math. Comput. 268 (2015), pp. 844–858.

- [20] C. Schneider and W. Werner. "Hermite interpolation: the barycentric approach". In: Computing 46.1 (1991), pp. 35–51.
- B. D. Welfert. "Generation of pseudospectral differentiation matrices. I". In: SIAM J. Numer. Anal. 34.4 (1997), pp. 1640–1657.