Reducing smooth functions to normal forms near critical points A.S. Orevkova

s15b3_orevkova@179.ru; Moscow State Univercity, Moscow Center of Fundamental and Applied Mathematics

Abstract

The paper is devoted to "uniform" reduction of smooth functions on 2-manifolds to canonical form near critical points by some coordinate changes in some neighbourhoods of these points. For singularity types E_6 , E_8 and A_n , we explicitly construct such coordinate changes and estimate from below (in terms of C^r -norm of the function) the radius of a required neighbourhood.

Keywords: right equivalence of smooth functions, ADE-singularities, normal form of singularities, uniform reducing to normal form. *MSC:* 58K05, 37C15.

1 Introduction

Definition. A smooth function $f = f(u_1, u_2)$ has a singularity type E_k (k = 6, 7, 8) at its critical point $P \in \mathbb{R}^2$ if

- (i) the first and second differentials df(P) = 0, $d^2f(P) = 0$, and the third differential $d^3f(P) \neq 0$ and is a perfect cube;
- (ii) one of the following coefficients of the Taylor series of f at P does not vanish: $f_{y^4}^{(4)}(P)$, $f_{xy^3}^{(4)}(P)$ and $f_{y^5}^{(5)}(P)$, where $(u_1, u_2) \rightarrow (x, y)$ is a linear coordinate change such that $d^3f(P) = (dx)^3$. More specifically: the singularity type is E_6 if $f_{y^4}^{(4)}(P) \neq 0$ (equivalently, there exists a tangent vector $v \in \text{Ker } d^3f(P)$ at P such that $v^4f \neq 0$, where v^4f denotes the fourth derivative of f along the vector v); the singularity type is E_7 if $f_{y^4}^{(4)}(P) = 0$ and $f_{xy^3}^{(4)}(P) \neq 0$; the singularity type is E_8 if $f_{y^4}^{(4)}(P) = f_{xy^3}^{(4)}(P) = 0$ and $f_{y^5}^{(5)}(P) \neq 0$.

From definition of E_k we have $f_{x^a y^b}^{(a+b)}(P) = 0$, $0 < \frac{a}{3} + \frac{2b}{k+2} < 1$, $f_{x^3}''(P) = 6$ (k = 6, 7, 8) and $f_{u^{1+k/2}}^{(1+k/2)}(P) \neq 0$ (k = 6, 8). We will assume that $P = \mathbf{0} = (0, 0)$ in the coordinates x, y.

Assumption 1. For singularities E_k (k = 6, 8), assume that $f_{y^{1+k/2}}^{(1+k/2)}(\mathbf{0}) = \pm (1 + \frac{k}{2})!$.

Theorem 1 (Reducing E_k to normal form [1]). Let a function $f(u_1, u_2)$ have a singularity E_k (k = 6, 7, 8) at a critical point P. Then, in some neighborhood of P, there is a local coordinate system \tilde{x}, \tilde{y} in which the point P is the origin, and the function has the normal form $f = f(P) + \tilde{x}^3 \pm \tilde{y}^{1+k/2}$ for k = 6, 8, $f = f(P) + \tilde{x}^3 + \tilde{x}\tilde{y}^3$ for k = 7.

In [1], the existence of a coordinate change was proved using the Tougeron theorem [13]. In view of this, obtaining a formula for the corresponding coordinate change requires solving the Cauchy problem for a system of ODE's. We construct our coordinate change explicitly, without using the Tougeron theorem.

Lemma 1 ([1]). In the case of a singularity E_k (k = 6, 8), under the hypotheses of Assumption 1, there exist coordinate changes $(x, y) \rightarrow (x_1 = x + d_1y^2, y) \rightarrow (x_1, y_1 = y + d_2x_1) \rightarrow (x_2 = x_1 + d_3y_1^{k/2-1}, y_1) \rightarrow (x_2, y_2 = y_1 + d_4x_2^2)$ where $d_j \in \mathbb{R}$ and $d_1 = 0$ for k = 6, such that $f_{x_1^a y_2^b}^{(a+b)}(\mathbf{0}) = 0$ for all $a, b \in \mathbb{Z}_+$ with a < 3 and $b < 1 + \frac{k}{2}$.

Theorem 2 (Estimating the radius of a neighborhood for the coordinate change). In the case of singularities E_k (k = 6, 8), under the hypotheses of Assumption 1, let $(x, y) \to (x_2, y_2)$ be the coordinate change from Lemma 1. Suppose that, in a neighborhood $U_0 = \{(x_2, y_2) \mid \max(|x_2|, |y_2|) < R_0\}$ of **0**, the following estimates hold: $C_{\alpha\beta} = \sup_{U_0} |f_{x_2^{\alpha}y_2^{\beta}}^{(\alpha+\beta)}(x_2, y_2)| \leq M$ for $(\alpha, \beta) \in \{(0, 5), (1, 4), (3, 1), (3, 2), (3, 3), (4, 0), (4, 1), (4, 2), (4, 3)\}$ if k = 6, for $(\alpha, \beta) \in \{(0, 6), (1, 5), (3, 1), (3, 2), (3, 3), (4, 0), (4, 1), (4, 2), (4, 3)\}$ if k = 8, where $R_0 > 0$, $M \geq 0$. Then, in the neighbourhood $U = \{(x_2, y_2) \mid \max(|x_2|, |y_2|) < R\}$, with $R = \min\{R_0, \frac{2}{M+2}\}$, there is a coordinate change of the form $\phi : (x_2, y_2) \to (\tilde{x} = x_2\sqrt[3]{h(x_2, y_2)}, \tilde{y} = y_2(g(x_2, y_2))^{\frac{2}{2+k}})$ that reduces f to the normal form $f = f(P) + \tilde{x}^3 \pm \tilde{y}^{1+k/2}$ of E_k . In detail:

- (a) the functions $h(\mathbf{x})$ and $g(\mathbf{x})$ are positive in U, thus the change $\phi|_U$ is well-defined and is C^{∞} -smooth;
- (b) $\|\phi'(\boldsymbol{x}) I\| < C < 1$ for all $\boldsymbol{x} \in U$, where $C = \frac{2}{5}$, thus $\phi|_U$ is C^1 -close to the identity;
- (c) the coordinate change $\phi|_U$ is injective and regular, i.e., it is an embedding and det $|\phi'(\boldsymbol{x})| \neq 0$ for all $\boldsymbol{x} \in U$, moreover $\phi(U)$

contains the open disk of radius (1 - C)R centred at **0**.

Our coordinate change $\phi|_U$ from Theorem 2 provides a "uniform" reduction of the function f at a singular point of type E_k , k = 6, 8, to the canonical form $f = f(P) + \tilde{x}^3 \pm \tilde{y}^{1+k/2}$ in the sense that the neighbourhood radius and the coordinate change we constructed in this neighbourhood (as well as all partial derivatives of the coordinate change) continuously depend on the function f and its partial derivatives. A uniform reduction of smooth functions near critical points to a canonical form was known earlier for several singularity types [2, 10–12].

The uniform Morse lemma [3] was applied in [4–7] for studying topology of the spaces of Morse functions on surfaces and decomposition of these spaces into classes of topological equivalence. Our results have similar applications for stydying topology of the spaces of smooth functions [8] and gradient-like flows [9] with prescribed *ADE*-singularities.

The author is grateful to Elena Kudryavtseva for stating the problem and useful discussions. The author is a Fellow of the Theoretical Physics and Mathematics Advancement Foundation "BASIS".

2 Key lemmas

Lemma 2. Let $f_{x^k y^l}^{(k+l)}(\mathbf{0}) = 0$ for all $k, l \in \mathbb{Z}_+$ with k < m and l < n, where $m, n \in \mathbb{N}$. Then the function $f|_U$ has the form $f(x, y) = x^m h(x, y) + y^n g(x, y)$ for some functions $h, g \in C^{\infty}(U)$, where U is a neighborhood of $\mathbf{0}$ in \mathbb{R}^2 .

Proof. We represent f as a function of y with a parameter x. We write down the Taylor formula at the point y = 0 with the remainder term in the integral form:

$$f(x,y) = S_n(x,y) + R_n(x,y) = x^m h(x,y) + y^n g(x,y),$$

$$S_n(x,y) = f(x,0) + f'_y(x,0)y + f''_{y^2}(x,0)\frac{y^2}{2!} + \dots + f^{(n-1)}_{y^{n-1}}(x,0)\frac{y^{n-1}}{(n-1)!} = x^m h(x,y),$$

$$R_n(x,y) = \frac{1}{(n-1)!} \int_0^y f^{(n)}_{y^n}(x,t)(y-t)^{n-1} dt = \frac{y^n}{(n-1)!} \int_0^1 f^{(n)}_{y^n}(x,sy)(1-s)^{n-1} ds = y^n g(x,y)$$

Here each summand of $S_n(x, y)$ is the product of y^i and a smooth function of x, which can be written as $x^m h_i(x)$, for some smooth functions $h_i(x)$. The lemma is proved.

Lemma 3. Let $\phi : U \to \mathbb{R}^n$ be a smooth mapping, where U is a convex open subset of \mathbb{R}^n . Let the differential of ϕ have the form $\phi'(\mathbf{x}) = I + A(\mathbf{x})$, where I is the unit matrix of dimension n, ||A|| < c, 0 < c < 1. Then ϕ is injective and det $|\phi'(\mathbf{x})| \neq 0$ at every point $\mathbf{x} \in U$, i.e., ϕ is a diffeomorphism to its image $\phi(U)$. Moreover, $\langle \phi(\mathbf{x}) - \phi(\mathbf{y}), \mathbf{x} - \mathbf{y} \rangle \geq (1-c)|\mathbf{x} - \mathbf{y}|^2$ for any pair of points $\mathbf{x}, \mathbf{y} \in U$.

Proof. Take any two points $\boldsymbol{a}, \boldsymbol{b} \in U$, $\boldsymbol{a} \neq \boldsymbol{b}$, and consider the mapping ϕ on the segment $[\boldsymbol{a}, \boldsymbol{b}]$. The velocity vector

$$v_t = \frac{d}{dt} (\phi(\boldsymbol{a} + t(\boldsymbol{b} - \boldsymbol{a}))).$$

Let us look at the projection of the velocity vector v_t onto $\boldsymbol{b} - \boldsymbol{a}$:

$$\langle v_t, \mathbf{b} - \mathbf{a} \rangle = \langle \frac{d}{dt} [\phi(\mathbf{a} + t(\mathbf{b} - \mathbf{a})) - \phi(\mathbf{a})], \mathbf{b} - \mathbf{a} \rangle = \langle \frac{d}{dt} \phi(\mathbf{a} + t(\mathbf{b} - \mathbf{a})) \rangle, \mathbf{b} - \mathbf{a} \rangle = \\ = \langle \phi'(\mathbf{a} + t(\mathbf{b} - \mathbf{a}))(\mathbf{b} - \mathbf{a}), \mathbf{b} - \mathbf{a} \rangle = \langle (I + A(\mathbf{a} + t(\mathbf{b} - \mathbf{a}))) \cdot (\mathbf{b} - \mathbf{a}), \mathbf{b} - \mathbf{a} \rangle = \\ = \langle \mathbf{b} - \mathbf{a}, \mathbf{b} - \mathbf{a} \rangle + \langle A(\mathbf{a} + t(\mathbf{b} - \mathbf{a})) \cdot (\mathbf{b} - \mathbf{a}), (\mathbf{b} - \mathbf{a}) \rangle.$$

Let us find an upper bound for the absolute value of the second term. Note that by the Cauchy-Schwarz inequality and by the definition of the matrix norm:

$$\left| \langle A(\boldsymbol{a}+t(\boldsymbol{b}-\boldsymbol{a})) \cdot (\boldsymbol{b}-\boldsymbol{a}), (\boldsymbol{b}-\boldsymbol{a}) \rangle \right| \leq \|A(\boldsymbol{a}+t(\boldsymbol{b}-\boldsymbol{a})) \cdot (\boldsymbol{b}-\boldsymbol{a})\| \cdot \|\boldsymbol{b}-\boldsymbol{a}\| \leq \|A\| \|\boldsymbol{b}-\boldsymbol{a}\|^2.$$

Let us go back to the estimation of $\langle v_t, \boldsymbol{b} - \boldsymbol{a} \rangle$:

$$\langle v_t, \boldsymbol{b} - \boldsymbol{a} \rangle = \langle \boldsymbol{b} - \boldsymbol{a}, \boldsymbol{b} - \boldsymbol{a} \rangle + \langle A(\boldsymbol{a} + t(\boldsymbol{b} - \boldsymbol{a})) \cdot (\boldsymbol{b} - \boldsymbol{a}), \boldsymbol{b} - \boldsymbol{a} \rangle \geq \\ \geq \|\boldsymbol{b} - \boldsymbol{a}\|^2 - |\langle A(\boldsymbol{a} + t(\boldsymbol{b} - \boldsymbol{a})) \cdot (\boldsymbol{b} - \boldsymbol{a}), \boldsymbol{b} - \boldsymbol{a} \rangle| \geq \\ \geq \|\boldsymbol{b} - \boldsymbol{a}\|^2 - \|A\| \|\boldsymbol{b} - \boldsymbol{a}\|^2 > \|\boldsymbol{b} - \boldsymbol{a}\|^2 \cdot (1 - c) > 0,$$

from which det $|\phi'(\boldsymbol{a})| \neq 0$.

By hypothesis ||A|| < c < 1. Let us look at the dot product $\langle \phi(\boldsymbol{b}) - \phi(\boldsymbol{a}), \boldsymbol{b} - \boldsymbol{a} \rangle$. As $\langle \boldsymbol{x}(t), \boldsymbol{a} \rangle'_t = \left[\sum x_i a_i\right]'_t = \sum x'_t a_i = \langle \boldsymbol{x}', \boldsymbol{a} \rangle$, we have

$$\langle \phi(\boldsymbol{b}) - \phi(\boldsymbol{a}), \boldsymbol{b} - \boldsymbol{a} \rangle = \int_0^1 \langle v_t, \boldsymbol{b} - \boldsymbol{a} \rangle \, dt > (1 - c) \int_0^1 \|\boldsymbol{b} - \boldsymbol{a}\|^2 \, dt = (1 - c) \, \|\boldsymbol{b} - \boldsymbol{a}\|^2 > 0.$$

In other words, the injectivity condition is satisfied: $\phi(\mathbf{a}) \neq \phi(\mathbf{b})$ for each point $\mathbf{b} \in U \setminus \{\mathbf{a}\}$. Lemma 3 is proved. **Lemma 4.** The following inequality is true: $|f_{x^ky^l}^{(k+l)}(x,y) - f_{x^ky^l}^{(k+l)}(0,0)| \le R(C_{k+1,l} + C_{k,l+1}) \le 2MR.$

Proof.

$$\begin{split} |f_{x^{k}y^{l}}^{(k+l)}(x,y) - f_{x^{k}y^{l}}^{(k+l)}(0,0)| &= |f_{x^{k}y^{l}}^{(k+l)}(x,y) - f_{x^{k}y^{l}}^{(k+l)}(0,y) - f_{x^{k}y^{l}}^{(k+l)}(0,0) + f_{x^{k}y^{l}}^{(k+l)}(0,y)| \leq \\ &\leq |f_{x^{k}y^{l}}^{(k+l)}(x,y) - f_{x^{k}y^{l}}^{(k+l)}(0,y)| + |f_{x^{k}y^{l}}^{(k+l)}(0,y) - f_{x^{k}y^{l}}^{(k+l)}(0,0)| = \\ &= \left| \int_{0}^{x} f_{x^{k+1}y^{l}}^{(k+l+1)}(t,y) \, dt \right| + \left| \int_{0}^{y} f_{x^{k}y^{l+1}}^{(k+l+1)}(0,t) \, dt \right|. \end{split}$$

The lemma is proved.

3 Proof of Theorem 2

By Lemma 1, after the change $(x, y) \to (x_2, y_2)$, we have $f_{x_2^a y_2^b}^{(a+b)}(\mathbf{0}) = 0$ for all $a, b \in \mathbb{Z}_+$ with a+b>0, a<3 and $b<1+\frac{k}{2}$. By Lemma 2, $f = f(P) + x_2^3h(x_2, y_2) \pm y_2^{1+k/2}g(x_2, y_2)$ for some functions $h, g \in C^{\infty}(U_0)$. This immediately reduces the function to the required normal form $f = f(P) + \tilde{x}^3 \pm \tilde{y}^{1+k/2}$ by the coordinate change $\phi : \mathbb{R}^2_{x_2,y_2} \to \mathbb{R}^2_{\tilde{x},\tilde{y}}$ with $\tilde{x} = x_2 \sqrt[3]{h(x_2,y_2)}, \tilde{y} = y_2(g(x_2,y_2))^{\frac{1}{1+k/2}}$. From our proof of Lemma 2, we obtain explicite formulas for h, g in terms of $S_{1+k/2}$ and $R_{1+k/2}$, namely: $f = \sum_{i=0}^{k/2} f_{y_2^i}^{(i)}(x_2,0) \frac{y_2^i}{i!} + R_{1+k/2}(x_2,y_2) = f(P) + x_2^3h(x_2,y_2) \pm y_2^{1+k/2}g(x_2,y_2) = f(P) + \tilde{x}^3 \pm \tilde{y}^{1+k/2}$.

It remains to apply Lemma 3 to the coordinate transformation ϕ . From the above formulas for the coordinate change ϕ , the bound $C_{\alpha\beta} < M$ and Lemma 4, using and the Taylor expansion formula with a remainder in the Lagrange form or an integral remainder, we will obtain the required bound $\|\phi'(\boldsymbol{x}) - I\| = \|A(\boldsymbol{x})\| < C < 1$ for each point $\boldsymbol{x} \in U$. We will estimate the norm of the matrix in terms of its elements: $\|A\| \leq \sqrt{\sum a_{ij}^2}, \quad A = \{a_{ij}\}_{i,j=1}^n$.

Let us proceed with detailed estimations, separately for the cases of E_6 and E_8 . By abusing notations, we will denote (x_2, y_2) by (x, y).

3.1 The case of E_6

We compute the elements of the Jacobi matrix of ϕ :

$$\frac{\partial \tilde{x}}{\partial x} = h^{\frac{1}{3}} + x \cdot \frac{1}{3} h^{-\frac{2}{3}} \cdot h'_x,\tag{1}$$

$$\frac{\partial \tilde{x}}{\partial y} = x \cdot \frac{1}{3} h^{-\frac{2}{3}} \cdot h'_y,\tag{2}$$

$$\frac{\partial \tilde{y}}{\partial x} = y \cdot \frac{1}{4} g^{-\frac{3}{4}} \cdot \left(\frac{1}{6} \int_{0}^{1} f_{xy^{4}}^{(5)}(x, sy)(1-s)^{3} \, ds\right),\tag{3}$$

$$\frac{\partial \tilde{y}}{\partial y} = g^{\frac{1}{4}} + y \cdot \left(\frac{1}{4}g^{-\frac{3}{4}} \cdot \frac{1}{6}\int_{0}^{1} sf_{y^{5}}^{(5)}(x,sy)(1-s)^{3}\,ds\right).$$
(4)

If M = 0, then $\phi = \text{id}$ and everything is proved. Let further M > 0, and therefore R < 1. By Assumption 1, we have h(0,0) = 1, g(0,0) = 1. Thus, the Jacobi matrix at **0** is the unit matrix I.

By using the above formulas (1)–(4) for Jacobi matrix' elements, let us estimate the elements of the Jacobi matrix $\frac{\partial(\tilde{x}-x,\tilde{y}-y)}{\partial(x,y)}$ and prove item (a) for the case of E_6 .

Step 1. Here we find an upper bound for $|1 - \frac{\partial \tilde{x}}{\partial x}|$. Remind that:

$$\tilde{x} = x\sqrt[3]{h(x,y)} = \sqrt[3]{f(x,0) + yf'_y(x,0) + \frac{y^2}{2}f''_{y^2}(x,0) + \frac{y^3}{6}f'''_{y^3}(x,0)}.$$

Denote $\tilde{h} = \tilde{h}(x, y) = h(x, y) - 1.$

Let us estimate $\tilde{h}(x, y)$. In the expression for h(x, y), we apply the Taylor expansion formula in x with a remainder in the Lagrange form to the coefficients of powers of y:

$$\begin{aligned} |h(x,y)-1| &= \left| \frac{f_{x^3}^{\prime\prime\prime}(0,0)}{6} + x \frac{f_{x^4}^{(4)}(c_0,0)}{24} + \sum_{k=1}^3 \left(\frac{f_{x^3}^{\prime\prime\prime\prime}(c_k,0)}{6} \right)_{y^k}^{(k)} \frac{y^k}{k!} - 1 \right| = \\ &= \left| 1 + x \frac{f_{x^4}^{(4)}(c_0,0)}{24} + y \frac{f_{x^3y}^{(4)}(c_1,0)}{6} + y^2 \frac{f_{x^3y^2}^{(5)}(c_2,0)}{12} + y^3 \frac{f_{x^3y^3}(c_3,0)}{36} - 1 \right| \le \\ &\leq \frac{MR}{12} \left(\frac{5}{2} + R + \frac{R^2}{3} \right) < \frac{5}{24} \cdot \frac{MR}{1-R} \le \frac{5}{24} \cdot \frac{2M}{M+2} \cdot \frac{1}{1-2/(M+2)} = \frac{5}{24} \cdot 2 < \frac{1}{2}. \end{aligned}$$

Hence $h(x, y) \in (0.5, 1.5)$. Therefore $\sqrt[3]{h(x, y)} \in (0.79, 1.21)$.

Item (a) of Theorem 2 is proved for the function h(x, y). Set c = 0.5. Then $|\tilde{h}(x, y)| < c$. By the formula (3) we have

$$\frac{\partial \tilde{x}}{\partial x} = h^{\frac{1}{3}} + x \cdot \frac{1}{3} h^{-\frac{2}{3}} \cdot h'_x = (1 + \tilde{h})^{\frac{1}{3}} + x \cdot \frac{1}{3} (1 + \tilde{h})^{-\frac{2}{3}} \cdot h'_x.$$

Let us estimate $|h'_x|$. We use Taylor' formula with an integral remainder:

$$\begin{split} |h'_{x}| &= \left| \left(\sum_{k=0}^{3} \frac{y^{k}}{k!} \frac{f_{y^{k}}^{(k)}(x,0)}{x^{3}} \right)'_{x} \right| = \left| \left(\sum_{k=0}^{3} \frac{y^{k}}{k!} \frac{1}{2} \int_{0}^{1} f_{y^{k}x^{3}}^{(k+3)}(sx,0)(1-s)^{2} \, ds \right)'_{x} \right| = \\ &= \left| \sum_{k=0}^{3} \frac{y^{k}}{k! \cdot 2} \int_{0}^{1} s f_{y^{k}x^{4}}^{(k+4)}(sx,0)(1-s)^{2} \, ds \right| \le \sum_{k=0}^{3} \frac{R^{k}M}{k! \cdot 2} \int_{0}^{1} (s-2s^{2}+s^{3}) \, ds < \\ &< \frac{M}{24} \left(1+R+\frac{R^{2}}{2}+\frac{R^{3}}{6} \right) < \frac{1}{24} \cdot \frac{M}{1-R} \le \frac{1}{12R}. \end{split}$$

Set $c_x = \frac{1}{12R}$. We obtain a bound for $\left|\frac{\partial \tilde{x}}{\partial x} - 1\right|$ using the estimate of $\sqrt[3]{h}$:

$$\begin{split} \left|\frac{\partial \tilde{x}}{\partial x} - 1\right| &= \left|\sqrt[3]{h} - 1 + x \cdot \frac{1}{3}(1 + \tilde{h})^{-\frac{2}{3}}h'_{x}\right| \le \left|\sqrt[3]{h} - 1\right| + \left|x \cdot \frac{1}{3}(1 + \tilde{h})^{-\frac{2}{3}}h'_{x}\right| < \\ &< 0.21 + \frac{c_{x}R}{3}(1 - c)^{-\frac{2}{3}} = 0.21 + \frac{2^{\frac{2}{3}}}{3 \cdot 12} < 0.21 + 0.05 = 0.26. \end{split}$$

Step 2. Let us estimate $\frac{\partial \tilde{x}}{\partial y}$ from the formula (2). a) First we estimate $|h'_y|$ when $x \neq 0$. We use Taylor' formula with an integral remainder:

$$\begin{split} |h'_{y}| &= \left| (f(x,0) + yf'_{y}(x,0) + \frac{y^{2}}{2}f''_{y^{2}}(x,0) + \frac{y^{3}}{6}f'''_{y^{3}}(x,0))'_{y}/x^{3} \right| = \\ &= \left| (f'_{y}(x,0) + yf''_{y^{2}}(x,0) + \frac{y^{2}}{2}f'''_{y^{3}}(x,0))/x^{3} \right| = \\ &= \left| \frac{1}{2} \int_{0}^{1} \left(f^{(4)}_{x^{3}y}(sx,0) + yf^{(5)}_{x^{3}y^{2}}(sx,0) + \frac{y^{2}}{2}f^{(6)}_{x^{3}y^{3}}(sx,0) \right) (1-s)^{2} ds \right| \le \\ &\leq \frac{M}{6} (1+R+R^{2}) < \frac{1}{6} \cdot \frac{M}{1-R} \le \frac{1}{3R}. \end{split}$$

Set $c_y = \frac{1}{3R}$.

b) In the formula for $|\frac{\partial \tilde{x}}{\partial y}|$ we expand $h^{-\frac{2}{3}}$ in a Taylor series in \tilde{h} and take $c_{\tilde{h}} = c_{\tilde{h}}(x,y) \in$ $[0, \tilde{h}(x, y)]$ from the Taylor-Lagrange formula:

$$\begin{aligned} \left|\frac{\partial \tilde{x}}{\partial y}\right| &= \left|x \cdot \frac{1}{3} \left(1 + \tilde{h}\right)^{-\frac{2}{3}} \cdot h'_{y}\right| = \left|\frac{x}{3} \left(1 - \frac{2}{3}\tilde{h} + \frac{5}{9}(1 + c_{\tilde{h}})^{-\frac{8}{3}}\tilde{h}^{2}\right)h'_{y}\right| \leq \\ &\leq \frac{R}{3} \left(1 + \frac{2}{3}c + \frac{5}{9}(1 + c_{\tilde{h}})^{-\frac{8}{3}}c^{2}\right)c_{y} < \frac{c_{y}R}{3} \left(1 + \frac{2}{3}c + \frac{5}{9}(1 - c)^{-\frac{8}{3}}c^{2}\right) = \\ &= \frac{1}{9} \left(1 + \frac{1}{3} + \frac{5}{9}2^{\frac{8}{3}}/4\right) < 0.25. \end{aligned}$$

Step 3. Let us estimate $\frac{\partial \tilde{y}}{\partial x}$ from the formula (3). Let us first estimate separately the factors of this expression.

a) Auxiliary Assessment. By Lemma 4, we have

$$\left| \int_{0}^{1} \left(f_{y^{4}}^{(4)}(x,sy) - f_{y^{4}}^{(4)}(0,0) \right) (1-s)^{3} \, ds \right| \leq \int_{0}^{1} 2MR(1-s)^{3} \, ds = \frac{MR}{2} < 1$$

b) By Assumption 1, we have $f_{y^4}^{(4)}(0,0) = 24$. We get the following lower bound:

$$\left| \int_{0}^{1} f_{y^{4}}^{(4)}(x,sy)(1-s)^{3} ds \right| = \left| \int_{0}^{1} \left(f_{y^{4}}^{(4)}(x,sy) - f_{y^{4}}^{(4)}(0,0) + f_{y^{4}}^{(4)}(0,0) \right) (1-s)^{3} ds \right|$$
$$\geq 6 - \frac{MR}{2} > 5 > 0.$$

Item (a) of Theorem 2 has been completely proved for E_6 . Set $\tilde{c} = 1 - \frac{MR}{12}$. Then

$$\begin{aligned} \left|\frac{\partial \tilde{y}}{\partial x}\right| &= \left|y\frac{1}{4}\left(\frac{1}{6}\int_{0}^{1}f_{y^{4}}^{(4)}(x,sy)(1-s)^{3}\,ds\right)^{-\frac{3}{4}}\left(\frac{1}{6}\int_{0}^{1}f_{xy^{4}}^{(5)}(x,sy)(1-s)^{3}\,ds\right)\right| \leq \\ &\leq R\frac{1}{4}\tilde{c}^{-\frac{3}{4}}\frac{M}{24} < \frac{MR}{96}\left(1-\frac{MR}{12}\right)^{-\frac{3}{4}} < \frac{1}{48}\left(1-\frac{1}{6}\right)^{-\frac{3}{4}} < 0.03. \end{aligned}$$

Step 4. Let us estimate $|1 - \frac{\partial \tilde{y}}{\partial y}|$ from formula (4).

$$\begin{split} \left| \frac{\partial \tilde{y}}{\partial y} - 1 \right| &= \left| \sqrt[4]{\frac{1}{6} \int_{0}^{1} f_{y^{4}}^{(4)}(x, sy)(1-s)^{3} ds} - 1 + \right. \\ &+ \left. \frac{y}{4} \left(\frac{1}{6} \int_{0}^{1} f_{y^{4}}^{(4)}(x, sy)(1-s)^{3} ds \right)^{-\frac{3}{4}} \frac{1}{6} \int_{0}^{1} s f_{y^{5}}^{(5)}(x, sy)(1-s)^{3} ds} \right| \leq \\ &< \left(1 + \frac{MR}{12} \right)^{\frac{1}{4}} - 1 + \frac{MR}{80} \left(1 - \frac{MR}{12} \right)^{-\frac{3}{4}} < \\ &< \left(1 + \frac{1}{6} \right)^{\frac{1}{4}} - 1 + \frac{1}{40} \left(1 - \frac{1}{6} \right)^{-\frac{3}{4}} < 0.07. \end{split}$$

For each point $\boldsymbol{x} \in U$, we have:

 $\|\phi'(\boldsymbol{x}) - I\| = \|A(\boldsymbol{x})\| \leq \sqrt{0.26^2 + 0.25^2 + 0.03^2 + 0.07^2} = \sqrt{0.1359} < 0.4 = C < 1$, that proves item (b) of the Theorem 2 for $C = \frac{2}{5}$. By Lemma 3 the coordinate change ϕ is injective in U, that proves item (c), except for the properties of $\phi(U)$.

From the last assertion of Lemma 3 and [4, Cor. 8.3, Step 1], we conclude that $\phi(U)$ contains the open disk of radius (1 - C)R centred at the origin. This completes our proof of Theorem 2 for the case of a singularity E_6 .

3.2 The case of E_8

Consider the coordinate change $\tilde{x} = x\sqrt[3]{h(x,y)} = \sqrt[3]{\sum_{k=0}^{4} \frac{y^{k}}{k!} f_{y^{k}}^{(k)}(x,0)}, \quad \tilde{y} = y\sqrt[5]{g(x,y)} = y\left[\frac{1}{24}\int_{0}^{1} f_{y^{5}}^{(5)}(x,sy)(1-s)^{4} ds\right]^{\frac{1}{5}}.$

Step 1. As in the case of E_6 , we find a bound for the term $\tilde{h}(x, y)$, in order to estimate $|1 - \frac{\partial \tilde{x}}{\partial x}|$. The difference from the case of E_6 is an additional term in the sum of absolute values that can be bound by $\frac{MR^4}{144}$, then the sum can be bound by the same geometric progression. In detail:

$$\begin{aligned} |h(x,y)-1| &= \left| \frac{f_{x^3}^{\prime\prime\prime}(0,0)}{6} + x \frac{f_{x^4}^{(4)}(c_0,0)}{24} + \sum_{k=1}^4 \left(\frac{f_{x^3}^{\prime\prime\prime\prime}(c_k,0)}{6} \right)_{y^k}^{(k)} \frac{y^k}{k!} - 1 \right| \le \\ &\le \frac{RC_{40}}{24} + \frac{RC_{31}}{6} + \frac{R^2C_{32}}{12} + \frac{R^3C_{33}}{36} + \frac{R^4C_{34}}{144} \le \frac{MR}{12} \left(\frac{5}{2} + R + \frac{R^2}{3} + \frac{R^3}{12} \right) < \frac{5}{24} \cdot 2 < \frac{1}{24} \cdot 2 \end{aligned}$$

The bound from before holds: $h(x, y) \in (0.5, 1.5)$ or $|\tilde{h}| < 0.5$. Then $\sqrt[3]{h(x, y)} \in (0.79, 1.21)$.

Similar arguments work for the estimation of the term $|h'_x|$. Thus we can bound it as we did it in the case of E_6 :

$$|h'_x| < \frac{1}{12R}.$$

Set $c_x = \frac{1}{12R}$. The element $\frac{\partial \tilde{x}}{\partial x}$ has the same representation in terms of the function h as in the case E_6 and the same estimation for every term. Hence the following bound holds:

$$\left|\frac{\partial \tilde{x}}{\partial x} - 1\right| < 0.26.$$

Step 2. a) For estimating the term $|h'_y|$ when $x \neq 0$, we can use the bound from the case of E_6 :

$$|h'_{y}| = \Big| \left(\sum_{k=0}^{4} \frac{y^{k}}{k!} \frac{f_{y^{k}}^{(k)}(x,0)}{x^{3}} \right)'_{y} \Big| \le \frac{1}{3R}.$$

b) The element $\frac{\partial \tilde{x}}{\partial y}$ has the same representation in terms of the function h as in the case of E_6 . Hence the following bound holds:

$$\left|\frac{\partial \tilde{x}}{\partial y}\right| < 0.25.$$

Step 3. In this step, we have orders of partial derivatives and a constant \tilde{c} that differ from the case of E_6 .

a) Auxiliary Assessment. By Lemma 4, we have

$$\left| \int_{0}^{1} \left(f_{y^{5}}^{(5)}(x,sy) - f_{y^{5}}^{(5)}(0,0) \right) (1-s)^{4} ds \right| \leq \int_{0}^{1} R(C_{15} + C_{06})(1-s)^{4} ds = \frac{(C_{15} + C_{06})R}{5} \leq \frac{2MR}{5}$$

b) By Assumption 1, we have $f_{y^5}^{(5)}(0,0) = 120$. We get the following lower bound:

$$\left| \int_{0}^{1} f_{y^{5}}^{(5)}(x,sy)(1-s)^{4} \, ds \right| = \left| \int_{0}^{1} \left(f_{y^{5}}^{(5)}(x,sy) - f_{y^{5}}^{(5)}(0,0) + f_{y^{5}}^{(5)}(0,0) \right) (1-s)^{4} \, ds \right| \ge 24 - \frac{2MR}{5}.$$

Set $\tilde{c} = 1 - \frac{MR}{60}$. Then

$$\begin{split} \left|\frac{\partial \tilde{y}}{\partial x}\right| &= \left|y \cdot \frac{1}{5} \left(\frac{1}{24} \int_{0}^{1} f_{y^{5}}^{(5)}(x, sy)(1-s)^{4} \, ds\right)^{-\frac{4}{5}} \cdot \left(\frac{1}{24} \int_{0}^{1} f_{xy^{5}}^{(6)}(x, sy)(1-s)^{4} \, ds\right)\right| \leq \\ &\leq R \frac{1}{5} \tilde{c}^{-\frac{4}{5}} \frac{C_{15}}{120} < \frac{MR}{600} \left(1 - \frac{MR}{60}\right)^{-\frac{4}{5}} < \frac{1}{300} \left(1 - \frac{1}{30}\right)^{-\frac{4}{5}} < 0.004 < 0.03. \end{split}$$

Step 4. Let us estimate $|1 - \frac{\partial \tilde{y}}{\partial y}|$:

$$\begin{split} \left|\frac{\partial\tilde{y}}{\partial y} - 1\right| &= \left|\sqrt[5]{\frac{1}{24}} \int_{0}^{1} f_{y^{5}}^{(5)}(x,sy)(1-s)^{4} \, ds - 1 + \right. \\ &+ \left. \frac{y}{5} \left(\frac{1}{24} \int_{0}^{1} f_{y^{5}}^{(5)}(x,sy)(1-s)^{4} \, ds\right)^{-\frac{4}{5}} \frac{1}{24} \int_{0}^{1} s f_{y^{6}}^{(6)}(x,sy)(1-s)^{4} \, ds \right| \leq \\ &\leq \left| \left(\frac{1}{24} \int_{0}^{1} \left(f_{y^{5}}^{(5)}(x,sy) - f_{y^{5}}^{(5)}(0,0) + f_{y^{5}}^{(5)}(0,0)\right)(1-s)^{4} \, ds\right)^{\frac{1}{5}} - 1 \right| + |\dots| \leq \\ &\leq \left| \left(1 - \frac{(C_{15} + C_{06})R}{120}\right)^{\frac{1}{5}} - 1 \right| + R\tilde{c}^{-\frac{4}{5}} \frac{C_{06}}{5 \cdot 24 \cdot 30} < 1 - \left(1 - \frac{MR}{60}\right)^{\frac{1}{5}} + \frac{MR}{3600} \left(1 - \frac{MR}{60}\right)^{-\frac{4}{5}} < \\ &< 1 - \left(1 - \frac{1}{30}\right)^{\frac{1}{5}} + \frac{1}{1800} \left(1 - \frac{1}{30}\right)^{-\frac{4}{5}} < 0.008 < 0.07. \end{split}$$

Thus we found the required bound for a singularity of the type E_8 : $\|\phi'(\boldsymbol{x}) - I\| < \sqrt{0.26^2 + 0.25^2 + 0.004^2 + 0.008^2} < 0.4 < 1.$

Theorem 2 is completely proved.

4 The case of A_n

Definition. A smooth function $f(u_1, u_2)$ has a singularity A_n $(n \ge 1)$ at its critical point $P \in \mathbb{R}^2$ if

- (i) the first differential df(P) = 0 and the second differential $d^2f(P) \neq 0$ and is a perfect square;
- (ii) some codition on coefficients of the Taylor series of f at P holds (see below).

Consider a linear change $(u_1, u_2) \to (x, y)$ such that $d^2 f(P) = \pm (dx)^2$. By using a non-linear change $(x, y) \to (x_1 = x + y^2 Q(y), y_1 = y)$ for some polynomial Q(y), one can achive that $f_{x_1^a y_1^b}^{(a+b)}(P) = 0$ for all $a, b \in \mathbb{Z}_+$ with $0 < \frac{a}{2} + \frac{b}{n+1} < 1$. Notice that $f_{x_1^2}'(P) = \pm 2$. The condition on the Taylor coefficients is as follows: $f_{y_1^{n+1}}^{(n+1)}(P) \neq 0$.

We will assume that P = (0, 0) = 0 in the coordinates x, y.

Assumption 2. For singularities A_n $(n \ge 1)$, assume that $f_{y_1^{n+1}}^{(n+1)}(0,0) = \pm (n+1)!$.

Theorem 3 (Estimating the radius of a neighborhood for the coordinate change). In the case of a singularity A_n $(n \ge 1)$, under the hypotheses of Assumption 2, suppose that, in a neighbourhood $U_0 = \{(x_1, y_1) \mid \max(|x_1|, |y_1|) < R_0\}$ of **0**, the following estimates hold: $C_{\alpha\beta} = \sup_{U_0} \left| f_{x_1^{\alpha}y_1^{\beta}}^{(\alpha+\beta)}(x_1, y_1) \right| \le M$ for all $(\alpha, \beta) \in \{(0, n+2), (0, n+1), (2, i), (3, 0), (3, i) \mid i =$ $0, 1, \ldots, k\}$, where $R_0 > 0$, $M \ge 0$. Then, in the neighborhood $U = \{(x_1, y_1) \mid \max(|x_1|, |y_1|) < R\}$, with $R = \min\{R_0, \frac{3}{4M+3}\}$, there is a coordinate change of the form $\phi : (x_1, y_1) \to (\tilde{x} =$ $x_1\sqrt{h(x_1, y_1)}, \tilde{y} = y_1(g(x_1, y_1))^{\frac{1}{k+1}}$) that reduces f to the normal form $f = f(P) \pm \tilde{x}^2 \pm \tilde{y}^{n+1}$ of A_n . In detail: the coordinate change $\phi|_U$ satisfies the conditions (a), (b), (c) from Theorem 2 with C = 0.93.

Proof. By abusing notations, we will denote (x_1, y_1) by (x, y). Consider the coordinate chage

$$\tilde{x} = x\sqrt{h(x,y)} = \sqrt{\sum_{k=0}^{n} \frac{y^k}{k!} f_{y^k}^{(k)}(x,0)}, \quad \tilde{y} = y \sqrt[n+1]{g(x,y)} = y \left[\frac{1}{n!} \int_0^1 f_{y^{n+1}}^{(n+1)}(x,sy)(1-s)^n \, ds\right]^{\frac{1}{n+1}}$$

We extend our proof of Theorem 2 (about singularities E_6, E_8) to the case of singularities A_n .

Step 1. a) Let us estimate $\tilde{h}(x, y)$:

$$\begin{split} |h(x,y)-1| &= \left| \frac{f_{x^2}''(0,0)}{2} + x \frac{f_{x^3}''(c_0,0)}{6} + \sum_{k=1}^n \frac{f_{x^2y^k}^{(k+2)}(c_k,0)}{2} \frac{y^k}{k!} - 1 \right| \leq \\ &\leq \frac{RC_{30}}{6} + \sum_{k=1}^n \frac{C_{2k}R^k}{2 \cdot k!} \leq \frac{2MR}{3(1-R)} \leq \frac{1}{2}. \end{split}$$

Thus $h(x,y) \in (0.5,1.5)$, or $|\tilde{h}| < 0.5 =: c$. Then $\sqrt{h(x,y)} \in (0.7,1.3)$. We have

$$\frac{\partial \tilde{x}}{\partial x} = h^{\frac{1}{2}} + x \cdot \frac{1}{2} h^{-\frac{1}{2}} \cdot h'_x = (1 + \tilde{h})^{\frac{1}{2}} + x \cdot \frac{1}{2} (1 + \tilde{h})^{-\frac{1}{2}} \cdot h'_x.$$

Let us estimate $|h'_x|$:

$$\begin{aligned} |h'_{x}| &= \left| \left(\sum_{k=0}^{n} \frac{y^{k}}{k!} \frac{f_{y^{k}}^{(k)}(x,0)}{x^{2}} \right)'_{x} \right| = \left| \left(\sum_{k=0}^{n} \frac{y^{k}}{k!} \int_{0}^{1} f_{y^{k}x^{2}}^{(k+2)}(sx,0)(1-s) \, ds \right)'_{x} \right| = \\ &= \left| \sum_{k=0}^{n} \frac{y^{k}}{k!} \int_{0}^{1} s f_{y^{k}x^{3}}^{(k+3)}(sx,0)(1-s) \, ds \right| \le \sum_{k=0}^{n} \frac{R^{k}C_{3k}}{k!} \int_{0}^{1} (s-s^{2}) \, ds \le \\ &\le \frac{M}{6(1-R)} \le \frac{1}{8R}. \end{aligned}$$

Set $c_x = \frac{1}{8R}$. Let us estimate $|\frac{\partial \tilde{x}}{\partial x} - 1|$:

$$\left|\frac{\partial \tilde{x}}{\partial x} - 1\right| = \left|\sqrt{h} - 1 + x \cdot \frac{1}{2}(1 + \tilde{h})^{-\frac{1}{2}}h'_{x}\right| \le \left|\sqrt{h} - 1\right| + \left|x \cdot \frac{1}{2}(1 + \tilde{h})^{-\frac{1}{2}}h'_{x}\right| < 0.3 + \frac{c_{x}R}{2}(1 - c)^{-\frac{1}{2}} < 0.3 + \frac{2^{\frac{1}{2}}}{2 \cdot 8} < 0.3 + 0.1 = 0.4.$$

Step 2. a) Let us estimate $|h'_y|$ when $x \neq 0$:

$$\begin{aligned} |h'_{y}| &= \Big| \left(\sum_{k=0}^{n} \frac{y^{k}}{k!} \frac{f_{y^{k}}^{(k)}(x,0)}{x^{2}} \right)'_{y} \Big| = \Big| \left(\sum_{k=0}^{n-1} \frac{y^{k}}{k!} \frac{f_{y^{k+1}}^{(k+1)}(x,0)}{x^{2}} \right) \Big| = \\ &= \left| \int_{0}^{1} \left(\sum_{k=0}^{n-1} \frac{y^{k}}{k!} f_{x^{2}y^{k+1}}^{(k+3)}(sx,0) \right) (1-s) \, ds \right| \le \frac{1}{2} \sum_{k=0}^{n-1} \frac{R^{k} C_{2,k+1}}{k!} \le \frac{M}{2(1-R)} \le \frac{3}{8R}. \end{aligned}$$

Set $c_y = \frac{3}{8R}$. b) Let us estimate $|\frac{\partial \tilde{x}}{\partial y}|$:

$$\begin{aligned} \left|\frac{\partial \tilde{x}}{\partial y}\right| &= \left|x \cdot \frac{1}{2} \left(1 + \tilde{h}\right)^{-\frac{1}{2}} \cdot h'_{y}\right| = \left|\frac{x}{2} \left(1 - \frac{1}{2}\tilde{h} + \frac{3}{8}(1 + c_{\tilde{h}})^{-\frac{5}{2}}\tilde{h}^{2}\right)h'_{y}\right| \leq \\ &\leq \frac{R}{2} \left(1 + \frac{1}{2}c + \frac{3}{8}(1 + c_{\tilde{h}})^{-\frac{5}{2}}c^{2}\right)c_{y} < \frac{c_{y}R}{2} \left(1 + \frac{1}{2}c + \frac{3}{8}(1 - c)^{-\frac{5}{2}}c^{2}\right) = \\ &= \frac{3}{16} \left(1 + \frac{1}{4} + \frac{3}{8}2^{\frac{5}{2}}/4\right) < 0.4. \end{aligned}$$

Step 3. a) Auxiliary Assessment. By Lemma 4:

$$\left| \int_{0}^{1} \left(f_{y^{n+1}}^{(n+1)}(x,sy) - f_{y^{n+1}}^{(n+1)}(0,0) \right) (1-s)^{n} ds \right| \leq \int_{0}^{1} R(C_{1,n+1} + C_{0,n+2}) (1-s)^{n} ds \leq \frac{2MR}{n+1}.$$

b) By assumption 3 we have $f_{y^{n+1}}^{(n+1)}(0,0) = (n+1)!$. We get the following lower bound:

$$\left| \int_{0}^{1} f_{y^{n+1}}^{(n+1)}(x,sy)(1-s)^{n} \, ds \right| = \left| \int_{0}^{1} \left(f_{y^{n+1}}^{(n+1)}(x,sy) - f_{y^{n+1}}^{(n+1)}(0,0) + f_{y^{n+1}}^{(n+1)}(0,0) \right) (1-s)^{n} \, ds \right| \ge n! - \frac{2MR}{n+1}$$

Set $\tilde{c} = 1 - \frac{2MR}{(n+1)!}$. Then the following sequence of inequalities holds:

$$\begin{split} \left|\frac{\partial \tilde{y}}{\partial x}\right| &= \left|y \cdot \frac{1}{n+1} \left(\frac{1}{n!} \int_{0}^{1} f_{y^{n+1}}^{(n+1)}(x, sy)(1-s)^{n} \, ds\right)^{-\frac{n}{n+1}} \cdot \left(\frac{1}{n!} \int_{0}^{1} f_{xy^{n+1}}^{(n+2)}(x, sy)(1-s)^{n} \, ds\right)\right| &\leq \\ &\leq R \frac{1}{n+1} \tilde{c}^{-\frac{n}{n+1}} \frac{C_{1,n+1}}{(n+1)!} \leq \frac{MR}{(n+1)(n+1)!} \left(1 - \frac{2MR}{(n+1)!}\right)^{-\frac{n}{n+1}} < \frac{3}{4 \cdot 4} \left(1 - \frac{3}{2(n+1)!}\right)^{-\frac{n}{n+1}} \leq \frac{3}{8}. \end{split}$$

Let us prove the latter inequality in this sequence. For n = 1 this inequality is in fact an equality. For n > 1 we have $\tilde{c}^{-\frac{n}{n+1}} < \tilde{c}^{-1}$, moreover the function $\tilde{c}^{-1}|_{MR=\frac{3}{4}}$ of n is monotone decreasing. Since $\tilde{c}^{-1}|_{MR=\frac{3}{4}} < 2$ for n = 2, we have the same for all n > 2. This proves the desired inequality.

Step 4. Let us estimate $|1 - \frac{\partial \tilde{y}}{\partial y}|$:

$$\begin{split} \left|\frac{\partial \tilde{y}}{\partial y} - 1\right| &= \left| \left| \sqrt[n+1]{1} \int_{0}^{1} f_{y^{n+1}}^{(n+1)}(x, sy)(1-s)^{n} ds - 1 + \right. \\ &+ \frac{y}{n+1} \left(\frac{1}{n!} \int_{0}^{1} f_{y^{n+1}}^{(n+1)}(x, sy)(1-s)^{n} ds \right)^{-\frac{n}{n+1}} \frac{1}{n!} \int_{0}^{1} sf_{y^{n+2}}^{(n+2)}(x, sy)(1-s)^{n} ds \right| \leq \\ &\leq \left| \left(\frac{1}{n!} \int_{0}^{1} \left(f_{y^{n+1}}^{(n+1)}(x, sy) - f_{y^{n+1}}^{(n+1)}(0, 0) + f_{y^{n+1}}^{(n+1)}(0, 0) \right)(1-s)^{n} ds \right)^{\frac{1}{n+1}} - 1 \right| + |\dots| \leq \\ &\leq \left| \left(1 - \frac{(C_{1,n+1} + C_{0,n+2})R}{(n+1)!} \right)^{\frac{1}{n+1}} - 1 \right| + R\tilde{c}^{-\frac{n}{n+1}} \frac{C_{0,n+2}}{(n+2)(n+1)(n+1)!} < \\ &< 1 - \left(1 - \frac{2MR}{(n+1)!} \right)^{\frac{1}{n+1}} + \dots = 1 - \tilde{c}^{\frac{1}{n+1}} + \frac{MR\tilde{c}^{-\frac{n}{n+1}}}{(n+2)(n+1)(n+1)!} < \\ &< \frac{3}{2 \cdot (n+1)!} + \frac{3 \cdot 2}{4 \cdot 3 \cdot 2 \cdot 2} \leq \frac{1}{2} \leq \frac{5}{8} \end{split}$$

if n > 1. And when n = 1, we get $1 - \tilde{c}^{\frac{1}{2}} \le \frac{1}{2}$, therefore $|\frac{\partial \tilde{y}}{\partial y} - 1| < \frac{1}{2} + \frac{1}{8} = \frac{5}{8}$. Thus for A_n we have $\|\phi'(\boldsymbol{x}) - I\| < \sqrt{0.4^2 + 0.4^2 + \frac{3^2}{8} + \frac{5^2}{8}} < 0.93 < 1$ for all $n \ge 1$. Theorem 3 is proved.

References

- Arnol'd V. I. Normal forms for functions near degenerate critical points, the Weyl groups of A_k, D_k, E_k and Lagrangian singularities // Funct. Anal. Appl. – 1972. – V. 6, No. 4. – P. 254–272.
- [2] Brodersen H. M-t topologically stable mappings are uniformly stable // Math. Scand. 1983. – V. 52. – P. 61–68.
- [3] Kudryavtseva E. A. Uniform Morse lemma and isotope Morse functions on surfaces // Moscow Univ. Math. Bull. - 2009. - V. 64, No. 4. - P. 150-158.
- [4] Kudryavtseva E. A., Permyakov D. A. Framed Morse functions on surfaces // Sbornik Math. - 2010. - V. 201, No. 4. - P. 501-567.
- [5] Kudryavtseva E. A. The Topology of Spaces of Morse Functions on Surfaces // Math. Notes - 2012. - V. 92, No. 2. - P. 219-236. http://arxiv.org/abs/1104.4792.
- [6] Kudryavtseva E. A. Special framed Morse functions on surfaces // Moscow Univ. Math. Bull. - 2012. - V. 67, No. 4. - P. 151-157. http://arxiv.org/abs/1106.3116.
- [7] Kudryavtseva E. A. On the homotopy type of spaces of Morse functions on surfaces // Sb. Math. - 2013. - V. 204, No. 1. - P. 75-113. http://arxiv.org/abs/1104.4796.
- [8] Kudryavtseva E. A. Topology of the spaces of functions with prescribed singularities on surfaces // Doklady Mathematics. - 2016. - V. 93, No. 3. - P. 264-266.
- [9] Kudryavtseva E. A. Topology of the spaces of functions and gradient-like flows with prescribed singularities on surfaces // Arxiv - 2021. http://arxiv.org/abs/2106.03017.
- [10] Mather J. N. Infinitesimal stability implies stability // Ann. of Math. 1969. V. 89. -P. 254-291.
- [11] Samoilenko A. M. The equivalence of a smooth function to a Taylor polynomial in the neighborhood of a finite-type critical point // Funct. Anal. Appl. 1968. V. 2, No. 4. P. 318–323.
- [12] Sergeraert F. Un théorème de fonctions implicites sur certains espaces de Fréchet et quelques applications // Annales scientifiques de l'É.N.S. 4e série. – 1972. – V. 5, No. 4. – P. 599–660.
- [13] Tougeron J. C. Ideaux de fonctions differentiables. I // Ann. Inst. Fourier. 1968. V. 18, No. 1. – P. 177–240.