LABELLED COSPAN CATEGORIES AND PROPERADS

JONATHAN BEARDSLEY AND PHILIP HACKNEY

ABSTRACT. We prove Steinebrunner's conjecture on the biequivalence between (colored) properads and labelled cospan categories. The main part of the work is to establish a 1-categorical, strict version of the conjecture, showing that the category of properads is equivalent to a category of strict labelled cospan categories via the symmetric monoidal envelope functor.

Contents

1. Background	4
1.1. Labelled cospan categories	4
1.2. The category of level graphs	6
1.3. Properads as Segal L-presheaves	8
2. The symmetric monoidal envelope of a properad	13
2.1. Congruences of level graphs	13
2.2. The category CX	15
2.3. Monoidal structure	18
2.4. The symmetry	20
3. Properads give labelled cospan categories	25
4. Strict labelled cospan categories	30
4.1. The inert part	31
4.2. The active part	32
5. Comparison of constructions	47
6. 2-Categorical structures	52
6.1. Equivalences of labelled cospan categories	53
6.2. Strict labelled cospan categories form a 2-category	54
6.3. Comparison with labelled cospan categories	57
References	61

One way to encode the data of a cobordism is as a cospan $M_1 \to P \leftarrow M_2$, where P is a manifold and M_1 and M_2 are its "left" and "right" boundaries, respectively. By taking connected components, each such cospan gives a cospan of finite sets. This

Date: April 27, 2023.

²⁰²⁰ Mathematics Subject Classification. 18M85, 18B10, 18F20, 18M60, 55P48, 55U10, 05C20. Key words and phrases. cospan, properad, labelled cospan category, Segal condition.

This work was supported by grants from the Simons Foundation (#853272, JB; #850849, PH). This material is partially based upon work supported by the National Science Foundation under Grant No. DMS-1928930 while the second author participated in a program supported by the Mathematical Sciences Research Institute. The program was held in the summer of 2022 in partnership with the Universidad Nacional Autónoma de México.




FIGURE 1. Properadic composition of type 124; 345

functor from cobordisms to cospans of finite sets precisely relates decompositions of objects and morphisms between the two categories. An abstract version of this appears under the name *labelled cospan category* in [Ste]. Therein, Steinebrunner develops the general theory of labelled cospan categories and uses it to show, among other things, that the classifying space of the category of 2-dimensional cobordisms is rationally equivalent to S^1 .

The cobordism category is a prop in the sense of Adams and Mac Lane. It is in fact the free prop on a *properad* of connected cobordisms. Properads, introduced in [Val07] and independently under the name compact symmetric polycategory in [Dun06], are like props but without horizontal composition. Operations may have multiple inputs and outputs, and can be composed by attaching some (nonzero number) of inputs of one to the outputs of another. We do not give the precise definition (for that, see [HRY15, Chapter 3] or [YJ15, 11.7]), but rather an illustration in Figure 1 where the letters represent colors in the properad. In the example of cobordisms, the connected cobordisms form a properad and then become a prop when we allow disjoint unions. Steinebrunner conjectured that a similar connection between properads and labelled cospan categories holds in generality. Our main result is the following:

Theorem A. The 2-category of properads is biequivalent to the 2-category of labelled cospan categories.

This appears as Corollary 6.16 below, and establishes the first part of Conjecture 2.31 of [Ste] (we do not address the second part of the conjecture which concerns ∞ -properads). The underlying functor of 1-categories from properads to labelled cospan categories is not an equivalence, so 2-categorical structure is essential. It is also the case (see Proposition 6.13) that if a symmetric monoidal category admits the structure of a labelled cospan category, then this structure is unique up to equivalence. Thus another interpretation of Theorem A is that it provides a faithful inclusion of properads into symmetric monoidal categories and identifies its image.

The main effort of this paper consists of a careful analysis of the symmetric monoidal envelope, which takes properads to symmetric monoidal categories; this functor extends the classical envelope of an operad [LV12, 5.4.1], which goes back to Boardman–Vogt. The envelope of the terminal properad is the category of cospans of finite sets, and applying the envelope to the unique map from a properad to the terminal properad yields a labelled cospan category. This construction actually produces a more rigid kind of object, which we have called a *strict labelled cospan category*. In the last section we establish a biequivalence (Theorem 6.15) between

labelled cospan categories and strict labelled cospan categories, which we combine with the following to obtain the main result.

Theorem B (Corollary 5.9 and Theorem 6.7). There is a strict 2-equivalence between the 2-category of properads and the 2-category of strict labelled cospan categories.

In particular, this yields an equivalence of categories between the underlying 1-categories. Though our focus in this introduction has been on 2-categories, it is this 1-categorical equivalence (Corollary 5.9) that is the core of the result.

A key ingredient in this paper is an efficient description of the symmetric monoidal envelope of a properad P. We use an alternative description of properads as Segal presheaves on a category of level graphs **L**. That is, we use an equivalence of categories $\mathbf{Ppd} \simeq \mathrm{Seg}(\mathbf{L})$; which appears below as Proposition 1.15 (the hard work of establishing this equivalence was already done in [HRY15] and [CH22]). The upshot is that a Segal **L**-presheaf looks much more like a symmetric monoidal category than a properad does (in particular, it is important that the category **L** contains disconnected graphs, as these account for the monoidal structure). To form the envelope is then a relatively straightforward quotienting process, though managing symmetry isomorphisms still requires some care.

It is much more involved to go back in the other direction. Namely, given a strict labelled cospan category C, we would like to produce a Segal **L**-presheaf that it comes from. There is an active-inert factorization system on **L**, and there is essentially only one possible choice of an \mathbf{L}_{int} -presheaf that could be the restriction of our desired **L**-presheaf. We must then extend the \mathbf{L}_{int} -presheaf by defining its action on active maps. Here we make use of additional simplicial structure: by using heights of level graphs, **L** is fibered over the simplicial category $\boldsymbol{\Delta}$. The main thing that is missing are the inner face and degeneracy operators arising from this additional simplicial structure, and coskeletality arguments allow us to extend from low simplicial degree (using the composition and identities in C) to arbitrary simplicial degree.

Remark (Related work). As mentioned above, we can describe a properad by giving its envelope, along with map to the envelope of the terminal properad. The same thing can be done for operads, though we are not aware of any classical literature which pursues this idea. However, this is precisely the approach that is taken by recent work of Haugseng–Kock [HK] for ∞ -operads (see also [BHS]). Following the initial public version of this paper, the preprints [KM] and [BS] appeared and offered related theorems to ours. The work by Barkan–Steinebrunner on ∞ -properads is related to the Haugseng–Kock approach for ∞ -operads. Some important results of [BS] are ∞ -categorical versions our theorems, and by restricting to discrete objects they recover the (2, 1)-categorical version of Theorem A (which is also an immediate consequence of Theorem A). Their elegant theory relies on the idea of 'equifibered maps of \mathbb{E}_{∞} -monoids,' and we highly recommend [BS] for an important alternative perspective. They also give information concerning the relationship with the 'hereditary unique factorization categories' of Kaufmann–Monaco [KM].

Acknowledgements. We are grateful to Jan Steinebrunner for feedback on an earlier version of this manuscript, as well as a number of interesting suggestions. We are thankful for several suggestions and comments from anonymous referees,

which have helped us improve the presentation and correct several oversights. We also thank Joe Moeller, Marcy Robertson, and Donald Yau for helpful discussions.


1. BACKGROUND

The category **FinSet** is the category of finite sets and arbitrary functions between them. Let $\mathbf{F} \subseteq \mathbf{FinSet}$ be the full subcategory whose objects are the ordered sets $\underline{k} = \{1, 2, \dots, k\}$ for $k \ge 0$. We will write $\boldsymbol{\Delta}$ for the (topologist's) simplicial category, whose objects are the ordered sets $[n] = \{0 < 1 < \dots < n\}$ for $n \ge 0$ and morphisms preserve the \leq relation. Note the shift in cardinality if you regard $\boldsymbol{\Delta}$ as a subcategory of \mathbf{F} .

The category **Csp** has objects the same as **FinSet**, and morphisms from A to B are equivalence classes of cospans $A \to C \leftarrow B$, where two are identified if there is an isomorphism on the middle term:



Composition of morphisms in Csp is given by pushout



Choices of coproducts of finite sets yields a monoidal structure on Csp with \emptyset as the monoidal unit. This is not a cocartesian monoidal category, however.

1.1. Labelled cospan categories. This work is concerned with Steinebrunner's notion of labelled cospan categories from [Ste], which are certain symmetric monoidal categories living over Csp. We briefly recall some definitions.


Definition 1.1. Let $\pi: C \to \mathbf{Csp}$ be a symmetric monoidal functor. For the moment, it is convenient to write

$$\begin{array}{ccc} \pi c & \pi d \\ & & \swarrow_{rf} \\ & & mf \end{array}$$

for $\pi(f: c \to d)$.

- An object $c \in C$ is *connected* if $\pi(c)$ has cardinality one.
- A morphism $f: c \to d$ is connected if m(f) has cardinality one.
- A morphism $f: c \to d$ is reduced if the cospan $\pi(f)$ is jointly surjective.
- hom^c(c, d) ⊂ hom(c, d) denotes the connected morphisms and hom^r(c, d) ⊂ hom(c, d) denotes the reduced morphisms.

By examining the diagrams



whose diamonds are pushouts, we see that the sets of connected morphisms and reduced morphisms are not closed under composition.

If $A \to X \leftarrow B$ represents an isomorphism in **Csp**, then both legs of the cospan are bijections. Thus every isomorphism in C is reduced, and every isomorphism involving a connected object is connected.

Definition 1.2 (Steinebrunner). A *labelled cospan category* is a symmetric monoidal functor $\pi: C \to \mathbf{Csp}$ satisfying the following:

- (1) If $\pi(c)$ has cardinality *n*, then we can find *n* objects c_1, \ldots, c_n which are connected so that *c* is isomorphic to $c_1 \otimes \cdots \otimes c_n$.
- (2) If **1** is the tensor unit of C, then the abelian monoid hom(**1**, **1**) is freely generated by the set hom^c(**1**, **1**) of connected morphisms.
- (3) The map

$$\operatorname{hom}^{\mathrm{r}}(c,d) \times \operatorname{hom}(\mathbf{1},\mathbf{1}) \xrightarrow{\otimes} \operatorname{hom}(c,d)$$

is a bijection.

(4) For each four objects c, d, c', d' in C, the following square is cartesian.

A map of labelled cospan categories is a symmetric monoidal functor and a choice of monoidal natural isomorphism making the triangle over **Csp** commute.




Remark 1.3. For a fixed symmetric monoidal functor $f: C \to C'$, there is at most one monoidal natural isomorphism as displayed above; in other words, the forgetful functor from labelled cospan categories to symmetric monoidal categories is faithful. Uniqueness of the monoidal natural isomorphism follows from Definition 1.2(1) and the fact that there are unique isomorphisms in **Csp** between objects of cardinality zero or one. Given two such natural isomorphisms, the following diagram of isomorphisms commutes for each of them, where $c \cong c_1 \otimes \cdots \otimes c_n$ is the decomposition into connected objects guaranteed by (1).

Since the top map is the same for both natural isomorphisms, so too is the bottom map.

Remark 1.4 (2-categorical structure). We can regard the collection of labelled cospan categories as a 2-category, where a map between morphisms (f, α) and (g, β) having the same source and target



is a monoidal natural transformation $\gamma\colon f\Rightarrow g$ so that the composite natural transformation




is an isomorphism. By Remark 1.3, we then have this composite is α . Steinebrunner only needs this when γ is an isomorphism, that is, he considers labelled cospan categories as forming a (2, 1)-category.

We will return to this 2-categorical structure of labelled cospan categories in Section 6.

1.2. The category of level graphs. In this section we recall the category of level graphs \mathbf{L} from [CH22, §2.1]. As mentioned below [CH22, Lemma 2.1.9], this category is closely related to the *double* category of cospans, an enhancement of **Csp**. In Section 2 will see how to rederive **Csp** (up to equivalence) from \mathbf{L} .

The category \mathscr{L}^n has, as its objects, pairs (i, j) with $0 \leq i \leq j \leq n$, and as morphisms, unique maps $(i, j) \to (k, \ell)$ whenever $0 \leq k \leq i \leq j \leq \ell \leq n$. This presentation is as in [CH22, Definition 2.16], though it is also true that \mathscr{L}^n is isomorphic to the twisted arrow category of $[n] = \{0 \to 1 \to \cdots \to n\}$. Note that every square in \mathscr{L}^n commutes and is a pushout. Here is \mathscr{L}^3 :



A level graph of height n is a functor $G: \mathscr{L}^n \to \mathbf{F}$ that sends every square to a pushout.¹ In particular, level graphs of height 0 are just objects of \mathbf{F} , and level graphs of height 1 are just cospans in \mathbf{F} . We write G_{ij} or $G_{i,j}$ for the value of G on

¹This is slightly different from [CH22], where the target category was all finite sets and the functor only concerned the top layers $\mathscr{L}_0^n \to \mathbf{FinSet}$. This makes no substantial difference, and we arrive at an equivalent category of level graphs below; compare with [CH22, 2.1.19].





FIGURE 2. A level graph of height 2

the object (i, j) and do not label the structural maps $G_{ij} \to G_{k\ell}$. Figure 2 gives a pictorial example of a particular height 2 level graph the following shape:



We consider the subcategory \mathbf{L}_n of $\operatorname{Fun}(\mathscr{L}^n, \mathbf{F})$ whose objects are the level graphs and whose morphisms are the natural transformations $G \Rightarrow H$ satisfying the following two properties for each $0 \leq i \leq j \leq n$:


- (1) The map $G_{ij} \to H_{ij}$ is a monomorphism.
- (2) The naturality square

$$\begin{array}{ccc} G_{ij} & \longrightarrow & H_{ij} \\ \downarrow & {}^{\neg} & & \downarrow \\ G_{0n} & \longrightarrow & H_{0n} \end{array}$$

is cartesian.

By the pasting law for pullbacks, the second condition is equivalent to the natural transformation being cartesian, as was required in [CH22, Definition 2.1.16]. This defines a functor $\mathbf{L}_{\bullet} \colon \Delta^{\mathrm{op}} \to \mathbf{Cat}$, and we write $\mathbf{L} \to \Delta$ for the associated Grothendieck fibration. (The version of \mathbf{L} appearing in [CH22] is a skeleton of this one.)

Some remarks on the simplicial category \mathbf{L}_{\bullet} will make later proofs easier to understand (see [CH22] for complete details). First we note that, in fact, the categories \mathscr{L}^n assemble into a cosimplicial category. Given a morphism $\alpha \colon [m] \to [n]$ in Δ the functor $\alpha^* \colon \mathscr{L}^m \to \mathscr{L}^n$ takes the pair (i, j) to the $(\alpha i, \alpha j)$. For instance all of the induced functors $\mathscr{L}^0 \to \mathscr{L}^n$ take the unique object (0, 0) to some object (i, i)in \mathscr{L}^n (and never objects of the form (i, j) with $i \neq j$). One can also consider the three coface morphisms $\mathscr{L}^1 \to \mathscr{L}^2$. These can be visualized with color as follows:



These induce functors $\mathbf{L}^2 \to \mathbf{L}^1$ that, respectively, truncate level 2 of a height 2 level graph, contract level 1 of a height 2 level graph, and truncate level 0 of a height 2 level graph. More generally, on a height *n* level graph $G: \mathscr{L}^n \to \mathbf{F}$, a morphism $\alpha: [m] \to [n]$ in $\mathbf{\Delta}$ induces a functor $\mathbf{L}^n \to \mathbf{L}^m$ which precomposes with $\mathscr{L}^m \to \mathscr{L}^n$, i.e. $(\alpha^* G)_{ij} = G_{\alpha i, \alpha j}$ for each $(i, j) \in \mathscr{L}^m$.

If $G \to H$ is a map in \mathbf{L}_n , then it is uniquely determined by $\{G_{ii} \to H_{ii}\}_{0 \le i \le n}$ and $\{G_{i-1,i} \to H_{i-1,i}\}_{1 \le i \le n}$, since the functors G and H are pushout-preserving. This implies that $\mathbf{L}_n \to \mathbf{L}_k \times_{\mathbf{L}_0} \mathbf{L}_{n-k}$ is fully faithful. Further, this functor is surjective on objects, hence an equivalence of categories. It is not injective on objects (unless k = 0 or n).

1.3. **Properads as Segal L-presheaves.** If G is a level graph of height m and H is a level graph of height n, then an *active map* $G \to H$ is one whose image in Δ is active (preserves the top and bottom elements) and where $G_{0m} \to H_{0n}$ is a bijection [CH22, Remark 2.1.24]. Inert maps $G \to H$ are precisely those whose image in Δ is inert (distance preserving).

Definition 1.5 (Elementary objects). The category \mathbf{L} contains several *elementary* objects. These consist of the edge $\mathfrak{e} \in \mathbf{L}_0$, which corresponds to $\underline{1}$ using $\mathrm{ob}(\mathbf{L}_0) = \mathrm{ob}(\mathbf{F})$, and a collection of m, n-corollas $\mathfrak{c}_{m,n} \in \mathbf{L}_1$ for non-negative integers m and n. The graph $\mathfrak{c}_{m,n}$ is the following:

$$\underbrace{\underline{m}}_{\substack{\boldsymbol{\searrow}} \ \boldsymbol{\swarrow}} \underbrace{\underline{n}}_{\underline{1}}$$

Definition 1.6. The above allows to define three subcategories of **L**:

- Write \mathbf{L}_{int} for the wide subcategory of \mathbf{L} on the inert morphisms.
- Write L_{el} for the full subcategory of L_{int} spanned by the elementary graphs of Definition 1.5.

• Write L_{act} for the wide subcategory of L on the active morphisms.

Remark 1.7. The three subcategories of Definition 1.6 make \mathbf{L}^{op} into an algebraic pattern in the sense of [CH21, Definition 2.1].

Definition 1.8 (Segal objects). Let $\mathbf{L}_{/G}^{\text{el}}$ denote the category whose objects are inert maps $E \rightarrow G$ with E elementary, and whose morphisms are commutative triangles with all maps inert. A presheaf $X \in \text{Psh}(\mathbf{L})$ is said to be *Segal* if

$$X_G \to \lim_{E \in (\mathbf{L}_{/G}^{\mathrm{el}})^{\mathrm{op}}} X_E$$

is a bijection for all $G \in \mathbf{L}$. We write $\text{Seg}(\mathbf{L}) \subseteq \text{Psh}(\mathbf{L})$ for the full subcategory consisting of the Segal presheaves.

Definition 1.9 (Segal core). Given $G \in \mathbf{L}$, the *Segal core* is the following colimit in $Psh(\mathbf{L})$

$$\operatorname{Sc}(G) \coloneqq \operatorname{colim}_{E \in \mathbf{L}_{/G}^{\operatorname{el}}} \mathfrak{L}(E) \to \mathfrak{L}(G)$$

which comes equipped with a map to the representable object $\&(G) = \hom_{\mathbf{L}}(-, G)$.

A presheaf $X \in Psh(\mathbf{L})$ is Segal if and only if it is local with respect to the Segal core inclusions, that is, if and only if

$$\hom(\mathfrak{L}(G), X) \to \hom(\mathrm{Sc}(G), X)$$

is a bijection for all graphs G.

Remark 1.10. Suppose $X \in Psh(\mathbf{L})$ is Segal. We list explicit consequences. The first two are related to the 'simplicial' direction of \mathbf{L} ; the second of these will play a role in the proof of Proposition 2.15. The third of these will be used repeatedly, and is about the behavior in the fibers.

• If $G \in \mathbf{L}_n$ and 0 < k < n, let $\alpha : [k] \rightarrow [n]$ be the inert inclusion into the first part of the interval $(\alpha(t) = t), \beta : [n - k] \rightarrow [n]$ the inert inclusion into the last part of the interval $(\beta(t) = t + k)$, and $k : [0] \rightarrow [n]$ pick out k. Then

$$X_G \to X_{\alpha^*(G)} \times_{X_{k^*(G)}} X_{\beta^*(G)}$$

is an isomorphism. At the level of presheaves, we have a commutative square

$$\begin{array}{c} \operatorname{Sc}(\alpha^{*}(G)) \amalg_{\operatorname{Sc}(k^{*}(G))} \operatorname{Sc}(\beta^{*}(G)) \longrightarrow \& (\alpha^{*}(G)) \amalg_{\& (k^{*}(G))} \& (\beta^{*}(G)) & \\ & \downarrow & \\ & \downarrow & \\ & \operatorname{Sc}(G) \longrightarrow \& (G) \end{array}$$

whose left edge is an isomorphism. The claimed statement follows by applying hom(-, X) (which transforms pushouts to pullbacks) and using 2-of-3 for isomorphisms.

• We can iterate the previous observation. Suppose $G \in \mathbf{L}$ is a height n level graph and for $1 \leq i \leq n$ that $\rho_i: [1] \mapsto [n]$ in $\mathbf{\Delta}$ is the inert map picking out i-1, i, and for $0 \leq i \leq n$ the map $\kappa_i: [0] \mapsto [n]$ picks out i. If $\mathfrak{c}_{p,q} \mapsto G$ is an inert map in \mathbf{L} , then it factors uniquely through $\rho_i^* G$ (which has height 1) for some i. Likewise, any $\mathfrak{e} \mapsto G$ factors uniquely through some $\kappa_i^* G$ (which has height 0). Further,

$$X_G \to X_{\rho_1^*G} \times_{X_{\kappa_1^*G}} X_{\rho_2^*G} \times_{X_{\kappa_2^*G}} \cdots \times_{X_{\kappa_{n-1}^*G}} X_{\rho_n^*G}$$

is a bijection. See [CH22, Proposition 3.2.9].

• Suppose $G, H \in \mathbf{L}_n$ are two height n level graphs. We can define a new height n level graph G + H with $(G + H)_{ij} = G_{ij} + H_{ij}$, which is a coproduct in the fiber \mathbf{L}_n (see Section 2.3 where this will be used more). Then $\mathbf{L}_{/(G+H)}^{\text{el}} = \mathbf{L}_{/G}^{\text{el}} \amalg \mathbf{L}_{/H}^{\text{el}}$, so it follows that the left hand map in the following square is an isomorphism:


$$Sc(G) \amalg Sc(H) \longrightarrow \& (G) \amalg \& (H) \\ \downarrow \cong \qquad \qquad \downarrow \\ Sc(G+H) \longrightarrow \& (G+H).$$

Applying hom(-, X) and using 2-of-3 for isomorphisms, we see that $X_{G+H} \rightarrow X_G \times X_H$ is a bijection:

$$\begin{array}{ccc} X_{G+H} & & \longrightarrow & X_G \times X_H \\ & & & \downarrow \cong & \\ & & \downarrow \cong & \\ \lim_{E \in (\mathbf{L}_{/(G+H)}^{\mathrm{el}})^{\mathrm{op}}} X_E & \stackrel{\cong}{\longrightarrow} & \left(\lim_{E \in (\mathbf{L}_{/G}^{\mathrm{el}})^{\mathrm{op}}} X_E\right) \times \left(\lim_{E \in (\mathbf{L}_{/H}^{\mathrm{el}})^{\mathrm{op}}} X_E\right) \end{array}$$

We also have the full subcategory $\mathbf{L}_c \subset \mathbf{L}$ whose objects are the connected level graphs (a level graph G of height m is connected just when G_{0m} is a point). This subcategory inherits an active-inert factorization system from the larger category \mathbf{L} . The elementary objects in \mathbf{L}_c are precisely the elementary objects of \mathbf{L} and there is a corresponding notion of Segal \mathbf{L}_c -presheaf. Finally, there is the graphical category defined in [HRY15] whose objects are certain connected graphs (see [HRY15, Definition 6.46] or [CH22, Definition 2.2.11]). Following [CH22], we call this category \mathbf{G} , and it has its own notion of Segal presheaf. We will not need any low-level details about this category or its algebraic pattern structure.

There is a zig-zag of functors over **FinSet**^{op}_{*}



where the downward functors take sets of vertices.

Proposition 1.11. The functors $\iota^* \colon Psh(\mathbf{L}) \to Psh(\mathbf{L}_c)$ and $\tau^* \colon Psh(\mathbf{G}) \to Psh(\mathbf{L}_c)$ restrict to functors $Seg(\mathbf{L}) \to Seg(\mathbf{L}_c) \leftarrow Seg(\mathbf{G})$ which are equivalences of categories.

One can prove that ι^* is an equivalence on Segal presheaves by imitating the proof of Proposition 3.2.29 of [CH22], which is an ∞ -categorical analogue. Similarly, one can imitate the proof of Theorem 5.1.4 of [CH22] to show that τ^* : Seg(**G**) \rightarrow Seg(**L**_c) is an equivalence. However, the latter proof is very involved. Instead, we deduce both of these results by recognizing that Seg(**X**) is equivalent to the subcategory of *discrete objects* of the ∞ -categorical version Seg^{∞}(**X**) \subseteq Psh^{∞}(**X**) \coloneqq Fun(**X**^{op}, S) when **X** \in {**L**, **L**_c, **G**}. Recall from [Lur09, 5.5.6.2] that an object x in a quasicategory is called *discrete* if, for every object y, the Kan complex Map(y, x) is equivalent to a set. We utilize the more general notion of algebraic pattern from [CH21, Definition 2.1]; in that source Seg^{∞}(**X**) would be denoted by Seg_{**X**op}(S) and Seg(**X**) would be denoted by Seg_{**X**op}(Set).

Lemma 1.12. Let X be a category equipped with an active-inert factorization system and a class of elementary objects so that \mathbf{X}^{op} , together with this structure, is an algebraic pattern. There is a fully faithful functor $\operatorname{Seg}(\mathbf{X}) \hookrightarrow \operatorname{Seg}^{\infty}(\mathbf{X})$ whose essential image is the full subcategory of discrete objects.

Proof. Let Set \hookrightarrow be the fully-faithful inclusion of the ∞ -category of sets into that of spaces. As postcomposition with a fully faithful functor is fully faithful (see Remark 1.13 below), there is a fully faithful composite $Seg(\mathbf{X}) \hookrightarrow Psh(\mathbf{X}) \hookrightarrow$ $\operatorname{Psh}^{\infty}(\mathbf{X})$. This functor factors through $\operatorname{Seg}^{\infty}(\mathbf{X})$ since $\mathbf{Set} \hookrightarrow S$ preserves limits [Lur09, 5.5.6.18]. It is automatic that $\text{Seg}(\mathbf{X}) \hookrightarrow \text{Seg}^{\infty}(\mathbf{X})$ is fully faithful. The essential image of the functor consists of those Segal objects $F: \mathbf{X}^{\mathrm{op}} \to \mathcal{S}$ so that F(x) is discrete for all $x \in \mathbf{X}$. We now show this is the same thing as $F \in \operatorname{Seg}^{\infty}(\mathbf{X})$ being a discrete object.

Suppose $G: \mathbf{X}^{\mathrm{op}} \to S$ is an arbitrary presheaf, and write $G \simeq \operatorname{colim}_{\alpha} \mathfrak{L}(x_{\alpha})$, where $\mathfrak{k} \colon \mathbf{X} \to \mathrm{Psh}^{\infty}(\mathbf{X})$ is the Yoneda embedding. Then there are equivalences


$$\lim F(x_{\alpha}) \simeq \lim \operatorname{Map}(\mathcal{L}(x_{\alpha}), F) \simeq \operatorname{Map}(G, F) \simeq \operatorname{Map}(LG, F)$$

where L: $Psh^{\infty}(\mathbf{X}) \to Seg^{\infty}(\mathbf{X})$ is the localization functor guaranteed by [CH21, Lemma 2.11(iii)]. If F(x) is discrete for all $x \in \mathbf{X}$, then since **Set** is closed under limits, we see that Map(G, F) is discrete hence $F \in Seg^{\infty}(\mathbf{X})$ is discrete. Conversely, if F is a discrete object of $\text{Seg}^{\infty}(\mathbf{X})$ and $x \in \mathbf{X}$ is arbitrary, taking $G = \mathfrak{k}(x)$ in the string of equivalences above implies that F(x) is discrete. We have thus identified the essential image of $\text{Seg}(\mathbf{X}) \hookrightarrow \text{Seg}^{\infty}(\mathbf{X})$ with the discrete objects of $\text{Seg}^{\infty}(\mathbf{X})$.

Remark 1.13. It is well known that if $\mathcal{C} \to \mathcal{D}$ is a fully faithful functor of quasi-categories, then the postcomposition functor $\operatorname{Fun}(\mathcal{B},\mathcal{C}) \to \operatorname{Fun}(\mathcal{B},\mathcal{D})$ is also fully faithful. One way to verify involves first observing that [RV20, 5.6] and [RV22, 3.5.6(iv)] together imply that being fully faithful as a functor of quasicategories is equivalent to being fully faithful as a functor in the ∞ -cosmos of quasi-categories. Since this ∞ -cosmos is cartesian closed, the assertion follows from the characterization given in [RV22, 3.5.6(iii)].

Proof of Proposition 1.11. As a result of [Lur09, 5.5.6.28], an equivalence of quasicategories preserves and reflects discrete objects. Therefore it restricts to an equivalence of categories between the full subcategories of its domain and codomain. Proposition 3.2.29 and Theorem 5.1.4 of [CH22] show that ι and τ induce equivalences $\operatorname{Seg}^{\infty}(\mathbf{L}) \to \operatorname{Seg}^{\infty}(\mathbf{L}_{c}) \leftarrow \operatorname{Seg}^{\infty}(\mathbf{G})$. The result now follows from Lemma 1.12. \Box


Let $\chi_3: \mathbf{G} \to \mathbf{Ppd}$ be the functor which takes a graph to a properad freely generated by it; see [HRY15, §5.1.2]. We let $\chi_2 = \chi_3 \circ \tau$, and now describe χ_1 appearing in the following diagram.



If G is a height m level graph, we can decompose G into connected components $G \cong \coprod_{x \in G_{0m}} G_x$ (in Definition 4.6 we will fix a particular such isomorphism). The functor $\chi_1: \mathbf{L} \to \mathbf{Ppd}$ is defined by sending G to $\coprod \chi_2(G_x) = \coprod \chi_3 \tau(G_x)^2$.

²Let us describe χ_1 applied to a morphism $f: G \to H \cong \coprod_{y \in H_{0n}} H_y$ lying over $\alpha: [m] \to [n]$ in Δ . For $x \in G_{0m}$ write \bar{x} for the image of x under the composite $G_{0m} \to H_{\alpha(0)\alpha(m)} \to H_{0n}$. Then

We obtain from these functors the following commutative diagram:



with $N_i(P)_G = \hom(\chi_i(G), P)$; this commutes since if G is a connected level graph, then

$$\iota^* N_1(P)_G = N_1(P)_{\iota G} = \hom(\chi_1(\iota G), P) = \hom(\chi_2(G), P) = N_2(P)_G$$

and likewise for the other triangle. By [HRY15, Lemma 7.38], N_3 takes values in the subcategory of Segal objects.

Since we know that τ^* takes Segal objects to Segal objects and $N_2 = \tau^* \circ N_3$, we have that N_2 lands in the subcategory of Segal objects. This same argument does not apply to N_1 , so we must check the following:

Lemma 1.14. If P is a properad, then $N_1(P)$ is Segal.

Proof. Let $G \in \mathbf{L}$ be a height m level graph, and for $x \in G_{0m}$ let $G_x \in \mathbf{L}_c$ be the corresponding connected level graph with $\coprod_x G_x = G$ in the fiber \mathbf{L}_m . Notice that the canonical map $N_1(P)_G \to \prod_x N_1(P)_{G_x}$ is an isomorphism, using our description of χ_1 :

$$N_1(P)_G = \hom(\chi_1(G), P) = \hom\left(\coprod_x \chi_2(G_x), P\right) = \prod_x N_1(P)_{G_x}.$$

The map $\coprod_x \operatorname{Sc}(G_x) \to \operatorname{Sc}(G)$ is an isomorphism of presheaves, where $\operatorname{Sc}(G)$ is the Segal core of G. In the commutative square

we know that the bottom map becomes an isomorphism after applying $hom(-, N_1P)$. Hence to see that $N_1(P)$ is Segal it is enough to check that $hom(\mathfrak{k}(H), N_1P) \rightarrow hom(\operatorname{Sc} H, N_1P)$ is an isomorphism for connected graphs H. But this is true: if $H \in \mathbf{L}_c$ is connected, then $\iota_! \mathfrak{k}(H) = \mathfrak{k}(\iota H)$ and $\iota_! \operatorname{Sc}(H) = \operatorname{Sc}(\iota H)$, and by Segality of $N_2(P) = \iota^* N_1(P)$ the bottom map in the following square

is an isomorphism.

the composite $G_x \to G \to H$ factors through a map between connected level graphs $f_x \colon G_x \to H_{\bar{x}}$, and $\chi_1(f) \colon \coprod \chi_2(G_x) \to \coprod \chi_2(H_y)$ is induced from the properad maps $\chi_2(f_x)$.

Proposition 1.15. The functors N_1 and N_2 induces equivalences of categories

$$N_1: \mathbf{Ppd} \simeq \mathrm{Seg}(\mathbf{L})$$

 $N_2: \mathbf{Ppd} \simeq \mathrm{Seg}(\mathbf{L}_c).$

Proof. Each N_i lands in the full subcategory of Segal objects (using Lemma 1.14 for N_1), hence we have the commutative diagram

By Proposition 1.11, ι^* and τ^* are equivalences, and N_3 is an equivalence by [HRY15]. By 2-of-3, N_1 and N_2 are equivalences as well.

We will have no further need for L_c or G in this paper.

2. The symmetric monoidal envelope of a properad

The envelope of a properad is the prop freely generated by it. In this section we give a detailed description of this structure, in a way that will make our later comparisons more transparent. We take as an input a Segal **L**-presheaf, relying on the equivalence of categories from Proposition 1.15.

Remark 2.1. In [HRY17, §4.A], a left adjoint to the forgetful functor from props to properads is exhibited. We caution the reader that this is not quite the same as what we are doing here, since the kind of prop used in that paper is slightly weaker than the original version [Mac65, HR15]. See [BB17, Remark 10.5] and [HR17, Remark 3.5] for details.

2.1. Congruences of level graphs.

Definition 2.2. A congruence in \mathbf{L}_n is an isomorphism $G \to H$ so that for $0 \le i \le n$, the map $G_{ii} \to H_{ii}$ is an identity. We denote such a congruence by

$$G \xrightarrow{=} H.$$


We say that G and H are *congruent*, denoted $G \equiv H$, if there is a congruence between them.

If $G \in \mathbf{L}_0$, then the only congruence involving G is the identity. The same is true for the corollas $\mathfrak{c}_{n,m} \in \mathbf{L}_1$.

Lemma 2.3. If $\alpha : [m] \to [n]$ is a map in Δ and $G \to H$ is a congruence in \mathbf{L}_n , then $\alpha^*G \to \alpha^*H$ is a congruence in \mathbf{L}_m .

Proof. If $0 \leq i \leq j \leq m$, then $(\alpha^* G)_{ij} \to (\alpha^* H)_{ij}$ is equal to $G_{\alpha i,\alpha j} \to H_{\alpha i,\alpha j}$. When i = j this map is an identity by definition of congruence.

Remark 2.4. Suppose $G \in \mathbf{L}_1$ is a graph where every vertex has an input or output. Then the only congruence $G \xrightarrow{\equiv} G$ is the identity, as the hypothesis tells us the horizontal morphisms in the following diagram are epimorphisms



which implies that $G_{01} \to G_{01}$ is the identity as well. This is not to say that G is not involved in any congruence at all, only that if one permutes the elements of G_{01} then one must also alter at least one of the functions $G_{00} \to G_{01}$ and $G_{11} \to G_{01}$. Indeed, if $G_{01} = \underline{n}$, then there are n! different congruences with domain G.

Definition 2.5 (Congruence category). Let $\mathbf{K}_n \subseteq \mathbf{L}_n$ denote the wide subcategory consisting of the congruences. This defines a functor $\mathbf{K}_{\bullet} : \Delta^{\mathrm{op}} \to \mathbf{Gpd} \subseteq \mathbf{Cat}$, and we write $\mathbf{K} \to \Delta$ for the associated (right) fibration.

It will be useful to collapse further:


Definition 2.6. Let $W \rightarrow \Delta$ be the discrete fibration associated to the composite

$$\Delta^{\operatorname{op}} \overset{\mathbf{K}_{ullet}}{\longrightarrow} \operatorname{\mathbf{Gpd}} \overset{\pi_0}{\longrightarrow} \operatorname{\mathbf{Set}}.$$

That is, $\mathbf{W}_n = \pi_0(\mathbf{K}_n)$ is the set of height *n* level graphs modulo congruence. Note that \mathbf{W}_n is a discrete category, but **W** itself is not.

Remark 2.7. The simplicial set \mathbf{W}_{\bullet} turns out to be isomorphic to the nerve of a skeleton of **Csp**. Elements of \mathbf{W}_0 can be identified with the set $\{\underline{k}\}$ for $k \in \mathbb{N}$. Elements of \mathbf{W}_1 are congruence classes of height 1 level graphs, and this relation means these are exactly the same thing as morphisms in **Csp** between the sets appearing in $\{\underline{k}\}_{k\in\mathbb{N}}$. In Corollary 2.9 below we will formally show that \mathbf{W}_{\bullet} is Segal, and by inspection one can see that the compositions coincide in \mathbf{W}_{\bullet} and $N\mathbf{Csp}$. See Example 2.17 and Example 2.34 for more details.

The zig-zags $\mathbf{L}_n \hookrightarrow \mathbf{K}_n \twoheadrightarrow \pi_0(\mathbf{K}_n) = \mathbf{W}_n$ as *n* varies constitute natural transformations



(considering Set \subseteq Gpd \subseteq Cat). This yields the following commutative diagram of fibrations over Δ .

$$\mathbf{W} \underbrace{\overset{\pi}{\longleftarrow} \mathbf{K} \xrightarrow{i} \mathbf{L}}_{q \longrightarrow \mathbf{L}} \underbrace{\overset{\gamma}{\longleftarrow} \mathbf{L}}_{\mathbf{\Delta}}$$

We saw above that $\mathbf{L}_{\bullet} : \Delta^{\mathrm{op}} \to \mathbf{Cat}$ is Segal, meaning that the Segal morphisms $\mathbf{L}_n \to \mathbf{L}_1 \times_{\mathbf{L}_0} \cdots \times_{\mathbf{L}_0} \mathbf{L}_1$ are all equivalences of categories. The same holds for \mathbf{K}_{\bullet} .

Proposition 2.8. The simplicial category \mathbf{K}_{\bullet} is Segal. Moreover, the Segal morphisms are surjective on objects.

Proof. Let n > 1, 1 < k < n, and consider the pushout square in Δ_{int} whose vertical maps preserve last elements and horizontal maps preserve first elements.

$$\begin{bmatrix} 0 \\ \downarrow^k \\ [k] \xrightarrow{\alpha} \\ \end{bmatrix} \begin{bmatrix} n-k \\ \downarrow^{\beta} \\ [n] \end{bmatrix}$$

We must show

$$\alpha^* \times \beta^* \colon \mathbf{K}_n \to \mathbf{K}_k \times_{\mathbf{K}_0} \mathbf{K}_{n-k}$$

is an equivalence of categories. We already know that the map is surjective on objects since $\mathbf{L}_n \to \mathbf{L}_k \times_{\mathbf{L}_0} \mathbf{L}_{n-k}$ is. Since $\mathbf{L}_n \to \mathbf{L}_k \times_{\mathbf{L}_0} \mathbf{L}_{n-k}$ is fully faithful, it is enough to show that if $G, H \in \mathbf{K}_n$ are height *n* level graphs and $\alpha^* G \to \alpha^* H$ and $\beta^* G \to \beta^* H$ are congruences, then they define a congruence $G \to H$. But this is clear, since

$$(G_{ii} \to H_{ii}) = \begin{cases} (\alpha^* G)_{ii} \to (\alpha^* H)_{ii} & \text{if } 0 \le i \le k \\ (\beta^* G)_{i-k,i-k} \to (\beta^* H)_{i-k,i-k} & \text{if } k \le i \le n \end{cases}$$

which are identities.

Though $\pi_0: \mathbf{Gpd} \to \mathbf{Set}$ does not preserve pullbacks in general, we still have the following result.


Corollary 2.9. The simplicial set W_{\bullet} is Segal.

Proof. We use the same notation as in the previous proof. We are to show that the bottom map in the following square is a bijection.

$$\begin{array}{c} \mathbf{K}_n \xrightarrow{\simeq} & \mathbf{K}_k \times_{\mathbf{K}_0} \mathbf{K}_{n-k} \\ \downarrow & \downarrow \\ \pi_0(\mathbf{K}_n) \xrightarrow{\alpha^* \times \beta^*} & \pi_0(\mathbf{K}_k) \times_{\pi_0(\mathbf{K}_0)} \pi_0(\mathbf{K}_{n-k}) \end{array}$$

The bottom map is surjective, since the top-right composite is surjective on objects. Suppose $w, v \in \mathbf{W}_n$ map to the same element in $\mathbf{W}_k \times_{\mathbf{W}_0} \mathbf{W}_{n-k}$. Let $G, H \in \mathbf{K}_n$ be graphs in these path components. The assumption is that $\alpha^* G \equiv \alpha^* H$ and $\beta^* G \equiv \beta^* H$. Choose congruences witnessing these relations, giving a morphism $(\alpha^* G, \beta^* G) \to (\alpha^* H, \beta^* H)$ in $\mathbf{K}_k \times_{\mathbf{K}_0} \mathbf{K}_{n-k}$. Since the top map in the square above is an equivalence, this comes from a congruence in \mathbf{K}_n . Hence $G \equiv H$, that is, w = v.

2.2. The category CX. Our next task is to describe the envelope of a properad. For now, we content ourselves in producing this as a category, rather than as a symmetric monoidal category. More specifically, we produce a functor S as in the following diagram by using restriction along i and left Kan extension along $q\pi = pi$,



and show that it takes Segal presheaves to Segal presheaves. We write $C: \text{Seg}(\mathbf{L}) \to \mathbf{Cat}$ for the composite of S with the left adjoint of the nerve functor N.

Lemma 2.10. Given a commutative triangle of small categories



If q is a fibration and r is a discrete fibration, then p is a fibration.

Proof. If $b \in B$ is any object, then $B_{/b} \to A_{/rb}$ is an isomorphism. Indeed, it is bijective on objects since any $a \to rb$ has a unique lift $\tilde{a} \to b$. It is also fully faithful since a string $a_0 \to a_1 \to rb$ has a unique lift $\tilde{a}_0 \to \tilde{a}_1 \to b$ in B; but the composite $\tilde{a}_0 \to b$ is the unique lift of $a_0 \to rb$.

To show that p is a fibration it is enough to exhibit a right adjoint right inverse to $E_{/e} \rightarrow B_{/pe}$ for every object e (see [Gra66, Theorem 2.10] or [LR20, Theorem 2.2.2]). But since q is a fibration, for each e the functor $E_{/e} \rightarrow A_{/qe}$ admits a right adjoint right inverse. So choose one and this defines the desired functor $B_{/pe} \cong A_{/rpe} \rightarrow E_{/e}$.

Since $q\pi \colon \mathbf{K} \to \mathbf{\Delta}$ is a fibration and q is a discrete fibration, we have:

Lemma 2.11. The functor $\pi \colon \mathbf{K} \to \mathbf{W}$ is a Grothendieck fibration.

It is classical that left Kan extension along an opfibration may be computed by taking the colimit over the fibers (see [na]), which gives the following.

Lemma 2.12. If A is a **K**-presheaf, then its left Kan extension $\pi_!A$ is given by

$$(\pi_! A)_w = \operatorname{colim}_{G \in \mathbf{K}_w^{\operatorname{op}}} A_G$$

where $\mathbf{K}_w \subseteq \mathbf{K}$ is the relevant connected component of $\mathbf{K}_{q(w)}$. Likewise, if B is a **W**-presheaf, then

$$(q_!B)_n = \sum_{w \in \mathbf{W}_n} B_w.$$

Definition 2.13. Let S denote the composition

 $\operatorname{Psh}(\mathbf{L}) \xrightarrow{i^*} \operatorname{Psh}(\mathbf{K}) \xrightarrow{q_!\pi_!} \operatorname{Psh}(\mathbf{\Delta}).$

That is, if $X \in Psh(\mathbf{L})$, define $SX \in Psh(\mathbf{\Delta})$ as

$$SX = (q\pi)_! i^* X.$$

In light of Lemma 2.12, this means that

(1)
$$SX_n = \sum_{w \in \mathbf{W}_n} \operatorname{colim}_{G \in \mathbf{K}_w^{\operatorname{op}}} X_G.$$

We write \overline{X}_G for $\operatorname{colim}_{G \in \mathbf{K}_w^{\operatorname{op}}} X_G$, which is a quotient of X_G by the action of selfcongruences of G. We occasionally write \overline{X}_w for this set, where $w = [G] = \pi(G) \in \mathbf{W}$.

Remark 2.14. If every vertex of $G \in \mathbf{L}_1$ has an input or output, then the canonical map $X_G \to \overline{X}_G = \operatorname{colim}_{G \in \mathbf{K}_w^{\operatorname{op}}} X_G$ is an isomorphism by Remark 2.4.

Proposition 2.15. If $X \in \text{Seg}(\mathbf{L})$, then $SX \in \text{Seg}(\mathbf{\Delta})$.

Proof. Suppose we have a pushout square of non-identity maps in Δ_{int} whose vertical maps preserve last elements and horizontal maps preserve first elements.

$$\begin{array}{ccc} [0] & \searrow & [\ell] \\ \downarrow_{k} & & & \downarrow_{\beta} \\ [k] & \searrow & & [n] \end{array}$$

It is enough to show that SX sends pushouts of this form to pullbacks.

We use the description of SX_n from (1) above. Suppose we have two elements s, s' of SX_n , represented by (w, G, x) and (w', G', x'), which are sent to the same element of $SX_k \times_{SX_0} SX_\ell$ (Here G is a graph in the path component w of \mathbf{K}_n and x is an element of X_G with s the image of x under $X_G \to \operatorname{colim} X_H \subseteq SX_n$). Then w = w' by Corollary 2.9. Write $\alpha^* x \in X_{\alpha^* G}$, $\beta^* x \in X_{\beta^* G}$, $\alpha^* x' \in X_{\alpha^* G'}$, and $\beta^* x' \in X_{\beta^* G'}$ for the images of x and x'. Since $\alpha^* x$ and $\alpha^* x'$ represent the same element in the colimit

$$\operatorname{colim}_{H \in \mathbf{K}_{\alpha^* w}^{\operatorname{op}}} X_H$$

there is a congruence $\gamma: \alpha^* G \to \alpha^* G'$ with $\gamma^*(\alpha^* x') = \alpha^* x$. Likewise, there is a congruence $\delta: \beta^* G \to \beta^* G'$ with $\delta^*(\beta^* x') = \beta^* x$. By Proposition 2.8, there is a unique congruence $\chi: G \to G'$ with $\alpha^*(\chi) = \gamma$ and $\beta^*(\chi) = \delta$. This implies, in particular, that $G_{kk} = G'_{kk}$, and we write \underline{m} for this common value. The following diagram commutes and consists of bijections since X is a Segal **L**-presheaf (see Remark 1.10).

$$\begin{array}{cccc} X_{G'} & \longrightarrow & X_{\alpha^*G'} \times_{X_{\underline{m}}} X_{\beta^*G'} \\ \chi^* & & \downarrow \\ \chi_G & \longrightarrow & X_{\alpha^*G} \times_{X_m} X_{\beta^*G} \end{array}$$

It follows that $\chi^* x' = x$. Since $x \in X_G$ and $x' \in X_{G'}$ represent the same element in the colimit colim_{$H \in \mathbf{K}_w^{\text{op}} X_H$, it follows that s = s' in SX_n . We have thus shown the Segal map is injective.}

We now turn to surjectivity. Suppose $(s, s') \in SX_k \times_{SX_0} SX_\ell$, with s represented by (w, G, x) and s' represented by (w', G', x'). We have $G_{kk} = G'_{00} = \underline{m}$. Choose a height n level graph H with $\alpha^* H = G$ and $\beta^* H = G'$. Since X is a Segal **L**-presheaf, there is a (unique) element $y \in X_H$ which maps to (x, x') under

$$\alpha^* \times \beta^* \colon X_H \xrightarrow{\cong} X_G \times_{X_m} X_{G'}.$$

Now the element of SX_n represented by $(\pi(H), H, y)$ is sent to (s, s') by the Segal map.

Definition 2.16. If $X \in \text{Seg}(\mathbf{L})$, we write $CX = \tau_1(SX) \in \mathbf{Cat}$ for the associated category (where $\tau_1: \mathbf{sSet} \to \mathbf{Cat}$ is left adjoint to N). This defines a functor $C: \text{Seg}(\mathbf{L}) \to \mathbf{Cat}$ with NCX = SX.

The following example has previously been discussed in Remark 2.7, but it is extremely important for the rest of the paper and there is no harm in revisiting it. We will return to the monoidal structure in Example 2.34 below.

Example 2.17 (Canonical inclusion). If $* \in Psh(\mathbf{L})$ is the terminal presheaf, then there is an injective-on-objects equivalence $C(*) \to \mathbf{Csp}$. From the formula following Definition 2.13, we see S(*) is isomorphic to the simplicial set \mathbf{W}_{\bullet} . Then the objects of C(*) are identified with the height 0 level graphs, that is, with $ob(\mathbf{F}) = \{\underline{0}, \underline{1}, \underline{2}, \ldots\}$. The morphisms of C(*) are the elements of \mathbf{W}_1 , which under the identification on objects are precisely the same as the morphisms between these objects in \mathbf{Csp} . Finally, compositions in both categories are given by pushouts of equivalence classes of cospans.

2.3. Monoidal structure. In this section, we explain why CX is a strict monoidal category. It turns out to also be symmetric monoidal (see Theorem 2.33), but this is more difficult to show since it requires working at several simplicial levels of SX. The main result of this section is that if X is Segal, then SX is a (strict) monoid in **sSet**. This means (see Proposition 2.22) that CX is a monoid in **Cat**, that is, a strict monoidal category.

For each $n \geq 0$, the category $\mathbf{L}_n \subseteq \operatorname{Fun}(\mathscr{L}^n, \mathbf{F})$ is monoidal, using the ordinal sum in \mathbf{F} . More specifically, if G and H are height n level graphs, then G + H is the height n level graph with $(G+H)_{ij} = G_{ij} + H_{ij}$. The maps $G_{ij} \hookrightarrow G_{ij} + H_{ij} \leftrightarrow H_{ij}$ are order-preserving, and every element in G_{ij} appears before every element in H_{ij} . In fact, \mathbf{L}_n is actually a permutative category since \mathbf{F} is (i.e. a symmetric strict monoidal category [May74, Definition 4.1]). From Remark 1.10 we know that a Segal presheaf X takes sums of height n level graphs to products of sets, which we use in the following.

Definition 2.18. Suppose $X \in \text{Seg}(\mathbf{L})$, G and H are height n level graphs, and let $i: G \to G + H$ and $j: H \to G + H$ be the inclusions. Then define

$$-\otimes -: X_G \times X_H \to X_{G+H}$$

to be the inverse of the bijection $i^* \times j^* \colon X_{G+H} \to X_G \times X_H$.

We will make use of the following lemma in Section 2.4.

Lemma 2.19. Let $G, H \in \mathbf{L}_n$ and let $\sigma: H + G \to G + H$ be the flip map. If $x \in X_G$ and $y \in X_H$, then $\sigma^*(x \otimes y) = y \otimes x$.

Proof. The diagram

$$H \xrightarrow{i_L} H + G$$

$$\downarrow G$$

$$i_R \xrightarrow{i_R} G + H \xleftarrow{i_L} G$$

commutes in \mathbf{L}_n , which implies the diagram

$$\begin{array}{ccc} X_G \times X_H & \underset{i_L \times i_R^*}{\overset{\cong}{\underset{i_L \times i_R^*}}} X_{G+H} \\ & \downarrow^{\tau} & \downarrow^{\sigma^*} \\ X_H \times X_G & \underset{i_L \times i_R^*}{\overset{\cong}{\underset{i_L \times i_R^*}}} X_{H+G} \end{array}$$

commutes, where τ is the flip map. The result follows.

Lemma 2.20. Let $X \in \text{Seg}(\mathbf{L})$ and suppose that $G, H, K \in \mathbf{L}_n$. If $x \in X_G, y \in X_H$, and $z \in X_K$, then

$$(x \otimes y) \otimes z = x \otimes (y \otimes z)$$

in $X_{(G+H)+K} = X_{G+(H+K)}$. If \emptyset_n is the empty height n level graph and * is the unique element of X_{\emptyset_n} , then

$$* \otimes x = x = x \otimes *$$

in $X_{\varnothing_n+G} = X_G = X_{G+\varnothing_n}$.

Proof. Associativity follows from commutativity of the following rectangle.

This uses that the monoidal structure on \mathbf{L}_n is strict, so (G+H)+K = G+(H+K). For the second statement, the Segal map at \mathcal{O}_n

$$X_{\varnothing_n} \xrightarrow{\cong} \lim_{K \in (\mathbf{L}_{\mathrm{el}}^{\mathrm{op}})_{\varnothing_n/}} X_K$$

identifies X_{\varnothing_n} with a limit over the empty category. Hence there is indeed a unique element * in X_{\varnothing_n} . The Segal maps are

$$X_G = X_{\varnothing_n + G} \longrightarrow X_{\varnothing_n} \times X_G \qquad \qquad X_G = X_{G + \varnothing_n} \longrightarrow X_G \times X_{\varnothing_n}$$
$$x \longmapsto (*, x) \qquad \qquad x \longmapsto (x, *)$$

which establishes the result.

Lemma 2.21. Suppose $X \in \text{Seg}(\mathbf{L})$ and $n \ge 0$. The collection of functions \otimes from Definition 2.18 descend to a function

$$-\otimes -: SX_n \times SX_n \to SX_n$$

Proof. Recall the description of SX_n from (1) on page 16. Let $\bar{x} \in \overline{X}_w \subseteq SX_n$ and $\bar{x}' \in \overline{X}_{w'} \subseteq SX_n$. Choose a representative $x \in X_G$ and $x' \in X_{G'}$ for these elements and define $\bar{x} \otimes \bar{x}' \in SX_n$ to be the image of (x, x') under the following composite:

$$X_G \times X_{G'} \xleftarrow{\cong} X_{G+G'} \longrightarrow \overline{X}_{G+G'} \longleftrightarrow SX_n.$$

The leftward map is a bijection since X is Segal (Remark 1.10). This definition does not depend on our choice of representatives: indeed, suppose we have $(y, y') \in$ $X_H \times X_{H'}$ along with congruences $f: H \to G$ and $g: H' \to G'$ so that $f^*(x) = y$ and $g^*(x') = y'$. We then have a commutative diagram

$$\begin{array}{cccc} X_G \times X_{G'} & \stackrel{\cong}{\longleftarrow} & X_{G+G'} & \longrightarrow & \overline{X}_{G+G'} & \longrightarrow & SX_n \\ & & & & & & & \\ & & & & & & & \\ f^* \times g^* & & & & & & \\ & & & & & & & & \\ & X_H \times X_{H'} & \stackrel{\cong}{\longleftarrow} & X_{H+H'} & \longrightarrow & \overline{X}_{H+H'} & \longmapsto & SX_n \end{array}$$

which establishes that $\bar{x} \otimes \bar{x}'$ does not depend on choice of representative.

Proposition 2.22. If $X \in \text{Seg}(\mathbf{L})$, then CX is a strict monoidal category.

Proof. The assignment \otimes on objects and morphisms from Lemma 2.21 of CX is a functor $CX \times CX \to CX$. This follows because the simplicial structure maps $\alpha^* \colon \mathbf{L}_m \to \mathbf{L}_n$ preserve +. The tensor unit is the image of the unique element of X_{\varnothing_0} in $SX_0 = \operatorname{ob} CX$, and the tensor is strictly associative and unital by Lemma 2.20.

Essentially by definition of the monoidal structure, if $X \to Y$ is a map in Seg(L) then $CX \to CY$ is a strict monoidal functor.


2.4. The symmetry. In order to most efficiently produce the symmetry isomorphism and prove its properties, we first produce a more general twisted tensor of any two morphisms in CX, whose codomain is the opposite from the normal tensor. That is, if $f: x \to x'$ and $g: y \to y'$ are two morphisms, then we construct $f \otimes g: x \otimes y \to y' \otimes x'$. The symmetry isomorphism is then given in Definition 2.31 as $\operatorname{id}_x \otimes \operatorname{id}_y: x \otimes y \to y \otimes x$.

As an auxiliary useful construction, we first define twisted sums of height n graphs. Though these are defined in general, we only need them for $n \leq 2$ which can be illustrated directly. For $G, H \in \mathbf{L}_n$ height n level graphs and an integer t, define a new graph $G + H \in \mathbf{L}_n$ by

$$(G + H)_{ij} = \begin{cases} G_{ij} + H_{ij} & i < t \\ H_{ij} + G_{ij} & i \ge t. \end{cases}$$

The structure maps are constructed as sums and flips of sums of those appearing in G and H.

Example 2.23. Special cases include G + H = H + G and G + H = G + H. This covers all cases when n = 0; when n = 1 there is one interesting case G + H




while when n = 2 we have G + H

and G + H

Define the isomorphism

$$G \underset{t}{+} H \xrightarrow{\sigma_t} G + H$$

which is the identity in the ij component for i < t and is the flip map for $i \ge t$. We denote the inclusions by i_t and j_t , and the following diagram commutes in \mathbf{L}_n .



The following is an essential lemma about the behavior of this structure with respect to coface maps.

Lemma 2.24. If k < t, then $d_k(G + H) = (d_kG) + (d_kH)$ and the following diagram in **L** commutes.

$$\begin{aligned} (d_k G) &+ (d_k H) \xrightarrow{=} d_k (G + H) \xrightarrow{d^k} G + H \\ & \downarrow^{\sigma_{t-1}} & \downarrow^{d_k (\sigma_t)} & \downarrow^{\sigma_t} \\ d_k G + d_k H \xrightarrow{=} d_k (G + H) \xrightarrow{d^k} G + H \end{aligned}$$

If $k \geq t$, then $d_k(G + H) = (d_k G) + (d_k H)$ and the following diagram commutes.


More concisely,

$$\sigma_t d^k = \begin{cases} d^k \sigma_{t-1} & \text{if } k < t \\ d^k \sigma_t & \text{if } k \ge t. \end{cases}$$

Proof. This is a direct verification from the definitions.

Next is a special case which will be used in Lemma 2.29.

Lemma 2.25. If $G, H \in \mathbf{L}_n$ for $n \ge 1$, then the following diagram commutes.



The dashed map is the isomorphism $\sigma_2^{-1}\sigma_1$.

Proof. The first part of Lemma 2.24 with k = 1 < 2 = t tells us that $d^1\sigma_1 = \sigma_2 d^1$, while the second part with t = k = 1 tells us that $d^1\sigma_1 = \sigma_1 d^1$, and both maps having domain $d_1G + d_1H$.

Definition 2.26. Suppose $G, H \in \mathbf{L}_1, x \in X_G$, and $y \in X_H$. Let $x \otimes y \in X_{G+H}$ be the element defined by the following diagram

That is, $x \otimes y = (i_1^* \times j_1^*)^{-1}(x, y) = \sigma_1^*(x \otimes y).$

Proposition 2.27. The preceding definition descends to a well-defined function

 $-\tilde{\otimes} -: \operatorname{mor} CX \times \operatorname{mor} CX \to \operatorname{mor} CX.$

Proof. If $G' \xrightarrow{\equiv} G$ and $H' \xrightarrow{\equiv} H$ are congruences in \mathbf{K}_1 , then there is an associated congruence $G' + H' \xrightarrow{\equiv} G + H$ as well, and the following diagram commutes.

Thus we may define $[x] \otimes [y] \coloneqq [x \otimes y]$ with no ambiguity about choice of representatives.

Lemma 2.28. If $f: x_0 \to x_1$ and $g: y_0 \to y_1$ are morphisms in CX, then $f \otimes g$ is a map $x_0 \otimes y_0 \to y_1 \otimes x_1$.

Proof. We may prove the result by working with representatives. The diagram

commutes, using $d_0(G + H) = d_0G + d_0H = d_0H + d_0G$ along with Lemma 2.19 and Lemma 2.24. The two outer maps $X_G \times X_H \to SX_0$ are equal, implying that $\operatorname{cod}[x \ \tilde{\otimes} y] = (\operatorname{cod}[y]) \otimes (\operatorname{cod}[x])$. A similar, but simpler diagram can be constructed for d_1 to see dom $[x \ \tilde{\otimes} y] = (\operatorname{dom}[x]) \otimes (\operatorname{dom}[y])$; it is simpler since $d_1(G + H) = d_1G + d_1H = d_1G + d_1H$.

Lemma 2.29. Given morphisms

$$\begin{array}{cccc} x_0 & \xrightarrow{f_1} & x_1 & \xrightarrow{f_2} & x_2 \\ y_0 & \xrightarrow{g_1} & y_1 & \xrightarrow{g_2} & y_2 \end{array}$$

in CX, we have

$$(f_2 \ \tilde{\otimes} \ g_2) \circ (f_1 \otimes g_1) = (f_2 f_1) \ \tilde{\otimes} \ (g_2 g_1) = (g_2 \otimes f_2) \circ (f_1 \ \tilde{\otimes} \ g_1).$$

Proof. The composable pairs given are elements in SX_2 , and hence we can find height 2 level graphs G and H and elements $x \in X_G$ and $y \in X_H$ so that $d_2x \in X_{d_2G}$ represents f_1 and $d_0x \in X_{d_0G}$ represents f_2 , and likewise for y and g_1, g_2 . We have $d_1(G + H) = (d_1G) + (d_1G) = d_1(G + H)$. Let $\sigma_1 \colon G + H \to G + H$ and

 $\sigma_2\colon G+H\to G+H$ be the partial flipping morphisms from above. By Lemma 2.25 we have

$$d_1\sigma_1^*(x \otimes y) = d_1\sigma_2^*(x \otimes y) \in X_{d_1(G_1+H)} = X_{d_1(G_2+H)}.$$

We claim that this element represents all three maps; we use freely the identities from Lemma 2.24. First, the equalities

 $d_1\sigma_1^*(x\otimes y) = \sigma_1^*d_1(x\otimes y) = \sigma_1^*((d_1x)\otimes (d_1y))$

show this element represents $(f_2 f_1) \otimes (g_2 g_1)$. The equations (the first of which uses $\sigma_0 = \sigma$ and Lemma 2.19)

$$d_0\sigma_1^*(x\otimes y) = \sigma_0^*d_0(x\otimes y) = \sigma_0^*(d_0x\otimes d_0y) = \sigma^*(d_0x\otimes d_0y) = d_0y\otimes d_0x$$
$$d_2\sigma_1^*(x\otimes y) = \sigma_1^*d_2(x\otimes y) = \sigma_1^*(d_2x\otimes d_2y) = d_2x\tilde{\otimes}d_2y$$

tell us our element represents $(g_2 \otimes f_2) \circ (f_1 \otimes g_1)$. Finally, we have

$$d_0\sigma_2^*(x\otimes y) = \sigma_1^*d_0(x\otimes y) = \sigma_1^*(d_0x\otimes d_0y) = d_0x\,\tilde{\otimes}\,d_0y$$
$$d_2\sigma_2^*(x\otimes y) = \sigma_2^*d_2(x\otimes y) = \sigma_2^*(d_2x\otimes d_2y) = d_2x\otimes d_2y$$

so the element represents $(f_2 \otimes g_2) \circ (f_1 \otimes g_1)$.

Lemma 2.30. Given morphisms

$$\begin{array}{cccc} x_0 \stackrel{f_1}{\longrightarrow} x_1 \stackrel{f_2}{\longrightarrow} x_2 \\ y_0 \stackrel{g_1}{\longrightarrow} y_1 \stackrel{g_2}{\longrightarrow} y_2 \end{array}$$

in CX, we have

$$(g_2 \otimes f_2) \circ (f_1 \otimes g_1) = (f_2 f_1) \otimes (g_2 g_1).$$

Proof. Use the same notation from the beginning of the previous proof. We introduce a new graph G + H as displayed:

Define $\tilde{\sigma}: \tilde{G+H} \to G + H$ which is the flip map in components 11 and 12, and the identity elsewhere. Our strategy is to show that

$$d_1\tilde{\sigma}^*(x\otimes y)\in X_{d_1(G\tilde{+}H)}$$

represents both maps in the statement of the lemma.

Notice that we have $d_0(G\tilde{+}H) = (d_0H) + (d_0G)$, $d_1(G\tilde{+}H) = d_1G + d_1H$, and $d_2(G\tilde{+}H) = (d_2G) + (d_2H)$. Furthermore, the following diagrams commute:

$$\begin{array}{c} (d_0H) + d_0G \xrightarrow{\sigma_1} d_0H + d_0G \xrightarrow{\sigma} d_0G + d_0H \\ \downarrow^{d^0} & \downarrow^{d^0} \\ G\tilde{+}H \xrightarrow{\tilde{\sigma}} G + H \end{array}$$

$$\begin{array}{cccc} d_1G + d_1H & & d_2G + d_2H \xrightarrow{\sigma_1} d_2G + d_2H \\ & & \downarrow^{d^1} & & \downarrow^{d^2} & & \downarrow^{d^2} \\ G\tilde{+}H \xrightarrow{\tilde{\sigma}} & G + H & & G\tilde{+}H \xrightarrow{\tilde{\sigma}} & G + H \end{array}$$

Now this comes down to computation: from the above diagrams and Lemma 2.19,

$$d_0\tilde{\sigma}^*(x\otimes y) = \sigma_1^*\sigma^*d_0(x\otimes y) = \sigma_1^*\sigma^*(d_0x\otimes d_0y) = \sigma_1^*(d_0y\otimes d_0x) = d_0y\,\tilde{\otimes}\,d_0x$$
$$d_2\tilde{\sigma}^*(x\otimes y) = \sigma_1^*d_2(x\otimes y) = \sigma_1^*(d_2x\otimes d_2y) = d_2x\,\tilde{\otimes}\,d_2y$$

so the element indeed represents $(g_2 \otimes f_2) \circ (f_1 \otimes g_1)$. On the other hand,

 $d_1\tilde{\sigma}^*(x\otimes y) = d_1(x\otimes y) = d_1x\otimes d_1y$

so the element represents $(f_2f_1) \otimes (g_2g_1)$.

Definition 2.31. If $x, y \in \text{ob} CX$, define $\tau_{x,y} \colon x \otimes y \to y \otimes x$ to be $\text{id}_x \otimes \text{id}_y$.

Proposition 2.32. For any $x, y \in \text{ob } CX$, we have $\tau_{y,x} \circ \tau_{x,y} = \text{id}_{x \otimes y}$. If $f: x \to x'$ and $g: y \to y'$ are two morphisms in CX, then the diagram

$$\begin{array}{c} x \otimes y \xrightarrow{\tau_{x,y}} y \otimes x \\ f \otimes g \downarrow & \downarrow g \otimes f \\ x' \otimes y' \xrightarrow{\tau_{x',y'}} y' \otimes x' \end{array}$$

commutes.

Proof. The first statement follows from Lemma 2.30, since

$$\tau_{y,x} \circ \tau_{x,y} = (\mathrm{id}_y \otimes \mathrm{id}_x) \circ (\mathrm{id}_x \otimes \mathrm{id}_y) = (\mathrm{id}_x \mathrm{id}_x) \otimes (\mathrm{id}_y \mathrm{id}_y) = \mathrm{id}_{x \otimes y}.$$

The second statement similarly follows from Lemma 2.29 since both maps are equal to $f \otimes g$. Indeed, we first have

$$\tau_{x',y'}(f \otimes g) = (\mathrm{id}_{x'} \ \tilde{\otimes} \ \mathrm{id}_{y'}) \circ (f \otimes g) = (\mathrm{id}_{x'} \ f) \ \tilde{\otimes} \ (\mathrm{id}_{y'} \ g)$$

utilizing the first equality, and then we have

$$(g \otimes f)\tau_{x,y} = (g \otimes f) \circ (\mathrm{id}_x \ \tilde{\otimes} \ \mathrm{id}_y) = (f \ \mathrm{id}_x) \ \tilde{\otimes} \ (g \ \mathrm{id}_y)$$

using the second equality.

Theorem 2.33. The functor $C: \text{Seg}(\mathbf{L}) \to \mathbf{Cat}$ factors through the category **Perm** of small permutative categories and strict symmetric monoidal functors. More precisely, this uses the monoidal structure on CX from Proposition 2.22 and the symmetry isomorphism from Definition 2.31.

Proof. By Proposition 2.22 and Proposition 2.32, CX is a permutative category for every $X \in \text{Seg}(\mathbf{L})$. But the definitions imply that both the monoidal structure and the symmetry are natural in X.

Example 2.34. The equivalence $C(*) \to \mathbf{Csp}$ from Example 2.17 is a symmetric monoidal functor. To see this, recall that objects of C(*) have been identified with objects of \mathbf{F} and morphisms are equivalence classes of cospans in \mathbf{F} . Direct inspection shows that the symmetry isomorphisms in C(*) and \mathbf{F} coincide. The monoidal constraint for $C(*) \to \mathbf{Csp}$ is then inherited from that for $\mathbf{F} \to \mathbf{FinSet}$, and we conclude that $C(*) \to \mathbf{Csp}$ is a symmetric monoidal functor. That \mathbf{Csp} is equivalent to a skeletal permutative category is well known; see [Lac04, 5.4].

Lack shows that C(*) is the prop for special Frobenius monoids [CF17, Proposition 6.1], also known as commutative separable algebras [Car91]. The introduction of [RSW06] contains a nice overview of these structures.

It seems to be a folklore result that special Frobenius monoids are also the algebras for the terminal properad. This is consistent with the fact that C gives the free prop generated by a properad.

3. PROPERADS GIVE LABELLED COSPAN CATEGORIES

In this section, we show that the envelope of a properad is a labelled cospan category. More precisely, if P is a properad then the composite

$$C(N_1(P)) \to C(*) \xrightarrow{\simeq} \mathbf{Csp}$$

is a labelled cospan category. Since N_1 is an equivalence of categories between **Ppd** and Seg(**L**), it is enough to prove the following. Below, we will verify the axioms from Definition 1.2 in a series of lemmas.

Proposition 3.1. If $X \in \text{Seg}(\mathbf{L})$, then $C(X) \to \mathbf{Csp}$ is a labelled cospan category. *Proof.* Combine Lemma 3.2, Lemma 3.4, Lemma 3.7, and Lemma 3.8.

Suppose $X \in \text{Seg}(\mathbf{L})$, and consider functor $CX \to C(*)$. The set

$$X_{\mathfrak{e}} = X_{\underline{1}} \subseteq \sum_{k \in \mathbb{N}} X_{\underline{k}} = \sum_{w \in \mathbf{W}_0} \overline{X}_w = \operatorname{ob} CX$$

is the set of *connected objects* of CX in the sense of Definition 1.1. Likewise, the set

$$\sum_{n,m\geq 0} X_{\mathfrak{c}_{n,m}} \subseteq \sum_{w\in \mathbf{W}_1} \overline{X}_w = \operatorname{mor} CX$$

is the set of *connected morphisms*. These displays use that the component of $\underline{k} \in \mathbf{K}_0$ and the component of $\mathbf{c}_{n,m} \in \mathbf{K}_1$ are discrete. The *reduced morphisms* are those on (congruence classes) of height 1 level graphs so that each vertex has either an input or an output. Alternatively, these graphs are those without a \mathbf{c}_{00} -summand.

Lemma 3.2. Let c be an object of CX so that $\pi(c) = \underline{n}$. Then $c = c_1 \otimes \cdots \otimes c_n$ for some connected objects c_1, \ldots, c_n .

Proof. Recall that CX is a strict monoidal category, and on objects the *n*-fold tensor product is given as the inverse of the Segal map

$$X_n \xrightarrow{\cong} X_1 \times \cdots \times X_1$$

Thus given an object c with $\pi(c) = \underline{n}$, we see that c is the tensor product of n connected objects c_1, \ldots, c_n .

Remark 3.3 (Uniqueness of decomposition). We emphasize that as the *n*-fold tensor product is given by the inverse of the Segal map $X_{\underline{n}} \xrightarrow{\cong} X_{\underline{1}} \times \cdots \times X_{\underline{1}}$ there is actually a *unique* list of connected objects c_1, \ldots, c_n with $c = c_1 \otimes \ldots \otimes c_n$ in the previous lemma.


See also Definition 4.2 below.

Lemma 3.4. The abelian monoid hom(1, 1) is freely generated by the set hom^c(1, 1) of connected morphisms.

Proof. The set of objects of L_1 with empty input and empty output is in bijection with \mathbb{N} , and we write temporarily abbreviate the object corresponding with n by

$$n\mathfrak{c}_{00} \coloneqq \underbrace{\mathfrak{c}_{00} + \cdots + \mathfrak{c}_{00}}_{n}$$

This object is not isomorphic, hence not congruent, to any other object in \mathbf{L}_1 . Every automorphism of $n\mathfrak{c}_{00}$ is a congruence, and the group of automorphisms is the symmetric group Σ_n . For $\gamma \in \Sigma_n$, the left diagram below commutes



hence so too does the right diagram, where π_t is the *t*th projection. This implies that the following commutes

$$\begin{array}{ccc} X_{n\mathfrak{c}_{00}} & \stackrel{\cong}{\longrightarrow} & \prod_{i=1}^{n} X_{\mathfrak{c}_{00}} \\ & & \uparrow \\ & & \uparrow \\ & & \uparrow \\ X_{n\mathfrak{c}_{00}} & \stackrel{\cong}{\longrightarrow} & \prod_{i=1}^{n} X_{\mathfrak{c}_{00}} \end{array}$$

where the right-hand map is the standard right action on lists, that is $(x_1, \ldots, x_n) \mapsto (x_1, \ldots, x_n) \cdot \gamma = (x_{\gamma(1)}, \ldots, x_{\gamma(n)})$. Further, the monoidal structure comes from the bottom right isomorphism in the following diagram.

Thus we see that hom(1, 1) as a monoid is isomorphic to

$$\sum_{n\geq 0} \left(\prod_{i=1}^n X_{\mathfrak{c}_{00}}\right) / \Sigma_n^{\mathrm{op}}$$

which is the free abelian monoid on $X_{\mathfrak{c}_{00}} = \hom^{c}(\mathbf{1}, \mathbf{1}).$

Let $R = \operatorname{mor}^{r} C(*) \subseteq \mathbf{W}_{1}$ be the congruence classes of graphs where each vertex has an input or output. The following lemma, concerning the connected components of the *q*-fibers in the congruence category **K** appearing in Lemma 2.12, will be used in the proofs of Lemma 3.7 and Proposition 3.9. If $r \in R$, then by Remark 2.4 the category \mathbf{K}_{r} is a preorder, and $\overline{X}_{r} \cong X_{G}$ if [G] = r (see Remark 2.14).

Lemma 3.5. If $r \in R \subseteq \mathbf{W}_1$ is reduced and $w \in \mathbf{W}_1$ is arbitrary, then

(2)
$$\mathbf{K}_r \times \mathbf{K}_w \to \mathbf{K}_{r \otimes w}$$


sending (G, H) to G+H is an equivalence of categories. It follows that if $X \in \text{Seg}(\mathbf{L})$, then $\overline{X}_r \times \overline{X}_w \to \overline{X}_{r\otimes w}$ is a bijection.

Proof. By the definition in Lemma 2.21, the class $r \otimes w \in \mathbf{W}_1$ contains some graph G+H where [G] = r and [H] = w. Thus (2) is essentially surjective. It is also clearly faithful. We now prove that it is full. We must show that if $G, G' \in r$ and $H, H' \in w$ then any congruence $f: G + H \equiv G' + H'$ restricts to congruences $f^L: G \equiv G'$ and $f^R: H \equiv H'$ with $f = f^L + f^R$. For i = 0, 1 we know that $\mathrm{id}_{G_{ii}}: G_{ii} \to G'_{ii}$ and $\mathrm{id}_{H_{ii}}: H_{ii} \to H'_{ii}$ sum to $f_{ii} = \mathrm{id}_{G_{ii}+H_{ii}}: (G + H)_{ii} \to (G + H')_{ii}$, so this part restricts. Since every vertex of G or G' has either an input or an output, the first horizontal maps in the following commutative rectangle are surjective.

It follows that f_{01} restricts to a function $f_{01}^L: G_{01} \to G'_{01}$. We must still check that f^L is a congruence, and for this simply note that $(f^{-1})_{01}$ restricts to a function $G'_{01} \to G_{01}$ by the same argument, providing an inverse to f_{01}^L . Since f^L is an isomorphism, it is a congruence.

Now the composite $H \to G + H \to G' + H'$ must land in H', since f is a monomorphism and $f^L: G \to G'$ is an epimorphism. Thus f restricts to a map $f^R: H \to H'$, which must be an epimorphism since f is. We conclude that (2) is full, hence an equivalence.

For the conclusion, suppose X is Segal. In the following diagram, we use the result to deduce the isomorphism at the bottom of the triangle; the others come from Segality.



We conclude that the dotted arrow is an isomorphism.

Remark 3.6. The proof of Lemma 3.5 truly depended on one of the elements being reduced. The simplest issue was apparent in Lemma 3.4 and its proof, since

$$X_{\mathfrak{c}_{00}} \times X_{\mathfrak{c}_{00}} = \overline{X}_{\mathfrak{c}_{00}} \times \overline{X}_{\mathfrak{c}_{00}} \to \overline{X}_{\mathfrak{c}_{00}+\mathfrak{c}_{00}} = X_{\mathfrak{c}_{00}+\mathfrak{c}_{00}} / \Sigma_2^{\mathrm{op}} \cong (X_{\mathfrak{c}_{00}} \times X_{\mathfrak{c}_{00}}) / \Sigma_2^{\mathrm{op}}$$

is not typically a bijection. But this also occurs for other congruence classes. For example, if $G = \mathfrak{c}_{10} + \mathfrak{c}_{00}$ and $H = \mathfrak{c}_{01} + \mathfrak{c}_{00}$, then the non-trivial endo-congruence




FIGURE 3. Graph decomposition from Lemma 3.7

on $G + H = \mathfrak{c}_{10} + \mathfrak{c}_{00} + \mathfrak{c}_{01} + \mathfrak{c}_{00}$ does not restrict in the way of the proof. Further, the automorphism group of G + H in \mathbf{K}_1 is Σ_2 , whereas G and H have trivial automorphism groups.

Lemma 3.7. The map

$$\hom^{\mathrm{r}}(c,d) \times \hom(\mathbf{1},\mathbf{1}) \xrightarrow{\otimes} \hom(c,d)$$

is a bijection.

Proof. Let $R = \operatorname{mor}^{r} C(*) \subseteq \mathbf{W}_{1}$ be the congruence classes of graphs where each vertex has an input or output, and $T = \{n\mathbf{c}_{00} \mid n \in \mathbb{N}\} \subseteq \mathbf{W}_{1}$ be the set of graphs with empty inputs and outputs (see proof of Lemma 3.4). Every element $w \in \mathbf{W}_{1}$ decomposes uniquely as $w = r \otimes t$ where $r \in R$ and $t \in T$. Indeed, to get this decomposition, if G is an arbitrary height 1 level graph, then we can form new graphs G^{r} and G^{t} and morphisms in \mathbf{L}_{1}


$$G^r \stackrel{f}{\longleftrightarrow} G \stackrel{g}{\longleftrightarrow} G^t$$

so that every vertex of G^r has an input or output, $f_{ii}: G^r_{ii} \to G_{ii}$ is an identity, $g_{ii}: \underline{0} \to G_{ii}$ is the unique map, and f_{01}, g_{01} are order-preserving. See Figure 3 for an illustration. There is a unique congruence $G^r + G^t \equiv G$ taking f and gto the usual inclusions into the sum. If $H^r + H^t \equiv G$ is some other congruence with $[H^r] \in R$ and $[H^t] \in T$, then the composite $H^r + H^t \to G \to G^r + G^t$ takes every vertex in H^r to a vertex in G^r and every vertex in H^t to a vertex in G^t , hence gives congruences $H^r \equiv G^r$ and $H^t \equiv G^t$. This establishes uniqueness of the decomposition, and we conclude that $\mathbf{W}_1 \cong R \times T$. Coupling this with Lemma 3.5, we conclude that the following

$$\operatorname{mor}^{\mathrm{r}} C(X) \times \operatorname{hom}(\mathbf{1}, \mathbf{1}) = \sum_{R} \overline{X}_{r} \times \sum_{T} \overline{X}_{t} = \sum_{R \times T} \overline{X}_{r} \times \overline{X}_{t} \xrightarrow{\cong} \sum_{\mathbf{W}_{1}} \overline{X}_{w} = \operatorname{mor} C(X)$$

is a bijection.

The large rectangle and bottom square in the following are pullbacks,



hence so is the upper square. The result follows.

Lemma 3.8. For each four objects c, d, c', d' in C, the following square is cartesian.

This follows from a different result.

Proposition 3.9. If $X \to Y$ is a morphism in Seg(L), then

is cartesian.

Proof. Let $R = \operatorname{mor}^{\mathrm{r}} C(*) \subseteq \mathbf{W}_1$ be the congruence classes of height 1 level graphs where each vertex has an input or output, so that

$$\operatorname{mor}^{\mathrm{r}} C(X) = \sum_{r \in R} \overline{X}_r.$$

By Lemma 3.5, there is a canonical bijection

$$\operatorname{mor}^{\mathbf{r}} C(X) \times \operatorname{mor}^{\mathbf{r}} C(X) = \sum_{r \in R} \overline{X}_r \times \sum_{r' \in R} \overline{X}_{r'} \cong \sum_{(r,r') \in R \times R} \overline{X}_r \times \overline{X}_{r'} \cong \sum_{(r,r') \in R \times R} \overline{X}_{r \otimes r'}.$$

Thus it suffices to observe that the following square,

$$\begin{array}{ccc} \sum\limits_{r,r'\in R} \overline{X}_{r\otimes r'} & \longrightarrow & \sum\limits_{s\in R} \overline{X}_s \\ & & \downarrow & & \downarrow \\ & & \downarrow & & \downarrow \\ \sum\limits_{r,r'\in R} \overline{Y}_{r\otimes r'} & \longrightarrow & \sum\limits_{s\in R} \overline{Y}_s \end{array}$$


whose horizontal legs are induced by $\otimes : R \times R \to R$, is cartesian.

Proof of Lemma 3.8. Apply the previous lemma in the case when Y = *, using that $C(*) \to \mathbf{Csp}$ is fully faithful. Suppose $f: \pi c \to \pi d$, $g: \pi c' \to \pi d'$, and $h: c \otimes c' \to d \otimes d'$ are reduced morphisms with $\pi(h) = f \otimes g$. By the previous lemma, we know there are unique maps $\tilde{f}: a \to b$ and $\tilde{g}: a' \to b'$ with $\tilde{f} \otimes \tilde{g} = h$, $\pi(\tilde{f}) = f$, and $\pi(\tilde{g}) = g$. The only thing to verify is that a = c, b = d, a' = c' and b' = d', but this follows from the uniqueness of decompositions of objects of CX into connected objects (as was observed in Remark 3.3).

Proposition 3.10. The functor $C: \operatorname{Seg}(\mathbf{L}) \to \mathbf{Cat}$ is faithful.

Proof. If $X, Y \in \text{Seg}(\mathbf{L})$, then a map $X \to Y$ is uniquely determined by its action on elementary objects $X_{\mathfrak{e}} \to Y_{\mathfrak{e}}$ and $X_{\mathfrak{c}_{p,q}} \to Y_{\mathfrak{c}_{p,q}}$ (as p and q vary). If $f_0, f_1 \colon X \to Y$

are two maps in Seg(L), then the following diagrams commute for i = 0, 1.



Thus if $C(f_0) = C(f_1)$, then $f_0 = f_1$.

This proposition implies that the functors from $Seg(\mathbf{L})$ to permutative categories and labelled cospan categories are also faithful.

4. Strict labelled cospan categories

In this section we identify a special kind of labelled cospan category, and show that to each we may associate a Segal L-presheaf (and hence a properad). This will provide an inverse to our previous construction: the colors of this properad will be the connected objects, and the operations in the associated properad will be the connected morphisms. Unfortunately, actually proving this gives a properad comes with substantial challenges.

We begin with an explanatory example.

Example 4.1. From the category C(*), create a new category C with a single additional object $\underline{1}'$ which is isomorphic to $\underline{1} \in C(*)$. As a permutative category, it will satisfy the equation $\underline{1}' \otimes \underline{n} = \underline{1} \otimes \underline{n}$ for $n \ge 1$ and the equation $\underline{1}' \otimes \underline{1}' = \underline{2}$. There is an evident functor $C \to C(*) \to \mathbf{Csp}$ sending the new object to <u>1</u>. By Remark 3.3 this labelled cospan category is not C(X) for any $X \in \text{Seg}(\mathbf{L})$, since $\underline{2} = \underline{1} \otimes \underline{1} = \underline{1}' \otimes \underline{1} = \underline{1} \otimes \underline{1}' = \underline{1}' \otimes \underline{1}'$ admits several distinct decompositions into connected objects.

Thus not all labelled cospan categories arise from properads (though they do up to equivalence). We now isolate the image of the functor C. Recall that a strict map of permutative categories is a strict monoidal functor which is compatible with the symmetries.

Definition 4.2. A strict labelled cospan category is a strict map of permutative categories $\pi: C \to C(*)$ satisfying the conditions of Definition 1.2 along with

(1) If $\pi(c) = \underline{n}$, then there are unique connected objects c_1, \ldots, c_n with c = $c_1 \otimes \cdots \otimes c_n$.

Maps are strict maps of permutative categories so that the triangle

$$C \xrightarrow{} C'$$

$$C(*)$$

commutes. We write **SLCC** for the associated category.

Alternatively, (1) can be rephrased as follows, which means that each strict labelled cospan category is a (colored) prop of a special type [HR15].

(1") The monoid ob(C) is the free monoid on the set of connected objects $ob^{c}(C)$.

One can show (see Proposition 6.13) that being a strict labelled cospan category is a *property* of a permutative category, rather than a *structure*. Indeed, if C is a permutative category, there is at most one functor $\pi: C \to C(*)$ making C into a strict labelled cospan category. Nevertheless, we will often require the map π , so we do not attempt a definition free from it.

Corollary 5.9 states that the category of properads is equivalent to **SLCC**. We know that $\mathbf{Ppd} \simeq \mathrm{Seg}(\mathbf{L})$ by Proposition 1.15, and we have already exhibited a functor $\text{Seg}(\mathbf{L}) \rightarrow \mathbf{SLCC}$ by Proposition 3.1 and Remark 3.3. Proposition 3.10 implies that this functor is faithful. It remains to show this functor is full and essentially surjective.

Suppose $C \xrightarrow{\pi} C(*)$ is in **SLCC**. We will define a presheaf $P = P(\pi) \in Psh(\mathbf{L})$ from this data, and later show that $CP \to C(*)$ is isomorphic to $C \to C(*)$, proving that $Seg(\mathbf{L}) \to SLCC$ is essentially surjective.

4.1. The inert part. We first define the restriction of the presheaf to L_{el} ; it is slightly more convenient to define this also on arbitrary height zero level graphs. We define $P_{\underline{n}}$ and $P_{\mathfrak{c}_{m,n}}$ as subsets of the objects of C and the *connected* morphisms of C so that the following diagrams are pullbacks.

Notice that $P_{\mathfrak{e}} = P_{\underline{1}}$ is precisely the set of connected objects in C, and there is a bijection

(4)
$$P_{\underline{n}} \xrightarrow{\cong} P_{\underline{1}} \times \cdots \times P_{\underline{1}}$$

that takes an object to the list of connected objects from (1') in Definition 4.2.

We must still define the action of our restriction on inert morphisms. Between the elementary objects of **L**, there are the following inert morphisms:

• For $1 \le i \le m$ the following composite

$$\underline{1} \xrightarrow{\neg i \neg} \underline{m} = d_1(\mathfrak{c}_{m,n}) \xrightarrow{d^1} \mathfrak{c}_{m,n}$$

which we call ℓ_i .

• For $1 \le j \le n$ the following composite

$$\underline{1} \stackrel{\lceil j \rceil}{\longrightarrow} \underline{n} = d_0(\mathfrak{c}_{m,n}) \stackrel{d^0}{\longrightarrow} \mathfrak{c}_{m,n}$$

which we call r_i .

• Automorphisms $f = (\gamma, \chi) : \mathfrak{c}_{m,n} \to \mathfrak{c}_{m,n}$, where $\gamma = d_1(f) : \underline{m} \to \underline{m}$ and $\chi = d_0(f) : \underline{n} \to \underline{n}$ are bijections.

The only relations among the above these morphisms are the following:

- $(\gamma, \chi) \circ \ell_i = \ell_{\gamma(i)}$
- $(\gamma, \chi) \circ r_j = r_{\chi(j)}$ $(\gamma, \chi) \circ (\gamma', \chi') = (\gamma \gamma', \chi \chi').$

Notation 4.3. If $\sigma: \underline{k} \to \underline{k}$ is a bijection, and c_1, \ldots, c_k are connected objects of C then we write $\hat{\sigma}: c_1 \otimes \cdots \otimes c_k \to c_{\sigma^{-1}(1)} \otimes \cdots \otimes c_{\sigma^{-1}(k)}$ for the map determined by the symmetry isomorphism. Then $\hat{\gamma} \circ \hat{\sigma} = \widehat{\gamma \circ \sigma}$.

Definition 4.4. Suppose that $p: c_1 \otimes \ldots \otimes c_m \to b_1 \otimes \ldots \otimes b_n$ is in $P_{\mathfrak{c}_{m,n}} \subseteq \operatorname{mor}^c C$. We define $\ell_i^*(p) = c_i \in P_{\mathfrak{c}}$ and $r_j^*(p) = b_j \in P_{\mathfrak{c}}$. If $f = (\gamma, \chi): \mathfrak{c}_{m,n} \to \mathfrak{c}_{m,n}$ is an isomorphism, we define $f^*p = (\gamma, \chi)^*(p)$ to be the following composite of p with symmetry isomorphisms.

$$c_{\gamma(1)} \otimes \ldots \otimes c_{\gamma(m)} \qquad b_{\chi(1)} \otimes \ldots \otimes b_{\chi(n)} \\ \downarrow^{\hat{\gamma}} \qquad \hat{\chi}^{-1} \uparrow \\ c_1 \otimes \ldots \otimes c_m \xrightarrow{p} b_1 \otimes \ldots \otimes b_n$$

that is,

$$f^*p = (\gamma, \chi)^*(p) = \hat{\chi}^{-1} \circ p \circ \hat{\gamma} = \widehat{d_0(f)}^{-1} \circ p \circ \widehat{d_1(f)}$$

We then have

$$(\gamma', \chi')^* (\gamma, \chi)^* (p) = (\gamma', \chi')^* (\hat{\chi}^{-1} p \hat{\gamma}) = (\hat{\chi}')^{-1} \hat{\chi}^{-1} p \hat{\gamma} \hat{\gamma}' = \widehat{\chi \chi'}^{-1} p \widehat{\gamma \gamma'}$$

which is $(\gamma \gamma', \chi \chi')^*(p)$, as desired. Likewise,

$$\ell_i^*(\gamma, \chi)^*(p) = c_{\gamma(i)} = \ell_{\gamma(i)}^*(p)$$

$$r_j^*(\gamma, \chi)^*(p) = b_{\chi(j)} = r_{\chi(j)}^*(p)$$

so we conclude that $P|_{\mathbf{L}_{el}} \colon \mathbf{L}_{el}^{op} \to \mathbf{Set}$ is indeed a functor.

Notice that our definition of ℓ_i^* actually factors through a map $d_1: P_{\mathfrak{c}_{m,n}} \to P_{\underline{m}}$ taking p to $c_1 \otimes \cdots \otimes c_m$ which appeared in (3), and then following by the *i*th projection from (4). A similar situation occurs for r_j^* . We cannot do the same for $(\gamma, \chi)^*$.

Definition 4.5. We define $P|_{\mathbf{L}_{int}}$ to be the right Kan extension of $P|_{\mathbf{L}_{el}}$ along the fully faithful inclusion $\mathbf{L}_{el}^{op} \hookrightarrow \mathbf{L}_{int}^{op}$.

Since the functor is fully faithful, we can take the restriction of $P|_{\mathbf{L}_{int}}$ to be exactly equal to $P|_{\mathbf{L}_{el}}$. Further, by (4) we can arrange things so our previously-defined $P_{\underline{n}}$ agrees with the new one. By [CH21, Lemma 2.9], if we can show that $P|_{\mathbf{L}_{int}}$ is the restriction of some **L**-presheaf P, then P will automatically be Segal.

4.2. The active part. We have defined $P|_{\mathbf{L}_{int}}$ in Definition 4.5. In particular, this values of this presheaf are defined on every level graph G and on every isomorphism. Since $(\mathbf{L}_{act}, \mathbf{L}_{int})$ is an orthogonal factorization system on \mathbf{L} , it remains to define P on every active map. But each active map is isomorphic to $\alpha^*G \to G$ for $\alpha: [m] \to [n]$ an active map of $\boldsymbol{\Delta}$, so we will only extend to maps of this form. Further if G is a height n level graph, we can focus on the maps $d^i: d_i G \to G$ for 0 < i < n and $s^i: s_i G \to G$ for $0 \leq i \leq n$, since $\boldsymbol{\Delta}_{act}$ is generated by such maps.

In this section it will be helpful to have the following explicit version of splitting a graph G into its connected components. A looser version of this was used in Section 1.3 in proving Lemma 1.14.

Definition 4.6 (Canonical splitting of G). If G is a height m level graph with $G_{0m} = \underline{k}$, then for $x \in G_{0m}$ we define G_x to be the connected height m level graph

where for each i, j the following is a pullback whose top map ι_x is order-preserving.

$$(G_x)_{ij} \xrightarrow{\iota_x} G_{ij}$$
$$\downarrow \qquad \downarrow \qquad \downarrow$$
$$\underline{1} \xrightarrow{x} G_{0m}$$

Taking the sum of the ι_x , we have an isomorphism $\iota \colon \sum_{x=1}^k G_x \to G$ in \mathbf{L}_m .

Construction 4.7. For each height 1 level graph $G \in \mathbf{L}_1$, we define a function $\mu = \mu_G : P_G \to \text{mor } C$ so that the following diagram commutes.

$$\begin{array}{cccc} P_{d_1G} & \xleftarrow{d_1} & P_G & \xrightarrow{d_0} & P_{d_0G} \\ & & \downarrow^{\subseteq} & & \downarrow^{\mu_G} & & \downarrow^{\subseteq} \\ & & \text{ob} \ C & \xleftarrow{s} & \text{mor} \ C & \xrightarrow{t} & \text{ob} \ C. \end{array}$$

When $G = \mathfrak{c}$ is a corolla, μ is just the standard inclusion $P_{\mathfrak{c}} \subseteq \operatorname{mor}^{\mathfrak{c}} C \subseteq \operatorname{mor} C$ from the beginning of Section 4.1. More generally, suppose that G has n input-edges $(d_1G = G_{00} = \underline{n}), m$ output-edges $(d_0G = G_{11} = \underline{m}), \text{ and } k$ vertices $(G_{01} = \underline{k})$. If $p \in P_G$, then we must define $\mu_G(p)$ to be a map

$$d_1(p) = c_1 \otimes \cdots \otimes c_n \to e_1 \otimes \cdots \otimes e_m = d_0(p)$$

where c_i and e_i are connected objects. Recall the canonical splitting

$$\iota\colon \sum_{x=1}^k G_x \xrightarrow{\cong} G$$

from Definition 4.6. We write $\iota^0 = d_0(\iota) = \iota_{11}$ and $\iota^1 = d_1(\iota) = \iota_{00}$. If $p \in P_G$, write $p_x := \iota_x^*(p) \in P_{G_x} \subseteq \operatorname{mor}^c C$, which is a map

$$p_x \colon c_{i_1} \otimes \cdots \otimes c_{i_r} \to e_{j_1} \otimes \cdots \otimes e_{j_r}$$

where $\{i_1, \ldots, i_r\} = \iota_x((G_x)_{00})$ and $\{j_1, \ldots, j_{r'}\} = \iota_x((G_x)_{11})$. Tensoring these maps together, we have

$$p_1 \otimes \cdots \otimes p_k \colon c_{\iota^1(1)} \otimes \cdots \otimes c_{\iota^1(n)} \to e_{\iota^0(1)} \otimes \cdots \otimes e_{\iota^0(m)}.$$

Then $\mu_G(p)$ is defined as the following composite:

$$\begin{array}{ccc} c_{\iota^{1}(1)} \otimes \cdots \otimes c_{\iota^{1}(n)} & \xrightarrow{p_{1} \otimes \cdots \otimes p_{k}} & e_{\iota^{0}(1)} \otimes \cdots \otimes e_{\iota^{0}(m)} \\ & & \uparrow^{(\iota^{1})^{-1}} & & \downarrow^{\iota^{0}} \\ & c_{1} \otimes \cdots \otimes c_{n} & \xrightarrow{\mu_{G}(p)} & e_{1} \otimes \cdots \otimes e_{m} \end{array}$$

As a special case, we have the following.

Lemma 4.8. Let $G \in \mathbf{L}_1$ be a graph with both legs isomorphisms, as follows.

$$\frac{\underline{n}}{\alpha} \underbrace{\cong}_{\alpha} \underbrace{\cong}_{\beta} \underbrace{\stackrel{\underline{n}}{\swarrow}_{\beta}}_{\underline{n}}$$

Suppose c_1, \ldots, c_n are connected objects of C and $p \in P_G$ is the element which goes to $(\mathrm{id}_{c_1}, \mathrm{id}_{c_2}, \ldots, \mathrm{id}_{c_n})$ under the Segal map $P_G \to \prod_{i=1}^n P_{\mathfrak{c}_{11}}$. Then $\mu_G(p)$ is the following composite

$$c_{\alpha^{-1}(1)} \otimes \cdots \otimes c_{\alpha^{-1}(n)} \xrightarrow{\widehat{\alpha}} c_1 \otimes \cdots \otimes c_n \xrightarrow{\widehat{\beta}^{-1}} c_{\beta^{-1}(1)} \otimes \cdots \otimes c_{\beta^{-1}(n)}. \quad \Box$$

Lemma 4.9. Suppose that G and G' are in \mathbf{L}_1 and $i: G \to G + G'$, $j: G' \to G + G'$ are the inclusions. If $p \in P_{G+G'}$, then $\mu_{G+G'}(p) = \mu_G(i^*p) \otimes \mu_{G'}(j^*p)$.

Proof. The canonical decomposition for H = G + G'

$$\sum_{z=1}^{k+\ell} H_z = \sum_{x=1}^k G_x + \sum_{y=1}^\ell G'_y \xrightarrow{\iota = \alpha + \beta} G + G' = H$$

is the sum of the canonical decompositions (denoted by α and β) for G and G'. Write $\iota^1 \coloneqq d_1(\iota)$ and $\iota^0 \coloneqq d_0(\iota)$ and likewise for α and β . Then if $p_z = \iota_z^* p$, we have

$$\mu_{H}(p) = (i^{\widehat{1}})^{-1} \circ (p_{1} \otimes \dots \otimes p_{k} \otimes p_{k+1} \otimes \dots \otimes p_{k+\ell}) \circ i^{\widehat{0}}$$

$$= (\widehat{\alpha^{1}} \otimes \widehat{\beta^{1}})^{-1} \circ (p_{1} \otimes \dots \otimes p_{k} \otimes p_{k+1} \otimes \dots \otimes p_{k+\ell}) \circ (\widehat{\alpha^{0}} \otimes \widehat{\beta^{0}})$$

$$= ((\widehat{\alpha^{1}})^{-1} \otimes (\widehat{\beta^{1}})^{-1}) \circ (\alpha_{1}^{*}p \otimes \dots \otimes \alpha_{k}^{*}p \otimes \beta_{1}^{*}p \otimes \dots \otimes \beta_{\ell}^{*}p) \circ (\widehat{\alpha^{0}} \otimes \widehat{\beta^{0}})$$

$$= ((\widehat{\alpha^{1}})^{-1} \circ (\alpha_{1}^{*}p \otimes \dots \otimes \alpha_{k}^{*}p) \circ \widehat{\alpha^{0}}) \otimes ((\widehat{\beta^{1}})^{-1} \circ (\beta_{1}^{*}p \otimes \dots \otimes \beta_{\ell}^{*}p) \circ \widehat{\beta^{0}})$$

$$= \mu_{G}(i^{*}p) \otimes \mu_{G'}(j^{*}p).$$

Lemma 4.10. Suppose G and G' are isomorphic height 1 level graphs, and let μ_G and $\mu_{G'}$ be as in Construction 4.7. Let $f: G' \to G$ be an isomorphism. If $p \in P_G$, then

$$\mu_{G'}f^*(p) = \widehat{d_0(f)}^{-1} \circ \mu_G(p) \circ \widehat{d_1(f)}.$$

In particular, if f is a congruence then $\mu_{G'}f^*(p) = \mu_G(p)$, since $d_0(f)$ and $d_1(f)$ are identities.

Proof. If an isomorphism f satisfies the lemma, then so does f^{-1} . Further, if two composable isomorphisms f, f' satisfy the lemma, then so does their composition. We use these facts to reduce the proof to several special cases.

If G and G' are connected, that is, if they are both a corolla \mathfrak{c} , then μ is the canonical inclusion into mor^c C and the equation already appeared in Definition 4.4. We can boost this up to the case where we have isomorphisms of connected graphs $f^i: H^i \to K^i$ for $1 \leq i \leq m$, and we consider their sum

$$f\colon G'=\sum_{i=1}^m H^i\to \sum_{i=1}^m K^i=G.$$

(Note that $H^i = K^i$ since these are isomorphic corollas.) Letting $\iota'_i : H^i \to G'$ and $\iota_i : K^i \to G$ be the inclusions and using Lemma 4.9 and the result for corollas, we

have

$$\mu_{G'}(f^*(p)) = (\iota'_1)^*(f^*(p)) \otimes \cdots \otimes (\iota'_m)^*(f^*(p))$$

= $(f^1)^*(\iota_1^*(p)) \otimes \cdots \otimes (f^m)^*(\iota_m^*(p))$
= $[(\widehat{d_0f^1})^{-1} \circ \iota_1^*(p) \circ \widehat{d_1f^1}] \otimes \cdots \otimes [(\widehat{d_0f^m})^{-1} \circ \iota_m^*(p) \circ \widehat{d_1f^m}]$
= $\widehat{d_0f^{-1}} \circ (\iota_1^*(p) \otimes \cdots \otimes \iota_m^*(p)) \circ \widehat{d_1f}$
= $\widehat{d_0f^{-1}} \circ \mu_G(p) \circ \widehat{d_1f}.$

On the other hand, suppose we have

$$f: G' = \sum_{i=1}^{m} G'_i \to \sum_{k=1}^{m} G_i = G.$$

with f acting as an automorphism on \underline{m} but as the identity on each component, that is, $f \circ \iota'_i = \iota_{fi} \colon G_{fi} = G'_i \to G' \to G$. Then

$$\mu_{G'}f^*(p) = (\iota'_1)^*f^*(p) \otimes \cdots \otimes (\iota'_m)^*f^*(p)$$

= $\iota^*_{f1}(p) \otimes \cdots \otimes \iota^*_{fm}(p)$
= $\widehat{d_0f^{-1}} \circ (\iota^*_1(p) \otimes \cdots \otimes \iota^*_m(p)) \circ \widehat{d_1f}$

where these are now block permuations.


In the special case when f is the canonical decomposition from Definition 4.6

$$\iota\colon G'=\sum G_x\to G$$

the required formula follows from the definition of μ_G in Construction 4.7. That is, if $p_k = \iota_k^* p$ then we have

$$\mu_G(p) = \widehat{d_0\iota} \circ (p_1 \otimes \cdots \otimes p_k) \circ \widehat{d_1\iota}^{-1} = \widehat{d_0\iota} \circ (\mu_{G'}\iota^*(p)) \circ \widehat{d_1\iota}^{-1}.$$

We can combine all of these cases as follows




The vertical maps are canonical decompositions, the first map on the top rearranges factors, and the second map on top is a sum of isomorphisms between corollas. The result now follows for f by the facts in the first paragraph of the proof.

Proposition 4.11. Let $G \in \mathbf{L}_1$ and $\pi: C \to C(*)$. Then $\pi(\mu_G(p)) \in \mathbf{W}_1 = \text{mor } C(*)$ is the congruence class [G] of G.

Proof. In the connected case, that is, when $G = \mathfrak{c}_{n,m}$ is a corolla, this is true by definition of $P_{\mathfrak{c}_{n,m}}$. If G is empty, then $\mu_G(p) = \mathbf{1}$, so the result follows. If the result is true for G and G', then applying Lemma 4.9 we have

$$\pi\mu_{G+G'}(p) = \pi(\mu_G(i^*p) \otimes \mu_{G'}(j^*p)) = \pi\mu_G(i^*p) \otimes \pi\mu_{G'}(j^*p) = [G] \otimes [G'] = [G+G'].$$

Finally, suppose the result is true for G' and that $f: G' \to G$ is an isomorphism. Let $H \in \mathbf{L}_3$ denote following the height 3 level graph



where the unlabelled maps are the structure maps. This graph has the property that $d_1d_2H = G$ and $d_0d_3H = G'$, while d_2d_3G and d_0d_0G are of the form in Lemma 4.8. Let $p \in P_G$ be an element with $d_1(p) = c_1 \otimes \cdots \otimes c_n$ and $d_0(p) = e_1 \otimes \cdots \otimes e_m$. Using the Segal condition, define an element q of P_H so that

$$P_H \xrightarrow{d_2 d_3} P_{d_2 d_3 H} \xrightarrow{\cong} \prod_{G_{00}} P_{\mathfrak{c}_{11}}$$

sends q to $(\mathrm{id}_{c_1},\ldots,\mathrm{id}_{c_n}),$

$$P_H \xrightarrow{d_0 d_0} P_{d_0 d_0 H} \xrightarrow{\cong} \prod_{G_{11}} P_{\mathfrak{c}_{11}}$$

sends q to $(\mathrm{id}_{e_1},\ldots,\mathrm{id}_{e_m})$, and $d_0d_3: P_H \to P_{G'}$ sends q to $f^*(p)$. By Lemma 4.8,

$$\mu_{d_2d_3H}(d_2d_3q) = \widehat{d_1(f)}^{-1} \qquad \& \qquad \mu_{d_0d_0H}(d_0d_0q) = \widehat{d_0(f)}$$

and notice that π sends these to $[d_2d_3H]$ and $[d_0d_0H],$ respectively. By Lemma 4.10 we also have

$$\mu_G(p) = \widehat{d_0(f)} \circ \mu_{G'} f^*(p) \circ \widehat{d_1(f)}^{-1},$$


which is sent to

$$[d_0d_0H]\circ[d_0d_3H]\circ[d_2d_3H]$$


by π , since our assumption is $\pi(\mu_{G'}f^*(p)) = [G']$. But this element is equal to $[d_1d_2H] = [G]$. Thus $\pi(\mu_G(p)) = [G]$.

Since every graph is isomorphic to a finite sum of connected graphs, the general result follows. $\hfill \Box$

Definition 4.12 (Inner face in lowest dimension). Suppose G is a height 2 level graph. If G is connected, then d_1G is a corolla. In this case we get a map $P_G \to P_{d_1G}$ by using composition in C.



For an arbitrary height 2 level graph, define $d_1: P_G \to P_{d_1G}$ by examining the following square.



Here, each G_x is connected and the bottom map is an isomorphism. Then applying P we see d_1 is the dashed map in the following.

$$\begin{array}{ccc} P_G & \stackrel{\cong}{\longrightarrow} & \prod_x P_{G_x} \\ & & & & \downarrow \\ & & & \downarrow \\ P_{d_1G} & \stackrel{\cong}{\longrightarrow} & \prod_x P_{d_1G_x} \end{array}$$

Lemma 4.13. Suppose $f: G' \to G$ is an isomorphism between connected graphs in L_2 and $d_1(f) = g: d_1G' \to d_1G$. Then the diagram

$$P_{G} \xrightarrow{f^{*}} P_{G'}$$

$$\downarrow_{d_{1}} \qquad \downarrow_{d_{1}}$$

$$P_{d_{1}G} \xrightarrow{g^{*}} P_{d_{1}G}$$

commutes.

Proof. Write

$$\alpha \coloneqq d_2(f) \colon H' = d_2G' \to d_2G = H$$
$$\beta \coloneqq d_0(f) \colon K' = d_0G' \to d_0G = K$$

for the induced isomorphisms between height 1 level graphs. For i = 0, 1, we also write $\alpha_i \coloneqq d_i(\alpha)$ and $\beta_i \coloneqq d_i(\beta)$. If $p \in P_G$, then by definition

$$d_1(p) = \mu_K(d_0p) \circ \mu_H(d_2p)$$

and

$$d_1(f^*p) = \mu_{K'}(d_0f^*p) \circ \mu_{H'}(d_2f^*p) = \mu_{K'}(\beta^*d_0p) \circ \mu_{H'}(\alpha^*d_2p)$$

By Lemma 4.10 this means

$$d_1(f^*p) = \hat{\beta}_0^{-1} \circ \mu_K(d_0p) \circ \hat{\beta}_1 \circ \hat{\alpha}_0^{-1} \circ \mu_H(d_2p) \circ \hat{\alpha}_1.$$

Now $\alpha_0 = d_0 d_2(f) = d_1 d_0(f) = \beta_1$, so


$$d_1(f^*p) = \hat{\beta}_0^{-1} \circ \mu_K(d_0p) \circ \mu_H(d_2p) \circ \hat{\alpha}_1 = \hat{\beta}_0^{-1} \circ d_1(p) \circ \hat{\alpha}_1.$$

The result now follows from Definition 4.4 since $\alpha_1 = d_1 d_2(f) = d_1 d_1(f) = d_1(g)$ and $\beta_0 = d_0 d_0(f) = d_0 d_1(f) = d_0(g)$, implying $\hat{\beta}_0^{-1} \circ d_1(p) \circ \hat{\alpha}_1 = g^* d_1(p)$. \Box


Proposition 4.14. If $G \in \mathbf{L}_2$, then

(1)
$$d_0d_1: P_G \to P_{d_1G} \to P_{d_0d_1G}$$
 is equal to $d_0d_0: P_G \to P_{d_0G} \to P_{d_0d_0G}$.
(2) $d_1d_1: P_G \to P_{d_1G} \to P_{d_1d_1G}$ is equal to $d_1d_2: P_G \to P_{d_2G} \to P_{d_2d_1G}$.

Proof. We prove only the first equality, as the second is dual. Suppose G is connected.



The above diagram commutes, where the two upward maps are projections onto the second factor, so $d_0d_1 = d_0d_0$ in this case. The general case follows from the connected case:



Now that we have defined $d^1: P_G \to P_{d_1G}$ for each $G \in \mathbf{L}_2$ and shown it is compatible at the ends, we can use the Segal condition to define inner faces at arbitrary height graphs. It is convenient to use the following uniform notation for the generating outer face maps.

Convention 4.15. If X is a simplicial set, we write $d_{\perp} \colon X_n \to X_{n-1}$ for the bottom face map d_0 and $d_{\top} \colon X_n \to X_{n-1}$ for the top face map d_n . We use the same symbols when G is a height n level graph to write $d_{\perp} = d_0 \colon P_G \to P_{d_{\perp}G} = P_{d_0G}$ and $d_{\top} = d_n \colon P_G \to P_{d_{\top}G} = P_{d_nG}$.

If G is a height n level graph, then for 0 < i < n we can define d_i by means of the following diagram, which is well-formed by Proposition 4.14.

$$P_{G} \xrightarrow{=} P_{d_{\top}^{n-i+1}G} \underset{P_{d_{\bot}^{i-1}d_{\top}^{n-i+1}G}}{\times} P_{d_{\bot}^{i-1}d_{\top}^{n-i-1}G} \underset{P_{d_{\bot}^{i+1}d_{\top}^{n-i-1}G}}{\times} P_{d_{\bot}^{i+1}d_{\top}^{n-i-1}G} \overset{(5)}{\downarrow} \overset{\downarrow}{}_{id \times d_{1} \times id} \overset{\downarrow}{\downarrow} \overset{\downarrow}{}_{id \times d_{1} \times id} \overset{\downarrow}{P_{d_{\bot}^{i}G}} \underset{P_{d_{\bot}^{i-1}d_{\top}G}}{\times} P_{d_{\bot}^{i-1}d_{\top}^{n-i-1}d_{i}G} \underset{P_{d_{\bot}^{i}d_{\top}G}}{\times} P_{d_{\bot}^{i-1}d_{\top}^{n-i-1}d_{i}G} \overset{(5)}{\downarrow} \overset{\downarrow}{}_{id \times d_{1} \times id} \overset{\downarrow}{P_{d_{\bot}^{i-1}d_{\top}G}} \overset{\downarrow}{P_{d_{\bot}^{i-1}d_{\top}G}} \overset{\downarrow}{P_{d_{\bot}^{i-1}d_{\top}G}} \overset{\downarrow}{P_{d_{\bot}^{i-1}d_{\top}G}} \overset{\downarrow}{P_{d_{\bot}^{i-1}d_{\top}G}} \overset{\downarrow}{P_{d_{\bot}^{i-1}d_{\top}G}} \overset{\downarrow}{P_{J_{\bot}^{i-1}d_{\top}G}} \overset{\downarrow}{P_{J_{\bot}^{i-1}d_{\bot}G}} \overset{\downarrow}{P_{J_$$

In other words, inner face maps are defined as follows.

Definition 4.16 (Inner face operators). If $G \in \mathbf{L}_n$, 0 < i < n, and $x \in P_G$, then $d_i(x) \in P_{d_iG}$ is the unique element so that

• $d_{\perp}^{n-i}(d_i x) = d_{\perp}^{n-i+1}(x)$ • $d_{\perp}^{i-1} d_{\perp}^{n-i-1}(d_i x) = d_1(d_{\perp}^{i-1} d_{\perp}^{n-i-1} x)$, and • $d_{\perp}^i(d_i x) = d_{\perp}^{i+1}(x)$.

In order to utilize this definition effectively, we now examine how an inner face d_i interacts with arbitrary inert simplicial operators. The identity in the following lemma also holds in an arbitrary simplicial object.

Lemma 4.17. Suppose G is a height n level graph and $x \in P_G$. If $k + \ell \leq n - 1$, then

$$d_{\perp}^{k} d_{\top}^{\ell} d_{i}(x) = \begin{cases} d_{\perp}^{k} d_{\top}^{\ell+1}(x) & n-\ell \leq i \\ d_{i-k} d_{\perp}^{k} d_{\top}^{\ell}(x) & k+1 \leq i \leq n-\ell-1 \\ d_{\perp}^{k+1} d_{\top}^{\ell}(x) & i \leq k. \end{cases}$$

Proof. If $i \ge n - \ell$, then by Definition 4.16

$$d_{\top}^{\ell}d_{i}(x) = d_{\top}^{\ell-n+i}d_{\top}^{n-i}d_{i}(x) = d_{\top}^{\ell-n+i}d_{\top}^{n-i+1}(x) = d_{\top}^{\ell+1}(x),$$

while if $i \leq k$ then

$$d_{\perp}^{k}d_{i}(x) = d_{\perp}^{k-i}d_{\perp}^{i}d_{i}(x) = d_{\perp}^{k-i}d_{\perp}^{i+1}(x) = d_{\perp}^{k+1}(x),$$

so the first and last options hold. The interesting case is the second option. Notice that 0 < i - k < s, and we use the definition of the inner face map d_{i-k} from Definition 4.16. We make the following three computations

$$\begin{split} d_{\top}^{(n-k-\ell)-(i-k)}(d_{\perp}^{k}d_{\top}^{\ell}d_{i}x) &= d_{\perp}^{k}d_{\top}^{n-i}d_{i}(x) \\ &= d_{\perp}^{k}d_{\top}^{n-i+1}x = d_{\top}^{(n-k-\ell)-(i-k)+1}(d_{\perp}^{k}d_{\top}^{\ell}(x)) \\ d_{\perp}^{(i-k)-1}d_{\top}^{(n-k-\ell)-(i-k)-1}(d_{\perp}^{k}d_{\top}^{\ell}d_{i}x) &= d_{\perp}^{i-1}d_{\top}^{n-i-1}d_{i}(x) \\ &= d_{1}(d_{\perp}^{i-1}d_{\top}^{n-i-1}x) \\ &= d_{1}(d_{\perp}^{(i-k)-1}d_{\top}^{(n-k-\ell)-(i-k)-1}(d_{\perp}^{k}d_{\top}^{\ell}(x))) \end{split}$$

$$\begin{aligned} d_{\perp}^{i-k}(d_{\perp}^{k}d_{\top}^{\ell}d_{i}x) &= d_{\top}^{\ell}d_{\perp}^{i}d_{i}x \\ &= d_{\top}^{\ell}d_{\perp}^{i+1}x = d_{\perp}^{(i-k)+1}(d_{\perp}^{k}d_{\top}^{\ell}(x)) \end{aligned}$$

and conclude that the element $d_{\perp}^k d_{\perp}^\ell d_i x$ in $P_{d_{\perp}^k d_{\perp}^\ell d_i G} = P_{d_{i-k} d_{\perp}^k d_{\perp}^\ell d_{\perp}^\ell G}$ is equal to $d_{i-k} (d_{\perp}^k d_{\perp}^\ell x)$.

Proposition 4.18. Suppose $f: G' \to G$ is a map in \mathbf{L}_n , and 0 < i < n. Let $d_i(f) = g: d_iG' \to d_iG$. Then the diagram

$$\begin{array}{ccc} P_G & & \stackrel{f^*}{\longrightarrow} & P_{G'} \\ & & \downarrow_{d_i} & & \downarrow_{d_i} \\ P_{d_iG} & \stackrel{g^*}{\longrightarrow} & P_{d_iG'} \end{array}$$

commutes.


Proof. By Definition 4.16, since f and g are inert, it suffices to prove the result when n = 2 and i = 1. Further, the diagram commutes if and only if the outer rectangle in

$$P_{G} \xrightarrow{f^{*}} P_{G'} \xrightarrow{} P_{G'_{x}}$$

$$\downarrow^{d_{1}} \qquad \downarrow^{d_{1}} \qquad \downarrow^{d_{1}} \qquad \downarrow^{d_{1}}$$

$$P_{d_{1}G} \xrightarrow{g^{*}} P_{d_{1}G'} \xrightarrow{} P_{d_{1}G'_{x}}$$

commutes for every $x \in G'_{02} = (d_1G')_{01}$, since d_1 is defined as a product over the connected pieces. Since the cube



commutes, it is enough to show that

$$\begin{array}{ccc} P_G & \longrightarrow & P_{G_{fx}} & \longrightarrow & P_{G'_x} \\ & \downarrow^{d_1} & \downarrow^{d_1} & \downarrow^{d_1} \\ P_{d_1G} & \longrightarrow & P_{d_1G_{fx}} & \longrightarrow & P_{d_1G'_x} \end{array}$$

commutes. The left square commutes by definition of d_1 on disconnected graphs, and $G'_x \to G_{fx}$ is an isomorphism since any map in \mathbf{L}_n between connected graphs is an isomorphism. Thus it suffices to establish the result only in the case when $f: G' \to G$ is an isomorphism between height 2 connected graphs, which we already did in Lemma 4.13.

Lemma 4.19. If $G \in \mathbf{L}_2$, then

$$\begin{array}{ccc} P_{G} \xrightarrow{\cong} & P_{d_{\top} \times d_{\perp}} & P_{d_{\top}G} \times_{P_{d_{\top}d_{\perp}G}} & P_{d_{\perp}G} & \xrightarrow{\mu_{d_{\top}G} \times \mu_{d_{\perp}G}} & \operatorname{mor} C \times_{\operatorname{ob} C} & \operatorname{mor} C = NC_{2} \\ & \downarrow_{d_{1}} & & \downarrow_{\circ} \\ P_{d_{1}G} & \xrightarrow{\mu_{d_{1}G}} & & \operatorname{mor} C = NC_{1} \end{array}$$

commutes.

Proof. The result holds when G is connected; in fact, it is the definition of d_1 . In a moment we will induct on the number of connected components of G. First, suppose the result is true for G' and that $f: G' \to G$ is an isomorphism. Then

$$\mu_{d_1G'}(d_1f^*p) = \mu_{d_\perp G'}(d_\perp f^*p) \circ \mu_{d_\top G'}(d_\top f^*p).$$

By Proposition 4.18 on the left and the fact that P is a presheaf on \mathbf{L}_{int} , the preceding equation can be written as

$$\mu_{d_{\perp}G'}((d_{\perp}f)^*d_{1}p) = \mu_{d_{\perp}G'}((d_{\perp}f)^*d_{\perp}p) \circ \mu_{d_{\top}G'}((d_{\top}f)^*d_{\top}p).$$

We now apply Lemma 4.10 to see that

$$\mu_{d_1G'}((d_1f)^*d_1p) = \widehat{d_0d_1f}^{-1} \circ \mu_{d_1G}(d_1p) \circ \widehat{d_1d_1f}$$

and

$$\begin{split} & \mu_{d_{\perp}G'}((d_{\perp}f)^*d_{\perp}p) \circ \mu_{d_{\top}G'}((d_{\top}f)^*d_{\top}p) \\ = & \widehat{d_0d_{\perp}f^{-1}} \circ \mu_{d_{\perp}G}(d_{\perp}p) \circ \widehat{d_1d_{\perp}f} \circ \widehat{d_0d_{\top}f^{-1}} \circ \mu_{d_{\top}G}(d_{\top}p) \circ \widehat{d_1d_{\top}f} \\ = & \widehat{d_0d_{\perp}f^{-1}} \circ \mu_{d_{\perp}G}(d_{\perp}p) \circ \mu_{d_{\top}G}(d_{\top}p) \circ \widehat{d_1d_{\top}f}, \end{split}$$


using that $d_0 d_{\top} = d_0 d_2 = d_1 d_0 = d_1 d_{\perp}$. But also $d_0 d_{\top} = d_0 d_1$ and $d_1 d_{\top} = d_1 d_2 = d_1 d_1$, so we see that $\mu_{d_1 G}(d_1 p) = \mu_{d_{\perp} G}(d_{\perp} p) \circ \mu_{d_{\top} G}(d_{\top} p)$.

Now suppose the result is known for graphs with strictly fewer than n connected components, and let $H + K \to G$ be an isomorphism where H and K are non-empty graphs. By the preceding paragraph, it is enough to prove the result for the graph H + K. Let $h: H \to H + K$ and $k: K \to H + K$ be the inclusions. Write $h_i = d_i(h)$ and $k_i = d_i(k)$. Then using the induction hypothesis and Lemma 4.9, we have

$$\begin{split} & \mu_{d_1(H+K)}(d_1p) \\ &= \mu_{d_1H}(h_1^*d_1p) \otimes \mu_{d_1K}(k_1^*d_1p) \\ &= \mu_{d_1H}(d_1h^*p) \otimes \mu_{d_1K}(d_1k^*p) \\ &= [(\mu_{d_0H}(d_0h^*p)) \circ (\mu_{d_2H}(d_2h^*p))] \otimes [(\mu_{d_0K}(d_0k^*p)) \circ (\mu_{d_2K}(d_2k^*p))] \\ &= [(\mu_{d_0H}(d_0h^*p)) \otimes (\mu_{d_0K}(d_0k^*p))] \circ [(\mu_{d_2H}(d_2h^*p)) \otimes (\mu_{d_2K}(d_2k^*p))] \\ &= [(\mu_{d_0H}(h_0^*d_0p)) \otimes (\mu_{d_0K}(k_0^*d_0p))] \circ [(\mu_{d_2H}(h_2^*d_2p)) \otimes (\mu_{d_2K}(k_2^*d_2p))] \\ &= (\mu_{d_0(H+K)}(d_0p)) \circ (\mu_{d_2(H+K)}(d_2p)). \end{split}$$

Lemma 4.20. If G is a height 3 level graph and $x \in P_G$, then $d_1d_2(x) = d_1d_1(x)$.

Proof. For space reasons, in the following two diagrams we do not notate the fiber products, nor do we label the μ . We use the notation $NC_3 = \operatorname{mor} C \times_{\operatorname{ob} C} \operatorname{mor} C \times_{\operatorname{ob} C} \operatorname{mor} C$ so that $d_2 = (\operatorname{id} \times \circ)$ and $d_1 = (\circ \times \operatorname{id})$. In each diagram the two rectangles commute by Lemma 4.19.



If G is connected, then so are d_1d_2G and d_1d_1G , and the bottom map of both diagrams is an inclusion, so the result holds in the connected case. The general case follows from the connected case and Proposition 4.18.

Proposition 4.21. If G is a height n level graph, $x \in P_G$, and $0 \le i < j \le n$, then $d_i d_j(x) = d_{j-1} d_i(x)$.

Proof. If $i \leq j-2$, then using Lemma 4.17 one can compute that $d_{\perp}^m d_{\top}^{n-3-m} d_i d_j$ and $d_{\perp}^m d_{\top}^{n-3-m} d_{j-1} d_i$ are both equal to

$$\begin{cases} d_{\perp}^{m} d_{\top}^{n-1-m} & m \leq i-2 \\ d_{\perp} d_{\perp}^{m} d_{\top}^{n-2-m} & m = i-1 \\ d_{\perp}^{m+1} d_{\top}^{n-2-m} & i \leq m \leq j-3 \\ d_{1} d_{\perp}^{m+1} d_{\top}^{n-3-m} & m = j-2 \\ d_{\perp}^{m+2} d_{\top}^{n-3-m} & m \geq j-1. \end{cases}$$

By the Segal condition, this shows that $d_i d_j(x) = d_{j-1} d_i(x)$ in this case.

It remains to consider the case when i = j - 1. In this case we still have for $m \leq i - 2$ that

$$d^m_\perp d^{n-3-m}_\top d_i d_{i+1} = d^m_\perp d^{n-1-m}_\top = d^m_\perp d^{n-3-m}_\top d_i d_i$$

and for $m \geq i$ that

$$d_{\perp}^{m}d_{\top}^{n-3-m}d_{i}d_{i+1} = d_{\perp}^{m+2}d_{\top}^{n-3-m} = d_{\perp}^{m}d_{\top}^{n-3-m}d_{i}d_{i}$$

For the remaining value m = i - 1 = j - 2, we must verify that $d_{\perp}^{i-1} d_{\top}^{n-2-i} d_i d_{i+1} = d_{\perp}^{i-1} d_{\top}^{n-2-i} d_i d_i$. By Lemma 4.17 we have $d_{\perp}^{i-1} d_{\top}^{n-2-i} d_i d_{i+1} = d_1 d_2 d_{\perp}^{i-1} d_{\top}^{n-2-i}$ and $d_{\perp}^{i-1} d_{\top}^{n-2-i} d_i d_i = d_1 d_1 d_{\perp}^{i-1} d_{\top}^{n-2-i}$. These are equal by Lemma 4.20.

Definition 4.22 (Degeneracy in lowest dimension). If \underline{n} is height 0 level graph, define $s_0 \colon P_{\underline{n}} \to P_{s_0(\underline{n})}$ by declaring that

$$P_{\underline{n}} \to P_{s_0(\underline{n})} \xrightarrow{\cong} \prod_{i=1}^n P_{\mathfrak{c}_{11}}$$

takes $c_1 \otimes \ldots \otimes c_n$ to $(\mathrm{id}_{c_1}, \ldots, \mathrm{id}_{c_n})$.

Lemma 4.23. If $x \in P_{\underline{n}}$, then $d_0s_0(x) = x = d_1s_0(x)$.

Proof. For i = 0, 1 the full composite of the diagram

takes $c_1 \otimes \cdots \otimes c_n$ to (c_1, \ldots, c_n) . Hence $d_i s_0 = id$.

Lemma 4.24. If $f: \underline{n} \to \underline{m}$ is a map in \mathbf{L}_0 and $g = s_0(f)$, then

$$s_0 f^* = g^* s_0 \colon P_m \to P_{s_0(n)}$$

Proof. By Segality, it is enough to show that the the equality holds after postcomposing with $P_{s_0(\underline{n})} \to P_{\mathfrak{c}_{11}}$ for each of the *n* inert maps $\mathfrak{c}_{11} \to s_0(\underline{n})$. Since the two

squares on the right in the following diagram commute,

$$P_{\underline{m}} \xrightarrow{f^*} P_{\underline{n}} \longrightarrow \prod_{i=1}^n P_{\mathfrak{e}} \longrightarrow P_{\mathfrak{e}}$$

$$\downarrow^{s_0} \qquad \downarrow^{s_0} \qquad \downarrow \qquad \qquad \downarrow$$

$$P_{s_0(\underline{m})} \xrightarrow{g^*} P_{s_0(\underline{n})} \longrightarrow \prod_{i=1}^n P_{\mathfrak{c}_{11}} \longrightarrow P_{\mathfrak{c}_{11}}$$

it thus enough to prove the result for $\underline{n} = \underline{1}$. But if $f: \underline{1} \to \underline{m}$ hits an element k, then $s_0 f^*(c_1 \otimes \cdots \otimes c_m) = s_0(c_k) = \mathrm{id}_{c_k}$, and by Definition 4.22 we have $g^* s_0(c_1 \otimes \cdots \otimes c_m) = \mathrm{id}_{c_k}$.

Definition 4.25 (Degeneracy operators). If G is a height n level graph, then $s_j: P_G \to P_{s_jG}$ is defined so that $s_j(x)$ is the unique element satisfying

•
$$d_{\top}^{n+1-j}(s_j x) = d_{\top}^{n-j}(x)$$

• $d_{\perp}^j d_{\top}^{n-j}(s_j x) = s_0 (d_{\perp}^j d_{\top}^{n-j} x)$, and
• $d_{\perp}^{j+1}(s_j x) = d_{\perp}^j(x)$.

In other words, s_j is defined so the following diagram commutes.

$$P_{G} \xrightarrow{\cong} P_{d_{\tau}^{n-j}G} \underset{P_{d_{\perp}^{j}d_{\tau}^{n-j}G}}{\overset{\times}{\underset{f}{\longrightarrow}}} P_{d_{\perp}^{j}d_{\tau}^{n-j}G} \underset{P_{d_{\perp}^{j}d_{\tau}^{n-j}G}}{\overset{\times}{\underset{f}{\longrightarrow}}} P_{d_{\perp}^{j}d_{\tau}^{n-j}G} \underset{P_{d_{\perp}^{j}d_{\tau}^{n-j}G}}{\overset{\times}{\underset{f}{\longrightarrow}}} P_{d_{\perp}^{j}d_{\tau}^{n-j}S_{j}G} \underset{P_{d_{\perp}^{j}d_{\tau}^{n-j}S_{j}G}}{\overset{\times}{\underset{f}{\longrightarrow}}} P_{d_{\perp}^{j+1}d_{\tau}^{n-j}S_{j}G} \underset{P_{d_{\perp}^{j+1}d_{\tau}^{n-j}S_{j}G}}{\overset{\times}{\underset{f}{\longrightarrow}}} P_{d_{\perp}^{j+1}S_{j}G}$$

In order to utilize this definition effectively, we now examine how a degeneracy s_j interacts with arbitrary inert simplicial operators. The identity in the following lemma also holds in an arbitrary simplicial object.

Lemma 4.26. Suppose G is a height n level graph and $x \in P_G$. If $k + \ell \leq n + 1$, then

$$d_{\perp}^{k} d_{\top}^{\ell} s_{j}(x) = \begin{cases} d_{\perp}^{k} d_{\top}^{\ell-1}(x) & n-\ell+1 \leq j \\ s_{j-k} d_{\perp}^{k} d_{\top}^{\ell}(x) & k \leq j \leq n-\ell \\ d_{\perp}^{k-1} d_{\top}^{\ell}(x) & j \leq k-1. \end{cases}$$

Proof. If $j \ge n - \ell + 1$, then by Definition 4.25

$$d_{\top}^{\ell}s_{j}(x) = d_{\top}^{\ell-n+j-1}d_{\top}^{n+1-j}s_{j}(x) = d_{\top}^{\ell-n+j-1}d_{\top}^{n-j}(x) = d_{\top}^{\ell-1}(x),$$

while if $j \leq k - 1$ then

$$d^{k}_{\perp}s_{j}(x) = d^{k-j-1}_{\perp}d^{j+1}_{\perp}s_{j}(x) = d^{k-j-1}_{\perp}d^{j}_{\perp}(x) = d^{k-1}_{\perp}(x),$$

so the first and last options hold. We now use the definition of degeneracy operators from Definition 4.25. We make the following three computations

$$d_{\top}^{(n-k-\ell)+1-(j-k)}(d_{\perp}^{k}d_{\top}^{\ell}s_{j}x) = d_{\perp}^{k}d_{\top}^{n-j+1}s_{j}(x)$$

= $d_{\perp}^{k}d_{\top}^{n-j}x = d_{\top}^{(n-k-\ell)-(j-k)}(d_{\perp}^{k}d_{\top}^{\ell}(x))$

$$\begin{split} d_{\perp}^{j-k} d_{\top}^{(n-k-\ell)-(j-k)} (d_{\perp}^{k} d_{\top}^{\ell} s_{j} x) &= d_{\perp}^{j} d_{\top}^{n-j} s_{j}(x) \\ &= s_{0} (d_{\perp}^{j} d_{\top}^{n-j} x) \\ &= s_{0} (d_{\perp}^{j-k} d_{\top}^{(n-k-\ell)-(j-k)} (d_{\perp}^{k} d_{\top}^{\ell}(x)) \end{split}$$

$$\begin{aligned} d_{\perp}^{j-k+1}(d_{\perp}^k d_{\top}^\ell s_j x) &= d_{\top}^\ell d_{\perp}^{j+1} s_j x \\ &= d_{\top}^\ell d_{\perp}^j x = d_{\perp}^{j-k}(d_{\perp}^k d_{\top}^\ell (x)) \end{aligned}$$

and conclude that the element $d_{\perp}^k d_{\top}^\ell s_j x$ in $P_{d_{\perp}^k d_{\top}^\ell s_j G} = P_{s_{j-k} d_{\perp}^k d_{\top}^\ell G}$ is equal to $s_{j-k} (d_{\perp}^k d_{\top}^\ell x)$.

Proposition 4.27. Suppose $f: G' \to G$ is a map in \mathbf{L}_n , and $0 \leq j \leq n$. Let $s_j(f) = g: s_jG' \to s_jG$. Then the diagram

$$\begin{array}{c} P_{G} \xrightarrow{f^{*}} P_{G'} \\ \downarrow^{s_{j}} & \downarrow^{s_{j}} \\ P_{s_{j}G} \xrightarrow{g^{*}} P_{s_{j}G'} \end{array}$$

commutes.

Proof. By Segality it is enough to show that $d_{\perp}^m d_{\top}^{n-m} s_j f^*(x) = d_{\perp}^m d_{\top}^{n-m} g^* s_j(x)$ for all $x \in P_G$ and each $0 \le m \le n$. Set

$$h = d_{\perp}^{m} d_{\top}^{m-m}(g) = d_{\perp}^{m} d_{\top}^{m-m} s_{j}(f) = \begin{cases} d_{\perp}^{m} d_{\top}^{m-m-1}(f) & m+1 \leq j \\ s_{0} d_{\perp}^{m} d_{\top}^{n-m}(f) & j = m \\ d_{\top}^{m-1} d_{\top}^{n-m}(f) & j \leq m-1. \end{cases}$$

By Lemma 4.26, for $x \in P_G$ we have

$$d_{\perp}^{m} d_{\top}^{n-m} s_{j} f^{*} x = \begin{cases} d_{\perp}^{m} d_{\top}^{n-m-1} f^{*} x & m+1 \leq j \\ s_{0} d_{\perp}^{m} d_{\top}^{n-m} f^{*} x & j = m \\ d_{\top}^{m-1} d_{\top}^{n-m} f^{*} x & j \leq m-1. \end{cases}$$

Since $P|_{\mathbf{L}_{int}}$ is a presheaf we have the first equality below, while the second follows from Lemma 4.26.

$$d_{\perp}^{m} d_{\top}^{n-m} g^{*} s_{j} x = h^{*} d_{\perp}^{m} d_{\top}^{n-m} s_{j} x = \begin{cases} h^{*} d_{\perp}^{m} d_{\top}^{n-m-1} x & m+1 \leq j \\ h^{*} s_{0} d_{\perp}^{m} d_{\top}^{n-m} x & j = m \\ h^{*} d_{\top}^{m-1} d_{\top}^{n-m} x & j \leq m-1 \end{cases}$$

Again using that $P|_{\mathbf{L}_{int}}$ is a presheaf, we have the desired equality for $m \neq j$. When m = j, we have

$$s_0 d_{\perp}^m d_{\top}^{n-m} f^* x = s_0 (d_{\perp}^m d_{\top}^{n-m}(f))^* d_{\perp}^m d_{\top}^{n-m}(x).$$

But

$$s_0(d_{\perp}^m d_{\top}^{n-m}(f))^* = (s_0 d_{\perp}^m d_{\top}^{n-m}(f))^* s_0 = h^* s_0$$

by Lemma 4.24, so

$$d^j_\perp d^{n-j}_\top s_j f^*(x) = d^j_\perp d^{n-j}_\top g^* s_j(x)$$

holds as well.

Proposition 4.28. Suppose that G is a height n level graph and $x \in P_G$. If $i \leq j$, then $s_i s_j(x) = s_{j+1} s_i(x)$.

Proof. We have $s_i s_j G = s_{j+1} s_i G$ in \mathbf{L}_{n+2} . By Segality, it is enough to verify, for $0 \leq m \leq n+1$ and $x \in P_G$, that the operator $d_{\perp}^m d_{\top}^{n+1-m}$ takes the two elements $s_i s_j(x)$ and $s_{j+1} s_i(x)$ to the same element. Applying Lemma 4.26 several times, and also using Lemma 4.23, one computes that $d_{\perp}^m d_{\top}^{n+1-m} s_i s_j(x)$ and $d_{\perp}^m d_{\top}^{n+1-m} s_{j+1} s_i(x)$ are both equal to

$$\begin{cases} d_{\perp}^{m} d_{\top}^{n-m-1} x & m \leq i-1 \\ s_{0} d_{\perp}^{m} d_{\top}^{n-m} x & m = i \\ d_{\perp}^{m-1} d_{\top}^{m-m} x & i+1 \leq m \leq j \\ s_{0} d_{\perp}^{m-1} d_{\top}^{n+1-m} x & m = j+1 \\ d_{\perp}^{m-2} d_{\top}^{n+1-m} x & m \geq j+2. \end{cases}$$

Thus $s_i s_j(x) = s_{j+1} s_i(x)$.

Proposition 4.29. If G is a height 1 level graph and $x \in P_G$, then

$$d_1 s_0 x = x = d_1 s_1 x.$$

Proof. If we can prove the relation for a corolla \mathfrak{c} , then it will follow for an arbitrary $G \in \mathbf{L}_1$ using Proposition 4.18, Proposition 4.27, and Segality. Indeed, if $\iota : \mathfrak{c} \to G$ is a component of G, then for $x \in P_G$ and k = 0, 1, we have

$$\iota^* d_1 s_k x = d_1 (s_k \iota)^* s_k x = d_1 s_k \iota^* x = \iota^* x$$

using the corolla case, so by Segality $d_1 s_k x = x$.

Let $x \in P_{\mathfrak{c}}$ where \mathfrak{c} is a corolla. By Definition 4.12, d_1 applied to the element $s_0 x \in P_{s_0 \mathfrak{c}}$ is

$$\mu_{d_0s_0\mathfrak{c}}(d_0s_0x)\circ\mu_{d_2s_0\mathfrak{c}}(d_2s_0x)$$

which is equal to

$$\mu_{d_0s_0\mathfrak{c}}(x)\circ\mu_{d_2s_0\mathfrak{c}}(s_0d_1x)=x\circ\mathrm{id}=x$$

by Definition 4.25. Likewise, d_1s_1x is

$$\mu_{d_0s_1\mathfrak{c}}(d_0s_1x) \circ \mu_{d_2s_1\mathfrak{c}}(d_2s_1x) = \mu_{d_0s_1\mathfrak{c}}(s_0d_0x) \circ \mu_{d_2s_1\mathfrak{c}}(x) = \mathrm{id}\, \circ x = x. \qquad \Box$$

The proof of the following lemma uses the same idea as the proofs of Proposition 4.21 and Proposition 4.28.

Proposition 4.30. If G is a height n level graph and $x \in P_G$, then

$$d_i s_j x = \begin{cases} s_{j-1} d_i x & i < j \\ x & i = j, j+1 \\ s_j d_{i-1} x & i > j+1. \end{cases}$$

Proof. We will apply the operator $d_{\perp}^m d_{\top}^{n-m-1}$ for $0 \le m \le n-1$ to both sides of the equation, and show that our answers coincide. Since P is Segal, that implies that equation holds.

Using Lemma 4.17, Lemma 4.26, and Proposition 4.29, one computes that

$$d_{\perp}^{m} d_{\top}^{n-m-1} d_{i} s_{j} x = d_{\perp}^{m} d_{\top}^{n-m-1} x$$

whenever $m \leq \min(i-2, j-1), m \geq \max(i, j), m = j = i-1$, or m = i-1 = j-1. If i = j or i = j+1, this covers all possible values of m, so the formula holds in these cases. Otherwise, one computes using Lemma 4.17 and Lemma 4.26 that

$$d_{\perp}^{m} d_{\top}^{n-m-1} d_{i} s_{j} x = \begin{cases} d_{\perp}^{m+1} d_{\top}^{n-m-2} x & i \leq m \leq j-2 \\ d_{1} d_{\perp}^{m} d_{\top}^{n-m-2} x & i-1=m \leq j-2 \\ s_{0} d_{\perp}^{m+1} d_{\top}^{n-m-1} x & i \leq m=j-1 \\ d_{\perp}^{m-1} d_{\top}^{n-m} x & j+1 \leq m \leq i-2 \\ d_{1} d_{\perp}^{m-1} d_{\top}^{n-1-m} x & j+1 \leq m=i-1 \\ s_{0} d_{\perp}^{m} d_{\top}^{n-m} x & m=j \leq i-2. \end{cases}$$

When i < j, one computes

$$d_{\perp}^{m} d_{\top}^{n-m-1} s_{j-1} d_{i} x = \begin{cases} d_{\perp}^{m} d_{\top}^{n-m-1} x & m \leq i-2 \text{ or } j \leq m \\ d_{1} d_{\perp}^{m} d_{\top}^{n-m-2} x & m = i-1 \\ d_{\perp}^{m+1} d_{\top}^{n-m-2} x & i \leq m \leq j-2 \\ s_{0} d_{\perp}^{m+1} d_{\top}^{n-m-1} x & m = j-1 \end{cases}$$

which implies $d_i s_j x = s_{j-1} d_i x$ by the previous calculation. When i > j + 1, one computes

$$d_{\perp}^{m} d_{\top}^{n-m-1} s_{j} d_{i-1} x = \begin{cases} d_{\perp}^{m} d_{\top}^{n-m-1} x & m \leq j-1 \text{ or } i \leq m \\ s_{0} d_{\perp}^{m} d_{\top}^{n-m} x & m = j \\ d_{\perp}^{m-1} d_{\top}^{n-m} x & j+1 \leq m \leq i-2 \\ d_{1} d_{\perp}^{m-1} d_{\top}^{n-m-1} x & m = i-1, \end{cases}$$

implying $d_i s_j x = s_j d_{i-1} x$.

Theorem 4.31. The \mathbf{L}_{int} -presheaf $P|_{\mathbf{L}_{int}} \in \text{Seg}(\mathbf{L}_{int})$ extends to an \mathbf{L} -presheaf $P \in \text{Seg}(\mathbf{L})$ with inner face maps and degeneracies defined as in Definition 4.16 and Definition 4.25. Further, the assignment $C \rightsquigarrow P$ is a functor $\mathbf{SLCC} \rightarrow \text{Seg}(\mathbf{L})$.

Proof. Every map $G \to H$ in **L** factors uniquely as follows:

$$G \xrightarrow{f} \alpha^* H \xrightarrow{\alpha} H$$

where α is a map in Δ and $f \in \mathbf{L}_n$ where *n* is the height of *G*. We have an existing map $f^* \colon P_{\alpha^*H} \to P_G$ using the \mathbf{L}_{int} -presheaf structure, and we define $\alpha^* \colon P_H \to P_{\alpha^*H}$ by decomposing into face and degeneracy operators; by Proposition 4.21, Proposition 4.28, and Proposition 4.30 the result does not depend on any choices of how to do this decomposition. On such a composite, we define $(\alpha \circ f)^* \colon P_H \to P_G$ to be the composite $P_H \to P_{\alpha^*H} \to P_G$. To see that composition is preserved, consider a pair of composable morphisms in \mathbf{L} presented in this way:

$$G \xrightarrow{f} \alpha^* H \xrightarrow{\alpha} H \xrightarrow{g} \beta^* K \xrightarrow{\beta} K$$

$$\stackrel{h}{\longrightarrow} \alpha^* \beta^* K \xrightarrow{\alpha} \beta \alpha$$

Then setting $h \coloneqq \alpha^*(g) \colon \alpha^* H \to \alpha^* \beta^* K$, the composite is presented as in the dashed maps. Then

$$(\alpha \circ f)^* (\beta \circ g)^* = f^* \alpha^* g^* \beta^* = f^* h^* \alpha^* \beta^*$$

where the last equality uses Proposition 4.18, Proposition 4.27, and that P is an \mathbf{L}_{int} -presheaf. But then $f^*h^* = (hf)^*$ since P is an \mathbf{L}_{int} -presheaf and $\alpha^*\beta^* = (\beta\alpha)^*$ by Propositions 4.21, 4.28 and 4.30. Thus $(\alpha \circ f)^*(\beta \circ g)^* = ((\beta \circ g) \circ (\alpha \circ f))^*$, so P is an \mathbf{L} -presheaf.

All constructions from this section were functorial on maps in **SLCC**: the construction of the \mathbf{L}_{el} -presheaf, the right Kan extension $Psh(\mathbf{L}_{el}) \rightarrow Psh(\mathbf{L}_{int})$, and finally the definition of the inner face operators (Definitions 4.12 and 4.16) and degeneracy operators (Definitions 4.22 and 4.25). Thus we have constructed a functor **SLCC** \rightarrow Seg(**L**).

5. Comparison of constructions

We now have two constructions: the first is the functor $\text{Seg}(\mathbf{L}) \to \mathbf{SLCC}$ that takes a presheaf X to the image of the unique map $X \to *$ under the envelope functor C, and the second is the functor $\mathbf{SLCC} \to \text{Seg}(\mathbf{L})$ that takes a strict labelled cospan category $\pi: C \to C(*)$ to the presheaf $P = P(\pi)$. In this section, we will show that these are part of an equivalence of categories $\text{Seg}(\mathbf{L}) \simeq \mathbf{SLCC}$ (Theorem 5.8). We first show that the composite $\mathbf{SLCC} \to \text{Seg}(\mathbf{L}) \to \mathbf{SLCC}$ is isomorphic to the identity.

Let $\pi: C \to C(*)$ be a strict labelled cospan category, and $P = P(\pi)$ be the associated Segal **L**-presheaf. Our goal is to compare C with the envelope CP of P.

The set of connected objects of CP is $P_{\mathfrak{c}}$, which is exactly the set of connected objects of C. (See the beginning of Section 3 and the beginning of Section 4.1). Since these are both strict labelled cospan categories, the set of objects of each is the free monoid on this set, hence the sets of objects can be identified. Under this identification the tensor units coincide. Moreover, the set of connected morphisms of CP is $\sum P_{\mathfrak{c}_{n,m}}$, which is naturally in bijection with the set of connected morphisms of C (each $P_{\mathfrak{c}_{n,m}}$ is a subset of mor^c C, see (3) on page 31). The abelian monoid hom(1, 1) is the same in both categories, that is, it is freely generated by the identical sets hom^c_C(1, 1) = $P_{\mathfrak{c}_{00}} = \hom^{c}_{CP}(1, 1)$ of connected endomorphisms (see Definition 1.2(2)).

In Construction 4.7, we defined, for each $G \in \mathbf{L}_1$, a function $\mu_G \colon P_G \to \text{mor } C$. We now identify the image of this function.

Lemma 5.1. If $G \in \mathbf{L}_1$, then $\mu_G \colon P_G \to \operatorname{mor} C$ factors through \overline{P}_G and the map $\overline{P}_G \to \operatorname{mor} C$ is injective.

Proof. If $f: H \stackrel{\equiv}{\longrightarrow} G$ is a congruence between height 1 graphs, then by Lemma 4.10 we have $\mu_H f^*(p) = \mu_G(p)$ for all $p \in P_G$. This implies that μ_G factors through $\overline{P}_G = \operatorname{colim}_{H \in \mathbf{K}_{[G]}^{\operatorname{op}}} P_H$. For injectivity, we may assume that G has the property that all vertices without inputs or outputs come at the end; that is, we utilize a congruence

$$H = \left(G' + \sum_{i=1}^{k-j} \mathfrak{c}_{00}\right) \xrightarrow{\equiv} G$$

where G' has j vertices and is reduced, and $G' \to G$ is order-preserving at each level. Now $\overline{P}_G \to \text{mor } C$ is equal to $\overline{P}_H \to \text{mor } C$, so we replace G with H. Let $p, p' \in P_G$, where $G \in \mathbf{L}_1$ has j vertices which have an input or output

Let $p, p' \in P_G$, where $G \in \mathbf{L}_1$ has j vertices which have an input or output and k - j vertices which have no input or output, and the former come before the latter in the order on G_{01} . Assume $\mu_G(p) = \mu_G(p')$; we wish to show that p and p' represent the same element of \overline{P}_G . We use similar notation as in Construction 4.7; specifically, we let

$$\iota = \sum \iota_x \colon \sum_{x=1}^k G_x \xrightarrow{\cong} G$$

be the canonical splitting from Definition 4.6, $p_x = \iota_x^* p$, $p'_x = \iota_x^* p'$, $\iota^0 = d_0(\iota)$, and $\iota^1 = d_1(\iota)$. Then

$$\mu(p) = \widehat{\iota^0} \circ (p_1 \otimes \cdots \otimes p_k) \circ (\widehat{\iota^1})^{-1}$$
$$\mu(p') = \widehat{\iota^0} \circ (p'_1 \otimes \cdots \otimes p'_k) \circ (\widehat{\iota^1})^{-1}$$

Our assumption was $\mu(p) = \mu(p')$, hence

$$p_1 \otimes \cdots \otimes p_k = p'_1 \otimes \cdots \otimes p'_k$$

with all p_x , p'_x connected morphisms of C. By Definition 1.2(3) we have $p_1 \otimes \cdots \otimes p_j = p'_1 \otimes \cdots \otimes p'_j$ and $p_{j+1} \otimes \cdots \otimes p_k = p'_{j+1} \otimes \cdots \otimes p'_k$. By (4) of Definition 1.2, $p_x = p'_x$ for $0 \leq x \leq j$. In light of Definition 1.2(2), there is an automorphism γ' of the set $\{j+1, j+2, \ldots, k\}$ so that $p_x = p'_{\gamma'(x)}$ for $j+1 \leq x \leq k$. We extend γ' to an automorphism γ of \underline{k} by letting $\gamma(x) = x$ for $1 \leq x \leq j$, and this comprises a congruence $f: G \xrightarrow{\equiv} G$ with $f_{01} = \gamma$. We next observe that $f^*(p') = p$. By Segality, it is enough to show $\iota_x^* f^*(p') = \iota_x^*(p)$ for $1 \leq x \leq k$. Notice that $\iota_x^* f^*(p') = \iota_{\gamma(x)}^*(p')$. For $1 \leq x \leq j$, we have $\iota_{\gamma(x)}^*(p') = \iota_x^*(p') = p'_x = p_x = \iota_x^*(p)$. For $j+1 \leq x \leq k$, we have $\iota_{\gamma(x)}^*(p') = \iota_{\gamma'(x)}^*(p) = \iota_x^*(p)$.

We have shown that if $p, p' \in P_G$ are two elements with $\mu(p) = \mu(p')$, then they are identified in \overline{P}_G . Since $P_G \to \overline{P}_G$ is surjective, this proves that $\overline{P}_G \to \text{mor } C$ is injective.

As we saw at the beginning of the preceding proof, the functions $\mu_G \colon P_G \to \text{mor } C$ give well defined functions $\overline{P}_w \to \text{mor } C$ where $w \in \mathbf{W}_1 = \text{mor } C(*)$. Thus we can make the following definition.

Definition 5.2 (Comparison of morphisms). Let


$$\bar{\mu}$$
: mor $CP = SP_1 = \sum_{w \in \mathbf{W}_1} \overline{P}_w \to \operatorname{mor} C$

be the function induced from the collection of functions μ_G (for $G \in \mathbf{L}_1$).

Lemma 5.3. The function $\bar{\mu}$: mor $CP \rightarrow mor C$ is compatible with sources and targets of morphisms.

Proof. For each graph $G \in \mathbf{L}_1$, the outer square and all inner chambers except the inner rectangle of the following diagram are known to commute (including using

Construction 4.7).



It follows from surjectivity of $P_G \to \overline{P}_G$ that the inner rectangle commutes as well, so the function mor $CP \to \text{mor } C$ preserves sources of maps. Compatibility with targets is the evident modification of this argument.

The next proof uses that $\bar{\mu}$ preserves identities between connected objects, which is immediate from Definition 4.22, since $s_0: P_{\underline{1}} \to P_{s_0(\underline{1})} = P_{\mathfrak{c}_{11}}$ takes c to $\mathrm{id}_c \in P_{\mathfrak{c}_{11}} \subseteq \mathrm{mor} C$.

Lemma 5.4. The function $\bar{\mu}$: mor $CP \rightarrow mor C$ strictly preserves the monoidal product of morphisms, and also preserves the symmetry isomorphisms.

Proof. Recall from Lemma 2.21 and Definition 2.18 that the tensor product of morphisms in CP is induced as in the left rectangle in the following diagram, while the right square commutes by Lemma 4.9.

Thus $\bar{\mu}$ preserves monoidal product.

To see that $\bar{\mu}$ preserves the symmetry, it is enough to check that it is true for a pair of connected objects c, d. Let G be the following graph

$$\frac{2}{2} \frac{1}{2} \frac{1}{2} \frac{1}{2} \frac{2}{2} \frac{1}{2} \frac{1}$$

(using the notation of Section 2.4, this means $G = \mathfrak{c}_{11} + \mathfrak{c}_{11}$). Let $p \in P_G$ be the unique element which is sent to $(\mathrm{id}_c, \mathrm{id}_d)$ under the isomorphism $P_G \to P_{\mathfrak{c}_{11}} \times P_{\mathfrak{c}_{11}}$. By Definition 2.26 and Definition 2.31, the symmetry $\tau_{c,d} : c \otimes d \to d \otimes c$ of CPis the image of p under $P_G \to \overline{P}_G \subset \mathrm{mor}\, CP$. The definition of $\overline{\mu}$ implies that $\overline{\mu}(\tau_{c,d}) = \mu_G(p)$. By Lemma 4.8, $\mu_G(p) : c \otimes d \to d \otimes c$ is the symmetry isomorphism of C. Hence $\overline{\mu}$ preserves the symmetry isomorphism.

Lemma 5.5. The function $\bar{\mu}$: mor $CP \to \text{mor } C$ is bijective.

Proof. Suppose $G \neq G'$, $p \in P_G$, and $p' \in P_{G'}$. Then using Proposition 4.11 we have $\pi \mu_G(p) = [G] \neq [G'] = \pi \mu_{G'}(p')$ in mor $C(*) = \mathbf{W}_1$, hence $\mu_G(p) \neq \mu_{G'}(p')$ in mor C. Combining this with Lemma 5.1 we conclude that mor $CP \to \text{mor } C$ is injective. We now turn to surjectivity.

We already know that $\bar{\mu}$ restricts to a bijection of connected morphisms mor^c $CP = \sum P_{\mathfrak{c}_{nm}} \to \operatorname{mor}^{c} C$ and that $\operatorname{hom}_{CP}(\mathbf{1},\mathbf{1}) \to \operatorname{hom}_{C}(\mathbf{1},\mathbf{1})$ is an isomorphism. From Lemma 5.4 we know that $\bar{\mu}$ preserves the monoidal product, so by Definition 1.2(3), if mor^r $CP \to \operatorname{mor}^{r} C$ is a surjection then the same is true of mor $CP \to \operatorname{mor} C$. Let $f: c_1 \otimes \cdots \otimes c_n \to e_1 \otimes \cdots \otimes e_m$ be a reduced morphism of C other than the identity of the tensor unit, and let $G \in \mathbf{L}_1$ be a graph which represents $\pi(f) \in \operatorname{mor} C(*) = \mathbf{W}_1$. For specificity, we take $G_{01} = \underline{k}$ and let $\iota: G' = \sum_{x \in \underline{k}} G_x \to G$ be the canonical splitting from Definition 4.6. Define a new morphism of C by

$$f' = (\iota^0)^{-1} \circ f \circ \iota^1 \colon c_{\iota^1(1)} \otimes \cdots \otimes c_{\iota^1(n)} \to e_{\iota^0(1)} \otimes \cdots \otimes e_{\iota^0(m)}$$

where $\iota^1 = d_1(\iota): \underline{n} \to \underline{n}$ and $\iota^0 = d_0(\iota): \underline{m} \to \underline{m}$. For each $x \in \underline{k}$ we also have $\iota^1_x = d_1(\iota_x): \underline{n_x} \to \underline{n_x}$ and $\iota^0_x = d_0(\iota_x): \underline{m_x} \to \underline{m_x}$ where n_x is the number of inputs and m_x is the number of outputs of $\overline{G_x}$. By (4) of Definition 1.2 the following diagram is cartesian.

Thus there is an element (f'_1, \ldots, f'_k) in the upper left corner with f'_x living over $[G_x]$ and $f'_1 \otimes \cdots \otimes f'_k = f'$. The functor $\pi \colon C \to C(*)$ is strict monoidal, hence f' lives over $[\sum G_x] = [G']$. Since each G_x is a corolla, by definition of P_{G_x} we have $f'_x \in P_{G_x} \subseteq \text{mor}^c C$. These assemble to an element $p' \in P_{G'} = P_{\sum G_x}$ so that $\mu_{G'}(p') = f'$. Then the element $p = (\iota^{-1})^*(p')$ will be sent by μ_G to f, since Lemma 4.10 gives the last equality in the following

$$(\widehat{\iota^0})^{-1} \circ f \circ \widehat{\iota^1} = f' = \mu_{G'}(p') = \mu_{G'}\iota^*(p) = (\widehat{\iota^0})^{-1} \circ \mu_G(p) \circ \widehat{\iota^1}.$$

surjective.

Thus $\bar{\mu}$ is surjective.

Theorem 5.6. The function $\bar{\mu}$: mor $CP \to \text{mor } C$ gives an isomorphism of strict labelled cospan categories $CP \cong C$.

Proof. By Lemma 4.19, $\bar{\mu}$ preserves composition. Lemma 5.3 and Lemma 5.5 imply that $\hom_{CP}(x, y) \to \hom_C(x, y)$ is a bijection for all x, y. Preservation of identities follows from surjectivity of these functions and the fact that $\bar{\mu}$ preserves composition. Hence $CP \to C$ is a functor. Since it is bijective on objects and fully-faithful, it is an isomorphism of categories. By Lemma 5.4 this is a map of permutative categories, hence an isomorphism of such. By Proposition 4.11 this is a map over C(*), hence an isomorphism in **SLCC**.

Lemma 5.7. The functor $Seg(\mathbf{L}) \rightarrow SLCC$ is full.

Proof. Suppose we are given $X, Y \in \text{Seg}(\mathbf{L})$ and a map of strict labelled cospan categories $f: CX \to CY$. From the description near the beginning of §3, since the

permutative functor f lives over C(*), we have induced functions $f_{\underline{k}} \colon X_{\underline{k}} \to Y_{\underline{k}}$ and $f_{\mathfrak{c}_{m,n}} \colon X_{\mathfrak{c}_{m,n}} \to Y_{\mathfrak{c}_{m,n}}$; we will first show these give a map $X|_{\mathbf{L}_{\mathrm{el}}} \to Y|_{\mathbf{L}_{\mathrm{el}}}$ of \mathbf{L}_{el} -presheaves. Three types of maps are indicated in §4.1 that we must contend with, namely $\ell_i, r_j \colon \underline{1} \to \mathfrak{c}_{m,n}$ and $(\gamma, \chi) \colon \mathfrak{c}_{m,n} \to \mathfrak{c}_{m,n}$. The diagram for ℓ_i is listed just below, where the top composite is $\ell_i^* \colon X_{\mathfrak{c}_{m,n}} \to X_{\underline{1}}$ and similarly for the bottom composite. The two left squares commute since f preserves domains and the monoidal product.

We similarly have $f_{\underline{1}}r_j^* = r_j^* f_{\mathfrak{c}_{m,n}}$. For the bipermutations (γ, χ) acting on $\mathfrak{c}_{m,n}$ it is enough to consider the cases where one of γ or χ is an adjacent transposition and the other is an identity. Let H be the height 2 level graph

$$\underbrace{\overset{m}{\cong}}_{\underset{\underline{m}'}{\cong}} \underbrace{\overset{m}{\swarrow}}_{\underset{\underline{1}}{\swarrow}} \underbrace{\overset{n}{\searrow}}_{\underline{1}} \underbrace{\overset{n}{\swarrow}}_{\underline{1}}$$


where the leftmost bijection τ interchanges i and i + 1 and acts as the identity otherwise. There is an isomorphism $H \cong s_0 \mathfrak{c}_{m,n}$ which is $\tau \colon H_{00} \to (s_0 \mathfrak{c}_{m,n})_{00}$, and the identity elsewhere. Then the composite

$$\mathfrak{c}_{m,n} \xrightarrow{d^1} H \xrightarrow{\cong} s_0 \mathfrak{c}_{m,n} \xrightarrow{s^0} \mathfrak{c}_{m,n}$$


is the automorphism of $c_{m,n}$ which interchanges the *i*th and (i + 1)st inputs. There is a congruence as to the left of the following commutative square (see Section 2.4).

$$\begin{array}{ccc} d_2H & & \cong & & d_2s_0\mathfrak{c}_{m,n} \\ & & \downarrow \equiv & & \downarrow = \\ s_0(\underline{i-1}) + (\mathfrak{c}_{11}+\mathfrak{c}_{11}) + s_0(\underline{n-i-1}) & \xrightarrow{\mathrm{id}+\sigma_1+\mathrm{id}} & s_0\underline{m} \end{array}$$

so that $X_{s_0\underline{n}} \to X_{d_2H}$ sends the element representing $\mathrm{id}_{x_1\otimes\cdots\otimes x_m}$ to the element representing $\mathrm{id}_{x_1\otimes\cdots\otimes x_{i-1}}\otimes\tau_{x_i,x_{i+1}}\otimes\mathrm{id}_{x_{i+2}\otimes\cdots\otimes x_m}$ in CX. Commutativity of the following diagram then gives that a connected morphism $\phi: x_1\otimes\cdots\otimes x_m \to$ $x'_1\otimes\cdots\otimes x'_n$ is sent to $\phi\circ(\mathrm{id}_{x_1\otimes\cdots\otimes x_{i-1}}\otimes\tau_{x_i,x_{i+1}}\otimes\mathrm{id}_{x_{i+2}\otimes\cdots\otimes x_m})$.




All of this structure is compatible with $f: CX \to CY$, so we conclude that



commutes. The case when an adjacent transposition acts on an output is similar. It follows that f defines a map $X|_{\mathbf{L}_{el}} \to Y|_{\mathbf{L}_{el}}$.

Since Y is Segal, $Y|_{\mathbf{L}_{int}}$ is the right Kan extension of $Y|_{\mathbf{L}_{el}}$, hence we obtain a unique extension $X|_{\mathbf{L}_{int}} \to Y|_{\mathbf{L}_{int}}$. In particular, we have functions $f_G \colon X_G \to Y_G$ for every level graph G. It remains only to check that these maps are compatible with degeneracies and inner faces. By (5) above Definition 4.16, we only need to check compatibility with the inner faces $d_1 \colon X_G \to X_{d_1G}$ where $G \in \mathbf{L}_2$, and by (6) below Definition 4.25 we only need to check compatibility with the degeneracies $s_0 \colon X_{\underline{n}} \to X_{s_0(\underline{n})}$. Again by Segality we may further reduce to the cases where G is connected (for d_1) and n = 1 (for s_0). For d_1 , all squares of the following cube are known to commute with the exception of the back, which then must commute since $Y_{d_1G} \to SY_1$ is injective (using that d_1G is a corolla).



The map $s_0: X_{\mathfrak{e}} \to X_{\mathfrak{c}_{1,1}} \subset \operatorname{mor}^{c}(CX)$ takes a connected object x to id_x , so $f(s_0x) = f(\operatorname{id}_x) = \operatorname{id}_{fx} = s_0(fx)$, as desired. We conclude that $X|_{\mathbf{L}_{\operatorname{int}}} \to Y|_{\mathbf{L}_{\operatorname{int}}}$ is, in fact, a map $X \to Y$. By construction, this map is sent to f by C.

Theorem 5.8. The functor $Seg(L) \rightarrow SLCC$ is an equivalence of categories.

Proof. This functor is fully faithful by Proposition 3.10 and Lemma 5.7, and is essentially surjective by Theorem 5.6. \Box

Combining the previous theorem with Proposition 1.15, we have:

Corollary 5.9. The category of properads is equivalent to the category of strict labelled cospan categories. \Box

6. 2-CATEGORICAL STRUCTURES

In this section we turn our attention to the biequivalence between properads and labelled cospan categories, Corollary 6.16, which is the main result of this paper. The functor from strict labelled cospan categories into labelled cospan categories is not an equivalence of 1-categories, as it is only surjective up to equivalence, not up to isomorphism. Thus to connect properads to labelled cospan categories, we will need to view **Ppd** as a 2-category, rather than as a 1-category. The following definition is introduced in [Hac], where it is related to the Boardman–Vogt-style tensor product of properads.

Definition 6.1 (Natural transformation of properad maps). If $F, G: P \to Q$ are two maps of properads with the same source and target, then a *natural transformation* from F to G is nothing but a polynatural transformation between the underlying map of polycategories/dioperads, in the sense of [JY21, Definition 2.5.16]. This means that a natural transformation is given by a collection of unary morphisms $\gamma_c: F(c) \to$




FIGURE 4. Relation satisfied by a properadic natural transformation

G(c), indexed by the colors of P, so that, for each $p \in P(c_1, \ldots, c_n; d_1, \ldots, d_m)$, we have

(7)
$$(Gp)(\gamma_{c_1},\ldots,\gamma_{c_n}) = (\gamma_{d_1},\ldots,\gamma_{d_m})(Fp)$$

in $Q(Fc_1,\ldots,Fc_n;Gd_1,\ldots,Gd_m)$.

As in [JY21], the notation in (7) is meant to represent iterated dioperadic composition, see Figure 4.

Remark 6.2. In [HRY15, Section 4.4], a closed monoidal structure is given for those properads whose operations all have at least one input and at least one output, so in particular this subcategory becomes a 2-category. But this structure is not extended there to the category of all properads. Definition 6.1 is the extension of this 2-category to **Ppd**. (See [Dun06, Definition 6.3] for a different notion.)


6.1. Equivalences of labelled cospan categories. Let LCC denote the 2-category of labelled cospan categories, equipped with the 2-cells of Remark 1.4.

Proposition 6.3. Suppose $(f, \alpha): C \to D$ is a 1-morphism of labelled cospan categories, as depicted below.




The map (f, α) is an equivalence in the 2-category of labelled cospan categories if and only if $f: C \to D$ is an equivalence of categories.

Proof. Suppose $f: C \to D$ is an equivalence of categories. Then f is an equivalence in the 2-category of symmetric monoidal categories, symmetric monoidal functors, and monoidal natural transformations. Choose $g: D \to C$ along with monoidal natural isomorphisms $\eta: \operatorname{id}_C \cong gf$ and $\varepsilon: fg \cong \operatorname{id}_D$. Define a (monoidal) natural transformation $\beta \colon \pi g \Rightarrow \mu$ as the following composite.



Then $(g,\beta): D \to C$ is a 1-morphism of labelled cospan categories. Since the following pasting composites are isomorphisms,



by Remark 1.4 we have $(g,\beta) \circ (f,\alpha) \cong \mathrm{id}_{\pi}$ and $(f,\alpha) \circ (g,\beta) \cong \mathrm{id}_{\mu}$ in LCC. \Box

Note that if f is an *isomorphism* of categories, then (f, α) is an isomorphism of labelled cospan categories.

6.2. Strict labelled cospan categories form a 2-category. We now endow SLCC with the structure of a 2-category, which will serve as an intermediary between **Ppd** (considered as a 2-category) and **LCC**.

Definition 6.4. Given two maps



of strict labelled cospan categories (that is, commutative triangles with f and g strict maps of permutative categories), we define a 2-morphism from one to the other to be a monoidal natural transformation $\gamma: f \Rightarrow g$ so that the whiskering




is the identity natural transformation on $\pi: C \to C(*)$.

If the whiskering (8) is an isomorphism, then it is automatically an identity. This follows from Remark 1.3 and the fact that the canonical inclusion $C(*) \rightarrow \mathbf{Csp}$ is injective-on-objects and faithful.

There is a functor from the 1-category of strict labelled cospan categories to the 2-category LCC which is given by composition with $C(*) \rightarrow \mathbf{Csp}$.

Lemma 6.5. The usual inclusion $SLCC \rightarrow LCC$ extends to a 2-functor.

Proof. Suppose we are given a 2-morphism γ between 1-morphisms f and g in **SLCC** as in Definition 6.4. We then obtain the middle equality in the following chain of equalities of symmetric monoidal natural transformations between monoidal functors.



Thus γ is a 2-morphism between the 1-morphisms of labelled cospan categories (f, id) and (g, id) as in Remark 1.4.

We will return to the relationship between **SLCC** and **LCC** in Section 6.3, but first we will give more detail to facilitate the comparison with natural transformations of properads. Given a 2-morphism γ of **SLCC** as in Definition 6.4, the underlying data consists of *connected* morphisms

$$\gamma_x \colon f(x) \to g(x)$$

for each connected object x of C. We know these maps must all be connected, as $\pi'(\gamma_x) = (\mathrm{id}_{\pi})_x = \mathrm{id}_{\pi(x)} = \mathrm{id}_{\underline{1}}$ is a connected map in C(*). These maps are also the full extent of the data: if c is an arbitrary object of C, it can be uniquely written as $c = x_1 \otimes \cdots \otimes x_n$ where the x_i are connected objects. Since γ is a monoidal natural transformation and f and g are strict monoidal functors, the diagram

commutes, that is, $\gamma_c = \gamma_{x_1} \otimes \cdots \otimes \gamma_{x_n}$. We now observe that one only needs to check compatibility of γ with *connected morphisms*, rather than arbitrary morphisms.

Proposition 6.6. Suppose $f, g: C \to C'$ are two 1-morphisms of strict labelled cospan categories, and for each connected object x of C we have a connected morphism $\gamma_x: f(x) \to g(x)$. If the diagram

commutes for every connected morphism $p: x_1 \otimes \cdots \otimes x_n \to y_1 \otimes \cdots \otimes y_m$ of C, then the collection $\{\gamma_{x_1 \otimes \cdots \otimes x_n} \coloneqq \gamma_{x_1} \otimes \cdots \otimes \gamma_{x_n}\}$ constitutes a 2-morphism $f \Rightarrow g$.

Proof. Any endmorphism p of the tensor unit $\mathbf{1} \in C$ can be written as a composition of connected morphisms by Definition 1.2(2). Hence our assumption implies that $g(p)\gamma_{\mathbf{1}} = \gamma_{\mathbf{1}}f(p)$ for all such endomorphisms; since $\gamma_{\mathbf{1}}$ is the identity on the tensor unit of C', this implies that f(p) = g(p). Now by Definition 1.2(3), any morphism $p: c \to d$ can be written uniquely as $p' \otimes p'': c \otimes \mathbf{1} \to d \otimes \mathbf{1}$ where p' is a reduced morphism and p'' is an endomorphism of **1**, so it remains to show that $\gamma_d f(p') = g(p')\gamma_c$ for reduced morphisms p'.

Let σ be a bijection on $\underline{n} = \{1, \ldots, n\}$ and x_1, \ldots, x_n be connected objects in C. Naturality of symmetry implies that the diagram

$$\begin{array}{c} f(x_{\sigma(1)}) \otimes \cdots \otimes f(x_{\sigma(n)}) \xrightarrow{\gamma_{x_{\sigma(1)}} \otimes \cdots \otimes \gamma_{x_{\sigma(n)}}} g(x_{\sigma(1)}) \otimes \cdots \otimes g(x_{\sigma(n)}) \\ & \downarrow^{\hat{\sigma}} & \downarrow^{\hat{\sigma}} \\ f(x_1) \otimes \cdots \otimes f(x_n) \xrightarrow{\gamma_{x_1} \otimes \cdots \otimes \gamma_{x_n}} g(x_1) \otimes \cdots \otimes g(x_n) \end{array}$$

commutes in the permutative category C'. Since f and g are maps of permutative categories, the preceding diagram is equal to the following,

$$\begin{array}{ccc} f(x_{\sigma(1)} \otimes \cdots \otimes x_{\sigma(n)}) & \xrightarrow{\gamma_{x_{\sigma(1)}} \otimes \cdots \otimes x_{\sigma(n)}} g(x_{\sigma(1)} \otimes \cdots \otimes x_{\sigma(n)}) \\ & & \downarrow^{f(\hat{\sigma})} & & \downarrow^{g(\hat{\sigma})} \\ f(x_1 \otimes \cdots \otimes x_n) & \xrightarrow{\gamma_{x_1} \otimes \cdots \otimes x_n} g(x_1 \otimes \cdots \otimes x_n) \end{array}$$

where $\hat{\sigma}: x_{\sigma(1)} \otimes \cdots \otimes x_{\sigma(n)} \to x_1 \otimes \cdots \otimes x_n$ is the permutation map in C. Thus the proposed natural transformation is compatible with permutation maps.

Suppose $p: x_1 \otimes \cdots \otimes x_n \to y_1 \otimes \cdots \otimes y_m$ is an arbitrary reduced morphism of C lying over $\underline{n} \to \underline{k} \leftarrow \underline{m}$ in C(*). By choosing appropriate bijections σ of \underline{n} and ρ of \underline{m} , we can arrange things so that $\hat{\rho}^{-1} \circ p \circ \hat{\sigma}$ is sent by π to a k-fold tensor product of connected morphisms in C(*) (see Lemmas 5.1 and 5.5). Once we've done this, by Definition 1.2(4) we can find unique connected morphisms p_1, \ldots, p_k of C so that the diagram

$$\begin{array}{cccc} x_{\sigma(1)} \otimes \cdots \otimes x_{\sigma(n)} \xrightarrow{p_1 \otimes \cdots \otimes p_k} y_{\rho(1)} \otimes \cdots \otimes y_{\rho(m)} \\ & & & \downarrow^{\hat{\sigma}} & & \downarrow^{\hat{\rho}} \\ & & & x_1 \otimes \cdots \otimes x_n \xrightarrow{p} y_1 \otimes \cdots \otimes y_m \end{array}$$

commutes. By assumption each of the connected morphisms p_i is compatible with γ , hence so is $p_1 \otimes \cdots \otimes p_k$. Combining this with the previous paragraph, we see that

$$p = \hat{
ho} \circ (p_1 \otimes \cdots \otimes p_k) \circ \hat{\sigma}^{-1} = \hat{
ho} \circ (p_1 \otimes \cdots \otimes p_k) \circ \hat{\sigma}^{-1}$$

is also compatible with γ .

We have a composite of equivalences of 1-categories

$$\mathbf{Ppd} \xrightarrow{\simeq} \operatorname{Seg}(\mathbf{L}) \xrightarrow{\simeq} \mathbf{SLCC}$$

by Proposition 1.15 and Theorem 5.8. This extends to a strict 2-equivalence between 2-categories.

Theorem 6.7. The 1-functor $\mathbf{Ppd} \rightarrow \mathbf{SLCC}$ extends to a 2-functor. This 2-functor induces isomorphisms on hom-categories, hence is a strict 2-equivalence.

The last conclusion uses the previously established fact that the functor is surjective on objects up to isomorphism.

Proof. Suppose $F, G: P \to P'$ are two maps of properads, and $f, g: C \to C'$ are the associated maps of strict labelled cospan categories. The colors of P are precisely the connected objects of C, and the operations of P are precisely the connected

56

morphisms of C. Under this correspondence, the diagram from Proposition 6.6 is exactly the condition for a family of unary operations in P' to constitute a natural transformation $F \Rightarrow G$ of properad maps (Definition 6.1). It follows that

$$\mathbf{Ppd}(P, P')(F, G) \to \mathbf{SLCC}(C, C')(f, g)$$


is a bijection, that is, $\mathbf{Ppd} \to \mathbf{SLCC}$ is locally fully faithful. But we already know that $\mathbf{Ppd}(P, P') \to \mathbf{SLCC}(C, C')$ is bijective on objects, hence is an isomorphism of categories.

6.3. Comparison with labelled cospan categories. In this section we show that the inclusion of strict labelled cospan categories into all labelled cospan categories is a biequivalence of 2-categories.

Notation 6.8. If $\pi: C \to C(*)$ is a strict labelled cospan category, we write

$$\tilde{\pi}: C \to C(*) \to \mathbf{Csp}$$

for the composition of π with the canonical inclusion $C(*) \rightarrow \mathbf{Csp}$ from Examples 2.17 and 2.34. Given a 1-morphism of strict labelled cospan categories, that is a commutative triangle below left



where f is a strict permutative functor, we have the 1-morphism $\tilde{f} \coloneqq (f, \mathrm{id}_{\tilde{\pi}})$ of labelled cospan categories above right. Finally, if $\gamma \colon f \Rightarrow g$ is a 2-morphism as in Definition 6.4, we write $\tilde{\gamma} \colon (f, \mathrm{id}_{\tilde{\pi}}) \Rightarrow (g, \mathrm{id}_{\tilde{\pi}})$ for the same natural transformation, but now thought of as a 2-morphism in **LCC**.


We already observed in Lemma 6.5 that $SLCC \rightarrow LCC$ is a 2-functor under these assignments.

Lemma 6.9. The 2-functor $SLCC \rightarrow LCC$ is locally fully faithful.

 $\mathit{Proof.}$ Suppose $f,g\colon C\to C'$ are two 1-morphisms of labelled cospan categories. We wish to show that

$$\mathbf{SLCC}(C, C')(f, g) \to \mathbf{LCC}(C, C')(\tilde{f}, \tilde{g})$$

is a bijection. It is automatically injective since the elements on both sides are just certain natural transformations $f \Rightarrow g$. Suppose we have a 2-morphism $\tilde{f} \Rightarrow \tilde{g}$ as in Remark 1.4, that is a monoidal natural transformation $\gamma: f \Rightarrow g$ so that the composite natural transformation




is $\mathrm{id}_{\tilde{\pi}}$. This composite is just the whiskering of $\tilde{\pi}'$ with γ . To show that γ is a 2-morphism of **SLCC**, we need to show that its whiskering with π' is the identity on π

$$C \xrightarrow[g]{f} C' \xrightarrow{\pi'} C(*) \longrightarrow \mathbf{Csp}$$

which follows since $C(*) \to \mathbf{Csp}$ is faithful.

Lemma 6.10. The 2-functor SLCC \rightarrow LCC is locally an equivalence of categories.

Proof. Let $\pi: C \to C(*)$ and $\pi': C' \to C(*)$ be strict labelled cospan categories. From the previous lemma we know that $\mathbf{SLCC}(C, C') \to \mathbf{LCC}(C, C')$ is fully faithful, so it remains to prove that it is essentially surjective. Suppose (f, α) as depicted below left is 1-morphism in **LCC**.



Here f is a symmetric monoidal functor (which does not need to be a strict map of permutative categories) and α is a monoidal natural isomorphism. Since $C(*) \rightarrow \mathbf{Csp}$ is fully faithful, there is a (unique) monoidal natural isomorphism β as displayed above right, whose whiskering with $C(*) \rightarrow \mathbf{Csp}$ is α . Generically write $\Phi: f(c_1) \otimes \cdots \otimes f(c_n) \rightarrow f(c_1 \otimes \cdots \otimes c_n)$ for the structural isomorphisms of f(including the case n = 0 for the tensor unit).

We define a new functor of permutative categories $g: C \to C'$. On a general object $x_1 \otimes \cdots \otimes x_n$ of C (with each x_i connected), g is given by

$$g(x_1 \otimes \cdots \otimes x_n) \coloneqq f(x_1) \otimes \cdots \otimes f(x_n)$$

If $p: x_1 \otimes \cdots \otimes x_n \to y_1 \otimes \cdots \otimes y_m$ is any morphism of C, define g(p) to sit in the following commutative square.

$$\begin{array}{ccc} f(x_1) \otimes \cdots \otimes f(x_n) & \xrightarrow{g(p)} & f(y_1) \otimes \cdots \otimes f(y_m) \\ & & & & \\ & & & & \\ & & & & \\ & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & \\ & & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & & \\$$

This g is automatically a functor, and the commutativity of the diagram

implies that the unique dashed map must be both $g(p \otimes p')$ and $g(p) \otimes g(p')$, hence g is a strict monoidal functor. A similar diagram shows how to infer from f being symmetric monoidal functor that the same is true for g.

We will return to checking that g is a 1-morphism in **SLCC** in a moment. First, we define $\gamma: g \Rightarrow f$ to be the monoidal natural isomorphism given by Φ . That is, if $c = x_1 \otimes \cdots \otimes x_n$, then γ_c is the map

$$g(c) = f(x_1) \otimes \cdots \otimes f(x_n) \to f(x_1 \otimes \cdots \otimes x_n) = f(c).$$

We now calculate the following composite.

(9)
$$C \xrightarrow{\gamma \downarrow} C' \xrightarrow{\pi'} C(*)$$

Since π and π' are strict monoidal functors and β_{x_i} is the identity on $\underline{1}$ for a connected object x_i , commutativity of the following square

tells us that $\beta_{x_1 \otimes \cdots \otimes x_n}$ is the inverse of $\pi'(\Phi_{x_1,\ldots,x_n})$. Thus $\beta_c \circ \pi'(\gamma_c) = \mathrm{id}_{\pi c}$ for all objects $c \in C$, so it follows that the composite of (9) is the identity 2-morphism on the 1-morphism π . We then have $\pi'g = \pi$ and so g is a 1-morphism in **SLCC**, and γ is an isomorphism in **LCC**(C, C') between $(g, \mathrm{id}_{\tilde{\pi}})$ and (f, α) .

It remains to prove that $SLCC \rightarrow LCC$ is surjective up to equivalence. We begin with a special case.

Lemma 6.11. Suppose that C is a permutative category whose set of objects is a free monoid on the set S under the monoidal product, and that $\pi: C \to \mathbf{Csp}$ is a symmetric monoidal functor. If π is a labelled cospan category such that the set of connected objects is precisely $S \subset \operatorname{ob} C$, then there is a strict labelled cospan category $\pi': C \to C(*)$ such that the labelled cospan category $\tilde{\pi}'$ is isomorphic to π .

Proof. In this proof we write $i: C(*) \to \mathbf{Csp}$ for the canonical inclusion, and, whenever k + j = n, write

$$\Psi_{k,j} \colon i\underline{k} \otimes ij \to i(\underline{k}+j) = i\underline{n}$$

for the natural monoidal structure isomorphism of *i*. We similarly write $\Phi_{c,d}: \pi c \otimes \pi d \to \pi(c \otimes d)$ for the structure isomorphism of the symmetric monoidal functor π . We now aim to simultaneously define a more strict version of π called $\kappa: C \to \mathbf{Csp}$, along with a natural isomorphism $\beta: \pi \cong \kappa$. On objects, κ takes a word $c = [x_1x_2 \dots x_n]$ of length n (where each $x_i \in S$) to $i\underline{n}$. We now inductively, based on word length, define isomorphisms $\beta_c: \pi c \to \kappa c$. For length zero and length one words, we declare that $\beta_{[]}: \pi[] \to \kappa[] = \emptyset$ is the identity, and $\beta_{[x]}: \pi[x] \to \kappa[x] = i\underline{1}$ is the unique isomorphism. Suppose c has length k > 0 and d has length j > 0, and

that β_c and β_d have been defined. We then define $\beta_{c\otimes d}$ as the unique map fitting into the following square

$$\pi c \otimes \pi d \xrightarrow{\beta_c \otimes \beta_d} i\underline{k} \otimes i\underline{j}$$

$$\cong \downarrow \Phi_{c,d} \qquad \cong \downarrow \Psi_{\underline{k},\underline{j}}$$

$$\pi (c \otimes d) \xrightarrow{\beta_{c \otimes d}} i(k+j).$$

One must check that this is well-defined, that is, for any positive-length words $c, d, e \in$ ob C that $\beta_{(c\otimes d)\otimes e} = \beta_{c\otimes (d\otimes e)}$. This follows by using the hexagon constraints for Φ and Ψ and the fact that C is strict monoidal.

We define κ on morphisms $p: c \to d$ by declaring

$$\kappa(p) \coloneqq \beta_d \circ \pi(p) \circ \beta_c^{-1}$$

So defined, κ is automatically a functor and β is a natural transformation. Since *i* is fully faithful, there is a unique functor $\pi' \colon C \to C(*)$ so that $i\pi' = \kappa$. Using the defining equations, one can show that π' is a strict monoidal functor, which is also symmetric monoidal. From the diagram above one concludes that β is a monoidal natural transformation $\pi \Rightarrow i\pi'$.

The existence of the monoidal natural isomorphism β allows one to check that κ is a labelled cospan category. Further, π' is a strict labelled cospan category and (id_C, β) is an isomorphism $\tilde{\pi}' = \kappa \cong \pi$ of labelled cospan categories.

Lemma 6.12. The 2-functor $SLCC \rightarrow LCC$ is surjective up to equivalence.

Proof. Suppose $\pi: C \to \mathbf{Csp}$ is a labelled cospan category. A variation on the proof of [May74, Proposition 4.2] lets us define a permutative category D and a symmetric monoidal equivalence $f: D \to C$ so that $\operatorname{ob} D$ is the free monoid on the set $\operatorname{ob}^c C$ of connected objects of the labelled cospan category C. The functor f takes an object $[x_1x_2\cdots x_{n-1}x_n]$ (where each $x_i \in \operatorname{ob}^c C$) to $x_1 \otimes (x_2 \otimes (\cdots \otimes (x_{n-1} \otimes x_n) \cdots))$). One can check that $\pi f: D \to \mathbf{Csp}$ is a labelled cospan category. By Proposition 6.3, π and πf are equivalent objects in **LCC**, and by Lemma 6.11, πf is isomorphic to a strict labelled cospan category. Hence π is equivalent to a strict labelled cospan category.


We now have enough machinery to establish the following uniqueness result for (strict) labelled cospan categories with a given domain.

Proposition 6.13 (Barkan–Steinebrunner). If D is a permutative category, there is at most one strict labelled cospan category $D \to C(*)$. If C is a symmetric monoidal category, then any two labelled cospan categories $C \to \mathbf{Csp}$ are equivalent in LCC.

Proof. Suppose $\pi, \mu: C \to \mathbf{Csp}$ are two labelled cospan categories with the same source category; we will reduce the existence of an equivalence $\pi \simeq \mu$ to the first statement. Note that $\pi(c)$ has cardinality zero if and only if $\mu(c)$ does by Definition 1.2(1); then if a π -connected object c decomposes as $c \cong c_1 \otimes \cdots \otimes c_n$ for μ -connected objects c_i , we must have n = 1. It follows that the permuative category D and the functor f from the proof of Lemma 6.12 are the same for both π and μ . Applying Lemma 6.11 we have $\pi \simeq \pi f \cong \tilde{\pi}'$ and $\mu \simeq \mu f \cong \tilde{\mu}'$ for strict labelled cospan categories $\pi', \mu': D \to C(*)$. These last two maps will turn out to be equal, establishing the second claim.

Given two strict labelled cospan categories $\pi', \mu': D \to C(*)$ with the same domain permutative category D, we now show that $\pi' = \mu'$. Since free monoids have unique generating sets and the set of objects of D is the free monoid on the set of connected objects, the connected objects of π' and μ' are the same. Hence, on objects, the maps π' and μ' coincide; further, since π' and μ' are strict maps of permutative categories, they agree on the maps $\hat{\sigma}: x_{\sigma(1)} \otimes \cdots \otimes x_{\sigma(n)} \to x_1 \otimes \cdots \otimes x_n$ for each permutation σ of n. A free abelian monoid has a unique generating set, so the connected endomorphisms of **1** coincide for π' and μ' by Definition 1.2(2). Since π', μ' : hom $(1, 1) \to hom(0, 0)$ send the generators to the unique generator $\underline{0} \rightarrow \underline{1} \leftarrow \underline{0}$, these maps coincide on endomorphisms of the tensor unit. Suppose that $p: c \to d$ is a morphism of D which is connected with respect to π' . We can write $p = \hat{\sigma} \circ (p_1 \otimes \cdots \otimes p_k) \circ \hat{\rho}$ where each p_i is connected with respect to μ' . Then since no $\pi'(p_i)$ is the identity on 0, we must have k = 1, so p is also connected with respect to μ' . Since π' and μ' agree on objects and have the same connected morphisms, they agree on connected morphisms. But then $\pi'(p) = \mu'(p)$ for arbitrary morphisms p, since we can write any such p as $\hat{\sigma} \circ (p_1 \otimes \cdots \otimes p_k) \circ \hat{\rho}$ with the p_i connected, and π' and μ' are strict maps of permutative categories. Thus $\pi' = \mu'$. \square

Remark 6.14. Suppose $f: C \to C'$ is a symmetric monoidal equivalence and $\pi: C \to \mathbf{Csp}, \pi': C' \to \mathbf{Csp}$ are labelled cospan categories. Then f automatically determines a 1-morphism in **LCC**. Indeed, in this situation it is automatic that $\pi' f$ is a labelled cospan category, so unraveling the proof of Proposition 6.13 we obtain a natural isomorphism $\pi' f \cong \pi$ as the following pasting composite.



Here, the functor $D \to C(*)$ is the strict labelled cospan category constructed in the first paragraph of the proof.

Theorem 6.15. The 2-functor $SLCC \rightarrow LCC$ is a biequivalence.

Proof. This follows from Lemma 6.10 and Lemma 6.12.

We conclude by combining this theorem with the biequivalence of Theorem 6.7.

Corollary 6.16. The composite 2-functor $\mathbf{Ppd} \rightarrow \mathbf{LCC}$ is a biequivalence. \Box

References

- [BB17] M. A. Batanin and C. Berger, Homotopy theory for algebras over polynomial monads, Theory Appl. Categ. 32 (2017), Paper No. 6, 148–253. MR 3607212
- [BHS] Shaul Barkan, Rune Haugseng, and Jan Steinebrunner, Envelopes for algebraic patterns, preprint, arXiv:2208.07183 [math.CT].
- [BS] Shaul Barkan and Jan Steinebrunner, *The equifibered approach to* ∞ *-properads*, preprint, arXiv:2211.02576 [math.AT].
- [Car91] Aurelio Carboni, Matrices, relations, and group representations, J. Algebra 136 (1991), no. 2, 497–529. MR 1089310
- [CF17] Brandon Coya and Brendan Fong, Corelations are the prop for extraspecial commutative Frobenius monoids, Theory Appl. Categ. 32 (2017), Paper No. 11, 380–395. MR 3633708

- [CH21] Hongyi Chu and Rune Haugseng, Homotopy-coherent algebra via Segal conditions, Adv. Math. 385 (2021), Paper No. 107733, 95. MR 4256131
- [CH22] Hongyi Chu and Philip Hackney, On rectification and enrichment of infinity properads,
 J. Lond. Math. Soc. (2) 105 (2022), no. 1, 1418–1517. MR 4389306
- [Dun06] Ross Duncan, Types for quantum computing, Ph.D. thesis, Oxford University, 2006.
- [Gra66] John W. Gray, Fibred and cofibred categories, Proc. Conf. Categorical Algebra (La Jolla, Calif., 1965), Springer, New York, 1966, pp. 21–83. MR 0213413
- [Hac] Philip Hackney, *The 2-category of properads*, (working title), in preparation.
- [HK] Rune Haugseng and Joachim Kock, ∞ -operads as symmetric monoidal ∞ -categories, Forthcoming in Publ. Mat., arXiv:2106.12975 [math.CT].
- [HR15] Philip Hackney and Marcy Robertson, On the category of props, Appl. Categ. Structures 23 (2015), no. 4, 543–573. MR 3367131
- $[\mathrm{HR17}] \quad \underbrace{\qquad}_{835-902.} \text{ MR 3649609} \text{ , final props, Israel J. Math. 219 (2017), no. 2,}$
- [HRY15] Philip Hackney, Marcy Robertson, and Donald Yau, Infinity properads and infinity wheeled properads, Lecture Notes in Mathematics, vol. 2147, Springer, Cham, 2015. MR 3408444
- [HRY17] _____, A simplicial model for infinity properads, High. Struct. 1 (2017), no. 1, 1–21. MR 3912049
- [JY21] Niles Johnson and Donald Yau, 2-dimensional categories, Oxford University Press, Oxford, 2021. MR 4261588
- [KM] Ralph M. Kaufmann and Michael Monaco, Plus constructions, plethysm, and unique factorization categories with applications to graphs and operad-like theories, preprint, arXiv:2209.06121 [math.CT].
- [Lac04] Stephen Lack, Composing PROPS, Theory Appl. Categ. 13 (2004), No. 9, 147–163. MR 2116328
- [LR20] Fosco Loregian and Emily Riehl, Categorical notions of fibration, Expo. Math. 38 (2020), no. 4, 496–514. MR 4177953
- [Lur09] Jacob Lurie, Higher topos theory, Annals of Mathematics Studies, vol. 170, Princeton University Press, Princeton, NJ, 2009. MR 2522659
- [LV12] Jean-Louis Loday and Bruno Vallette, Algebraic operads, Grundlehren der mathematischen Wissenschaften, vol. 346, Springer, Heidelberg, 2012. MR 2954392
- [Mac65] Saunders Mac Lane, Categorical algebra, Bull. Amer. Math. Soc. 71 (1965), 40–106. MR 171826
- [na] nLab authors, Kan extension along (op)fibration, https://ncatlab.org/nlab/show/ Kan+extension#AlongFibrations.
- [RSW06] R. Rosebrugh, N. Sabadini, and R. F. C. Walters, Generic commutative separable algebras and cospans of graphs, Theory Appl. Categ. 15 (2005/06), No. 6, 164–177. MR 2210579
- $[RV20] Emily Riehl and Dominic Verity, On the construction of limits and colimits in \infty$ categories, Theory Appl. Categ.**35**(2020), Paper No. 30, 1101–1158. MR 4127725
- [RV22] _____, Elements of ∞-category theory, Cambridge Studies in Advanced Mathematics, vol. 194, Cambridge University Press, Cambridge, 2022. MR 4354541
- [Ste] Jan Steinebrunner, *The surface category and tropical curves*, preprint, arXiv:2111.14757v1 [math.AT].
- [Val07] Bruno Vallette, A Koszul duality for PROPs, Trans. Amer. Math. Soc. 359 (2007), no. 10, 4865–4943. MR 2320654 (2008e:18020)
- [YJ15] Donald Yau and Mark W. Johnson, A foundation for PROPs, algebras, and modules, Mathematical Surveys and Monographs, vol. 203, American Mathematical Society, Providence, RI, 2015. MR 3329226

 $\label{eq:linear} \begin{array}{l} \text{Department of Mathematics and Statistics, University of Nevada, Reno, USA} \\ Email \ address: \ \texttt{jbeardsley@unr.edu} \\ URL: \ \texttt{https://www.jonathanbeardsley.com/} \end{array}$

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF LOUISIANA AT LAFAYETTE, USA *Email address:* philip@phck.net *URL:* http://phck.net