

# DEFERRED NORLUND STATISTICAL CONVERGENCE IN PROBABILITY, MEAN AND DISTRIBUTION FOR SEQUENCES OF RANDOM VARIABLES

KULDIP RAJ, SWATI JASROTIA AND REZA SAADATI\*

ABSTRACT. We introduce and study deferred Nörlund statistical convergence in probability, mean of order  $r$ , distribution and study the interrelation among them. Based upon the proposed method to illustrate the findings, we presented new Korovkin type theorems for the sequence of random variables via deferred Nörlund statistically convergence and present compelling examples to demonstrate the effectiveness of the results.

## 1. Introduction and Preliminaries

Fast [5] and also Schoenberg [7] studied the concept of statistical convergence and continued by Rath–Tripathy [8] and Gadjiev–Orhan [20].

Suppose that  $(x_m)$  and  $(y_m)$  are the sequences of non-negative integers fulfilling

$$(1.1) \quad x_m < y_m, \forall m \in \mathbb{N} \quad \text{and} \quad \lim_{x \rightarrow \infty} y_m = \infty.$$

Further, let  $(e_m)$  and  $(g_m)$  be two sequences of non-negative real numbers such that

$$(1.2) \quad \mathcal{E}_m = \sum_{n=x_m+1}^{y_m} e_n \quad \text{and} \quad \mathcal{F}_m = \sum_{n=x_m+1}^{y_m} g_n.$$

The convolution of (1.2) is defined as

$$\mathcal{R}_m = \sum_{v=x_m+1}^{y_m} e_v g_{y_m-v}.$$

As introduced by Srivastava et al. in [24], the deferred Nörlund ( $DN$ ) mean is defined as

$$t_m = \frac{1}{\mathcal{R}_m} \sum_{n=x_m+1}^{y_m} e_{y_m-n} g_n y_n.$$

Suppose that  $(x_m)$  and  $(y_m)$  are the sequences fulfilling conditions (1.1) and  $(e_m), (g_m)$  are sequences satisfying (1.2). A sequence  $(Y_m)$  is called as deferred Nörlund statistically convergent to  $Y$  if  $\forall \varepsilon > 0$ , the set

$$\{n : n \leq \mathcal{R}_m, \text{ and } e_{y_m-n} g_n |Y_m - Y| \geq \varepsilon\}$$

---

2010 *Mathematics Subject Classification.* 40A05, 40A30.

*Key words and phrases.* Probability convergence, deferred Nörlund, mean convergence, distribution convergence, statistical convergence.

has zero deferred Nörlund density, i.e. if

$$\lim_{m \rightarrow \infty} \frac{1}{\mathcal{R}_m} \left| \left\{ n : n \leq \mathcal{R}_m \text{ and } e_{y_m-n} g_n |Y_m - Y| \geq \varepsilon \right\} \right| = 0.$$

We write it as

$$St_{DN} \lim Y_m = Y.$$

Suppose that  $(x_m)$  and  $(y_m)$  are the sequences fulfilling conditions (1.1) and  $(e_m), (g_m)$  are sequences satisfying (1.2). A sequence  $(Y_m)$  is called as deferred Nörlund statistically probability (or  $St_{DNP}$ -) convergent to a random variable  $Y$ , if  $\forall \varepsilon > 0$  and  $\delta > 0$ , the set

$$\{n : n \leq \mathcal{R}_m \text{ and } e_{y_m-n} g_n P(|Y_m - Y| \geq \varepsilon) \geq \delta\}$$

has  $DN$ -density zero, i.e.,

$$\lim_{m \rightarrow \infty} \frac{1}{\mathcal{R}_m} \left| \left\{ n : n \leq \mathcal{R}_m \text{ and } e_{y_m-n} g_n P(|Y_m - Y| \geq \varepsilon) \geq \delta \right\} \right| = 0$$

or

$$\lim_{m \rightarrow \infty} \frac{1}{\mathcal{R}_m} \left| \left\{ n : n \leq \mathcal{R}_m \text{ and } 1 - e_{y_m-n} g_n P(|Y_m - Y| \leq \varepsilon) \geq \delta \right\} \right| = 0,$$

and it is denoted as

$$St_{DNP} \lim_{m \rightarrow \infty} e_{y_m-n} g_n P(|Y_m - Y| \geq \varepsilon) = 0$$

or

$$St_{DNP} \lim_{m \rightarrow \infty} e_{y_m-n} g_n P(|Y_m - Y| \leq \varepsilon) = 1.$$

## 2. DEFERRED NÖRLUND STATISTICALLY PROBABILITY CONVERGENCE

In this section, we study deferred Nörlund statistically probability convergence, for a historical review and basic concept we refer [3], [4], [10], [12], [19] [9], [15], [23], [16], [17] and [27].

**Theorem 2.1.** *Suppose that  $(Y_m)$  and  $(Z_m)$  are sequences of random variables and consider two random variables  $Y$  and  $Z$ . Then the following assertions are satisfied*

- (1)  $St_{DNP} Y_m \rightarrow Y$  and  $St_{DNP} Y_m \rightarrow Z \Rightarrow P(Y = Z) = 1$ ,
- (2)  $St_{DNP} Y_m \rightarrow y \Rightarrow St_{DNP} Y_m^2 = y^2$ ,
- (3)  $St_{DNP} Y_m \rightarrow y$  and  $St_{DNP} Z_m \rightarrow z \Rightarrow St_{DNP} Y_m Z_m \rightarrow yz$ ,
- (4)  $St_{DNP} Y_m \rightarrow y$  and  $St_{DNP} Z_m \rightarrow z \Rightarrow St_{DNP} \frac{Y_m}{Z_m} \rightarrow \frac{y}{z}$ ,  $z \neq 0$ ,
- (5)  $St_{DNP} Y_m \rightarrow Y$  and  $St_{DNP} Z_m \rightarrow Z \Rightarrow St_{DNP} Y_m Z_m \rightarrow YZ$ ,
- (6) if  $St_{DNP} Y_m \rightarrow Y \forall \varepsilon, \delta > 0$ , then  $\exists a \in \mathbb{N}$  s.t.

$$d(\{n : n \leq \mathcal{R}_m \text{ and } e_{y_m-n} g_n P(|Y_m - Y_a| \geq \varepsilon) \geq \delta\}) = 0.$$

*Proof.* Let  $\epsilon$  and  $\delta$  be positively small real numbers. Also consider  $(x_m)$  and  $(y_m)$  are the sequences fulfilling conditions (1.1) and  $(e_m), (g_m)$  are sequences satisfying (1.2).

- (1) Suppose that  $a \in \left\{ n : n \leq \mathcal{R}_m \text{ and } e_{y_m-n} g_n P\left(|Y_m - Y| \geq \frac{\epsilon}{2}\right) < \frac{\delta}{2} \right\} \cap \left\{ n : n \leq \mathcal{R}_m \text{ and } e_{y_m-n} g_n P\left(|Y_m - Z| \geq \frac{\epsilon}{2}\right) < \frac{\delta}{2} \right\}$  (as the limit density of both the sets is 1). Then,  $e_{y_m-n} g_n P\left(|Y - Z| \geq \epsilon\right) \leq e_{y_m-n} g_n P\left(|Y_a - Y| \geq \frac{\epsilon}{2}\right) + e_{y_m-n} g_n P\left(|Y_a - Z| \geq \frac{\epsilon}{2}\right) < \delta$ . It means

$$P\{Y = Z\} = 1.$$

- (2) If  $St_{DNP} Y_m \rightarrow 0$ , then  $St_{DNP} Y_m^2 \rightarrow 0$ . Here, we see that  $a \in \{n : n \leq \mathcal{R}_m \text{ and } e_{y_m-n} g_n P(|Y_m - 0| \geq \epsilon) > \delta\} = a \in \{n : n \leq \mathcal{R}_m \text{ and } e_{y_m-n} g_n P(|Y_m^2 - 0| \geq \epsilon > \delta)\}$ . Now, take  $Y_m^2 = (Y_m - y)^2 + 2y(Y_m - y) + y^2$ . Thus,  $St_{DNP} Y_m^2 \rightarrow y^2$ .
- (3) Suppose that  $St_{DNP} Y_m \rightarrow y$  and  $St_{DNP} Z_m \rightarrow z$ . As  $St_{DNP} Y_m Z_m = St_{DNP} \frac{1}{4} \{(Y_m + Z_m)^2 - (Y_m - Z_m)^2\} = \frac{1}{4} \{(y_m + z_m)^2 - (y_m - z_m)^2\} = yz$ .
- (4) Suppose that  $R$  and  $S$  be two events correspond  $|Z_m - z| < |z|$  and  $\left|\frac{1}{Z_m} - \frac{1}{z}\right| \geq \epsilon$ . We have

$$\left|\frac{1}{Z_m} - \frac{1}{z}\right| = \frac{|Z_m - z|}{|z Z_m|} = \frac{|Z_m - z|}{|z| \cdot |z + (Z_m - z)|} \leq \frac{|Z_m - z|}{|z| \cdot (|z| - |Z_m - z|)}.$$

If the events  $R$  and  $S$  occurs at same time, then

$$|Z_m - z| \geq \frac{\epsilon |z|^2}{1 + \epsilon |z|}.$$

Further, let  $\epsilon_0 = \epsilon |z|^2 / (1 + \epsilon |z|)$  and  $A$  be the event such that  $|Z_m - z| \geq \epsilon_0$ . Thus,

$$RS \subseteq A \Rightarrow P(S) \leq P(A) + P(R^c).$$

Thus,

$$\left\{ n : n \leq \mathcal{R}_m \text{ and } e_{y_m-n} g_n P\left(\left|\frac{1}{Z_m} - \frac{1}{z}\right| \geq \epsilon\right) \geq \delta \right\} \subseteq \left\{ n : n \leq \mathcal{R}_m \text{ and } \right.$$

$$\left. e_{y_m-n} g_n P(|Z_m - z| \geq \epsilon_0) \geq \frac{\delta}{2} \right\} \cup \left\{ n : n \leq \mathcal{R}_m \text{ and } e_{y_m-n} g_n P(|Z_m - z| \geq |z|) \geq \frac{\delta}{2} \right\}.$$

Therefore,  $St_{DNP} \frac{1}{Z_m} \rightarrow \frac{1}{z}$ . Hence, we write  $St_{DNP} \frac{Y_m}{Z_m} \rightarrow \frac{y}{z}$ ,  $z \neq 0$ .

- (5) Suppose that  $St_{DNP} Y_m \rightarrow Y$  and  $X$  be a random variable such that  $Y_m X \rightarrow YX$ . Since  $X$  is a random variable such that  $\forall \epsilon > 0, \exists \delta > 0$  and  $e_{y_m-n} g_n P(|X| > \delta) \leq \frac{\epsilon}{2}$ . Next,  $\forall \epsilon' > 0$ ,

$$\begin{aligned} e_{y_m-n} g_n P(|Y_m X - YX| \geq \epsilon') &= e_{y_m-n} g_n P(|Y_m - Y||X| \geq \epsilon', |Z| > \delta) \\ &\quad + e_{y_m-n} g_n P(|Y_m - Y||X| \geq \epsilon', |Z| \leq \delta) \leq \frac{\epsilon}{2} \\ &\quad + e_{y_m-n} g_n P(|Y_m - Y| \geq \frac{\epsilon'}{\delta}). \end{aligned}$$

Which implies,  $\{n : n \leq \mathcal{R}_m \text{ and } e_{y_m-n}g_nP(|Y_mX - YX| \geq \varepsilon')\} \subseteq \{n : n \leq \mathcal{R}_m \text{ and } e_{y_m-n}g_nP(|Y_m - y| \geq \frac{\varepsilon'}{\delta}) \geq \frac{\varepsilon}{2}\}$ . Therefore,

$$St_{DNP}(Y_m - Y)(Z_m - Z) \rightarrow 0.$$

Thus,

$$St_{DNP}Y_mZ_m \rightarrow YZ.$$

(6) Suppose that  $(x_m)$  and  $(y_m)$  be two non-negative sequences such that

$$e_{y_m-n}g_nP\left(|Y_m - Y| \geq \frac{\varepsilon}{2}\right) < \frac{\delta}{2}$$

and

$$\left\{n : n \leq \mathcal{R}_m \text{ and } e_{y_m-n}g_nP\left(|Y_m - Y| \geq \frac{\varepsilon}{2}\right) < \frac{\delta}{2}\right\} = 1.$$

Now,

$\{n : n \leq \mathcal{R}_m \text{ and } e_{y_m-n}g_nP(|Y_m - Y| \geq \varepsilon) \geq \delta\} \subseteq \{n : n \leq \mathcal{R}_m \text{ and } e_{y_m-n}g_nP(|Y_m - Y| \geq \frac{\varepsilon}{2}) < \frac{\delta}{2}\} = 1$ . Which implies that

$$d(\{n : n \leq \mathcal{R}_m \text{ and } e_{y_m-n}g_nP(|Y_m - Y| \geq \varepsilon) \geq \delta\}) = 0.$$

□

**Theorem 2.2.** Suppose that  $f : \mathbb{R} \rightarrow \mathbb{R}$  is uniform continuous on  $\mathbb{R}$  and  $St_{DNP}Y_m \rightarrow Y$ . Then  $St_{DNP}f(Y_m) \rightarrow f(Y)$ .

*Proof.* Let us consider a random variable  $Y$  such that for each  $\delta > 0$ ,  $\exists \beta \in \mathbb{R}$  such that  $P(Y > \beta) \leq \delta/2$ . Since,  $f$  is uniformly continuous on  $[\beta, \beta] \forall \varepsilon > 0$ ,  $\exists \delta_0$  such that

$$|f(y_m) - f(y)| < \varepsilon \text{ whenever } |y_m - y| < \delta_0.$$

Thus,

$$\begin{aligned} P(|f(Y_m) - f(Y)| \geq \varepsilon) &\leq P(|Y_m - Y| \geq \delta_0) + P(|Y| > \beta) \\ &\leq P(|Y_m - Y| \geq \delta_0) + \delta/2. \end{aligned}$$

However, from the definition of  $St_{DNP}$ -convergence, we have

$$\begin{aligned} &\left\{n : n \leq \mathcal{R}_m \text{ and } e_{y_m-n}g_nP(|f(Y_m) - f(Y)| \geq \varepsilon) \geq \delta\right\} \\ &\subseteq \left\{n : n \leq \mathcal{R}_m \text{ and } e_{y_m-n}g_nP(|Y_m - Y| \geq \delta_0) < \frac{\delta}{2}\right\}. \end{aligned}$$

□

### 3. DEFERRED NÖRLUND STATISTICAL MEAN CONVERGENCE

**Definition 3.1.** Suppose that  $r \geq 1$  be a fixed number. A sequence  $(Y_m)$  is  $r^{th}$  mean convergent to  $Y$ , if

$$\lim_{m \rightarrow \infty} E(|Y_m - Y|^r) = 0.$$

**Definition 3.2.** A sequence  $(Y_m)$  is statistically  $r^{th}$  mean convergent ( $MC$ ) to a random variable  $Y$ , where  $Y : S \rightarrow \mathbb{R}$  if,

$$\lim_{m \rightarrow \infty} \frac{1}{m} \left| n : n \leq m \text{ and } E(|Y_m - Y|^r \geq \varepsilon) \right| = 0$$

for any  $\varepsilon > 0$ .

We write it as

$$St_{MC} \lim_{m \rightarrow \infty} E(|Y_m - Y|^r) = 0.$$

**Definition 3.3.** Suppose that  $(x_m)$  and  $(y_m)$  are the sequences fulfilling conditions (1.1) and  $(e_m), (g_m)$  are sequences satisfying (1.2). A sequence  $(Y_m)$  is said to be deferred Nörlund statistically  $r^{\text{th}}$  ( $r \geq 1$ ) mean convergent to  $Y$  ( $Y : S \rightarrow \mathbb{R}$ ), if for  $\varepsilon > 0$ ,

$$\lim_{m \rightarrow \infty} \frac{1}{\mathcal{R}_m} \left| \left\{ n : n \leq \mathcal{R}_m \text{ and } e_{y_m-n} g_n E(|Y_m - Y|^r \geq \varepsilon) \right\} \right| = 0.$$

It is denoted as

$$St_{DNM} \lim_{m \rightarrow \infty} E(|Y_m - Y|^r) = 0.$$

**Theorem 3.4.** Let  $St_{DNM} \lim_{m \rightarrow \infty} E(|Y_m - Y|^r) = 0$  for  $r \geq 1$ , then  $St_{DNM} \lim_{m \rightarrow \infty} P(|Y_m - Y| \geq \varepsilon) = 0$ .

*Proof.* For every  $\varepsilon > 0$ , we have from Markov's inequality

$$\begin{aligned} St_{DNM} \lim_{m \rightarrow \infty} P(|Y_m - Y| \geq \varepsilon) &= St_{DNM} \lim_{m \rightarrow \infty} P(|Y_m - Y|^r \geq \varepsilon^r) \quad (r \geq 1) \\ &\leq St_{DNM} \lim_{m \rightarrow \infty} \frac{E(|Y_m - Y|^r)}{\varepsilon^r} = 0. \end{aligned}$$

From definition of statistically deferred Nörlund mean convergence

$$St_{DNM} \lim_{m \rightarrow \infty} E(|Y_m - Y|^r) = 0,$$

it implies that

$$St_{DNP} \lim_{m \rightarrow \infty} P(|Y_m - Y| \geq \varepsilon) = 0.$$

□

We now present an example to show that a sequence of random variables is statistically probability convergent but not statistically  $r^{\text{th}}$ -mean convergent.

**Example 1:** Suppose that  $x_m = 2m - 1, y_m = 4m - 1$ . Also, suppose that  $e_{y_m-m} = 2m$  and  $g_m = 1$ . Further, consider a sequence  $(z_m)$  of random variables such that

$$Y_m = \begin{cases} m, & \text{with probability } \frac{1}{\sqrt{m}} \\ 0, & \text{with probability } 1 - \frac{1}{\sqrt{m}}. \end{cases}$$

Then the statistically deferred Nörlund convergence of  $Y_m$  is given as

$$\begin{aligned} \lim_{m \rightarrow \infty} \frac{1}{2m} \left| \left\{ n : n \leq \mathcal{R}_m \text{ and } 2mP(|Y_m - 0| \geq \varepsilon) \right\} \right| &= \lim_{m \rightarrow \infty} P(Y_m = m) \\ &= \lim_{m \rightarrow \infty} \frac{1}{\sqrt{m}} \\ &= 0. \end{aligned}$$

However, statistically deferred Nörlund mean convergence, for  $r \geq 1$ , is

$$\begin{aligned} \lim_{m \rightarrow \infty} \frac{1}{2m} \left| \left\{ n : n \leq \mathcal{R}_m \text{ and } 2mE(|Y_m - 0|^r) \right\} \right| &= \lim_{m \rightarrow \infty} \left( m^r \left( \frac{1}{\sqrt{m}} \right) + 0 \left( 1 - \frac{1}{\sqrt{m}} \right) \right) \\ &= \lim_{m \rightarrow \infty} m^{r-1/2} \\ &= \infty. \end{aligned}$$

This implies that the sequence  $(Y_m)$  is  $St_{DNP}$ -convergent but not  $St_{DNM}$ -convergent.

## 4. STATISTICAL DISTRIBUTION CONVERGENCE VIA DEFERRED NÖRLUND

**Definition 4.1.** The sequence of random variables  $(Y_m)$  is said to be distribution convergent (or convergent in distribution) to  $Y$ , if

$$\lim_{m \rightarrow \infty} F_{Y_m}(y) = F_Y(y)$$

for all  $y \in \mathbb{R}$  at which  $F_Y(y)$  is continuous.

Thoroughout the paper  $(F_{Y_m}(y))$  is the sequence of distribution functions of  $(Y_m)$  and  $F_Y(y)$  is the distribution function of  $Y$ .

**Definition 4.2.** The sequence  $(F_{Y_m}(y))$  is called as statistically distribution convergent (or  $St_{DC}$ ), if there exists  $F_Y(y)$  of random variable  $Y$  such that for each  $\varepsilon > 0$ ,

$$\lim_{m \rightarrow \infty} \frac{1}{m} \left| \left\{ n : n \leq m \text{ and } |F_{Y_m}(y) - F_Y(y)| \geq \varepsilon \right\} \right| = 0.$$

We may write this as

$$St_{DC} \lim_{m \rightarrow \infty} F_{Y_m}(y) = F_Y(y).$$

**Definition 4.3.** The sequence  $(F_{Y_m}(y))$  of distribution functions is called as deferred Nörlund statistically distribution convergent (or  $St_{DNDC}$ ), if there exists  $F_Y(y)$  of  $Y$  such that for each  $\varepsilon > 0$ ,

$$\lim_{m \rightarrow \infty} \frac{1}{\mathcal{R}_m} \left| \left\{ n : n \leq \mathcal{R}_m \text{ and } e_{y_m - n} g_n |F_{Y_m}(y) - F_Y(y)| \geq \varepsilon \right\} \right| = 0.$$

In this case, we say

$$St_{DNDC} \lim_{m \rightarrow \infty} F_{Y_m}(y) = F_Y(y).$$

**Theorem 4.4.** Suppose that  $St_{DNP} \lim_{m \rightarrow \infty} P(|Y_m - Y| \geq \varepsilon) = 0$ , then

$$St_{DNDC} \lim_{m \rightarrow \infty} F_{Y_m}(y) = F_Y(y).$$

*Proof.* Suppose that  $(F_{Y_m}(y))$  is distribution functions of  $(Y_m)$ , and  $F_Y(y)$  be the distribution function of  $Y$ . For  $i, j \in \mathbb{R}$  such that  $i < j$ , we have

$$(Y \leq i) = (Y_m \leq j, Y \leq i) + (Y_m \geq j, Y \leq i).$$

Further,

$$(Y_m \leq j, Y \leq i) \subseteq (Y_m \leq j),$$

which implies that

$$(4.1) \quad (Y \leq i) \subseteq (Y_m \leq j) + (Y_m \geq j, Y \leq i).$$

Let us take the probability to left hand side and right hand side of equation (4.1)

$$\begin{aligned} P(Y \leq i) &\leq P\{(Y_m \leq j) + (Y_m \geq j, Y \leq i)\} \\ &\leq P(Y_m \leq j) + P(Y_m \geq j, Y \leq i). \end{aligned}$$

It means that

$$(4.2) \quad F_{Y_m}(j) \geq F_Y(i) - P(Y_m \geq j, Y \leq i).$$

If  $Y_m \geq j, Y \leq i$ , then  $Y_m \geq j, -Y \geq -i$ , so that  $Y_m - Y > j - i$ , that is,

$$(Y_m \geq j, Y \leq i) \subseteq (Y_m - Y > j - i) \subseteq (|Y_m - Y| > j - i).$$

This means

$$P(Y_m \geq j, Y \leq i) \leq P(|Y_m - Y| > j - i).$$

As we know that  $i < j$  and  $St_{DNP} Y_m \rightarrow Y$ , we obtain

$$St_{DNP} \lim_{m \rightarrow \infty} P(Y_m \geq j, Y \leq i) = 0.$$

From (4.2) we get

$$St_{DNDC} \lim_{m \rightarrow \infty} F_{Y_m}(j) \geq F_Y(i).$$

Similarly, if  $j < a$  for any real constant  $a$ , then

$$(Y \leq j) = (Y \leq a, Y_m \leq j) + (Y > a, Y_m \leq j).$$

Consequently,

$$F_{Y_m}(j) \leq F_Y(a) + P(Y > a, Y_m \leq j)$$

and

$$St_{DNDC} \lim_{m \rightarrow \infty} P(Y > a, Y_m \leq j) = 0.$$

Therefore, we get

$$St_{DNDC} \lim_{m \rightarrow \infty} F_{Y_m}(j) \leq F_Y(a).$$

Thus, with  $i < j < a$ , we have

$$St_{DNDC} \lim_{m \rightarrow \infty} F_{Y_m}(j) = F_Y(i).$$

□

**Example 2:** Consider the random variables  $((Y_m), Y)$  of two dimensions as  $\{(0, 0), (0, 1), (1, 0), (1, 1)\}$  such that

$$(Y_m, Y) = \begin{cases} 0, & [P(Y_m = 0, Y = 0) = 0 = P(Y_m = 1, Y = 1)] \\ \frac{1}{2}, & [P(Y_m = 1, Y = 0) = 0 = P(Y_m = 0, Y = 1)]. \end{cases}$$

The distribution function of  $Y_m$  is given by  $Y_m = (\lambda_1 = 0, 1)$ , with probability mass function

$$(p_{y_m, \lambda_1}) = P(Y_m = \lambda_1), \text{ where } p_{y_m, 0} = \frac{1}{2} = p_{y_m, 1}$$

and for  $Y = \lambda_2(\lambda_2 = 0, 1)$ , with probability mass function

$$(p_{y_m, \lambda_2}) = P(Y_m = \lambda_2), \text{ where } p_{y, 0} = \frac{1}{2} = p_{y, 1}.$$

If  $(F_{Y_m}(y))$  is distribution functions of  $(Y_m)$  and  $F_Y(y)$  is the distribution function of  $Y$ , then

$$F_Y(y) = \lim_{m \rightarrow \infty} F_{Y_m}(y) = \begin{cases} 0, & (y < 0) \\ \frac{1}{2}, & (0 \leq y < 1). \\ 1, & (z > 1). \end{cases}$$

Thus, we get

$$St_{DNDC} \lim_{m \rightarrow \infty} F_{Y_m}(y) = F_Y(y), \text{ where } x_m = 2m-1, y_m = 4m-1, e_{y_m-m} = 2m \text{ and } g_m = 1.$$

But, it is not  $St_{PC}$  for the sequence of random variables, i.e.

$$St_{DNPC} \lim_{m \rightarrow \infty} P(|Y_m - Y| \geq \varepsilon) \neq 0, \text{ where } x_m = 2m-1, y_m = 4m-1, e_{y_m-m} = 2m \text{ and } g_m = 1.$$

## 5. APPLICATIONS

The hypothesis of the Korovkin-type theorems have been studied by several researchers in various field in different ways such as in, summability theory, functional analysis and probability theory. Korovkin-type approximation theorems have been investigated by many mathematicians under various background, involving function spaces, Banach spaces, and so on. Recently, Mohiuddine and Alamri studied Korovkin and Voronovskaya type approximation theorems in [11]. Further, Hazarika et al. [6] studied Korovkin approximation theorem for Bernstein operator of rough statistical convergence of triple sequences. For detailed study on Korovkin approximation theorem one may refer [2], [13], [14], [18], [22]. By  $\mathcal{C}(Y)$ , we denote the space of all continuous probability functions defined on a compact subset  $Z \subset \mathbb{R}$ . The space  $\mathcal{C}(Y)$  is a Banach space with respect to the norm

$$\|f\|_\infty = \sup_{z \in Y} \{|f(z)|\}, \quad f \in \mathcal{C}(Y).$$

We say that  $\mathcal{Y}$  is a positive linear operator of sequence of random variables if

$$\mathcal{Y}(f, z) \geq 0 \text{ whenever } f \geq 0.$$

Throughout,  $\mathcal{Y}_n : \mathcal{C}(Y) \rightarrow \mathcal{C}(Y)$  be a sequence of random variables of positive linear operators.

**Theorem 5.1.** [26] *Let  $\mathcal{Y}_n : \mathcal{C}(Y) \rightarrow \mathcal{C}(Y)$ . Then for all  $f \in \mathcal{C}(Y)$ , we have*

$$(5.1) \quad St_{DNP} \lim_{n \rightarrow \infty} \|\mathcal{Y}_n(f, z) - f(z)\|_\infty = 0$$

*iff*

$$(5.2) \quad St_{DNP} \lim_{n \rightarrow \infty} \|\mathcal{Y}_n(1, z) - 1\|_\infty = 0,$$

$$(5.3) \quad St_{DNP} \lim_{n \rightarrow \infty} \|\mathcal{Y}_n(z, z) - z\|_\infty = 0,$$

$$(5.4) \quad St_{DNP} \lim_{n \rightarrow \infty} \|\mathcal{Y}_n(z^2, z) - z^2\|_\infty = 0.$$

**Theorem 5.2.** *Let  $\mathcal{Y}_n : \mathcal{C}(Y) \rightarrow \mathcal{C}(Y)$ . Then for all  $f \in \mathcal{C}(Y)$ , we have*

$$(5.5) \quad St_{DNM} \lim_{n \rightarrow \infty} \|\mathcal{Y}_n(f, z) - f(z)\|_\infty = 0$$

*iff*

$$(5.6) \quad St_{DNM} \lim_{n \rightarrow \infty} \|\mathcal{Y}_n(1, z) - 1\|_\infty = 0,$$

$$(5.7) \quad St_{DNM} \lim_{n \rightarrow \infty} \|\mathcal{Y}_n(z, z) - z\|_\infty = 0,$$

$$(5.8) \quad St_{DNM} \lim_{n \rightarrow \infty} \|\mathcal{Y}_n(z^2, z) - z^2\|_\infty = 0.$$

**Theorem 5.3.** *Let  $\mathcal{Y}_n : \mathcal{C}(Y) \rightarrow \mathcal{C}(Y)$ . Then for all  $f \in \mathcal{C}(Y)$ , we have*

$$(5.9) \quad St_{DNDC} \lim_{n \rightarrow \infty} \|\mathcal{Y}_n(f, z) - f(z)\|_\infty = 0$$

*iff*

$$(5.10) \quad St_{DNDC} \lim_{n \rightarrow \infty} \|\mathcal{Y}_n(1, z) - 1\|_\infty = 0,$$



$$(5.11) \quad St_{DNDC} \lim_{n \rightarrow \infty} \|\mathcal{Y}_n(z, z) - z\|_\infty = 0,$$

$$(5.12) \quad St_{DNDC} \lim_{n \rightarrow \infty} \|\mathcal{Y}_n(z^2, z) - z^2\|_\infty = 0.$$

**Example 3:** Let  $\mathcal{M}_m(f, y)$  be a Meyer-König and Zeller operators on  $\mathcal{C}[0, 1]$  and  $Z = [0, 1]$  as defined in [1] as

$$\mathcal{M}_m(f, y) = (1 - y)^{m+1} \sum_{t=0}^{\infty} f\left(\frac{t}{t+m+1}\right) \binom{m+t}{t} y^t.$$

Further, let us consider a sequence of operators  $\mathcal{Y}_n : \mathcal{C}[0, 1] \rightarrow \mathcal{C}[0, 1]$  and  $(Y_n)$  as defined in example 2.4 such that

$$(5.13) \quad \mathcal{Y}_n(f, y) = [1 + F_{Y_m}(y)]\mathcal{M}_n(f), \quad (f \in \mathcal{C}[0, 1]),$$

where  $(F_{Y_m}(y))$  is defined in Example 2. Now we observe that

$$\begin{aligned} \mathcal{Y}_n(1, y) &= [1 + F_{Y_m}(y)] \cdot 1 = [1 + F_{Y_m}(y)], \\ \mathcal{Y}_n(u, z) &= [1 + F_{Y_m}(y)] \cdot y = [1 + F_{Y_m}(y)] \cdot y \end{aligned}$$

and

$$\mathcal{Y}_n(u^2, z) = [1 + F_{Y_m}(y)] \cdot \left\{ y^2 \left( \frac{m+2}{m+1} \right) + \frac{y}{m+1} \right\}.$$

Therefore, we have

$$St_{DNDC} \lim_{n \rightarrow \infty} \|\mathcal{Y}_n(1, y) - 1\|_\infty = 0,$$

$$St_{DNDC} \lim_{n \rightarrow \infty} \|\mathcal{Y}_n(y, y) - y\|_\infty = 0,$$

$$St_{DNDC} \lim_{n \rightarrow \infty} \|\mathcal{Y}_n(y^2, y) - y^2\|_\infty = 0.$$

Hence,  $\mathcal{Y}_n(f, y)$  fulfills (5.10), (5.11) and (5.12). Thus, from Theorem 5.3

$$St_{DNDC} \lim_{n \rightarrow \infty} \|\mathcal{Y}_n(f, y) - f\|_\infty = 0.$$

Hence, it is  $(DNDC)$ -convergent. However,  $(Y_m)$  is neither  $(DN)$ -statistical convergent nor  $(DN)$ -convergent. Thus, we can exhibit that the work in [24] does not hold for our operators described in (5.13). Hence, our Theorem 5.3 is stronger than the theorem proved in [24].

## 6. CONCLUSION

Upon prior analysis, our interest is to modify the studies of Srivastava et al. [25] and introduce various aspects of statistical convergence for the sequences of random variables and sequences of real numbers via deferred Norlund summability mean. We first study various results presenting the connection by using fundamental limit concepts of sequences of random variables. As an applications of our findings, we present new Krorvkin-type approximation results and also demonstrated the effectiveness of the findings. As a future work one can obtain the corresponding results of the present paper using deferred Euler summability mean.

## Compliance with ethical standards

**Availability of data and material:** Not applicable.

**Conflict of interest:** The authors declare that they have no conflict of interest.

**Ethical approval:** This article does not contain any studies with human participants or animals performed by any of the authors.

**Acknowledgement:** The corresponding author thanks the Council of Scientific and Industrial Research (CSIR), India for partial support under Grant No. 25(0288)/18/EMR-II, dated 24/05/2018.

## REFERENCES

- [1] A. Altin, O. Doğru and F. Taşdelen *The generalization of Meyer-König and Zeller operators by generating functions*, J. Math. Anal. Appl., **312** (2005), 181–194.
- [2] H. Dutta, S. K. Paikray and B. B. Jena *On statistical deferred Cesàro summability*, Current Trends in Mathematical Analysis and Its Interdisciplinary Applications. Birkhäuser, Cham, **487(1)** (2019), 885–909.
- [3] A. Esi and E. Savas, *On lacunary statistically convergent triple sequences in probabilistic normed space*, Appl. Math. Inf. Sci., **9(5)** (2015), 2529–2534.
- [4] M. Et, P. Baliarsingh and H. Sengul, *Deferred statistical convergence and strongly deferred summable functions*, AIP Conf. Proc., **2183(1)** (2019).
- [5] H. Fast, *Sur la convergence statistique*, Colloquium Mathematicae, **2**(1951), 241–244.
- [6] B. Hazarika, N. Subramanian, and M. Mursaleen *Korovkin-type approximation theorem for Bernstein operator of rough statistical convergence of triple sequences*, Adv. Oper. Theory, **5(2)** (2020): 324–335.
- [7] I. J. Schoenberg, *The integrability of certain functions and related summability methods*, The American Mathematical Monthly, **66(5)** (1959), 361–775.
- [8] D. Rath and B. C. Tripathy, *Matrix maps on sequence spaces associated with sets of integers*, Indian J. Pure Appl. Math., **27(2)** (1996), 197–206.
- [9] B. B. Jena, S. K. Paikray and H. Dutta *On Various New Concepts of Statistical Convergence for Sequences of Random Variables via Deferred Cesàro Mean*, J. Math. Anal. Appl., **487(1)** (2020), 1239–50.
- [10] S. Jasrotia, U. P. Singh and K. Raj, *Applications of Statistical Convergence of order  $(\eta, \delta + \gamma)$  in difference sequence spaces of fuzzy numbers*, J. Intell. Fuzzy Systems, **40(3)**(2021), 4695–4703.
- [11] S. A. Mohiuddine and B. A. S. Alamri *Generalization of equi-statistical convergence via weighted lacunary sequence with associated Korovkin and Voronovskaya type approximation theorems*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM, **113** (2019): 1955–1973.
- [12] S. A. Mohiuddine, A. Alotaibi and M. Mursaleen, *Statistical convergence of double sequences in locally solid Riesz spaces*, Abstr. Appl. Anal., (2012), **2012**.
- [13] S. A. Mohiuddine, A. Alotaibi, and M. Mursaleen *Statistical summability  $(C, 1)$  and a Korovkin type approximation theorem*, J. Inequal. Appl., **2012(1)** (2012): 1–8.
- [14] M. Mursaleen, V. Karakaya, M. Ertürk and F. Gürsoy *Weighted statistical convergence and its application to Korovkin type approximation theorem*, Adv. Oper. Theory, **218(18)** (2012): 9132–9137.
- [15] M. Mursaleen, *Applied Summability Methods*, Springer Briefs. Springer, New York, 2014.
- [16] Jemima, D. Eunice, Srinivasan, V. *Norlund statistical convergence and Tauberian conditions for statistical convergence from statistical summability using Norlund means in non-Archimedean fields*. Journal of Mathematics and Computer Science, 24, no. 4 (2022), 299–307.
- [17] Khan, V. A., Fatima, H., Khan, M. D., Ahamd, A. *Spaces of neutrosophic  $\lambda$ -statistical convergence sequences and their properties*. Journal of Mathematics and Computer Science, 23, no. 1 (2021), 1–9.
- [18] K. Raj and A. Choudhary, *Relative modular uniform approximation by means of the power series method with applications*, Rev. Un. Mat. Argentina, **60** (2019), 187–208.
- [19] K. Raj and S. Pandoh *Some vector-valued statistical convergent sequence spaces*, Malaya J. Mat., **3** (2015): 161–167.
- [20] A. D. Gadjiev and C. Orhan, *Some approximation theorems via statistical convergence*, Rocky Mountain Journal of Mathematics, **32(1)**(2002), 129–138.

- [21] J. A. Fridy, *On statistical convergence*, Analysis, **5(4)** (1985), 301-313.
- [22] B.C. Tripathy, A. Esi and T. Balakrushna, *On a new type of generalized difference Cesàro sequence spaces*, Soochow J. Math. **31**, (2005) 333-340.
- [23] S. Ghosal, *Statistical convergence for a sequence of random variables and limit theorems*, Appl. Math. (Prague). **58**, (2013) 423-437.
- [24] H. M. Srivastava, B. B. Jena, S. K. Paikray and U. K. Misra *Generalized equi-statistical convergence of the deferred Nörlund summability and its applications to associated approximation theorems*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM, **112** (2018), 1487–1501.
- [25] Srivastava HM, Jena BB, Paikray SK. *Statistical probability convergence via the deferred Nörlund mean and its applications to approximation theorems*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM, 2020;114:1-14.
- [26] H. M. Srivastava, B. B. Jena, S. K. Paikray *Deferred Cesàro statistical probability convergence and its applications to approximation theorems*, J. Nonlinear Convex Anal., **20** (2019), 1777–1792.
- [27] H. M. Srivastava, B. B. Jena and S. K. Paikray, *A certain class of statistical probability convergence and its applications to approximation theorems*, Appl. Anal. Discrete Math., **14(3)** (2020), 579-598.

SCHOOL OF MATHEMATICS SHRI MATA VAISHNO DEVI UNIVERSITY, KATRA-182320, J & K (INDIA)

Email address: [kuldipraj68@gmail.com](mailto:kuldipraj68@gmail.com)

Email address: [swatijasrotial2@gmail.com](mailto:swatijasrotial2@gmail.com)

REZA SAADATI

SCHOOL OF MATHEMATICS, IRAN UNIVERSITY OF SCIENCE AND TECHNOLOGY, NARMAK, TEHRAN, IRAN

Email address: [rsaadati@eml.cc](mailto:rsaadati@eml.cc), [rsaadati@iust.ac.ir](mailto:rsaadati@iust.ac.ir)