

# GROUND STATES OF SCHRÖDINGER SYSTEMS WITH CHERN-SIMONS GAUGE FIELDS

YAHUI JIANG, TAIYONG CHEN, JIANJUN ZHANG, MARCO SQUASSINA, AND NOUF ALMOUSA

ABSTRACT. We are concerned with the following coupled nonlinear Schrödinger system

$$\begin{cases} -\Delta u + u + \left( \int_{|x|}^{\infty} \frac{h(s)}{s} u^2(s) ds + \frac{h^2(|x|)}{|x|^2} \right) u = |u|^{2p-2} u + b|v|^p |u|^{p-2} u, & x \in \mathbb{R}^2, \\ -\Delta v + \omega v + \left( \int_{|x|}^{\infty} \frac{g(s)}{s} v^2(s) ds + \frac{g^2(|x|)}{|x|^2} \right) v = |v|^{2p-2} v + b|u|^p |v|^{p-2} v, & x \in \mathbb{R}^2, \end{cases}$$

where  $\omega, b > 0, p > 1$ . By virtue of the variational approach, we show the existence of nontrivial ground state solutions. Precisely, the system above admits a positive ground state solution if  $p > 3$  and  $b > 0$  large enough or if  $p \in (2, 3]$  and  $b > 0$  small.

## 1. INTRODUCTION

In this paper, we consider the following coupled Schrödinger equations with Chern-Simons gauge fields

$$(1.1) \quad \begin{cases} -\Delta u + u + \left( \int_{|x|}^{\infty} \frac{h(s)}{s} u^2(s) ds + \frac{h^2(|x|)}{|x|^2} \right) u = |u|^{2p-2} u + b|v|^p |u|^{p-2} u, & x \in \mathbb{R}^2, \\ -\Delta v + \omega v + \left( \int_{|x|}^{\infty} \frac{g(s)}{s} v^2(s) ds + \frac{g^2(|x|)}{|x|^2} \right) v = |v|^{2p-2} v + b|u|^p |v|^{p-2} v, & x \in \mathbb{R}^2, \end{cases}$$

where  $\omega > 0, b > 0, p > 1$  and

$$h(s) = \int_0^s \frac{r}{2} u^2(r) dr, \quad g(s) = \int_0^s \frac{r}{2} v^2(r) dr.$$

When  $b = 0$ , then system (1.1) is uncoupled and it reduces to two equations of the same type. In recently years, a single nonlinear Schrödinger equations coupled with the Chern-Simons gauge field as follows has received much attention

$$(1.2) \quad \begin{cases} iD_0\phi + (D_1D_1 + D_2D_2)\phi = -f(\phi), \\ \partial_0 A_1 - \partial_1 A_0 = -\text{Im}(\bar{\phi} D_2 \phi), \\ \partial_0 A_2 - \partial_2 A_0 = \text{Im}(\bar{\phi} D_1 \phi), \\ \partial_1 A_2 - \partial_2 A_1 = -\frac{1}{2}|\phi|^2, \end{cases}$$

where  $i$  denotes the imaginary unit,  $\partial_0 = \frac{\partial}{\partial t}$ ,  $\partial_1 = \frac{\partial}{\partial x_1}$ ,  $\partial_2 = \frac{\partial}{\partial x_2}$ ,  $(t, x_1, x_2) \in \mathbb{R}^{1+2}$ ,  $\phi : \mathbb{R}^{1+2} \rightarrow \mathbb{C}$  is the complex scalar field,  $A_\mu : \mathbb{R}^{1+2} \rightarrow \mathbb{C}$  is the gauge field and  $D_\mu = -\partial_\mu + iA_\mu$  is the covariant derivative for  $\mu = 0, 1, 2$ . The Chern-Simons-Schrödinger system consists of Schrödinger equations augmented by the gauge field, which was first proposed and studied in [13, 14]. The

*Date:* May 2, 2023.

2000 *Mathematics Subject Classification.* 35B09, 35J50, 81T10.

*Key words and phrases.* Schrödinger systems, Ground states, Chern-Simons gauge fields, Variational methods.

(1) Corresponding author: marsquassina@pnu.edu.sa.

(2) J. J. Zhang was partially supported by NSFC(No.11871123).

model was proposed to study vortex solutions, which carry both electric and magnetic charges. This feature of the model is important for the study of the high-temperature superconductor, fractional quantum Hall effect and Aharonov-Bohm scattering. For more details about system (1.2), we refer the readers to [8, 10, 11]. System (1.2) is invariant under gauge transformation

$$\phi \rightarrow \phi e^{i\chi}, \quad A_\mu = A_\mu - \partial_\mu \chi,$$

for any arbitrary  $C^\infty$  function  $\chi$ .

Byeon, Hun and Seok [5] investigated the existence of standing wave solutions for system (1.2) with power type nonlinearity, that is  $f(u) = \lambda|u|^{p-2}u$  with  $p > 2$  and  $\lambda > 0$ . By using the ansatz

$$\begin{cases} \phi(t, x) = u(|x|)e^{i\omega t}, & A_0(t, x) = k(|x|), \\ A_1(t, x) = \frac{x_2}{|x|^2}h(|x|), & A_2(t, x) = -\frac{x_1}{|x|^2}h(|x|). \end{cases}$$

Byeon et al. got the following nonlocal semilinear elliptic equation

$$(1.3) \quad -\Delta u + (\omega + \xi)u + \left( \int_{|x|}^\infty \frac{h(s)}{s} u^2(s) ds + \frac{h^2(|x|)}{|x|^2} \right) u = f(u), \quad x \in \mathbb{R}^2,$$

where  $\xi$  is a constant and  $h(s)$  is defined as above. Byeon et al. showed that the existence and nonexistence of positive solutions for (1.3) were established depending on the range of  $p > 2$  and  $\lambda > 0$ . For the special case  $p = 4$ , there exist solutions if  $\lambda > 1$ . It seems hard to obtain the boundedness of Palais-Smale sequence when  $p \in (4, 6)$ . They constructed a Nehari-Pohozaev manifold to obtain the boundedness of Palais-Smale sequence. For  $p \in (2, 4)$ , Pomponio and Ruiz [19] proved the existence and nonexistence of positive solutions for (1.3) under the different range of the  $\omega$ . A series of existence and nonexistence results of solutions for (1.3) has been researched in [6, 12, 15, 19, 20, 25]. There has been increasing interest in studying the existence, multiplicity and the concentration behavior of the solutions for the problems above mentioned.

Problem (1.1) is a nonlocal problem due to the appearance of the term  $\int_{|x|}^\infty \frac{h(s)}{s} u^2(s) ds$ , which indicates that (1.1) is not a pointwise identity. This causes some mathematical difficulties which make the study of such a problem particularly interesting. System (1.1) is quite different from the following local scalar field system

$$(1.4) \quad \begin{cases} -\Delta u + u = |u|^{2q-2} + b|v|^q|u|^{q-2}u, & \text{in } \mathbb{R}^N, \\ -\Delta v + \omega^2 v = |v|^{2q-2} + b|u|^q|v|^{q-2}v, & \text{in } \mathbb{R}^N, \end{cases}$$

for  $\omega > 0$ ,  $b \in \mathbb{R}$ ,  $q \in (2, 2^*)$ , which does not depend on the nonlocal term any more. The coupled nonlinear Schrödinger system (1.4) has attracted considerable attention in the past fifteen years. Maia et al. [18] by using the variational methods and the ideas of Rabinowitz [21] investigated the existence of positive ground state solutions for system (1.4) under sufficient conditions on parameter  $b$  and  $\omega$ . For more progress in this aspect, we refer to [1, 2, 4, 16, 23] and the references therein.

The energy functional for system (1.1)  $I : E \rightarrow \mathbb{R}$  is defined by

$$I(u, v) = \frac{1}{2} \|(u, v)\|_E^2 + \frac{1}{2} (B(u) + B(v)) - \frac{1}{2p} F(u, v),$$

where

$$B(u) = \int_{\mathbb{R}^2} \frac{u^2}{|x|^2} \left( \int_0^{|x|} \frac{s}{2} u^2(s) ds \right)^2 dx,$$

$$F(u, v) = \int_{\mathbb{R}^2} (u^{2p} + v^{2p} + 2b|uv|^p) dx.$$

Here  $E$  denotes the subspace of radially symmetric functions in  $H_r^1(\mathbb{R}^2) \times H_r^1(\mathbb{R}^2)$  with the norm

$$\|(u, v)\|_E^2 := \int_{\mathbb{R}^2} (|\nabla u|^2 + u^2 + |\nabla v|^2 + \omega v^2) dx.$$

The functional  $I$  is of class  $C^1(E)$ , its critical point  $(u, v)$  is a weak solution of (1.1) and by standard regularity theory is a classical solution.

Motivated by [18], we try to study the existence of positive ground state solutions for coupled Schrödinger equations with Chern-Simons gauge fields (1.1) with suitable conditions on  $\omega, b$ .

One of the main difficulties is the boundedness of Palais-Smale sequences if we try to use directly by the mountain pass theorem to get the critical points of  $I$  in  $E$ . For  $p \geq 3$ , it is standard to show that Palais-Smale condition holds for  $I$ . For  $p \in (2, 3)$ , the functional  $I$  has the mountain-pass geometry. However, it seems hard to prove the Palais-Smale condition holds for the functional  $I$ . Motivated by [5], by using a constrained minimizer on Nehari-Pohozaev manifold we circumvent this obstacle.

Another problem is the existence of positive ground state for system (1.1), i.e., a minimal action solution  $(u, v)$  with both  $u > 0, v > 0$  nontrivial. We point out that the system (1.1) also possesses a trivial solution  $(0, 0)$  and semi-trivial solutions of type  $(u, 0)$  or  $(0, v)$ . A solution  $(u, v)$  of (1.1) is nontrivial if  $u \not\equiv 0$  and  $v \not\equiv 0$ . Here we overcome this obstacle by energy estimation.

We now state the main result of the paper. The constants  $b_1, b_2, b_3, b_4, b_\delta$  involved in the statement depends on the ground state of the single equation. We will give the corresponding expressions in Section 3.

**Theorem 1.1.** *Assume that one of the following conditions holds*

(i)  $p \in (2, 3]$  and  $b \in (0, b_\delta)$  sufficiently small,

(ii)  $p \in (3, 3 + \sqrt{6})$  and  $b > \max\{b_1, b_2\}$ ,

(iii)  $p \in [3 + \sqrt{6}, \infty)$  and  $b > \max\{b_3, b_4\}$ ,

*then the system (1.1) admits a positive vector ground state.*

Additionally, we prove also the following nonexistence result.

**Theorem 1.2.** *There exist  $\tilde{b} > 0$  sufficiently small and  $\tilde{\omega} > 0$  sufficiently large such that  $b \in (0, \tilde{b})$  and  $\omega > \tilde{\omega}$ , then the system (1.1) has only trivial solution if  $p \in (1, 2]$ .*

Compared with the case of  $p > 2$ , it seems that the case of  $p \in (1, 2]$  becomes more complicated and we will consider it in a forthcoming paper. The rest of this paper is organized as follows. In Section 2, we present some notations and preliminary results and prove the nonexistence result Theorem 1.2. Then we give the proof of the existence of a positive ground state in Theorem 1.1. The item (ii) and (iii) are proved in Section 3, and the item (i) is proved in Section 4.

## 2. PRELIMINARIES

To prove the main results, we use the following notations:

- $E := H_r^1(\mathbb{R}^2) \times H_r^1(\mathbb{R}^2)$  with norm

$$\|(u, v)\|_E^2 = \|u\|_{H_r^1(\mathbb{R}^2)}^2 + \|v\|_{H_r^1(\mathbb{R}^2)}^2.$$

- $L^{2p}(\mathbb{R}^2) \times L^{2p}(\mathbb{R}^2)$  for  $p > 1$  with the norm

$$\|(u, v)\|_{2p}^{2p} = \|u\|_{2p}^{2p} + \|v\|_{2p}^{2p}.$$

- $\mathbb{L}_{loc}^{2p}(\mathbb{R}^2) := L_{loc}^{2p}(\mathbb{R}^2) \times L_{loc}^{2p}(\mathbb{R}^2)$ ;

**Lemma 2.1.** [5] Suppose that a sequence  $u_n$  converges weakly to a function  $u$  in  $H_r^1(\mathbb{R}^2)$  as  $n \rightarrow \infty$ . Then for each  $\varphi \in H_r^1(\mathbb{R}^2)$ ,  $B'(u_n)$ ,  $B'(u_n)\varphi$  and  $B'(u_n)u_n$  converges up to a subsequence to  $B(u)$ ,  $B'(u)\varphi$  and  $B'(u)u$ , respectively, as  $n \rightarrow \infty$ .

**Lemma 2.2.** [5] For  $u \in H_r^1(\mathbb{R}^2)$ , the following inequality holds

$$\int_{\mathbb{R}^2} |u|^4 dx \leq 4 \left( \int_{\mathbb{R}^2} |\nabla u|^2 dx \right)^{\frac{1}{2}} \left( \int_{\mathbb{R}^2} \frac{u^2}{|x|^2} \left( \int_0^{|x|} \frac{s}{2} u^2(s) ds \right)^2 dx \right)^{\frac{1}{2}}.$$

Furthermore, the equality is attained by a continuum of functions

$$\left\{ u_l = \frac{\sqrt{8}l}{1 + |lx|^2} \in H_r^1(\mathbb{R}^2) \mid l \in (0, \infty) \right\},$$

$$\frac{1}{4} \int_{\mathbb{R}^2} |u_l|^4 dx = \int_{\mathbb{R}^2} |\nabla u_l|^2 dx = \int_{\mathbb{R}^2} \frac{u_l^2}{|x|^2} \left( \int_0^{|x|} \frac{s}{2} u_l^2(s) ds \right)^2 dx = \frac{16\pi l^2}{3}.$$

**Lemma 2.3.** If  $(u, v)$  is a solution of (1.1) then it satisfies the Pohozaev identity

$$(2.1) \quad \|u\|_2^2 + \omega \|v\|_2^2 + 2 \int_{\mathbb{R}^2} \left( \frac{h^2(|x|)}{|x|^2} u^2 + \frac{g^2(|x|)}{|x|^2} v^2 \right) dx = \frac{1}{p} \left( \|(u, v)\|_{2p}^{2p} + 2b \|uv\|_p^p \right).$$

*Proof.* We adopt some idea in [5]. Assume that  $(u, v) \in E$  is a weak solution for problem (1.1). Similar to [3] and [5], we know that  $\int_{|x|}^\infty \frac{h(s)}{s} u^2(s) ds, \int_{|x|}^\infty \frac{g(s)}{s} v^2(s) ds, \frac{h^2(|x|)}{|x|^2}, \frac{g^2(|x|)}{|x|^2} \in L^\infty(\mathbb{R}^2)$ . Thus, the standard elliptic estimates [9] imply that  $u, v \in C_{loc}^{1,\gamma}(\mathbb{R}^2)$  for some  $\gamma > 0$ . Then we obtain  $\int_{|x|}^\infty \frac{h(s)}{s} u^2(s) ds, \int_{|x|}^\infty \frac{g(s)}{s} v^2(s) ds, \frac{h^2(|x|)}{|x|^2}, \frac{g^2(|x|)}{|x|^2} \in C(\mathbb{R}^2)$ . Since  $u, v \in H_r^1(\mathbb{R}^2)$ , we deduce that  $u, v \in C^2(\mathbb{R}^2)$ . Then, multiplying the first equation in (1.1) by  $x \cdot \nabla u$  and integrating by

parts on a ball  $B_R = \{x \in \mathbb{R}^2 : |x| < R\}$ , then

$$\begin{aligned}
\int_{B_R} \Delta u (\nabla u \cdot x) dx &= \frac{R}{2} \int_{\partial B_R} |\nabla u|^2 dS_x, \\
\int_{B_R} u (\nabla u \cdot x) dx &= - \int_{B_R} u^2 dx + o_R(1), \\
\int_{B_R} |u|^{2p-2} u (\nabla u \cdot x) dx &= -\frac{1}{p} \int_{B_R} u^{2p} dx + o_R(1), \\
\int_{B_R} \left( \int_{|x|}^{\infty} \frac{h(s)}{s} u^2(s) ds \right) u (\nabla u \cdot x) dx &+ \int_{B_R} \frac{h^2(|x|)}{|x|^2} u (\nabla u \cdot x) dx = \pi h^2(R) u^2(R) \\
&+ \pi \left( \int_R^{+\infty} \frac{h(s)}{s} u^2(s) ds \right) u^2(R) R^2 - 2 \int_{\mathbb{R}^2} \frac{h^2(|x|)}{|x|^2} u^2 dx + o_R(1).
\end{aligned}$$

Thus, we deduce that

$$\begin{aligned}
(2.2) \quad & \frac{R}{2} \int_{\partial B_R} |\nabla u|^2 dS_x + \int_{B_R} u^2 dx - \pi h^2(R) u^2(R) - \pi \left( \int_R^{+\infty} \frac{h(s)}{s} u^2(s) ds \right) u^2(R) R^2 \\
& + 2 \int_{\mathbb{R}^2} \frac{h^2(|x|)}{|x|^2} u^2 dx = \frac{1}{p} \int_{B_R} u^{2p} dx - b \int_{B_R} |v|^p |u|^{p-2} u x \cdot \nabla u dx.
\end{aligned}$$

In a similar way,

$$\begin{aligned}
(2.3) \quad & \frac{R}{2} \int_{\partial B_R} |\nabla v|^2 dS_x + \int_{B_R} \omega v^2 dx - \pi g^2(R) v^2(R) - \pi \left( \int_R^{+\infty} \frac{g(s)}{s} v^2(s) ds \right) v^2(R) R^2 \\
& + 2 \int_{\mathbb{R}^2} \frac{g^2(|x|)}{|x|^2} v^2 dx = \frac{1}{p} \int_{B_R} v^{2p} dx - b \int_{B_R} |u|^p |v|^{p-2} v x \cdot \nabla v dx,
\end{aligned}$$

can be obtained. Then, summing up (2.2) and (2.3), we obtain

$$\begin{aligned}
& \int_{B_R} (u^2 + \omega v^2) dx + 2 \int_{\mathbb{R}^2} \left( \frac{h^2(|x|)}{|x|^2} u^2 + \frac{g^2(|x|)}{|x|^2} v^2 \right) dx - \frac{1}{p} \int_{B_R} (u^{2p} + v^{2p}) dx - \frac{2b}{p} \int_{B_R} |uv|^p dx \\
& = \pi \left( \int_R^{+\infty} \frac{h(s)}{s} u^2(s) ds \right) u^2(R) R^2 + \pi \left( \int_R^{+\infty} \frac{g(s)}{s} v^2(s) ds \right) v^2(R) R^2 + \pi h^2(R) u^2(R) \\
& + \pi g^2(R) v^2(R) - \frac{R}{2} \int_{\partial B_R} (|\nabla u|^2 + |\nabla v|^2) dS_x.
\end{aligned}$$

Arguing as in [5, Proposition 2.3.], there exists a suitable sequence  $R_n \rightarrow \infty$  on which the right hand side above tends to zero. Passing to the limit we get the identity. This completes the proof.  $\square$

*Proof of Theorem 1.2.* Let  $(u, v)$  be a solution of (1.1). By Lemma 2.2, we obtain

$$\begin{aligned}
0 &= \int_{\mathbb{R}^2} (|\nabla u|^2 + u^2 + |\nabla v|^2 + \omega v^2) dx + 3 \left( B(u) + B(v) \right) - \int_{\mathbb{R}^2} (u^{2p} + v^{2p} + 2b|uv|^p) dx \\
&\geq \int_{\mathbb{R}^2} \left( u^2 + \frac{1}{2} u^4 - (1+b)u^{2p} \right) dx + \int_{\mathbb{R}^2} \left( \omega v^2 + \frac{1}{2} v^4 - (1+b)v^{2p} \right) dx.
\end{aligned}$$

Denote

$$\begin{aligned} f_1(t) &= t^2 + \frac{1}{2}t^4 - (1+b)t^{2p}, \\ f'_1(t) &= 2t + 2t^3 - 2p(1+b)t^{2p-1} = 2t(1 + t^2 - 2p(1+b)t^{2p-2}), \end{aligned}$$

There exists  $\tilde{b} > 0$  small enough such that

$$1 + \left(p(1 + \tilde{b})\right)^{\frac{1}{2-p}} (p-1)^{\frac{p-1}{2-p}} (p-2) = 0.$$

Then, we have

$$f_1(t) \geq 0, \quad t \in \mathbb{R}, \quad b \in (0, \tilde{b}).$$

There exists a  $\tilde{\omega} > 0$  such that the function  $t \mapsto \omega t^2 + \frac{1}{2}t^4 - (1+b)t^{2p}$  is nonnegative and strictly increases as  $\omega > \tilde{\omega}, b \in (0, \tilde{b})$ . Hence,  $(u, v)$  must be identically  $(0, 0)$ . This completes the proof.  $\square$

### 3. PROOF OF (ii) AND (iii) OF THEOREM 1.1.

Consider the following problem

$$(3.1) \quad -\Delta u + \omega u + \left( \int_{|x|}^{\infty} \frac{h(s)}{s} u^2(s) ds + \frac{h^2(|x|)}{|x|^2} \right) u = |u|^{2p-2} u, \quad x \in \mathbb{R}^2,$$

By [5], when  $p \in (3, \infty)$ , problem (3.1) admits a positive ground state solution  $u_\omega$ . To be more precise, define the associated energy functional by

$$J_\omega = \frac{1}{2} \int_{\mathbb{R}^2} (|\nabla u|^2 + \omega u^2) dx + \frac{1}{2} B(u) - \frac{1}{2p} \int_{\mathbb{R}^2} u^{2p} dx.$$

Denote the ground state level by

$$E_\omega := \min \{ J_\omega(u) : u \in H_r^1(\mathbb{R}^2) \setminus \{0\}, J'_\omega(u) = 0 \},$$

Moreover,

$$E_\omega := \inf_{u \in \mathcal{N}_\omega} J_\omega(u),$$

where

$$\mathcal{N}_\omega := \left\{ u \in H_r^1(\mathbb{R}^2) \setminus \{0\} : \int_{\mathbb{R}^2} (|\nabla u|^2 + \omega u^2) dx + 3B(u) = \int_{\mathbb{R}^2} u^{2p} dx \right\}.$$

Define the Nehari manifold of problem (1.1) by

$$\mathcal{N} := \{ (u, v) \in E \setminus \{(0, 0)\} \mid \langle I'(u, v), (u, v) \rangle = 0 \}.$$

The corresponding groundstate energy is described as

$$c_{\mathcal{N}} := \inf_{(u, v) \in \mathcal{N}} I(u, v).$$

**Lemma 3.1.** (*Mountain-Pass geometry*) Assume  $b > 0$ , then the functional  $I$  satisfies the following conditions

- (i) There exists a positive constant  $r > 0$  such that  $I(u, v) > 0$  for  $\|(u, v)\|_E = r$ ;
- (ii) There exists  $(e_1, e_2) \in E$  with  $\|(e_1, e_2)\|_E > r$  such that  $I(e_1, e_2) < 0$ .

*Proof.* Since

$$2b \int |uv|^p dx \leq b(\|u\|_{2p}^{2p} + \|v\|_{2p}^{2p}),$$

and by the Sobolev embedding theorem, there exists a positive constant  $C$  such that

$$\begin{aligned} I(u, v) &= \frac{1}{2} \|(u, v)\|_E^2 + \frac{1}{2} (B(u) + B(v)) - F(u, v); \\ &\geq \frac{1}{2} \|(u, v)\|_E^2 - C \|(u, v)\|_E^{2p}. \end{aligned}$$

Hence, there exists  $r > 0$  such that

$$\inf_{\|(u, v)\|_E = r} I(u, v) > 0.$$

For any  $(u, v) \in E \setminus \{(0, 0)\}$  and  $t > 0$ ,

$$I(tu, tv) = \frac{t^2}{2} \|(u, v)\|_E^2 + \frac{t^6}{2} (B(u) + B(v)) - \frac{t^{2p}}{2p} F(u, v),$$

which implies that  $I(tu, tv) \rightarrow -\infty$  as  $t \rightarrow +\infty$ . This completes the proof.  $\square$

That is  $I$  satisfies the geometric conditions of the Mountain-Pass theorem. Define

$$c = \inf_{\gamma \in \Gamma} \max_{t \in [0, 1]} I(\gamma(t)),$$

where  $\Gamma = \{\gamma \in C([0, 1], E) : I(\gamma(0)) = 0, I(\gamma(1)) < 0\}$ .

**Lemma 3.2.** *For every  $(u, v) \in E \setminus \{(0, 0)\}$  there exists a unique  $\bar{t}_{uv} > 0$  such that  $\bar{t}_{uv}(u, v) \in \mathcal{N}$ . The maximum of  $I(tu, tv)$  for  $t \geq 0$  is achieved at  $t = \bar{t}_{uv}$ .*

*Proof.*  $\forall (u, v) \in E \setminus \{(0, 0)\}$  and  $t > 0$ , let

$$g(t) := I(tu, tv) = \frac{t^2}{2} \|(u, v)\|_E^2 + \frac{t^6}{2} (B(u) + B(v)) - \frac{t^{2p}}{2p} F(u, v).$$

By Lemma 3.1, we deduce that there exists  $\bar{t} = \bar{t}_{uv} > 0$  such that

$$g(\bar{t}) = \max_{t > 0} g(t).$$

Moreover

$$\begin{aligned} g'(t) &= t \|(u, v)\|_E^2 + 3t^5 (B(u) + B(v)) - t^{2p-1} F(u, v) \\ &= t^{2p-1} \left( \frac{1}{t^{2p-2}} \|(u, v)\|_E^2 + \frac{3}{t^{2p-6}} (B(u) + B(v)) - F(u, v) \right). \end{aligned}$$

As  $p > 3$ , thus  $g'(t)$  is strictly decreasing, the point  $t = \bar{t}_{uv}$  is the unique value of  $t > 0$  at which  $\bar{t}_{uv}(u, v) \in \mathcal{N}$ . This completes the proof.  $\square$

**Lemma 3.3.**  $c_{\mathcal{N}} = c$ .

*Proof.* Define

$$c_1 := \inf_{(u, v) \in E \setminus \{(0, 0)\}} \max_{t \geq 0} I(tu, tv).$$

By Lemma 3.2, we can obtain  $c_{\mathcal{N}} = c_1$ . Since  $I(tu, tv) < 0$  for any  $t$  large, it follows that  $c \leq c_1$ . Since every  $\gamma \in \Gamma$  intersects  $\mathcal{N}$ , so that  $c \geq c_{\mathcal{N}}$ . This completes the proof.  $\square$

Set

$$\begin{aligned} b_1 &= \left( \frac{(p-1)3^{\frac{p}{p-1}} \|u_1\|_{2p}^{2p}}{pE_\omega} \right)^{p-1}, & b_2 &= \left( \frac{(p-1)3^{\frac{p}{p-1}} \|u_\omega\|_{2p}^{2p}}{pE_1} \right)^{p-1}, \\ b_3 &= \left( \frac{(p-1)3^{\frac{3}{p-3}} \|u_1\|_{2p}^{2p}}{pE_\omega} \right)^{p-1}, & b_4 &= \left( \frac{(p-1)3^{\frac{3}{p-3}} \|u_\omega\|_{2p}^{2p}}{pE_1} \right)^{p-1}. \end{aligned}$$

**Lemma 3.4.** *Assume that one of the following conditions holds*

(i)  $p \in (3, 3 + \sqrt{6})$  and  $b > \max\{b_1, b_2\}$ ,

(ii)  $p \in [3 + \sqrt{6}, \infty)$  and  $b > \max\{b_3, b_4\}$ ,

then  $c_N < \min\{E_1, E_\omega\}$ , where  $E_1$  is a groundstate energy of problem (3.1) with  $\omega = 1$ .

*Proof.* Denote

$$\begin{aligned} a(u, v) &:= \|\nabla u\|_2^2 + \|\nabla v\|_2^2, \\ b(u, v) &:= \|u\|_2^2 + \omega \|v\|_2^2, \\ c(u, v) &:= B(u) + B(v). \end{aligned}$$

For fixed  $(u, v) \in E \setminus \{(0, 0)\}$ , we have

$$\begin{aligned} c_N &\leq \max_{t \geq 0} I(tu, tv) \\ &\leq \max_{t \geq 0} \left\{ \frac{t^2}{2} (a(u, v) + b(u, v)) + \frac{t^6}{2} c(u, v) - \frac{t^{2p}}{2p} F(u, v) \right\} \\ &\leq \max_{t \geq 0} \left\{ \frac{t^2}{2} (a(u, v) + b(u, v)) - \frac{t^{2p}}{4p} F(u, v) \right\} + \max_{t \geq 0} \left\{ \frac{t^6}{2} c(u, v) - \frac{t^{2p}}{4p} F(u, v) \right\} \\ &\leq \frac{p-1}{2p} \left( \frac{2(a(u, v) + b(u, v))^p}{F} \right)^{\frac{1}{p-1}} + \frac{p-3}{2p} \left( \frac{6c(u, v)^{\frac{p}{3}}}{F} \right)^{\frac{3}{p-3}}. \end{aligned}$$

Assume  $\omega \leq 1$ ,  $E_1 \geq E_\omega$ . Choosing  $(u_1, u_1)$  such that

$$c_N \leq \max_{t \geq 0} I(tu_1, tu_1) < E_\omega,$$

where  $u_1$  is a positive ground state solution of (3.1) with  $\omega = 1$ . Then

$$\begin{aligned} c_N &\leq \max_{t \geq 0} I(tu_1, tu_1) \\ &\leq \frac{p-1}{2p} \left( \frac{2(2\|u_1\|^2)^p}{2(1+b)\|u_1\|_{2p}^{2p}} \right)^{\frac{1}{p-1}} + \frac{p-3}{2p} \left( \frac{6(2B(u_1))^{\frac{p}{3}}}{2(1+b)\|u_1\|_{2p}^{2p}} \right)^{\frac{3}{p-3}} \\ &\leq \max\{2^{\frac{p}{p-1}}, 3^{\frac{3}{p-3}}\} \frac{p-1}{p} \|u_1\|_{2p}^{2p} \left( \frac{1}{1+b} \right)^{\frac{1}{p-1}} \\ &< \max\{2^{\frac{p}{p-1}}, 3^{\frac{3}{p-3}}\} \frac{p-1}{p} \|u_1\|_{2p}^{2p} \left( \frac{1}{b} \right)^{\frac{1}{p-1}} \end{aligned}$$



Due to

$$\max\{2^{\frac{p}{p-1}}, 3^{\frac{3}{p-3}}\} = \begin{cases} 3^{\frac{p}{p-1}}, & p \in [3 + \sqrt{6}, \infty), \\ 3^{\frac{3}{p-3}}, & p \in (3, 3 + \sqrt{6}), \end{cases}$$

we deduce that

$$b > \begin{cases} \left( \frac{(p-1)3^{\frac{p}{p-1}} \|u_1\|_{2p}^{2p}}{pE_\omega} \right)^{p-1}, & p \in [3 + \sqrt{6}, \infty), \\ \left( \frac{(p-1)3^{\frac{3}{p-3}} \|u_1\|_{2p}^{2p}}{pE_\omega} \right)^{p-1}, & p \in (3, 3 + \sqrt{6}). \end{cases}$$

In a similar fashion, if  $\omega \geq 1$ , it follows that

$$b > \begin{cases} \left( \frac{(p-1)3^{\frac{p}{p-1}} \|u_\omega\|_{2p}^{2p}}{pE_1} \right)^{p-1}, & p \in [3 + \sqrt{6}, \infty), \\ \left( \frac{(p-1)3^{\frac{3}{p-3}} \|u_\omega\|_{2p}^{2p}}{pE_1} \right)^{p-1}, & p \in (3, 3 + \sqrt{6}). \end{cases}$$

In conclusion,  $c_{\mathcal{N}} < \min\{E_1, E_\omega\}$  provided

$$b > \begin{cases} \max\{b_1, b_2\} & p \in [3 + \sqrt{6}, \infty), \\ \max\{b_3, b_4\} & p \in (3, 3 + \sqrt{6}). \end{cases}$$

This completes the proof.  $\square$

*Proof of (ii) and (iii) of Theorem 1.1.* We divide the proof in several steps.

**Step 1.** We show that the existence of ground state solutions of problem (1.1). By the Ekeland variational principle [24] that there exists a sequence  $\{(u_n, v_n)\} \in E$  such that

$$I(u_n, v_n) \rightarrow c, \quad I'(u_n, v_n) \rightarrow 0 \text{ in } E'.$$

By computing  $I(u_n, v_n) - \frac{1}{6}\langle I'(u_n, v_n), (u_n, v_n) \rangle$ , it is easy to get that  $\{(u_n, v_n)\}$  is bounded in  $E$ . Then there exists  $(u, v) \in E$  such that, up to a subsequence

$$\begin{aligned} (u_n, v_n) &\rightarrow (u, v) \text{ weakly in } E, \\ (u_n, v_n) &\rightarrow (u, v) \text{ strongly in } L^{2p}(\mathbb{R}^2) \times L^{2p}(\mathbb{R}^2) \text{ for } p \in (3, +\infty), \\ (u_n, v_n) &\rightarrow (u, v) \text{ a.e. in } \mathbb{R}^2. \end{aligned}$$

Then  $I'(u, v) = 0$  and  $I(u, v) \leq c$ . If  $(u, v) \neq (0, 0)$ , it follows that  $(u, v) \in \mathcal{N}$ ,  $I(u, v) \geq c$ . Then,  $I(u, v) = c$ . Now, it remains to prove  $(u, v) \neq (0, 0)$ . Assume by the contrary that  $(u, v) = (0, 0)$ , then

$$\begin{aligned} c + o_n(1) \|(u_n, v_n)\|_E &= I((u_n, v_n)) - \frac{1}{2}\langle I'(u_n, v_n), (u_n, v_n) \rangle \\ &= -(B(u + B(v)) + \left(\frac{1}{2} - \frac{1}{2p}\right)F(u, v) + o_n(1) \\ &= o_n(1), \end{aligned}$$

which is a contradiction. Thus  $(u, v)$  is a ground state solution of system (1.1).

**Step 2.** We show that  $u \neq 0$  and  $v \neq 0$ . Without loss of generality, we assume that  $u = 0$  and  $v \neq 0$ . Multiplying the first equation in (1.1) by  $u_n$ , we obtain

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^2} (|\nabla u_n|^2 + u_n^2) dx = 0,$$

Multiplying the second equation in (1.1) by  $v_n$ ,

$$(3.2) \quad \int_{\mathbb{R}^2} (|\nabla v_n|^2 + v_n^2) dx + 3B(v_n) = \int_{\mathbb{R}^2} v_n^{2p} dx + o_n(1).$$

Therefore, there exists  $t_n$  such that

$$(3.3) \quad \frac{1}{t_n^2} \int_{\mathbb{R}^2} (|\nabla v_n|^2 + v_n^2) dx + 3B(v_n) = t_n^{2p-6} \int_{\mathbb{R}^2} v_n^{2p} dx.$$

Combining (3.2) and (3.3), it follows that  $t_n \rightarrow 1$ , as  $n \rightarrow \infty$ . Hence

$$\lim_{n \rightarrow \infty} I(u_n, v_n) \rightarrow I(0, v) \geq E_\omega,$$

which contradicts the fact that  $c_{\mathcal{N}} < E_\omega$ . Similarly, if  $v = 0$  and  $u \neq 0$ , we can obtain  $I(u_n, v_n) \rightarrow I(u, 0)$ , as  $n \rightarrow \infty$ , which contradicts the fact that  $c_{\mathcal{N}} < E_1$ .

Therefore,  $u \neq 0$  and  $v \neq 0$ ,  $(u, v)$  is a nontrivial ground state solution of (1.1). In fact, since  $(|u|, |v|) \in \mathcal{N}$  and  $c_{\mathcal{N}} = I(|u|, |v|)$ , we conclude that  $(|u|, |v|)$  is a nonnegative solution of (1.1). Using the strong maximum principle, we infer that  $|u|, |v| > 0$ . Thus  $(|u|, |v|)$  is a positive least energy solution of (1.1). This completes the proof.  $\square$

#### 4. PROOF OF (i) OF THEOREM 1.1.

Given  $(u, v) \in E \setminus \{(0, 0)\}$ , consider the path

$$\gamma_{u,v}(t) := (t^\alpha u(t), t^\alpha v(t)), \quad t \geq 0,$$

where  $\alpha > 1$  such that  $\frac{1}{p-1} < \alpha < \frac{1}{3-p}$  for  $p \in (2, 3)$  and  $\alpha > 1$  for  $p = 3$ . Then

$$\begin{aligned} I(\gamma_{u,v}(t)) &= \frac{t^{2\alpha}}{2} (\|\nabla u\|_2^2 + \|\nabla v\|_2^2) + \frac{t^{2(\alpha-1)}}{2} (\|u\|_2^2 + \omega \|v\|_2^2) \\ &\quad + \frac{t^{6\alpha-4}}{2} (B(u) + B(v)) - \frac{t^{2p\alpha-2}}{2p} F(u, v). \end{aligned}$$

By differentiating both sides with respect to  $t$  at 1, we obtain the following constraint

$$\begin{aligned} J(u, v) &= \alpha (\|\nabla u\|_2^2 + \|\nabla v\|_2^2) + (\alpha - 1) (\|u\|_2^2 + \omega \|v\|_2^2) \\ &\quad + (3\alpha - 2) (B(u) + B(v)) - \frac{p\alpha - 1}{p} F(u, v). \end{aligned}$$

Define a constraint manifold of Pohozaev-Nehari mype

$$\mathcal{M}_b := \{(u, v) \in E \setminus \{(0, 0)\} \mid J(u, v) = 0\}.$$

The corresponding groundstate energy is described as

$$c_b := \inf_{(u,v) \in \mathcal{M}_b} I(u, v).$$

By [5], when  $p \in (2, 3]$ , problem (3.1) admits a positive ground state solution  $\tilde{u}_\omega$ . To be more precise, denote the ground state level by

$$\tilde{E}_\omega := \inf_{u \in \mathcal{M}_\omega} J_\omega(u)$$

where

$$\mathcal{M}_\omega := \left\{ u \in H_r^1(\mathbb{R}^2) \setminus \{0\} : \int_{\mathbb{R}^2} (\alpha |\nabla u|^2 + (\alpha - 1)\omega u^2) dx + (3\alpha - 2)B(u) = \frac{p\alpha - 1}{p} \int_{\mathbb{R}^2} u^{2p} dx \right\}.$$

Define the set of ground state solutions of (3.1) by

$$S_\omega := \{u \in H_r^1(\mathbb{R}^2) \setminus \{0\} : J'_\omega(u) = 0, J_\omega(u) = E_\omega\}.$$

$S_\omega$  is nonempty.

**Lemma 4.1.** *The set  $S_\omega$  is compact in  $H_r^1(\mathbb{R}^2)$ . More precisely, any sequence  $\{u_n\} \subset S_\omega$  has a subsequence  $\{u_{j_n}\} \subset S_\omega$  and  $u \in S_\omega$  such that  $u_{j_n} \rightarrow u$  strongly in  $H_r^1(\mathbb{R}^2)$  as  $n \rightarrow \infty$ .*

*Proof.* For any  $\{u_n\} \in S_\omega$ ,

$$\begin{aligned} E_\omega &= \frac{1}{2} \int_{\mathbb{R}^2} (|\nabla u_n|^2 + \omega u_n^2) dx + \frac{1}{2} B(u_n) - \frac{1}{2p} \int_{\mathbb{R}^2} u_n^{2p} dx \\ &= \left( \frac{1}{2} - \frac{\alpha}{2(p\alpha - 1)} \right) \int_{\mathbb{R}^2} |\nabla u_n|^2 dx + \left( \frac{1}{2} - \frac{\alpha - 1}{2(p\alpha - 1)} \right) \omega \int_{\mathbb{R}^2} u_n^2 dx + \left( \frac{1}{2} - \frac{3\alpha - 2}{2(p\alpha - 1)} \right) B(u_n) \\ &\geq \left( \frac{1}{2} - \frac{\alpha}{2(p\alpha - 1)} \right) \int_{\mathbb{R}^2} (|\nabla u_n|^2 + u_n^2) dx. \end{aligned}$$

Therefore  $\{u_n\}$  is bounded on  $S_\omega$ . There exists  $u \in H_r^1(\mathbb{R}^2)$ , such that, up to a subsequence,  $u_n \rightarrow u$  weakly in  $H_r^1(\mathbb{R}^2)$ , strongly in  $L^p(\mathbb{R}^2)$  for  $p \in (2, 3]$  as  $n \rightarrow \infty$ . Moreover, due to  $u_n \in \mathcal{M}_\omega$  for any  $n$ , we can obtain

$$(4.1) \quad \liminf_{n \rightarrow \infty} \|u_n\|_{H_r^1(\mathbb{R}^2)} > 0.$$

First we prove that  $u \neq 0$ . Indeed, if  $u = 0$ , by Lemma 2.1, we obtain  $u_n \rightarrow 0$  strongly in  $H_r^1(\mathbb{R}^2)$ , as  $n \rightarrow \infty$ , which contradicts (4.1). Thus  $u \neq 0$ . Next, we show that  $u \in S_\omega$ . Since  $u_n$  satisfies problem (3.1), we get that  $u$  is a solution of problem (3.1). By the semicontinuity of the norms,

$$\tilde{E}_\omega \leq J_\omega(u) \leq \liminf_{n \rightarrow \infty} J_\omega(u_n) = \tilde{E}_\omega.$$

Therefore  $u \in S_\omega$  and  $u_n \rightarrow u$  strongly in  $H_r^1(\mathbb{R}^2)$ , as  $n \rightarrow \infty$ . That is,  $S_\omega$  is compact in  $H_r^1(\mathbb{R}^2)$ .  $\square$

**Lemma 4.2.** *For given positive constants  $a, b, c, d$ , a function  $f(t) = at^{2\alpha} + bt^{2(\alpha-1)} + ct^{6\alpha-4} - dt^{2p\alpha-2}$  has exactly one critical point on  $(0, +\infty)$ , the maximum.*

*Proof.* The proof is similar to [5], we omit it here.  $\square$

**Lemma 4.3.** *For any  $(u, v) \in E \setminus \{(0, 0)\}$ , there exists a unique  $t_{u,v} > 0$  such that  $\gamma_{u,v}(t_{u,v}) \in \mathcal{M}_b$  and*

$$c_b = \inf_{(u,v) \in E \setminus \{(0,0)\}} \max_{t>0} I(\gamma_{u,v}(t)).$$

*Proof.* The proof is standard, we omit it here.  $\square$

**Lemma 4.4.** *If  $b > 0$ , then  $c_b < \tilde{E}_1 + \tilde{E}_\omega$ , where  $\tilde{E}_1$  is a groundstate energy of problem (3.1) with  $\omega = 1$ .*

*Proof.* Let  $\tilde{u}_1$  and  $\tilde{u}_\omega$  be a positive ground state solution associated to the level  $\tilde{E}_1$  and  $\tilde{E}_\omega$  respectively. Denote  $f(t) = I(t^\alpha \tilde{u}_1(t \cdot), t^\alpha \tilde{u}_\omega(t \cdot))$ , that is,

$$\begin{aligned} f(t) = & \frac{t^{2\alpha}}{2} (\|\nabla \tilde{u}_1\|_2^2 + \|\nabla \tilde{u}_\omega\|_2^2) + \frac{t^{2(\alpha-1)}}{2} (\|\tilde{u}_1\|_2^2 + \omega \|\tilde{u}_\omega\|_2^2) \\ & + \frac{t^{6\alpha-4}}{2} (B(\tilde{u}_1) + B(\tilde{u}_\omega)) - \frac{t^{2p\alpha-2}}{2p} F(\tilde{u}_1, \tilde{u}_\omega). \end{aligned}$$

According to Lemma 4.2, there exists unique  $t_0 > 0$  such that  $(t_0^\alpha \tilde{u}_1(t_0 \cdot), t_0^\alpha \tilde{u}_\omega(t_0 \cdot)) \in \mathcal{M}_b$ . Hence

$$\begin{aligned} c_b & \leq I(t_0^\alpha \tilde{u}_1(t_0 \cdot), t_0^\alpha \tilde{u}_\omega(t_0 \cdot)) = I(t_0 \tilde{u}_1(t_0 \cdot), 0) + I(0, t_0 \tilde{u}_\omega(t_0 \cdot)) - 2bt_0^{2p} \int_{\mathbb{R}^2} |\tilde{u}_1(t_0 x) \tilde{u}_\omega(t_0 x)|^p dx \\ & < I(t_0 \tilde{u}_1(t_0 \cdot), 0) + I(0, t_0 \tilde{u}_\omega(t_0 \cdot)) \\ & = \tilde{E}_1 + \tilde{E}_\omega. \end{aligned}$$

This completes the proof.  $\square$

**Lemma 4.5.**  $\liminf_{b \rightarrow 0} c_b > 0$ .

*Proof.* Suppose by contradiction the lemma does not hold. Then there exists  $\{b_k\}$  such that  $b_k \rightarrow 0$  and  $c_{b_k} \rightarrow 0$ , as  $k \rightarrow \infty$ . Moreover, there exists  $\{(u_k, v_k)\} \in \mathcal{M}_{b_k}$  such that  $I(u_k, v_k) \rightarrow 0$ , as  $k \rightarrow \infty$ , that is

$$\begin{aligned} o_k(1) & = \frac{1}{2} \int_{\mathbb{R}^2} (|\nabla u_k|^2 + u_k^2 + |\nabla v_k|^2 + \omega v_k^2) dx + \frac{1}{2} (B(u_k) + B(v_k)) - \frac{1}{2p} F(u_k, v_k) \\ & \geq \left( \frac{1}{2} - \frac{\alpha}{2(p\alpha - 1)} \right) \|(u_k, v_k)\|_E^2. \end{aligned}$$

We deduce that  $\|(u_k, v_k)\|_E \rightarrow 0$ , as  $k \rightarrow \infty$ . Since  $\{(u_k, v_k)\} \in \mathcal{M}_{b_k}$ ,

$$(\alpha - 1) \|(u_k, v_k)\|_E^2 \leq C \|(u_k, v_k)\|_E^{2p}.$$

Thus we have  $\|(u_k, v_k)\|_E \geq \left( \frac{\alpha-1}{C} \right)^{\frac{1}{2p-2}}$ , which is a contradiction. This completes the proof.  $\square$

Given  $\delta > 0$ , let

$$(S_\omega)^\delta := \{u \in H_r^1(\mathbb{R}^2) | u = \tilde{u} + \bar{u}, \tilde{u} \in S_\omega, \|\bar{u}\|_{H_r^1(\mathbb{R}^2)} \leq \delta\},$$

be the neighborhood of  $S_\omega$  of radius  $\delta$ .

**Lemma 4.6.** *For any  $\delta > 0$ , there exists  $b_\delta > 0$  such that for any  $b \in (0, b_\delta)$ , up to a subsequence, there exists  $(u_n^b, v_n^b) \in \mathcal{M}_b$  satisfying*

$$(4.2) \quad I(u_n^b, v_n^b) \rightarrow c_b, \quad I'(u_n^b, v_n^b) \rightarrow 0, \text{ as } n \rightarrow \infty,$$

and  $\{u_n^b\} \in (S_1)^\delta$  and  $\{v_n^b\} \in (S_\omega)^\delta$ .

*Proof.* We adopt some idea in [7]. Suppose by contradiction the lemma does not hold. Then for  $\delta_0 > 0$ , there exists  $\{b_k\} \in \mathbb{R}^+$  such that  $b_k \rightarrow 0$ , as  $k \rightarrow \infty$ , and for any  $\{(u_n^{b_k}, v_n^{b_k})\} \in \mathcal{M}_{b_k}$  satisfying (4.2), there holds  $\{u_n^{b_k}\} \in H_r^1(\mathbb{R}^2) \setminus (S_1)^{\delta_0}$  or  $\{v_n^{b_k}\} \in H_r^1(\mathbb{R}^2) \setminus (S_\omega)^{\delta_0}$ . For any  $k$ , there exists  $n_k$  such that

$$|I(u_{n_k}^{b_k}, v_{n_k}^{b_k}) - c_{b_k}| \leq 1/k.$$

Let  $\tilde{u}_k = u_{n_k}^{b_k}$  and  $\tilde{v}_k = v_{n_k}^{b_k}$ . By Lemma 4.4, we have

$$(4.3) \quad \limsup_{k \rightarrow \infty} I(\tilde{u}_k, \tilde{v}_k) \leq \limsup_{k \rightarrow \infty} c_{b_k} \leq \tilde{E}_1 + \tilde{E}_\omega.$$

Since  $\{(\tilde{u}_k, \tilde{v}_k)\} \in \mathcal{M}_{b_k}$ ,

$$\begin{aligned} c_{b_k} &= \frac{1}{2} \|(\tilde{u}_k, \tilde{v}_k)\|_E^2 + \frac{1}{2} (B(\tilde{u}_k) + B(\tilde{v}_k)) - \frac{1}{2p} F(\tilde{u}_k, \tilde{v}_k) \\ &\geq \left( \frac{1}{2} - \frac{\alpha}{2(p\alpha - 1)} \right) \|(\tilde{u}_k, \tilde{v}_k)\|_E^2, \end{aligned}$$

we deduce that  $\{(\tilde{u}_k, \tilde{v}_k)\}$  is bounded in  $E$ . Up to a subsequence,  $\tilde{u}_k \rightarrow u$  and  $\tilde{v}_k \rightarrow v$  weakly in  $H_r^1(\mathbb{R}^2)$ , strongly in  $L^{2p}(\mathbb{R}^2)$  for  $p \in (2, 3]$ , as  $k \rightarrow \infty$ . By Lemma 4.5, we have

$$\liminf_{k \rightarrow \infty} \min\{\|\tilde{u}_k\|_{H_r^1(\mathbb{R}^2)}, \|\tilde{v}_k\|_{H_r^1(\mathbb{R}^2)}\} > 0.$$

Noting that  $b_k > 0$  and

$$o_k(1) = \int_{\mathbb{R}^2} |\nabla \tilde{u}_k|^2 + \tilde{u}_k^2 + 3B(\tilde{u}_k) - \int_{\mathbb{R}^2} \tilde{u}_k^{2p},$$

there exists  $t_k$  such that

$$\frac{1}{t_k^{4\alpha-4}} \int_{\mathbb{R}^2} |\nabla \tilde{u}_k|^2 + \frac{1}{t_k^{4\alpha-2}} \int_{\mathbb{R}^2} \tilde{u}_k^2 + 3B(\tilde{u}_k) = t_k^{2+2p\alpha-6\alpha} \int_{\mathbb{R}^2} \tilde{u}_k^{2p},$$

that is  $t_k^\alpha \tilde{u}_k(t_k \cdot) \in \mathcal{N}_1$ . Similarly, there exists  $s_k$  such that  $s_k^\alpha \tilde{v}_k(t_k \cdot) \in \mathcal{N}_\omega$ .

**Step 1.** We claim that  $t_k \rightarrow 1$  and  $s_k \rightarrow 1$  as  $k \rightarrow \infty$ . We only give the proof of  $t_k \rightarrow 1$ , as the second convergence being similar. We consider two cases:

**Case I.**  $u \neq 0$ . If  $\limsup_{k \rightarrow \infty} t_k > 1$ , then we can assume that  $t_k > 1$  for all  $k$  we have

$$\begin{aligned} o_k(1) &= (t_k^{2+2p\alpha-6\alpha} - 1) \int_{\mathbb{R}^2} \tilde{u}_k^{2p} - \left( \frac{1}{t_k^{4\alpha-4}} - 1 \right) \int_{\mathbb{R}^2} |\nabla \tilde{u}_k|^2 - \left( \frac{1}{t_k^{4\alpha-2}} - 1 \right) \int_{\mathbb{R}^2} \tilde{u}_k^2 \\ &\geq (t_k^{2+2p\alpha-6\alpha} - 1) \int_{\mathbb{R}^2} \tilde{u}_k^{2p}, \end{aligned}$$

which yields  $t_k \rightarrow 1$  as  $k \rightarrow \infty$ . This is a contradiction. So  $\limsup_{k \rightarrow \infty} t_k \leq 1$ . Similarly,  $\liminf_{k \rightarrow \infty} t_k \geq 1$ . Then  $\lim_{k \rightarrow \infty} t_k = 1$ .

**Case II.**  $u = 0$ . If  $\limsup_{k \rightarrow \infty} t_k > 1$ , then we can assume that  $t_k > 1$  for all  $k$  we have  $\limsup_{k \rightarrow \infty} \|\tilde{u}_k\|_{H_r^1(\mathbb{R}^2)} = 0$ , which contradicts (4.3). So  $\limsup_{k \rightarrow \infty} t_k \leq 1$ . Similarly,  $\liminf_{k \rightarrow \infty} t_k \geq 1$ . Then  $\lim_{k \rightarrow \infty} t_k = 1$ .

**Step 2.** Let  $\bar{u}_k = t_k^\alpha \tilde{u}_k(t_k \cdot)$  and  $\bar{v}_k = s_k^\alpha \tilde{v}_k(t_k \cdot)$ , then  $\bar{u}_k \rightarrow u$  and  $\bar{v}_k \rightarrow v$  weakly in  $H_r^1(\mathbb{R}^2)$ , as  $k \rightarrow \infty$ . Next we show  $u \in S_1$ ,  $v \in S_\omega$  and  $\bar{u}_k \rightarrow u$ ,  $\bar{v}_k \rightarrow v$  in  $H_r^1(\mathbb{R}^2)$ , as  $k \rightarrow \infty$ . This will be a contradiction.

Since  $\tilde{u}_k \in H_r^1(\mathbb{R}^2)$ , there exist  $U_k \in C_0(\mathbb{R}^2)$  and  $V_k \in C_0(\mathbb{R}^2)$  such that

$$\int_{\mathbb{R}^2} |\nabla \tilde{u}_k - U_k|^2 dx < \varepsilon, \quad \int_{\mathbb{R}^2} |\tilde{u}_k - V_k|^2 dx < \varepsilon.$$

Therefore

$$\begin{aligned} & \|\nabla(\bar{u}_k - \tilde{u}_k)\|_2^2 \\ &= \int_{\mathbb{R}^2} |\nabla(t_k^\alpha \tilde{u}_k(t_k x) - \tilde{u}_k(x))|^2 dx \\ &\leq 2 \int_{\mathbb{R}^2} |\nabla(t_k^\alpha \tilde{u}_k(t_k x)) - U_k(x)|^2 dx + 2 \int_{\mathbb{R}^2} |\nabla \tilde{u}_k(x) - U_k(x)|^2 dx \\ &= 2 \int_{\mathbb{R}^2} |t_k^{\alpha+1} \nabla(\tilde{u}_k(t_k x)) - U_k(x)|^2 dx + 2 \int_{\mathbb{R}^2} |\nabla \tilde{u}_k(x) - U_k(x)|^2 dx \\ &\leq 4t_k^{2\alpha+2} \int_{\mathbb{R}^2} |U_k(t_k x) - U_k(x)|^2 dx + 2|t_k^{\alpha+1} - 1|^2 \int_{\mathbb{R}^2} U_k^2(x) dx + (4t_k^{2\alpha} + 2)\varepsilon \\ &= 12\varepsilon, \end{aligned}$$

and

$$\begin{aligned} & \|\bar{u}_k - \tilde{u}_k\|_2^2 \\ &= \int_{\mathbb{R}^2} |t_k^\alpha \tilde{u}_k(t_k x) - \tilde{u}_k(x)|^2 dx \\ &\leq 2 \int_{\mathbb{R}^2} |t_k^\alpha \tilde{u}_k(t_k x) - V_k(x)|^2 dx + 2 \int_{\mathbb{R}^2} |\tilde{u}_k(x) - V_k(x)|^2 dx \\ &\leq 4t_k^{2\alpha} \int_{\mathbb{R}^2} |V_k(t_k x) - V_k(x)|^2 dx + 2|t_k^\alpha - 1|^2 \int_{\mathbb{R}^2} V_k^2(x) dx + (4t_k^{2\alpha-2} + 2)\varepsilon \\ &= 12\varepsilon. \end{aligned}$$

It follows that  $\|\bar{u}_k - \tilde{u}_k\|_{H_r^1(\mathbb{R}^2)} \rightarrow 0$  and  $\|\bar{v}_k - \tilde{v}_k\|_{H_r^1(\mathbb{R}^2)} \rightarrow 0$  as  $k \rightarrow \infty$ . So

$$I(\tilde{u}_k, \tilde{v}_k) = I(\bar{u}_k, \bar{v}_k) + o_k(1) \geq \tilde{E}_1 + \tilde{E}_\omega + o_k(1).$$

Recalling that  $\limsup_{k \rightarrow \infty} I(\tilde{u}_k, \tilde{v}_k) \leq \tilde{E}_1 + \tilde{E}_\omega$ , we obtain

$$\lim_{k \rightarrow \infty} J_1(\bar{u}_k) = \tilde{E}_1, \quad \lim_{k \rightarrow \infty} J_\omega \bar{v}_k = \tilde{E}_\omega.$$

Arguing as in the proof of Lemma 4.1, we deduce that  $u \neq 0$ . Thanks to the lower semicontinuity of norms,

$$J_1(u) \leq \liminf_{k \rightarrow \infty} J_1(\bar{u}_k) = B_1$$

If  $J_1(u) = E_1$ , which yields that  $\bar{u}_k \rightarrow u$  strongly in  $H_r^1(\mathbb{R}^2)$  and  $u \in S_1$ . If not, we have

$$\|u\|_{H_r^1(\mathbb{R}^2)} < \liminf_{k \rightarrow \infty} \|\bar{u}_k\|_{H_r^1(\mathbb{R}^2)}.$$

It follows that  $u \notin \mathcal{M}_1$ . Then there exists a unique  $t_0 \in (0, 1)$  such that  $J(t_0^\alpha u(t_0 \cdot)) = 0$ . Thus, we have

$$J_1(t_0^\alpha u(t_0 \cdot)) < \lim_{n \rightarrow \infty} \left( \frac{t_0^{2\alpha}}{2} \|\nabla \bar{u}_k\|_2^2 + \frac{t_0^{2(\alpha-1)}}{2} \|\bar{u}_k\|_2^2 + \frac{t_0^{6\alpha-4}}{2} B(\bar{u}_k) - \frac{t_0^{2p\alpha-2}}{2p} \|\bar{u}_k\|_{2p}^{2p} \right).$$

Since  $J_1(t^\alpha \bar{u}_k(t))$  has the maximum value at  $t = 1$  for all  $k$ . It follows that

$$\begin{aligned} J_1(t_0^\alpha u(t_0)) &< \lim_{n \rightarrow \infty} \left( \frac{1}{2} \|\nabla \bar{u}_k\|_2^2 + \frac{1}{2} \|\bar{u}_k\|_2^2 + \frac{1}{2} B(\bar{u}_k) - \frac{1}{2p} \|\bar{u}_k\|_{2p}^{2p} \right) \\ &= \tilde{E}_1, \end{aligned}$$

which is a contradiction. That is  $u \in \mathcal{M}_1$  and  $J_1(u) = \tilde{E}_1$ , which yields  $u \in S_1$ , and  $\bar{u}_k \rightarrow u$  strongly in  $H_r^1(\mathbb{R}^2)$  as  $k \rightarrow \infty$ . Finally, we can similarly prove  $v \in S_\omega$  and  $\bar{v}_k \rightarrow v$  strongly in  $H_r^1(\mathbb{R}^2)$  as  $k \rightarrow \infty$ . By Step 1, we know that  $\tilde{u}_k \rightarrow u$  and  $\tilde{v}_k \rightarrow v$  strongly in  $H_r^1(\mathbb{R}^2)$ , as  $k \rightarrow \infty$ . This is a contradiction with the fact that  $\tilde{u}_k \in H_r^1(\mathbb{R}^2) \setminus (S_1)^{\delta_0}$  or  $\tilde{v}_k \in H_r^1(\mathbb{R}^2) \setminus (S_\omega)^{\delta_0}$ . This completes the proof.  $\square$

*Proof of (i) of Theorem 1.1.* We divide the proof into several steps.

**Step 1.**  $\mathcal{M}_b$  is nonempty. For each  $(u, v) \in E \setminus \{(0, 0)\}$ ,  $J(t^\alpha u(t), t^\alpha v(t))$  is of the form  $at^{2\alpha} + bt^{2(\alpha-1)} + ct^{6\alpha-4} - dt^{2p\alpha-2}$ , which is positive for small  $t$  and negative for large  $t$ . Thus, there exists  $\tilde{t}_{uv} > 0$  such that  $J(\tilde{t}_{uv}^\alpha u(\tilde{t}_{uv} \cdot), \tilde{t}_{uv}^\alpha v(\tilde{t}_{uv} \cdot)) = 0$ . Thus,  $\mathcal{M}_b$  is not empty.

**Step 2.**  $\mathcal{M}_b$  is bounded away from zero, i.e.  $(0, 0) \notin \partial \mathcal{M}_b$ . For each  $(u, v) \in \mathcal{M}_b$ ,

$$\begin{aligned} (4.4) \quad F(u, v) &= \frac{p}{p\alpha - 1} \left( \alpha a(u, v) + (\alpha - 1)b(u, v) + (3\alpha - 2)c(u, v) \right) \\ &\geq \frac{p}{p\alpha - 1} (\alpha - 1) \|(u, v)\|_E^2. \end{aligned}$$

By the Sobolev embedding theorem there exists a constant  $C > 0$  such that for any  $(u, v) \in \mathcal{M}_b$ ,  $\|(u, v)\|_E^{2p} \geq C \|(u, v)\|_E^2$ . Therefore,  $\|(u, v)\|_E \geq \rho > 0$  and the conclusion holds.

**Step 3.**  $c_b > 0$ . For each  $(u, v) \in \mathcal{M}_b$ , combining (4.4)

$$\begin{aligned} I(u, v) &= \frac{1}{2} \|(u, v)\|_E^2 + \frac{1}{2} (B(u) + B(v)) - \frac{1}{2p} F(u, v) \\ &= \frac{1}{2} \|(u, v)\|_E^2 + \frac{1}{2} (B(u) + B(v)) - \frac{1}{2(p\alpha - 1)} \left( \alpha a(u, v) + (\alpha - 1)b(u, v) + (3\alpha - 2)c(u, v) \right) \\ &= \left( \frac{1}{2} - \frac{\alpha}{2(p\alpha - 1)} \right) a(u, v) + \left( \frac{1}{2} - \frac{\alpha - 1}{2(p\alpha - 1)} \right) b(u, v) + \left( \frac{1}{2} - \frac{3\alpha - 2}{2(p\alpha - 1)} \right) c(u, v) \\ &\geq \left( \frac{1}{2} - \frac{\alpha}{2(p\alpha - 1)} \right) \|(u, v)\|_E^2. \end{aligned}$$

Then taking into account Step 2 and  $p\alpha - 1 > \alpha$ , we can obtain  $c_b > 0$ .

**Step 4:** If  $\{(u_n, v_n)\}$  is a minimizing sequence for  $I$  on  $\mathcal{M}_b$ , then it is bounded. Let  $\{(u_n, v_n)\} \in \mathcal{M}_b$  such that  $I(u_n, v_n) \rightarrow c_b$ . As in Step 3, we get

$$I(u_n, v_n) = \left( \frac{1}{2} - \frac{\alpha}{2(p\alpha - 1)} \right) a(u_n, v_n) + \left( \frac{1}{2} - \frac{\alpha - 1}{2(p\alpha - 1)} \right) b(u_n, v_n) + \left( \frac{1}{2} - \frac{3\alpha - 2}{2(p\alpha - 1)} \right) c(u_n, v_n).$$

Since the coefficients of  $a(u_n, v_n)$ ,  $b(u_n, v_n)$  and  $c(u_n, v_n)$  are positive, then

$$I(u_n, v_n) \geq \left( \frac{1}{2} - \frac{\alpha}{2(p\alpha - 1)} \right) \|(u_n, v_n)\|_E^2,$$

it follows that  $\{(u_n, v_n)\}$  is bounded in  $E$ . Thus, there exists  $(u, v) \in E$  such that, up to a subsequence

$$\begin{aligned} (u_n, v_n) &\rightarrow (u, v) \text{ weakly in } E, \\ (u_n, v_n) &\rightarrow (u, v) \text{ strongly in } L^{2p}(\mathbb{R}^2) \times L^{2p}(\mathbb{R}^2) \text{ for } p \in (2, 3], \\ (u_n, v_n) &\rightarrow (u, v) \text{ a.e. in } \mathbb{R}^2. \end{aligned}$$

If  $a(u, v) + b(u, v) = \liminf_{n \rightarrow \infty} a(u_n, v_n) + b(u_n, v_n)$ , then it follows  $(u_n, v_n) \rightarrow (u, v)$  strongly in  $E$  as  $n \rightarrow \infty$  and  $(u, v) \neq (0, 0)$ , then  $c_b$  is attained by  $(u, v)$ .

If  $a(u, v) + b(u, v) < \liminf_{n \rightarrow \infty} a(u_n, v_n) + b(u_n, v_n)$ , by the Lemma 2.1 and  $J(u_n, v_n) = 0$ , we deduce that  $J(u, v) < 0$ , then it follows that  $(u, v) \notin \mathcal{M}_b$  and  $(u, v) \neq (0, 0)$ . Then there exists a unique  $t_0 \in (0, 1)$  such that  $J(t_0^\alpha u(t_0 \cdot), t_0^\alpha v(t_0 \cdot)) = 0$ . Thus, we have

$$I(t_0^\alpha u(t_0 \cdot), t_0^\alpha v(t_0 \cdot)) < \lim_{n \rightarrow \infty} \left( \frac{t_0^{2\alpha}}{2} a(u_n, v_n) + \frac{t_0^{2(\alpha-1)}}{2} b(u_n, v_n) + \frac{t_0^{6\alpha-4}}{2} c(u_n, v_n) - \frac{t_0^{2p\alpha-2}}{2p} F(u_n, v_n) \right).$$

Since  $J(t^\alpha u_n(t \cdot), t^\alpha v_n(t \cdot))$  has the maximum value at  $t = 1$  for all  $n$ . It follows that

$$\begin{aligned} I(t_0^\alpha u(t_0 \cdot), t_0^\alpha v(t_0 \cdot)) &< \lim_{n \rightarrow \infty} \left( \frac{1}{2} a(u_n, v_n) + \frac{1}{2} b(u_n, v_n) + \frac{1}{2} c(u_n, v_n) - \frac{1}{2p} F(u_n, v_n) \right) \\ &= c_b, \end{aligned}$$

which is a contradiction.

**Step 5.** The minimizer  $(u, v)$  is a regular point of  $\mathcal{M}_b$ , i.e.  $J'(u, v) \neq 0$ . To the contrary, suppose that  $J'(u, v) = 0$ . For  $(u_t, v_t) = (t^\alpha u(tx), t^\alpha v(tx))$ , one has

$$\begin{aligned} J(u_t, v_t) &= t \frac{d}{dt} I(u_t, v_t), \\ \frac{d}{dt} J(u_t, v_t) &= \frac{d}{dt} I(u_t, v_t) + t \cdot \frac{d^2}{dt^2} I(u_t, v_t). \end{aligned}$$

Since  $\frac{d}{dt} \Big|_{t=1} J(u_t, v_t) = 0$ , it follows that

$$2\alpha^2 a(u, v) + 2(\alpha - 1)^2 b(u, v) + 2(3\alpha - 2)^2 c(u, v) - \frac{2(p\alpha - 1)^2}{p} F(u, v) = 0.$$

Then, combining with  $J(u, v) = 0$ , we get

$$\begin{aligned} 0 &= (\alpha^2 - \alpha(p\alpha - 1))a(u, v) + ((\alpha - 1)^2 - (\alpha - 1)(p\alpha - 1))b(u, v) \\ &\quad + ((3\alpha - 2)^2 - (3\alpha - 2)(p\alpha - 1))c(u, v) \end{aligned}$$

The coefficients of  $a(u, v)$ ,  $b(u, v)$  and  $c(u, v)$  in the above identity are negative, which is a contradiction.

**Step 6.**  $I'(u, v) = 0$ . Thanks to Lagrange multiplier rule, there exists  $\mu \in \mathbb{R}$  such that

$$(4.5) \quad I'(u, v) = \mu J'(u, v).$$

We claim  $\mu = 0$ . There holds

$$\begin{cases} \alpha a + (\alpha - 1)b + (3\alpha - 2)c - \frac{p\alpha - 1}{p}d = 0; \\ (1 - 2\alpha\mu)a + (1 - 2\mu(\alpha - 1))b + 3(1 - \mu(6\alpha - 4))c - (1 - \mu(2p\alpha - 2))d = 0; \\ (2\mu(\alpha - 1) - 1)b + 2(\mu(6\alpha - 4) - 1)c - \frac{\mu(2p\alpha - 2) - 1}{p}d = 0. \end{cases}$$



The first equation holds since  $J(u, v) = 0$ . The second one follows by multiplying (4.5) by  $(u, v)$  and integrating. The third one comes from Pohozaev equality. We get

$$0 = \mu \left( ((2\alpha^2 - (2p\alpha - 2)\alpha)a(u, v) + (2(\alpha - 1)^2 - 2(p\alpha - 1)(\alpha - 1))b(u, v) + (2(3\alpha - 2)^2 - 2(p\alpha - 1)(3\alpha - 2))c(u, v) \right).$$

All coefficients of  $a(u, v)$ ,  $b(u, v)$ ,  $c(u, v)$  in the above identity are negative. This implies that  $\mu = 0$ .

**Step 7.** Thanks to Lemma 4.6, the minimization  $\{(u_n, v_n)\}$  can be chosen in  $(S_1)^\delta \times (S_\omega)^\delta$ , where  $\delta > 0$  is small such that  $0 \notin S_1$  and  $0 \notin S_\omega$ . Hence,  $u \not\equiv 0$  and  $v \not\equiv 0$ ,  $(u, v)$  is a nontrivial ground state solution of (1.1). In fact, since  $(|u|, |v|) \in \mathcal{N}$  and  $c_b = I(|u|, |v|)$ , we conclude that  $(|u|, |v|)$  is a nonnegative solution of (1.1). Using the strong maximum principle, we infer that  $|u|, |v| > 0$ . Thus  $(|u|, |v|)$  is a positive ground state solution of (1.1). This completes the proof.  $\square$

## REFERENCES

- [1] A. Ambrosetti, and E. Colorado, Bound and ground states of coupled nonlinear Schrödinger equations, *C.R. Math.*, **342**(2006), 453-458.
- [2] A. Ambrosetti, and E. Colorado, Standing waves of some coupled nonlinear Schrödinger equations, *J. London Math. Soc.*, **75**(2007), 67-82.
- [3] A. Azzollini and A. Pomponio, Positive energy static solutions for the Chern-Simons-Schrödinger system under a large-distance fall-off requirement on the gauge potentials, *Calc. Var. Partial Differ. Equ.*, **60**(2021), 1-30.
- [4] T. Bartsch, and Z.-Q. Wang, Note on ground states of nonlinear Schrödinger systems, *J. Partial Differ. Equ.*, **19**(2006), 200-207.
- [5] J. Byeon, H. Huh, and J. Seok, Standing waves of nonlinear Schrödinger equations with the gauge field, *J. Funct. Anal.*, **263**(2012), 1575-1608.
- [6] J. Byeon, H. Huh, J. Seok, On standing waves with a vortex point of order  $N$  for the nonlinear Chern-Simons-Schrödinger equations, *J. Differ. Equat.*, **261**(2016) 1285-1316.
- [7] D. Cassani, H. Tavares and J.-J. Zhang, Bose fluids and positive solutions to weakly coupled systems with critical growth in dimension two, *J. Differ. Equat.*, **269**(2020) 2328-2385.
- [8] G. V. Dunne, Self-Dual Chern-Simons Theories, Springer, 1995.
- [9] D. Gilbarg and N. Trudinger, Elliptic Partial Differential Equations of Second Order, second edition, Grundlehren Math. Wiss., vol. 224, Springer, Berlin, 1983.
- [10] C. R. Hagen, A new gauge theory without an elementary photon, *Ann. Phys.*, **157**(1984), 342-359.
- [11] C. R. Hagen, Rotational anomalies without anyons, *Phys. Rev.*, **31**(1985), 2135-2136.
- [12] H. Huh, Standing waves of the Schrödinger equation coupled with the Chern-Simons gauge field, *J. Math. Phys.*, **53**(2012).
- [13] R. Jackiw and S.-Y. Pi, Classical and quantal nonrelativistic Chern-Simons theory, *Phys. Rev. D.*, **42**(1990), 3500-3513.
- [14] R. Jackiw and S.-Y. Pi, Soliton solutions to the gauged nonlinear Schrödinger equations on the plane, *Phys. Rev. Lett.*, **64**(1990), 2969-2972.
- [15] C. Ji, and F. Fang, Standing waves for the Chern-Simons-Schrödinger equation with critical exponential growth, *J. Math. Anal. Appl.*, **450**(2017), 578-591.
- [16] T.-C. Lin, and J.-C. Wei, Ground state of  $N$  coupled nonlinear Schrödinger equations in  $\mathbb{R}^n$ ,  $n \leq 3$ , *Comm. Math. Phys.*, **255**(2005), 629-653.
- [17] Z.-S. Liu, Z. Ouyang, and J.-J. Zhang, Existence and multiplicity of sign-changing standing waves for a gauged nonlinear Schrödinger equation in  $\mathbb{R}^2$ , *Nonlinearity*, **32**(2019), 3082-3111.
- [18] L. A. Maia, E. Montefusco, and B. Pellacci, Positive solutions for a weakly coupled nonlinear Schrödinger system, *J. Differ. Equat.*, **229**(2006).

- [19] A. Pomponio, and D. Ruiz, A variational analysis of a gauged nonlinear Schrödinger equation, *J. Eur. Math. Soc.*, **17**(2015), 1463-1486.
- [20] A. Pomponio, and D. Ruiz, Boundary concentration of a gauged nonlinear Schrödinger equation on large balls, *Calc. Var. Partial Differ. Equ.*, **53**(2015) 289-316.
- [21] P. H. Rabinowitz, On a class of nonlinear Schrödinger equations, *Z. Angew. Math. Phys.*, **43**, 270-291(1992).
- [22] D. Ruiz, The Schrödinger-Poisson equation under the effect of a nonlinear local term, *J. Funct. Anal.*, **237**(2006), 655-674.
- [23] B. Sirakov, Least energy solitary waves for a system of nonlinear Schrödinger equations in  $\mathbb{R}^n$ , *Comm. Math. Phys.*, **271**(2007), 199-221.
- [24] M. Willem, Minimax Theorems, Birkhäuser, Boston, 1996.
- [25] J. Zhang, W. Zhang, and X. Xie, Infinitely many solutions for a gauged nonlinear Schrödinger equation, *Appl. Math. Lett.*, **88**(2019), 21-27.

(Y. H. Jiang)

SCHOOL OF MATHEMATICS

CHINA UNIVERSITY OF MINING AND TECHNOLOGY

XUZHOU, 221116, CHINA

Email address: [18843111149@163.com](mailto:18843111149@163.com)

(T. Y. Chen)

SCHOOL OF MATHEMATICS

CHINA UNIVERSITY OF MINING AND TECHNOLOGY

XUZHOU, 221116, CHINA

Email address: [taiyongchencumt@163.com](mailto:taiyongchencumt@163.com)

(J. J. Zhang)

COLLEGE OF MATHEMATICS AND STATISTICS

CHONGQING JIAOTONG UNIVERSITY

CHONGQING 400074, CHINA

Email address: [zhangjianjun09@tsinghua.org.cn](mailto:zhangjianjun09@tsinghua.org.cn)

(M. Squassina)

COLLEGE OF SCIENCE

PRINCESS NOURAH BINT ABDUL RAHMAN UNIVERSITY

SAUDI ARABIA, RIYADH, PO BOX 84428

Email address: [marsquassina@pnu.edu.sa](mailto:marsquassina@pnu.edu.sa)

(N. Almousa)

COLLEGE OF SCIENCE

PRINCESS NOURAH BINT ABDUL RAHMAN UNIVERSITY

SAUDI ARABIA, RIYADH, PO BOX 84428

Email address: [nmalmousa@pnu.edu.sa](mailto:nmalmousa@pnu.edu.sa)

(M. Squassina)

DIPARTIMENTO DI MATEMATICA E FISICA

UNIVERSITÀ CATTOLICA DEL SACRO CUORE

VIA DEI MUSEI 41, BRESCIA, ITALY

Email address: [marco.squassina@unicatt.it](mailto:marco.squassina@unicatt.it)