# Baumslag-Solitar graphs through colours

Katie Buchhorn

June 3, 2022

Supervised by Sam Webster and Nathan Brownlowe

#### Abstract

Higher-rank graphs are, as the name suggests, higher-dimensional analogues of directed graphs which we will define using category theory. The whole idea of my project was to construct what we call a Baumslag-Solitar graph, a higher-rank graph. They're too abstract an idea to picture, but in building Baumslag-Solitar graphs through coloured graphs, we are able to do so. I modelled my work on constructing Baumslag-Solitar graphs off Hazlewood, Raeburn, Sims, and Webster's *Remarks on some fundamental results about higher-rank graphs and their*  $C^*$ -algebras [2]. This is a novel approach in understanding the generalisation of directed graphs.


## 1 Motivation in constructing k-graphs

#### 1.1 The basic idea

**Definition 1.1.** A directed graph  $E = (E^0, E^1, r, s)$  consists of two countable sets  $E^0, E^1$  and functions  $r, s : E^1 \to E^0$ .

The set  $E^0$  is the set of vertices,  $E^1$  is the set of edges, r is called the range map and s is called the source map.

For  $f \in E^1$ , we have



A path of length n is a sequence of edges  $\mu = \mu_1 \mu_2 \dots \mu_n$  such that  $s(\mu_i) = r(\mu_{i+1})$ . Denote by  $E^n$  the set of all paths of length n and define  $E^* := \bigcup_{n \in \mathbb{N}} E^n$ .

We will take this idea of a directed graph and generalise it to a higher-dimension, then an abstract semigroup. This generalised graph will be defined using categorical framework. **Definition 1.2.** A category C consists of two sets  $C^0$  and  $C^*$ , two maps  $r, s : C^* \to C^0$ , a partially defined product  $(f,g) \mapsto fg$  from  $\{(f,g) \in C^* \times C^* : s(f) = r(g)\}$  to  $C^*$ , and distinguished elements  $\{\iota_v \in C^* : v \in C^0\}$  satisfying

- (i) r(fg) = r(f) and s(fg) = s(g);
- (ii) (fg)h = f(gh) when s(f) = r(g) and s(g) = r(h);
- (iii)  $r(\iota_v) = v = s(\iota_v)$  and  $\iota_v f = f$ ,  $g\iota_v = g$  when r(f) = v and s(g) = v.

The similarity between directed graphs and categories is the reason why the framework of categories is the natural choice in defining the generalisation of directed graphs.

### 1.2 Generalisation of a directed graph in $\mathbb{N}^k$

Instead of the length of a path having values in  $\mathbb{N}$ , we will explore the higher-dimensional analogue of a directed graph, first with the length of a path defined in  $\mathbb{N}^k$  as was first done by Hazelwood, Raeburn, Sims and Webster [2], and secondly with the length of a path defined in the Baumslag-Solitar semigroup.

A k-coloured graph is a directed graph E together with a map  $c: E^1 \to \{c_1, c_2, ..., c_k\}$ . So a path  $x \in E^*$  has both colour and shape determined by  $q: \{c_1, c_2, ..., c_k\} \to \mathbb{N}^k$  where  $q(c_i) = e_i$  for all i. For  $x \in E^*$ , define  $d^*(x) = q(c(x))$ . We call  $d^*(x)$  the degree of a path  $x \in E^*$  which we can see is assigned some higher-dimensional value in  $\mathbb{N}^k$ .

**Definition 1.3.** For  $m \in (\mathbb{N} \cup \{\infty\})^k$ , define a k-coloured graph  $E_{k,m}$  by

$$E_{k,m}^{0} = \{n \in \mathbb{N}^{k} : 0 \le n \le m\}$$
  

$$E_{k,m}^{1} = \{n + v_{i} : n, n + e_{i} \in E_{k,m}^{0}\}$$
  

$$r(n + v_{i}) = n, s(n + v_{i}) = n + e_{i}, \text{and } c(n + v_{i}) = c_{i}$$

A graph morphism  $\psi$  from a directed graph E to a directed graph F consists of functions  $\psi^0 : E^0 \to F^0$  and  $\psi^1 : E^1 \to F^1$  such that  $r_F(\psi^1(e)) = \psi^0(r_E(e))$  and  $s_F(\psi^1(e)) = \psi^0(s_E(e))$  for all  $e \in E^1$ . A coloured-graph morphism is a graph morphism  $\psi$  such that  $c_E(e) = c_F(\psi(e))$  for every  $e \in E^1$ .

For a given coloured-graph morphism  $\lambda : E_{k,m} \to E$  we say  $\lambda$  has degree m and write  $d(\lambda) = m$ , and define  $r(\lambda) := \lambda(0)$  and  $s(\lambda) := \lambda(m)$ .

For  $p, q \in \mathbb{N}^k$  with  $p \leq q$ , define  $E_{k,[p,q]}$  to be the subgraph of  $E_{k,q}$  such that

$$\begin{split} E^0_{k,[p,q]} &= \{ n \in \mathbb{N}^k : p \le n \le q \} \\ E^1_{k,[p,q]} &= \{ x \in E^1_{k,q} : s(x), r(x) \in E^0_{k,[p,q]} \} \end{split}$$

Given a coloured-graph morphism  $\lambda : E_{k,m} \to E$  and  $p, q \in \mathbb{N}^k$  such that  $p \leq q \leq m$ , define  $\lambda|_{E_{k,[p,q]}}^* : E_{k,[p,q]} \to E$  by

$$\lambda|_{E_{k,[p,q]}}^*(a) = \lambda(p+a)$$

The star is to remind us that this non-standard restriction involves a translation.

Given any k-coloured graph E and distinct  $i, j \in \{1, ..., k\}$ , an  $\{i, j\}$ - square in E is a coloured directed graph morphism  $\lambda : E_{k,e_i+e_j} \to E$ . A collection C of squares in E is complete if each  $x \in F^*$  with  $c(x) = c_i c_j$  and  $i \neq j$ , there is a unique  $\phi \in C$  such that  $x = \phi(v_i)\phi(e_i + v_j)$ . We write  $\phi(v_i)\phi(e_i + v_j) \sim \phi(v_j)\phi(e_j + v_i)$ .

If  $\lambda : E_{k,m} \to E$  is a coloured-graph morphism and  $\phi$  is a square in E, then  $\phi$  occurs in  $\lambda$  if there exists  $n \in \mathbb{N}^k$  such that  $\phi(x) = \lambda(x+n)$  for all  $x \in E_{k,e_i+e_i}$ .

Let *E* be a *k*-coloured graph, and  $m \in \mathbb{N}^k \setminus \{0\}$ . Fix  $x \in E^*$  and a coloured-graph morphism  $\lambda : E_{k,m} \to E$ . We say *x* traverses  $\lambda$  if the shape q(c(x)) of *x* is equal to  $d^*(x)$ and  $\lambda(q(c(x_1x_2...x_{l-1})) + v_{c_{x_l}}) = x_l$  for all  $0 < l \leq |m|$ . If m = 0, then  $x \in E^0$  and  $dom(\lambda) = \{0\}$ , and we say *x* traverses  $\lambda$  if  $x = \lambda(0)$ .

#### 1.3 Some major results

**Theorem 1.4.** Let E be a 2-coloured graph and let C be a complete collection of squares in E. For every  $x \in E^*$  there is a unique C-compatible coloured-graph morphism  $\lambda_x : E_{2,d(x)} \to E$  such that x traverses  $\lambda_x$ .

*Proof.* We prove by induction over |x|. If |x| = 0, then  $x \in E^0$ . Define the colouredgraph morphism  $\lambda_x : E_{2,(0,0)} \to E$  by  $\lambda_x((0,0)) \mapsto x$ . Then  $d(\lambda_x) = q(c(x)) = (0,0)$  and  $\lambda_x((0,0)) = x$ . Hence x traverses  $\lambda_x$ .

 $\lambda_x$  maps a single point  $(0,0) \in E_{2,(0,0)}$  to  $x \in E$ , there is only one way this can be done hence  $\lambda_x$  is unique. No squares occur in  $\lambda_x$ , so it is trivial to see that  $\lambda_x$  is C-compatible.

Now suppose, for the inductive hypothesis, that for each  $y \in E^*$  with  $y \leq n$  there exists a unique coloured-graph morphism  $\lambda_y : E_{2,d(y)} \to E$  such that y traverses  $\lambda_y$ . Fix a path  $x \in E^*$  with |x| = n+1, and express x = yf where  $f \in E^1$  with  $c(f) = c_i$ . Let m := q(c(y)). By the inductive hypothesis, there exists a unique  $\mathcal{C}$ -compatible coloured-graph morphism  $\lambda_y$  such that y traverses  $\lambda_y$ .

Consider the set  $\{j \leq 2 : j \neq i, m_j > 0\}$  as colours in the path y that differ from  $c(f) = c_i$  where  $m = (m_1, m_2)$ . Note that  $m_j$  can be thought of as expressing how many edges of colour j exist in y. Since we are only dealing with a maximum of two colours, we have two cases: (1) edges in y are all colour i, or, (2) at least one edge in y is colour j.

That is,

$$|\{j \le 2 : j \ne i, m_j > 0\}| = \begin{cases} 0 : \text{Case 1} \\ 1 : \text{Case 2} \end{cases}$$

For Case 1, we can see that  $E_{2,d(x)} = E_{2,m} \cup E_{[m,m+e_i]}$ . Figure 1.5. Image of  $E_{2,d(x)}$  for i = 1

$$\underbrace{(0,0) + v_1}_{(0,0)} \underbrace{(1,0) + v_1}_{(1,0)} \underbrace{(2,0) + v_1}_{(2,0)} \cdots \underbrace{m + v_1}_{m + e_1}$$

So define the coloured graph morphism  $\lambda_x$  by the formulae

$$\lambda_x|_{E_{2,m}} = \lambda_y, \lambda_x(m+v_i) = f \text{ and } \lambda_x(m+e_i) = s(f)$$
(1)

Figure 1.6. Image of  $\lambda_x$  for i = 1

$$\begin{array}{cccc} y_1 & y_2 & y_n & f \\ \bullet & & \\ r(y_1) & s(y_1) = r(y_2) & s(y_2) = r(y_3) & s(y_n) = r(f) & s(f) \end{array}$$

Hence x traverses  $\lambda_x$ . To see that  $\lambda_x$  is unique, suppose x also traverses a colouredgraph morphism  $\mu$ . Then  $d(x) = q(c(x)) = d(\mu)$  and  $\mu(q(c(x_1x_2...x_{l-1})) + v_{c(x_l)}) = x_l$  for all  $0 < l \leq |n+1|$ . Which roughly says that  $\mu$  maps all successive edges in  $E_{2,d(x)}$  to the path  $x = yf \in E^*$ , so then  $\mu$  satisfies the formula (1) hence  $\mu = \lambda_x$ .

For Case 2 we have  $i \neq j$  satisfying  $m_j > 0$ . Then

$$E_{2,m+e_i} = E_{2,m} \cup (E_{2,m+e_i-e_j}) \cup (E_{2,[m-e_j,m+e_i]})$$

Claim 1: Let  $\phi^j$  be the unique square in  $\mathcal{C}$  traversed by  $\lambda_y((m-e_j)+v_j)f$ . Let  $g^j = \phi^j(0+v_i)$  and  $h^j = \phi^j(e_i+v_j)$ , so  $g^jh^j \sim \lambda_y((m-e_j)+v_j)f$ . Then there is a unique coloured-graph morphism  $\lambda^j : E_{2,m-e_j+e_i} \to E$  such that  $\lambda^j|_{E_{2,m-e_j}} = \lambda_y|_{E_{2,m-e_j}}$  and  $\lambda^j((m-e_j)+v_i) = g^j$ .

Fix a path  $z^j$  which traverses  $\lambda_y|_{E_{2,m-e_j}}$ . Observe that  $|z^jg^j| = n$ , so the inductive hypothesis gives a unique  $\mathcal{C}$ -compatible coloured-graph morphism  $\lambda^j : E_{2,m-e_j+e_i} \to E$  such that  $z^jg^j$  traverses  $\lambda^j$ . So then  $z^j$  traverses  $\lambda^j|_{E_{2,m-e_j}}$ . By the uniqueness of the  $\mathcal{C}$ -compatible coloured-graph morphism  $\lambda^j$ , we have  $\lambda^j|_{E_{2,m-e_j}} = \lambda_y|_{E_{2,m-e_j}}$  proving Claim 1.

Figure 1.7. Image of  $E_{2,m+e_i}$  for i = 1




Figure 1.8. Image of  $\phi^j$  for j = 2

$$s(\lambda_y((m - e_2) + v_2)) = r(f) \bullet (f) \bullet (f) \bullet (f) = s(h^2)$$

$$\lambda_y((m - e_2) + v_2) \bullet (h^2) \bullet (h^2) = s(g^2)$$

$$r(\lambda_y((m - e_2) + v_2)) = r(g^2) \bullet (g^2) \bullet (h^2) = s(g^2)$$

Define the function  $\lambda_x: E_{2,m+e_i} \to E$  via the formula

$$\lambda_x|_{E_{2,m}} = \lambda_y, \lambda_x|_{E_{2,m+e_i-e_j}} = \lambda^j \text{ and } \lambda_x|_{E_{2,[m-e_j,m+e_i]}} = \phi^j$$

This  $\lambda_x$  is well defined on its domain from Claim 1. Since  $\phi^j$  is a square in  $\mathcal{C}$ , with  $\lambda_y$  and  $\lambda^j \mathcal{C}$ -compatible, then  $\lambda_x$  is  $\mathcal{C}$ -compatible, and x = yf traverses  $\lambda_x$ .

For uniqueness, fix a C-compatible coloured-graph morphism  $\mu$  traversed by x. We wish to show that  $\mu = \lambda_x$ , we will show this piece by piece. Since x = yf traverses  $\mu$ , y traverses  $\mu|_{E_{2,m}}$ . The inductive hypothesis says that

$$\mu|_{E_{2,m}} = \lambda_y = \lambda_x|_{E_{2,m}}.$$

Now,  $z^j$  is a traversal of  $\lambda_y|_{E_{2,m-e_j}} = \lambda_x|_{E_{2,m-e_j}} = \mu|_{E_{2,m-e_j}}$ . Then  $z^j\lambda_y(m-e_j+v_j)$  traverses  $\lambda_y$ . So  $z^j\lambda_y((m-e_j)+v_j)f$  traverses  $\mu$ . Now  $\phi^j$  is the unique square in  $\mathcal{C}$  traversed by  $\lambda_y((m-e_j)+v_j)f$ . So

$$\mu|_{2,[m-e_j,m+e_1]}^* = \phi^j = \lambda_x|_{2,[m-e_j,m+e_1]}^*$$

Now  $z^j g^j$  traverses  $\mu|_{E_{2,m+e_i-e_j}}$  so the inductive hypothesis forces

$$\mu|_{E_{2,m+e_i-e_j}} = \lambda^j = \lambda_x|_{E_{2,m+e_i-e_j}}$$

Hence  $\mu = \lambda_x$ , completing the proof for the second case.

**Notation 1.9.** Fix a 2-coloured graph E and a complete collection of squares C in E. For each  $m \in \mathbb{N}^2$ , we write  $\Lambda^m$  for the set of all C-compatible coloured-graph morphisms  $\lambda : E_{2,m} \to E$ . For  $v \in E^0$  we define  $\lambda_v : E_{2,0} \to E$  by  $\lambda_v(0) = v$ . Let

$$\Lambda = \bigcup_{m \in \mathbb{N}^2} \Lambda^m.$$

We have previously defined  $d: \Lambda \to \mathbb{N}^2$  and  $r, s: \Lambda \to \Lambda^0$ .

**Theorem 1.10.** If  $\mu : E_{2,m} \to E$  and  $\nu : E_{2,n} \to E$  are *C*-compatible coloured-graph morphisms such that  $s(\mu) = r(\nu)$ , then there exists a unique *C*-compatible coloured-graph morphism  $\mu\nu : E_{m+n} \to E$  such that  $(\mu\nu)|_{E_{2,m}} = \mu$  and  $(\mu\nu)|_{E_{2,[m,m+n]}}^* = \nu$ . Under this composition map, the set  $\Lambda$  is a category.

Proof. Fix  $x, y \in E^*$  such that x traverses  $\mu$  and y traverses  $\nu$ . Lemma lem2 implies that xy traverses a unique  $\mathcal{C}$ -compatible coloured-graph morphism  $\mu\nu$ . Then x traverses  $(\mu\nu)|_{E_{2,m}}$ , and y traverses  $(\mu\nu)|_{E_{2,[m,m+n]}}$ , so the uniqueness of the coloured-graph morphism in Theorem - gives  $(\mu\nu)|_{E_{2,m}} = \mu$  and  $(\mu\nu)|_{E_{2,[m,m+n]}} = \nu$ . That same uniqueness property forces any other coloured-graph morphism  $\lambda$  with  $\lambda|_{E_{2,m}} = \mu$  and  $\lambda|_{E_{2,[m,m+n]}} = \nu$  to equal  $\mu\nu$ , because xy traverse both  $\lambda$  and  $\mu\nu$ .

It remains to show that  $\Lambda$  is a category. For composable  $\mu$ ,  $\nu$  we have

$$s(\mu\nu) = (\mu\nu)(d(\mu\nu)) = (\mu\nu)|_{E_{2,[d(\mu),d(\mu)+d(\nu)]}}^*(d(\nu)) = \nu(d(\nu)) = s(\nu)$$

Similarly,  $r(\mu\nu) = r(\mu)$ . So condition (i) in Definition 2.27 holds.

For condition (ii) in Definition 2.27 of associativity, let  $\lambda : E_{2,l} \to E$ ,  $\mu : E_{2,m} \to E$ and  $\nu : E_{2,n} \to E$  be  $\mathcal{C}$ -compatible coloured-graph morphisms such that  $s(\lambda) = r(\mu)$ and  $s(\mu) = r(\nu)$ . Fix  $x_{\lambda}, x_{\mu}, x_{\nu} \in E^*$  such that  $x_{\lambda}$  traverses  $\lambda, x_{\mu}$  traverses  $\mu$  and  $x_{\nu}$ traverses  $\nu$ . Then  $x_{\lambda}x_{\mu}$  traverses  $\lambda\mu$ . Hence  $x_{\lambda}x_{\mu}x_{\nu} = (x_{\lambda}x_{\mu})x_{\nu}$  traverses  $(\lambda\mu)\nu$ . Similarly,  $x_{\lambda}x_{\mu}x_{\nu} = x_{\lambda}(x_{\mu}x_{\nu})$  traverses  $\lambda(\mu\nu)$ . So Theorem 1.4 implies that  $(\lambda\mu)\nu = \lambda(\mu\nu)$ 

Now for condition (iii) in Definition 2.27, for  $v \in E^0$ , we have  $r(\lambda_v) = \lambda_v(0) = v$  and  $s(\lambda_v) = \lambda_v(d(\lambda_v)) = \lambda_v(0) = v$  by definition. Further, if  $r(\mu) = v = s(\lambda_v)$ , then  $\lambda_v \mu = \mu$  and if  $s(\nu) = v = r(\lambda_v)$ , then  $\nu = \lambda_v \nu$ .

# 2 Generalisation of a directed graph in the Baumslag-Solitar semigroup

**Definition 2.1.** For each pair of non-zero integers *m* and *n*, there is a corresponding group

$$BS(m,n) = \langle a,b \mid ab^m = b^n a \rangle$$

Denote  $BS(m,n)^+$  as the semigroup of all reduced elements in BS(m,n) that contain no inverses  $a^{-1}$  or  $b^{-1}$ . We will only be dealing with the semigroup  $BS(2,1)^+$  for the remainder of this report, hence the semigroup relation is  $ab^2 = ba$ .


Fix a 2-coloured graph E. We have the colour map  $c: E^1 \to \{c_a, c_b\}$ . Let  $c_a := red$  and  $c_b := blue$ . Define  $q: \{c_a, c_b\} \to BS(2, 1)^+$  with  $q(c_l) = l$  for  $l \in \{a, b\}$ . Hence the degree of a path  $x \in E^*$  is now  $d^*(x) = q(c(x)) \in BS(2, 1)^+$ .

**Definition 2.2.** For  $w \in BS(2,1)^+$ , we define the 2-coloured graph  $E_w$  by

$$E_w^0 = \{z \in BS(2,1)^+ : z \le w\}, \quad E_w^1 = \{(z,za) : z, za \in E_w^0\} \bigcup \{(z,zb) : z, zb \in E_w^0\} = \{(z,zl) : z, zl \in E_w^0 \text{ and } l = a, b\}$$

With the maps r(z, zl) = z, s(z, zl) = zl and  $c(z, zl) = c_l$ .

Figure 2.3. The 2- coloured graph  $E_{ba}$ 



**Definition 2.4.** For a given coloured-graph morphism  $\lambda : E_w \to E$  we say  $\lambda$  has degree  $w \in BS(2,1)^+$  and write  $d(\lambda) = w$ . Define  $r(\lambda) := \lambda(e)$  and  $s(\lambda) := \lambda(w)$ . For  $v \in E^0$  we define  $\lambda_v : E_e \to E$  by  $\lambda_v(e) = v$ .

Let E be a 2 coloured-graph. A collection of squares C in E is complete if:

1. for each  $x \in E^*$  with  $c(x) = c_a c_b c_b$ , there exists a unique  $\varphi \in \mathcal{C}$  such that  $x = \varphi(e, a)\varphi(a, ab)\varphi(ab, ab^2)$ ; and

2. for each  $x \in E^*$  with  $c(x) = c_b c_a$ , there exists a unique  $\varphi \in \mathcal{C}$  such that  $x = \varphi(e, b)\varphi(b, ba)$ .

If  $\lambda: E_w \to E$  is a coloured-graph morphism and  $\varphi: E_{ba} \to E$  is a square in E, then  $\varphi$  occurs in  $\lambda$  if:

- 1. there exists  $m \in BS(2,1)^+$  such that  $\lambda(zm) = \varphi(z)$  for all  $z \in E_{ba}^0$ , and,
- 2.  $\lambda(zm, zml) = \varphi(z, zl)$  for all  $(z, zl) \in E_{ba}^1$

A coloured-graph morphism  $\lambda : E_w \to E$  is *C*-compatible if every square occurring in  $\lambda$  belongs to the collection  $\mathcal{C}$ .

For a path  $x \in E^*$  and coloured-graph morphism  $\lambda : E_w \to E$ , we say x traverses  $\lambda$  if  $d(\lambda) = d^*(x)$  and if  $\lambda(q(c(x_1x_2...x_{n-1})), q(c(x_1x_2...x_{n-1}))q(c(x_n))) = x_n$  for all  $0 < n \le |x|$ .

For  $w_1, w_2 \in BS(2,1)^+$  with  $w_2 = w_1w'$  for  $w' \in BS(2,1)^+$ , define  $E_{[w_1,w_2]}$  to be the subgraph of  $E_{w_2}$  such that


$$E^{0}_{[w_{1},w_{2}]} = \{ w \in BS(2,1)^{+} : w_{1} \text{ is a subword of } w, w \text{ is a subword of } w_{2} \}$$
$$E^{*}_{[w_{1},w_{2}]} = \{ x \in E^{1}_{w_{2}} : s(x), r(x) \in E^{0}_{[w_{1},w_{2}]} \}$$

Given a coloured-graph morphism  $\lambda : E_w \to E$  and  $w_1, w_2 \in BS(2,1)^+$  such that  $w_2 = w_1w'$  and  $w = w_2w''$  for  $w', w'' \in BS(2,1)^+$  define  $\lambda|_{[w_1,w_2]}^* : E_{w_1^{-1}w_2} \to E$  by

$$\lambda|_{[w_1,w_2]}^*(a) = \lambda(w_1a)$$


**Example 2.5.** Let E be the 2-coloured graph as in Figure 2.6.

Figure 2.6.



Squares in E are  $\varphi_1$  and  $\varphi_2$  in Figure 2.7.


Figure 2.7.



We can see that E has two blue-blue-red paths, and they are kkf and ggh, represented only once in the collection  $\mathcal{C} = \{\varphi_1, \varphi_2\}$ , as are the red-blue paths hk and fg in E. So  $\{\varphi_1, \varphi_2\}$  form a complete collection.

Let  $\lambda: E_{b^2a^2} \to E$  be defined as in Figure 2.8





So  $\varphi_1$  occurs in  $\lambda$  because  $\lambda^*_{[b,b^2a]} = \varphi_1$ . In fact, every square occurring in  $\lambda$  is either  $\varphi_1$  or  $\varphi_2$ , so  $\lambda$  is  $\mathcal{C}$ -compatible. Also, for example, the path  $ggfh \in E^*$  traverses  $\lambda$ .

Remark 2.9. Every  $w \in BS(2,1)^+$  can be expressed as

$$b^{m_1}a^{n_1}ba^{n_2}ba^{n_3}b...ba^{n_0}$$
 and  $a^nb^m$ 

where  $n_i \neq 0 \ \forall i \neq 0$ .

**Definition 2.10.** Let E be a 2-coloured graph, and let  $\lambda : E_w \to E$  be a coloured-graph morphism, with  $x \in E^*$  a traversal of  $\lambda$ . Define  $\lfloor x \rfloor \in E^*$  to be a traversal of  $\lambda$  such that |x| is minimised. Expressing  $w = b^m a^{n_1} b a^{n_2} b a^{n_3} b \dots b a^{n_0}$ , then the shortest path traversing  $\lambda$  is of the form:

$$\begin{split} \lfloor x \rfloor = &\lambda(e, b)\lambda(b, b^{2})...\lambda(b^{m-1}, b^{m}) \text{ or } \\ \lfloor x \rfloor = &\lambda(e, a)\lambda(a, a^{2})...\lambda(a^{n_{1}-1}, a^{n_{1}})\lambda(a^{n_{1}}, a^{n_{1}}b)\lambda(a^{n_{1}}b, a^{n_{1}}ba)...\lambda(a^{n_{1}}ba^{n_{2}-1}, a^{n_{1}}ba^{n_{2}}) \\ &\lambda(a^{n_{1}}ba^{n_{2}-1}, a^{n_{1}}ba^{n_{2}}b)...\lambda(b^{m}a^{n_{1}}ba^{n_{2}}ba^{n_{3}}b...a^{n_{k}}, b^{m}a^{n_{1}}ba^{n_{2}}ba^{n_{3}}b...a^{n_{k}}b) \text{ or } \\ \lfloor x \rfloor = &\lambda(e, a)\lambda(a, a^{2})...\lambda(a^{n_{1}-1}, a^{n_{1}})\lambda(a^{n_{1}}, a^{n_{1}}b)\lambda(a^{n_{1}}b, a^{n_{1}}ba)...\lambda(a^{n_{1}}ba^{n_{2}-1}, a^{n_{1}}ba^{n_{2}}) \\ &\lambda(a^{n_{1}}ba^{n_{2}-1}, a^{n_{1}}ba^{n_{2}}b)...\lambda(b^{m}a^{n_{1}}ba^{n_{2}}ba^{n_{3}}b...a^{n_{k}}ba^{n_{0}-1}, b^{m}a^{n_{1}}ba^{n_{2}}ba^{n_{3}}b...a^{n_{k}}ba^{n_{0}}) \end{split}$$

Define  $\lceil x \rceil \in E^*$  to be a traversal of  $\lambda$  such that |x| is maximised. Expressing  $w = a^n b^m$ , we have the longest path traversing  $\lambda$  is:

$$\begin{split} \lceil x \rceil = &\lambda(e, b)\lambda(b, b^2)...\lambda(b^{m-1}, b^m) \text{ or} \\ \lceil x \rceil = &\lambda(e, a)\lambda(a, a^2)...\lambda(a^{n-1}, a^n) \text{ or} \\ \lceil x \rceil = &\lambda(e, a)\lambda(a, a^2)...\lambda(a^n, a^nb)...\lambda(b^{m-1}, b^m) \end{split}$$

This implies that every coloured-graph morphism has a traversal.

**Example 2.11.** Let  $\lambda : E_{b^2a^2} \to E$  be a coloured-graph morphism. Let  $x \in E^*$  traverse  $\lambda$ . Then the shortest path traversing  $\lambda$  is |x| with

$$d^*(\lfloor x \rfloor) = q(c(\lfloor x \rfloor))$$
$$= q(c_b c_b c_a c_a)$$
$$= b^2 a^2$$

The longest path traversing  $\lambda$  is [x] with

$$d^*(\lceil x \rceil) = q(c(\lceil x \rceil))$$
  
=  $q(c_a c_a c_b c_b c_b c_b c_b c_b c_b c_b c_b)$   
=  $a^2 b^8$ 

Note that  $b^2 a^2 = a^2 b^8 \in BS(2,1)^+$ , this illustrates how paths of the same degree can differ in length.

**Lemma 2.12.** Let E be a 2 coloured-graph and let  $\lambda : E_w \to E$  be a coloured-graph morphism where  $w \in BS(2,1)^+$ . Write  $w = w_1w_2$  where  $w_1, w_2 \in BS(2,1)^+$ . If  $x \in E^*$ traverses  $\lambda|_{E_{[e,w_1]}}$  and  $y \in E^*$  traverses  $\lambda|_{E_{[w_1,w_1w_2]}}^*$ , then xy traverses  $\lambda$ .

$$\begin{aligned} Proof. & \text{We have } d^*(\lambda) = w = w_1 w_2 = q(c(x))q(c(y)) = q(c(xy)) \text{ and for } m \leq |xy|, \text{ we have} \\ \lambda(d^*((xy)_1...(xy)_{m-1}), d^*((xy)_1...(xy)_{m-1})q(c(xy_m))) \\ &= \begin{cases} \lambda|_{E_{[e,w_1]}}(d^*((xy)_1...(xy)_{m-1}), d^*((xy)_1...(xy)_{m-1})q(c(xy_m)))) & \text{ if } m \leq |x| \\ \lambda|_{E_{[w_1,w_1w_2]}}^*(d((xy)_1...(xy)_{m-|x|-1}), d((xy)_1...(xy)_{m-|x|-1})q(c(xy_{m-|x|})))) & \text{ otherwise} \end{cases} \\ &= \begin{cases} x_m & \text{ if } m \leq |x| \\ y_{m-|x|} & \text{ otherwise} \end{cases} \end{aligned}$$

**Theorem 2.13.** Let E be a 2-coloured graph and let C be a complete collection of squares in E. For every  $x \in E^*$  there is a unique C-compatible coloured-graph morphism  $\lambda_x : E_{d^*(x)} \to E$ .

We will prove by induction over |x|. If |x| = 0 so that x is a vertex, then  $\lambda_x : E_e \to E$ with  $\lambda_x(e) \mapsto x$  is the unique coloured-graph morphism such that x traverses  $\lambda_x$ . That  $\lambda_x$ is  $\mathcal{C}$ -compatible is trivial given that no squares exist in  $\lambda_x$ .

Suppose, for the inductive hypothesis, that for every  $y \in E^*$  with  $|y| \leq n$ , the path y traverses a unique coloured graph morphism  $\lambda_y : E_{d^*(y)} \to E$ . Fix a path  $x \in E^*$  with |x| = n + 1. Express x = yf where  $y \in E^*$  and  $f \in E^1$  and  $c(f) = c_l$  where  $l \in \{a, b\}$ .


Let  $w := d^*(y) = q(c(y))$ . Express w in its shortest form  $w = b^m a^{n_1} b a^{n_2} b a^{n_3} b \dots a^{n_k} b a^{n_0}$ where  $n_i \neq 0 \ \forall i \neq 0$ . Then either

Case 1: 
$$n_0 = 0$$
 so that  $w = b^m a^{n_1} b a^{n_2} b a^{n_3} b \dots a^{n_k} b$  where  $n_i \neq 0 \ \forall i \neq 0$ 

Case 2:  $n_0 \neq 0$  so that  $w = b^m a^{n_1} b a^{n_2} b a^{n_3} b \dots a^{n_k} b a^{n_0}$  where  $n_i \neq 0 \ \forall i$ 

For Case 1:  $E_w$  contains, by definition, all  $w' \in BS(2,1)^+$  such that w = w'w'' for some  $w'' \in BS(2,1)^+$ , and edges whose source and range belong to the set of vertices and differ by a basis element, characterising  $E_w$  by the blue edge  $(wb^{-1}, w)$  seen in Figure 2.14.


**Figure 2.14.** Image of  $E_w$  as the domain of  $\lambda_y$ 



Suppose first that  $c(f) = c_b$ . We have |y| = n, so  $|\lfloor y \rfloor| \leq n$ . We know now that that path  $\lfloor y \rfloor$  ends in a red then a blue edge. Write  $\lfloor y \rfloor = \lfloor y \rfloor_1 \lfloor y \rfloor_2 \dots \lfloor y \rfloor_k$  and define  $y' = [y]_1[y]_2...[y]_{k-2}$  so that  $|y'| \le n-2$ . Hence y' traverses  $\lambda_y|_{E_{wb^{-1}a^{-1}}}$ . Let  $\phi$  be the unique square in  $\mathcal{C}$  traversed by  $\lambda(wb^{-1}a^{-1}, wb^{-1})\lambda(wb^{-1}, w)f$  so

 $\lambda(wb^{-1}a^{-1},wb^{-1})\lambda(wb^{-1},w)f\sim \phi(wb^{-1}a^{-1},wba^{-1})\phi(wb,wba^{-1})$ 

Figure 2.15. Image of  $\phi$ 



Let  $x' = y'\phi(wb^{-1}a^{-1}, wba^{-1})\phi(wb, wba^{-1})$ . We have  $|x'| \leq n$  with  $d^*(x') = wb$ , so the inductive hypothesis implies that x' traverses a unique  $\mathcal{C}$ -compatible coloured-graph morphism  $\lambda_{x'} : E_{wb} \to E$ .

**Lemma 2.16.** Suppose  $z \in E^n$  has  $c(z_{n-1}z_n) = c_bc_a$  and z traverses a C-compatible 2coloured graph-morphism  $\lambda : E_{d^*(z)} \to E$ . Then

 $z_1...z_{n-2}\lambda(d^*(z)a^{-1}b^{-1}, d^*(z)a^{-1}b^{-1}a)\lambda(d^*(z)a^{-1}b^{-1}a, d^*(z)a^{-1}b^{-1}ab)\lambda(d^*(z)a^{-1}b^{-1}ab, d^*(z)a^{-1}b^{-1}ab^2)$ 

also traverses  $\lambda$ .

*Proof.* Suppose  $z = z_1 z_2 \dots z_n$  traverses  $\lambda$  as hypothesised above. Then  $d(\lambda) = q(c(z))$  and

$$\lambda(q(c(z_1z_2...z_{m-1})), q(c(z_1z_2...z_{m-1}))l_m) = z_m$$

for all  $0 < m \le |z|$ .

Let

$$\begin{aligned} z_{n-1}' &= \lambda(d(z)a^{-1}b^{-1}, d(z)a^{-1}b^{-1}a) \\ z_n' &= \lambda(d(z)a^{-1}b^{-1}a, d(z)a^{-1}b^{-1}ab) \\ z_{n+1}' &= \lambda(d(z)a^{-1}b^{-1}ab, d(z)a^{-1}b^{-1}ab^2) \\ \text{and } z' &= z_1...z_{n-2}z_{n-1}'z_n'z_{n+1}' \end{aligned}$$

Then

$$q(c(z')) = q(c(z_1...z_{n-2})c(z'_{n-1}z'_nz'_{n+1})$$
  
=  $q(c(z_1...z_{n-2})c_ac_bc_b)$   
=  $q(c(z_1...z_{n-2}))ab^2$   
=  $q(c(z_1...z_{n-2}))ba$   
=  $q(c(z))$ 

Because z traverses  $\lambda$ , we know

$$\lambda(q(c(z_1z_2...z_{m-1})), q(c(z_1z_2...z_{m-1}))l_m) = z_m \text{ for all } 0 < m \le |z| - 2.$$
  

$$\lambda(q(c(z_1z_2...z_{n-2})), q(c(z_1z_2...z_{n-2}))a) = z'_{n-1}$$
  

$$\lambda(q(c(z_1z_2...z_{n-2}z'_{n-1})), q(c(z_1z_2...z_{n-2}z'_{n-1}))b) = z'_n$$
  


$$\lambda(q(c(z_1z_2...z_{n-2}z'_{n-1}z'_n)), q(c(z_1z_2...z_{n-2}z'_{n-1}z'_n))b) = z'_{n+1}$$

Hence z' traverses  $\lambda$ .

proof. of Theorem 2.13 continued: Because  $|x'| \leq n$  and  $c(x') = c_{\lfloor y \rfloor_1} c_{\lfloor y \rfloor_2} \dots c_b c_a$ , Lemma 2.16 give the path  $\lfloor y \rfloor f$  as a traversal of  $\lambda'$ . Since  $\lfloor y \rfloor \sim y$ , then  $\lfloor y \rfloor f \sim yf$ . Hence x = yf traverses the unique C-compatible graph morphism  $\lambda_{x'}$ .

Suppose  $c(f) = c_a$ . Write  $y = y_1 y_2 \dots y_k = y' y_k$  so that |y'| = n - 1. Let  $\phi'$  be the unique square in  $\mathcal{C}$  traversed by  $y_k f$ . Let  $f' = \phi'(wb^{-1}, wb^{-1}a)$ .

Figure 2.17. Image of  $\phi'$ 



We have |y'f'| = n, so the inductive hypothesis gives a unique *C*-compatible colouredgraph morphism  $\lambda_{y'f'} : E_{d^*(y'f')} \to E$  with y'f' traversing  $\lambda_{y'f'}$ . Note that

$$E_{d^*(x)} = E_{d^*(y)} \cup E_{d^*(y'f')} \cup E_{[wb^{-1},wa]}.$$


So we can define the coloured-graph morphism  $\lambda_x$  by the equations

$$\begin{split} \lambda_x|_{E_{d^*(y'f')}} &= \lambda_{y'f'}\\ \lambda_x|_{E_{d^*(y)}} &= \lambda_y\\ \lambda_x|_{E_{[wb^{-1},wa]}} &= \phi' \end{split}$$

such that x = yf traverses  $\lambda_x$ . Since  $\lambda_{y'f'}$  and  $\lambda_y$  are both  $\mathcal{C}$ -compatible, and  $\phi'$  belongs to  $\mathcal{C}$ ,  $\lambda_x$  is also  $\mathcal{C}$ -compatible. To see that  $\lambda_x$  is unique, fix a  $\mathcal{C}$ -compatible coloured-graph morphism  $\mu$  traversed by x. Then y traverses  $\mu|_{E_{d(y)}}$ , so the inductive hypothesis forces  $\mu|_{E_{d(y)}} = \lambda_y = \lambda_x|_{E_{d(y)}}$ . That  $\mu$  is  $\mathcal{C}$ -compatible forces  $\mu|_{E_{[wb^{-1},wa]}} = \phi'$ , so that y'f'traverses  $\mu|_{E_{d(y'f')}} = \lambda_{y'f'} = \lambda_x|_{E_{d(y'f')}}$ . This completes the proof for Case 1.

For Case 2:

**Figure 2.18.** Image of the 2-coloured graph  $E_w$ , and domain of the coloured-graph morphism  $\lambda_y$ 



Suppose  $c(f) = c_b$ . We have  $E_{wb} = E_w \cup E_{[w,wb]}$ . Define the coloured-graph morphism  $\lambda_x$  by the formulae


$$\lambda_x|_{E_w} = \lambda_y|_{E_w}, \ \lambda_x(w, wb) = f \text{ and } \lambda_x(wb) = s(f)$$

By construction, x traverses  $\lambda_x$ . Given  $\lambda_y$  is C-compatible and unique, that  $\lambda_x$  is C-compatible and unique is trivial.

Now suppose that  $c(f) = c_a$ . Write d(y) = w in its longest form, so that  $w = a^N b^M$  for some  $N, M \in \mathbb{N}$ .


Let  $\phi_1$  be the unique square in  $\mathcal{C}$  traversed by the blue-red path  $\lambda_y(wb^{-1}, w)f$ .

Figure 2.19. Image of  $\phi_1$ 



Define recursively the squares  $\phi_n \in \mathcal{C}$  where  $\phi_n$  is uniquely determined by the traversal  $\lambda_y(wb^{-n}, wb^{-n+1})\phi_{n-1}(wb^{-n+1}, wb^{-n+1}a)$  for  $1 < n \leq M$ .

Figure 2.20. Image of  $\phi_n$ 



Define the coloured-graph morphism  $\lambda_x$  by the formulae

$$\begin{split} \lambda_x|_{E_{d^*(y)}} &= \lambda_y \\ \lambda_x|_{E_{[wb^{-n},wb^{-n+1}a]}} &= \phi_n \text{ for } 1 \leq n \leq M \end{split}$$

Then x traverses  $\lambda_x$ , and  $\lambda_x$  is C-compatible because  $\lambda_y$  is C-compatible and  $\phi_n \in C$  for  $1 \leq n \leq M$ .

To see that  $\lambda_x$  is unique, suppose there exists a C-compatible coloured-graph morphism  $\mu$  such that x traverses  $\mu$ . Then y traverses  $\mu|_{E_{d^*(y)}}$ , so the inductive hypothesis forces  $\lambda_x|_{E_{d^*(y)}} = \mu|_{E_{d^*(y)}} = \lambda_y$ . Now we have

$$\lambda_y(wb^{-n}, wb^{-n+1}) = \lambda_x(wb^{-n}, wb^{-n+1}) = \mu(wb^{-n}, wb^{-n+1}) \text{ for all } 1 \le n \le M.$$

Hence  $\mu|_{E_{[wb^{-n},wb^{-n+1}a]}} = \phi_n$  for all  $1 \le n \le M$ . So  $\mu = \lambda_x$ .

**Corollary 2.21.** Let E be a 2-coloured graph and let C be a complete collection of squares in E. If  $\mu : E_{w_1} \to E$  and  $\nu : E_{w_2} \to E$  are C-compatible coloured-graph morphisms such that  $s(\mu) = r(\nu)$ , then there exists a unique C-compatible coloured-graph morphism  $\mu\nu : E_{w_1w_2} \to E$ , called the composition of  $\mu$  and  $\nu$  such that  $(\mu\nu)|_{E_{w_1}} = \mu$  and  $(\mu\nu)|_{E_{[w_1,w_1w_2]}}^* = \nu$ .

Proof. Fix  $x, y \in E^*$  such that x traverses  $\mu$  and y traverses  $\nu$ . Then s(x) = r(y) and Theorem 2.13 implies that xy traverses a unique C-compatible coloured-graph morphism  $\mu\nu$ . Restricting the domain gives that x traverses  $(\mu\nu)|_{E_{w_1}}$ , because  $d^*(x) = w_1$ , and y traverses  $(\mu\nu)|_{E_{[w_1,w_1w_2]}}$ , because  $d^*(y) = w_2$ . So the uniqueness of  $(\mu\nu)|_{E_{w_1}}$  and  $(\mu\nu)|_{E_{[w_1,w_1w_2]}}^*$  by Theorem 2.13 give  $(\mu\nu)|_{E_{w_1}} = \mu$  and  $(\mu\nu)|_{E_{[w_1,w_1w_2]}}^* = \nu$ .

So we have a unique C-compatible coloured-graph morphism  $\mu\nu$  given traversals x of  $\mu$ and y of  $\nu$ . We wish to show that  $\mu\nu$  is unique given  $\mu$  and  $\nu$ . Suppose that we start with any other coloured-graph morphism, say  $\lambda$ , such that  $\lambda|_{E_{w_1}} = \mu$  and  $\lambda|_{E_{[w_1,w_1w_2]}}^* = \nu$ . So xtraverses  $\lambda|_{E_{w_1}}$  and y traverses  $\lambda|_{E_{[w_1,w_1w_2]}}^*$ . Then Lemma 2.12 shows that xy traverses  $\lambda$ . Since xy traverses both  $\lambda$  and  $\mu\nu$ , the uniqueness in Theorem 2.13 forces  $\lambda = \mu\nu$ . Hence the composition of  $\mu$  and  $\nu$  is unique.

Remark 2.22. Corollary 2.21 implies that  $\mu := \lambda|_{E_{w_1}}$  and  $\nu := \lambda|_{E_{[w_1,w_1w_2]}}^*$  satisfy  $\lambda = \mu\nu$ . We wish to show that  $\mu$  and  $\nu$  are unique. Suppose that  $\mu' : E_{w_1} \to E$  and  $\nu' : E_{w_2} \to E$  are two other  $\mathcal{C}$ -compatible coloured-graph morphisms such that  $\mu'\nu' = \lambda$ . Now,  $\mu$  and  $\mu'$  are coloured-graph morphisms which both have the domain  $E_{w_1}$  forcing  $\mu' = \lambda|_{E_{w_1}} = \mu$ . Similarly,  $\nu' = \lambda|_{E_{[w_1,w_1w_2]}}^* = \nu$ . So  $\mu$  and  $\nu$  are unique coloured-graph morphisms such that  $d(\mu) = w_1$  and  $d(\nu) = w_2$ , and  $\lambda = \mu\nu$ .

#### **2.1** Defining a $BS(2,1)^+$ -graph

In this section we will be constructing a Higher-rank graph, called the  $BS(2,1)^+$ -graph, given a set of coloured-graph morphisms.

**Definition 2.23.** Let  $\mathcal{C} = (\mathcal{C}^0, \mathcal{C}^*, r_{\mathcal{C}}, s_{\mathcal{C}})$  and  $\mathcal{D} = (\mathcal{D}^0, \mathcal{D}^*, r_{\mathcal{D}}, s_{\mathcal{D}})$  be categories. Then  $F = (F^0 : \mathcal{C}^0 \to \mathcal{D}^0, F^* : \mathcal{C}^* \to \mathcal{D}^*)$  is a functor if:

- (i)  $s_{\mathcal{D}}(F^*(f)) = F^0(s_{\mathcal{C}}(f)) \ \forall f \in \mathcal{C}^*$
- (ii)  $r_{\mathcal{C}}(F^*(f)) = F^0(r_{\mathcal{D}}(f)) \ \forall f \in \mathcal{C}^*$
- (iii)  $F^*(fg) = F^*(f)F^*(g) \ \forall f, g \in \mathcal{C}^*$
- (iv)  $F^*(\iota_v) = \iota_{F^0(v)} \ \forall v \in \mathcal{C}^0$

So a functor between categories maps morphisms in  $\mathcal{C}$  to morphisms in  $\mathcal{D}$ , maps objects in  $\mathcal{C}$  to objects in  $\mathcal{D}$ , and preserves the domain and codomain maps, composition, and respects the identity morphisms.

Such an F is also called a *covariant functor*. If we replace (iii) by  $F^*(fg) = F^*(g)F^*(f)$  $\forall f, g \in \mathcal{C}^*$  then we call F a *contravariant functor*.

**Definition 2.24.** Let E be a 2-coloured graph and C be a complete collection of squares in E. For each  $w \in BS(2,1)^+$ , write  $\Lambda^w$  for the set of all C-compatible coloured-graph morphisms  $\lambda : E_w \to E$ . Let  $\Lambda = \bigcup_{w \in BS(2,1)^+} \Lambda^w$ .

**Definition 2.25.** A  $BS(2,1)^+$ -graph is a pair  $(\Lambda_{BS}, d)$ , where  $\Lambda_{BS}$  is a countable category and d is a functor  $d : \Lambda_{BS} \to BS(m,n)^+$  satisfying the Factorisation property: For every  $\lambda \in Mor(\Lambda_{BS})$  and  $w_1, w_2 \in BS(2,1)^+$  with  $d(\lambda) = w_1w_2$ , there are unique elements  $\mu, \nu \in Mor(\Lambda_{BS})$  such that  $\lambda = \mu\nu$ ,  $d(\mu) = w_1$  and  $d(\nu) = w_2$ .

**Theorem 2.26.** [Major Theorem] Fix a 2-coloured graph E and a complete collection of squares C in E. Define  $\Lambda$  as in Definition 2.24, endowed with the structure maps  $r, s : \Lambda \to \Lambda^0$  and with composition as defined in Corollary 2.21. Let  $d : \Lambda \to BS(2,1)^+$  be as defined in Definition 2.4, then  $(\Lambda, d)$  is a  $BS(2,1)^+$ -graph.

## **2.2** Constructing $BS(2,1)^+$ -graphs

**Theorem 2.27.** Fix a 2-coloured graph E and a complete collection of squares C in E. The set  $\Lambda$ , endowed with the structure maps  $r, s : \Lambda \to \Lambda^0$  and with composition as defined in Corollary 2.21, is a category.

Proof. Fix C-compatible coloured-graph morphisms  $\lambda : E_{w_1} \to E, \ \mu : E_{w_2} \to E$ , and  $\nu : E_{w_3} \to E$  with  $s(\lambda) = r(\mu)$  and  $s(\mu) = r(\nu)$ . Then the composition of  $\mu$  and  $\nu$  is a unique C-compatible coloured-graph morphism  $\mu\nu : E_{w_2w_3} \to E$ , such that  $(\mu\nu)|_{E_{w_2}} = \mu$  and  $(\mu\nu)|_{E_{[w_2,w_2w_3]}}^* = \nu$ . We have

$$s(\mu\nu) = \mu\nu(w_2w_3) = \mu\nu|_{E_{[w_2,w_2w_3]}}^*(w_3) = \nu(w_3) = s(\nu), \text{ and}$$
$$r(\mu\nu) = \mu\nu(e) = \mu\nu|_{E_{w_2}}(e) = \mu(e) = r(\mu)$$

so condition (i) holds.

For condition (ii), fix  $x_{\lambda}, x_{\mu}, x_{\nu} \in E^*$  such that  $x_{\lambda}$  traverses  $\lambda$ ,  $x_{\mu}$  traverses  $\mu$  and  $x_{\nu}$  traverses  $\nu$ . Corollary 2.21 gives the composition of  $\lambda$  and  $\mu$  as a unique *C*-compatible coloured-graph morphism  $\lambda \mu$  with  $(\lambda \mu)|_{E_{w_1}} = \lambda$  and  $(\lambda \mu)|_{E_{w_1,w_1w_2}}^* = \mu$ . Then Lemma 2.12 gives that  $x_{\lambda}x_{\mu}$  traverses  $\lambda \mu$ . A repeated application of Corollary 2.21 and Lemma 2.12 on traversals  $x_{\lambda}x_{\mu}$  of  $\lambda \mu$  and  $x_{\nu}$  of  $\nu$  gives

$$x_{\lambda}x_{\mu}x_{\nu} = (x_{\lambda}x_{\mu})x_{\nu}$$
 traverses  $(\lambda\mu)\nu$ 

Similarly,  $x_{\mu}x_{\nu}$  traverses  $\mu\nu$ , and

$$x_{\lambda}x_{\mu}x_{\nu} = x_{\lambda}(x_{\mu}x_{\nu}) \text{ traverses } \lambda(\mu\nu)$$

By Theorem 2.13, we conclude that  $(\lambda \mu)\nu = \lambda(\mu \nu)$ .

Finally, for  $v \in E^0$ , we have previously defined the identity morphism  $\lambda_v : E_e \to E$ by  $\lambda_v(e) = v$ . We have  $r(\lambda_v) = \lambda_v(e) = v$  and  $s(\lambda_v) = \lambda_v(d(\lambda_v)) = \lambda_v(e) = v$ . Now suppose  $r(\mu) = \mu(e) = : v$ , we wish to show that  $\mu = \lambda_v \mu$ . We have  $\mu|_{E_e} : E_e \to E$  with  $\mu|_{E_e}(e) = \mu(e) = v = \lambda_v(e)$ . So  $\lambda_v = \mu|_{E_e}$ . Also,  $\mu|_{E_{[e,w_2]}}^* : E_{[e,w_2]} \to E$ . By definition,  $E_{[e,w_2]} = E_{w_2}$  and  $\mu|_{E_{[e,w_2]}}^*(a) = \mu(ea) = \mu(a)$ . So then  $\mu = \mu|_{E_{[e,w_2]}}^*$ . Remark 2.22 gives  $\mu = \mu|_{E_e}\mu|_{E_{[e,w_2]}}^*$ , hence  $\mu = \lambda_v\mu$ .

For  $s(\nu) = \nu(d(\nu)) = \nu(w_3) := \nu$ , we wish to show  $\nu\lambda_v = \nu$ . Restricting the domain of  $\nu$  gives both  $\nu|_{E_{w_3}} : E_{w_3} \to E$  and  $\nu|_{E_{[w_3,w_3]}}^* : E_{[w_3,w_3]} \to E$ . It is trivial to see that  $\nu|_{E_{w_3}} = \nu$ , and that the subgraph  $E_{[w_3,w_3]}$  of  $E_{w_3}$  is  $E_e$ . Then  $\nu|_{E_{[w_3,w_3]}}^*(e) = \nu(w_3e) = \nu(w_3) = v$ . So  $\nu|_{E_{[w_3,w_3]}}^* = \lambda_v$ . Again, Remark 2.22 gives  $\nu = \nu|_{E_{w_3}}\nu|_{E_{[w_3,w_3]}}^*$ . Thus  $\nu\lambda_v = \nu$ . Condition (iii) holds hence  $\Lambda$  is a category.

**Lemma 2.28.**  $BS(2,1)^+$  is a category with  $Obj(BS(2,1)^+) = \{\star\}$  and  $\iota_{\star} = e$ ,  $Mor(BS(2,1)^+) = BS(2,1)^+$ , maps  $r, s : w \mapsto \star$  for  $w \in BS(2,1)^+$  and composition  $w_1 \cdot w_1 := w_1 w_2$ .

*Proof.* Fix morphisms  $w_1, w_2, w_3 \in BS(2, 1)^+$ . There is only one object in  $BS(2, 1)^+$ , so the source and range of any morphism in  $BS(2, 1)^+$  is precisely the object  $\star$ . It follows that

$$s(w_1w_2) = \star = s(w_2)$$
  

$$r(w_1w_2) = \star = r(w_1) \ \forall w_1, w_2 \in BS(2, 1)^+$$

So condition (i) holds. For condition (ii), we know that  $(w_1w_2)w_3 = w_1(w_2w_3) \forall w_1, w_2, w_3 \in BS(2,1)^+$  given that associativity is inherited from the property that makes  $BS(2,1)^+$  a semigroup. We also have  $r(\iota_{\star}) = r(e) = \star = s(e) = s(\iota_{\star})$  and  $\iota_{\star}w_1 = ew_1 = w_1$  and  $w_1\iota_{\star} = w_1e = w_1 \forall w_1 \in BS(2,1)^+$ , proving condition (iii) for all  $w_1 \in BS(2,1)^+$ . Hence,  $BS(2,1)^+$  is a category.

Proof. of Theorem 2.26

We have that  $\Lambda = (\Lambda^0, \Lambda^*, r_\Lambda, s_\Lambda)$  is a category, directly from Theorem 2.27 and that  $BS(2, 1)^+ = (\{\star\}, BS(2, 1)^+, r_{BS}, s_{BS})$  is a category from Lemma 2.28.

All that remains to show is that  $d : \Lambda \to BS(2,1)^+$  is a functor which satisfies the factorisation property. We have

$$d^{0}: \Lambda^{0} \to \star$$
$$d: \Lambda^{*} \to BS(2,1)^{+}$$

Since  $BS(2,1)^+$  as a category has only one object  $\star$ , d respects  $r_{BS}$  and  $s_{BS}$ 

$$s_{BS}(d(\lambda)) = \star = d^0(s_\Lambda)$$
, and,  
 $r_{BS}(d(\lambda)) = \star = d^0(r_\Lambda)$ 

for all  $\lambda \in \Lambda^*$ .

To see that d respects composition, fix traversals  $x_{\lambda_1}$  of  $\lambda_1$  and  $x_{\lambda_2}$  of  $\lambda_2$ . Then

$$d(\lambda_1 \lambda_2) = d^*(x_{\lambda_1} x_{\lambda_2})$$
  
=  $d^*(x_{\lambda_1}) d^*(x_{\lambda_2})$  by Corollary 2.21  
=  $d(\lambda_1) d^*(\lambda_2)$ 

We must now show that  $d(\lambda_v) = \lambda_{d^0(v)}$  for all  $v \in \Lambda^0$ . We have

$$d(\lambda_v) = e = \iota_\star = \lambda_{d^0(v)} \ \forall v \in \Lambda^0$$

So  $d : \Lambda \to BS(2,1)^+$  is a functor. That d satisfies the factorisation property follows directly from Corollary 2.21. Hence  $(\Lambda, d)$  is a  $BS(2,1)^+$ -graph.

# 3 References

[1] N. Brownlowe, A. Sims, and S. T. Vittadello, *Co-universal C<sup>\*</sup>-algebras associated to generalised graphs*, Israel J. Math. **193** (2013), 399–440.

[2] R. Hazlewood, I. Raeburn, A. Sims, and S. B. G. Webster, *Remarks on some fundamental results about higher-rank graphs and their*  $C^*$ -algebras, Proc. Edinburgh Math. Soc. **56** (2013), 575–597.

[3] A. Kumjian and D. Pask, *Higher rank graph C*<sup>\*</sup>-algebras, New York J. Math. 6 (2000), 1–20.