

THE ERROR TERM IN THE TRUNCATED PERRON FORMULA FOR THE LOGARITHM OF AN L -FUNCTION

STEPHAN RAMON GARCIA, JEFFREY LAGARIAS, AND ETHAN SIMPSON LEE

ABSTRACT. We improve upon the traditional error term in the truncated Perron formula for the logarithm of an L -function. All our constants are explicit.

1. INTRODUCTION

The truncated Perron formula relates the summatory function of an arithmetic function to a contour integral that may be estimated using techniques from complex analysis. Let $F(s) = \sum_{n=1}^{\infty} f(n)n^{-s}$ be absolutely convergent on $\operatorname{Re} s > c_F$; examples include the Riemann zeta-function, Dirichlet L -functions, the Dedekind zeta-function associated to a number field, and Artin L -functions. The truncated Perron formula tells us that if $x > 0$ is not an integer, $T \geq 1$, and $c > c_F$, then

$$\sum_{n \leq x} f(n) = \frac{1}{2\pi i} \int_{c-iT}^{c+iT} F(s) \frac{x^s}{s} ds + O^* \left(\sum_{n=1}^{\infty} \left(\frac{x}{n} \right)^c |f(n)| \min \left\{ 1, \frac{1}{T |\log \frac{x}{n}|} \right\} \right), \quad (1)$$

in which $O^*(g(x)) = h(x)$ means $|h(x)| \leq g(x)$; see [11, Ex. 4.4.15], [8, Ch. 7], [15, Sect. II.2], and [10, Sect. 5.1]. We let T depend on x and let $c = c_F + 1/\log x$, so that $x^c = ex^{c_F}$. A variation of (1) improves the order of the error term by truncating the integral at $\pm T^*$ for an unknown $T^* \in [T, O(T)]$ [3], although this is inconvenient if one must avoid T^* that correspond to the ordinates of non-trivial zeros of $F(s)$. The authors of [3] have also informed us in a personal communication that their paper inherited an unfortunate typo from another paper, so the error term in their variation of the truncated Perron formula could be worse by a factor of $\log x$; this means that our main result (Theorem 1) will be comparable in strength *and* more straightforward to apply when compared against the outcome of their result.

For $\operatorname{Re} s > 1$, the logarithm of the Riemann zeta function $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$ is $\log \zeta(s) = \sum_{n=1}^{\infty} \Lambda(n)(\log n)^{-1}n^{-s}$, in which $\Lambda(n)$ is the von Mangoldt function. The logarithm of a typical L -function is of the form $\sum_{n=1}^{\infty} \Lambda(n)a_n(\log n)^{-1}n^{-s}$, in which the a_n are easily controlled. For example, $|a_n| \leq 1$ for Dirichlet L -functions and $|a_n| \leq d$ for Artin L -functions of degree d ; see [7, Ch. 5]. In these cases, the error term in (1) with $c = 1/\log x$ is on the order of

$$\sum_{n=2}^{\infty} \left(\frac{x}{n} \right)^{\frac{1}{\log x}} \frac{\Lambda(n)}{n \log n} \min \left\{ 1, \frac{1}{T |\log \frac{x}{n}|} \right\} = O \left(\frac{\log x}{T} \right). \quad (2)$$

Granville and Soundararajan used (2) with Dirichlet L -functions to study large character sums [6, (8.1)]. Cho and Kim applied it to Artin L -functions to obtain asymptotic bounds on Dedekind zeta residues [1, Prop. 3.1]. A bilinear relative of (2) appears in Selberg's work on primes in short intervals [14, Lem. 4]. Analogous sums arise with the logarithmic derivative of an L -function [4, p. 106], [12, p. 44].

2010 *Mathematics Subject Classification.* 11M06, 11R42, 11M99, 11S40.

Key words and phrases. Perron formula, L -function, zeta function, Lambert function.

SRG supported by NSF Grant DMS-2054002. ESL thanks the Heilbronn Institute for Mathematical Research for their support. We also thank the referee for their suggestions.

We improve upon (2) asymptotically and explicitly. Moreover, our result has a wide and explicit range of applicability.

Theorem 1. *If $x \geq 3.5$ is a half integer and $T \geq (\log \frac{3}{2})^{-1} > 2.46$, then*

$$\sum_{n=2}^{\infty} \left(\frac{x}{n}\right)^{\frac{1}{\log x}} \frac{\Lambda(n)}{n \log n} \min \left\{ 1, \frac{1}{T |\log \frac{x}{n}|} \right\} \leq \frac{R(x)}{T}, \quad (3)$$

in which

$$R(x) = 40.23 \log \log x + 58.12 + \frac{3.87}{\log x} + \frac{5.22 \log x}{\sqrt{x}} - \frac{1.84}{\sqrt{x}}. \quad (4)$$

The following corollary employs (1) with $T = x$ and $c = 1/\log x$. Since one can use analytic techniques to see the integral below is asymptotic to $\log L(1, \chi)$, one can relate $\log L(1, \chi)$ to a short sum. We hope to do so explicitly in the future.

Corollary 2. *Let $L(s, \chi)$ be an entire Artin L -function of degree d such that*

$$L(s, \chi) = \prod_p \prod_{i=1}^d \left(1 - \frac{\alpha_i(p)}{p^s} \right)^{-1} \quad \text{for } \operatorname{Re} s > 1$$

with $a(p^k) = \alpha_1(p)^k + \cdots + \alpha_d(p)^k$ for prime p . Then with $R(x)$ as in (4)

$$\sum_{1 < n < x} \frac{\Lambda(n) a(n)}{n \log n} = \frac{1}{2\pi i} \int_{\frac{1}{\log x} - ix}^{\frac{1}{\log x} + ix} \frac{x^s}{s} \log L(1 + s, \chi) ds + O^* \left(\frac{d R(x)}{x} \right).$$

2. PRELIMINARIES

Here we establish several lemmas needed for the proof of Theorem 1.

Lemma 3. *If $\sigma > 0$, then $\log \zeta(1 + \sigma) \leq -\log \sigma + \gamma \sigma$.*

Proof. For $s > 1$, we have $\zeta(s) \leq e^{\gamma(s-1)}/(s-1)$ [13, Lem. 5.4]. Let $s = 1 + \sigma$ and take logarithms to obtain the desired result. \square

For real z, w , the equation $z = we^w$ can be solved for w if and only if $z \geq -e^{-1}$. There are two branches for $-e^{-1} \leq z < 0$. The lower branch defines the *Lambert $W_{-1}(z)$ function* [2], which decreases to $-\infty$ as $z \rightarrow 0^-$; see Figure 1a. For $n \geq 6 > 2e$, we define the strictly increasing sequence

$$y_n = \frac{-n}{2} W_{-1} \left(\frac{-2}{n} \right) \quad \text{for } n \geq 6. \quad (5)$$

Lemma 4. *For $n \geq 8$, we have $\frac{2y_n}{\log y_n} = n$ and $y_n \geq \frac{n}{2} \log n$.*

Proof. For $n \geq 6$, the definition of W_{-1} and (5) confirm that $\frac{2y_n}{\log y_n} = n$. Thus, the desired inequality is equivalent to $W_{-1} \left(\frac{-2}{n} \right) \leq -\log n$. Since $f(w) = we^w$ decreases on $(-\infty, -1]$ (Figure 1b) and $-\frac{1}{e} < -\frac{2}{n} < 0$, the desired inequality is equivalent to


$$-\frac{2}{n} \geq f(-\log n) = (-\log n) e^{-\log n} = -\frac{\log n}{n},$$

which holds whenever $\log n \geq 2$. This occurs for $n \geq e^2 \approx 7.38906$. \square

Remark 5. For all $-e^{-1} \leq x < 0$, the bound $W_{-1}(x) \leq \log(-x) - \log(-\log(-x))$ is valid; see [9, (8), (39)]. It follows from this observation and (5) that

$$y_n \geq \frac{n}{2} \left(\log \left(\frac{1}{2} \log \frac{n}{2} \right) + \log n \right),$$


which also implies Lemma 4 for $n \geq 15$.



(A) The two branches of $z = we^w$. The lower branch (blue) is the Lambert function $W_{-1}(z)$, the upper branch (gold) is $W_0(z)$.

(B) $f(w) = we^w$ decreases on $(-\infty, -1]$.

FIGURE 1. Graphs relevant to the construction of the sequence y_n .



(A) $F_1(x)$ is bounded above by C (red).

(B) $F_2(x)$ is bounded above by C (red).

FIGURE 2. The functions $F_1(x)$ and $F_2(x)$ behave erratically.

The next lemma is needed later to handle a few exceptional primes.

Lemma 6. *Let $x > 1$ be a half integer and let $C = \frac{1284699552}{444215525} = 2.89206\dots$*

(a) *Let $p_{-8} < p_{-7} < \dots < p_{-1} < x$ denote the largest eight primes (if they exist) in the interval $(\frac{x}{2}, x)$. We have the sharp bound*

$$F_1(x) = \sum_{1 \leq n \leq 8} \frac{1}{x - p_{-n}} \leq C; \quad (6)$$

see Figure 2a. The corresponding summand in (6) is zero if p_{-n} does not exist.

(b) *Let $x < p_1 < p_2 < \dots < p_8$ denote the smallest eight primes (if they exist) in the interval $(x, \frac{3x}{2})$. We have the sharp bound*

$$F_2(x) = \sum_{1 \leq n \leq 8} \frac{1}{p_n - x} \leq C; \quad (7)$$

see Figure 2b. The corresponding summand in (7) is zero if p_n does not exist.

Proof. (a) If $x \geq 10.5$, then $2, 3, 5 \notin (\frac{x}{2}, x)$. Computation confirms that $F_1(x) \leq F_1(3.5) = \frac{8}{3} = 2.66\dots$ for $x \leq 9.5$. Let $x \geq 10.5$. Then any prime in $(\frac{x}{2}, x)$ is congruent to one of $1, 7, 11, 13, 17, 19, 23, 29 \pmod{30}$. There are finitely many patterns modulo 30 that the $p_{-8}, p_{-7}, \dots, p_{-1}$ may assume. Among these, computation confirms that $F_1(x)$ is maximized if

$$\begin{aligned} p_{-1} &= \lfloor x \rfloor \equiv 19 \pmod{30}, & p_{-5} &= \lfloor x \rfloor - 12 \equiv 7 \pmod{30}, \\ p_{-2} &= \lfloor x \rfloor - 2 \equiv 17 \pmod{30}, & p_{-6} &= \lfloor x \rfloor - 18 \equiv 1 \pmod{30}, \\ p_{-3} &= \lfloor x \rfloor - 6 \equiv 13 \pmod{30}, & p_{-7} &= \lfloor x \rfloor - 20 \equiv 29 \pmod{30}, \\ p_{-4} &= \lfloor x \rfloor - 8 \equiv 11 \pmod{30}, & p_{-8} &= \lfloor x \rfloor - 26 \equiv 23 \pmod{30}, \end{aligned}$$




FIGURE 3. Proof of Lemma 7.

which yields the desired upper bound C . This prime pattern first occurs for $x = 88819.5$; see <https://oeis.org/A022013>.

(b) If $x \geq 5.5$, then $2, 3, 5 \notin (x, \frac{3x}{2})$. Observe that $F_2(x) \leq 2$ for $x \leq 4.5$ (attained at $x = 1.5, 2.5, 4.5$). Let $x \geq 5.5$. As in (a), any prime in $(x, \frac{3x}{2})$ is congruent to one of $1, 7, 11, 13, 17, 19, 23, 29 \pmod{30}$. Then $F_2(x)$ is maximized if

$$\begin{aligned} p_1 &= \lceil x \rceil \equiv 11 \pmod{30}, & p_5 &= \lceil x \rceil + 12 \equiv 23 \pmod{30}, \\ p_2 &= \lceil x \rceil + 2 \equiv 13 \pmod{30}, & p_6 &= \lceil x \rceil + 18 \equiv 29 \pmod{30}, \\ p_3 &= \lceil x \rceil + 6 \equiv 17 \pmod{30}, & p_7 &= \lceil x \rceil + 20 \equiv 1 \pmod{30}, \\ p_4 &= \lceil x \rceil + 8 \equiv 19 \pmod{30}, & p_8 &= \lceil x \rceil + 26 \equiv 7 \pmod{30}. \end{aligned}$$

which yields the desired upper bound C . Although this prime pattern occurs for $x = 10.5$, not all eight primes lie in $(x, \frac{3}{2}x)$. The first admissible value is $x = 15760090.5$; see <https://oeis.org/A022011>. \square

We also need an elementary estimate on k th powers in intervals.

Lemma 7. *Let $X > 1$ be a noninteger, $h > 1$, and $k \geq 2$.*

(a) *There are at most $N_k + 1$ perfect k th powers in $[X, X+h)$, in which $N_k \leq \frac{h}{k\sqrt[k]{X}}$.*

(b) *The shortest gap between k th powers in $[X, X+h)$ (if they exist) is $G_k \geq k\sqrt[k]{X}$.*

Proof. We may assume that X is so large that $N_k \geq 1$. Let $m = \lceil X^{\frac{1}{k}} \rceil$ so that m^k is the first k th power larger than X . Consider the gaps g_1, g_2, \dots, g_{N_k} between the N_k consecutive k th powers in $[X, X+h)$; see Figure 3. Then

$$G_k = \min\{g_1, g_2, \dots, g_{N_k}\} = g_1 = (m+1)^k - m^k \geq km^{k-1} \geq kX^{\frac{k-1}{k}} \geq k\sqrt[k]{X}.$$

The desired inequality follows since $N_k G_k \leq g_1 + g_2 + \dots + g_{N_k} \leq h$. \square

Finally, we need an estimate on the n th harmonic number $H_n = \sum_{j=1}^n \frac{1}{j}$:

$$\frac{1}{2n + \frac{2}{5}} < H_n - \log n - \gamma < \frac{1}{2n + \frac{1}{3}} \leq \frac{3}{7} \quad \text{for } n \geq 1, \quad (8)$$

in which γ is the Euler–Mascheroni constant [16]. We require the upper bound

$$\begin{aligned} \sum_{\ell=1}^n \frac{1}{2\ell-1} &= \left(1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{2n}\right) - \left(\frac{1}{2} + \frac{1}{4} + \dots + \frac{1}{2n}\right) = H_{2n} - \frac{1}{2}H_n \\ &\leq \left(\log 2n + \gamma + \frac{1}{4n + \frac{1}{3}}\right) - \frac{1}{2} \left(\log n + \gamma + \frac{1}{2n + \frac{2}{5}}\right) \\ &\leq \frac{1}{2} \log n + \frac{1}{2} \gamma + \log 2 + \frac{1}{4n + \frac{1}{3}} - \frac{1}{4n + \frac{4}{5}} \\ &= \frac{1}{2} \log n + \frac{1}{2} \gamma + \log 2 + \frac{7}{240n^2 + 68n + 4} \\ &\leq \frac{1}{2} \log n + \frac{1}{2} \gamma + \log 2 + \frac{7}{312} \quad \text{for } n \geq 1. \end{aligned} \quad (9)$$




FIGURE 4. Graphs relevant to the derivation of (11).

3. PROOF OF THEOREM 1

In what follows, $x \in \mathbb{N} + \frac{1}{2} = \{\frac{3}{2}, \frac{5}{2}, \frac{7}{2}, \dots\}$ and $c = \frac{1}{\log x}$. Minor improvements below are possible; these were eschewed in favor of a final estimate of simple shape.

3.1. When n is very far from x . Suppose that $n \leq \frac{x}{2}$ or $\frac{3x}{2} \geq n$. Then $\log \frac{x}{n} \leq -\log \frac{3}{2}$ or $\log \frac{3}{2} < \log 2 \leq \log \frac{x}{n}$, so $|\log \frac{x}{n}| \geq \log \frac{3}{2}$. If $T \geq (\log \frac{3}{2})^{-1} > 2.46$, then

$$\min \left\{ 1, \frac{1}{T |\log \frac{x}{n}|} \right\} \leq \frac{1}{T |\log \frac{x}{n}|} \leq \frac{1}{T \log \frac{3}{2}}.$$

For such T , the previous inequality and Lemma 3 imply (recall that $c = \frac{1}{\log x}$)

$$\begin{aligned} \sum_{\substack{n \leq \frac{x}{2} \text{ or} \\ n \geq \frac{3x}{2}}} \left(\frac{x}{n} \right)^c \frac{\Lambda(n)}{n \log n} \min \left\{ 1, \frac{1}{T |\log \frac{x}{n}|} \right\} &\leq x^c \sum_{\substack{n \leq \frac{x}{2} \text{ or} \\ n \geq \frac{3x}{2}}} \frac{1}{n^{1+c}} \left(\frac{\Lambda(n)}{\log n} \right) \left(\frac{1}{T \log \frac{3}{2}} \right) \\ &\leq \frac{x^c}{T \log \frac{3}{2}} \log \zeta(1+c) \\ &\leq \frac{x^c}{T \log \frac{3}{2}} (-\log c + \gamma c) \\ &= \frac{e}{T \log \frac{3}{2}} \left(\log \log x + \frac{\gamma}{\log x} \right). \end{aligned} \quad (10)$$

3.2. Reduction to a sum over prime powers. Suppose that $\frac{x}{2} < n < \frac{3x}{2}$. Let $z = 1 - \frac{n}{x}$ and observe that $|z| < \frac{1}{2}$. Then

$$\log \frac{x}{n} = -\log(1-z) = z \left(-\frac{\log(1-z)}{z} \right),$$

in which the function in parentheses is positive and achieves its minimum value $2 \log \frac{3}{2} = 0.81093\dots$ on $|z| < \frac{1}{2}$ at its left endpoint $-\frac{1}{2}$; see Figure 4a. Then

$$|\log(1-z)| > \left(2 \log \frac{3}{2} \right) |z| \quad \text{for } |z| < \frac{1}{2}, \quad (11)$$

whose validity is illustrated in Figure 4b. Therefore,

$$\left| \log \frac{x}{n} \right| > \left(2 \log \frac{3}{2} \right) \left| 1 - \frac{n}{x} \right| \quad \text{for } \frac{x}{2} < n < \frac{3x}{2}, \quad (12)$$

and hence

$$\begin{aligned}
& \sum_{\frac{x}{2} < n < \frac{3x}{2}} \left(\frac{x}{n}\right)^c \frac{\Lambda(n)}{n \log n} \min \left\{ 1, \frac{1}{T |\log \frac{x}{n}|} \right\} \\
& \leq \frac{x^c}{T} \sum_{\frac{x}{2} < n < \frac{3x}{2}} \left(\frac{\Lambda(n)}{n^{1+c} \log n} \right) \left(\frac{1}{|\log \frac{x}{n}|} \right) \\
& \leq \frac{x^c}{T} \sum_{\frac{x}{2} < n < \frac{3x}{2}} \left(\frac{\Lambda(n)}{n^{1+c} \log n} \right) \frac{1}{(2 \log \frac{3}{2}) |1 - \frac{n}{x}|} \quad (\text{by (12)}) \\
& \leq \frac{x^c}{T(2 \log \frac{3}{2})} \sum_{\frac{x}{2} < n < \frac{3x}{2}} \frac{2}{x} \left(\frac{\Lambda(n)}{n^c \log n} \right) \frac{1}{|1 - \frac{n}{x}|} \quad (\text{since } \frac{x}{2} < n) \\
& \leq \frac{x^c}{T \log \frac{3}{2}} \sum_{\frac{x}{2} < n < \frac{3x}{2}} \left(\frac{\Lambda(n)}{n^c \log n} \right) \frac{1}{|x - n|} \\
& \leq \frac{x^c}{T \log \frac{3}{2}} \sum_{\frac{x}{2} < p^k < \frac{3x}{2}} \left(\frac{\log p}{(p^k)^c k \log p} \right) \frac{1}{|x - p^k|} \quad (\text{def. of } \Lambda) \\
& \leq \frac{e}{T \log \frac{3}{2}} \sum_{\frac{x}{2} < p^k < \frac{3x}{2}} \frac{1}{k |x - p^k|}, \quad (\text{since } c = \frac{1}{\log x}) \quad (13)
\end{aligned}$$

in which the final two sums run over all prime powers p^k in the stated interval.

The remainder of the proof uses ideas from [5, Lem. 2] to estimate

$$\sum_{\frac{x}{2} < p^k < \frac{3x}{2}} \frac{1}{k |x - p^k|} = \underbrace{\sum_{\frac{x}{2} < p < \frac{3x}{2}} \frac{1}{|x - p|}}_{S_{\text{prime}}(x)} + \underbrace{\sum_{\substack{\frac{x}{2} < p^k < \frac{3x}{2} \\ k \geq 2}} \frac{1}{k |x - p^k|}}_{S_{\text{power}}(x)}. \quad (14)$$

3.3. The sum over primes. First observe that

$$S_{\text{prime}}(x) \leq \underbrace{\sum_{\frac{x}{2} < p < x} \frac{1}{x - p}}_{S_{\text{prime}}^-(x)} + \underbrace{\sum_{x < p < \frac{3x}{2}} \frac{1}{p - x}}_{S_{\text{prime}}^+(x)}.$$

We require the Brun–Titchmarsh theorem (see Montgomery–Vaughan [10, Cor. 2]):

$$\pi(X + Y) - \pi(X) \leq \frac{2Y}{\log Y} \quad \text{where } \pi(x) = \sum_{p \leq x} 1, \quad X > 0, \text{ and } Y > 1. \quad (15)$$

3.3.1. The lower sum over primes. Let $p_{-k} < p_{-(k-1)} < \cdots < p_{-2} < p_{-1}$ be the primes in $(\frac{x}{2}, x)$; note that $k \leq \frac{x}{2}$. Apply (15) with $X = x - y_n$ and $Y = y_n$ to get

$$0 \leq \pi(x) - \pi(x - y_n) \leq \frac{2y_n}{\log y_n} = n \quad \text{for } 6 \leq n \leq k$$

by Lemma 4, so $(x - y_n, x]$ contains at most n primes. Thus, $p_{-(n+1)} \leq x - y_n$ and

$$\frac{1}{x - p_{-(n+1)}} \leq \frac{1}{y_n} \quad \text{for } 6 \leq n \leq k - 1. \quad (16)$$

Then Lemma 4, which requires $k \geq 8$, and the integral test provide

$$\sum_{\frac{x}{2} < p < x} \frac{1}{x - p} = \sum_{1 \leq n \leq 8} \frac{1}{x - p_{-n}} + \sum_{9 \leq n \leq k} \frac{1}{x - p_{-n}}$$

$$\begin{aligned}
&\leq F_1(x) + \sum_{8 \leq n \leq k-1} \frac{1}{y_n} && \text{(by (6) and (16))} \\
&\leq C + 2 \sum_{7 < n \leq \frac{x}{2}} \frac{1}{n \log n} && \text{(by Lemma 4)} \\
&\leq C + 2 \log \log x,
\end{aligned}$$

which is valid for $k \leq 7$ since Lemma 6a shows that the sum is majorized by C .

3.3.2. The upper sum over primes. Let $p_1 < p_2 < \dots < p_k$ denote the primes in $(x, \frac{3x}{2})$ and note that $k \leq \frac{x}{2}$. Then (15) with $X = x$ and $Y = y_n$ ensures that

$$0 \leq \pi(x + y_n) - \pi(x) \leq \frac{2y_n}{\log y_n} = n \quad \text{for } 6 \leq n \leq k$$

by Lemma 4, so $(x, x + y_n]$ contains at most n primes. Thus, $p_{n+1} \geq x + y_n$ and

$$\frac{1}{p_{n+1} - x} \leq \frac{1}{y_n} \quad \text{for } 6 \leq n \leq k. \quad (17)$$

An argument similar to that above reveals that

$$\sum_{x < p < \frac{3x}{2}} \frac{1}{p - x} \leq \sum_{1 \leq n \leq 8} \frac{1}{p_n - x} + \sum_{9 \leq n \leq k} \frac{1}{p_n - x} \leq C + 2 \log \log x.$$

3.3.3. Final bound over primes. For $x \in \mathbb{N} + \frac{1}{2}$, the previous inequalities yield

$$S_{\text{prime}}(x) = \sum_{\frac{x}{2} < p < \frac{3x}{2}} \frac{1}{|x - p|} \leq 2C + 4 \log \log x. \quad (18)$$

3.4. The sum over prime powers. We now majorize

$$S_{\text{power}}(x) = \sum_{\substack{\frac{x}{2} < p^k < \frac{3x}{2} \\ k \geq 2}} \frac{1}{k|x - p^k|}.$$

3.4.1. Initial reduction. To bound $S_{\text{power}}(x)$ it suffices to majorize

$$S_{\text{sqf}}(x) = \sum_{\substack{\frac{x}{2} < n^k < \frac{3x}{2} \\ k \geq 2 \\ n \geq 2 \text{ sq. free}}} \frac{1}{k|x - n^k|}, \quad (19)$$

in which the prime powers p^k are replaced with the powers n^k of square free $n \geq 2$. The square-free restriction ensures that powers such as $2^6 = (2^2)^3 = (2^3)^2$ are not counted multiple times in (19). If $\frac{x}{2} < n^k < \frac{3x}{2}$ and $k \geq 2$, then (since $n \geq 2$)

$$k \leq \frac{\log \frac{3x}{2}}{\log 2} \leq \lfloor 2.4 \log x \rfloor \quad \text{for } x \geq 3.5. \quad (20)$$

3.4.2. Nearest-power sets. The largest contributions to $S_{\text{sqf}}(x)$ come from the powers closest to x . We handle those summands separately and split the sum (19) accordingly. For each $k \geq 2$, the inequalities $\lfloor x^{\frac{1}{k}} \rfloor^k < x < \lceil x^{\frac{1}{k}} \rceil^k$ exhibit the two k th powers nearest to x . Define

$$\mathcal{N}_k \subseteq \{ \lfloor x^{\frac{1}{k}} \rfloor^k, \lceil x^{\frac{1}{k}} \rceil^k \} \quad (21)$$

according to the following rules:

- \mathcal{N}_k contains $\lfloor x^{\frac{1}{k}} \rfloor^k$ if it is square free and belongs to $(\frac{x}{2}, \frac{3x}{2})$, and
- \mathcal{N}_k contains $\lceil x^{\frac{1}{k}} \rceil^k$ if it is square free and belongs to $(\frac{x}{2}, \frac{3x}{2})$.

Consequently, \mathcal{N}_k , if nonempty, contains only powers that satisfy the restrictions in (19). The square-free condition ensures that $\mathcal{N}_j \cap \mathcal{N}_k = \emptyset$ for $j \neq k$.

Write $S_{\text{sqf}}(x) = S_{\text{near}}(x) + S_{\text{far}}(x)$, in which

$$S_{\text{near}}(x) = \sum_{\substack{\frac{x}{2} < n^k < \frac{3x}{2} \\ k \geq 2 \\ n \geq 2 \text{ sq. free} \\ n^k \in \mathcal{N}_k}} \frac{1}{k|x - n^k|} \quad \text{and} \quad S_{\text{far}}(x) = \sum_{\substack{\frac{x}{2} < n^k < \frac{3x}{2} \\ k \geq 2 \\ n \geq 2 \text{ sq. free} \\ n^k \notin \mathcal{N}_k}} \frac{1}{k|x - n^k|}. \quad (22)$$

3.4.3. *Near Sum.* For $x \geq 3.5$, a nearest-neighbor overestimate provides

$$\begin{aligned} S_{\text{near}}(x) &= \sum_{\substack{\frac{x}{2} < n^k < \frac{3x}{2} \\ k \geq 2 \\ n \geq 2 \text{ sq. free} \\ n^k \in \mathcal{N}_k}} \frac{1}{k|x - n^k|} & \quad (\text{by (22)}) \\ &= \sum_{k=2}^{\lfloor 2.4 \log x \rfloor} \sum_{m \in \mathcal{N}_k} \frac{1}{k|x - m|} & \quad (\text{by (20)}) \\ &\leq \frac{1}{2} \sum_{k=2}^{\lfloor 2.4 \log x \rfloor} \sum_{m \in \mathcal{N}_k} \frac{1}{|x - m|} & \quad (\text{since } k \geq 2) \\ &\leq \frac{1}{2} \sum_{j=0}^{\lfloor 2.4 \log x \rfloor - 2} \left(\frac{1}{x - (\lfloor x \rfloor - j)} + \frac{1}{(\lceil x \rceil + j) - x} \right) & \quad (\text{see below}) \\ &\leq \frac{1}{2} \sum_{\ell=1}^{\lfloor 2.4 \log x \rfloor - 1} \frac{2}{\ell - \frac{1}{2}} < 2 \sum_{\ell=1}^{\lfloor 2.4 \log x \rfloor} \frac{1}{2\ell - 1} \\ &\leq \log(\lfloor 2.4 \log x \rfloor) + \gamma + 2 \log 2 + \frac{14}{312} & \quad (\text{by (9)}) \\ &< \log \log x + \gamma + 2 \log 2 + \log 2.4 + \frac{7}{156}. \end{aligned} \quad (23)$$

Let us elaborate on a crucial step above. Consider the at most $\lfloor 2 \log x \rfloor - 1$ pairs of values $|x - m|$ that arise as m ranges over each \mathcal{N}_k with $2 \leq k \leq \lfloor 2 \log x \rfloor$ (since $\mathcal{N}_j(x) \cap \mathcal{N}_k = \emptyset$ for $j \neq k$, no m appears more than once). Replace these values with the absolute deviations of x from its $2 \times (\lfloor 2 \log x \rfloor - 1)$ nearest neighbors $\lfloor x \rfloor - j$ (to the left) and $\lceil x \rceil + j$ (to the right), in which $0 \leq j \leq \lfloor 2 \log x \rfloor - 2$. Since $x \in \mathbb{N} + \frac{1}{2}$, these deviations are of the form $\ell - \frac{1}{2}$ for $1 \leq \ell \leq \lfloor 2 \log x \rfloor - 1$.

3.4.4. *Splitting the second sum.* From (22), the second sum in question is

$$S_{\text{far}}(x) = \sum_{\substack{\frac{x}{2} < n^k < \frac{3x}{2} \\ k \geq 2 \\ n \geq 2 \text{ sq. free} \\ n^k \notin \mathcal{N}_k}} \frac{1}{k|x - n^k|} \leq \sum_{\substack{\frac{x}{2} < n^k < \frac{3x}{2} \\ n, k \geq 2 \\ n^k \notin \mathcal{N}_k}} \frac{1}{k|x - n^k|} = S_{\text{far}}^-(x) + S_{\text{far}}^+(x),$$

in which

$$S_{\text{far}}^-(x) = \sum_{k \geq 2} \sum_{\substack{\frac{x}{2} < n^k < x \\ n \geq 2, n^k \notin \mathcal{N}_k}} \frac{1}{k|x - n^k|} \quad \text{and} \quad S_{\text{far}}^+(x) = \sum_{k \geq 2} \sum_{\substack{x < n^k < \frac{3x}{2} \\ n \geq 2, n^k \notin \mathcal{N}_k}} \frac{1}{k|x - n^k|}. \quad (24)$$

For $k \geq 2$, Lemma 7 with $X = h = \frac{x}{2}$, then with $X = x$ and $h = \frac{x}{2}$, implies that

$$G_k^- \geq \frac{k\sqrt{x}}{\sqrt{2}}, \quad N_k^- \leq \frac{\sqrt{x}}{2\sqrt{2}}, \quad \text{and} \quad G_k^+ \geq k\sqrt{x}, \quad N_k^+ \leq \frac{\sqrt{x}}{4} \quad (25)$$



FIGURE 5. Analysis of k th powers in $[\frac{x}{2}, \frac{3x}{2}]$, in which $m_- = \lfloor x^{1/k} \rfloor$ and $m_+ = \lceil x^{1/k} \rceil$ are excluded from consideration. There are at most N_k^- admissible k th powers in $[\frac{x}{2}, x)$, with minimal gap size G_k^- , and at most N_k^+ admissible k th powers in $[x, \frac{3x}{2})$, with minimal gap size G_k^+ .

are admissible in Figure 5. For $1 \leq j \leq N_k^-$ and $1 \leq j \leq N_k^+$, respectively,

$$|x - (m_- - j)^k| \geq \frac{jk\sqrt{x}}{\sqrt{2}} \quad \text{and} \quad |x - (m_+ + j)^k| \geq jk\sqrt{x}.$$

Let $N_k^\pm \geq 1$, since otherwise the corresponding sum estimated below is zero. Then

$$\sum_{\substack{\frac{x}{2} < n^k < x \\ n \geq 2, n^k \notin \mathcal{N}_k}} \frac{1}{k|x - n^k|} = \sum_{j=1}^{N_k^-} \frac{1}{k|x - (m_- - j)^k|} \leq \frac{\sqrt{2}H_{N_k^-}}{k^2\sqrt{x}} \quad (26)$$

and

$$\sum_{\substack{x < n^k < \frac{3x}{2} \\ n \geq 2, n^k \notin \mathcal{N}_k}} \frac{1}{k|x - n^k|} = \sum_{j=1}^{N_k^+} \frac{1}{k|x - (m_+ + j)^k|} \leq \frac{H_{N_k^+}}{k^2\sqrt{x}}. \quad (27)$$

Therefore,

$$\begin{aligned} S_{\text{far}}^-(x) &= \sum_{k \geq 2} \sum_{\substack{\frac{x}{2} < n^k < x \\ n \geq 2, n^k \notin \mathcal{N}_k}} \frac{1}{k|x - n^k|} \leq \frac{\sqrt{2}}{\sqrt{x}} \sum_{k \geq 2} \frac{H_{N_k^-}}{k^2} \quad (\text{by (24) and (26)}) \\ &\leq \frac{\sqrt{2}}{\sqrt{x}} \left(\frac{1}{2} \log x - \frac{3}{2} \log 2 + \gamma + \frac{3}{7} \right) \sum_{k \geq 2} \frac{1}{k^2} \quad (\text{by (8) and (25)}) \\ &= \frac{\pi^2 - 6}{3\sqrt{2x}} \left(\frac{1}{2} \log x - \frac{3}{2} \log 2 + \gamma + \frac{3}{7} \right) \quad (\text{since } \zeta(2) - 1 = \frac{\pi^2 - 6}{6}) \quad (28) \end{aligned}$$

and

$$\begin{aligned} S_{\text{far}}^+(x) &= \sum_{k \geq 2} \sum_{\substack{x < n^k < \frac{3x}{2} \\ n \geq 2, n^k \notin \mathcal{N}_k}} \frac{1}{k|x - n^k|} \leq \frac{1}{\sqrt{x}} \sum_{k \geq 2} \frac{H_{N_k^+}}{k^2} \quad (\text{by (24) and (27)}) \\ &\leq \frac{1}{\sqrt{x}} \left(\frac{1}{2} \log x - 2 \log 2 + \gamma + \frac{3}{7} \right) \sum_{k \geq 2} \frac{1}{k^2} \quad (\text{by (8) and (25)}) \\ &= \frac{\pi^2 - 6}{6\sqrt{x}} \left(\frac{1}{2} \log x - 2 \log 2 + \gamma + \frac{3}{7} \right) \quad (\text{since } \zeta(2) - 1 = \frac{\pi^2 - 6}{6}) \quad (29) \end{aligned}$$

3.4.5. *Final prime-power estimate.* Using (23), (28), and (29), we can bound

$$S_{\text{power}}(x) \leq S_{\text{sqf}}(x) = S_{\text{near}}(x) + S_{\text{far}}^-(x) + S_{\text{far}}^+(x).$$

We postpone doing this explicitly until the finale below.

3.5. Conclusion. For $x \geq 3.5$, with $T \geq (\log \frac{3}{2})^{-1}$, the sum (3) is bounded by

$$\begin{aligned}
& \underbrace{\frac{e}{T \log \frac{3}{2}} \left(\log \log x + \frac{\gamma}{\log x} \right)}_{\text{by (10)}} + \underbrace{\frac{e}{T \log \frac{3}{2}} \sum_{\frac{x}{2} < p^k < \frac{3x}{2}} \frac{1}{k|x - p^k|}}_{\text{by (13)}} \\
& \leq \frac{e}{T \log \frac{3}{2}} \left(\log \log x + \frac{\gamma}{\log x} \right) + \frac{e}{T \log \frac{3}{2}} \underbrace{(S_{\text{prime}}(x) + S_{\text{sqf}}(x))}_{\text{by (14) and (19)}} \\
& \leq \frac{e}{T \log \frac{3}{2}} \left(\log \log x + \frac{\gamma}{\log x} \right) + \frac{e}{T \log \frac{3}{2}} \left[\underbrace{2C + 4 \log \log x}_{S_{\text{prime}}(x) \text{ bounded by (18)}} \right. \\
& \quad + \underbrace{\left(\log \log x + \gamma + 2 \log 2 + \log 2.4 + \frac{7}{156} \right)}_{S_{\text{near}}(x) \text{ bounded by (23)}} + \underbrace{\frac{\pi^2 - 6}{3\sqrt{2x}} \left(\frac{1}{2} \log x - \frac{3}{2} \log 2 + \gamma + \frac{3}{7} \right)}_{S_{\text{far}}^-(x) \text{ bounded by (28)}} \\
& \quad \left. + \underbrace{\frac{\pi^2 - 6}{6\sqrt{x}} \left(\frac{1}{2} \log x - 2 \log 2 + \gamma + \frac{3}{7} \right)}_{S_{\text{far}}^+(x) \text{ bounded by (29)}} \right] \\
& < \frac{1}{T} \left(40.22465 \log \log x + 58.11106 + \frac{3.86972}{\log x} + \frac{5.21918 \log x}{\sqrt{x}} - \frac{1.85268}{\sqrt{x}} \right). \quad \square
\end{aligned}$$

REFERENCES

1. P. J. Cho and H. H. Kim, *Extreme residues of Dedekind zeta functions*, Math. Proc. Cambridge Philos. Soc. **163** (2017), no. 2, 369–380. MR 3682635
2. R. M. Corless, G. H. Gonnet, D. E. G. Hare, D. J. Jeffrey, and D. E. Knuth, *On the Lambert W function*, Adv. Comput. Math. **5** (1996), no. 4, 329–359. MR 1414285
3. M. Cully-Hugill and D. R. Johnston, *on the error term in the explicit formula of Riemann–von Mangoldt*, Int. J. Number Theory (2022).
4. H. Davenport, *Multiplicative number theory*, third ed., Graduate Texts in Mathematics, vol. 74, Springer-Verlag, New York, 2000, Revised and with a preface by Hugh L. Montgomery. MR 1790423
5. D. A. Goldston, *On a result of Littlewood concerning prime numbers. II*, Acta Arith. **43** (1983), no. 1, 49–51. MR 730847
6. A. Granville and K. Soundararajan, *Large character sums*, J. Amer. Math. Soc. **14** (2001), no. 2, 365–397. MR 1815216
7. H. Iwaniec and E. Kowalski, *Analytic number theory*, American Mathematical Society Colloquium Publications, vol. 53, American Mathematical Society, Providence, RI, 2004. MR 2061214
8. D. Koukoulopoulos, *The distribution of prime numbers*, Graduate Studies in Mathematics, vol. 203, American Mathematical Society, Providence, RI, [2019] ©2019. MR 3971232
9. L. Lóczy, *Guaranteed- and high-precision evaluation of the Lambert W function*, Appl. Math. Comput. **433** (2022), Paper No. 127406, 22. MR 4456416
10. H. L. Montgomery and R. C. Vaughan, *The large sieve*, Mathematika **20** (1973), 119–134. MR 374060
11. M. R. Murty, *Problems in analytic number theory*, Graduate Texts in Mathematics, Springer New York, 2008.
12. S. J. Patterson, *An introduction to the theory of the Riemann zeta-function*, Cambridge Studies in Advanced Mathematics, vol. 14, Cambridge University Press, Cambridge, 1988. MR 933558
13. O. Ramaré, *An explicit density estimate for Dirichlet L-series*, Math. Comp. **85** (2016), no. 297, 325–356. MR 3404452
14. A. Selberg, *On the normal density of primes in small intervals, and the difference between consecutive primes*, Arch. Math. Naturvid. **47** (1943), no. 6, 87–105. MR 12624

15. G. Tenenbaum, *Introduction to analytic and probabilistic number theory*, Cambridge Studies in Advanced Mathematics, vol. 46, Cambridge University Press, Cambridge, 1995, Translated from the second French edition (1995) by C. B. Thomas. MR 1342300
16. L. Tóth and S. Mare, *E* 3432, Amer. Math. Monthly **98** (1991), no. 3, 264.

DEPARTMENT OF MATHEMATICS AND STATISTICS, POMONA COLLEGE, 610 N. COLLEGE AVE., CLAREMONT, CA 91711, USA

Email address: `stephan.garcia@pomona.edu`

URL: `http://pages.pomona.edu/~sg064747`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, 530 CHURCH STREET, ANN ARBOR, MI 48109, USA

Email address: `lagarias@umich.edu`

URL: `https://dept.math.lsa.umich.edu/~lagarias`

UNIVERSITY OF BRISTOL, SCHOOL OF MATHEMATICS, FRY BUILDING, WOODLAND ROAD, BRISTOL, BS8 1UG

Email address: `ethan.lee@bristol.ac.uk`

URL: `https://sites.google.com/view/ethansleemath/home`