

Baireness of the space of pointwise stabilizing functions of the first Baire class

Alexander V. Osipov

*Krasovskii Institute of Mathematics and Mechanics,
Ural Federal University, Ural State University of Economics, Yekaterinburg, Russia*

Abstract

A topological space X is *Baire* if the Baire Category Theorem holds for X , i.e., the intersection of any sequence of open dense subsets of X is dense in X . In this paper, we have obtained that the space $B_1^{st}(X)$ of pointwise stabilizing Baire-one functions is Baire if the space $B_1(X)$ of Baire-one functions is so. This answers a question posed recently by T. Banach and S. Gabrielyan.

Keywords: stable convergence, stable Baire class, Baire property, function space, Baire-one function

2010 MSC: 54C08, 54H05, 26A21

1. Introduction

For a topological space X , we denote by $C_p(X)$ the family $C(X)$ of all continuous function from X to the real line \mathbb{R} endowed with the topology of pointwise convergence.

A real-valued function f on a space X is a *Baire-one function* (or a *function of the first Baire class*) if f is a pointwise limit of a sequence of continuous functions on X . Let $B_1(X)$ denote the family of all Baire-one real-valued functions on a space X endowed with the topology of pointwise convergence.

We say that a sequence $\{f_n\}_{n \in \omega} \subseteq Y^X$ *stable (or discretely) converges* to a function $f \in Y^X$ if for every $x \in X$ the set $\{n \in \omega : f_n(x) \neq f(x)\}$ is finite. Let $B_1^{st}(X)$ be the family of all functions $f : X \rightarrow \mathbb{R}$ which are limits of sequence $\{f_n\}_{n \in \omega} \subseteq C_p(X)$ stably converging to f . Functions in the

Email address: OAB@list.ru (Alexander V. Osipov)

family $B_1^{st}(X)$ are called the *functions of stable first Baire class (or pointwise stabilizing functions of the first Baire class)*.

A space is *meager* (or *of the first Baire category*) if it can be written as a countable union of closed sets with empty interior. A topological space X is *Baire* if the Baire Category Theorem holds for X , i.e., the intersection of any sequence of open dense subsets of X is dense in X . Clearly, if X is Baire, then X is not meager.

Being a Baire space is an important topological property for a space and it is therefore natural to ask when function spaces are Baire. The Baire property for continuous mappings was first considered in [6]. Then a paper [7] appeared, where various aspects of this topic were considered. In [7], necessary and, in some cases, sufficient conditions on a space X were obtained under which the space $C_p(X)$ is Baire. The problem about a characterization of Baireness for $C_p(X)$ was solved independently by Pytkeev [3], Tkachuk [4] and van Douwen [5].

Theorem 1.1. (*Pytkeev-Tkachuk-van Douwen*) *The space $C_p(X)$ is Baire if, and only if, every pairwise disjoint sequence of non-empty finite subsets of X has a strongly discrete subsequence.*

A collection \mathcal{G} of subsets of X is *discrete* if each point of X has a neighborhood meeting at most one element of \mathcal{G} , and is *strongly discrete* if for each $G \in \mathcal{G}$ there is an open superset U_G of G such that $\{U_G : G \in \mathcal{G}\}$ is discrete.

A Coz_δ -subset of X containing x is called a *Coz_δ neighborhood* of x .

Definition 1.2. [9] A set $A \subseteq X$ is called *strongly Coz_δ -disjoint*, if there is a pairwise disjoint collection $\{F_a : F_a \text{ is a } Coz_\delta \text{ neighborhood of } a, a \in A\}$ such that $\{F_a : a \in A\}$ is *completely Coz_δ -additive*, i.e. $\bigcup_{b \in B} F_b \in Coz_\delta$ for each $B \subseteq A$.

A disjoint sequence $\{\Delta_n : n \in \mathbb{N}\}$ of (finite) sets is *strongly Coz_δ -disjoint* if the set $\bigcup\{\Delta_n : n \in \mathbb{N}\}$ is strongly Coz_δ -disjoint.

In [9], we have obtained a characterization when $B_1(X)$ is Baire.

Theorem 1.3. *Let X be a topological space. The following assertions are equivalent:*

1. $B_1(X)$ is Baire;
2. every pairwise disjoint sequence of non-empty finite subsets of X has a strongly Coz_δ -disjoint subsequence.

In [2], T.Banakh and S.Gabrielyan considers the following question (Problem 6.10 and Problem 9.11): *Is there a topological space X such that $B_1(X)$ is Baire and $B_1^{st}(X)$ is meager?*

In this paper, we get the following theorem which answers the Banakh-Gabrielyan question.

Theorem 1.4. *Let X be a topological space. The following assertions are equivalent:*

1. $B_1(X)$ is Baire;
2. $B_1^{st}(X)$ is Baire;
3. *Every pairwise disjoint sequence $\{\Delta_n : n \in \mathbb{N}\}$ of non-empty finite subsets of X has a subsequence $\{\Delta_{n_k} : k \in \mathbb{N}\}$ such that there is a pairwise disjoint family $\{F_a : F_a \text{ is a zero-set neighborhood of } a, a \in \bigcup \Delta_{n_k}\}$ and $\{F_a : a \in \bigcup \Delta_{n_k}\}$ is completely Coz_δ -additive.*

2. Main definitions and notation

Throughout this paper, all spaces are assumed to be Tychonoff. The set of positive integers is denoted by \mathbb{N} and $\omega = \mathbb{N} \cup \{0\}$. Let \mathbb{R} be the real line, we put $\mathbb{I} = [0, 1] \subset \mathbb{R}$, and let \mathbb{Q} be the rational numbers. Let $f : X \rightarrow \mathbb{R}$ be a real-valued function, then $\|f\| = \sup\{|f(x)| : x \in X\}$, $S(g, \epsilon) = \{f : \|g - f\| < \epsilon\}$, $B(g, \epsilon) = \{f : \|g - f\| \leq \epsilon\}$, where g is a real valued function and $\epsilon > 0$. Let $V = \{f \in \mathbb{R}^X : f(x_i) \in V_i, i = 1, \dots, n\}$ where $x_i \in X$, $V_i \subseteq \mathbb{R}$ are bounded intervals for $i = 1, \dots, n$, then $suppV = \{x_1, \dots, x_n\}$, $diamV = \max\{diamV_i : 1 \leq i \leq n\}$.

We recall that a subset of X that is the complete preimage of zero for a certain function from $C(X)$ is called a *zero-set*. A subset $O \subseteq X$ is called a *cozero-set* (or *functionally open*) of X if $X \setminus O$ is a zero-set of X . It is easy to check that zero sets are preserved by finite unions and countable intersections. Hence cozero sets are preserved by finite intersections and countable unions. Countable unions of zero sets will be denoted by Zer_σ (or $Zer_\sigma(X)$), countable intersection of cozero sets by Coz_δ (or $Coz_\delta(X)$). It is easy to check that Zer_σ -sets are preserved by countable unions and finite intersections. Note that any zero set is Coz_δ and any cozero-set is Zer_σ . It is well known that f is of the first Baire class if and only if $f^{-1}(U) \in Zer_\sigma$ for every open $U \subseteq \mathbb{R}$ (see Exercise 3.A.1 in [8]).

Lemma 2.1. (Lemma 3.3 in [9]) Let X be a Hausdorff space and let $\{F_i : i \in \mathbb{N}\}$ forms a disjoint completely Coz_δ -additive system. Then any family $\{L_i : L_i \subseteq F_i, i \in \mathbb{N}\}$ of Coz_δ subsets of X is a completely Coz_δ -additive system.

3. Proof of Theorem 1.4

Proof. (2) \Rightarrow (1). Since $B_1^{st}(X)$ is a dense subset of $B_1(X)$, then $B_1(X)$ is Baire if $B_1^{st}(X)$ is so.

(1) \Rightarrow (3). By Theorem 1.3, every pairwise disjoint sequence of non-empty finite subsets of X has a strongly Coz_δ -disjoint subsequence. Let $\{\Delta_n : n \in \mathbb{N}\}$ be a pairwise disjoint sequence of non-empty finite subsets of X and $\{\Delta_{n_k} : k \in \mathbb{N}\}$ be a strongly Coz_δ -disjoint subsequence. Then, there is a pairwise disjoint collection $\{F_a : F_a \text{ is a } \text{Coz}_\delta \text{ neighborhood of } a, a \in \bigcup_{k \in \mathbb{N}} \Delta_{n_k}\}$ such that $\{F_a : a \in \bigcup_{k \in \mathbb{N}} \Delta_{n_k}\}$ is completely Coz_δ -additive. For every $a \in F_a$ there is a zero-set W_a of X such that $a \in W_a \subseteq F_a$. By Lemma 2.1, the family $\{W_a : a \in \bigcup_{k \in \mathbb{N}} \Delta_{n_k}\}$ is completely Coz_δ -additive.

(3) \Rightarrow (2). Assume contrary. Let $B_1^{st}(X) = \bigcup_{n \in \mathbb{N}} F_n$, where F_n is nowhere dense in $B_1(X)$ and $F_n \subseteq F_{n+1}$ for every $n \in \mathbb{N}$.

Claim 1. By Theorem 3.5 in ([9], Claim 1), there are a sequence $\{\Delta_i : i \in \mathbb{N}\}$ of pairwise disjoint finite subsets of X , a sequence $\{\gamma_i : i \in \mathbb{N}\}$ of finite families of basis open sets in \mathbb{R}^X , and a sequence $\{m_i : i \in \mathbb{N}\} \subseteq \mathbb{N}$ such that the following conditions hold for every $i \in \mathbb{N}$:

(a') $1 \leq m_1$ and $m_i + 2 \leq m_{i+1}$;

(b') $\overline{U}^{\mathbb{R}^X} \cap F_i = \emptyset$ and $\text{supp}(U) \subseteq \bigcup_{j=1}^i \Delta_j$ for every $U \in \gamma_i$;

(c') if $f \in U \in \gamma_i$, then $|f(x)| \leq m_i$ for every $x \in \text{supp}(U)$;

(d') if $\varphi : A_i := \bigcup_{j=1}^i \Delta_j \rightarrow \mathbb{R}$ is such that $\|\varphi\|_{A_i} \leq m_i$, then there is a continuous function $f \in \bigcup \gamma_{i+1}$ such that $\|\varphi - f\|_{A_i} < \frac{1}{i}$.

Now we redefine the sequences in Claim 1 to make simpler and clearer their usage in what follows.

By assumption the sequence $\{\Delta_n : n \in \mathbb{N}\}$ constructed in Claim 1 contains a strongly Coz_δ -disjoint subsequence $\{\Delta_{n_k} : k \in \mathbb{N}\}$, where $1 < n_1 < n_2 < \dots$. Put

$$R_1 := F_1, \Omega_1 := \bigcup_{i=1}^{n_1-1} \Delta_i, \mu_1 := \bigcup_{i=1}^{n_1-1} \gamma_i, l_1 := m_{n_1-1},$$

and for every $k \in \mathbb{N}$, we define

$$\begin{aligned} R_{2k} &:= F_{n_k}, \Omega_{2k} := \Delta_{n_k}, \mu_{2k} := \gamma_{n_k}, l_{2k} := m_{n_k} \text{ and} \\ R_{2k+1} &:= F_{n_{k+1}}, \Omega_{2k+1} := \bigcup_{i=n_k+1}^{n_{k+1}-1} \Delta_i, \mu_{2k+1} := \bigcup_{i=n_k+1}^{n_{k+1}-1} \gamma_i, \text{ and} \\ l_{2k+1} &:= m_{n_{k+1}-1}. \end{aligned}$$

It is clear that $\{R_i : i \in \mathbb{N}\}$ is an increasing sequence of nowhere dense sets in $B_1^{st}(X)$ such that $B_1^{st}(X) = \bigcup_{n \in \mathbb{N}} R_n$.

Claim 2. With respect to the sequence $\{R_i : i \in \mathbb{N}\}$, the pairwise disjoint sequence $\{\Omega_i, i \in \mathbb{N}\}$, and the sequences $\{\mu_i : i \in \mathbb{N}\}$ and $\{l_i : i \in \mathbb{N}\}$ satisfy the following conditions ($i \in \mathbb{N}$):

- (a) $1 \leq l_1$ and $l_i + 2 \leq l_{i+1}$;
- (b) $\overline{U}^{\mathbb{R}^X} \cap R_i = \emptyset$ and $\text{supp}(U) \subseteq \bigcup_{j=1}^i \Omega_j$ for every $U \in \mu_i$;
- (c) if $f \in U \in \mu_i$, then $|f(x)| \leq l_i$ for every $x \in \text{supp}(U)$;
- (d) if $\varphi : A_i := \bigcup_{j=1}^i \Omega_j \rightarrow \mathbb{R}$ is such that $\|\varphi\|_{A_i} \leq l_i$, then there is a continuous function $f \in \bigcup \mu_{i+1}$ such that $\|\varphi - f\|_{A_i} < \frac{1}{i}$.

Moreover, the sequence $\{\Omega_{2i} : i \in \mathbb{N}\}$ is strongly Coz_δ -disjoint.

Proof of Claim 2. By construction, $\{\Omega_i : i \in \mathbb{N}\}$ is a sequence of pairwise disjoint finite subsets of X , all families μ_i are finite, and the sequence $\{\Omega_{2i} : i \in \mathbb{N}\}$ is strongly Coz_δ -disjoint by the choice of $\{\Delta_{n_k} : k \in \mathbb{N}\}$. The conditions (a) and (c) are satisfied by (a') and (c'), respectively. Since $F_i \subseteq F_j$ for every $i \leq j$, we have $\text{supp}(U) \subseteq \bigcup_{j=1}^i \Omega_j$ for every $U \in \mu_i$, and hence the condition (b) holds true. The condition (d) is satisfied by the definition of μ_i and the condition (d') for the sets γ_i . The claim is proved. \square

Since, by Claim 2, the sequence $\{\Omega_{2i} : i \in \mathbb{N}\}$ is strongly Coz_δ -disjoint, there is a disjoint completely Coz_δ -additive family $\{W_x : x \in \bigcup_{k \in \mathbb{N}} \Omega_{2k}\}$ of zero-sets of X such that $x \in W_x$ for all $x \in \bigcup_{k \in \mathbb{N}} \Omega_{2k}$, and moreover, the sets

$$W(i) := \bigcup \{W_x : x \in \Omega_{2i}\} \quad (i \in \mathbb{N})$$

are zero-sets in X .

Let $F := X \setminus \bigcup_{i \in \mathbb{N}} W(i)$. Then, $F \in \text{Zer}_\sigma(X) \cap \text{Coz}_\delta(X)$.

Claim 3. By Theorem 3.5 in ([9], Claim 4), there are a sequence $\{f_i : i \in \mathbb{N}\} \subset B_1(X)$, a strictly increasing sequence $\{r_i : i \in \mathbb{N}\} \subseteq \mathbb{N}$ with $r_1 = 1$, a sequence $\{b_i : i \in \mathbb{N}\}$, and a sequence $\{U_i : i \in \mathbb{N}\}$ of basic open sets in \mathbb{R}^X such that

- (e) $U_i \in \mu_{2r_i}$ for every $i \in \mathbb{N}$;
- (f) $\|f_i\|_X \leq l_{2r_i}$ for every $i \in \mathbb{N}$;

- (g) $\|f_{i+1} - f_i\|_{C_i} \leq \frac{1}{2^{b_i}}$ for every $i \in \mathbb{N}$, where $C_i := F \cup \bigcup_{j=1}^{r_i} W(j)$;
- (h) $f_j \in U_i$ for every $j \geq i \geq 1$;
- (k) $b_{i+1} - b_i \rightarrow \infty$.

It follows from Claim 4(g) that the sequence $\{f_i : i \in \mathbb{N}\} \subseteq B_1(X)$ converges pointwise to some function $f : X \rightarrow \mathbb{R}$.

Since $F \in \text{Zer}_\sigma(X)$, then $F = \bigcup_{i \in \mathbb{N}} Q_i$ where $Q_i \subseteq Q_{i+1}$ and Q_i is a zero-set of X for every $i \in \mathbb{N}$.

Since $\{Q_i, \{W_z : z \in \bigcup_{j=1}^i \Omega_{2j}\}\}$ is a pairwise disjoint finite family of zero-sets of X , one can define a continuous function $g_i : X \rightarrow \mathbb{R}$ such that $g_i(x) = 0$ for $x \in Q_i$ and $g_i(x) = f(z)$ for every $x \in W_z$ where $z \in \bigcup_{j=1}^i \Omega_{2j}$.

The sequence $\{g_i : i \in \mathbb{N}\} \subseteq C_p(X)$ converges pointwise to the function $g : X \rightarrow \mathbb{R}$ where

$$g(x) := \begin{cases} 0, & \text{if } x \in F; \\ f(z), & \text{if } x \in W_z \text{ for some } z \in \bigcup_{i \in \mathbb{N}} \Omega_{2i}. \end{cases}$$

Note that $g \in B_1^{st}(X)$.

By Claim 3(h), $g \in \overline{U_i}$ for each $i \in \mathbb{N}$. But, then, by Claim 2(b), $g \notin \bigcup_{i=1}^{\infty} F_i = B_1^{st}(X)$, a contradiction. □

Let $\mathfrak{b} = \min\{|X| : X \text{ has a countable pseudocharacter but } X \text{ is not a } \lambda\text{-space}\}$ (see [10], p.149).

By Theorem 1.4, Proposition 3.15 and Example 4.8 in [2], we get the following results:

- If X is metrizable and $|X| < \mathfrak{b}$ then $B_1^{st}(X)$ is Baire.
- (CH) If X is a Luzin set then $B_1^{st}(X)$ is meager.
- It is consistent with *ZFC* there is a zero-dimensional metrizable separable space X with $|X| = \mathfrak{b}$ such that $B_1^{st}(X)$ is Baire but $B_1^{st}(X)$ is not Choquet (see the definition of a *Choquet* space in [2]).

References

- [1] R. Engelking, General Topology, Revised and completed edition, Heldermann Verlag Berlin (1989).

- [2] T. Banach, S. Gabrielyan, Baire category of some Baire type function spaces, *Topology and its Applications*, **272**, 2020, 107078.
- [3] E.G. Pytkeev, Baire property of spaces of continuous functions. *Mathematical Notes of the Academy of Science of the USSR* **38** (1985), 908–915.(Translated from *Matematicheskie Zametki*, **38**:5, 726–740).
- [4] V.V. Tkachuk, Characterization of the Baire property in $C_p(X)$ by the properties of the space X , *Research papers, Topology-Maps and Extensions of Topological Spaces (Ustinov)*(1985), 21–27.
- [5] J.van Mill ed, Eric K. van Douwen, *Collected Papers*, vol. 1, North-Holland, Amsterdam (1994).
- [6] G. Vidossich, On topological spaces whose functions space is of second category, *Invent.Math.*, **8**:2 (1969), 111–113.
- [7] D.J. Zutter, R.A. McCoy, Category in function spaces, *Pacif. J. Math.*, **90**:1 (1980), 145–168.
- [8] J. Lukeš, J. Malý, L. Zajíček, *Fine Topology Methods in Real Analysis and Potential Theory*, *Lecture Notes in Mathematics*, vol. 1189, Springer, Berlin, 1986.
- [9] A.V. Osipov, Baire property of space of Baire-one functions, [arXiv:2110.15496](https://arxiv.org/abs/2110.15496).
- [10] K. Kunen, J.E. Vaughan, *Handbook of set-theoretic topology*, Elsevier Sc. Publ. B.V., 1984.