# LINEAR MAPS PRESERVING PRODUCTS EQUAL TO PRIMITIVE IDEMPOTENTS OF AN INCIDENCE ALGEBRA

JORGE J. GARCÉS AND MYKOLA KHRYPCHENKO

ABSTRACT. Let A, B be algebras and  $a \in A$ ,  $b \in B$  a fixed pair of elements. We say that a map  $\varphi : A \to B$  preserves products equal to a and b if for all  $a_1, a_2 \in A$  the equality  $a_1a_2 = a$  implies  $\varphi(a_1)\varphi(a_2) = b$ . In this paper we study bijective linear maps  $\varphi : I(X, F) \to I(X, F)$  preserving products equal to primitive idempotents of I(X, F), where I(X, F) is the incidence algebra of a finite connected poset X over a field F. We fully characterize the situation, when such a map  $\varphi$  exists, and whenever it does,  $\varphi$  is either an automorphism of I(X, F) or the negative of an automorphism of I(X, F).

## Contents

Introduction	1
1. Preliminaries	2
1.1. Rings and algebras	2
1.2. Posets	3
1.3. Incidence algebras	3
2. Zero product preservers of $I(X, F)$	4
3. Preservers of products equal to primitive idempotents of $I(X, F)$	4
Acknowledgements	11
References	12

## INTRODUCTION

There are several ways to generalize the notion of a homomorphism  $\varphi : A \to B$ between two algebras. One of them deals with a generalization of the equality

$$\varphi(a_1 a_2) = \varphi(a_1)\varphi(a_2),\tag{1}$$

leading, for instance, to Jordan and Lie homomorphisms. The other one consists in a restriction of the set of pairs  $(a_1, a_2)$  for which (1) holds. Perhaps, the most known generalizations in this direction are zero product preservers  $(a_1a_2 = 0)$ , orthogonality preservers  $(a_1a_2 = a_2a_1 = 0)$  and idempotent preservers  $(a_1 = a_2 \in E(A))$ . Inspired by the notion of a zero product preserver, Chebotar, Ke, Lee and Shiao [10] proposed to study maps that preserve "constant products", i.e. satisfying

$$a_1 a_2 = a_3 a_4 \Rightarrow \varphi(a_1)\varphi(a_2) = \varphi(a_3)\varphi(a_4). \tag{2}$$

<sup>2010</sup> Mathematics Subject Classification. Primary: 16S50, 15A86; secondary: 17C27.

Key words and phrases. Incidence algebra; product preserver; primitive idempotent.

A variant of this property is the following:

$$a_1 a_2 = a \Rightarrow \varphi(a_1)\varphi(a_2) = b, \tag{3}$$

where  $a \in A$  and  $b \in B$  are fixed. In this case we say that  $\varphi$  preserves products equal to a and b. If a and b are zero, we get a zero product preserver. If a and b are the identity elements, we get a map preserving the inverses.

Chebotar, Ke, Lee and Shiao [10] proved that a bijective additive map on a division ring satisfying (2) with  $a_1a_2 = a_3a_4$  being the identity element is either an automorphism multiplied by a central element or an anti-automorphism multiplied by a central element. This generalizes an old result [1, Theorem 1.15] on inverse preservers of division rings. Catalano [4] extended the above description by Chebotar, Ke, Lee and Shiao to the case, where  $a_1a_2$  is an arbitrary invertible element of a division ring. Later, this was generalized in [7] to maps on the ring  $M_n(D)$  of matrices (with usual or Jordan product) over a division ring D. The next natural question was to study maps that preserve products equal to non-invertible elements of  $M_n(D)$ . The case, where such products are rank-one idempotents of  $M_n(\mathbb{C})$ , was treated by Catalano in [5], and the corresponding maps were proved to be  $\pm$  homomorphisms. Bijective maps on  $M_n(\mathbb{C})$  preserving products equal to nilpotent matrices were later characterized in [6] by Catalano and Chang-Lee. Catalano and Julius [8] solved the problem, where the products are diagonalizable matrices over  $\mathbb C$  with the same eigenvalues. Perhaps, the most general situation was considered in [9]: one of the products is a matrix from  $M_n(\mathbb{C})$  of rank at most n-2, or both products have the same rank n-1. The corresponding product preservers are again of the standard form, i.e. automorphisms multiplied by scalars. When  $M_n(\mathbb{C})$  is equipped with the Lie bracket [a, b] = ab - ba, the description slightly differs, but still reminds the description of commutativity preservers (i.e. zero product preservers with respect to [-, -]), see [14, 15]. In the setting of C<sup>\*</sup>-algebras the problem of studying linear mappings that are \*-homomorphisms at a fixed point has been considered in [3].

Notice that the incidence algebra I(X, F) of a finite poset X over a field F can be seen as a subalgebra of the algebra  $T_n(F)$  of upper triangular matrices, where n = |X|. The primitive idempotents of I(X, F) (which correspond to rank-one idempotent matrices from  $T_n(F)$ ) were described in [17] and the automorphisms of I(X, F) have been studied by various authors (see, for example, [19, 2, 12, 16]), so it is natural to ask whether the Catalano's result [5] holds for maps on incidence algebras. Under the assumption that X is connected, we show in Theorem 3.17 that a bijective linear map  $\varphi : I(X, F) \to I(X, F)$  preserving the products equal to fixed primitive idempotents does not always exist, and its existence depends on the existence of an automorphism  $\lambda$  of X mapping one fixed element  $x \in X$ to another fixed element  $y \in X$ . If such  $\lambda \in \operatorname{Aut}(X)$  exists, then  $\varphi$  is either an automorphism of I(X, F) or the negative of an automorphism of I(X, F). Our result can be applied, in particular, to the upper triangular matrix algebra  $T_n(F)$ (see Corollary 3.18).

### 1. Preliminaries

1.1. Rings and algebras. An element a of a ring R satisfying  $a^2 = a$  is called an *idempotent*. The set of idempotents of R will be denoted by E(R).

Two elements  $a, b \in R$  are orthogonal if ab = ba = 0. An idempotent  $e \in E(R)$  is primitive if  $e = e_1 + e_2$  with  $e_1, e_2 \in E(R)$  orthogonal implies  $e_1, e_2 \in \{0, e\}$ .

1.2. **Posets.** A poset is a pair  $(X, \leq)$ , where X is a set and  $\leq$  is a partial order (i.e., a reflexive, transitive and anti-symmetric binary relation) on X. All the posets in this article will be finite. The *interval* from x to y in a poset X is the subset  $[x, y] := \{z \in X \mid x \leq z \leq y\}$ . A chain in X is a non-empty subset  $C \subseteq X$  such that for all  $x, y \in C$  either  $x \leq y$  or  $y \leq x$ . The length of a chain  $C \subseteq X$  is defined to be |C|-1. The length of a non-empty poset X is  $l(X) := \max\{l(C) \mid C \text{ is a chain in } X\}$ . We write l(x, y) for l([x, y]). A walk in X is a sequence  $x_0, x_1, \ldots, x_m$  of elements of X, such that for all  $i = 0, \ldots, m-1$  either  $x_i \leq x_{i+1}$  and  $l(x_i, x_{i+1}) = 1$  or  $x_{i+1} \leq x_i$  and  $l(x_{i+1}, x_i) = 1$ . A poset X is connected if for all  $x, y \in X$  there exists a walk  $x = x_0, \ldots, x_m = y$ .

Given a subset  $A \subseteq X^2$ , denote by  $A_{\leq}$  (resp.  $A_{<}$ ) the set  $\{(x, y) \in A \mid x \leq y\}$  (resp.  $\{(x, y) \in A \mid x < y\}$ ).

A map  $\lambda : X \to Y$  between two posets is *order-preserving* whenever  $x \leq y \Rightarrow \lambda(x) \leq \lambda(y)$  for all  $x, y \in X$ . An *(order)* automorphism of a finite poset X is an order-preserving bijection of X.<sup>1</sup>

1.3. Incidence algebras. The incidence algebra I(X, F) of a finite poset X over a field F is the F-vector space with basis  $\{e_{xy} \mid x \leq y\}$  (called the standard basis) and multiplication

$$e_{xy}e_{uv} = \begin{cases} e_{xv}, & y = u, \\ 0, & y \neq u. \end{cases}$$

For an element  $f \in I(X, F)$  we write  $f = \sum_{x \le y} f(x, y) e_{xy}$ , so that

$$(fg)(x,y) = \sum_{x \le z \le y} f(x,z)g(z,y).$$

The algebra I(X, F) is associative and unital, where the identity element of I(X, F) is  $\delta := \sum_{x \in X} e_{xx}$ .

Given  $f \in I(X, F)$ , define  $f_D = \sum_{x \in X} f(x, x) e_{xx}$  and  $f_U = \sum_{x < y} f(x, y) e_{xy}$ , so that  $f = f_D + f_U$ . An element  $f \in I(X, F)$  is said to be diagonal whenever  $f = f_D$ . The diagonal elements form a commutative subalgebra of I(X, F), denoted by D(X, F). The elements f with  $f = f_U$  (equivalently,  $f_D = 0$ ) form an ideal which coincides with the Jacobson radical of I(X, F), denoted J(I(X, F)) (see [18, Theorem 4.2.5]). The set  $\mathcal{B} := \{e_{xy} \mid x < y\}$  is clearly a basis of J(I(X, F)). Thus,  $I(X, F) = D(X, F) \oplus J(I(X, F))$  as vector spaces. Invertible elements of I(X, F)are exactly those  $f \in I(X, F)$  such that  $f(x, x) \in F^*$  for all  $x \in X$  by [18, Theorem 1.2.3]. The center of I(X, F) coincides with  $\{r\delta \mid r \in F\}$  by [18, Corollary 1.3.15], provided that X is connected. For any  $Y \subseteq X$  the incidence algebra I(Y, F) can be seen as a subspace of I(X, F) closed under multiplication, where Y is assumed to be endowed with the induced partial order. Observe that we avoid using the term "subalgebra", because the identity element of I(Y, F) is different from that of I(X, F) whenever  $Y \neq X$ .

We will write  $e_x := e_{xx}$  and, more generally,  $e_A := \sum_{a \in A} e_{aa}$  for  $A \subseteq X$ . Then  $\{e_x \mid x \in X\}$  is a set of pairwise orthogonal primitive idempotents of I(X, F) which

<sup>&</sup>lt;sup>1</sup>For an infinite X one should also require that the inverse bijection be order-preserving, which holds automatically in the finite case.

form a basis of D(X, F). Observe that I(X, F) also admits a basis formed by the idempotents  $\{e_x\}_{x \in X} \sqcup \{e_x + e_{xy}\}_{x < y}$ .

Recall from [19, 2, 16] the description of automorphisms of I(X, F). Any order automorphism  $\lambda$  of X induces an automorphism  $\widehat{\lambda}$  of I(X, F) by means of

$$\widehat{\lambda}(e_{xy}) = e_{\lambda(x)\lambda(y)},\tag{4}$$

where  $x \leq y$ . Moreover, a map  $\sigma : X_{\leq}^2 \to F^*$  with

$$\sigma(x, y)\sigma(y, z) = \sigma(x, z) \tag{5}$$

for all  $x \leq y \leq z$  defines the multiplicative automorphism  $M_{\sigma}$  of I(X, F) acting as follows:

$$M_{\sigma}(e_{xy}) = \sigma(x, y)e_{xy},\tag{6}$$

where  $x \leq y$ . Any automorphism of I(X, F) is the composition of an inner automorphism of I(X, F) and automorphisms  $\lambda$  and  $M_{\sigma}$  described in (4) and (6).

## 2. Zero product preservers of I(X, F)

Recall that a map  $\varphi: A \to B$  between two algebras preserves zero products (or is a zero product preserver) if  $\varphi(a)\varphi(b) = 0$  in B whenever ab = 0 in A. It turns out that the description of zero product preservers of I(X, F) by their action on the natural basis will be useful in the following section.

**Proposition 2.1.** Let X be finite and  $\varphi: I(X, F) \to A$  a linear map, where A is an algebra. Then  $\varphi$  is a zero product preserver if and only if

$$\varphi(e_{xy})\varphi(e_{uv}) = 0 \text{ for all } x \le y \text{ and } u \le v \text{ with } e_{xy}e_{uv} = 0, \tag{7}$$

$$\varphi(e_x)\varphi(e_{xy}) = \varphi(e_{xz})\varphi(e_{zy}) = \varphi(e_{xy})\varphi(e_y) \text{ for all } x < z < y.$$
(8)

*Proof.* Let  $\varphi$  be a zero product preserver. Then (7) is immediate. Since  $(e_x + e_y)$  $(e_{xz})(e_{zy} - e_{xy}) = 0$  for all  $x < z \le y$ , then using (7) we have

$$0 = \varphi(e_x)\varphi(e_{zy}) - \varphi(e_x)\varphi(e_{xy}) + \varphi(e_{xz})\varphi(e_{zy}) - \varphi(e_{xz})\varphi(e_{xy})$$
  
=  $\varphi(e_{xz})\varphi(e_{zy}) - \varphi(e_x)\varphi(e_{xy}),$ 

proving (8).

Conversely, assume (7) and (8). Take arbitrary  $f, g \in I(X, F)$  and calculate

$$\begin{split} \varphi(f)\varphi(g) &= \sum_{x \le y} \sum_{u \le v} f(x,y)g(u,v)\varphi(e_{xy})\varphi(e_{uv}) = \sum_{x \le z \le y} f(x,z)g(z,y)\varphi(e_{xz})\varphi(e_{zy}) \\ &= \sum_{x \le y} \left( \sum_{x \le z \le y} f(x,z)g(z,y) \right) \varphi(e_x)\varphi(e_{xy}) = \sum_{x \le y} (fg)(x,y)\varphi(e_x)\varphi(e_{xy}). \end{split}$$
  
So, if  $fg = 0$ , then  $\varphi(f)\varphi(g) = 0$ .

So, if fg = 0, then  $\varphi(f)\varphi(g) = 0$ .

### 3. Preservers of products equal to primitive idempotents of I(X, F)

Our goal is to study the bijective linear maps  $\varphi : I(X,F) \to I(X,F)$  which preserve products equal to a fixed pair of primitive idempotents  $\epsilon, \eta \in I(X, F)$ , i.e. satisfying (3) with  $a = \epsilon$  and  $b = \eta$ . By [17, Lemma 1] there exist  $x, y \in X$ 

and inner automorphisms  $\psi_1, \psi_2$  of I(X, F), such that  $\psi_1(e_x) = \epsilon$  and  $\psi_2(\eta) = e_y$ . Then, replacing  $\varphi$  by  $\psi_2 \circ \varphi \circ \psi_1$ , we may assume that  $\varphi$  satisfies

$$fg = e_x \Rightarrow \varphi(f)\varphi(g) = e_y \tag{9}$$

for all  $f, g \in I(X, F)$ .

We first list here easy consequences of (9) as in [5].

Lemma 3.1. The following equalities hold:

 $\begin{array}{ll} (i) \ \varphi(e_x)^2 = e_y; \\ (ii) \ (a) \ \varphi(e_x)\varphi(e_{uv}) = 0 \ for \ all \ x \neq u \leq v; \\ (b) \ \varphi(e_{uv})\varphi(e_x) = 0 \ for \ all \ u \leq v \neq x; \\ (iii) \ (a) \ \varphi(e_z)\varphi(e_{ux}) = 0 \ for \ all \ x \neq u \leq x; \\ (b) \ \varphi(e_x)\varphi(e_x) = 0 \ for \ all \ x \leq v \neq z; \\ (iv) \ \varphi(e_u)\varphi(e_v) = 0 \ for \ all \ u \leq v; \\ (v) \ (a) \ \varphi(e_{ux})\varphi(e_x) - \varphi(e_u)\varphi(e_{ux}) = r\varphi(e_{ux})^2 \ for \ all \ u < x \ and \ r \in F^*; \\ (b) \ \varphi(e_{ux})\varphi(e_{xv}) - \varphi(e_{xv})\varphi(e_v) = r\varphi(e_{xv})^2 \ for \ all \ v > x \ and \ r \in F^*; \\ (vi) \ (a) \ \varphi(e_{ux})\varphi(e_z) = 0 \ for \ all \ u \leq x, \ z \notin \{u, x\}; \end{array}$ 

(b) 
$$\varphi(e_z)\varphi(e_{xv}) = 0$$
 for all  $x \le v, z \notin \{x, v\}$ .

*Proof.* (i) follows from  $e_x^2 = e_x$ ; (ii)a follows from (i) and  $e_x(e_x + e_{uv}) = e_x$  for  $x \neq u \leq v$ ; (ii)b follows from (i) and  $(e_x + e_{uv})e_x = e_x$  for  $u \leq v \neq x$ .

It suffices to prove (iii)a for u < x, since u = x is a particular case of (ii)b. So, assume u < x and write  $\varphi(e_x)\varphi(e_{ux}) = 0$  by (ii)a. In particular, this proves (iii)a for z = x. Thus, take  $z \neq x$ . Then  $(e_x + e_z)(e_x + e_{ux}) = e_x$ . Combining this with  $\varphi(e_z)\varphi(e_x) = 0$  (by (ii)b) and (i), we obtain (iii)a. Similarly, (iii)b follows from  $\varphi(e_{xv})\varphi(e_x) = 0$  for x < v,  $(e_x + e_{xv})(e_x + e_z) = e_x$  for  $z \neq x$  and (i).

If  $x \in \{u, v\}$ , then (iv) is a particular case of (ii)a or (ii)b. Otherwise, consider the product  $(e_x + e_u)(e_x + e_v) = e_x$  and apply (i), (ii)a and (ii)b.

For (v)a write  $(e_x + e_u + re_{ux})(e_x - re_{ux}) = e_x$  and use (i), (ii)a and (ii)b. Item (v)b is similar and follows from  $(e_x - re_{xv})(e_x + e_v + re_{xv}) = e_x$ .

For (vi)a write  $(e_x + e_u + e_{ux})(e_x - e_{ux} + e_z) = e_x$  and use (ii)a, (iv) and (v)a. Analogously, (vi)b is a consequence of (ii)b, (iv) and (v)b.

## Corollary 3.2. Let $z \in X \setminus \{y\}$ .

- (i) If  $\varphi(e_u)(z, z) = 0$  for some u < x, then  $\varphi(e_{ux})(z, z) = 0$ .
- (ii) If  $\varphi(e_v)(z, z) = 0$  for some v > x, then  $\varphi(e_{xv})(z, z) = 0$ .

*Proof.* Let us prove (i). Using Lemma 3.1 (i) we have  $\varphi(e_x)(z, z)^2 = e_y(z, z) = 0$ , so  $\varphi(e_x)(z, z) = 0$ . Therefore,

$$r\varphi(e_{ux})(z,z)^2 = \varphi(e_{ux})(z,z)\varphi(e_x)(z,z) - \varphi(e_u)(z,z)\varphi(e_{ux})(z,z) = 0$$

by Lemma 3.1 (v)a, whence  $\varphi(e_{ux})(z, z) = 0$ . The proof of (ii) similarly follows from Lemma 3.1 (v)b.

From now on we view  $I(X \setminus \{x\}, F)$  and  $I(X \setminus \{y\}, F)$  as subspaces of I(X, F) closed under multiplication. Their identity elements are  $e_{X \setminus \{x\}}$  and  $e_{X \setminus \{y\}}$ , respectively.

**Lemma 3.3.** Let  $f \in I(X, F)$ . If  $f^2 = e_y$ , then  $f = \pm e_y + g$  for some  $g \in J(I(X \setminus \{y\}, F))$  with  $g^2 = 0$ .

*Proof.* It immediately follows that  $f_D = \pm e_y$ . Now take u < y and write

$$\begin{split} 0 &= e_y(u,y) = f^2(u,y) = f(u,y)f(y,y) + \sum_{u < v < y} f(u,v)f(v,y) \\ &= \pm f(u,y) + \sum_{u < v < y} f(u,v)f(v,y), \end{split}$$

whence

$$f(u,y) = \mp \sum_{u < v < y} f(u,v)f(v,y).$$

$$\tag{10}$$

If l(u, y) = 1, then there is no  $v \in X$  with u < v < y, so the sum on the right-hand side of (10) vanishes, and f(u, y) = 0. If l(u, y) > 1, then l(v, y) < l(u, y) for all u < v < y, so the obvious induction argument implies f(u, y) = 0.

Similarly, for all y < v one has

$$\begin{split} 0 &= e_y(y,v) = f^2(y,v) = f(y,y)f(y,v) + \sum_{y < u < v} f(y,u)f(u,v) \\ &= \pm f(y,v) + \sum_{y < u < v} f(y,u)f(u,v), \end{split}$$

whence f(y, v) = 0 by induction on l(y, v).

Thus,  $f = \pm e_y + g$ , where  $g = f_U$ . Observe that  $e_y g = \sum_{y < v} f(y, v) e_{yv} = 0$  and  $ge_y = \sum_{u < y} f(u, y) e_{uy} = 0$ . Therefore,  $e_y = f^2 = e_y \pm e_y g \pm g e_y + g^2 = e_y + g^2$ , whence  $g^2 = 0$ .

**Corollary 3.4.** We have  $\varphi(e_x) = \pm e_y + g$  for some  $g \in J(I(X \setminus \{y\}, F))$  with  $g^2 = 0$ .

**Lemma 3.5.** We have  $\varphi(I(X \setminus \{x\}, F)) \subseteq I(X \setminus \{y\}, F)$ . Moreover,  $\varphi$  is a zero product preserver from  $I(X \setminus \{x\}, F)$  to  $I(X \setminus \{y\}, F)$ .

*Proof.* Let  $u, v \in X \setminus \{x\}$ ,  $u \leq v$ . Then  $\varphi(e_x)\varphi(e_{uv}) = 0$  by Lemma 3.1 (ii)a. Multiplying this by  $\varphi(e_x)$  on the left and using Lemma 3.1 (i) we obtain  $e_y\varphi(e_{uv}) = 0$ . Similarly  $\varphi(e_{uv})e_y = 0$  by (i) and (ii)b of Lemma 3.1. Therefore,  $\varphi(e_{uv})(a, y) = \varphi(e_{uv})(y, b)$  for all  $a \leq y \leq b$ . This proves the desired inclusion.

Now take  $f, g \in I(X \setminus \{x\}, F)$  such that fg = 0. It is easily seen that  $(e_x + f)(e_x + g) = e_x$  in I(X, F). Hence,  $(\varphi(e_x) + \varphi(f))(\varphi(e_x) + \varphi(g)) = e_y$  thanks to (9). The latter is equivalent to  $\varphi(f)\varphi(g) = 0$  by (i), (ii)a and (ii)b of Lemma 3.1.  $\Box$ 

Observe that dim $(I(X \setminus \{x\}, F)) = |(X \setminus \{x\})_{\leq}^2|$  and dim $(I(X \setminus \{y\}, F)) = |(X \setminus \{y\})_{\leq}^2|$ , so we have the following immediate consequence of Lemma 3.5.

**Corollary 3.6.** If  $|(X \setminus \{x\})_{\leq}^2| > |(X \setminus \{y\})_{\leq}^2|$ , then there is no bijective linear map  $\varphi : I(X, F) \to I(X, F)$  satisfying (9).

Thus, in what follows we assume  $|(X \setminus \{x\})^2_{\leq}| \leq |(X \setminus \{y\})^2_{\leq}|$ .

**Corollary 3.7.** We have  $\varphi(J(I(X \setminus \{x\}, F))) \subseteq J(I(X \setminus \{y\}, F))$ .

*Proof.* Indeed, for any  $u, v \in X \setminus \{x\}, u < v$ , it follows from  $e_{uv}^2 = 0$  and Lemma 3.5 that  $\varphi(e_{uv})^2 = 0$ , so  $\varphi(e_{uv}) \in J(I(X \setminus \{y\}, F))$ .

**Lemma 3.8.** The element  $\varphi(e_{X \setminus \{x\}})$  is invertible in  $I(X \setminus \{y\}, F)$ .

*Proof.* For all  $u \in X \setminus \{x\}$  denote  $Z_u = \{z \in X \mid \varphi(e_u)(z, z) \neq 0\} \subseteq X \setminus \{y\}$ . Observe that  $Z_u \cap Z_v = \emptyset$  for  $u \neq v$  thanks to Lemma 3.1 (iv). Since  $e_{X \setminus \{x\}} = \sum_{u \in X \setminus \{x\}} e_u$ , we have

$$Z := \left\{ z \in X \mid \varphi\left(e_{X \setminus \{x\}}\right)(z, z) \neq 0 \right\} = \bigsqcup_{u \in X \setminus \{x\}} Z_u.$$

Assume that  $\varphi(e_{X\setminus\{x\}}) \notin I(X\setminus\{y\}, F)^*$ . Then  $Z \neq X\setminus\{y\}$ , so there exists  $z_0 \in X\setminus\{y\}$  such that  $z_0 \notin Z$ . It follows that  $\varphi(e_u)(z_0, z_0) = 0$  for all  $u \in X\setminus\{x\}$ . By Corollary 3.2 we conclude that  $\varphi(e_{ux})(z_0, z_0) = 0$  for all u < x and  $\varphi(e_{xv})(z_0, z_0) = 0$  for all v > x. Furthermore,  $\varphi(e_{uv})(z_0, z_0) = 0$  for all  $u, v \in X\setminus\{x\}, u < v$ , by Corollary 3.7. Finally,  $\varphi(e_x)(z_0, z_0) = 0$  by Corollary 3.4. Thus,  $\varphi(f)(z_0, z_0) = 0$  for all  $f \in I(X, F)$ , a contradiction with  $e_{z_0} \in \varphi(I(X, F))$ .

**Lemma 3.9.** We have  $\varphi(e_x) = \pm e_y$  and there exist a monomorphism  $\psi : I(X \setminus \{x\}, F) \to I(X \setminus \{y\}, F)$  and  $c \in I(X \setminus \{y\}, F)^*$  such that  $\varphi(f) = c\psi(f)$  for all  $f \in I(X \setminus \{x\}, F)$ .

Proof. Recall that  $I(X \setminus \{x\}, F)$  has a basis formed by idempotents. Therefore, since  $\varphi : I(X \setminus \{x\}, F) \to I(X \setminus \{y\}, F)$  preserves zero products by Lemma 3.5 and  $\varphi(e_{X \setminus \{x\}})$  is invertible in  $I(X \setminus \{y\}, F)$ , it follows from [11, Theorem 2.6(vi)] that there are a homomorphism  $\psi : I(X \setminus \{x\}, F) \to I(X \setminus \{y\}, F)$  and  $c = \varphi(e_{X \setminus \{x\}}) \in$  $I(X \setminus \{y\}, F)^*$  such that  $\varphi(f) = c\psi(f)$  for all  $f \in I(X \setminus \{x\}, F)$ . The homomorphism  $\psi$  is injective, because  $\varphi$  is injective and c is invertible. Recall from Corollary 3.4 that  $\varphi(e_x) = \pm e_y + g$ , where  $g \in I(X \setminus \{y\}, F)$ . Now,  $\varphi(e_x)c = \varphi(e_x)\varphi(e_{X \setminus \{x\}}) = 0$ by Lemma 3.1 (iv). On the other hand,  $\varphi(e_x)c = \pm e_yc + gc = gc$ . Hence, gc = 0. But c is invertible in  $I(X \setminus \{y\}, F)$ , so g = 0. Thus,  $\varphi(e_x) = \pm e_y$ .

**Lemma 3.10.** There exist  $\beta \in I(X, F)^*$ ,  $\{\alpha_z\}_{z \in X} \subseteq F^*$  and a bijection  $\lambda : X \to X$  with  $\lambda(x) = y$  such that

$$\beta\varphi(e_z)\beta^{-1} = \alpha_z e_{\lambda(z)} \tag{11}$$

for all  $z \in X$ . Moreover,

$$\beta e_y \beta^{-1} = e_y \tag{12}$$

and  $\alpha_x = 1 \Leftrightarrow \varphi(e_x) = e_y, \ \alpha_x = -1 \Leftrightarrow \varphi(e_x) = -e_y.$ 

*Proof.* By Lemma 3.9 for any  $z \in X \setminus \{x\}$  we have  $\varphi(e_z) = c\psi(e_z)$ , where  $c \in I(X \setminus \{y\}, F)^*$  and  $\psi$  is a monomorphism  $\psi : I(X \setminus \{x\}, F) \to I(X \setminus \{y\}, F)$ . Since  $\{\psi(e_z)\}_{z \in X \setminus \{x\}}$  is a set of orthogonal idempotents and  $|X \setminus \{x\}| = |X \setminus \{y\}|$ , there exists a bijection  $\lambda' : X \setminus \{x\} \to X \setminus \{y\}$  such that  $\psi(e_z)_D = e_{\lambda'(z)}$ . Hence, by [13, Lemma 5.4] there is  $\beta' \in I(X \setminus \{y\}, F)^*$  such that  $\beta'\psi(e_z)\beta'^{-1} = e_{\lambda'(z)}$  for all  $z \in X \setminus \{x\}$ . Then

$$\beta'\varphi(e_z)\beta'^{-1} = \beta'c\psi(e_z)\beta'^{-1} = \beta'c\beta'^{-1}e_{\lambda'(z)}.$$
(13)

Denote  $\gamma = \beta' c \beta'^{-1} \in I(X \setminus \{y\}, F)^*$ . Let u < v in  $X \setminus \{y\}$  and  $a, b \in X \setminus \{x\}$  such that  $\lambda'(a) = u$  and  $\lambda'(b) = v$ . Then it follows from Lemma 3.1 (iv) and (13) that

$$0 = \varphi(e_a)\varphi(e_b) = \gamma e_u \gamma e_v = \gamma \cdot \gamma(u, v) e_{uv}.$$

Since  $\gamma$  is invertible in  $I(X \setminus \{y\}, F)$ , then  $\gamma(u, v)e_{uv} = 0$ , whence  $\gamma(u, v) = 0$ . Thus,  $\gamma$  is diagonal and we define  $\alpha_z = \gamma(z, z) \in F^*$  for all  $z \in X \setminus \{x\}$ , so that

$$\beta'\varphi(e_z)\beta'^{-1} = \alpha_z e_{\lambda'(z)} \tag{14}$$

for such z.

We now extend  $\lambda'$  to a bijection  $\lambda: X \to X$  by  $\lambda(x) = y$  and define  $\beta = e_y + \beta'$ ,

$$\alpha_x = \begin{cases} 1, & \varphi(e_x) = e_y, \\ -1, & \varphi(e_x) = -e_y. \end{cases}$$

Then  $\beta \in I(X, F)^*$  and using (14) we obtain

$$\beta\varphi(e_z)\beta^{-1} = (e_y + \beta')\varphi(e_z)(e_y + {\beta'}^{-1}) = \beta'\varphi(e_z)\beta'^{-1} = \alpha_z e_{\lambda'(z)} = \alpha_z e_{\lambda(z)}$$

for all  $z \in X \setminus \{x\}$  and

$$\beta\varphi(e_x)\beta^{-1} = (e_y + \beta')(\pm e_y)(e_y + \beta'^{-1}) = \pm e_y = \alpha_x e_{\lambda(x)}.$$

So, (11) holds for all  $z \in X$ . Moreover,

$$\beta e_y \beta^{-1} = (e_y + \beta')e_y(e_y + \beta'^{-1}) = e_y,$$

proving (12).

Observe from Lemma 3.10 and (9) and (11) that if  $fg = e_x$  then

$$\beta\varphi(f)\beta^{-1}\beta\varphi(g)\beta^{-1} = \beta\varphi(f)\varphi(g)\beta^{-1} = \beta e_y\beta^{-1} = e_y,$$

so, without loss of generality, we may assume that, for all  $z \in X$ ,

$$\varphi(e_z) = \alpha_z e_{\lambda(z)},\tag{15}$$

where  $\alpha_z \in F^*$ ,  $\alpha_x = 1 \Leftrightarrow \varphi(e_x) = e_y$ ,  $\alpha_x = -1 \Leftrightarrow \varphi(e_x) = -e_y$  and  $\lambda : X \to X$  is a bijection with  $\lambda(x) = y$ .

**Lemma 3.11.** The bijection  $\lambda$  is an automorphism of X.

*Proof.* Let  $u, v \in X \setminus \{x\}$  with  $u \leq v$ . Then  $e_u, e_v, e_{uv} \in I(X \setminus \{x\}, F)$ . By Lemma 3.9 and (15)

$$0 \neq \varphi(e_{uv}) = c\psi(e_{uv}) = c\psi(e_u e_{uv} e_v) = c\psi(e_u)\psi(e_{uv})\psi(e_v) = ce_{\lambda(u)}\psi(e_{uv})e_{\lambda(v)},$$

whence  $\lambda(u) \leq \lambda(v)$ . So,  $\lambda|_{X \setminus \{x\}}$  is order-preserving. Now let u < x. By Lemma 3.1 (iii)a and (15) for any  $z \neq u$  we have

$$0 = \varphi(e_z)\varphi(e_{ux}) = \alpha_z e_{\lambda(z)}\varphi(e_{ux}),$$

so  $e_{\lambda(z)}\varphi(e_{ux}) = 0$  for all  $z \neq u$ . It follows that

$$\varphi(e_{ux}) = \delta \cdot \varphi(e_{ux}) = \left(\sum_{z \in X} e_{\lambda(z)}\right) \varphi(e_{ux}) = e_{\lambda(u)} \varphi(e_{ux}).$$
(16)

Now by (15) and (16) and Lemma 3.1 (v)a (with r = 1) we have

$$0 = \varphi(e_{ux})\alpha_x e_{\lambda(x)} - \alpha_u e_{\lambda(u)}\varphi(e_{ux}) - e_{\lambda(u)}\varphi(e_{ux})e_{\lambda(u)}\varphi(e_{ux})$$
  
$$= e_{\lambda(u)}\varphi(e_{ux})\alpha_x e_{\lambda(x)} - \alpha_u\varphi(e_{ux}) - \varphi(e_{ux})(\lambda(u),\lambda(u))e_{\lambda(u)}\varphi(e_{ux})$$
  
$$= \alpha_x e_{\lambda(u)}\varphi(e_{ux})e_{\lambda(x)} - (\alpha_u + \varphi(e_{ux})(\lambda(u),\lambda(u)))\varphi(e_{ux}).$$
(17)

8

On the other hand, by (15) and Lemma 3.1 (vi)a we have  $\varphi(e_{ux})e_{\lambda(z)} = 0$  for all  $z \notin \{u, x\}$ , so combining this with (16) we get

$$\varphi(e_{ux}) = \varphi(e_{ux}) \cdot \delta = \varphi(e_{ux}) \cdot \left(\sum_{z \in X} e_{\lambda(z)}\right) = \varphi(e_{ux})e_{\lambda(u)} + \varphi(e_{ux})e_{\lambda(x)}$$
$$= e_{\lambda(u)}\varphi(e_{ux})e_{\lambda(u)} + e_{\lambda(u)}\varphi(e_{ux})e_{\lambda(x)}$$
$$= \varphi(e_{ux})(\lambda(u), \lambda(u))e_{\lambda(u)} + e_{\lambda(u)}\varphi(e_{ux})e_{\lambda(x)}.$$
(18)

Suppose that  $e_{\lambda(u)}\varphi(e_{ux})e_{\lambda(x)} = 0$ . Then thanks to (17) and  $\varphi(e_{ux}) \neq 0$  we obtain  $\varphi(e_{ux})(\lambda(u), \lambda(u)) = -\alpha_u$ . Therefore, (15) and (18) give  $\varphi(e_{ux}) = -\alpha_u e_{\lambda(u)} = -\varphi(e_u) = \varphi(-e_u)$ , whence  $e_{ux} = -e_u$ , a contradiction. Thus,  $e_{\lambda(u)}\varphi(e_{ux})e_{\lambda(x)} \neq 0$ , i.e.  $\lambda(u) < \lambda(x)$  and  $\varphi(e_{ux})(\lambda(u), \lambda(x)) \neq 0$ .

Similarly one proves that  $\lambda(x) < \lambda(v)$  for all x < v.

Thus,  $\lambda$  is order-preserving. Since X is finite,  $\lambda^{-1}$  is also order-preserving. Thus,  $\lambda$  is an automorphism of X.

**Lemma 3.12.** We have  $\alpha_u = \alpha_x$  and

$$\varphi(e_{ux}) = \varphi(e_{ux})(\lambda(u), \lambda(x))e_{\lambda(u)\lambda(x)}$$
(19)

for all u < x. Similarly,  $\alpha_v = \alpha_x$  and

$$\varphi(e_{xv}) = \varphi(e_{xv})(\lambda(x), \lambda(v))e_{\lambda(x)\lambda(v)}$$
(20)

for all v > x.

*Proof.* Substituting the right-hand side of (18) to (17), we obtain

$$0 = (\alpha_x - \alpha_u - \varphi(e_{ux})(\lambda(u), \lambda(u)))\varphi(e_{ux})(\lambda(u), \lambda(x))e_{\lambda(u)\lambda(x)} - (\alpha_u + \varphi(e_{ux})(\lambda(u), \lambda(u)))\varphi(e_{ux})(\lambda(u), \lambda(u))e_{\lambda(u)}.$$

If  $\varphi(e_{ux})(\lambda(u), \lambda(u)) \neq 0$ , then  $\alpha_u + \varphi(e_{ux})(\lambda(u), \lambda(u)) = 0$ , so  $\varphi(e_{ux})(\lambda(u), \lambda(x)) = 0$ , a contradiction with  $e_{\lambda(u)}\varphi(e_{ux})e_{\lambda(x)} \neq 0$  proved in Lemma 3.11. Consequently,  $\varphi(e_{ux})(\lambda(u), \lambda(u)) = 0$  and  $\alpha_u = \alpha_x$ . Then (19) follows from (18). The proof of  $\alpha_v = \alpha_x$  and (20) for all v > x is similar.

Corollary 3.13. We have  $\varphi(J(I(X,F)) = J(I(X,F)))$ .

*Proof.* Observe that  $\varphi(e_{ux}), \varphi(e_{xv}) \in J(I(X, F))$  for all u < x < v by Lemma 3.12. Together with Corollary 3.7 this results in  $\varphi(J(I(X, F))) \subseteq J(I(X, F))$ . Since  $\varphi$  is injective, we have  $\varphi(J(I(X, F))) = J(I(X, F))$ .

Let  $\widehat{\lambda}$  be the automorphism of I(X, F) induced by  $\lambda$  as in (4). Replacing  $\varphi$  by  $(\widehat{\lambda})^{-1} \circ \varphi$ , we may assume that x = y and, for all  $z \in X$ ,

$$\varphi(e_z) = \alpha_z e_z,\tag{21}$$

where  $\alpha_z \in F^*$  with  $\alpha_x = 1 \Leftrightarrow \varphi(e_x) = e_x$  and  $\alpha_x = -1 \Leftrightarrow \varphi(e_x) = -e_x$ .

**Lemma 3.14.** For all u < v in X we have

$$\varphi(e_{uv}) = \sigma(u, v)e_{uv} \tag{22}$$

for some  $\sigma(u, v) \in F^*$ .

*Proof.* It suffices to prove that  $\varphi(e_{uv}) = e_u \varphi(e_{uv}) = \varphi(e_{uv})e_v$  for all u < v, because in this case

$$\varphi(e_{uv}) = e_u \varphi(e_{uv}) = e_u \varphi(e_{uv}) e_v = \varphi(e_{uv})(u, v) e_{uv},$$

so we define  $\sigma(u, v) = \varphi(e_{uv})(u, v) \in F^*$ .

Case 1.  $u, v \in X \setminus \{x\}$ . Then  $\varphi(e_x)\varphi(e_{uv}) = 0$  by Lemma 3.1 (ii)a, which implies  $e_x\varphi(e_{uv}) = 0$  by (21). Similarly,  $\varphi(e_z)\varphi(e_{uv}) = 0$  for all  $z \neq u, x$  by Lemma 3.5, so  $e_z\varphi(e_{uv}) = 0$  for all  $z \neq u, x$ . It follows that

$$\varphi(e_{uv}) = \delta \cdot \varphi(e_{uv}) = \left(\sum_{z \in X} e_z\right) \varphi(e_{uv}) = e_u \varphi(e_{uv}).$$

In a similar way one proves using Lemma 3.1 (ii)b, Lemma 3.5 and (21) that  $\varphi(e_{uv}) = \varphi(e_{uv})e_v$ ,

Case 2. x = u < v. Then  $\varphi(e_{uv})\varphi(e_z) = \varphi(e_{xv})\varphi(e_z) = 0$  for all  $z \neq v$  by Lemma 3.1 (iii)b, whence  $\varphi(e_{uv})e_z = 0$  for all  $z \neq v$  in view of (21). Therefore,

$$\varphi(e_{uv}) = \varphi(e_{uv}) \cdot \delta = \varphi(e_{uv})e_v. \tag{23}$$

Now,  $\varphi(e_z)\varphi(e_{uv}) = \varphi(e_z)\varphi(e_{xv}) = 0$  for  $z \neq x, v$  by Lemma 3.1 (vi)b, so  $e_z\varphi(e_{uv}) = 0$  for  $z \neq u, v$  by (21). Consequently, using (23) and Corollary 3.13, we get

$$\varphi(e_{uv}) = \delta \cdot \varphi(e_{uv}) = e_u \varphi(e_{uv}) + e_v \varphi(e_{uv}) = e_u \varphi(e_{uv}) + e_v \varphi(e_{uv}) e_v = e_u \varphi(e_{uv}),$$

as needed.

Case 3. u < v = x. This case is similar to Case 2.

**Lemma 3.15.** Let X be connected. Then  $\varphi(\delta) = \pm \delta$ .

*Proof.* Using (21) we write

$$\varphi(\delta) = \sum_{z \in X} \varphi(e_z) = \sum_{z \in X} \alpha_z e_z.$$

It suffices to show that  $\alpha_z = \alpha_x$  for all  $z \in X$ . We know by Lemma 3.12 that  $\alpha_u = \alpha_x$  for all u < x and  $\alpha_v = \alpha_x$  for all v > x. Moreover, for all u < v, in view of Lemma 3.14 and (21) we obtain

$$\alpha_u \varphi(e_{uv}) = \sigma(u, v) \alpha_u e_{uv} = \sigma(u, v) \alpha_u e_u \cdot e_{uv} = \sigma(u, v) \varphi(e_u) \cdot e_{uv} = \varphi(e_u) \varphi(e_{uv}).$$
  
Similarly

$$\alpha_v \varphi(e_{uv}) = \sigma(u, v) \alpha_v e_{uv} = \sigma(u, v) e_{uv} \cdot \alpha_v e_v = \sigma(u, v) e_{uv} \varphi(e_v) = \varphi(e_{uv}) \varphi(e_v).$$

However,  $\varphi(e_u)\varphi(e_{uv}) = \varphi(e_{uv})\varphi(e_v)$  for all u < v from  $X \setminus \{x\}$  by Lemma 3.5 and Proposition 2.1. Hence,  $\alpha_u = \alpha_v$ . Since X is connected, we are done.

Replacing  $\varphi$  by  $-\varphi$ , if necessary, we may assume that

$$\varphi(\delta) = \delta. \tag{24}$$

Then, for all  $z \in X$ , we have

$$\varphi(e_z) = e_z. \tag{25}$$

In particular,  $c = \varphi(e_{X \setminus \{x\}}) = e_{X \setminus \{y\}}$ , so  $\varphi = \psi$  on  $I(X \setminus \{x\}, F)$ . We will now show that these conditions guarantee that  $\varphi$  is a multiplicative automorphism (6) of I(X, F).

**Lemma 3.16.** Let X be connected. Then  $\varphi$  is a multiplicative automorphism of I(X, F).

*Proof.* Consider  $\sigma : X_{\leq}^2 \to F^*$  from Lemma 3.14 and extend it to  $\sigma : X_{\leq}^2 \to F^*$  by means of  $\sigma(z, z) = 1$  for all  $z \in X$ . Observe that (22) now holds for all  $u \leq v$  thanks to (25). Thus, it only remains to prove that  $\sigma$  satisfies (5). Take arbitrary  $u \leq v \leq w$ . We need to show that  $\sigma(u, v)\sigma(v, w) = \sigma(u, w)$ . Since  $\sigma(z, z) = 1$  for all  $z \in X$ , it suffices to assume u < v < w. Consider the following cases.

Case 1.  $x \notin \{u, v, w\}$ . Then  $\sigma(u, v)\sigma(v, w) = \sigma(u, w)$  because  $\varphi = \psi$  on  $I(X \setminus \{x\}, F)$  and  $\psi(e_{uw}) = \psi(e_{uv})\psi(e_{vw})$ .

Case 2. x = u. Since  $(e_x + e_{xv} - e_{xw})(\delta - e_{xv} + e_{vw}) = e_x$ , by (22) and (24) we have

$$(e_x + \sigma(x, v)e_{xv} - \sigma(x, w)e_{xw})(\delta - \sigma(x, v)e_{xv} + \sigma(v, w)e_{vw}) = e_x,$$

which gives  $\sigma(x, v)\sigma(v, w) = \sigma(x, w)$ .

Case 3. x = w. This case is proved similarly to Case 2 using the product  $(\delta - e_{vx} + e_{uv})(e_x + e_{vx} - e_{ux}) = e_x$ .

Case 4. x = v. The result follows by applying  $\varphi$  to the product  $(e_x + e_{ux} + e_{xw} + e_{uw} - e_u)(\delta - e_{xw} + e_{ux} - e_u) = e_x$ .

We can finally prove the main result of our work.

**Theorem 3.17.** Let X be a finite connected poset and  $x, y \in X$ . Let  $\epsilon, \eta \in I(X, F)$  be primitive idempotents such that  $\epsilon_D = e_x$  and  $\eta_D = e_y$ . There exists a bijective linear map  $\varphi : I(X, F) \to I(X, F)$  preserving products equal to  $\epsilon$  and  $\eta$  if and only if there exists an automorphism of X mapping x to y, in which case  $\varphi$  is either an automorphism of I(X, F) or the negative of an automorphism of I(X, F).

*Proof.* Assume that  $\varphi : I(X, F) \to I(X, F)$  preserves products equal to  $\epsilon$  and  $\eta$ . By Lemmas 3.10, 3.11 and 3.16, up to an inner automorphism,  $\varphi$  is of the form  $\pm \widehat{\lambda} \circ M_{\sigma}$ , where  $\lambda$  is an automorphism of X with  $\lambda(x) = y$  and  $M_{\sigma}$  is a multiplicative automorphism of I(X, F).

Conversely, if  $\lambda$  is an automorphism of X with  $\lambda(x) = y$ , then  $\hat{\lambda} \in \operatorname{Aut}(I(X, F))$ preserves products equal to  $e_x$  and  $e_y$ . Composing  $\hat{\lambda}$  with appropriate inner automorphisms of I(X, F), we get  $\varphi \in \operatorname{Aut}(I(X, F))$  which preserves products equal to  $\epsilon$  and  $\eta$ .

**Corollary 3.18.** Let  $\epsilon, \eta \in T_n(F)$  be primitive idempotents. There exists a bijective linear map  $\varphi : T_n(F) \to T_n(F)$  preserving products equal to  $\epsilon$  and  $\eta$  if and only if  $\epsilon$  and  $\eta$  have the same diagonal, in which case  $\varphi$  is either an automorphism or the negative of an automorphism of  $T_n(F)$ .

*Proof.* For,  $T_n(F) \cong I(X, F)$ , where X is a chain of length n-1. Then,  $\epsilon$  and  $\eta$ , seen as elements of I(X, F), have diagonals  $e_x$  and  $e_y$  for some  $x, y \in X$ . Since  $\operatorname{Aut}(X) = \{\operatorname{id}\}$ , the result follows.

#### Acknowledgements

The first author was partially supported by Junta de Andalucía (grants FQM375 and PY20\_00255). The second author was partially supported by CNPq (process 404649/2018-1). We thank the referee for pointing out inaccuracies throughout the text and suggestions that improved its clarity.

#### References

- ARTIN, E. Geometric algebra, vol. 3 of Intersci. Tracts Pure Appl. Math. Interscience Publishers, New York, NY, 1957.
- [2] BACLAWSKI, K. Automorphisms and derivations of incidence algebras. Proc. Amer. Math. Soc. 36, 2 (1972), 351–356.
- [3] BURGOS, M. J., SÁNCHEZ, J. C., AND PERALTA, A. M. Linear maps between C\*-algebras that are \*-homomorphisms at a fixed point. *Quaest. Math.* 42, 2 (2019), 151–164.
- [4] CATALANO, L. On maps characterized by action on equal products. J. Algebra 511 (2018), 148–154.
- [5] CATALANO, L. On maps preserving products equal to a rank-one idempotent. *Linear Multi*linear Algebra 69, 4 (2021), 673–680.
- [6] CATALANO, L., AND CHANG-LEE, M. On maps preserving rank-one nilpotents. *Linear Multilinear Algebra 69*, 16 (2021), 3092–3098.
- [7] CATALANO, L., HSU, S., AND KAPALKO, R. On maps preserving products of matrices. *Linear Algebra Appl. 563* (2019), 193–206.
- [8] CATALANO, L., AND JULIUS, H. On maps preserving products equal to a diagonalizable matrix. Commun. Algebra 49, 10 (2021), 4334–4344.
- [9] CATALANO, L., AND JULIUS, H. On maps preserving products equal to fixed elements. J. Algebra 575 (2021), 220–232.
- [10] CHEBOTAR, M. A., KE, W.-F., LEE, P.-H., AND SHIAO, L.-S. On maps preserving products. Can. Math. Bull. 48, 3 (2005), 355–369.
- [11] CHEBOTAR, M. A., KE, W.-F., LEE, P.-H., AND WONG, N.-C. Mappings preserving zero products. Stud. Math. 155, 1 (2003), 77–94.
- [12] DROZD, Y., AND KOLESNIK, P. Automorphisms of incidence algebras. Comm. Algebra 35, 12 (2007), 3851–3854.
- [13] GARCÉS, J. J., AND KHRYPCHENKO, M. Potent preservers of incidence algebras. Linear Algebra Appl. 635 (2022), 171–200.
- [14] GINSBURG, V., JULIUS, H., AND VELASQUEZ, R. On maps preserving Lie products equal to a rank-one nilpotent. *Linear Algebra Appl. 593* (2020), 212–227.
- [15] JULIUS, H. On maps preserving Lie products equal to  $e_{11} e_{22}$ . Linear Multilinear Algebra 69, 14 (2021), 2620–2628.
- [16] KHRIPCHENKO, N. S. Automorphisms of finitary incidence rings. Algebra and Discrete Math. 9, 2 (2010), 78–97.
- [17] KHRIPCHENKO, N. S., AND NOVIKOV, B. V. Finitary incidence algebras. Comm. Algebra 37, 5 (2009), 1670–1676.
- [18] SPIEGEL, E., AND O'DONNELL, C. J. Incidence Algebras. New York, NY: Marcel Dekker, 1997.
- [19] STANLEY, R. Structure of incidence algebras and their automorphism groups. Bull. Am. Math. Soc. 76 (1970), 1236–1239.

DEPARTAMENTO DE MATEMÁTICA APLICADA A LA INGENIERÍA INDUSTRIAL, ETSIDI, UNI-VERSIDAD POLITÉCNICA DE MADRID, MADRID, SPAIN Email address: j.garces@upm.es

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE SANTA CATARINA, CAMPUS REITOR JOÃO DAVID FERREIRA LIMA, FLORIANÓPOLIS, SC, CEP: 88040-900, BRAZIL Email address: nskhripchenko@gmail.com