Cosserat micropolar elasticity: classical Eringen vs. dislocation form

Ionel-Dumitrel Ghiba¹, Gianluca Rizzi²,

Angela Madeo³ and Patrizio Neff⁴

June 7, 2022

Abstract

In this paper we do a comparative presentation of the linear isotropic Cosserat elastic model from two perspectives: the classical Mindlin-Eringen-Nowacki description in terms of a microrotation vector and a new formulation in terms of a skew-symmetric matrix and a curvature energy in dislocation form. We provide the reader with an alternative representation of the energy for the isotropic Cosserat model to ease the comparison with the relaxed micromorphic model and the geometrically nonlinear Cosserat elastic model.

Keywords: Cosserat micropolar model, relaxed micromorphic model, microrotation vector, Cosserat couple modulus, parameter identification, notations, dislocation density tensor, Nye's formula.

Contents

1	Introduction	2				
6 Constitutive assumptions in the new parameter set 1 6.1 Positive definiteness of the elastic energy density and well-posedness of the solution 1 6.2 Real plane waves in isotropic Cosserat elastic solids 1 6.3 Strong ellipticity (rank-one convexity, Legendre-Hadamard ellipticity) 1 conditions 1 7 Further identifications for micropolar constants 1 7.1 Eringen's initial notation: the notation used by Lakes 1 7.2 Mindlin's notation 1 7.3 Nowacki's notation 1 8 Linear Cosserat elasticity as a particular case of the relaxed 1 8.1 Identification of the relaxed micromorphic parameters in Mindlin's notation 1 9 Nonlinear isotropic Cosserat model 2		3				
3	Classical linear Cosserat model in Eringen notation	4				
4	Cosserat theory of isotropic elastic solids in dislocation format	7				
5 Identification of parameters in the dislocation formulation						
6	 6.1 Positive definiteness of the elastic energy density and well-posedness of the solution	13				
7	7.1 Eringen's initial notation: the notation used by Lakes	16				
8	micromorphic model					
9	Nonlinear isotropic Cosserat model	20				
10) Conclusions	21				
R	eferences	4 7 10 12 12 12 12 13 15 16 16 16 16 18 18 19 19 20 21 24				
	¹ Department of Mathematics, Alexandru Ioan Cuza University of Iaşi, Blvd. Carol I, no. 11, 700506 Iaşi, Romania; and Oo	ctav				

Mayer Institute of Mathematics of the Romanian Academy, Iaşi Branch, 700505 Iaşi

²Chair of Architecture and Civil Engineering, Technische Universität Dortmund, August-Schmidt-Str. 8, 44227 Dortmund, Germany

³Head of Chair of Continuum Mechanics, Faculty of Architecture and Civil Engineering, Technische Universität Dortmund, August-Schmidt-Str. 8, 44227 Dortmund, Germany

 $^{^4 {\}rm Head}$ of Chair for Nonlinear Analysis and Modelling, Fakultät für Mathematik, Universität Duisburg-Essen, Thea-Leymann-Straße 9, 45127 Essen, Germany

1 Introduction

Classical continuum mechanics considers material continua as simple point-continua with points having three displacement-degrees of freedom, and the response of a material to the displacement of its points is characterized by a symmetric Cauchy force-stress tensor presupposing that the transmission of loads through surface elements is uniquely determined by a force vector, neglecting couples. Such a model may be insufficient for the description of certain physical phenomena. Non-classical behaviour due to microstructural effects is observed mostly in regions of high strain gradients, e.g. at notches, holes or cracks. The Cosserat model is one of the best-known generalized continuum models. In that model, the classical translational degrees of freedom are augmented with a so-called microrotation, providing three additional degrees of freedom.

The Cosserat model has emerged from the seminal work of the brothers Francois and Eugene Cosserat at the turn of the last century ("Theorie des corps deformables." 1909, Review 1912, english translation by D. Delphenich 2007) [7]. They attempted to unify field theories embracing mechanics, optics and electro-dynamics through a common principle of least action (Euclidean action). Their main aim was to produce the correct general form of the energy for the variational problem. Postulating the invariance of energy under Euclidean transformations they were able to derive the equations of balance of forces and balance of angular momentum in a geometrically nonlinear format. However, they never wrote down any constitutive equations and never considered a linearized model as we will do here.

Compared to classical linear elasticity the linear Cosserat model features three additional, independent degrees of freedom, related to the rotation of each particle which need not coincide with the macroscopic rotation of the continuum at the same point. In the simplest isotropic case, one coupling constant, here called Cosserat couple modulus $\mu_c > 0$ and three internal length scale parameters need to be determined/measured in addition to the two classical elastic Lamé-constants. In the following, let us concentrate first on the static linear setting with quadratic free energy. In this linear setting, the model is in fact already given by the german scientist W. Voigt in 1887 [66]. Also P. Duhem in 1893 [9] had noticed that various phenomena which seemed incompatible with classical continuum mechanics could be described as effects of direction, and he suggested that materials be visualized as sets of points having vectors attached to them, i.e., oriented or polar media. However, for historical precision, the date of birth of a polar continuum is the year 1686, when Jakob Bernoulli introduced angular momentum as a postulate, independent of balance of momentum, see the work by Truesdell [65].

One of the essential features of polar continua is that the force stress tensor is not necessarily symmetric¹, and the balance of angular momentum equation has to be modified accordingly. All theories in which the stress tensor may not be symmetric can be regarded as polar-continua. The non-symmetry of the stress tensor may also appears also if higher order deformation gradients are included in the free energy, instead of only the first order gradients. Both such theories typically predict a size-effect, meaning that smaller samples of the same material behave relatively stiffer than larger samples. This is an often observed experimental fact, but completely neglected in the classical approach. It implies that some of the additional parameters in the Cosserat model define a length-scale present in the material.

The linear static Cosserat model may be posed in a variational format as a two-field minimization problem for the usual displacement u and the three entries of the infinitesimal microrotation \mathbf{A} , which is an element of the Lie-algebra $\mathfrak{so}(3)$ of skew symmetric matrices. The mathematical analysis of linear micropolar models is fairly well established with a wealth of analytical solutions for boundary value problems, existence and uniqueness theorems and continuous dependence results. It is usually based on a uniform positivity assumption on the free energy which sets it apart from linear elasticity in that Korn's inequality is not needed.

As often the case, notation is a nightmare. Unfortunately, in earlier works of Eringen and others, the interpretation of elastic constants have been misleading with the consequence of giving erroneous parameter ranges for positive definiteness. This has been corrected by S.C. Cowin [8] and the more recent book by Eringen [14].

Since any skew-symmetric matrix $\mathbf{A} \in \mathfrak{so}(3)$ can be identified with its axial vector $\vartheta = \operatorname{axl}(\mathbf{A})$, practically all previous developments for the linear Cosserat model have automatically opted for a presentation of the model in terms of the displacement $u \in \mathbb{R}^3$ and the microrotation vector $\vartheta \in \mathbb{R}^3$. The curvature expression is then simply a quadratic form of the second order tensor $\mathcal{D}\vartheta \in \mathbb{R}^{3\times 3}$. The advantage of a concise formulation is, however, bought at the expense of transparency of the modelling. Moreover, the advantage of using a microrotation

¹But it can still be chosen to be symmetric in the nonlinear Cosserat model by setting the Cosserat couple modulus $\mu_c = 0$ [47].

vector is immediately lost when considering a geometrically nonlinear Cosserat model in which there appears an orthogonal matrix (the trièdre mobil) $\overline{\mathbf{R}} \in SO(3)$. Here, the extraction of a microrotation vector is algebraically difficult and completely unnatural. Therefore, we advocate a presentation of the linear Cosserat model fully in terms of skew-symmetric matrices $\mathbf{A} \in \mathfrak{so}(3)$. The remaining question is how to concisely express the curvature energy, now naively a quadratic term in the third order tensor $\mathbf{DA} \in \mathbb{R}^{3\times3\times3}$. However, using \mathbf{DA} can be avoided by taking recourse to the so called Nye's formula [59], expressing $\mathbf{D}\vartheta = \mathrm{Daxl}(\mathbf{A})$ in terms of the matrix Curl \mathbf{A} , i.e.,

$$-\operatorname{Curl} \mathbf{A} = (\operatorname{D}\operatorname{axl} \mathbf{A})^T - \operatorname{tr} \left[(\operatorname{D}\operatorname{axl} \mathbf{A})^T \right] \cdot \mathbb{1}, \qquad \qquad \operatorname{D}\operatorname{axl} \mathbf{A} = -(\operatorname{Curl} \mathbf{A})^T + \frac{1}{2} \operatorname{tr} \left[(\operatorname{Curl} \mathbf{A})^T \right] \cdot \mathbb{1}, \qquad (1.1)$$

and using

$$(\mathbf{D}\mathbf{A})_{ijl} = A_{ij,l} = -\epsilon_{ijk} \operatorname{axl}(\mathbf{A})_{k,l} = -\epsilon_{ijk} (\operatorname{Daxl}(\mathbf{A}))_{kl}.$$
(1.2)

We note that Curl **A** is a second order tensor which effectively controls all partial derivatives of **A**, see [52]. For reasons connected to plasticity theory, we will call Curl **A** the dislocation density tensor. Thus, our new Cosserat formulation can be fully expressed in the triplet $(u, \mathbf{A}, \operatorname{Curl} \mathbf{A})$ and the main aim of the current contribution is a complete comparison of the microrotation vector approach in $(u, \vartheta, \operatorname{D} \vartheta)$ versus the new representation. We can already note that in terms of $(u, \mathbf{A}, \operatorname{Curl} \mathbf{A})$ there is a straightforward extension to the geometrically nonlinear Cosserat model as well as a transparent way to relate to the family of micromorphic models. Especially, the newly developed relaxed micromorphic model practically uses the same curvature expression as our new formulation. The linear Cosserat model can then be obtained as a singular limit of the relaxed micromorphic model.

Cosserat media may serve as a model for the prediction of size-effects in foam like structures (like bones) or cellular materials. This approach has been championed by Lakes [26, 67, 24, 27, 28, 29, 30]. He determines Cosserat parameters by careful size experiments. His values seem to be the only consistent choice of Cosserat parameters ever given for the linear isotropic model. Interestingly, his values have been rejected by a prominent proponent of the micropolar model because Lakes values make the Cosserat free energy only positive semi-definite instead of some supposed positive definiteness in the curvature term. A careful mathematical inspection of analytical solution reveals, however, that Lakes parameter range is a must in order to avoid certain unphysical stiffening behaviour of the Cosserat model for very small samples [50, 28].

After a short section introducing the main notations of this paper, in Section 3 we present the linear Cosserat model for isotropic elastic materials in the Eringen's microrotation vector notation. In Section 4 we give an alternative formulation of the linear Cosserat model for isotropic elastic materials in terms of a new strain measure, the dislocation tensor. In Section 5 we provide the identification between all constitutive parameters involved in the classical formulation given by Eringen and the constitutive parameters involved in our formulation. In Section 6 we express some well known constitutive requirements, e.g., positive definiteness of the internal energy density, real plane waves propagation, Legendre-Hadamard ellipticity (strong ellipticity) condition, in terms of the constitutive coefficients considered in our new formulation. In Section 7 we compare the constitutive parameters with those used in other formulations, i.e., Nowacki's formulation and Eringen's initial formulation. In Section 8 we show that our formulation is a particular case of a formulation given in a more general model, i.e., the relaxed micromorphic model. In Section 9 we exhibit that our linear formulation follows directly from a nonlinear model for isotropic elastic Cosserat solids. In Section 10 we give a concluding table which facilitates a translation of the experimental results interpreted in terms of the coefficients from other formulation to the constitutive parameters used in our description.

2 Notation

We consider that the mechanical behaviour of a body accupying the unbounded regular region of three dimensional Euclidean space is modelled with the help of the Cosserat theory of linear isotropic elastic materials. We denote by n the outward unit normal on $\partial\Omega$. The body is referred to a fixed system of rectangular Cartesian axes $Ox_i(i = 1, 2, 3)$, $\{e_1, e_2, e_3\}$ being the unit vectors of these axes. Throughout this paper (when we do not specify else) Latin subscripts take the values 1, 2, 3. Everywhere we adopt the Einstein convention of summation over repeated indices if not differently specified.

In the following, we recall some useful notations for the present work. For $a, b \in \mathbb{R}^{3\times3}$ we let $\langle a, b \rangle_{\mathbb{R}^3}$ denote the scalar product on \mathbb{R}^3 with associated vector norm $||a||^2 = \langle a, a \rangle$. We denote by $\mathbb{R}^{3\times3}$ the set of real 3×3 second order tensors, written with capital letters. Matrices will be denoted by bold symbols, e.g. $\mathbf{X} \in \mathbb{R}^{3\times3}$, while X_{ij} will denote its component. The standard Euclidean product on $\mathbb{R}^{3\times3}$ is given by $\langle \mathbf{X}, \mathbf{Y} \rangle_{\mathbb{R}^{3\times3}} = \operatorname{tr}(\mathbf{X} \mathbf{Y}^T)$, and thus, the Frobenious tensor norm is $||\mathbf{X}||^2 = \langle \mathbf{X}, \mathbf{X} \rangle_{\mathbb{R}^{3\times3}}$. In the following we omit the index $\mathbb{R}^3, \mathbb{R}^{3\times3}$. The identity tensor on $\mathbb{R}^{3\times3}$ will be denoted by $\mathbb{1}$, so that $\operatorname{tr}(\mathbf{X}) = \langle \mathbf{X}, \mathbb{1} \rangle$. We let Sym denote the set of symmetric tensors. We adopt the usual abbreviations of Lie-algebra theory, i.e., $\mathfrak{so}(3) := \{\mathbf{A} \in \mathbb{R}^{3\times3} | \mathbf{A}^T = -\mathbf{A}\}$ is the Lie-algebra of skew-symmetric tensors and $\mathfrak{sl}(3) := \{\mathbf{X} \in \mathbb{R}^{3\times3} | \operatorname{tr}(\mathbf{X}) = 0\}$ is the Lie-algebra of traceless tensors. For all $\mathbf{X} \in \mathbb{R}^{3\times3}$ we set sym $\mathbf{X} = \frac{1}{2}(\mathbf{X}^T + \mathbf{X}) \in \operatorname{Sym}(3)$, skew $\mathbf{X} = \frac{1}{2}(\mathbf{X} - \mathbf{X}^T) \in \mathfrak{so}(3)$ and the deviatoric (trace-free) part dev $\mathbf{X} = \mathbf{X} - \frac{1}{3}\operatorname{tr}(\mathbf{X}) \mathbb{1} \in \mathfrak{sl}(3)$ and we have the orthogonal Cartan-decomposition of the Lie-algebra $\mathfrak{gl}(3) = \{\mathfrak{sl}(3) \cap \operatorname{Sym}(3)\} \oplus \mathfrak{so}(3) \oplus \mathbb{R} \cdot \mathbb{1}$,

$$\mathbf{X} = \operatorname{dev} \operatorname{sym} \mathbf{X} + \operatorname{skew} \mathbf{X} + \frac{1}{3} \operatorname{tr}(\mathbf{X}) \mathbb{1}.$$
(2.3)

We use the canonical identification of \mathbb{R}^3 with $\mathfrak{so}(3)$, and, for

$$\mathbf{A} = \begin{pmatrix} 0 & -a_3 & a_2 \\ a_3 & 0 & -a_1 \\ -a_2 & a_1 & 0 \end{pmatrix} \in \mathfrak{so}(3)$$
(2.4)

we define the operators axl $: \mathfrak{so}(3) \to \mathbb{R}^3$ and $\mathbf{Anti} : \mathbb{R}^3 \to \mathfrak{so}(3)$ through

$$\operatorname{axl} \mathbf{A} := (a_1, a_2, a_3)^T, \qquad \mathbf{A} \cdot v = (\operatorname{axl} \mathbf{A}) \times v, \qquad (\mathbf{Anti}(v))_{ij} = -\epsilon_{ijk} v_k, \qquad \forall v \in \mathbb{R}^3,$$
$$(\operatorname{axl} \mathbf{A})_k = -\frac{1}{2} \epsilon_{ijk} \mathbf{A}_{ij} = \frac{1}{2} \epsilon_{kij} A_{ji}, \qquad A_{ij} = -\epsilon_{ijk} (\operatorname{axl} \mathbf{A})_k =: \mathbf{Anti}(\operatorname{axl} \mathbf{A})_{ij}, \qquad (2.5)$$

where ϵ_{ijk} is the totally antisymmetric third order permutation tensor.

For a regular enough function $f(t, x_1, x_2, x_3)$, $f_{,t}$ denotes the derivative with respect to the time t, while $\frac{\partial f}{\partial x_i}$ and $f_{,i}$ denotes the *i*-component of the gradient Df. For vector fields $u = (u_1, u_2, u_3)^T$ with $u_i \in H^1(\Omega) = \{u_i \in L^2(\Omega) \mid Du_i \in L^2(\Omega)\}, i = 1, 2, 3$, we define $Du := (Du_1 \mid Du_2 \mid Du_3)^T$.

The corresponding Sobolev-space will be also denoted by $\mathrm{H}^{1}(\Omega)$. For vector fields u with components in $\mathrm{H}^{1}(\Omega)$ and tensor fields P with rows in $\mathrm{H}(\mathrm{curl};\Omega)$, resp. $\mathrm{H}(\mathrm{div};\Omega)$, i.e.,

$$u = \begin{pmatrix} u_1 \\ u_2 \\ u_3 \end{pmatrix}, u_i \in \mathrm{H}^1(\Omega), \quad P = \begin{pmatrix} P_{11} & P_{12} & P_{13} \\ P_{21} & P_{22} & P_{23} \\ P_{31} & P_{32} & P_{33} \end{pmatrix} \quad P^T e_i \in \mathrm{H}(\mathrm{curl}\,;\Omega) \quad \text{resp.} \quad P^T e_i \in \mathrm{H}(\mathrm{div}\,;\Omega) \quad (2.6)$$

we define

$$\operatorname{curl} u = \begin{pmatrix} u_{3,2} - u_{2,3} \\ u_{1,3} - u_{3,1} \\ u_{2,3} - u_{3,1} \end{pmatrix} = (\epsilon_{ijk} u_{k,j})_{i=1,2,3}, \qquad \operatorname{D} u = \begin{pmatrix} u_{1,1} & u_{1,2} & u_{1,3} \\ u_{2,1} & u_{2,2} & u_{2,3} \\ u_{3,1} & u_{3,2} & u_{3,3} \end{pmatrix} = (u_{i,j})_{i,j=1,2,3}, \qquad (2.7)$$
$$\operatorname{Curl} \mathbf{P} = \begin{pmatrix} [\operatorname{curl} (P_{11} & P_{12} & P_{13})^T]^T \\ [\operatorname{curl} (P_{21} & P_{22} & P_{23})^T]^T \end{pmatrix} = \begin{pmatrix} P_{13,2} - P_{12,3} & P_{11,3} - P_{13,1} & P_{12,3} - P_{13,1} \\ P_{23,2} - P_{22,3} & P_{21,3} - P_{23,1} & P_{22,3} - P_{23,1} \end{pmatrix} = (\epsilon_{jmn} P_{in,m})_{i,j=1,2,3},$$

$$\left(\begin{bmatrix} \operatorname{curl} \left(P_{31} \ P_{32} \ P_{33} \right)^T \right)^T \right)^T \left(P_{33,2} - P_{32,3} \ P_{31,3} - P_{33,1} \ P_{32,3} - P_{33,1} \right)^T \right)$$

$$\operatorname{Div} \mathbf{P} = \begin{pmatrix} \operatorname{div} \left(P_{11} \ P_{12} \ P_{13} \right)^T \\ \operatorname{div} \left(P_{21} \ P_{22} \ P_{23} \right)^T \\ \operatorname{div} \left(P_{31} \ P_{32} \ P_{33} \right)^T \end{pmatrix} = \begin{pmatrix} P_{11,1} + P_{12,2} + P_{13,3} \\ P_{21,1} + P_{22,2} + P_{23,3} \\ P_{31,1} + P_{32,2} + P_{33,3} \end{pmatrix} = \left(P_{ij,j} \right)_{i,j=1,2,3}.$$

3 Classical linear Cosserat model in Eringen notation

In the isotropic linear Cosserat model the kinematics is described through a vector field, the displacement $u: [0,T] \times \Omega \subset \mathbb{R}^3 \to \mathbb{R}^3$, and a skew-symmetric tensor, the micro-rotation tensor $\mathbf{A}: [0,T] \times \Omega \subset \mathbb{R}^3 \to \mathfrak{so}(3)$. Being skew-symmetric, the micro-rotation tensor of \mathbf{A} is fully determined by its axial vector, i.e.,

$$\vartheta := \operatorname{axl} \mathbf{A}, \qquad \operatorname{Anti} \vartheta = \mathbf{A},$$
(3.1)

which is called the *micro-rotation vector*.

The linear strain tensors considered in the Eringen formulation [14, Page 104] are

$$\mathbf{e}_{ji}^* = u_{i,j} + \epsilon_{ijm} \,\vartheta_m, \qquad \qquad \mathbf{\hat{s}}_{ji} = \vartheta_{j,i}. \tag{3.2}$$

In a compact matrix form these strain tensors are

$$\mathbf{e}^* = (\mathbf{D}u)^T + \mathbf{Anti}\,\vartheta = (\mathbf{D}u)^T + \mathbf{A}, \qquad \mathbf{\hat{\kappa}} = \mathbf{D}\vartheta = \mathbf{D}(\mathbf{axl}\mathbf{A}). \tag{3.3}$$

Following Eringen [14, Page 111], the internal density energy for isotropic materials is

$$W = \frac{1}{2} \Big[(\mu^* + \varkappa) \, \mathbf{e}_{ji}^* \mathbf{e}_{ji}^* + \mu^* \, \mathbf{e}_{ji}^* \mathbf{e}_{ij}^* + \lambda \, \mathbf{e}_{ii}^* \mathbf{e}_{jj}^* + \gamma \, \mathbf{\mathfrak{K}}_{ji} \, \mathbf{\mathfrak{K}}_{ji} + \beta \, \mathbf{\mathfrak{K}}_{ji} \, \mathbf{\mathfrak{K}}_{ij} + \alpha \, \mathbf{\mathfrak{K}}_{ii} \, \mathbf{\mathfrak{K}}_{jj} \Big].$$
(3.4)

In a compact form, without involving the notations in indices, the expression of the internal density energy is

$$W = \frac{1}{2} \Big[(\mu^* + \varkappa) \langle \mathbf{e}^*, \mathbf{e}^* \rangle + \mu^* \langle \mathbf{e}^*, \mathbf{e}^{*T} \rangle + \lambda \left(\operatorname{tr}(\mathbf{e}^*) \right)^2 + \gamma \langle \mathbf{\mathfrak{K}}, \mathbf{\mathfrak{K}} \rangle + \beta \langle \mathbf{\mathfrak{K}}, \mathbf{\mathfrak{K}}^T \rangle + \alpha \left(\operatorname{tr}(\mathbf{\mathfrak{K}}) \right)^2 \Big].$$
(3.5)

Let us remark that

$$\begin{split} W &= \frac{1}{2} \Big[(\mu^* + \varkappa) \langle (\operatorname{sym} \mathbf{e}^* + \operatorname{skew} \mathbf{e}^*), (\operatorname{sym} \mathbf{e}^* + \operatorname{skew} \mathbf{e}^*) \rangle \\ &+ \mu^* \langle (\operatorname{sym} \mathbf{e}^* + \operatorname{skew} \mathbf{e}^*), (\operatorname{sym} \mathbf{e}^* - \operatorname{skew} \mathbf{e}^*) + \lambda \left(\operatorname{tr}(\operatorname{sym} \mathbf{e}^*) \right)^2 \\ &+ \gamma \langle (\operatorname{sym} \mathbf{\mathfrak{K}} + \operatorname{skew} \mathbf{\mathfrak{K}}), (\operatorname{sym} \mathbf{\mathfrak{K}} + \operatorname{skew} \mathbf{\mathfrak{K}}) \rangle \\ &+ \beta \langle (\operatorname{sym} \mathbf{\mathfrak{K}} + \operatorname{skew} \mathbf{\mathfrak{K}}), (\operatorname{sym} \mathbf{\mathfrak{K}} - \operatorname{skew} \mathbf{\mathfrak{K}}) \rangle + \alpha \left(\operatorname{tr}(\operatorname{sym} \mathbf{\mathfrak{K}}) \right)^2 \Big] \end{split} (3.6) \\ &= \frac{1}{2} \Big[(\mu^* + \varkappa) \| \operatorname{sym} \mathbf{e}^* \|^2 + (\mu^* + \varkappa) \| \operatorname{skew} \mathbf{e}^* \|^2 + \mu^* \| \operatorname{sym} \mathbf{e}^* \|^2 - \mu^* \| \operatorname{skew} \mathbf{e}^* \|^2 + \lambda \left(\operatorname{tr}(\operatorname{sym} \mathbf{e}^*) \right)^2 \\ &+ \gamma \| \operatorname{sym} \mathbf{\mathfrak{K}} \|^2 + \gamma \| \operatorname{skew} \mathbf{\mathfrak{K}} \|^2 + \beta \| \operatorname{sym} \mathbf{\mathfrak{K}} \|^2 - \beta \| \operatorname{skew} \mathbf{\mathfrak{K}} \|^2 + \alpha \left(\operatorname{tr}(\operatorname{sym} \mathbf{\mathfrak{K}}) \right)^2 \Big] \\ &= \frac{1}{2} \Big[(2 \, \mu^* + \varkappa) \| \operatorname{sym} \mathbf{e}^* \|^2 + \varkappa \| \operatorname{skew} \mathbf{e}^* \|^2 + \lambda \left(\operatorname{tr}(\operatorname{sym} \mathbf{\mathfrak{K}}) \right)^2 \\ &+ (\gamma + \beta) \| \operatorname{sym} \mathbf{\mathfrak{K}} \|^2 + (\gamma - \beta) \| \operatorname{skew} \mathbf{\mathfrak{K}} \|^2 + \alpha \left(\operatorname{tr}(\operatorname{sym} \mathbf{\mathfrak{K}}) \right)^2 \Big] . \end{aligned}$$

Since

$$2\operatorname{axl}(\operatorname{skew} \operatorname{D} u) = \operatorname{curl} u, \qquad 2(\operatorname{skew} \operatorname{D} u) = \operatorname{Anti}(\operatorname{curl} u), \tag{3.7}$$

and

$$\|\operatorname{skew} \mathrm{D}u - \mathbf{A}\|^{2} = 2\|\operatorname{axl}(\operatorname{skew} \mathrm{D}u - \operatorname{Anti}\vartheta)\|^{2} = 2\|\frac{1}{2}\operatorname{curl} u - \vartheta\|^{2} = \frac{1}{2}\|\operatorname{curl} u - 2\,\vartheta\|^{2}, \qquad (3.8)$$
$$\|\operatorname{skew} \mathrm{D}\vartheta\|^{2} = 2\|\operatorname{axl}(\operatorname{skew} \mathrm{D}\vartheta\|^{2} = 2\|\frac{1}{2}\operatorname{curl}\vartheta\|^{2} = \frac{1}{2}\|\operatorname{curl}\vartheta\|^{2}, \\\|\mathbf{\hat{\kappa}}\|^{2} = \|\mathrm{D}\vartheta\|^{2} = \|\mathrm{Daxl}(\mathbf{A})\|^{2} = \frac{1}{2}\|\mathrm{D}\mathbf{A}\|^{2}$$

we obtain the following equivalent alternative form of the internal energy density

$$W = \left[(\mu^* + \frac{\varkappa}{2}) \|\operatorname{sym} \operatorname{D} u\|^2 + \frac{\varkappa}{4} \|\operatorname{curl} u - 2\vartheta\|^2 + \frac{\lambda}{2} (\operatorname{tr}(\operatorname{D} u))^2 + \frac{(\gamma + \beta)}{2} \|\operatorname{sym} \operatorname{D} \vartheta\|^2 + \frac{(\gamma - \beta)}{4} \|\operatorname{curl} \vartheta\|^2 + \frac{\alpha}{2} (\operatorname{tr}(\operatorname{D} \vartheta))^2 \right]$$

$$= \left[(\mu^* + \frac{\varkappa}{2}) \|\operatorname{sym} \operatorname{D} u\|^2 + \frac{\varkappa}{4} \|\operatorname{curl} u - 2\vartheta\|^2 + \frac{\lambda}{2} (\operatorname{tr}(\operatorname{D} u))^2 + \frac{(\gamma + \beta)}{2} \|\operatorname{dev} \operatorname{sym} \operatorname{D} \vartheta\|^2 + \frac{(\gamma - \beta)}{4} \|\operatorname{curl} \vartheta\|^2 + \frac{3\alpha + \gamma + \beta}{6} (\operatorname{tr}(\operatorname{D} \vartheta))^2 \right]$$

$$= \left[(\mu^* + \frac{\varkappa}{2}) \|\operatorname{sym} \operatorname{D} u\|^2 + \frac{\varkappa}{4} \|\operatorname{curl} u - 2\vartheta\|^2 + \frac{\lambda}{2} (\operatorname{div} u)^2 + \frac{(\gamma + \beta)}{2} \|\operatorname{dev} \operatorname{sym} \operatorname{D} \vartheta\|^2 + \frac{(\gamma - \beta)}{4} \|\operatorname{curl} \vartheta\|^2 + \frac{3\alpha + \gamma + \beta}{6} (\operatorname{div} \vartheta)^2 \right].$$
(3.9)

In indices notation [14, Page 111] we have the following expression of the non-symmetric force stress and the couple stress tensor definition

$$\sigma_{ji} = \frac{\partial W}{\partial \mathbf{e}_{ji}^*} = (\mu^* + \varkappa) \, \mathbf{e}_{ji}^* + \mu^* \mathbf{e}_{ij}^* + \lambda \, \mathbf{e}_{kk}^* \delta_{ji} = (\mu^* + \varkappa) (u_{i,j} + \epsilon_{ijm} \vartheta_m) + \mu^* (u_{j,i} + \epsilon_{jim} \vartheta_m) + \lambda \, u_{k,k} \delta_{ji},$$
$$m_{ji}^* = \frac{\partial W}{\partial \mathfrak{K}_{ji}} = \beta \, \mathfrak{K}_{ij} + \gamma \, \mathfrak{K}_{ji} + \alpha \, \mathfrak{K}_{kk} \delta_{ji} = \beta \, \vartheta_{j,i} + \gamma \, \vartheta_{i,j} + \alpha \, \vartheta_{k,k} \delta_{ji}.$$
(3.10)

In a compact matrix description, the force stress and the couple stress tensor definition in the *classical form* are given by²

$$\boldsymbol{\sigma} = \mathbf{D}_{\mathbf{e}^*} W = (\mu^* + \varkappa) \mathbf{e}^* + \mu^* \mathbf{e}^{*T} + \lambda \operatorname{tr}(\mathbf{e}^*) \mathbb{1}$$

$$= (\mu^* + \varkappa) (\operatorname{sym} \mathbf{e}^* + \operatorname{skew} \mathbf{e}^*) + \mu^* (\operatorname{sym} \mathbf{e}^* - \operatorname{skew} \mathbf{e}^*) + \lambda \operatorname{tr}(\mathbf{e}^*) \mathbb{1}$$

$$= (2 \,\mu^* + \varkappa) \operatorname{sym} \mathbf{e}^* + \varkappa \operatorname{skew} \mathbf{e}^* + \lambda \operatorname{tr}(\mathbf{e}^*) \mathbb{1}$$

$$= (2 \,\mu^* + \varkappa) \operatorname{sym} \operatorname{D} u + \varkappa (\operatorname{skew}(\operatorname{D} u)^T + \operatorname{Anti} \vartheta) + \lambda \operatorname{tr}(\operatorname{D} u) \mathbb{1}$$

$$= (2 \,\mu^* + \varkappa) \operatorname{sym} \operatorname{D} u - \varkappa (\operatorname{skew}(\operatorname{D} u) - \operatorname{Anti} \vartheta) + \lambda \operatorname{tr}(\operatorname{D} u) \mathbb{1}$$

$$= (2 \,\mu^* + \varkappa) \operatorname{sym} \operatorname{D} u - \frac{\varkappa}{2} \operatorname{Anti}(\operatorname{curl} u - 2 \,\vartheta) + \lambda \operatorname{tr}(\operatorname{D} u) \mathbb{1},$$

$$\boldsymbol{m}^* = \mathbf{D}_{\mathbf{f}} W = \gamma \,\mathbf{f} + \beta \,\mathbf{f}^T + \alpha \operatorname{tr}(\mathbf{f}) \,\mathbb{1} = \beta \operatorname{D} \vartheta + \gamma (\operatorname{D} \vartheta)^T + \alpha \operatorname{tr}(\operatorname{D} \vartheta) \,\mathbb{1}.$$
(3.12)

By considering the action functional

$$\mathcal{A} = \int_0^T \int_\Omega \left(\frac{1}{2} \rho \| \dot{u} \|^2 + \rho j \| \dot{\vartheta} \|^2 - W \right) dv \, dt, \tag{3.13}$$

where j is a micro-inertia coefficient, we obtain in the absence of body loads, the motion equations in the *classical form*

$$\varrho u_{i,tt} = \sigma_{ji,j}, \qquad i = 1, 2, 3, \qquad (3.14)$$

$$\varrho j \vartheta_{i,tt} = m_{ji,j}^* + \epsilon_{ijk} \sigma_{jk}, \qquad i = 1, 2, 3,$$

equivalently, in matrix (**Div-Div**) form,

$$\varrho \, u_{,tt} = \operatorname{Div} \boldsymbol{\sigma}^{T},$$

$$\varrho \, j \, \vartheta_{,tt} = \operatorname{Div} \boldsymbol{m}^{*T} - 2 \operatorname{axl} (\operatorname{skew} \boldsymbol{\sigma}).$$
(3.15)

 2 These expressions agree with the form considered by Eremeyev et al. [12, Eq. (4.68)], where the internal energy is written as

$$W = \frac{\lambda}{2} \left[\operatorname{tr}(\varepsilon) \right]^2 + \frac{\mu^* + \varkappa}{2} \operatorname{tr}\left(\varepsilon \,\varepsilon^T\right) + \frac{\mu^*}{2} \operatorname{tr}(\varepsilon \,\varepsilon) + \frac{\beta_1}{2} \left[\operatorname{tr}(\mathfrak{K}) \right]^2 + \frac{\beta_2}{2} \operatorname{tr}\left(\mathfrak{K} \,\mathfrak{K}^T\right) + \frac{\beta_3}{2} \operatorname{tr}(\mathfrak{K} \,\mathfrak{K}).$$
(3.11)

In the absence of external body forces and external body moment, the PDE-system of the model [14, Pages 118-119] is

$$\rho u_{i,tt} = (\mu^* + \varkappa) u_{i,jj} + (\lambda + \mu^*) u_{j,ij} + \varkappa \epsilon_{ijk} \vartheta_{k,j}, \qquad i = 1, 2, 3$$

$$\rho j \vartheta_{i,tt} = (\alpha + \beta) \vartheta_{i,ij} + \gamma \vartheta_{i,jj} + \varkappa \epsilon_{ijk} u_{k,j} - 2 \varkappa \vartheta_i, \qquad i = 1, 2, 3.$$
(3.16)

In a compact form, the above PDE-system reads

$$\rho \frac{\partial^2 u}{\partial t^2} = (\mu^* + \varkappa) \Delta u + (\mu^* + \lambda) \operatorname{D}(\operatorname{div} u) + \varkappa \operatorname{curl} \vartheta, \qquad i = 1, 2, 3, \tag{3.17}$$

$$\rho j \frac{\partial^2 \vartheta}{\partial t^2} = \gamma \Delta \vartheta + (\alpha + \beta) \operatorname{D}(\operatorname{div} \vartheta) + \varkappa \left(\operatorname{curl} u - 2 \vartheta\right), \qquad i = 1, 2, 3.$$

Using the well-known identity

$$0 = -\operatorname{curl}\left(\operatorname{curl} u\right) + \mathcal{D}(\operatorname{div} u) - \Delta u \tag{3.18}$$

the PDE-system can be rewritten as

$$\rho \frac{\partial^2 u}{\partial t^2} = \mu^* \Delta u + (\mu^* + \lambda + \varkappa) \mathcal{D}(\operatorname{div} u) - \varkappa \operatorname{curl} (\operatorname{curl} u - \vartheta), \qquad (3.19)$$
$$\rho j \frac{\partial^2 \vartheta_i}{\partial t^2} = \gamma \Delta \vartheta + (\alpha + \beta) \mathcal{D}(\operatorname{div} \vartheta) + \varkappa (\operatorname{curl} u - 2 \vartheta).$$

4 Cosserat theory of isotropic elastic solids in dislocation format

Now we propose another form of the internal energy density and we show that after an identification of the parameters it coincides with the internal energy density proposed by Eringen. However, the new form of the internal energy density has the advantage to have the same structure as in the case of a more generalised theory, i.e., the relaxed micromorphic model [49, 36, 35, 37, 48], see Subsection 8 and the nonlinear Cosserat model [46], see Subsection 9.

We keep the physical meaning of the unknowns u and $\vartheta = axl(\mathbf{A})$ from the Cosserat theory, but we propose a new form of the Cosserat internal energy density, namely

$$W = \mu_{\rm e} \| \text{sym} (\mathrm{D}u - \mathbf{A}) \|^{2} + \mu_{\rm c} \| \text{skew} (\mathrm{D}u - \mathbf{A}) \|^{2} + \frac{\lambda_{\rm e}}{2} [\operatorname{tr} (\mathrm{D}u - \mathbf{A})]^{2} + \frac{\mu_{\rm e} L_{\rm c}^{2}}{2} \left[a_{1} \| \operatorname{dev} \operatorname{sym} \operatorname{Curl} \mathbf{A} \|^{2} + a_{2} \| \text{skew} \operatorname{Curl} \mathbf{A} \| + \frac{a_{3}}{3} \operatorname{tr} (\operatorname{Curl} \mathbf{A})^{2} \right],$$
(4.1)

where $(\mu_{\rm e}, \lambda_{\rm e})$, $\mu_{\rm c}, L_{\rm c}$ and (a_1, a_2, a_3) are the elastic moduli representing the parameters related to the mesoscale, the Cosserat couple modulus $\mu_{\rm c}$, the characteristic length $L_{\rm c}$, and the three general isotropic curvature parameters (nondimensional weights), respectively.

The new energy is expressed in terms of the strain and curvature tensors

$$\mathbf{e} = \mathrm{D}u - \operatorname{Anti} \vartheta = \mathrm{D}u - \mathbf{A} \in \mathbb{R}^{3 \times 3}, \qquad \boldsymbol{\alpha} := -\mathrm{Curl} \, \mathbf{A} \in \mathbb{R}^{3 \times 3}. \tag{4.2}$$

The stress and moment definition in the *dislocation form* are

$$\boldsymbol{\sigma} = \mathbf{D}_{\mathbf{e}} W = 2\mu_{\mathrm{e}} \operatorname{sym} \mathbf{D}u + 2\mu_{\mathrm{c}} \operatorname{skew} \mathbf{D}u - \mathbf{A} + \lambda_{\mathrm{e}} \operatorname{tr} (\mathbf{D}u) \,\mathbb{1}$$

$$= 2\mu_{\mathrm{e}} \operatorname{sym} \mathbf{D}u + 2\mu_{\mathrm{c}} (\operatorname{skew}(\mathbf{D}u) - \operatorname{\mathbf{Anti}} \vartheta) + \lambda_{\mathrm{e}} \operatorname{tr}(\mathbf{D}u) \,\mathbb{1}$$

$$= 2\mu_{\mathrm{e}} \operatorname{sym} \mathbf{D}u + \mu_{\mathrm{c}} \operatorname{\mathbf{Anti}} (\operatorname{curl} u - 2\,\vartheta) + \lambda_{\mathrm{e}} \operatorname{tr}(\mathbf{D}u) \,\mathbb{1}, \qquad (4.3)$$

$$\boldsymbol{m} = \mathbf{D}_{\boldsymbol{\alpha}} W = \mu_{\mathrm{e}} L_{\mathrm{c}}^{2} \left(a_{1} \operatorname{dev} \operatorname{sym} \operatorname{Curl} \mathbf{A} + a_{2} \operatorname{skew} \operatorname{Curl} \mathbf{A} + \frac{a_{3}}{3} \operatorname{tr} (\operatorname{Curl} \mathbf{A}) \,\mathbb{1} \right).$$

Considering the action functional

$$\mathcal{A} = \int_0^T \int_\Omega \left(\frac{1}{2} \rho \| u_{,t} \|^2 + \frac{1}{2} \rho \eta \tau_c^2 \| \mathbf{A}_{,t} \|^2 - W \right) dv \, dt \tag{4.4}$$

with $\eta \tau_c^2$ an inertia coefficient, $\eta > 0$ a nondimensional weight parameter and τ_c the internal characteristic time [14, page 163], we obtain the Euler-Lagrange equation, i.e., the following system in **Div-Curl-form** (the dislocation formulation)

$$\rho \, u_{,tt} = \operatorname{Div} \boldsymbol{\sigma} \,, \tag{4.5}$$
$$\tau_c^2 \, \mathbf{A}_{,tt} = -\operatorname{skew} \operatorname{Curl} \boldsymbol{m} + \operatorname{skew} \boldsymbol{\sigma} \,,$$

which, expressed in terms of u and \mathbf{A} gives

$$\rho u_{,tt} = \operatorname{Div}\left[2\mu_{e}\operatorname{sym}\operatorname{D}u + 2\mu_{c}\operatorname{skew}\left(\operatorname{D}u - \mathbf{A}\right) + \lambda_{e}\operatorname{tr}\left(\operatorname{D}u\right)\mathbb{1}\right], \qquad (4.6)$$

$$\rho\eta \tau_{c}^{2} \mathbf{A}_{,tt} = -L_{c}^{2} \mu_{e}\operatorname{skew}\operatorname{Curl}\left[a_{1}\operatorname{dev}\operatorname{sym}\operatorname{Curl}\mathbf{A} + a_{2}\operatorname{skew}\operatorname{Curl}\mathbf{A} + \frac{a_{3}}{3}\operatorname{tr}\left(\operatorname{Curl}\mathbf{A}\right)\mathbb{1}\right] + 2\mu_{c}\operatorname{skew}\left(\operatorname{D}u - \mathbf{A}\right).$$

We recall the Curl-Daxl identities, (see [52], Nye's formula [59])

 $\rho\eta$

$$\boldsymbol{\alpha} = -\operatorname{Curl} \mathbf{A} = (\operatorname{D} \operatorname{axl} \mathbf{A})^T - \operatorname{tr} \left[(\operatorname{D} \operatorname{axl} \mathbf{A})^T \right] \cdot \mathbb{1} = \boldsymbol{\mathfrak{K}}^T - \operatorname{tr} \left(\boldsymbol{\mathfrak{K}}^T \right) \mathbb{1},$$
(4.7)
$$\boldsymbol{\mathfrak{K}} = \operatorname{D} \operatorname{axl} \mathbf{A} = -(\operatorname{Curl} \mathbf{A})^T + \frac{1}{2} \operatorname{tr} \left[(\operatorname{Curl} \mathbf{A})^T \right] \cdot \mathbb{1} = \boldsymbol{\alpha}^T - \frac{1}{2} \operatorname{tr} \left(\boldsymbol{\alpha}^T \right) \mathbb{1},$$

as well as its implications

$$sym \boldsymbol{\alpha} = sym \boldsymbol{\mathfrak{K}} - tr(\boldsymbol{\mathfrak{K}}) \mathbb{1}, \qquad dev sym \boldsymbol{\alpha} = dev sym \boldsymbol{\mathfrak{K}}, \qquad (4.8)$$
$$skew \boldsymbol{\alpha} = -skew \boldsymbol{\mathfrak{K}}, \qquad tr(\boldsymbol{\alpha}) = -2 tr(\boldsymbol{\mathfrak{K}}).$$

Due to this converting formulae, the action functional admits the equivalent form

$$\mathcal{A} = \int_0^T \int_\Omega \left(\frac{1}{2} \rho \| u_{,t} \|^2 + \rho \eta \tau_{\rm c}^2 \| (\text{axl } \mathbf{A})_{,t} \|^2 - W \right) dv \, dt,$$

where the internal energy density is expressed in the alternative form

$$W = \mu_{e} \|\text{sym} (\mathrm{D}u - \mathbf{A})\|^{2} + \mu_{c} \|\text{skew} (\mathrm{D}u - \mathbf{A})\|^{2} + \frac{\lambda_{e}}{2} [\text{tr}(\mathrm{D}u - \mathbf{A})]^{2} + \frac{\mu_{e} L_{c}^{2}}{2} \left[a_{1} \|\text{dev} \, \text{sym} \, \mathrm{D} \, \text{axl} \, \mathbf{A}\|^{2} + a_{2} \|\text{skew} \, \mathrm{D} \, \text{axl} \, \mathbf{A}\|^{2} + \frac{4 \, a_{3}}{3} [\text{tr} (\mathrm{D} \, \text{axl} \, \mathbf{A})]^{2} \right].$$
(4.9)

We also notice that

$$\|\mathbf{\mathfrak{K}}\|^{2} = \|\operatorname{dev}\operatorname{sym}\mathbf{\mathfrak{K}}\|^{2} + \|\operatorname{skew}\mathbf{\mathfrak{K}}\|^{2} + \frac{1}{3}[\operatorname{tr}\mathbf{\mathfrak{K}}]^{2} = \|\operatorname{dev}\operatorname{sym}\boldsymbol{\alpha}\|^{2} + \|\operatorname{skew}\boldsymbol{\alpha}\|^{2} + \frac{1}{12}[\operatorname{tr}(\boldsymbol{\alpha})]^{2}.$$
(4.10)

Since $H^1(\Omega) \cap \mathfrak{so}(3)$ coincides with $H(\operatorname{Curl}; \Omega) \cap \mathfrak{so}(3)$, see [52], taking variations over all $\mathbf{A} \in H^1(\Omega) \cap \mathfrak{so}(3)$ is equivalent to considering variations over all $\mathbf{A} \in H(\operatorname{Curl}; \Omega) \cap \mathfrak{so}(3)$, i.e. the obtained Euler-Lagrange equations obtained using variations in $\mathbf{A} \in H^1(\Omega) \cap \mathfrak{so}(3)$ and $\mathbf{A} \in H(\operatorname{Curl}; \Omega) \cap \mathfrak{so}(3)$, respectively, are equivalent. Taking variations $H(\operatorname{Curl}; \Omega) \cap \mathfrak{so}(3)$, the minimum action principle leads to the following system of partial differential equations

$$\rho u_{,tt} = \operatorname{Div}\left[\underbrace{2\mu_{e} \operatorname{sym} \operatorname{D}u + 2\,\mu_{c} \operatorname{skew}(\operatorname{D}u - \mathbf{A}) + \lambda_{e} \operatorname{tr}(\operatorname{D}u)}_{\sigma} \mathbb{1}\right],$$

$$2\,\rho\,\eta\,\tau_{c}^{2} \ (\operatorname{axl}\,\mathbf{A})_{,tt} = \frac{\mu_{e}L_{c}^{2}}{2}\operatorname{Div}\left[2\,a_{1}\operatorname{dev}\operatorname{sym}\left(\operatorname{D}\operatorname{axl}\,\mathbf{A}\right) + 2\,a_{2}\operatorname{skew}(\operatorname{D}\operatorname{axl}\,\mathbf{A}) + \frac{8\,a_{3}}{3}\operatorname{tr}(\operatorname{D}\operatorname{axl}\,\mathbf{A}) \cdot \mathbb{1}\right] - 4\,\mu_{c}\operatorname{axl}\left(\operatorname{skew}\operatorname{D}u - \mathbf{A}\right),$$

$$(4.11)$$

The above systems is in complete agreement to the equations proposed by Eringen in the linear Cosserat theory. To see this explicitly, let us write this PDE-system in indices, giving

$$\rho u_{i,tt} = (\mu_{\rm e} + \mu_{\rm c})u_{i,jj} + (\mu_{\rm e} - \mu_{\rm c} + \lambda_{\rm e})u_{j,ij} + 2\,\mu_{\rm c}\,\epsilon_{ijk}\vartheta_{k,j}, \qquad (4.12)$$

$$2\,\rho\,\eta\,\mu_{\rm e}\,\tau_{\rm c}^2\,\vartheta_{i,tt} = \mu_{\rm e}\frac{L_{\rm c}^2}{2}\left[(a_1 + a_2)\,\vartheta_{i,jj} + \frac{1}{3}\,(a_1 - 3\,a_2 + 2\,a_3))\vartheta_{j,ij}\right] + 2\,\mu_{\rm c}\,\varepsilon_{ijk}u_{k,j} - 4\,\mu_{\rm c}\,\vartheta_i\,.$$

Other alternative expressions of the energy, in which the quadratic form in terms of \mathbf{e} has a similar structure as the quadratic form in terms of \mathfrak{K} , are

$$W = \underbrace{\mu_{e} \|\operatorname{dev}\operatorname{sym} \mathbf{e}\|^{2} + \mu_{c} \|\operatorname{skew} \mathbf{e}\|^{2} + \frac{2\mu_{e} + 3\lambda_{e}}{6} [\operatorname{tr}(\mathbf{e})]^{2}}_{:=W_{1}(\mathbf{e})} + \underbrace{\frac{\mu_{e}L_{c}^{2}}{2} \left[\alpha_{1} \|\operatorname{dev}\operatorname{sym} \mathfrak{K}\|^{2} + \alpha_{2} \|\operatorname{skew} \mathfrak{K}\|^{2} + \frac{2\alpha_{1} + 3\alpha_{3}}{6} [\operatorname{tr}(\mathfrak{K})]^{2} \right]}_{:=W_{2}(\mathfrak{K})}$$

$$= \mu_{e} \|\operatorname{sym} (\operatorname{D} u - \mathbf{A})\|^{2} + \mu_{c} \|\operatorname{skew} (\operatorname{D} u - \mathbf{A})\|^{2} + \frac{\lambda_{e}}{2} [\operatorname{tr}(\operatorname{D} u - \mathbf{A})]^{2} + \frac{\mu_{e}L_{c}^{2}}{2} \left[\alpha_{1} \|\operatorname{dev}\operatorname{sym}\operatorname{Curl} \mathbf{A}\|^{2} + \alpha_{2} \|\operatorname{skew}\operatorname{Curl} \mathbf{A}\| + \frac{2\alpha_{1} + 3\alpha_{3}}{24} [\operatorname{tr}(\operatorname{Curl} \mathbf{A})]^{2} \right],$$

$$(4.13)$$

where we have used the new weight parameters $\alpha_1, \alpha_2, \alpha_3$

$$a_1 = \alpha_1, \qquad a_2 = \alpha_2, \qquad a_3 = \frac{2\alpha_1 + 3\alpha_3}{8}$$
 (4.14)

equivalently

$$\alpha_1 = a_1, \qquad \alpha_2 = a_2, \qquad \alpha_3 = \frac{2}{3}(4a_3 - a_1).$$
 (4.15)

This new form yields³

$$\boldsymbol{\sigma} := \mathbf{D}_{\mathbf{e}} W = 2\mu_{\mathbf{e}} \operatorname{sym} \mathbf{e} + 2\,\mu_{\mathbf{c}} \operatorname{skew} \mathbf{e} + \lambda_{\mathbf{e}} \operatorname{tr}(\mathbf{e})\,\mathbb{1},$$
$$\mathbf{m} := \mathbf{D}_{\mathfrak{K}} W = \frac{\mu_{\mathbf{e}} L_{\mathbf{c}}^2}{2} \left[2\,\alpha_1 \operatorname{sym} \mathfrak{K} + 2\,\alpha_2 \operatorname{skew} \mathfrak{K} + \alpha_3 \operatorname{tr}(\mathfrak{K})\,\mathbb{1} \right]. \tag{4.17}$$

Using this expression of the internal energy density, some calculations of constitutive requirements in terms of $\mu_{\rm e}, \mu_{\rm c}, \lambda_{\rm e}$ can immediately be extrapolated to similar relations in terms of $\alpha_1, \alpha_2, \alpha_3$, as e.g., the Legendre-Hadamard ellipticity conditions imposed on $W_1(\mathbf{e})$ and $W_2(\mathfrak{K})$, respectively.

Anticipating the importance of having the same structure of these two energy's parts, let us first treat the generic case of a quadratic form which can then be applied to the balance of linear and angular momentum system. The generic quadratic form we consider is

$$\widetilde{W}(\mathbf{D}\phi) := b_1 \|\text{sym}\,\mathbf{D}\phi\|^2 + b_2 \|\text{skew}\,\mathbf{D}\phi\|^2 + b_3 [\operatorname{tr}(\mathbf{D}\phi)]^2.$$
(4.18)

 3 This form agrees with that considered by Ehlers [10, Eqs. (46) and (47)], where the internal energy is

$$W = \frac{\lambda}{2} \left[\operatorname{tr}(\varepsilon) \right]^2 + \frac{\mu + \mu_c}{2} \operatorname{tr}(\varepsilon \varepsilon) + \frac{\mu - \mu_c}{2} \operatorname{tr}\left(\varepsilon^T \varepsilon\right) + \frac{\alpha}{2} \left[\operatorname{tr}(\mathbf{\hat{\kappa}}) \right]^2 + \frac{\beta + \gamma}{2} \operatorname{tr}(\mathbf{\hat{\kappa}} \mathbf{\hat{\kappa}}) + \frac{\beta - \gamma}{2} \operatorname{tr}\left(\mathbf{\hat{\kappa}}^T \mathbf{\hat{\kappa}}\right).$$
(4.16)

Replacing $D\phi$ by the rank one dyadic product $\xi \otimes \eta$ we obtain

$$\begin{aligned} D^{2}W(D\phi).(\xi \otimes \eta, \xi \otimes \eta) &= b_{1} \|\operatorname{sym} \xi \otimes \eta\|^{2} + b_{2} \|\operatorname{skew} \xi \otimes \eta\|^{2} + b_{3} [\operatorname{tr}(\xi \otimes \eta)]^{2} \\ &= \frac{b_{1}}{4} \|\xi \otimes \eta + \eta \otimes \xi\|^{2} + \frac{b_{2}}{4} \|\xi \otimes \eta - \eta \otimes \xi\|^{2} + b_{3} \langle \xi, \eta \rangle^{2} \\ &= \frac{b_{1}}{4} \|\xi \otimes \eta\|^{2} + 2 \langle \xi \otimes \eta, \eta \otimes \xi \rangle + \frac{b_{2}}{4} (2\|\xi \otimes \eta\|^{2} - 2 \langle \xi \otimes \eta, \eta \otimes \xi \rangle) + b_{3} \langle \xi, \eta \rangle^{2} \\ &= \frac{b_{1}}{4} (\|\xi\|^{2} \|\eta\|^{2} + 2 \langle \xi, \eta \rangle^{2}) + \frac{b_{2}}{4} (\|\xi\|^{2} \|\eta\|^{2} - 2 \langle \xi, \eta \rangle^{2}) + b_{3} \langle \xi, \eta \rangle^{2} \\ &= \frac{b_{1} + b_{2}}{2} \|\xi\|^{2} \|\eta\|^{2} + \frac{b_{1} - b_{2} + 2a_{3}}{2} \langle \xi, \eta \rangle^{2} \\ &= \frac{b_{1} + b_{2}}{2} \|\xi\|^{2} \|\eta\|^{2} (\sin^{2}\theta + \cos^{2}\theta) + \frac{b_{1} - b_{2} + 2b_{3}}{2} \|\xi\|^{2} \|\eta\|^{2} \cos^{2}\theta \\ &= \frac{b_{1} + b_{2}}{2} \|\xi\|^{2} \|\eta\|^{2} \sin^{2}\theta + (\frac{b_{1} + b_{2}}{2} + \frac{b_{1} - b_{2} + 2b_{3}}{2} \|\xi\|^{2} \|\eta\|^{2} \cos^{2}\theta \\ &= \frac{b_{1} + b_{2}}{2} \|\xi\|^{2} \|\eta\|^{2} \sin^{2}\theta + (b_{1} + b_{3}) \|\xi\|^{2} \|\eta\|^{2} \cos^{2}\theta, \end{aligned}$$

where θ represents the angle between ξ and η .

~ .

Thus, the bilinear form of the second derivative of \widetilde{W} reads

$$D^{2}\overline{W}(D\phi).(\xi \otimes \eta, \xi \otimes \eta) = (b_{1} + b_{2})\|\xi\|^{2}\|\eta\|^{2}\sin^{2}\theta + (b_{1} + b_{3})\|\xi\|^{2}\|\eta\|^{2}\cos^{2}\theta,$$
(4.20)

and we infer the necessary and sufficient conditions for strict Legendre-Hadamard ellipticity of the quadratic form \widetilde{W} , i.e.,

$$\mathsf{D}^{2}\widetilde{W}(\mathsf{D}\phi).(\xi\otimes\eta,\xi\otimes\eta)>0\qquad\forall\ \eta,\xi\in\mathbb{R}^{3},\ \|\eta\|=\|\xi\|=1,$$
(4.21)

to be

$$b_1 + b_2 > 0, \qquad b_1 + b_3 > 0.$$
 (4.22)

Applying this result to both the strain energy $W_1(\mathbf{e})$ and the curvature energy $W_2(\mathbf{\hat{s}})$, see (4.13), we obtain the Legendre-Hadamard ellipticity conditions for both energy's part, see Subsection 6.3

$$\mu_{\rm e} + \mu_{\rm c} > 0, \qquad 2\mu_{\rm e} + \lambda_{\rm e} > 0, \qquad (4.23)$$

and formally similar

$$\alpha_1 + \alpha_2 > 0, \qquad 2\alpha_1 + \alpha_3 > 0, \qquad (4.24)$$

respectively.

Using the weight parameters $\alpha_1, \alpha_2, \alpha_3$, the form of the system of partial differential equations is

$$\rho \ u_{i,tt} = (\mu_{\rm e} + \mu_{\rm c}) u_{i,jj} + (\mu_{\rm e} - \mu_{\rm c} + \lambda_{\rm e}) u_{j,ij} + 2 \,\mu_{\rm c} \,\epsilon_{ijk} \,\vartheta_{k,j}, \tag{4.25}$$

$$2 \,\rho \,\eta \,\mu_{\rm e} \,\tau_{\rm c}^2 \,\vartheta_{i,tt} = \frac{\mu_{\rm e} L_{\rm c}^2}{2} \,\left[(\alpha_1 + \alpha_2) \vartheta_{i,jj} + (\alpha_1 - \alpha_2 + \alpha_3) \vartheta_{j,ij} \right] + 2 \,\mu_{\rm c} \,\varepsilon_{ijk} \,u_{k,j} - 4 \,\mu_{\rm c} \,\vartheta_i \,.$$

5 Identification of parameters in the dislocation formulation

We immediately remark that $\mathbf{e} = \mathbf{e}^{*T}$ while $\boldsymbol{\alpha}$ is not directly related to the curvature stress tensor $\boldsymbol{\Re}$ considered by Eringen. However, due to the Curl-Daxl converting formulae (4.7), the internal energy density proposed

here admits the alternative expression

$$W = \mu_{e} \|\operatorname{sym} (\mathrm{D}u - \mathbf{A})\|^{2} + \mu_{c} \|\operatorname{skew} (\mathrm{D}u - \mathbf{A})\|^{2} + \frac{\lambda_{e}}{2} [\operatorname{tr} (\mathrm{D}u - \mathbf{A})]^{2} + \frac{\mu_{e} L_{c}^{2}}{2} \left[a_{1} \|\operatorname{dev} \operatorname{sym} \operatorname{Curl} \mathbf{A}\|^{2} + a_{2} \|\operatorname{skew} \operatorname{Curl} \mathbf{A}\| + \frac{a_{3}}{3} \operatorname{tr} (\operatorname{Curl} \mathbf{A})^{2} \right] = \mu_{e} \|\operatorname{dev} \operatorname{sym} \mathbf{e}^{*}\|^{2} + \mu_{c} \|\operatorname{skew} \mathbf{e}^{*}\|^{2} + \frac{2\mu_{e} + 3\lambda_{e}}{6} [\operatorname{tr} (\mathbf{e}^{*})]^{2} + \frac{\mu_{e} L_{c}^{2}}{2} \left[a_{1} \|\operatorname{dev} \operatorname{sym} \mathbf{\hat{\kappa}}\|^{2} + a_{2} \|\operatorname{skew} \mathbf{\hat{\kappa}}\|^{2} + \frac{4a_{3}}{3} [\operatorname{tr} (\mathbf{\hat{\kappa}})]^{2} \right].$$
(5.1)

Due to the orthogonal Cartan-decomposition of the Lie-algebra

$$\mathfrak{gl}(3) = \{\mathfrak{sl}(3) \cap \operatorname{Sym}(3)\} \oplus \mathfrak{so}(3) \oplus \mathbb{R} \cdot \mathbb{1}, \qquad \mathbf{X} = \operatorname{dev} \operatorname{sym} \mathbf{X} + \operatorname{skew} \mathbf{X} + \frac{1}{3} \operatorname{tr}(\mathbf{X}) \cdot \mathbb{1}$$
(5.2)

and the identity

$$\langle \mathbf{X}, \mathbf{Y} \rangle = \langle \operatorname{dev} \operatorname{sym} \mathbf{X}, \operatorname{dev} \operatorname{sym} \mathbf{Y} \rangle + \langle \operatorname{skew} \mathbf{X}, \operatorname{skew} \mathbf{Y} \rangle + \frac{1}{3} \operatorname{tr}(\mathbf{X}) \operatorname{tr}(\mathbf{Y})$$
 (5.3)

the expression of the internal energy density proposed by Eringen may be written as

$$W = \frac{1}{2} \Big[(\mu^* + \varkappa) \Big(\langle \operatorname{dev} \operatorname{sym} \mathbf{e}^*, \operatorname{dev} \operatorname{sym} \mathbf{e}^* \rangle + \langle \operatorname{skew} \mathbf{e}^*, \operatorname{skew} \mathbf{e}^* \rangle + \frac{1}{3} \operatorname{tr}(\mathbf{e}^*) \operatorname{tr}(\mathbf{e}^*) \Big) \\ + \mu^* \Big(\langle \operatorname{dev} \operatorname{sym} \mathbf{e}^*, \operatorname{dev} \operatorname{sym} \mathbf{e}^* \rangle - \langle \operatorname{skew} \mathbf{e}^*, \operatorname{skew} \mathbf{e}^* \rangle + \frac{1}{3} \operatorname{tr}(\mathbf{e}^*) \operatorname{tr}(\mathbf{e}^*) \Big) \Big) + \lambda_e \left(\operatorname{tr} \mathbf{e}^* \right)^2$$
(5.4)
$$+ \gamma \Big(\langle \operatorname{dev} \operatorname{sym} \mathbf{\hat{R}}, \operatorname{dev} \operatorname{sym} \mathbf{\hat{R}} \rangle + \langle \operatorname{skew} \mathbf{\hat{R}}, \operatorname{skew} \mathbf{\hat{R}} \rangle + \frac{1}{3} \operatorname{tr}(\mathbf{\hat{R}}) \operatorname{tr}(\mathbf{\hat{R}}) \Big) \\ + \beta \Big(\langle \operatorname{dev} \operatorname{sym} \mathbf{\hat{R}}, \operatorname{dev} \operatorname{sym} \mathbf{\hat{R}} \rangle - \langle \operatorname{skew} \mathbf{\hat{R}}, \operatorname{skew} \mathbf{\hat{R}} \rangle + \frac{1}{3} \operatorname{tr}(\mathbf{\hat{R}}) \operatorname{tr}(\mathbf{\hat{R}}) \Big) \\ + \beta \Big(\langle \operatorname{dev} \operatorname{sym} \mathbf{\hat{R}}, \operatorname{dev} \operatorname{sym} \mathbf{\hat{R}} \rangle - \langle \operatorname{skew} \mathbf{\hat{R}}, \operatorname{skew} \mathbf{\hat{R}} \rangle + \frac{1}{3} \operatorname{tr}(\mathbf{\hat{R}}) \operatorname{tr}(\mathbf{\hat{R}}) \Big) \\ + \beta \Big(\langle \operatorname{dev} \operatorname{sym} \mathbf{\hat{R}}, \operatorname{dev} \operatorname{sym} \mathbf{\hat{R}} \rangle - \langle \operatorname{skew} \mathbf{\hat{R}}, \operatorname{skew} \mathbf{\hat{R}} \rangle + \frac{1}{3} \operatorname{tr}(\mathbf{\hat{R}}) \operatorname{tr}(\mathbf{\hat{R}}) \Big) \\ + \alpha \left(\operatorname{tr} \mathbf{\hat{R}} \right)^2 \Big] \\ = \frac{1}{2} \Big[(2\,\mu^* + \varkappa) \| \operatorname{dev} \operatorname{sym} \mathbf{e}^* \|^2 + \varkappa \| \operatorname{skew} \mathbf{e}^* \|^2 + \frac{2\,\mu^* + \varkappa + 3\,\lambda}{3} [\operatorname{tr}(\mathbf{e}^*)]^2 \\ + (\gamma + \beta) \| \operatorname{dev} \operatorname{sym} \mathbf{\hat{R}} \|^2 + (\gamma - \beta) \| \operatorname{skew} \mathbf{\hat{R}} \|^2 + \frac{\beta + \gamma + 3\,\alpha}{3} [\operatorname{tr}(\mathbf{\hat{R}})]^2 \Big].$$

Using again the orthogonal Cartan-decomposition of the Lie-algebra $\mathfrak{gl}(3)$ and since \mathbf{e}^* and \mathfrak{K} are independent constitutive variables, we obtain the relations between the parameters in the two forms

$$\mu_{\rm e} = \mu^* + \frac{\varkappa}{2}, \qquad \mu_{\rm c} = \frac{\varkappa}{2}, \qquad \frac{2\mu_{\rm e} + 3\lambda}{6} = \frac{2\,\mu^* + \varkappa + 3\,\lambda}{6}, \\ \mu_{\rm e} L_{\rm c}^2 \,a_1 = \gamma + \beta, \qquad \mu_{\rm e} L_{\rm c}^2 \,a_2 = \gamma - \beta, \qquad \mu_{\rm e} L_{\rm c}^2 \,\frac{4\,a_3}{3} = \frac{\beta + \gamma + 3\,\alpha}{3}, \qquad 2\,\eta\,\mu_{\rm e}\,\tau_{\rm c}^2 = j, \tag{5.5}$$

which implies

$$\lambda_{\rm e} = \lambda, \qquad \mu_{\rm e} = \mu^* + \frac{\varkappa}{2}, \qquad \mu_{\rm c} = \frac{\varkappa}{2}, a_1 = \frac{\gamma + \beta}{L_{\rm c}^2 \left(\mu^* + \frac{\varkappa}{2}\right)}, \qquad a_2 = \frac{\gamma - \beta}{L_{\rm c}^2 \left(\mu^* + \frac{\varkappa}{2}\right)}, \qquad a_3 = \frac{3\alpha + \beta + \gamma}{4 L_{\rm c}^2 \left(\mu^* + \frac{\varkappa}{2}\right)}, \qquad \eta = \frac{j}{2 \left(\mu^* + \frac{\varkappa}{2}\right) \tau_{\rm c}^2}, \tag{5.6}$$

or equivalently, expressed in the weight parameters $\alpha_1, \alpha_2, \alpha_3$

$$\lambda_{\rm e} = \lambda, \qquad \mu_{\rm e} = \mu^* + \frac{\varkappa}{2}, \qquad \mu_{\rm c} = \frac{\varkappa}{2}, \alpha_1 = \frac{\gamma + \beta}{L_{\rm c}^2 \left(\mu^* + \frac{\varkappa}{2}\right)}, \qquad \alpha_2 = \frac{\gamma - \beta}{L_{\rm c}^2 \left(\mu^* + \frac{\varkappa}{2}\right)}, \qquad \alpha_3 = \frac{2\,\alpha}{L_{\rm c}^2 \left(\mu^* + \frac{\varkappa}{2}\right)}, \qquad \eta = \frac{j}{2\left(\mu^* + \frac{\varkappa}{2}\right)\tau_{\rm c}^2}, \tag{5.7}$$

And in the other way

$$\lambda = \lambda_{\rm e}, \qquad \mu^* = \mu_{\rm e} - \mu_{\rm c}, \qquad \varkappa = 2\,\mu_{\rm c}, \gamma = \frac{a_1 + a_2}{2}\,L_{\rm c}^2\,\mu_{\rm e} = \frac{\alpha_1 + \alpha_2}{2}\,L_{\rm c}^2\,\mu_{\rm e}, \qquad \beta = \frac{a_1 - a_2}{2}\,L_{\rm c}^2\,\mu_{\rm e} = \frac{\alpha_1 - \alpha_2}{2}\,L_{\rm c}^2\,\mu_{\rm e}\,,$$
(5.8)
$$\alpha = \frac{1}{3}L_{\rm c}^2\,\mu_{\rm e}(4\,a_3 - a_1) = \frac{1}{2}\,L_{\rm c}^2\,\mu_{\rm e}\alpha_3\,, \qquad j = 2\,\eta\,\mu_{\rm e}\,\tau_{\rm c}^2\,.$$

6 Constitutive assumptions in the new parameter set

6.1 Positive definiteness of the elastic energy density and well-posedness of the solution

Using the following equivalent alternative forms of the internal energy density, namely

$$\begin{split} W &= \mu_{\rm e} \, \| \operatorname{dev} \operatorname{sym} \mathbf{e} \|^{2} + \mu_{\rm c} \, \| \operatorname{skew} \mathbf{e} \|^{2} + \frac{2\mu_{\rm e} + 3\,\lambda_{\rm e}}{6} \, [\operatorname{tr}\left(\mathbf{e}\right)]^{2} \\ &+ \frac{\mu_{\rm e}L_{\rm c}^{2}}{2} \left[a_{1} \| \operatorname{dev} \operatorname{sym} \operatorname{Curl} \mathbf{A} \|^{2} + a_{2} \| \operatorname{skew} \operatorname{Curl} \mathbf{A} \| + \frac{a_{3}}{3} \, \operatorname{tr}\left(\operatorname{Curl} \mathbf{A}\right)^{2} \right] \end{split}$$
(6.1)

$$&= \mu_{\rm e} \, \| \operatorname{dev} \operatorname{sym} \mathbf{e} \|^{2} + \mu_{\rm c} \, \| \operatorname{skew} \mathbf{e} \|^{2} + \frac{2\mu_{\rm e} + 3\,\lambda_{\rm e}}{6} \, [\operatorname{tr}\left(\mathbf{e}\right)]^{2} \\ &+ \frac{\mu_{\rm e}L_{\rm c}^{2}}{2} \left[\alpha_{1} \| \operatorname{dev} \operatorname{sym} \operatorname{Curl} \mathbf{A} \|^{2} + \alpha_{2} \| \operatorname{skew} \operatorname{Curl} \mathbf{A} \| + \frac{2\,\alpha_{1} + 3\,\alpha_{3}}{6} \, \operatorname{tr}\left(\operatorname{Curl} \mathbf{A}\right)^{2} \right] \\ &= \mu_{\rm e} \, \| \operatorname{dev} \operatorname{sym} \mathbf{e}^{*} \|^{2} + \mu_{\rm c} \, \| \operatorname{skew} \mathbf{e}^{*} \|^{2} + \frac{2\mu_{\rm e} + 3\,\lambda_{\rm e}}{6} \, [\operatorname{tr}\left(\mathbf{e}^{*}\right)]^{2} \\ &+ \frac{\mu_{\rm e}L_{\rm c}^{2}}{2} \left[a_{1} \, \| \operatorname{dev} \operatorname{sym} \mathfrak{K} \|^{2} + a_{2} \, \| \operatorname{skew} \mathfrak{K} \|^{2} + \frac{4\,a_{3}}{3} \, [\operatorname{tr}\left(\mathfrak{K}\right)]^{2} \right] \\ &= \mu_{\rm e} \, \| \operatorname{dev} \operatorname{sym} \mathbf{e} \|^{2} + \mu_{\rm c} \, \| \operatorname{skew} \mathbf{e} \|^{2} + \frac{2\mu_{\rm e} + 3\,\lambda_{\rm e}}{6} \, [\operatorname{tr}\left(\mathbf{e}\right)]^{2} \\ &+ \frac{\mu_{\rm e}L_{\rm c}^{2}}{2} \left[\alpha_{1} \, \| \operatorname{dev} \operatorname{sym} \mathfrak{K} \|^{2} + \alpha_{2} \, \| \operatorname{skew} \mathfrak{K} \|^{2} + \frac{2\,\alpha_{1} + 3\,\alpha_{3}}{6} \, [\operatorname{tr}\left(\mathfrak{K}\right)]^{2} \right] , \end{aligned}$$

and due to the orthogonal Cartan-decomposition of the Lie-algebra $\mathfrak{gl}(3)$, the strict positive definiteness as function of the strain tensors $\mathbf{e} = \mathbf{D}u - \mathbf{A}$ and $\boldsymbol{\alpha} = -\operatorname{Curl} \mathbf{A}$, as well as function of the strain tensors $\mathbf{e}^* = (\mathbf{D}u)^T + \mathbf{A}$ and $\mathfrak{K} = \mathbf{D}\vartheta = \mathbf{D}(\operatorname{axl} \mathbf{A})$, is equivalent to

$$\mu_{\rm e} > 0, \qquad \mu_{\rm c} > 0, \qquad 2\mu_{\rm e} + 3\,\lambda_{\rm e} > 0, \tag{6.3}$$
$$a_1 > 0, \qquad a_2 > 0, \qquad a_3 > 0.$$

Moreover, $(6.3)_2$ is equivalent to

$$\alpha_1 > 0, \qquad \alpha_2 > 0, \qquad 2\,\alpha_1 + 3\,\alpha_3 > 0.$$
(6.4)

Note that in the linear Cosserat model it is not possible to set the Cosserat couple modulus $\mu_c = 0$ since otherwise the constitutive coupling of the fields is lost (a coupling may remain due to some non-standard boundary conditions, but this is a pathological case [25]).

Under the assumption of positive definiteness, it was proven that in both the dynamic and the static case, and for both formulation, in terms of CurlA and D(axlA), the obtained mathematical model has a unique solution.

However, due to some new Korn-type inequality [20, 32, 33, 34], the existence and uniqueness of the solution is still valid for

$$\mu_{\rm e} > 0, \qquad \mu_{\rm c} > 0, \qquad 2\mu_{\rm e} + 3\,\lambda_{\rm e} > 0, \qquad \alpha_1 > 0, \qquad \alpha_2 \ge 0, \qquad 2\,\alpha_1 + 3\,\alpha_3 \ge 0\,, \quad (6.5)$$

or equivalently

$$\mu_{\rm e} > 0, \qquad \mu_{\rm c} > 0, \qquad 2\mu_{\rm e} + 3\lambda_{\rm e} > 0, \qquad a_1 > 0, \qquad a_2 \ge 0, \qquad a_3 \ge 0, \qquad (6.6)$$

i.e., under weaker assumptions on the constitutive parameters, in comparison to the positive definiteness conditions of the internal energy density, see the last two inequalities which allow α_2 and $2\alpha_1 + 3\alpha_3$ to vanish and the curvature energy may be chosen as (**the conformal curvature case**)

$$\frac{\mu_{e}L_{c}^{2}}{2}\alpha_{1}\|\operatorname{dev}\operatorname{sym}\operatorname{Curl}\mathbf{A}\|^{2} = \frac{\mu_{e}L_{c}^{2}}{2}a_{1}\|\operatorname{dev}\operatorname{sym}\operatorname{Curl}\mathbf{A}\|^{2}$$
$$= \frac{\mu_{e}L_{c}^{2}}{2}a_{1}\|\operatorname{dev}\operatorname{sym}\boldsymbol{\Re}\|^{2} = \frac{\mu_{e}L_{c}^{2}}{2}\alpha_{1}\|\operatorname{dev}\operatorname{sym}\boldsymbol{\Re}\|^{2} = \frac{\mu_{e}L_{c}^{2}}{2}\alpha_{1}\|\operatorname{dev}\operatorname{sym}\operatorname{D}\vartheta\|^{2}.$$
(6.7)

This choice is mandatory for stable identification of Cosserat parameters in the sense discussed in [50].

6.2 Real plane waves in isotropic Cosserat elastic solids

. .

We say that there exists *real plane waves* in the direction $\xi = (\xi_1, \xi_2, \xi_3)$, $\|\xi\|^2 = 1$, if for every wave number k > 0 the system of partial differential equations (4.12) admits a solution in the form:

$$u(x_1, x_2, x_3, t) = \underbrace{\begin{pmatrix} \widehat{u}_1 \\ \widehat{u}_2 \\ \widehat{u}_3 \end{pmatrix}}_{:= \widehat{u}} e^{i \left(k \langle \xi, x \rangle_{\mathbb{R}^3} - \omega t\right)}, \qquad (6.8)$$
$$\vartheta(x_1, x_2, x_3, t) = \underbrace{i \begin{pmatrix} \widehat{\vartheta}_1 \\ \widehat{\vartheta}_2 \\ \widehat{\vartheta}_3 \end{pmatrix}}_{:= \widehat{\vartheta}} e^{i \left(k \langle \xi, x \rangle_{\mathbb{R}^3} - \omega t\right)}, \quad \widehat{u}, \widehat{\vartheta} \in \mathbb{C}^3, \quad (\widehat{u}, \widehat{\vartheta})^T \neq 0,$$

only for real frequencies $\omega \in \mathbb{R}$, where $i = \sqrt{-1}$ is the complex unit. The plane wave is called "real" since it is defined by real values of ω . Note that we take $i \hat{\vartheta}$ since this choice will lead us in the end to only real valued matrices. Otherwise, we would have to deal with complex valued matrices in the linear Cosserat theory.

There exist real plane waves if for every wave number k > 0 the following systems of equations (4.12) admit non-trivial solutions:

$$\begin{bmatrix} \mathbf{Q}_1(e_1, k) - \omega^2 \hat{1} \end{bmatrix} w = 0 \qquad w = (\hat{u}_1, \hat{u}_2, \hat{\vartheta}_3)^T,$$

$$\begin{bmatrix} \mathbf{Q}_2(e_1, k) - \omega^2 \tilde{1} \end{bmatrix} w = 0 \qquad w = (\hat{u}_3, \hat{\vartheta}_1, \hat{\vartheta}_2)^T$$
(6.9)

only for real frequencies $\omega \in \mathbb{R}$, where

$$\mathbf{Q}_{1}(e_{1},k) = \begin{pmatrix} k^{2}(2\,\mu_{e}+\lambda_{e}) & 0 & 0 \\ 0 & k^{2}(\mu_{e}+\mu_{c}) & -2k\,\mu_{c} \\ 0 & -2\,k\,\mu_{c} & k^{2}\,\mu_{e}\,L_{c}^{2}\,(\alpha_{1}+\alpha_{2})+4\,\mu_{c} \end{pmatrix}, \\ \mathbf{Q}_{2}(e_{1},k) = \begin{pmatrix} k^{2}(\mu_{e}+\mu_{c}) & 0 & 2k\,\mu_{c} \\ 0 & k^{2}\,\mu_{e}\,L_{c}^{2}\,(2\,\alpha_{1}+\alpha_{3})+4\mu_{c} & 0 \\ 2\,k\,\mu_{c} & 0 & k^{2}\,\mu_{e}\,L_{c}^{2}\,(\alpha_{1}+\alpha_{2})+4\,\mu_{c} \end{pmatrix},$$

$$\widehat{1} = \begin{pmatrix} \rho & 0 & 0 \\ 0 & \rho & 0 \\ 0 & 0 & \rho \, j \,\mu_{e}\,\tau_{c}^{2} \end{pmatrix}, \qquad \widetilde{1} = \begin{pmatrix} \rho & 0 & 0 \\ 0 & \rho \, j \,\mu_{e}\,\tau_{c}^{2} & 0 \\ 0 & 0 & \rho \, j \,\mu_{e}\,\tau_{c}^{2} \end{pmatrix}.$$
(6.10)

Then, for k > 0 and due to the isotropy, extrapolating to all directions of propagation [51], we have

Proposition 6.1. The necessary and sufficient conditions for existence of real planar waves in any direction $\xi \in \mathbb{R}^3$, $\xi \neq 0$, in the framework of the linear isotropic elastic Cosservat theory [51, 22] are

$$2\mu_{\rm e} + \lambda_{\rm e} > 0, \qquad \mu_{\rm e} > 0, \qquad \mu_{\rm c} > 0, \qquad 2\alpha_1 + \alpha_3 > 0, \qquad \alpha_1 + \alpha_2 > 0, \tag{6.11}$$

equivalently

$$2\mu_{\rm e} + \lambda_{\rm e} > 0, \qquad \mu_{\rm e} > 0, \qquad \mu_{\rm c} > 0, \qquad a_1 + 2a_3 > 0, \qquad a_1 + a_2 > 0. \tag{6.12}$$

Name	Expression	Dispersive waves/ Non-dispersive waves	
the velocity of the acoustic branch of transla- tional compression (longitudinal) plane wave	$\mathfrak{c}_p = \sqrt{rac{\lambda_e + 2\mu_e}{ ho}}$	non-dispersive	
the limit of the group/phase velocity of the acoustic branch of the shear–rotational wave at $\omega \to 0 \ (k \to 0)$	$\mathfrak{c}_t = \sqrt{rac{\mu_e}{ ho}}$	dispersive	
the limit of the group/phase velocity of the acoustic branch of the shear–rotational wave at $\omega \to \infty \ (k \to \infty)$	$\mathfrak{c}_s = \sqrt{rac{\mu_e + \mu_c}{ ho}}$	dispersive	
the group/phase velocity for the compressional rotational wave in the limit $\omega \to \infty$ $(k \to \infty)$	$\mathfrak{c}_{m,p} = \sqrt{rac{L_c^2(2lpha_1+lpha_3)}{ hoj au_c^2}}$	dispersive	
the limit of the group/phase velocity of the acoustic branch of the shear–rotational wave at $\omega \to \infty \ (k \to \infty)$	$\mathfrak{c}_{m,s} = \sqrt{\frac{L_c^2(\alpha_1 + \alpha_2)}{\rho j \tau_c^2}} = \sqrt{\frac{L_c^2 \gamma}{\rho j \tau_c^2}}$	dispersive	
the limit of the optical branch (compressional- rotational and shear-rotational) at the cut-off frequency $\omega = 2\sqrt{\frac{\mu_c}{\rho j \mu_e \tau_c^2}}, \ k = 0$	0 (group velocity) $/\infty$ (phase velocity)	dispersive	

Table 1: Group velocities $c = \frac{\omega}{k}$ or/and phase velocities $\frac{d\omega}{dk}$ and the cut-off frequency in linear Cosserat elasticity, see [22].

Remark 6.2. Using the notation from Table 1, we have the following interpretation:

- i) the first implication of the set of conditions (6.11) means that all these waves (compressional/shear-rotational waves, acoustic/optical branch) are real; Once this aspect is clarified, we can treat and interpret further the propagation of plane waves;
- ii) under conditions (6.11) all branches of waves are real for the entire range $[0,\infty)$ of the frequency;
- iii) we can also see directly from the first condition that the translational compressional wave is real;
- iv) the second means that the acoustic branch of shear-rotational wave is real at low frequencies and together with the third means that the acoustic branch of shear-rotational wave is real at high frequencies;
- v) the fourth implies that the optical branch of the shear-rotational wave is real at high frequencies;
- vi) the third one means that the optical branch of the shear-rotational wave at high frequencies has a larger velocity than the acoustic branch of the same wave at low frequencies (if they both exist, which is the case due to other conditions);
- vii) the fifth one expresses directly that the compressional rotational wave at high frequencies is real.

6.3 Strong ellipticity (rank-one convexity, Legendre-Hadamard ellipticity) conditions

In linear isotropic classical elasticity, the necessary and sufficient conditions for existence of real planar waves in any direction $\xi \in \mathbb{R}^3$, $\xi \neq 0$ are

$$2\mu_{\rm e} + \lambda_{\rm e} > 0, \qquad \mu_{\rm e} > 0, \tag{6.13}$$

and they are equivalent to the strong ellipticity conditions (Legendre-Hadamard ellipticity).

For the Cosserat (micropolar) model the necessary and sufficient conditions for existence of a real planar wave are slightly different compared to the strong ellipticity conditions (Legendre–Hadamard ellipticity), the strong ellipticity conditions being connected to acceleration waves. In our notation, the strong ellipticity condition for Cosserat media is represented by the inequality [1, 11, 63]

$$\frac{d^2}{d\tau^2}W(\mathbf{e}+\tau\,\boldsymbol{\xi}\otimes\boldsymbol{\eta},\boldsymbol{\mathfrak{K}}+\tau\,\boldsymbol{\zeta}\otimes\boldsymbol{\eta})\bigg|_{\tau=0} > 0 \qquad \forall \,\boldsymbol{\eta},\boldsymbol{\xi},\boldsymbol{\zeta}\in\mathbb{R}^3, \ \|\boldsymbol{\eta}\|=\|\boldsymbol{\xi}\|=\|\boldsymbol{\zeta}\|=1.$$
(6.14)

The Legendre–Hadamard ellipticity condition [1, 11, 63] is equivalent to the positive definiteness of the *acoustic tensor*

$$\widehat{\mathbf{Q}} = \begin{pmatrix} \widehat{\mathbf{Q}}_1 & 0\\ 0 & \widehat{\mathbf{Q}}_2 \end{pmatrix} \tag{6.15}$$

whose block matrices are defined through

$$\widehat{\mathbf{Q}}_{1}(\xi) = \frac{1}{2} \begin{pmatrix} (2\mu_{e} + \lambda_{e})\xi_{1}^{2} + (\mu_{e} + \mu_{c})(\xi_{2}^{2} + \xi_{3}^{2}) & (\mu_{e} - \mu_{c} + \lambda_{e})\xi_{1}\xi_{2} & (\mu_{e} - \mu_{c} + \lambda_{e})\xi_{1}\xi_{3} \\ (\mu_{e} - \mu_{c} + \lambda_{e})\xi_{1}\xi_{2} & (2\mu_{e} + \lambda_{e})\xi_{2}^{2} + (\mu_{e} + \mu_{c})(\xi_{3}^{2} + \xi_{1}^{2}) & (\mu_{e} - \mu_{c} + \lambda_{e})\xi_{2}\xi_{3} \\ (\mu_{e} - \mu_{c} + \lambda_{e})\xi_{1}\xi_{3} & (\mu_{e} - \mu_{c} + \lambda_{e})\xi_{2}\xi_{3} & (2\mu_{e} + \lambda_{e})\xi_{3}^{2} + (\mu_{e} + \mu_{c})(\xi_{1}^{2} + \xi_{2}^{2}) \end{pmatrix}$$

and

$$\widehat{\mathbf{Q}}_{2}(\zeta) = \frac{1}{2} \begin{pmatrix} (2\alpha_{1} + \alpha_{3})\zeta_{1}^{2} + (\alpha_{1} + \alpha_{2})(\zeta_{1}^{2} + \zeta_{3}^{2}) & (\alpha_{1} - \alpha_{2} + \alpha_{3})\zeta_{1}\zeta_{2} & (\alpha_{1} - \alpha_{2} + \alpha_{3})\zeta_{1}\zeta_{3} \\ (\alpha_{1} - \alpha_{2} + \alpha_{3})\zeta_{1}\zeta_{2} & (2\alpha_{1} + \alpha_{3})\zeta_{2}^{2} + (\alpha_{1} + \alpha_{2})(\zeta_{3}^{2} + \zeta_{1}^{2}) & (\alpha_{1} - \alpha_{2} + \alpha_{3})\zeta_{2}\zeta_{3} \\ (\alpha_{1} - \alpha_{2} + \alpha_{3})\zeta_{1}\zeta_{3} & (\alpha_{1} - \alpha_{2} + \alpha_{3})\zeta_{2}\zeta_{3} & (2\alpha_{1} + \alpha_{3})\zeta_{3}^{2} + (\alpha_{1} + \alpha_{2})(\zeta_{1}^{2} + \zeta_{2}^{2}) \end{pmatrix}$$

for any nonzero wave directions $\xi \in \mathbb{R}^3$ and $\zeta \in \mathbb{R}^3$.

The positive definiteness of the acoustic tensor, i.e., the Legendre-Hadamard ellipticity conditions, is studied in [63, 1, 11] and, see (4.23) and (4.24), it is satisfied if and only if

$$2\mu_{\rm e} + \lambda_{\rm e} > 0, \qquad \mu_{\rm e} + \mu_{\rm c} > 0, \qquad 2\alpha_1 + \alpha_3 > 0, \qquad \alpha_1 + \alpha_2 > 0, \qquad (6.16)$$

equivalently

$$2\mu_{\rm e} + \lambda_{\rm e} > 0, \qquad \mu_{\rm e} + \mu_{\rm c} > 0, \qquad a_1 + 2a_3 > 0, \qquad a_1 + a_2 > 0.$$
 (6.17)

The absence of a coupling between **e** and \mathfrak{K} in the strain energy leads to a simplification of the calculations. The conditions (6.16) (strong ellipticity conditions, Legendre-Hadamard ellipticity, the positive definiteness of the acoustic tensor) imply the existence of real translational compressional waves in the entire range of real frequencies, of real shear rotational waves (both branches) at high frequencies, and of real rotational compressional wave at high frequencies, but at lower frequencies the latter waves may not be real since (6.16) does not guaranty that $\mathfrak{c}_t = \frac{\mu_e}{\rho}$ is real, since the positivity of μ_e is not necessarily implied by (6.16). To the contrary, the conditions (6.11) imply that all these branches and types of plane wave are real, i.e., the group/phase velocities are real on the entire range of possible frequencies.

The strong ellipticity conditions (6.16) are weaker than the conditions (6.11) in the sense that they are implied by the necessary and sufficient conditions for existence of a real planar wave (i.e., they imply the strong ellipticity and, therefore, the considered PDEs system is not unstable) but not vice versa. However, the strong ellipticity conditions (6.16) are not sufficient for some applications, and we believe that they are also not suitable for any approach regarding the propagation of Rayleigh waves in Cosserat solids [22]. Of course positive definiteness or the weaker conditions (6.5) of the elastic energy should be sufficient for any application.

7 Further identifications for micropolar constants

7.1 Eringen's initial notation: the notation used by Lakes

The first notations and tensors used by Eringen [13] and then in the Lakes group's experiments [25, 28] are

$$\widetilde{\sigma}_{ji} = (2\,\mu^* + \varkappa)(u_{i,j} + u_{j,i}) + \varkappa \,\epsilon_{jim} \left(\underbrace{\frac{1}{2}\epsilon_{mks}u_{s,k}}_{r_m} - \vartheta_m\right) + \lambda \,u_{k,k}\delta_{ji},\tag{7.1}$$
$$m_{ji} = \beta \,\vartheta_{j,i} + \gamma \,\vartheta_{i,j} + \alpha \,\vartheta_{k,k}\delta_{ji}.$$

The vector $r_m := \frac{1}{2} \epsilon_{mks} \frac{\partial u_s}{\partial x_k} = \frac{1}{2} (\operatorname{curl} u)_m$, m = 1, 2, 3, is the macrorotation vector of the linear theory of elasticity which is kinematically distinct from the microrotation vector. The internal energy density was chosen in the form

$$W = \frac{1}{2} \Big[(2\mu^* + \kappa) \varepsilon_{kl} \varepsilon_{lk} + \lambda \varepsilon_{kk} \varepsilon_{ll} + 2 \varkappa (r_k - \varphi_k) (r_k - \vartheta_k) + \beta \vartheta_{k,l} \vartheta_{l,k} + \gamma \vartheta_{l,k} \vartheta_{l,k} + \alpha \vartheta_{k,k} \vartheta_{l,l} \Big]$$
(7.2)
$$= \frac{2\mu^* + \kappa}{2} \|\varepsilon\|^2 + \frac{\lambda}{2} [\operatorname{tr}(\varepsilon)]^2 + \varkappa \|r - \vartheta\|^2 + \frac{\gamma}{2} \|\mathfrak{K}\|^2 + \frac{\beta}{2} \langle \mathfrak{K}, \mathfrak{K}^T \rangle + \frac{\alpha}{2} [\operatorname{tr}(\mathfrak{K})]^2,$$

where $\boldsymbol{\varepsilon} = \operatorname{sym} \operatorname{D} u$ is the classical strain tensor from the linear theory of classical elasticity and $\boldsymbol{\mathfrak{K}} = \operatorname{D} \vartheta = \operatorname{D}(\operatorname{axl} \mathbf{A})$.

In any case, the notations used by Eringen for the constitutive parameters are coherent and consistent, since we have the alternative form of the stress-strain relations

$$\sigma_{ji} = \mu^* (u_{i,j} + u_{j,i}) + \varkappa (u_{i,j} - \epsilon_{jis}\vartheta_s) + \lambda u_{k,k}\delta_{ji},$$

$$m_{ji} = \beta \vartheta_{j,i} + \gamma \vartheta_{i,j} + \alpha \vartheta_{k,k}\delta_{ji}$$
(7.3)

which are agree with the force stress and the couple stress tensor definition (7.19). Therefore, we have the same identifications as those already given.

The set of these coefficients was used by Lakes in expressing the experimental results in terms of the following coefficients, called *the micropolar technical constants* by Lakes [23, Page 2576], i.e.,

Young's modulus
$$E = \frac{(2\mu^* + \kappa)(3\lambda + 2\mu^* + \kappa)}{2\lambda + 2\mu^* + \kappa},$$
shear modulus $G = \frac{2\mu^* + \kappa}{2},$ Poisson ratio $\nu = \frac{\lambda}{2\lambda + 2\mu^* + \kappa},$ characteristic length, torsion $\ell_t = \sqrt{\frac{\beta + \gamma}{2\mu^* + \kappa}},$ characteristic length, bending $\ell_b = \sqrt{\frac{\gamma}{2(2\mu^* + \kappa)}},$ coupling number $N = \sqrt{\frac{\kappa}{2(\mu^* + \kappa)}},$ polar ratio $\Psi = \frac{\beta + \gamma}{\alpha + \beta + \gamma}.$

Using the micropolar technical constants, for the materials considered in Lakes' experiments, we will identify the numerical values of the constitutive parameters considered in our formulation (see Table 4).

7.2 Mindlin's notation

Mindlin [39] has considered the following strain tensors

$$\varepsilon_{ij} = \frac{1}{2}(u_{i,j} + u_{j,i}), \qquad \gamma_{[ij]} = u_{[j,i]} - P_{[ij]}, \qquad \kappa_{i[jk]} = P_{[jk],i},$$
(7.5)

where

$$u_{[i,j]} = \frac{1}{2}(u_{i,j} - u_{j,i}) = (\text{skew } Du)_{ij}, \qquad P_{[ij]} = \frac{1}{2}(P_{ij} - P_{ji}) = (\text{skew } P)_{ij}, \tag{7.6}$$

i.e., in our notation,

$$\boldsymbol{\varepsilon} = (\varepsilon_{ij})_{i,j=1,2,3} = \operatorname{sym} \operatorname{D} u, \qquad \boldsymbol{\gamma} = (\gamma_{[ij]})_{i,j=1,2,3} = \operatorname{skew}(\operatorname{D} u - \mathbf{P}), \qquad (7.7)$$
$$\boldsymbol{\kappa} = (\kappa_{i[jk]})_{i,j,k=1,2,3} = \operatorname{D} \operatorname{skew} \mathbf{P},$$

where $\mathbf{P} \in \mathbb{R}^{3 \times 3}$ is the microdistortion tensor arising in the micromorphic theory. Therefore, using our notation and identifying $\mathbf{A} = \text{skew } \mathbf{P}$, we obtain that the strain tensors considered by Mindlin are actually

$$\boldsymbol{\varepsilon} = \operatorname{sym}(\operatorname{D} u - \mathbf{A}) = \operatorname{sym} \mathbf{e}, \qquad \boldsymbol{\gamma} = \operatorname{skew}(\operatorname{D} u - \mathbf{A}) = \operatorname{skew} \mathbf{e}, \qquad \boldsymbol{\kappa} = \operatorname{D} \mathbf{A}.$$
(7.8)

Note that here κ is a third order tensor. The isotropic internal energy density in the form considered by Mindlin reads

$$W = \mu^{M} \varepsilon_{ij} \varepsilon_{ij} + \frac{1}{2} \lambda^{M} \varepsilon_{ii} \varepsilon_{jj} + \mu_{c}^{M} \gamma_{[ij]} \gamma_{[ij]} + \beta_{1}^{M} \kappa_{i[ik]} \kappa_{j[jk]} + \beta_{2}^{M} \kappa_{i[jk]} \kappa_{i[jk]} + \beta_{3}^{M} \kappa_{i[jk]} \kappa_{j[ik]}$$

$$= \mu^{M} \|\varepsilon\|^{2} + \frac{\lambda^{M}}{2} (\operatorname{tr}(\varepsilon))^{2} + \mu_{c}^{M} \|\gamma\|^{2} + \beta_{1}^{M} \kappa_{i[ik]} \kappa_{j[jk]} + \beta_{2} \kappa_{i[jk]} \kappa_{i[jk]} + \beta_{3}^{M} \kappa_{i[jk]} \kappa_{j[ik]}$$

$$= \mu^{M} \|\operatorname{sym}(\operatorname{D} u - \mathbf{A})\|^{2} + \frac{\lambda^{M}}{2} (\operatorname{tr}(\operatorname{sym}(\operatorname{D} u - \mathbf{A})))^{2} + \mu_{c}^{M} \|\operatorname{skew}(\operatorname{D} u - \mathbf{A})\|^{2} + \beta_{1}^{M} \kappa_{i[ik]} \kappa_{j[jk]} + \beta_{2}^{M} \kappa_{i[jk]} \kappa_{j[ik]}.$$

$$(7.9)$$

The assumed isotropic format of the curvature energy is not immediately apparent, since Mindlin deals with a third order tensor.

The equilibrium equations in the Mindlin form are

$$0 = (\lambda^{M} + \mu^{M} - \mu_{c}^{M})u_{j,ji} + (\mu^{M} + \mu_{c}^{M})u_{i,jj} - 2\mu_{c}^{M}P_{[ji],j},$$

$$0 = (\beta_{1}^{M} + \beta_{3}^{M})(P_{[ki],kj} + P_{[jk],ki}) + 2\beta_{2}^{M}P_{[ij],kk} - 2\mu_{c}^{M}P_{[ij]} + \mu_{c}^{M}(u_{j,i} - u_{i,j}).$$
(7.10)

We note

$$\|\boldsymbol{\kappa}\|^{2} = \|\mathbf{D}\mathbf{A}\|^{2} = 2 \|\mathbf{D}\mathrm{axl}(\mathbf{A})\|^{2} = 2 \|\boldsymbol{\mathfrak{K}}\|^{2} = 2 \|\mathbf{D}\vartheta\|^{2}$$
(7.11)

and that

$$\kappa_{i[jk]} = \mathbf{A}_{[jk],i} = \frac{1}{2} \left(A_{jk,i} - A_{kj,i} \right) = \frac{1}{2} \left(-\epsilon_{jkl} \vartheta_{l,i} + \epsilon_{kjl} \vartheta_{l,i} \right) = \frac{1}{2} \left(\epsilon_{kjl} + \epsilon_{kjl} \right) \vartheta_{l,i} = \epsilon_{kjl} \vartheta_{l,i}.$$
(7.12)

Therefore, on one hand we deduce

$$\kappa_{i[ik]} = \epsilon_{kil} \vartheta_{l,i} = (\operatorname{curl} \vartheta)_k,$$

$$\kappa_{i[ik]} \kappa_{j[jk]} = \|\operatorname{curl} \vartheta\|^2 = 2 \|\operatorname{skew} \mathcal{D}\vartheta\|^2 = 2 \|\operatorname{skew} \mathfrak{K}\|^2.$$
(7.13)

On the other hand we obtain

$$\kappa_{i[jk]}\kappa_{i[jk]} = \epsilon_{kjl}\vartheta_{l,i}\epsilon_{kjm}\vartheta_{m,i} = \epsilon_{kjl}\epsilon_{kjm}\vartheta_{l,i}\vartheta_{m,i} = (\delta_{jj}\delta_{lm} - \delta_{jm}\delta_{jl})\vartheta_{l,i}\vartheta_{m,i}$$

$$= (3\,\delta_{lm} - \delta_{lm})\vartheta_{l,i}\vartheta_{m,i} = 2\,\delta_{lm}\vartheta_{l,i}\vartheta_{m,i} = 2\,\vartheta_{l,i}\vartheta_{l,i} = 2\|\mathbf{D}\vartheta\|^2 = 2\|\mathbf{R}\|^2$$

$$= 2\left(\|\text{sym}\,\mathbf{R}\|^2 + \|\text{skew}\,\mathbf{R}\|^2 + \frac{1}{3}[\text{tr}(\mathbf{R})]^2\right)$$
(7.14)

and

$$\kappa_{i[jk]}\kappa_{j[ik]} = \epsilon_{kjl}\vartheta_{l,i}\epsilon_{kim}\vartheta_{m,j} = \epsilon_{kjl}\epsilon_{kim}\vartheta_{l,j}\vartheta_{m,i}$$

$$= (\delta_{ji}\delta_{lm} - \delta_{jm}\delta_{li})\vartheta_{l,i}\vartheta_{m,j} = \vartheta_{m,i}\vartheta_{m,i} - \vartheta_{i,i}\vartheta_{j,j} = \|\mathbf{D}\vartheta\|^2 - [\mathrm{tr}(\mathbf{D}\vartheta)]^2$$

$$= \|\mathrm{sym}\,\mathbf{\mathfrak{K}}\|^2 + \|\mathrm{skew}\,\mathbf{\mathfrak{K}}\|^2 + \frac{1}{3}[\mathrm{tr}(\mathbf{\mathfrak{K}})]^2 - [\mathrm{tr}(\mathbf{D}\vartheta)]^2 \qquad (7.15)$$

$$= \|\mathrm{sym}\,\mathbf{\mathfrak{K}}\|^2 + \|\mathrm{skew}\,\mathbf{\mathfrak{K}}\|^2 - \frac{2}{3}[\mathrm{tr}(\mathbf{\mathfrak{K}})]^2.$$

Thus, the internal energy density considered by Mindlin can be rewritten in the form

$$\begin{split} W &= \mu^{M} \|\text{sym}(\mathrm{D}u - \mathbf{A})\|^{2} + \frac{\lambda^{M}}{2} \left(\text{tr}(\text{sym}(\mathrm{D}u - \mathbf{A})) \right)^{2} + \mu_{c} \|\text{skew}(\mathrm{D}u - \mathbf{A})\|^{2} \\ &+ 2\beta_{1}^{M} \|\text{skew}\,\mathbf{\mathfrak{K}}\|^{2} + 2\beta_{2}^{M} \left(\|\text{sym}\,\mathbf{\mathfrak{K}}\|^{2} + \|\text{skew}\,\mathbf{\mathfrak{K}}\|^{2} + \frac{1}{3}[\text{tr}(\mathbf{\mathfrak{K}})]^{2} \right) \\ &+ \beta_{3}^{M} \left(\|\text{sym}\,\mathbf{\mathfrak{K}}\|^{2} + \|\text{skew}\,\mathbf{\mathfrak{K}}\|^{2} - \frac{2}{3}[\text{tr}(\mathbf{\mathfrak{K}})]^{2} \right) \\ &= \mu^{M} \|\text{dev}\,\text{sym}\,\mathbf{e}\|^{2} + \mu_{c}^{M} \|\text{skew}\,\mathbf{e}\|^{2} + \frac{2\mu^{M} + 3\lambda^{M}}{6} [\text{tr}(\mathbf{e})]^{2} \\ &+ (2\beta_{2}^{M} + \beta_{3}^{M})\|\text{sym}\,\mathbf{\mathfrak{K}}\|^{2} + (2\beta_{1}^{M} + 2\beta_{2}^{M} + \beta_{3}^{M}) \|\text{skew}\,\mathbf{\mathfrak{K}}\|^{2} + \frac{2\beta_{2}^{M} - 2\beta_{3}^{M}}{3} [\text{tr}(\mathbf{\mathfrak{K}})]^{2} \\ &= \mu^{M} \|\text{dev}\,\text{sym}\,\mathbf{e}\|^{2} + \mu_{c}^{M} \|\text{skew}\,\mathbf{e}\|^{2} + \frac{2\mu^{M} + 3\lambda^{M}}{6} [\text{tr}(\mathbf{e})]^{2} \\ &+ (2\beta_{2}^{M} + \beta_{3}^{M})\|\text{dev}\,\text{sym}\,\mathbf{\mathfrak{K}}\|^{2} + (2\beta_{1}^{M} + 2\beta_{2}^{M} + \beta_{3}^{M}) \|\text{skew}\,\mathbf{\mathfrak{K}}\|^{2} + \frac{4\beta_{2}^{M} - \beta_{3}^{M}}{3} [\text{tr}(\mathbf{\mathfrak{K}})]^{2}. \end{split}$$
(7.16)

By comparing the Mindlin's internal energy density to the form (4.13) considered by us in the new formulation, we obtain the identification of the parameters

$$\lambda_{\rm e} = \lambda^{\rm M}, \qquad \mu_{\rm e} = \mu^{\rm M}, \qquad \mu_{\rm c} = \mu_{\rm c}^{\rm M}, \qquad \mu_{\rm c} = \mu_{\rm c}^{\rm M}, \qquad \alpha_{\rm 1} = \frac{2}{L_{\rm c}^2 \mu^{\rm M}} (2 \beta_2^{\rm M} + \beta_3^{\rm M}), \qquad \alpha_{\rm 2} = \frac{2}{L_{\rm c}^2 \mu^{\rm M}} (2 \beta_1^{\rm M} + 2 \beta_2^{\rm M} + \beta_3^{\rm M}), \qquad \alpha_{\rm 3} = \frac{4}{3 L_{\rm c}^2 \mu^{\rm M}} (2 \beta_2^{\rm M} - 3 \beta_3^{\rm M}) \qquad (7.17)$$

and, vice versa,

$$\lambda^{M} = \lambda_{e}, \qquad \mu^{M} = \mu_{e}, \qquad \mu_{c}^{M} = \mu_{c}, \qquad \beta_{1}^{M} = \frac{L_{c}^{2} \mu_{e}}{4} (\alpha_{2} - \alpha_{1}), \qquad \beta_{2}^{M} = \frac{3 L_{c}^{2} \mu_{e}}{4} (2 \alpha_{1} + \alpha_{3}), \qquad \beta_{3}^{M} = \frac{L_{c}^{2} \mu_{e}}{8} (\alpha_{1} - \frac{3}{2} \alpha_{3}). \qquad (7.18)$$

7.3Nowacki's notation

In [18] the Nowacki notation [58] is used and another set of parameters is presented. The constitutive relations are given in indices as^4

$$\sigma_{ji} = (\mu^{\mathrm{N}} + \varkappa^{\mathrm{N}}) (u_{i,j} - \epsilon_{jim} \vartheta_m) + (\mu^{\mathrm{N}} - \varkappa^{\mathrm{N}}) (u_{j,i} - \epsilon_{ijm} \vartheta_m) + \lambda^{\mathrm{N}} u_{k,k} \delta_{ji},$$

$$m_{ji}^* = (\gamma^{\mathrm{N}} + \beta^{\mathrm{N}}) \vartheta_{i,j} + (\gamma^{\mathrm{N}} - \beta^{\mathrm{N}}) \vartheta_{j,i} + \alpha^{\mathrm{N}} \vartheta_{k,k} \delta_{ji},$$
(7.19)

This implies the following identifications

$$\mu^* + \varkappa = \mu^{N} + \varkappa^{N}, \qquad \mu^* = \mu^{N} - \varkappa^{N}, \qquad \lambda = \lambda^{N}, \qquad (7.20)$$
$$\beta = \gamma^{N} - \beta^{N}, \qquad \gamma = \gamma^{N} + \beta^{N}, \qquad \alpha = \alpha^{N},$$

and

$$\mu^{N} = \mu^{*} + \frac{\varkappa}{2}, \qquad \varkappa^{N} = \frac{\varkappa}{2}, \qquad \lambda^{N} = \lambda,$$

$$\gamma^{N} = \frac{\beta + \gamma}{2}, \qquad \beta^{N} = \frac{\gamma - \beta}{2}, \qquad \alpha^{N} = \alpha.$$
(7.21)

$$=rac{eta+\gamma}{2}, \qquad eta^{\mathrm{N}}=rac{\gamma-eta}{2}, \qquad lpha^{\mathrm{N}}=lpha\,.$$

Vice versa, it holds

$$\mu^* = \mu^{N} - \varkappa^{N}, \qquad \varkappa = 2\varkappa^{N}, \qquad \lambda = \lambda^{N}, \qquad (7.22)$$

$$\beta = \gamma^{N} - \beta^{N}, \qquad \gamma = \gamma^{N} + \beta^{N}, \qquad \alpha = \alpha^{N}.$$

⁴The tensors σ_{ji} and m_{ji}^* must be *transposed* before the parameters' identification with the ones proposed in this paper.

By using the identifications given in Section 5, we obtain the comparison between the constitutive parameters used by Nowacki and the parameters used in the dislocation format of the internal energy density, i.e.

$$\lambda_{\rm e} = \lambda^{\rm N}, \qquad \mu_{\rm e} = \mu^{\rm N}, \qquad \mu_{\rm c} = \varkappa^{\rm N}, \alpha_1 = \frac{2}{L_c^2 \,\mu^{\rm N}} \gamma^{\rm N}, \qquad \alpha_2 = \frac{2}{L_c^2 \,\mu^{\rm N}} \beta^{\rm N}, \qquad \alpha_3 = \frac{2}{L_c^2 \,\mu^{\rm N}} \alpha^{\rm N}.$$
(7.23)

8 Linear Cosserat elasticity as a particular case of the relaxed micromorphic model

The form of the internal energy proposed here keeps the form of the internal energy proposed in a more general theory, namely, the relaxed micromorphic theory. In the micromorphic theory, the microdistortion tensor $\mathbf{P} = (\mathbf{P}_{ij}) : \Omega \times [0,T] \to \mathbb{R}^{3\times 3}$ describes the substructure of the material which can rotate, stretch, shear and shrink, while $u = (u_i) : \Omega \times [0,T] \to \mathbb{R}^3$ is the displacement of the macroscopic material points. In the relaxed micromorphic model, in which the appearance of a strictly positive Cosserat modulus $\mu_c > 0$ is related to the isotropic Eringen-Claus model for dislocation dynamics [5, 15, 6], the free energy is given by

$$W_{\text{relax}} = \mu_{\text{e}} \|\text{sym} (\mathrm{D}u - \mathbf{P})\|^{2} + \mu_{\text{c}} \|\text{skew} (\mathrm{D}u - \mathbf{P})\|^{2} + \frac{\lambda_{\text{e}}}{2} [\text{tr}(\mathrm{D}u - \mathbf{P})]^{2} + \mu_{\text{micro}} \|\text{sym}\,\mathbf{P}\|^{2} + \frac{\lambda_{\text{micro}}}{2} [\text{tr}(\mathbf{P})]^{2} + \frac{\mu L_{\text{c}}^{2}}{2} \left[a_{1} \|\text{dev}\,\text{sym}\,\text{Curl}\,\mathbf{P}\|^{2} + a_{2} \|\text{skew}\,\text{Curl}\,\mathbf{P}\| + \frac{a_{3}}{3} \,\text{tr}(\text{Curl}\,\mathbf{P})^{2}\right],$$
(8.1)

where $(\mu_{\rm e}, \lambda_{\rm e})$, $(\mu_{\rm micro}, \lambda_{\rm micro})$, $\mu_{\rm c}$, $L_{\rm c}$ and (a_1, a_2, a_3) are the elastic moduli representing the parameters related to the meso-scale, the parameters related to the micro-scale, the Cosserat couple modulus $\mu_{\rm c} \geq 0$, the characteristic length $L_{\rm c}$, and the three general isotropic curvature parameters (non dimensional weights), respectively.

Due to the generalized Korn-type inequalities [56, 53, 55, 54, 3, 4], the relaxed micromorphic model is well-posed for either

$$\mu_{e} > 0, \qquad 3 \lambda_{e} + 2 \mu_{e} > 0, \qquad \mu_{c} \ge 0,$$

$$\mu_{\text{micro}} > 0, \qquad 3 \lambda_{\text{micro}} + 2 \mu_{\text{micro}} > 0, \qquad (8.2)$$

$$a_{1} > 0, \qquad a_{2} \ge 0, \qquad a_{3} \ge 0$$

or

$$\mu_{e} > 0, \qquad 3\lambda_{e} + 2\mu_{e} > 0, \qquad \mu_{c} \ge 0,$$

$$\mu_{\text{micro}} > 0, \qquad 3\lambda_{\text{micro}} + 2\mu_{\text{micro}} = 0,$$

$$a_{1} > 0, \qquad a_{2} > 0, \qquad a_{3} > 0.$$
(8.3)

Note that here, setting the Cosserat couple modulus $\mu_c = 0$ is possible since the two independent fields u and \mathbf{P} are still constitutively coupled via their strain tensors sym Du and sym P (this strain tensor coupling is missing in the Cosserat framework). The linear Cosserat model appears as a singular limit of the relaxed micromorphic model for

$$\mu_{\rm micro}, \qquad 3\lambda_{\rm micro} + 2\mu_{\rm micro} \to \infty.$$
 (8.4)

In this sense one can say that the relaxed micromorphic model uses a straightforward extension of the curvature energy in the Cosserat model (once the Cosserat model is represented in the appropriate second order dislocation tensor format). The Cosserat model is often supposed to describe a "rigid microstructure". This interpretation perfectly fits with the limit (8.4) constraining effectively the affine microdistortion P to infinitesimal rigid motions $\mathbf{P} = \mathbf{A} \in \mathfrak{so}(3)$ and giving back the Cosserat model.

8.1 Identification of the relaxed micromorphic parameters in Mindlin's notation

As a final identification, we report the correspondence between the coefficients in the relaxed micromorphic model and in the classical Mindlin's notation for linear elasticity with microstructure [41]. The isotropic Mindlin

formulation features 7+15 = 22 number of coefficients, while the isotropic relaxed micromorphic model displays altogether 5+3 = 8 coefficients. The relaxed micromorphic model is nevertheless a subclass of Mindlin's formulation. The identification for the coefficients of the linear part of the energy is (see [44, 40])

$$\begin{split} \mu^{\mathrm{M}} &= \mu_{\mathrm{micro}} \,, & \lambda^{\mathrm{M}} &= \lambda_{\mathrm{micro}} \,, & g_{1}^{\mathrm{M}} &= -\lambda_{\mathrm{micro}} \,, \\ b_{1}^{\mathrm{M}} &= \lambda_{\mathrm{e}} + \lambda_{\mathrm{micro}} \,, & b_{2}^{\mathrm{M}} &= \mu_{\mathrm{e}} + \mu_{\mathrm{micro}} + \mu_{\mathrm{c}} \,, & b_{3}^{\mathrm{M}} &= \mu_{\mathrm{e}} + \mu_{\mathrm{micro}} \,, \\ \end{split}$$

while for the curvature parameters we have (see [61, 40])

$$a_{1,2,3,5,8,11,14,15}^{\rm M} = 0, \qquad a_4^{\rm M} = \mu L_c^2 \frac{2a_3 - a_1}{3}, \qquad a_{10}^{\rm M} = \mu L_c^2 \frac{a_1 + a_2}{2}, \qquad a_{13}^{\rm M} = \mu L_c^2 \frac{a_1 - a_2}{2}.$$
(8.5)

9 Nonlinear isotropic Cosserat model

The nonlinear deformation of the body occupying the domain Ω is described by a map φ (called deformation) and by a *microrotation* orthogonal tensor field $\overline{\mathbf{R}}$,

$$\varphi: \Omega \subset \mathbb{R}^3 \to \mathbb{R}^3, \qquad \overline{\mathbf{R}}: \Omega \subset \mathbb{R}^3 \to \mathrm{SO}(3).$$
 (9.1)

We denote the current configuration by $\Omega := \varphi(\Omega) \subset \mathbb{R}^3$. The deformation and the microrotation is solution of the following *geometrically nonlinear minimization problem* posed on Ω :

$$I(\varphi, \overline{\mathbf{R}}) = \int_{\Omega} \left[W_{\rm mp}(\overline{\mathbf{U}}) + W_{\rm curv}(\boldsymbol{\alpha}) \right] dV \to \text{ min. w.r.t. } (\varphi, \overline{\mathbf{R}}), \qquad (9.2)$$

where

$$\begin{split} \mathbf{F} &:= \mathbf{D}\varphi \in \mathbb{R}^{3 \times 3} & \text{(the deformation gradient),} \\ \overline{\mathbf{U}} &:= \overline{\mathbf{R}}^T \mathbf{F} \in \mathbb{R}^{3 \times 3} & \text{(the non-symmetric Biot-type stretch tensor),} \\ \boldsymbol{\alpha} &:= \overline{\mathbf{R}}^T \operatorname{Curl} \overline{\mathbf{R}} \in \mathbb{R}^{3 \times 3} & \text{(the second order dislocation density tensor),} \\ \end{split}$$

$$W_{\rm mp}(\overline{\mathbf{U}}) := \mu_{\rm e} \left\| \operatorname{sym}(\overline{\mathbf{U}} - \mathbb{1}_3) \right\|^2 + \mu_{\rm c} \left\| \operatorname{skew}(\overline{\mathbf{U}} - \mathbb{1}_3) \right\|^2 + \frac{\lambda_{\rm e}}{2} \left[\operatorname{tr}(\operatorname{sym}(\overline{\mathbf{U}} - \mathbb{1}_3)) \right]^2 \quad \text{(physically linear)},$$

$$W_{\text{curv}}(\boldsymbol{\alpha}) := \mu_{\text{e}} \frac{L_{\text{c}}^{2}}{2} \left(a_{1} \| \text{dev} \operatorname{sym} \boldsymbol{\alpha} \|^{2} + a_{2} \| \text{skew} \, \boldsymbol{\alpha} \|^{2} + \frac{a_{3}}{3} [\operatorname{tr}(\boldsymbol{\alpha})]^{2} \right)$$
(curvature energy),

and dV denotes the volume element in the $\Omega\text{-configuration.}$

The total elastically stored energy $W = W_{\rm mp} + W_{\rm curv}$ depends on the deformation gradient $\mathbf{F} = \mathbf{D}\varphi$ and microrotations $\mathbf{\overline{R}}$ together with their spatial derivatives. In general, the *Biot-type stretch tensor* $\mathbf{\overline{U}}$ is not symmetric (the first Cosserat deformation tensor [7]). The parameters $\mu_{\rm e}$ and $\lambda_{\rm e}$ are the *Lamé constants* of classical isotropic elasticity, $\frac{2\mu_{\rm e}+3\lambda_{\rm e}}{3}$ is the *infinitesimal bulk modulus*, a_1, a_2, a_3 are *non-dimensional constitutive* curvature coefficients (weights), $\mu_{\rm c} \geq 0$ is called the Cosserat couple modulus and $L_{\rm c} > 0$ introduces an *internal* length which is characteristic for the material, e.g., related to the grain size in a polycrystal. The internal length $L_{\rm c} > 0$ is responsible for size effects in the sense that smaller samples are relatively stiffer than larger samples. The poplinger Cosserat model is length to be well paged [46, 45, 42, 21] for

The nonlinear Cosserat model is known to be well-posed [46, 45, 43, 31] for

$$\mu_{\rm e} > 0, \qquad 2\mu_{\rm e} + 3\lambda_{\rm e} > 0, \qquad \mu_{\rm c} > 0, \qquad a_1 > 0, \qquad a_2 > 0, \qquad a_3 > 0.$$
(9.4)

For the rotation tensor $\overline{\mathbf{R}} \in SO(3)$ there exists a unique skew-symmetric matrix

$$\mathbf{A} = \mathbf{Anti}(\vartheta_1, \vartheta_2, \vartheta_3) := \begin{pmatrix} 0 & -\vartheta_3 & \vartheta_2 \\ \vartheta_3 & 0 & -\vartheta_1 \\ -\vartheta_2 & \vartheta_1 & 0 \end{pmatrix} \in \mathfrak{so}(3),$$
(9.5)

such that

$$\overline{\mathbf{R}} = \exp \mathbf{A} = \sum_{k=0}^{\infty} \frac{1}{k!} \mathbf{A}^{k} = \mathbb{1}_{3} + \mathbf{A} + \text{h.o.t.}$$
(9.6)

Considering the linearisation, i.e. for situations of small deformations and small curvature,

$$\varphi = \mathrm{id} + u + \mathrm{h.o.t.}, \qquad \overline{\mathbf{R}} = \exp \mathbf{A} = \mathbb{1}_3 + \mathbf{A} + \mathrm{h.o.t.}$$
(9.7)

with $u: \Omega \to \mathbb{R}^3$ the infinitesimal displacement and the tensor field **A** the infinitesimal microrotation, the linear Cosserat model is obtained. Here, "h.o.t" stands for terms of order higher than linear with respect to u and **A**.

Note that, due to the results obtained in [52], the curvature tensor $\boldsymbol{\alpha} = \overline{\mathbf{R}}^T \operatorname{Curl} \overline{\mathbf{R}} \in \mathbb{R}^{3\times 3}$ controls all partial derivatives⁵ of $\overline{\mathbf{R}}$. It is important to realize that, contrary to the linear Cosserat model, the nonlinear Cosserat model allows to set the Cosserat couple modulus $\mu_c = 0$. In this case the curvature energy needs to be modified to allow a control of $\boldsymbol{\alpha} = \overline{\mathbf{R}}^T \operatorname{Curl} \overline{\mathbf{R}} \in \mathbb{R}^{3\times 3}$ in $L^p(\Omega), p > n$, i.e.

$$W_{\text{curv}}(\boldsymbol{\alpha}) := \mu_{\text{e}} \left[\frac{L_{\text{c}}^2}{2} \left(a_1 \| \text{dev sym}\,\boldsymbol{\alpha} \|^2 + a_2 \| \text{skew}\,\boldsymbol{\alpha} \|^2 + \frac{a_3}{3} \, [\text{tr}(\boldsymbol{\alpha})]^2 \right) \right]^{\frac{p}{2}}.$$
(9.9)

We also note that for $\mu_c > 0$ the quadratic nonlinear Cosserat model is **redundant** in the sense of Romano et al. [62]. Redundancy is connected to the way rigid body movements are penalised in the elastic energy. Certainly, zero elastic energy occurs if and only if the body undergoes a rigid body movement. In the case of the Cosserat model, however, a part of the elastic energy being zero already suffices to constrain the movement to a rigid motion. In this sense, **the typical Cosserat model is over-constraining rigid movements**. More precisely, for positive Cosserat couple modulus $\mu_c > 0$ it holds

$$W_{\rm mp}(\overline{\mathbf{U}}) = 0 \quad \Rightarrow \quad W_{\rm curv}(\boldsymbol{\alpha}) = 0$$

$$\tag{9.10}$$

and the strain measure already determines the curvature measure in zero. To see this, observe that for $\mu_c > 0$

$$W_{\rm mp}(\overline{\mathbf{U}}) = 0 \quad \Rightarrow \quad \overline{\mathbf{R}}^T \mathbf{F} - \mathbb{1} = 0 \quad \Rightarrow \quad \overline{\mathbf{R}} = \mathbf{F} = \mathbf{D}\varphi$$

$$\tag{9.11}$$

and by taking the Curl-operator on both sides we obtain

$$W_{\rm mp}(\overline{\mathbf{U}}) = 0 \quad \Rightarrow \quad \operatorname{Curl} \overline{\mathbf{R}} = 0.$$
 (9.12)

Since $\overline{\mathbf{R}} \in SO(3)$, this implies that $\overline{\mathbf{R}}$ is constant [52] and yields

$$W_{\rm mp}(\overline{\mathbf{U}}) = 0 \quad \Rightarrow \quad \boldsymbol{\alpha} = 0 \qquad (\boldsymbol{\mathfrak{K}} = 0).$$
 (9.13)

The same redundancy is true for the linear Cosserat model (in which $\mu_c > 0$ a priori). In this case we note that

$$\mathbf{e} = 0 \Rightarrow \mathbf{D}u = \mathbf{A} \Rightarrow 0 = \operatorname{Curl} \mathbf{A} = \alpha \Rightarrow \mathbf{A} = \operatorname{constant} \Rightarrow \operatorname{Daxl}(\mathbf{A}) = 0 \quad (\mathbf{\mathfrak{K}} = 0). \quad (9.14)$$

Note that the linear relaxed micromorphic model is non-redundant for $\mu_c = 0$, a choice which is permitted for well-posedness.

10 Conclusions

In the literature on linear isotropic Cosserat or micropolar solids many different abbreviations and definitions are frequently encountered. For convenience of the reader, especially for interpreting the experimental results [67, 23, 25, 24, 60, 2, 27, 28], based on the analysis presented in the present paper, we collect these technical constants in Table 2. We mention that further numerical values are proposed in [16, 64, 19].

The Cosserat theory of isotropic elastic solid written in the dislocation format (our proposal) represents a direct particular case of both the relaxed micromorphic model and of the geometrically nonlinear isotropic

$$\|\overline{\mathbf{R}}^{T} \mathrm{D}\overline{\mathbf{R}}\|^{2} = \|\mathrm{D}\overline{\mathbf{R}}\|^{2} = 2\left(\|\operatorname{dev}\operatorname{sym}\boldsymbol{\alpha}\|^{2} + \|\operatorname{skew}\boldsymbol{\alpha}\|^{2} + \frac{1}{12}\left[\operatorname{tr}(\boldsymbol{\alpha})\right]^{2}\right).$$
(9.8)

 $^{^5\}mathrm{For}$ example, we can express the isotropic term

Name	classical elasticity	Our notations	Lakes (Eringen)	Mindlin	Hassanpour & Heppler (Nowacki)
G (shear modulus)	μ	$\mu_{ m e}$	$\mu^* + \frac{\varkappa}{2}$	μ^{M}	$\mu^{ m N}$
Lamé first parameter	λ	$\lambda_{ m e}$	λ	$\lambda^{ m M}$	$\lambda^{\rm N}$
ν (classical Poisson ratio)	ν	$\frac{\lambda_{\rm e}}{2(\mu_{\rm e}+\lambda_{\rm e})}$	$\frac{\lambda}{2(\lambda + \mu^* + \frac{\varkappa}{2})}$	$\frac{\lambda^{\rm M}}{2(\mu^{\rm M}+\lambda^{\rm M})}$	$\frac{\lambda^{\rm N}}{2(\mu^{\rm N}+\lambda^{\rm N})}$
E (classical Young's modulus)	E	$\frac{\mu_{\rm e}(2\mu_{\rm e}+3\lambda_{\rm e})}{\mu_{\rm e}+\lambda_{\rm e}}$	$\frac{(2\mu^* + \varkappa)(3\lambda + 2\mu^* + \varkappa)}{2\lambda + 2\mu^* + \varkappa}$	$\frac{\mu^{\mathrm{M}}(2\mu^{\mathrm{M}}+3\lambda^{\mathrm{M}})}{\mu^{\mathrm{M}}+\lambda^{\mathrm{M}}}$	$\frac{\mu^{\mathrm{N}}(2\mu^{\mathrm{N}}+3\lambda^{\mathrm{N}})}{\mu^{\mathrm{N}}+\lambda^{\mathrm{N}}}$
κ (classical bulk modulus)	ĸ	$\lambda_{\rm e} + \frac{2}{3}\mu_{\rm e}$	$\lambda + \frac{2}{3} \left(\mu^* + \frac{\varkappa}{2} \right)$	$\lambda^{\rm M} + \frac{2}{3} \mu^{\rm M}$	$\lambda^{\rm N} + \frac{2}{3} \mu^{\rm N}$
mass density	ρ	ρ	ρ	ρ	ρ
Cosserat couple modulus	_	$\mu_{ m c}$	$\frac{\varkappa}{2}$	$\mu_{ m c}^{ m M}$	\mathcal{H}^{N}
first Cosserat twist coefficient	-	$\alpha_1 = a_1$	$\frac{\gamma + \beta}{L_{\rm c}^2 \left(\mu^* + \frac{\varkappa}{2}\right)}$	$\frac{2}{\mu^{\rm M} L_{\rm c}^2} (2\beta_2^{\rm M} + \beta_3^{\rm M})$	$\frac{2\gamma^{\rm N}}{\mu^{\rm N} L_{\rm c}^2}$
second Cosserat twist coefficient	_	$\alpha_2 = a_2$	$\frac{\gamma - \beta}{L_{\rm c}^2 \left(\mu^* + \frac{\varkappa}{2}\right)}$	$\frac{2}{\mu^{\rm M} L_{\rm c}^2} (2(\beta_1^{\rm M} + \beta_2^{\rm M}) + \beta_3^{\rm M})$	$\frac{2\beta^{\rm N}}{\mu^{\rm N} L_{\rm c}^2}$
third Cosserat twist coefficient	_	$\alpha_3 = \frac{2}{3} \left(4a_3 - a_1 \right)$	$\frac{2\alpha}{L_{\rm c}^2\left(\mu^* + \frac{\varkappa}{2}\right)}$	$\frac{2}{\mu^{\rm M} L_{\rm c}^2} \frac{4(\beta_2^{\rm M} - \beta_3^{\rm M})}{3}$	$\frac{2\alpha^{\rm N}}{\mu^{\rm N} L_{\rm c}^2}$
Ψ (dimensionless polar ratio)	-	$\frac{2\alpha_1}{\alpha_3 + 2\alpha_1} = \frac{3a_1}{2a_1 + 4a_3}$	$\frac{\beta + \gamma}{\alpha + \beta + \gamma}$	$\frac{3(2\beta_2^{\mathrm{M}}+\beta_3^{\mathrm{M}})}{10\beta_2^{\mathrm{M}}-\beta_3^{\mathrm{M}}}$	$\frac{2\gamma^{\rm N}}{\alpha^{\rm N}+2\gamma^{\rm N}}$
ℓ_t (characteristic length for torsion)	_	$L_{\rm c}\sqrt{\frac{\alpha_1}{2}} = L_{\rm c}\sqrt{\frac{a_1}{2}}$	$\sqrt{\frac{\beta+\gamma}{2\mu^*+\varkappa}}$	$\sqrt{\frac{2\beta_2^{\rm M}+\beta_3^{\rm M}}{\mu^{\rm M}}}$	$\sqrt{rac{\gamma^{ m N}}{\mu^{ m N}}}$
ℓ_b (characteristic length for bending)	-	$L_{\rm c}\sqrt{\frac{\alpha_1+\alpha_2}{8}} = L_{\rm c}\sqrt{\frac{a_1+a_2}{8}}$	$\sqrt{\frac{\gamma}{2\left(2\mu^*+\varkappa\right)}}$	$\sqrt{\frac{\beta_1^{\rm M}+2\beta_2^{\rm M}+\beta_3^{\rm M}}{2\mu^{\rm M}}}$	$\sqrt{\frac{\beta^{\rm N}+\gamma^{\rm N}}{4\mu^{\rm N}}}$
N (Cosserat coupling number)	-	$\sqrt{rac{\mu_{ m c}}{\mu_{ m e}+\mu_{ m c}}}$	$\sqrt{\frac{\varkappa}{2\left(\mu^*+\varkappa\right)}}$	$\sqrt{\frac{\mu_{\rm c}^{\rm M}}{\mu^{\rm M}+\mu_{\rm c}^{\rm M}}}$	$\sqrt{\frac{\varkappa^{\rm N}}{\mu^{\rm N}+\varkappa^{\rm N}}}$
ξ (micropolar twist Poisson's ration)	-	$\frac{\alpha_3}{2(\alpha_1 + \alpha_3)}$	$\frac{\alpha}{2\alpha+\beta+\gamma}$	$\frac{2(\beta_3^{\mathrm{M}}-\beta_2^{\mathrm{M}})}{\beta_3^{\mathrm{M}}-10\beta_2^{\mathrm{M}}}$	$\frac{\alpha^{\rm N}}{2(\gamma^{\rm N}+\alpha^{\rm N})}$
\mathcal{E} (micropolar tortile or torsional modulus)	_	$\frac{\mu_{\rm e} L_{\rm c}^2}{2} \frac{\alpha_1(2\alpha_1 + 3\alpha_3)}{\alpha_1 + \alpha_3}$	$\frac{(\beta+\gamma)(3\alpha+\beta+\gamma)}{2\alpha+\beta+\gamma}$	$6\left(\frac{\beta_{3}^{M^{2}}-8\beta_{2}^{M^{2}}-2\beta_{2}^{M}\beta_{3}^{M}}{\beta_{3}^{M}-10\beta_{2}^{M}}\right)$	$\frac{\gamma^{N}(2\gamma^{N}+3\alpha^{N})}{\gamma^{N}+\alpha^{N}}$
\mathcal{B} (micropolar tortile or torsional bulk modulus)	_	$\frac{\mu_{\rm e} L_{\rm c}^2}{2} \frac{2\alpha_1 + 3\alpha_3}{3}$	$\frac{(3\alpha+\beta+\gamma)}{3}$	$\frac{2}{3}(4\beta_2^{\mathrm{M}}-\beta_3^{\mathrm{M}})$	$\alpha^{\rm N} + \frac{2}{3}\gamma^{\rm N}$
microinertia den- sity	-	η	$\frac{j}{\tau_{\rm c}^2(\mu^*+\frac{\varkappa}{2})}$	$rac{j}{ au_{ m c}^2 \mu^{ m M}}$	$rac{j}{ au_{ m c}^2\mu^{ m N}}$

Table 2: A concluding table for converting all the needed quantities in terms of the new constitutive parameters. The terminology is that used in [18]. Note that $\Psi = \frac{3}{2}$ if and only if $a_3 = 0$.

Cosserat model which uses the second order dislocation density tensor $\boldsymbol{\alpha} = \overline{\mathbf{R}}^T \operatorname{Curl} \overline{\mathbf{R}} \in \mathbb{R}^{3 \times 3}$ as curvature measure.

We remark that $0 \le N \le 1$ and the limit case N = 1 ($\mu_c \to \infty$) corresponds to the case of the couple stress theory [17, 38, 42, 57]. The technical parameter Ψ (polar ratio) is positive if the necessary and sufficient conditions (6.4) for positive definiteness of the internal energy density are satisfied. However Ψ is not positive if the conditions (6.11) for existence of real planar waves in any direction are satisfied and neither when the

Material	E [MPa]	G [MPa]	ν[-]	N^2 [-]	$\ell_t \; [\mathrm{mm}]$	$\ell_b \; [\mathrm{mm}]$	Ψ [-]
Human bone	12000	4000	0.5	0.5	0.22	0.45	3/2
@0.2mm	12000	1000	0.0	0.0	0.22	0.40	0/2
Graphite @1.6mm	4500	2122.64	0.06	1	1.6	2.8	$(3/2)^*$
(H237)	4000	2122.04	0.00	1	1.0	2.0	(0/2)
Foam @1mm	1.28	0.6	0.07	0.09	3.8	5	3/2
(0.6 PS)	1.20	0.0	0.01	0.03	9.0	0	5/2
Foam @0.18mm	300	104	0.4	0.04	0.62	0.33	3/2
(dense polyurethane)							
Foam @0.15mm	2758	1033	0.34	0.1	0.065	0.0325	3/2
(dense syntactic)	2100	1000	0.04	0.1	0.000	0.0320	5/2

Table 3: In accordance with Lakes [27] (Table 1 at the end), Foam @1mm (0.6 PS) according to Lakes [23] (pp. 2576-2577), and Foam @0.15mm (dense syntactic) according to [24] (p. 60). For Foam @0.15mm (dense syntactic) we have used $\ell_b = 0.325$ [24], instead of $\ell_b = 0.33$ included in [27, Table 1], since otherwise the positive semi-definiteness is violated as well as the identified condition in bending experiment for this material [27, p. 60], i.e., $\beta/\gamma = 1$. * Note that for Graphite the Cosserat couple modulus $\mu_c \to \infty$, a constraint that reduces the Cosserat model to its particular case, i.e., the couple stress model. In this case, the moment part of the stress tensor is trace free, consistent with $\Psi = 3/2$ ($a_3 = 0$).

Material	$\mu_{\rm e}~[{\rm MPa}]$	$\lambda_{\rm e}$ [MPa]	$\mu_{\rm c}$ [MPa]	$\mu_{\mathbf{e}} L_{\mathbf{c}}^2 \alpha_1 [\mathbf{N}]$ $= \mu_{\mathbf{e}} L_{\mathbf{c}}^2 a_1 [\mathbf{N}]$	$ \begin{aligned} & \mu_{\mathbf{e}} L_{\mathbf{c}}^2 \alpha_2 \ [\mathrm{N}] \\ &= \boldsymbol{\mu}_{\mathbf{e}} L_{\mathbf{c}}^2 \boldsymbol{a}_2 \ [\mathrm{N}] \end{aligned} $	$\mu_{\rm e} L_{\rm c}^2 \alpha_3 \left[{\rm N} \right]$	$\mu_{\mathrm{e}}L_{\mathrm{c}}^2a_3~[\mathrm{N}]$
Human bone @0.2mm	4000	$\infty~(\nu=1/2)$	4000	387.2	6092.8	-258.133	0
Graphite @1.6mm (H237)	2122.64	289.451	∞	10867.9	122264	(-16301.85)*	(0) *
Foam @1mm (0.6 PS)	0.6	0.0923077	0.0593407	17.328	102.672	-11.552	0
Foam @0.18mm (dense polyurethane)	104	797.333	4.33333	79.9552	10.6496	-53.3035	0
Foam @0.15mm (dense syntactic)	1033	2096.29	114.778	8.72885	0	-5.81923	0

Table 4: The numerical experimental values of the new constitutive parameters according to Lakes [27]. *It is possible to evaluate α_3 since $\mu_c \to \infty$ implies that the moment part of the stress tensor is trace free, i.e., $\Psi = 3/2$. The last row corresponds to conformal curvature.

Legendre-Hadamard ellipticity condition (6.16) are satisfied, since both these constitutive conditions imply the positivity of $2 \alpha_1 + \alpha_3$ but they do not imply the positivity of α_1 .

Therefore, assuming the internal energy to be positive definite (the same remains true if $\alpha_2 \ge 0$) the following one-to-one relation between our constitutive parameters and the constitutive technical constants considered by Lakes hold

$$\lambda_{e} = -\frac{G(E - 2G)}{E - 3G}, \qquad \mu_{e} = G, \qquad \mu_{c} = \frac{GN^{2}}{1 - N^{2}},$$

$$\alpha_{1} = \frac{2\ell_{t}^{2}}{L_{c}^{2}}, \qquad \alpha_{2} = \frac{8\ell_{b}^{2} - 2\ell_{t}^{2}}{L_{c}^{2}}, \qquad \alpha_{3} = \frac{4\ell_{t}^{2}(1 - \psi)}{\psi L_{c}^{2}}, \qquad (10.15)$$

$$a_{1} = \frac{2\ell_{t}^{2}}{L_{c}^{2}}, \qquad a_{2} = \frac{8\ell_{b}^{2} - 2\ell_{t}^{2}}{L_{c}^{2}}, \qquad a_{3} = \frac{4\ell_{t}^{2}(3 - 2\psi)}{\psi L_{c}^{2}}.$$

In the conformal curvature case [20, 21]

$$\alpha_2 = 0 \Leftrightarrow a_2 = 0 \Leftrightarrow \ell_t = 2\,\ell_b \Leftrightarrow \beta = \gamma$$

and
$$2\alpha_1 + 3\alpha_3 = 0 \Leftrightarrow a_3 = 0 \Leftrightarrow \Psi = \frac{3}{2},$$
 (10.16)

the internal energy density is

$$W = \mu_{\rm e} \|\operatorname{dev} \operatorname{sym} \mathbf{e}\|^{2} + \mu_{\rm c} \|\operatorname{skew} \mathbf{e}\|^{2} + \frac{2\mu_{\rm e} + 3\,\lambda_{\rm e}}{6} [\operatorname{tr}(\mathbf{e})]^{2} + \frac{\mu_{\rm e}L_{\rm c}^{2}}{2} a_{1} \|\operatorname{dev} \operatorname{sym} \operatorname{Curl} \mathbf{A}\|^{2}$$

$$= \mu_{\rm e} \|\operatorname{dev} \operatorname{sym} \mathbf{e}\|^{2} + \mu_{\rm c} \|\operatorname{skew} \mathbf{e}\|^{2} + \frac{2\mu_{\rm e} + 3\,\lambda_{\rm e}}{6} [\operatorname{tr}(\mathbf{e})]^{2} + \frac{\mu_{\rm e}L_{\rm c}^{2}}{2} \alpha_{1} \|\operatorname{dev} \operatorname{sym} \operatorname{Curl} \mathbf{A}\|^{2}$$
(10.17)
$$= \mu_{\rm e} \|\operatorname{dev} \operatorname{sym} \mathbf{e}\|^{2} + \mu_{\rm c} \|\operatorname{skew} \mathbf{e}\|^{2} + \frac{2\mu_{\rm e} + 3\,\lambda_{\rm e}}{6} [\operatorname{tr}(\mathbf{e})]^{2} + \frac{\mu_{\rm e}L_{\rm c}^{2}}{2} \alpha_{1} \|\operatorname{dev} \operatorname{sym} \mathfrak{K}\|^{2}.$$

Therefore, the conformal curvature case [50] is characterised by the technical constants⁶

$$\ell_t = 2\,\ell_b \qquad (\beta = \gamma) \qquad \text{and} \qquad \Psi = \frac{3}{2}.$$
 (10.18)

This is the case of Foam @0.15mm (dense syntactic) and nearly satisfied for Foam @0.18mm. We also note that $\Psi = \frac{3}{2}$ corresponds to the curvature energy $\frac{\mu_e L_e^2}{2} \alpha_1 \| \text{dev sym Curl } \mathbf{A} \|^2$ and is sufficient for bounded stiffness in torsion [20], while $\beta = \gamma$ ($a_2 = 0$) is necessary for bounded stiffness in bending. Both limit cases violate uniform positivity of the curvature but are allowed by the weaker requirements in (6.5).

Under uniform positive definiteness of the energy, the admitted range for Ψ is (see Table 2)

$$0 < \Psi < \frac{3}{2}$$
 (together with $\beta - \gamma > 0$). (10.19)

According to our weaker requirements (8.2) the admitted range for Ψ is

$$0 < \Psi \le \frac{3}{2}$$
 (a₃ = 0) (and $\beta - \gamma = 0$, too). (10.20)

The latter limit value $\Psi = \frac{3}{2} (a_3 = 0)$ has been consistently taken in the identification by Lakes, see Tables 3 and 4.

If only one of the two conditions (10.16) characterising the conformal curvature case is satisfied we have: for $\alpha_2 = 0 \Leftrightarrow a_2 = 0 \Leftrightarrow \ell_t = 2 \ell_b \Leftrightarrow \beta = \gamma$ the internal energy density is

$$W = \mu_{e} \|\operatorname{dev}\operatorname{sym} \mathbf{e}\|^{2} + \mu_{c} \|\operatorname{skew} \mathbf{e}\|^{2} + \frac{2\mu_{e} + 3\lambda_{e}}{6} [\operatorname{tr}(\mathbf{e})]^{2} + \frac{\mu_{e}L_{c}^{2}}{2} \left[a_{1}\|\operatorname{dev}\operatorname{sym}\operatorname{Curl} \mathbf{A}\|^{2} + \frac{a_{3}}{3} \operatorname{tr}(\operatorname{Curl} \mathbf{A})^{2}\right] \\ = \mu_{e} \|\operatorname{dev}\operatorname{sym} \mathbf{e}\|^{2} + \mu_{c} \|\operatorname{skew} \mathbf{e}\|^{2} + \frac{2\mu_{e} + 3\lambda_{e}}{6} [\operatorname{tr}(\mathbf{e})]^{2} + \frac{\mu_{e}L_{c}^{2}}{2} \left[\alpha_{1} \|\operatorname{dev}\operatorname{sym} \mathfrak{K}\|^{2} + \frac{2\alpha_{1} + 3\alpha_{3}}{6} [\operatorname{tr}(\mathfrak{K})]^{2}\right],$$

while for $2\alpha_1 + 3\alpha_3 = 0 \Leftrightarrow a_3 = 0 \Leftrightarrow \Psi = \frac{3}{2}$ the internal energy density is

$$W = \mu_{\mathbf{e}} \|\operatorname{dev}\operatorname{sym} \mathbf{e}\|^{2} + \mu_{\mathbf{c}} \|\operatorname{skew} \mathbf{e}\|^{2} + \frac{2\mu_{\mathbf{e}} + 3\lambda_{\mathbf{e}}}{6} [\operatorname{tr}(\mathbf{e})]^{2} + \frac{\mu_{\mathbf{e}}L_{\mathbf{c}}^{2}}{2} \left[a_{1}\|\operatorname{dev}\operatorname{sym}\operatorname{Curl} \mathbf{A}\|^{2} + a_{2}\|\operatorname{skew}\operatorname{Curl} \mathbf{A}\|\right]$$
$$= \mu_{\mathbf{e}} \|\operatorname{dev}\operatorname{sym} \mathbf{e}\|^{2} + \mu_{\mathbf{c}} \|\operatorname{skew} \mathbf{e}\|^{2} + \frac{2\mu_{\mathbf{e}} + 3\lambda_{\mathbf{e}}}{6} [\operatorname{tr}(\mathbf{e})]^{2} + \frac{\mu_{\mathbf{e}}L_{\mathbf{c}}^{2}}{2} \left[\alpha_{1} \|\operatorname{dev}\operatorname{sym}\mathfrak{K}\|^{2} + \alpha_{2} \|\operatorname{skew}\mathfrak{K}\|^{2}\right].$$

Using these direct identification, in Table 4 we provide the numerical values of the material parameters considered in our formulation of the Cosserat theory of isotropic elastic solid, according to the experimental results obtained in [27] (see Table 3).

Acknowledgements. The authors are indebted to Roderic Lakes (Wisconsin Distinguished Professor, University of Wisconsin-Madison) for the motivation to provide this comprehensive comparison. The work of I.D. Ghiba was supported by a grant of the Romanian Ministry of Research and Innovation, CNCS–UEFISCDI, Project no. PN-III-P1-1.1-TE-2019-0348, Contract No. TE 8/2020, within PNCDI III. Angela Madeo and Gianluca Rizzi acknowledges support from the European Commission through the funding of the ERC Consolidator Grant META-LEGO, N° 101001759. Angela Madeo and Gianluca Rizzi acknowledge funding from the French Research Agency ANR, "METASMART" (ANR-17CE08-0006). Angela Madeo thanks IUF (Institut Universitaire de France) for its support. Patrizio Neff acknowledges support in the framework of the DFG-Priority Programme 2256 "Variational Methods for Predicting Complex Phenomena in Engineering Structures and Materials", Neff 902/10-1, Project-No. 440935806.

⁶These two conditions are independent.

References

- H. Altenbach, V. Eremeyev, L. Lebedev, and L.A. Rendón. Acceleration waves and ellipticity in thermoelastic micropolar media. Archive of Applied Mechanics, 80(3):217–227, 2010.
- W.B. Anderson and R.S. Lakes. Size effects due to Cosserat elasticity and surface damage in closed-cell polymethacrylimide foam. Journal of Materials Science, 29:6413–6419, 1994.
- [3] S. Bauer, P. Neff, D. Pauly, and G. Starke. New Poincaré-type inequalities. Comptes Rendus Mathematique, 352(2):163–166, 2014.
- [4] S. Bauer, P. Neff, D. Pauly, and G. Starke. Dev-Div- and DevSym-DevCurl-inequalities for incompatible square tensor fields with mixed boundary conditions. ESAIM: Control, Optimisation and Calculus of Variations, 22(1):112–133, 2016.
- [5] W.D. Claus and A.C. Eringen. Three dislocation concepts and micromorphic mechanics. In Developments in Mechanics, Proceedings of the 12th Midwestern Mechanics Conference, volume 6, pages 349–358. Midwestern, 1969.
- [6] W.D. Claus and A.C. Eringen. Dislocation dispersion of elastic waves. International Journal of Engineering Science, 9:605–610, 1971.
- [7] E. $\operatorname{Cosserat}$ and F. Cosserat. Théorie desdéformables. Librairie Sciencorpsavailable Hermann Fils D. Delphenich 2007tifique Α. et (engl. translation by pdf at http://www.mathematik.tu-darmstadt.de/fbereiche/analysis/pde/staff/neff/patrizio/Cosserat.html), reprint 2009 by Hermann Librairie Scientifique, ISBN 978 27056 6920 1, Paris, 1909.
- [8] S. Cowin. An incorrect inequality in micropolar elasticity theory. Zeitschrift f
 ür Angewandte Mathematik und Physik, 21:494– 497, 1970.
- [9] P. Duhem. Le potentiel thermodynamique et la pression hydrostatique. In Annales scientifiques de l'École Normale Supérieure, volume 10, pages 183–230, 1893.
- [10] W. Ehlers and S. Bidier. Cosserat media. In H. Altenbach and A. Öchsner, editors, Encyclopedia of Continuum Mechanics, pages 436–446. Springer, 2020.
- [11] V.A. Eremeyev. Acceleration waves in micropolar elastic media. Doklady Physics, 50:204–206, 2005.
- [12] V.A. Eremeyev, L.P. Lebedev, and H. Altenbach. Foundations of Micropolar Mechanics. Springer Science & Business Media, 2012.
- [13] A.C. Eringen. Theory of Micropolar Elasticity. In H. Liebowitz, editor, Fracture. An advanced treatise., volume II, pages 621–729. Academic Press, New York, 1968.
- [14] A.C. Eringen. Microcontinuum Field Theories. Springer, Heidelberg, 1999.
- [15] A.C. Eringen and W.D. Claus. A micromorphic approach to dislocation theory and its relation to several existing theories. In J.A. Simmons, R. de Wit, and R. Bullough, editors, *Fundamental Aspects of Dislocation Theory.*, volume 1 of *Nat. Bur. Stand. (U.S.), Spec. Publ.*, pages 1023–1040. Spec. Publ., 1970.
- [16] R.D. Gauthier and W.E. Jahsman. A quest for micropolar constants. ASME Journal of Applied Mechanics, 42:369–374, 1975.
- [17] I.D. Ghiba, P. Neff, A. Madeo, and I. Munch. A variant of the linear isotropic indeterminate couple stress model with symmetric local force-stress, symmetric nonlocal force-stress, symmetric couple-stresses and complete traction boundary conditions. *Mathematics and Mechanics of Solids*, 22: 1221-1266, 2017.
- [18] S. Hassanpour and G.R. Heppler. Micropolar elasticity theory: a survey of linear isotropic equations, representative notations, and experimental investigations. *Mathematics and Mechanics of Solids*, 22(2):224–242, 2017.
- [19] R. Izadi, M. Tuna, P. Trovalusci, and E. Ghavanloo. Torsional characteristics of carbon nanotubes: Micropolar elasticity models and molecular dynamics simulation. *Nanomaterials*, 11(2):453, 2021.
- [20] J. Jeong and P. Neff. Existence, uniqueness and stability in linear Cosserat elasticity for weakest curvature conditions. Mathematics and Mechanics of Solids, 15(1):78–95, 2010.
- [21] J. Jeong, H. Ramezani, I. Münch, and P. Neff. A numerical study for linear isotropic Cosserat elasticity with conformally invariant curvature. Zeitschrift für Angewandte Mathematik und Mechanik, 89(7):552–569, 2009.
- [22] H. Khan, I.D. Ghiba, A. Madeo, and P. Neff. Existence and uniqueness of Rayleigh waves in isotropic elastic Cosserat materials and algorithmic aspects. Wave Motion, page 102898, 2022.
- [23] R.S. Lakes. Size effects and micromechanics of a porous solid. Journal of Materials Science, 18:2572–2580, 1983.
- [24] R.S. Lakes. Experimental microelasticity of two porous solids. International Journal of Solids and Structure, 22:55–63, 1985.
- [25] R.S. Lakes. A pathological example in micropolar elasticity. ASME Journal of Applied Mechanics, 52:234–235, 1985.
- [26] R.S. Lakes. Foam structure with negative Poisson's ration. Science, 235:1038–1040, 1987.
- [27] R.S. Lakes. Experimental methods for study of Cosserat elastic solids and other generalized elastic continua. http://silver.neep.wisc.edu/~lakes/CossRv.pdf. In H.B. Mühlhaus, editor, Continuum Models for Materials with Microstructure., pages 1–25. Wiley, 1995.
- [28] R.S. Lakes. Physical meaning of elastic constants in Cosserat, void, and microstretch elasticity. Journal of Mechanics of Materials and Structures, 11(3):217–229, 2016.
- [29] R.S. Lakes. Stability of Cosserat solids: size effects, ellipticity and waves. Journal of Mechanics of Materials and Structures, 13(1):83–91, 2018.

- [30] R.S. Lakes. Softening of Cosserat sensitivity in a foam: Warp effects. International Journal of Mechanical Sciences, 192:106125, 2021.
- [31] J. Lankeit, P. Neff, and D. Pauly. Uniqueness of integrable solutions to $\nabla \zeta = G \zeta$, $\zeta|_{\Gamma} = 0$ for integrable tensor coefficients G and applications to elasticity. *Zeitschrift für Angewandte Mathematik und Physik*, 64:1679–1688, 2013.
- [32] P. Lewintan, S. Müller, and P. Neff. Korn inequalities for incompatible tensor fields in three space dimensions with conformally invariant dislocation energy. *Calculus of Variations and Partial Differential Equations*, 60(4):1–46, 2021.
- [33] P. Lewintan and P. Neff. L^p-trace-free generalized Korn inequalities for incompatible tensor fields in three space dimensions. Proceedings of the Royal Society of Edinburgh Section A: Mathematics, pages 1–32, 2021.
- [34] P. Lewintan and P. Neff. Nečas–Lions lemma revisited: An L^p -version of the generalized Korn inequality for incompatible tensor fields. *Mathematical Methods in the Applied Sciences*, 44(14):11392–11403, 2021.
- [35] A. Madeo, P. Neff, I. D. Ghiba, L. Placidi, and G. Rosi. Band gaps in the relaxed linear micromorphic continuum. Zeitschrift für Angewandte Mathematik und Mechanik, 95(9):880–887, 2015.
- [36] A. Madeo, P. Neff, I. D. Ghiba, L. Placidi, and G. Rosi. Wave propagation in relaxed linear micromorphic continua: modelling metamaterials with frequency band-gaps. *Continuum Mechanics and Thermodynamics*, 27:551–570, 2015.
- [37] A. Madeo, P. Neff, I.D. Ghiba, and G. Rosi. Reflection and transmission of elastic waves in non-local band-gap metamaterials: a comprehensive study via the relaxed micromorphic model. *Journal of the Mechanics and Physics of Solids*, 95:441–479, 2016.
- [38] A. Madeo, I.D. Ghiba, P. Neff, and I. Munch. A new view on boundary conditions in the Grioli-Koiter-Mindlin-Toupin indeterminate couple stress model. *European Journal of Mechanics A/Solids*, 59: 294-322, 2016.
- [39] R.D. Mindlin. Stress functions for a Cosserat continuum. International Journal of Solids and Structures, 1(3):265–271, 1965.
- [40] R.D. Mindlin. Microstructure in linear elasticity. Archive for Rational Mechanics and Analysis, 16: 51-78, 1964.
- [41] R.D. Mindlin and H.F. Tiersten. Effects of couple stresses in linear elasticity. Archive for Rational Mechanics and Analysis, 11:415–447, 1962.
- [42] I. Münch, P. Neff, A. Madeo, and I.D. Ghiba. The modified indeterminate couple stress model: Why Yang et al.'s arguments motivating a symmetric couple stress tensor contain a gap and why the couple stress tensor may be chosen symmetric nevertheless, Zeitschrift für Angewandte Mathematik und Mechanik, 97: 1524-1554,2017.
- [43] P. Neff. Existence of minimizers for a geometrically exact Cosserat solid. Proceedings in Applied Mathematics and Mechanics, 4(1):548–549, 2004.
- [44] P. Neff. On material constants for micromorphic continua. Trends in Applications of Mathematics to Mechanics, STAMM Proceedings, Seeheim, pages 337–348, 2004.
- [45] P. Neff. Existence of minimizers for a finite-strain micromorphic elastic solid. Proceedings of the Royal Society of Edinburgh Section A, 136:997–1012, 2006.
- [46] P. Neff, M. Bîrsan, and F. Osterbrink. Existence theorem for geometrically nonlinear Cosserat micropolar model under uniform convexity requirements. *Journal of Elasticity*, 121(1):119–141, 2015.
- [47] P. Neff, A. Fischle, and I. Münch. Symmetric Cauchy-stresses do not imply symmetric Biot-strains in weak formulations of isotropic hyperelasticity with rotational degrees of freedom. Acta Mechanica, 197:19–30, 2008.
- [48] P. Neff, I.D. Ghiba, M. Lazar, and A. Madeo. The relaxed linear micromorphic continuum: well-posedness of the static problem and relations to the gauge theory of dislocations. *The Quarterly Journal of Mechanics and Applied Mathematics*, 68:53–84, 2015.
- [49] P. Neff, I.D. Ghiba, A. Madeo, L. Placidi, and G. Rosi. A unifying perspective: the relaxed linear micromorphic continuum. Continuum Mechanics and Thermodynamics, 26:639–681, 2014.
- [50] P. Neff, J. Jeong, and A. Fischle. Stable identification of linear isotropic Cosserat parameters: bounded stiffness in bending and torsion implies conformal invariance of curvature. Acta Mechanica, 211(3-4):237–249, 2010.
- [51] P. Neff, A. Madeo, G. Barbagallo, M.V. d'Agostino, R. Abreu, and I.D. Ghiba. Real wave propagation in the isotropic-relaxed micromorphic model. Proceedings of the Royal Society A: Mathematical, Physical, 473(2197):20160790, 2017.
- [52] P. Neff and I. Münch. Curl bounds Grad on SO(3). ESAIM: Control, Optimisation and Calculus of Variations, 14(1):148–159, 2008.
- [53] P. Neff, D. Pauly, and K.J. Witsch. A canonical extension of Korn's first inequality to H(Curl) motivated by gradient plasticity with plastic spin. Comptes Rendus Mathématique. Académie des Sciences. Paris, 349(23):1251–1254, 2011.
- [54] P. Neff, D. Pauly, and K.J. Witsch. Maxwell meets Korn: A new coercive inequality for tensor fields in $\mathbb{R}^{n \times n}$ with squareintegrable exterior derivative. *Mathematical Methods in the Applied Sciences*, 35(1):65–71, 2012.
- [55] P. Neff, D. Pauly, and K.J. Witsch. On a canonical extension of Korn's first and Poincaré's inequalities to H(Curl). Journal of Mathematical Sciences, 185(5):721–727, 2012.
- [56] P. Neff, D. Pauly, and K.J. Witsch. Poincaré meets Korn via Maxwell: Extending Korn's first inequality to incompatible tensor fields. *Journal of Differential Equations*, 258(4):1267–1302, 2015.
- [57] P. Neff, I. Münch, I.D. Ghiba, and A. Madeo. On some fundamental misunderstandings in the indeterminate couple stress model. A comment on recent papers of A.R. Hadjesfandiari and G.F. Dargush. *International Journal of Solids and Structures*, 81:233-243, 2016.

- [58] W. Nowacki. Theory of Micropolar Elasticity. Springer, 1972.
- [59] J.F. Nye. Some geometrical relations in dislocated crystals. Acta Metallurgica, 1:153–162, 1953.
- [60] H.C. Park and R.S. Lakes. Torsion of a micropolar elastic prism of square cross section. International Journal of Solids and Structure, 23:485–503, 1987.
- [61] G. Rizzi, G. Hütter, A. Madeo, and P. Neff. Analytical solutions of the cylindrical bending problem for the relaxed micromorphic continuum and other generalized continua. *Continuum Mechanics and Thermodynamics*, 33(4):1505–1539, 2021.
- [62] G. Romano, R. Barretta, and M. Diaco. Micromorphic continua: non-redundant formulations. Continuum Mechanics and Thermodynamics, 28(6):1659–1670, 2016.
- [63] M. Shirani, D.J. Steigmann, and P. Neff. The Legendre-Hadamard condition in Cosserat elasticity theory. The Quarterly Journal of Mechanics and Applied Mathematics, 73:293–303, 2020.
- [64] A. Taliercio. Torsion of micropolar hollow circular cylinders. Mechanics Research Communications, 37(4):406–411, 2010.
- [65] C. Truesdell. Zusammenfassender Bericht. Die Entwicklung des Drallsatzes. Zeitschrift f
 ür Angewandte Mathematik und Mechanik, 44(4-5):149–158, 1964.
- [66] W. Voigt. Theoretische Studien über die Elasticitätsverhältnisse der Krystalle. I Ableitung der Grundgleichungen aus der Annahme mit Polarität begabter Moleküle. Abh. königl. Gesell. Wiss. Göttingen (math. Klasse), 34(3), 1887.
- [67] J.F.C. Yang and R.S. Lakes. Transient study of couple stress effects in human compact bone. Journal of Biomechanical Engineering, 103:275–279, 1981.