Research Report

on

Assessment of Consumers' Awareness of Nutrition, Food Safety and Hygiene in the Life Cycle for Sustainable Development of Bangladesh: A Study on Sylhet City Corporation.

Professor Dr. Md. Nazrul Islam

Professor, Department of Business Administration

June-2021

University Research Centre

Shahjalal University of Science and Technology Sylhet-3114

Assessment of Consumers' Awareness of Nutrition, Food Safety and Hygiene in the Life Cycle for Sustainable Development of

Bangladesh: A Study on Sylhet City Corporation ¹

¹ Funded by University Research Centre, Shahjalal University of Science and Technology through Promotional Research Grant–One year



Submitted to

University Research Centre

Shahjalal University of Science and Technology, Sylhet-3114

Project code: SS/2020/1/11

Project category: Promotional Research Grant – One year

Submitted by

Principal Investigator: Professor Dr. Md. Nazrul Islam

Department of Business Administration,

Shahjalal University of Science and Technology, Sylhet-3114

Research Assistant

Fazlur Rahman

MS in Statistics, Department of Statistics, Shahjalal University of Science and Technology

June, 2021

Acknowledgement

The project was funded by University Research Centre, Shahjalal University of Science and Technology Sylhet-3114, so my first I show my highest gratitude to the Director of the center and also its members for making research opportunity by granting the fund. I am also indebted to the data collectors and research assistant their sincere cooperation in conducting the research.

(Professor Dr. Md. Nazrul Islam)

Professor, Department of Business Administration

and

Principal Investigator of the project.

Assessment of Consumers' Awareness of Nutrition, Food Safety and Hygiene in the Life Cycle for Sustainable Development of Bangladesh: A Study on Sylhet City Corporation

Abstract

The objective of the study is to explore the knowledge of consumers on nutrition, food safety and hygiene of the sampled respondents in the Sylhet City Corporation (SCC) in Bangladesh in terms of different demographic characteristics. This study includes the different types of consumers in the life cycle viz. baby, child, adolescent, young and old so that an overall awareness level can be measured in the urban area of Bangladesh. The study is confined to SCC area in which all types of respondents has been included and findings from this study will be used generally for Bangladesh in making policy In conducting the study the population has been divided into six group as: (i) Baby; (ii) child; (iii) adolescent; (iv) parental; (v) unmarried adult young and (vi) married adult matured. We find that the average score of awareness of food nutrition and hygiene of unmarried adult is higher than that of married adults. The study suggested it is needed to increase awareness in of the parents for feeding babies. The average awareness of parents to their child's eating behavior between 5-9 years is 3.36 out of 5. The awareness is around 67% so we should be more careful in this regard. The average awareness adolescent (10-19 years) food habit is 1.89 on three points scales which about 63% only. Therefore, the consciousness of adolescent (10-19 years) has been to increase in taking food. The average feeding styles of parents is 4.24 out of 5 to their children up to 9 years and in percentage it is 84%.

Executive Summary

The food systems approach initiated in the 2011 First Country Investment Plan (CIP-1) under its third component of food utilization through program later in the Second Country Investment Plan which focused on Nutrition Sensitive Food Systems reflects the multi-sector collaboration for ensuring Nutrition, Food safety and Hygiene (NFSH). Achieving adequate nutrition and food security for all entails that the right foods are available, and that people are able to physically and financially access those foods. But an underlying assumption is that people are willing to eat the right foods and prepare and to eat them in a manner that conserves their nutritious value and that is hygienic. Thus, Bangladesh still consumes far more rice per capita than recommended by dietary guidelines. Feeding and hygiene practices are often inadequate with dire consequences for small children's nutritional status and development. This is true across socio-economic categories. The World Food Summit-1996 emphasized on three important components for ensuring food security viz. availability of adequate food, stability in food supplies, and access to food and nutrition security. The present government of Bangladesh has given emphasis on poverty alleviation and elimination including all forms of disparity at the forefront of country's development strategy. The ending of poverty and hunger is one of the prime objectives of the Sustainable Development Goals (SDGs) to be implemented by 2030. In consonance with the SDGs, the government of Bangladesh has adopted Seventh Five Year Plan (SFYP), in which one of the targets has been fixed to bring down extreme poverty to around 8.9% by the year 2020. In the SFYP the government has identified that there is persistent micronutrient deficiencies, lack of public awareness, maternal under nutrition, acute malnutrition and lack of dietary diversity are among other challenges in the nutrition program sub-sector. In Bangladesh there are very few studies and all the studies are made on specific group on the awareness of food safety, nutrition and hygiene of the consumers. But the present study is an in-depth study which has included the different types of consumers in the life cycle viz. baby, child, adolescent, young and old so that an overall awareness level can be measured in the urban area of Bangladesh. The study is confined to SCC area in which all types of respondents has been included and findings from this study will be used generally for Bangladesh in making policy. The core objective of the study is to explore the knowledge of consumers on nutrition, food safety and hygiene of the sampled

respondents in the Sylhet City Corporation in Bangladesh in terms of different demographic characteristics. In conducting the study the population has been divided into six

group as: (i) Baby; (ii) child; (iii) adolescent; (iv) parental; (v) unmarried adult young and (vi) married adult matured. Separate questionnaire for each group has been used for collecting data. Total sample size is 234 under the six group as (i) Baby-40; (ii) child-40; (iii) adolescent-34; (iv) parental-34; (v) unmarried adult young -40 and (vi) married adult matured-40. The collected data has been analysed by different techniques of (Frequency distribution mean and standard deviation of the variables have been computed to get the picture of the data. Tables and Graphs) and inferential statistics (Principal Component analysis (PCA) and two sample t-test have been used for conclusion about the data) have also been used to present the data. The report is divided into three chapters where first chapter has comprehended the introductory aspects of the study such as problem statement, methodology and review of related literature. The second chapter comprises the analysis and presentation of the collected data with statistical methods and techniques. Finally the third chapter has depicted the findings and suggestions of the study.

In order to find the Baby's Eating Behaviour up to 4(Four Year) Responded by Parents 20 mother and 20 father were interviewed with a questionnaire with five point scale as Table-never =1 rarely=2 sometimes =3 "Baby often=4 and Always =5. The average frequency in percentage in the different scales are 23%, 30%, 16%,11% and 11% respectively. There were 18 scaling questionnaire and the average score of the items is 100.5 and the average score of the respondents is 2.51. It means the eating behaviour of baby lies between rarely and sometimes. Reliability of the questionnaire was checked by Cronbach's Alpha test which is 0.710 that means there is consistency of the items or variables. The PCA reduces 21 items into six factors significant for eating behaviours of parents. In assessing the Child's eating behaviour between years similarly 20 fathers and 20 mothers were given questionnaire with 35 items or variables in the five point scale like previously. The average percentage of frequency distribution in the five point scales are 20%,27%, 23%, 16% and 19% respectively. The average of items is 117.66 and the average of respondents is 3.36. That means the eating behaviour of children varied between sometimes and often. Through the PCA 35 items are reduced to 10 factors. The food habits of adolescent (10-19) is measured in terms of given items or variables with three point scale as Most like=2; like=1 and Neutral=0; with 23 items. The average

percentages of frequency in the scales are 37%, 55% and 8% respectively. The data showed that that the average of score of the items is 43.56 and the average score of respondents is 1.89. It means the choice varies between most like and like. The PCA reduces 21 items to eight factors of food habits of adolescent. Parental Feeding Style to his Children up to 9 years has been measured with five point scale 27 items. The average percentage of frequency distribution in the five point scales are 15%, 10%, 33%, 13% and 30% respectively. The Average of items is 114.52 and the average of respondents is 4.24 that mean the parental Feeding Style to his Children up to 9 years varies between often and always. The total variance explained by the PCA is 78.96% that means the PCA is accepted and based on Eigen values seven items are considered important. The food preference of adults unmarried (20-29) has also been measured by the same five pint scale with 62 items. The average percentages of frequency distribution in the different scales are 21%, 8%, 17% 31% and 24% respectively. The Average of items is 131.35 and the average of respondents is 2.12 that mean the food preference of adults (20-29 years) varies between neither like or dislike and like a little. The Food Preference for Adults (29-64) Married Male and Female has been measured by the same questionnaire as adult unmarried that means 62 items with five point scale. The average percentages of frequency distribution in the different scales are 37%, 16%, 16% 18% and 14% respectively The Average of items is 102.87 and the average of respondents is 1.66 that means the food preference of adults (29-64 years) varies between dislike a little and neither like nor dislike. A Comparison of food preference between adults of 20-29 unmarried and 29-64 married is made with two sample t-test to confirm whether there is any similarity between these two groups. We found that mean score of unmarried and married are 191.10 and 102.87 correspondingly with Std. deviation 89.873 and 39.889 for unmarried and married respectively. Therefore the average score of unmarried adult is higher than that of married adults. Levene's Test for Equality of Variances and t-test for Equality of means were used also. In the first test i.e. Levene's Test for Equality of Variances here our null hypothesis is that the both the variances are equal and alternative is that the variances are different for the two groups. The Levene's Test for Equality of Variances showed the p-value is 0.00 that means there are significant differences of variances of the two groups. The second test is t-test for Equality of Means which is presented in the second row and showed that the observed test statistics is 7.065 with p-value 0.000. In this case we can also reject the null hypothesis and can conclude that there is significant difference between the means of the two

groups. Finally we can conclude that there is significant difference between the food preference of unmarried adults and married adults and their preferences s are not same. The overall score of awareness of food nutrition and hygiene of six groups are: (i) Parents of baby up to 4 years = 2.51(in 5 point scale); (ii) Parents of Child eating behaviour between 5-9 years = 3.36(in 5 point scale); (iii) Adolescent (10-19 years) Food Habits= 1.89(in 3 point scale); (iv) Parental Feeding Style to his Children up to 9 years= 4.24(in 5 point scale);; (v) Food preference of adults unmarried (20-29)=2.12 (in 5 point scale); and (vi)Food Preference for Adults (29-64) Married Male and Female=1.61(in 5 point scale). The study suggested it is needed to increase awareness in of the parents for feeding babies. The average awareness of parents to their child's eating behaviour between 5-9 years is 3.36 out of 5. The awareness is around 67% so we should be more careful in this regard. The average awareness adolescent (10-19 years) food habit is 1.89 on three points scales which about 63% only. Therefore the consciousness of adolescent (10-19 years) is be to increase in taking food. The average feeding styles of parents is 4.24 out of 5 to their children up to 9 years and in percentage it is 84%. Therefore it is about to satisfactory.

Table of Contents

	Cover page	i					
	Acknowledgement	ii					
	Executive Summary	iii-ix					
	List of Tables						
	List of Figures	xiv					
	List of Acronyms	XV					
CHAPTER I	INTRODUCTORY ASPECTS OF THE STUDY	01-09					
1.1	An Overview of Haor Area in Bangladesh	01					
1.2	Sunamganj District	02					
1.3	Poverty in Bangladesh	03					
1.3.1	Measurement of the Incidence of Poverty in Bangladesh	04					
1.3.2	Trends of Poverty	04					
1.4	Statement of the Problem	05					
1.5	Objectives of the Study	07					
1.6	Research Questions and Corresponding Expected Results	07					
1.7	Relevance of the Project	08					
CHAPTER II	REVIEW OF RELATED LITERATURE	10-16					
2.	Literature Review	10					
CHAPTER III	METHODOLOGY OF THE STUDY	17-21					
3.1	Study Area	17					
3.2	Sample and Sample Design	18					
3.3	Questionnaire Design	19					
3.4	Collection of Data	19					
3.5	Analysis of Data	19					
CHAPTER IV	ANALYSIS OF DATA	22-45					

4.1	Demographic Profile of the Respondents	22
4.2	Demographic Profile of Household Members	22
4.3	Housing Information of the Households	25
4.4	Possession of Durable Assets by Households	25
4.5	Possession of Productive Assets by Households	26
4.5.1	Landholdings by the households	26
4.6	Households' Knowledge about Micro-credit Benefit	28
4.7	Factors of Exclusion from Micro-credit Program	28
4.8	Profile of Micro-credit Programs	30
4.9	Purpose of Loan	31
4.9.1	Descriptive on the major purposes of loan	31
4.9.2	Major Dimensional purpose of taking Loan	31
4.10	Expenditure and Investment Pattern of Loans	33
4.11	Socio-economic Impact of Micro-credit on borrowers:	34
4.11.1	Impact of informal micro-credit through economic performance	
	based on before-after comparison:	34
4.11.2	Impact of formal micro-credit through economic performance	
	based on before-after comparison:	35
4.11.3	Impact of micro-credit in terms of formal and informal credit	
	receiving households based on Difference-in-difference Method	
	(DID)	35
4.11.4	Percentage change of households' benefits using micro-credit	
	based on did method	35
4.11.5	Percentage distribution of households' food insecurity during the	
	study period	36
4.11.6	Percentage distribution of households self-assessed socio-	
	economic status	36
4.11.7	Impact of micro-credit on changing educational and healthcare	
	expenditure in 2019/20 compared to 2016/17	37
4.12	Causes of non-payment of loan	38
4.13	Attitude of Borrowers on Micro-credit	41
4.14	Causes of Not Over-coming from Vicious Cycle of Poverty	43
		_

CHAPTER V	FOCUS GROUP DISCUSSION	46-48
5.1	Issues of Focus Group Discussions (FGD)	46
5.2	Outcomes of FGD	46
CHAPTER VI	SUMMARY AND CONCLUSION OF THE STUDY	49-59
	References	60-69
	Annex-1: Appendix Tables	70-82
	Annex-2: Survey Questionnaire	83-90

List of Tables

Table No.	Table No. Title of the Tables					
Table 1.1	Distribution of <i>Haor</i> Areas in Bangladesh.	01				
Table 1.2	Incidence of Poverty in Bangladesh.	05				
Table 1.3	Cooperative Development Indicators of Bangladesh and	06				
	Sunamganj District.					
Table 1.4	Research Objectives, Research Questions and	07				
	Corresponding Expected Results.					
Table 1.5	Targets in Seventh Five Year Plan to Achieved through the	09				
	Study.					
Table 3.1	Sampling Frame of the Study.	19				
Table 4.2.1	Dropouts and Basic Educational Indicators of the	24				
	Household Population.					
Table 4.4	Possession of Durable Assets by Households Micro-credit	25				
	Status.					
Table 4.5.2	Non-land productive assets of the households.	27				
Table 4.7.1	Major dimensional factors of excluded from the micro-	30				
	credit benefits.					
Table 4.9.2	Major dimensional purpose of taking loan	33				
Table 4.10	Distribution of Loan among Different Heads of	34				
	Expenditure in 2019/20.					
Table 4.11.5	Percentage distribution of households' food insecurity for	36				
	the period of 2016/17 and 2019/20					
Table 4.11.6	Percentage distribution of households self-assessed socio-	37				
	economic status for the period of 2016/17 and 2019/20					
Table 4.11.7	Perceptions on change on educational and healthcare	38				
	expenditure in 2019 compared to 2016.					
Table 12.1	Causes of non-payment of loan in time.	38				
Table 4.12.2	Major dimensional causes of non-payment of loan timely	40				
Table 4.13.1	Major dimensional attitude of borrowers on micro-credit	43				
Table 4.14.1	Descriptive of the Causes of Not Over-coming from	44				

	Vicious Cycle of Poverty.				
Table 4.14.2	Major dimensional causes of not over-coming from vicious	45			
	cycle of poverty				
Appendix table 4.1	Demographic Characteristics of Respondents' by	70			
	Borrowing Status.				
Appendix table 4.2	Demographic Profile of Household (HH) Members of	71			
	Respondents.				
Appendix table 4.3	Housing Condition and Sanitation Facilities by Micro-	72			
	credit Status.				
Appendix table 4.5.1	Landholdings by Study Households.	73			
Appendix table 4.6	Household's Knowledge about Micro-credit Benefits.	74			
Appendix table 4.7	endix table 4.7 Causes of for Excluded from Micro-credit Program by				
	Non-borrower Participation.				
Appendix table 4.8	Profile of Micro-credit Programs.	76			
Appendix table 4.9.1	Purpose of Loan of the Study Households.				
Appendix table 4.11.1 Impact of Informal Micro-credit through Economic					
	Performance based on before-after Comparison				
Appendix table 4.11.2	Impact of Formal Micro-credit through Economic	79			
	Performance based on before-after Comparison.				
Appendix table 4.11.3	Impact of Micro-credit through Economic Performance in	80			
	terms of Formal and Informal Credit Receiving				
	Households based on Diff-in-diff Method.				
Appendix table 4.11.4	Percentage Change of Households' Economic Performance	81			
	due to Micro-credit Benefits based on Diff-in-diff Method.				
Appendix table 4.13	Descriptive statistics of the Attitude of Borrowers on	82			
	Micro-credits.				

List of Figures

Figure No.	Title of the Figures	Page no.
Figure 1.1	Haor Areas of Bangladesh.	02
Figure 1.2	Sunamganj District.	03
Figure 3.1	Study Area.	18
Figure 4.2.1	Age Structure of the Members of HHs.	23
Figure 4.2.2	Educational Status of the Members of HHs	23
Figure 4.2.3	Occupation of the Members of HHs.	24
Figure 4.1	Possession of Durable Assets.	26
Figure 4.5.1	Landholdings by Households	27
Figure 4.7	Scree Plot of Factors of Exclusion from Micro-credit Program.	29
Figure 4.9.2	Scree Plot of the Purpose of Taking Loan.	32
Figure 4.12	Causes of Non-payment of Loan.	39
Figure 4.13	The Scree Plot of Major Dimensional Attitude of Borrowers on Micro-	42
	credit.	
Figure 4.14	The Scree Plot of Major Dimensional Causes of Not Over-coming from	44
	Vicious Cycle of Poverty.	

List of Acronyms

BBS Bangladesh Bureau of Statistics **BHWDB** Bangladesh *Haor* and Wetland Development Board **CBN** Cost of Basic Needs **CEGIS** Center for Environment and Geographic Information Services Direct Calorie Intake DCI DID Difference-in-difference **FEI** Food Energy Intake **FGD** Focus Group Discussions GoB Government of Bangladesh НН Household HIES Household Income and Expenditure Survey LPL Lower Poverty Line LPL Lower Poverty Line MAS Multi-agent systems MC Micro-credit MDGs Millennium Development Goals MFI Micro Finance Institution MPI Multidimensional Poverty Index NGO Non-Government Organization **PCA Principal Component Analysis SACCOS** Savings and Credit Co-operative Societies **SFYP** Seventh Five Year Plan **SMCP** Savings and Micro Credit Program **SPSS** Statistical Package for Social Sciences

UP

Union Parishad

1.1.Inroduction: The food systems approach initiated in the 2011 First Country Investment Plan (CIP-1) under its third component of food utilization through program 10, 11 and 12 where total fund required were 736 billion US \$ (GoB, 2011). Later in the Second Country Investment Plan (CIP-2) which focused on Nutrition Sensitive Food Systems reflects the multi-sector collaboration for ensuring Nutrition, Food safety and Hygiene (NFSH). Achieving adequate nutrition and food security for all entails that the right foods are available, and that people are able to physically and financially access those foods. But an underlying assumption is that people are willing to eat the right foods and prepare and to eat them in a manner that conserves their nutritious value and that is hygienic. Thus, Bangladesh still consumes far more rice per capita than recommended by dietary guidelines. Feeding and hygiene practices are often inadequate (Manikam et al., 2017; Mahmud, 2016) with dire consequences for small children's nutritional status and development. This is true across socio-economic categories. All these issues materialize in poor nutritional outcomes where high rates of stunting coexist with fast rising levels of obesity. While efforts are made all round to ensure awareness of what adequate nutrition and safe practices are, there is only scattered evidence of the progress so far, which makes a nationwide coherent strategy difficult to develop. In the CIP-2 Investment Plan for Nutrition-Sensitive Food Systems, under its Program III.I proposes measures for 'Enhanced nutrition knowledge, promotion of good practices, and consumption of safe and nutritious diets'. The CIP-2 for this purpose requires investment of 89 million US \$. This strategy necessitates a thorough understanding of the situation of awareness of consumers on Nutrition, Food Safety and Hygiene.

1.2. Statement of the problem: The World Food Summit-1996 emphasized on three important components for ensuring food security viz. availability of adequate food, stability in food supplies, and access to food and nutrition security. The present government of Bangladesh has given emphasis on poverty alleviation and elimination including all forms of disparity at the forefront of country's development strategy. The ending of poverty and hunger is one of the prime objectives of the Sustainable Development Goals (SDGs) to be implemented by 2030. In consonance with the SDGs, the government of Bangladesh has adopted Seventh Five Year Plan (SFYP), in which one of the targets has been fixed to bring down extreme poverty to around

8.9% by the year 2020. In the SFYP the government has identified that there is persistent micronutrient deficiencies, lack of public awareness, maternal under nutrition, acute malnutrition, climate risks and adaptations, and lack of dietary diversity are among other challenges in the nutrition program sub-sector (GoB. SFYP, p. 519; Finance Division, 2014). In this plan nutrition and hygiene education has been incorporated in the education and ensure the availability of diverse foods quality (GoB. SFYP,p. 520). Food- borne illness is still an important issue in the world. Mishandling of food occur during food preparation, handling and storage and studies shows that consumers have inadequate knowledge about measures to prevent food borne illness in the home. The economic burden resulting from the diseases caused by foods on the health systems and the productivity losses they cause are at significant levels (Gil *et al.*, 2002; Banati, 2003).

In Bangladesh there is no general study on the awareness of NFSH. There are small studies on specific group viz. Karmakar (2017) made a study on a only 200 university students of *Noakhali* region of Bangladesh, Alam *et al.*, (2010) made a study on rural unmarried adolescent girls aged 13–18 years in 708 rural clusters to measure the nutritional status and dietary intake and relevant knowledge of adolescent girls in rural Bangladesh. Iqbal Kabir *et al.*,(2013) developed a Complementary Feeding Manual for Bangladesh and found that two- third of mothers did not have enough knowledge about nutrition. But one-third of them thought that vegetables, fish, egg, meat, water melon, banana, khichuri (one types mixed food of rice, dal, vegetables etc.), seasonal fruits contain a lot of vitamins. Therefore, in Bangladesh there are very few studies and all the studies are made on specific group on the awareness of food safety, nutrition and hygiene of the consumers. But the present study is an in-depth study which has included the different types of consumers in the life cycle viz. baby, child, adolescent, young and old so that an overall awareness level can be measured in the urban area of Bangladesh. Due to budget constraint the study is confined to SCC area in which all types of respondents has been included and findings from this study will be used generally for Bangladesh in making policy.

1.3. Review of literature: Safety food is the basic human right when billions of people are in risk of unsafe food and many millions become sick and hundreds of thousand die yearly (Fung, *et al.*, 2018). On an average 9.4 million people suffer from food –borne diseases illness and 1351 death in United States of America (Scallan, *et al.*, 2011). The estimated total economic loss

of foodborne illness ranged between US\$5.6 billion and US\$9.4 billion in USA and in Australia between \$487 million and \$1900 million (Cameron, et al., 1995). Food borne and water borne diseases kill about 2.2 million people in each year (WHO, 2004). In developing country like Bangladesh every year 2 million people including children are killed by food-borne disease (Daily star, 2015). The low income and minority people experience grater rate of food-borne illness (Quinlan, 2013). The foods available to consumers in Bangladesh are mostly contaminated as observed by the joint survey Institute of Public Health of Dhaka University and WHO (1994 and 2003). The water is also adulterated in Bangladesh (Saha, et al., 2011). Besides food adulteration irregular intake of breakfast and dinner, eating junk food and fried food regularly, lack of knowledge about balance nutrition and obesity, habits of eating outside of the residence causes illness (Karmakar, et al., 2016). The consumers' knowledge about food safety, nutrition and hygiene is very poor in Bangladesh and also around the world. The knowledge about food safety concern at all levels (family, institution, community and government) may reduce illness of people (Alam & Hossain, 2018; Karmakar, et al., 2016). The consumer must aware of product information, environmental impacts of their buying decisions, food handling behavior and knowing the temperature of the refrigerator and cooking temperature when removing the food from fridge you must reheat the food then only the bacteria disappear from the food (Alam & Zakaria, 2021; Subba rao, et al., 2007). Food safety is the most important predictor of attitude while health consciousness appears to be the least important motive in contrast to findings from some previous research and consumers ethically intent to purchase organic foods (Nina and Hassan, n.d) Consumer's awareness of food safety is very essential to avoid the risk of unsafe food and maintain good (Bhuvneswari and Muthukumar, 2015). Most of the peoples have unsatisfactory dietary habit and providing pupils with nutrition knowledge would promote healthy dietary behaviors (Doaa, et al., 2017). A study showed that overall dietary knowledge among the adolescent girls in rural Bangladesh is very low as compare to the semi-urban and small town areas (Islam, Alam, & Afzal, 2021) and more than half could not name the main food sources of energy and protein, and 36% were not aware of the importance of taking extra nutrients during adolescence for growth spurt (Nurul, et al., 2010). Even most of the doctors are lac of nutrition knowledge (Uddin, et al., 2008). The behavioral change communication should improve infant and young child nutrition knowledge (Hoddinott, 2017). It is important to measure the level of consumers' awareness on the safety of food to reduce food

-borne illness. The safety education campaign should raise knowledge about high risk foods and educate persons who will fall high-risk groups about their increased risk for food borne illness (Nesbitt, et al., 2014). In a study on complementary feeding practices living in high income countries showed that promoters influence complementary feeding practices including income, lack of knowledge and incorrect advice (Logan, et al., 2018). Due to ineffective communication of food safety message to consumers, causes of less awareness of food-borne illness and consumers fail to recognize foodborne disease rightly (Bruhn, 1997). The more consumers informed about food safety the more possibility of reducing risk from potential hazards of unsafe foods (Bruhn and Schutz, 1999). The Chinese people have high awareness of safe food but low knowledge on safe food, identification of safe food yet they have positive attitude on safe food (Liu, et al., 2013). Due to increase in awareness of unsafe food the demand of organic fruits and vegetables is increasing and dominating the basket of organic consumers and consumers are ready to pay more prices for organic products (Emmanuel, et al., 2005; Sedef, et al., 2007). There is a need to assess the food safety knowledge, attitude and practice of parents and food preparers as they are more likely to engage in risky eating behaviors (Nevin and Ece, 2012). The hotel and restaurants also include food safety and hygiene as they prepare food for consumers and it is given emphases by Jenny and Serli (2010). The consumers are associated food mostly often with safe food, and food safety is the topic most often discussed while tasty food is the most important food conversation topic (Van, et al., 2003). The assessment of consumers in this respect can be measured in terms of nutrition and food related Knowledge, Attitude and Practice (KAP) as per FAO (FAO, 2104). Therefore, the above literatures tell that the consumers are suffered from various diseases due to consumption of unsafe and un-hygienic food which causes the death of millions of people in Bangladesh and all over the world. This is because the consumers are not aware of unsafe food, required nutrition and hygiene environment. The awareness should be increased about unsafe food, nutrition food habit and hygiene for good health to all the stages of life cycle.

1.4. Rationale of the Study: The rationality of the study is discussed from the following perspectives:

(i) Filling the gap of existing knowledge: As we observed in the problem statement that there is very few studies on NFSH in Bangladesh and information on this issue is vital for good health of

the people in Bangladesh. Further there is no study on SCC and even Sylhet region. The findings of the study will add value to the existing store of knowledge in the field of research in Bangladesh and especially for the Sylhet region.

- (ii) Relationship with the Sustainable Development Goals (SDGs): The United Nations has set the 169 targets under 17 SDGs for transforming our world by 2030 for people, planet and prosperity. Among the 17 SDGs, SDG#1 is to end poverty in all its forms everywhere; and SDG# 2 is to end hunger, achieve food security and improved nutrition and promote sustainable agriculture and SDG# 3 is to ensure healthy lives and promote well-being for all at all ages. Through this project we will measure the level awareness of nutrition, food safety and hygiene of consumers so that government can prepare policy and program to ensure balanced nutrition, safety food and hygiene in food preparation and preservation. A large number of people are suffered from food borne ill as stated earlier and suffer from unbalanced nutrition which makes them ill and unhealthy. Further people do not properly make food and preserve them rightly for which they loss food value and suffer from unhygienic illness. The data from such type of an indepth study will surely help to policy makers in this regard which will help the government to achieve the SDGs 1, 2, and 3 directly. Further SDG# 12 is to ensure sustainable consumption and production patterns. The data will indirectly also help to achieve SDG# 12.
- (iii) Relevant to the Second Country Investment PlanCIP-2): In the CIP-2 under its Nutrition-Sensitive Food Systems in the Program III.1, has proposed to measures for 'Enhanced nutrition knowledge, promotion of good practices, and consumption of safe and nutritious diets'. Such strategy necessitates a thorough understanding of the baseline situation of the country. But due to limited fund the data of this in-depth study will help to make decision by the concerned authorities in this regard.
- (iv) Relevant to Seventh Five Year Plan (SFYP), vision 2021 and perspective plan (2010-2021): The Government seeks to create conditions whereby the people of Bangladesh have the opportunity to reach and maintain the highest attainable level of health. It is the vision of 2021 that recognizes health as a fundamental human right and, therefore, the need to promote health and to alleviate ill health and suffering in the spirit of social justice. The target of vision 2021 and perspective plan (2010-21) are: (a) 85% of the population have standard nutritional food; (b). poor people ensured a minimum of 2122 kilo calories of food; (c) all kinds of contagious

diseases eliminated (d)longevity increases to 70 years; and (e) infant mortality comes down to 15 from 54 per thousand at present. All these targets need to assess the awareness of NFSH of consumers in their life cycle.

1.5 . **Objectives of the study**: The core objectives of the study are to:

- (i) Explore the knowledge of consumers on nutrition, food safety and hygiene of the sampled respondents in the Sylhet City Corporation in Bangladesh in terms of different demographic characteristics.
- (ii) Compare knowledge level among the different groups of respondents ;and
- (iii) Identify the factors associated with the levels of awareness and suggestion for betterment of consumers on nutrition, food safety and hygiene.
- **1.6. Study Design**: The goal of the study was to make a base line survey on the consumers' awareness of nutrition, food safety and hygiene on different age groups of Sylhet City Corporation (SCC). To this aim the study has collected data from different types of consumers so that whole society is included in the survey. For this purpose the study has divided the population into six group as: (i) Baby; (ii) child; (iii) adolescent; (iv) parental; (v) unmarried adult young and (vi) married adult matured. Separate questionnaire for each group has been used for collecting data.
- **1.7. Sample Design**: Bangladesh Bureau of Statistics (BBS) developed primary sampling unit based on Population and Housing Census 2011 in 2012 to conduct various demographic and socio-economic surveys 61 in SCC. The PSU identified by BBS will be the base for the study where each PSU is comprised of 100 to 150 households. The list of PSU is available in the district's statistics office and also in BBS. For conducting the study 240 respondents have been interviewed from 61 PSU conveniently but in practice we collected 234 as given in table 1.1

Serial	Respondents	Number of	Scale of
No.		respondents	questionnaire
1.	Parents of baby up to 4 years	40	5.00
2.	Parents of Child eating behavior between 5-9 years	40	5.00
3.	Adolescent (10-19 years) Food Habits	34	3.00
4.	Parental Feeding Style to his Children up to 9 years	34	5.00
5.	Food preference of adults unmarried (20-29)	40	5.00
6.	Food Preference for Adults (29-64) Married Male and	40	5.00

Female		
Total	228	

The population up to 64 age is 95.06% of total population in Bangladesh (BBB, HIES, 2016)¹. The age up to 04 years is baby, age between 5 and 9 is children and age between 10 and 19 is adolescent (UNICEF, 2011)². More than 10% of target population is acceptable in a nation-wide survey.³

1.8. Questionnaire Design:

Construction of questionnaire: Our target respondents are six groups of people viz. (i) Baby; (ii) child; (iii) adolescent; (iv) parental; (v) adult young (vi) adult matured. The eating behaviour of baby and child will be received from mother and father, feeding behaviour of baby and child has been received by mother and father (parental behaviour). Habit of nutrition, safety food and hygiene of adolescent, young and adult old has been received from themselves respectively by both male and female.

<u>Variables of questionnaire</u>: Each questionnaire will be prepared to tap the factors associated with eating behaviours, feeding behaviours, food preference, food safety, nutrition knowledge, family eating and food buying behaviours, dietary practice, cooking practice, family hygiene knowledge and consciousness. The questionnaire has included demographic and socioeconomic factors of respondents. There are many sub-variables behind each variable so that total consumers awareness of nutrition, food safety and hygiene is explored in the Sylhet City Corporation.

<u>Validity of questionnaire</u>: Validity is characteristics of measurement concerned with the extent that a test measures what the researcher actually wishes to measure and that differences found with a measurement tool reflect true differences among participants drawn from the population (Cooper and Schindler, 2008)⁴. There are three types of validity viz. (i) content validity- it is done through a panel of experts; (ii) criterion related validity- it is done in terms of relevancy,

¹ BBS (2016), Household Income and Expenditure Survey, 2016.

² UNICEF (2011), The State of The World's Children

³ UNICEF (2005), Multiple indicator cluster survey manual, Monitoring the situation of children and woman

⁴ Cooper, Donald, R, Schindler & Pamela, S (2008), Business research methods, 9th Ed., New Delhi, The McGraw-Hill, p.24.

freedom from bias, reliability and availability; (iii) construct validity- is done by factor analysis and each factor is analysed individually using Principal Component Analysis(PCA). It is explained briefly under data analysis. Factor analysis groups' variables (i.e. single question) has been divided into factors based on their common correlation. Those variables which are correlated with other has been grouped together and such a group of variables is called a factor (Saraph *et al.*,1989)⁵. Each factor is assumed as a separate construct and must one or multidimensional.

<u>Test of internal consistency</u>: Testing internal consistency is to test the homogeneity among the items. This is one of the approaches to test reliability of the questionnaire. It is tested by Cronbach's alpha (\propto) values. A measure of scale reliability namely Cronbach's alpha (\propto) measures the internal consistency of responses, that is, how closely related a set of variables as a group. It can be written as a function of the number of test variables and the average inter correlation among the variables. Mathematically, it can be written as:

$$\alpha = \frac{N \, \bar{c}}{\bar{v} + (N-1)\bar{c}}$$

Where, N is the number of items in the data set, \bar{c} is the average inter-correlation among the items and \bar{v} is the average variance. The required value of Cronbach's alpha (\propto) for reliability is 0.70 (Nunnaly, 1967).

Measurement scale of questionnaire: Scaling is the procedure for the assignment of numbers or other symbols to a property of objects in order to impart some of the characteristics of numbers to the properties in question (Cooper and Schindler, 2008). To measure the intensity of the responses in respect of the statements pertaining to each of the criterions 5 point and 3 point rating scales have been used The Likert scales are probably more reliable and provide greater volume of data than many other scales (Edwards & Kenney, 1946). To answer the demographic question on characteristics, food choice, nutrition intake simple category scale, multiple choices, single response scale, and multiple response scale have also been used.

⁵ Saraph, J. V., Benson, P. G. and Schroeder, R. G. (1989) 'An instrument for measuring the critical factors of quality management'. Decision Sciences, Vol.20, No.4, pp.810-829.

⁶ Nunnaly, JC(1967), Psychometric theory, McGraw-Hill, New York.

⁷ Edwards, A. L., & Kenney, K. C. (1946). A comparison of the Thurstone and Likert techniques of attitude scale construction. Journal of Applied Psychology, 30, 72-83.

- **1.9. Analysis o:** Data has been analyzed in the following techniques:
- **1.91. Descriptive statistics:** Frequency distribution mean and standard deviation of the variables have been computed to get the picture of the data. Tables and Graphs have also been used to present the data.
- **1.9.2: Inferential statistics:** Principal Component analysis (PCA) and two sample t-test have been used for conclusion about the data.
- **1.10. Conclusion:** This chapter has presented the introductory aspects of the study viz. problem of the study, literature review and methodology of the study. The next chapter will present the analysis of data.

CHAPTER -TWO

ANALYSIS OF DATA

- **2. Introduction**: In the previous chapter we have discussed about the introductory aspects of the study such as statement of problem, related literature and methodology of the study. In this chapter we have analyzed the collected data following the methodology using different statistical techniques and methods. It is divided into eight sections and there are different sub sections under each section.
- **2.1. Analysis of Baby's Eating Behavior up to 4(Four Year) Responded by Parents:** The eating behavior of baby is analyzed by the dummy of their parents. The analysis is discussed through the following aspects.

2.1.1 Demographic profile and descriptive statistics of the respondents and the babies: Demographic profile and descriptive statistics of the respondents and the babies are presented in table 2.1.1

Table-2.1.1: Demographic profile and descriptive statistics of the respondents and the babies

Variable Number Variable		Minimum in	Maximum				
			years	years	Mean	Std. Deviation	
Sex of respondent	Sex of respondent Age of baby		0.3	4	3.035	1.1160	
Father	20	Descriptive S	tatistics of number of	of Baby			
Mother	20	Total baby	1	4	2.10	.955	
Total	40	Up to 4 years	1	2	1.10	.304	
		Descripti	ve statistics of pare	nts occupation			
Occ	cupation		Frequency	%	Valid %	Cumulative %	
	Business			42.5	42.5	42.5	
		Service	14	35.0	35.0	77.5	
t	oth service	and business	1	2.5	2.5	80.0	
Others	including	housekeeping	8	20.0	20.0	100.0	
		Total	40	100.0	100.0		
Descriptive Statistics	s of Family	's annual incom	e and expenditure				
				Maximum	Mean	Std. Dev.	
annual income(in thousand taka)			120	1200	502.00	249.134	
annual income(in thousand taka)			95	1000	390.40	217.485	
Descriptive Statistics	Descriptive Statistics of Education of Parents						
Education in terms	of Schoolin	ng year	8.0	17.0	13.675	2.9733	

Table 2.1.1 showed that out of 40 respondents 20 mother and 20 fathers. The age of the babies varied between 0.3 year and 4.00 and on average 3.035 with standard deviation 1.1160 .The number of total baby fluctuated is between 1 and 4 while number of baby under 4 years is ranged from 1 to 2. The maximum occupation of the parents are business (17) following service (14), others including housekeeping (8) and both service and business (1). The minimum, maximum and average of annual income of the responding household are 120 thousands, 1200 thousands and 502 thousands respectively and the minimum, maximum and average of education in terms of schooling year of the responding parents are 8, 17 and 14 respectively.


2.1.2.Frequency of items to find out Baby's' eating behavior: in order to assess the eating behavior of babies we have given 18 items of behavior as in the following table with 5 point Likert Scale 1=Never, 2=Rarely, 3=Sometimes, 4= often to 5=Always. The frequency of the variables is given in table 2.1.2

Table-2.1.2: Frequency of distribution baby's eating behavior

Sl.	Table-2.1.2. Trequency of distr	1=Ne	2=Rarely	3=Somet	4=	5=Al	Total
No.	Variables/ items of behavior	ver	2—Raiciy	imes	often	ways	Total
1	Baby seemed contented while feeding	, 61	2	20	8	10	40
2	Baby frequently wanted more milk than given	9	15	10	2	4	40
3	Baby loves milk	3	4	15	7	11	40
4	Baby had a big appetite	8	7	9	4		40
5	Baby finished feeding quickly	8	11	16	3	2	40
6	Baby became distressed while feeding	0	9	15	12	4	40
7	Baby got full up easily		6	15	11	8	40
8	If allowed to, my baby would take too much milk	18	11	5	6	0	40
9	Baby took more than 30 minutes to finish feeding	4	14	12	5	5	40
10	Baby got full before taking all the milk	1	4	17	12	4	40
11	Baby fed slowly	2	5	10	11	12	40
12	Even baby had just eaten he/she was happy to feed	20	10	7	3		40
	again if offered						
13	Baby found it difficult to manage a complete feed	2	15	12	4	7	40
14	Baby was always demanding a feed	12	4	13	9	2	40
15	Baby sucked more and more slowly during the	6	13	13	5	3	40
	course of a feed						
16	If given the chance, my baby would always be	12	10	8	6	4	40
	feeding						
17	Baby enjoyed feeding time	13	9	10	1	2	40
18	Baby could easily take a feed within 30 minutes of	10	18	8	4	0	40
	the last one						

Table-2.1.2 showed that highest frequency in the scales are never= 20, rarely=18, sometimes =20, often =12 and always =12. In percentage form the frequency is given following table:

Sl. No. of	1.Never	2. Rarely	3. Sometimes	4. often	5. Always	Total
items						
1	5%	50%	20%	25%	25%	100%
2	38%	25%	5%	10%	10%	100%
3	10%	38%	18%	28%	28%	100%
4	18%	23%	10%	0%	0%	100%
5	28%	40%	8%	5%	5%	100%
6	23%	38%	30%	10%	10%	100%
7	15%	38%	28%	20%	20%	100%
8	28%	13%	15%	0%	0%	100%
9	35%	30%	13%	13%	13%	100%
10	10%	43%	30%	10%	10%	100%
11	13%	25%	28%	30%	30%	100%
12	25%	18%	8%	0%	0%	100%
13	38%	30%	10%	18%	18%	100%
14	10%	33%	23%	5%	5%	100%
15	33%	33%	13%	8%	8%	100%
16	25%	20%	15%	10%	10%	100%
17	23%	25%	3%	5%	5%	100%
18	45%	20%	10%	0%	0%	100%
Average	23%	30%	16%	11%	11%	100%



In corresponding of table 2.1.2 we have determined the total point in terms of scaling point of eating behavior as in table 2.1.3.

Table-2.1.3: Total point of baby's eating behavior based on table 2.1.2

Sl. No. of	1.Never	2. Rarely	3. Sometimes	4. often	5. Always	Total
items						
1	0	4	60	40	50	154
2	9	30	30	16	20	105
3	3	8	45	44	55	155
4	8	14	27	0	0	49
5	8	22	48	8	10	96
6	0	18	45	16	20	99
7		12	45	32	40	129
8	18	22	15	0	0	55
9	4	28	36	20	25	113
10	1	8	51	16	20	96
11	2	10	30	48	60	150
12	20	20	21	0	0	61
13	2	30	36	28	35	131
14	12	8	39	8	10	77
15	6	26	39	12	15	98
16	12	20	24	16	20	92
17	13	18	30	8	10	79
18	10	36	24	0	0	70
	То	tal on avera	ige of 18 items			100.5
	overall A	verage scal	le of 40 respond	ents		2.51

Table 2.1.3 showed that the top five items are — my baby loved milk(155) following my baby seemed contented while feeding (154), my baby fed slowly(150), my baby found it difficult to manage a complete feed(131), my baby got full up easily(113) and my baby frequently wanted more milk than I provided(105). The average score of the items is 100.5 and the average score of the respondents is 2.51. It means the eating behavior lies between rarely and sometimes. Reliability of the questionnaire was checked by Cronbach's Alpha test—which is 0.710 that means there is consistency of the items or variables. The item-wise point is presented in the figure 2.1.3.





Figure-2.1.3: Item-wise point of baby's eating behavior.

2.1.3 Factors analysis of baby's eating behavior:

The researcher has identified some behavioral causes of bay's in eating by asking to the parents. There were eighteen variables or items—in the schedule and the respondents were asked to on 5-point scaling as never=1; rarely=2; sometimes =3 often=4 and always=5. The Principal Component Analysis (PCA) with Varimax Orthogonal rotation method explained 75.66% of total variation by the extracted factors with 0.615 Kaiser-Meyer-Olkin (KMO) measure of sampling adequacy (Appendix -1). The Bartlett's Test of Sphericity by Approx. Chi-Square 396.205 and significant level is 0.00. It means the item size is sufficient to explore the factors. Details of the factors or items are given in appendix table 1, 2 and 3 and from the PCA tables we identified six factors as shown in figure 2.1.3.1 based on Eigenvalue.

Figure-2.1.3.1: Scree plot of Factors



The component Transformation matrix is given in figure 2.1.3.1.

Table:2.1.3.1:Component Transformation Matrix									
Component	Component 1 2 3 4 5								
1	.560	.390	421	.147	451	364			
2	.114	.477	.266	.706	.244	.362			
3	357	.564	475	373	.018	.434			
4	.242	.423	.688	524	111	044			
5	.495	350	040	109	280	.735			
6	.493	002	226	234	.804	073			

Extraction Method: Principal Component Analysis.

Rotation Method: Varimax with Kaiser Normalization.

CD1 .	. •	•	•	•	. 11	^ 1		\sim
The component score	matrix	10	OIVED	1n	table	')	١ - ٢	٠,
The component score	mania	10	given	111	table	4.	ı .J.	

Table-2.1.3.2:Component Score Covariance Matrix										
Component	1	2	3	4	5	6				
1	1.000	.000	.000	.000	.000	.000				
2	.000	1.000	.000	.000	.000	.000				
3	.000	.000	1.000	.000	.000	.000				
4	.000	.000	.000	1.000	.000	.000				
5	.000	.000	.000	.000	1.000	.000				
6	.000	.000	.000	.000	.000	1.000				
Extraction Met	Extraction Method: Principal Component Analysis.									
Rotation Meth	od: Varimax	with Kaiser I	Normalizatio	n.						

The component Score Covariance Matrix shows that all the six items are significant as the p-value is 0.00 among each other. Therefore we have seen in the eating behavior of baby less than 4 years that there are six items of behavior as important statistically significant.

2.2. Analysis of Child eating behavior between 5-9 years: In this section an attempt is taken to analyze the child eating behavior between 5-9 years from different perspectives as follows:

2.2.1. Demographic profile and descriptive statistics of the respondents and the babies:

Table-2.1: Demographic profile and descriptive statistics of the respondents and the babies

Variable	Number	Variable	Minimum	Maximum			
			years	years	Mean	Std. Deviation	
Sex of respondent		Age of baby	5	9	6.98	1.527	
Father 20 Descriptive			tatistics of number	of Baby			
Mother	20	Total no. children	1	5	2.10	.955	
Total	40	4-9 years	1.0	2.0	1.200	.4051	
Descriptive statistics of parents occupation							
Occupation			Frequency	Percent	Valid %	Cumulative %	
Business			8	20	20	20	
Service			18	45	45	65	
	both servic	e and business	0	0	0	0	
Oth	ners including	housekeeping	14	35	35	100	
		Total	40	100.0	100.0		
Descriptive Statist	ics of Family	's annual income	and expenditure				
			Minimum	Maximum	Mean	Std. Dev.	
annual income(in	thousand tak	a)	240	1000	519.00	207.831	
annual income(in thousand taka)			180	900	394.75	161.959	
Descriptive Statist	ics of Educat	ion of Parents					
Fathers' Schoolin	g year	·	10	17	14.77	2.731	
Mothers Schoolin	ig year		8	17	12.50	2.873	

Table-2. 2.1 showed that out of 40 respondents 20 mother and 20 fathers. The age of the children varied between 4 year and 9.00 and on average 6.98 with standard deviation 1.57. The number of total children is fluctuated between 1 and 5 while number of children between 4-9 years is ranged from 1 to 2. The maximum occupation of the parents are service (18), following others including housekeeping (14), business (8) and both service and business (0). The minimum, maximum and average of annual income of the responding household are 240 thousands, 1000 thousands and 519 thousands respectively and the minimum, maximum and average of education in terms of schooling year of the responding father are 10, 17 and 14.77 respectively while mothers are 8, 17 and 12.50 respectively


2.2.2 Child's eating behavior: Childs' eating behavior is analyzed through the following sub sections using the five point scaling of behavior from Scale 1=Never, 2=Rarely, 3=Sometimes, 4= often to 5=Always. Frequency distribution of child's eating behavior: Childs' eating behavior is given in table -2.2.2.1:

Table-: 2. 2.2.1: Frequency distribution of child's eating behavior

S1.	Behavior items or variable	1=Never	2=Rarely	3=Som	4=	5=Alwa	Total
No				e-times	often	ys	
1	Child loves food	0	2	26	5	7	40
2	Child eats more when worried	8	22	9	1	0	40
3	Child has big appetite	9	6	15	6	4	40
4	Child finishes his/her meal quickly	10	11	5	10	4	40
5	Child is interested in food	6	16	11	7	0	40
6	Child is always asking for a drink	3	11	5	9	12	40
7	Child refuses new foods at first	10	8	10	3	9	40
8	Child eats slowly	15	3	2	2	18	40
9	Child eats less when angry	6	7	4	12	11	40
10	Child enjoys tasting new foods	8	17	9	3	3	40
11	Child eats less when s/he is tired	5	12	12	7	4	40
12	Child is always asking for food	3	2	15	10	10	40
13	Child eats more when annoyed	15	21	2	2	0	40
14	Allowed to, my child would eat too much	20	12	4	2	2	40
15	Child eats more when anxious	15	19	3	2	1	40
16	Child enjoys a wide variety of foods	1	3	18	9	9	40
17	Leaves food on his/her plate at the end of a meal	7	7	2	5	19	40
18	Child takes more than 30 minutes to finish a meal	8	5	7	14	6	40
19	The choice, child would eat most of the time	17	6	5	6	5	40
20	Child looks forward to mealtimes	10	8	17	2	3	40
21	Child gets full before his/her meal is finished	1	11	8	10	10	40
22	Child enjoys eating	0	2	22	8	7	40
23	Child eats more when s/he is happy	0	5	21	8	5	40
24	Child is difficult to please with meals	6	9	4	18	3	40
25	Child eats less when upset	10	5	8	10	9	40
26	Child gets full up easily	7	0	4	9	20	40
27	Child eats more when s/he has nothing else to do	14	10	10		6	40
28	Child is full finds room to eat his/her favorite food	7	8	2	4	19	40
29	If given the chance, child drink continuously the day	8	14	12	5	1	40
30	Child cannot eat a meal if has had a snack just before	6	1	6	12	15	40
31	The chance my child would always be having a drink	3	22	8	4	3	40
32	Interested in tasting food hasn't tasted before	4	3	13	4	16	40
33	Child that s/he doesn't like a food, without tasting	13	4	12	7	4	40
34	If given the chance, child have food in his mouth	12	10	6	0	12	40
35	My child eats more and more slowly during meal	6	9	10	7	8	40

The reliability analysis showed that Cronbach's Alpha is above 0.7 in the table of Reliability Statistics and according to Reliability Statistics if Cronbach's Alpha is above 0.7 then we can say that the data has good level of internal consistency. The table of Item Total statistics (Appendix-4) shows that all the items are important as deleting any of them reduces Cronbach's Alpha. The table of Scale Statistics shows the summary of Likert scale data that there was 35 items and the mean 67.775 and the variance is 158.025 with standard deviation is 12.5708. Table of Reliability Statistics showed that Cronbach's Alpha is 0.769. The table -2.2.2.1 showed that in the different highest frequency are never=20, rarely=19, sometimes =26, often= and always=20. The percentage distribution is given in the following table

SI. of Behavior items or variable.	1=Never	2=Rarely	3=Some-times	4= often	5=Always	Total
1	0%	5%	65%	13%	18%	100
2	20%	55%	23%	3%	0%	100
3	23%	15%	38%	15%	10%	100
4	25%	28%	13%	25%	10%	100
5	15%	40%	28%	18%	0%	100
6	8%	28%	13%	23%	30%	100
7	25%	20%	25%	8%	23%	100
8	38%	8%	5%	5%	45%	100
9	15%	18%	10%	30%	28%	100
10	20%	43%	23%	8%	8%	100
11	13%	30%	30%	18%	10%	100
12	8%	5%	38%	25%	25%	100
13	38%	53%	5%	5%	0%	100
14	50%	30%	10%	5%	5%	100
15	38%	48%	8%	5%	3%	100
16	3%	8%	45%	23%	23%	100
17	18%	18%	5%	13%	48%	100
18	20%	13%	18%	35%	15%	100
19	43%	15%	13%	15%	13%	100
20	25%	20%	43%	5%	8%	100
21	3%	28%	20%	25%	25%	100
22	0%	5%	55%	20%	18%	100
23	0%	13%	53%	20%	13%	100
24	15%	23%	10%	45%	8%	100
25	25%	13%	20%	25%	23%	100
26	18%	0%	10%	23%	50%	100
27	35%	25%	25%	0%	15%	100
28	18%	20%	5%	10%	48%	100
29	20%	35%	30%	13%	3%	100
30	15%	3%	15%	30%	38%	100
31	8%	55%	20%	10%	8%	100
32	10%	8%	33%	10%	40%	100
33	33%	10%	30%	18%	10%	100
34	30%	25%	15%	0%	30%	100
35	15%	23%	25%	18%	20%	100
Average	20%	22%	23%	16%	19%	100



The frequency of table 2.2.1 is converted into five point scale is presented in corresponding to the scale in table 2.2.2.2 below:

Table-2.2.2.2: Point of Eating behavior in in corresponding to the scale in table 2.2.2.1

Sl.No.	1=Never	2=Rarely	3=Some-times	4= often	5=Always	Total	Position
1	0	4	78	20	35	137	Fifth
2	8	44	27	4	0	83	
3	9	12	45	24	20	110	
4	10	22	15	40	20	107	
5	6	32	33	28	0	99	
6	3	22	15	36	60	136	Sixth
7	10	16	30	12	45	113	
8	15	6	6	8	90	125	
9	6	14	12	48	55	135	Seventh
10	8	34	27	12	15	96	
11	5	24	36	28	20	113	
12	3	4	45	40	50	142	Fourth
13	15	42	6	8	0	71	
14	20	24	12	8	10	74	
15	15	38	9	8	5	75	
16	1	6	54	36	45	142	Fourth
17	7	14	6	20	95	142	Fourth
18	8	10	21	56	30	125	
19	17	12	15	24	25	93	
20	10	16	51	8	15	100	
21	1	22	24	40	50	137	Fifth
22	0	4	66	32	35	137	Fifth
23	0	10	63	32	25	130	
24	6	18	12	72	15	123	
25	10	10	24	40	45	129	
26	7	0	12	36	110	165	First
27	14	20	30	0	30	94	
28	7	16	6	16	95	140	
29	8	28	36	20	5	97	
30	6	2	18	48	75	149	Second
31	3	44	24	16	15	102	
32	4	6	39	16	80	145	Third
33	13	8	36	28	20	105	
34	12	20	18	0	60	110	
35	0	4	78	20	35	137	
Average of	of items					117.66	
	of respondents					3.36	

The table 2.2.2.2 depicted the total point of each variable and the top seven items are as follows-My child gets full up easily(165), Child cannot eat a meal if s/he has had a snack just before(149), My child is interested in tasting food s/he hasn't tasted before(145), My child is always asking for food(142), My child enjoys a wide variety of foods(142), My child leaves food on his/her plate at the end of a meal(137), My child loves food(137), My child is always asking for a drink(136), My child eats less when angry(135). The average of items is 117.66 and the average of respondents is 3.36. That means the eating behavior of children varied between sometimes and often. The table 2.2.2.2 depicted the total point of each variable and the top seven items correspondingly based on the total point in figure 2.2.2.2 the total points are presented.




Figure-2.2.2.2: Total point of Child's eating behavior.

2.2.2.3. Factor Analysis: The Table of Scale Statistics shows that there are 35 items or variables of Childs 'eating behavior and to reduce the variables factor analysis is done with Principal Component Analysis with Varimax rotation. The communalities of the variables varied between 0.791 and 0.970 (Appendix -5). The total variance explained by the PCA is 88.388% that means the PCA is accepted and based on Eigen values ten items are considered important(Appendix-5). The Scree plot based on Eigen values is given in figure-2.2. 2.3




Figure-2.2.2.3: The Scree plot based on Eigen values of Child's Eating Behavior

Therefore the 35 variables are reduced to ten components explaining more than 88% of total variance. As there are more than 25 items the rotted component matrix is impossible to do.

2.3. Analysis of Adolescent (10-19) Food Habits: The food habits of adolescent are discussed below:

2.3.1. Socioeconomic profile of the respondents: The socioeconomic profile of the respondents are discussed through table 2.3.1.1 to 2.3.1.7:

Table-3.3.1.1: Respondents by Sex

Sex	Frequency	Valid Percent	Cumulative Percent
Male	17	50.0	50.0
Female	17	50.0	100.0
Total	34	100.0	

Table-2.3.1.1 showed that 50% respondents is male and 50% female i.e. the respondents by sex are equal .

Table-2.3.1.2: Education status of the respondents:

Schooling Year	Frequency	Valid Percent	Cumulative Percent
5.0	2	5.9	5.9
7.0	1	2.9	8.8
8.0	3	8.8	17.6
9.0	5	14.7	32.4
10.0	2	5.9	38.2
11.0	5	14.7	52.9
12.0	11	32.4	85.3
13.0	4	11.8	97.1
15.0	1	2.9	100.0
Total	34	100.0	

Table 2.3.1.2 depicted that education status of the respondents in terms of schooling year is classified from 5 years to 15 years and showed that 12 years education is 11(32.4%) following 9 years and 11 years each is 5(14.7%).

2.3.1.3. Mother's Education: Education statuses of the mother of the respondents are given in table 3.1.3:

Table-2.3.1.3: Mother's Education

Schooling Year	Frequency	Valid percent	Cumulative Percent
5.0	1	2.9	2.9
8.0	5	14.7	17.6
10.0	7	20.6	38.2
11.0	3	8.8	47.1
12.0	3	8.8	55.9
13.0	1	2.9	58.8
14.0	1	2.9	61.8
16.0	6	17.6	79.4
18.0	7	20.6	100.0
Total	34	100.0	

Table-2.3.1.3 education status of respondent's mother in terms of schooling year varied between 5 years and 18 years and maximum of them is 7 years (20.6) of schooling.

Table-2.3.1.4: Mother's Occupation

Twell Zieviv iv Ivietner a detupunen								
Occupation	Frequency	Valid Percent	Cumulative Percent					
service	1	2.9	2.9					
housekeeping	24	70.6	73.5					
Both of the above	8	23.5	97.1					
Business	1	2.9	100.0					
Total	34	100.0						

Table-2. 3.1.4 showed that lion portion (70.6%) of mother's occupation is housekeeping following both service and housekeeping(23.5%).

Table- 2. 3.1.5: Father's Education

Schooling Year	Frequency	Valid Percent	Cumulative Percent
1.0	1	2.9	2.9
10.0	5	14.7	17.6
11.0	1	2.9	20.6
12.0	8	23.5	44.1
13.0	1	2.9	47.1
15.0	1	2.9	50.0
16.0	5	14.7	64.7
17.0	2	5.9	70.6
18.0	10	29.4	100.0
Total	34	100.0	

Table-2.3.1.5 education status of respondent's father in terms of schooling year varied between 1 years and 18 years and maximum of them is 18 years (29.6%) of schooling following 12 years (23.5%), 16 years (14.7%).

Table-2.3.1.6: Father's Occupation

Occupation	Frequency	Valid Percent	Cumulative Percent
service	12	35.3	35.3
Business	13	38.2	73.5
Both	2	5.9	79.4
others	7	20.6	100.0
Total	34	100.0	

Table -2.3.1.6 showed that highest number (38.2%) of father's occupation is business following service(35.3%) and others (20.6%).

Table-2.3.1.7: Descriptive statistics annual income and expenditure of the respondent's family

Variables	Minimum	Maximum	Mean	Std. Deviation
annual income(Taka in thousand)	100	1450	479.12	301.507
annual expenditure(Taka in thousand)	95	1050	393.97	255.494

Table 2.3.1.7 showed that annual income is varied between 100 thousand and 1450 and on average 479.12 while annual expenditure fluctuated between 95 and 1050 with an average 393.97 thousand.

2.3.2. Descriptive Statistics of Adolescent Food Habits: The food habits of adolescent is measured in terms of given items or variables with three point scale **as Most** like=2; Neutral=0; and like=1. The descriptive statistics of adolescents' food habits is given in the following tables

2.3.2.1: Frequency Distribution of food adolescent food habits: Frequency Distribution of food adolescent food habits is given in table 2.3.2.1

Table-2.3.2.1: Frequency Distribution of food adolescent food habits

sl.no	Items or variables	Most	like=1	Neutral=	Total
		like=2		0	
1	If I having lunch away from home, I often choose a low-fat option.	11	20	3	34
2	I usually avoid eating fried foods.	19	15	0	34
3	I usually eat a dessert or pudding if there is one available.	5	29	0	34
4	I make sure I eat at least one serving of fruit a day.	15	19	0	34
5	I try to keep my overall fat intake down.	11	23		34
6	If I am buying crisps, I often choose a low-fat brand.	12	11	11	34
7	I avoid eating lots of sausages and burgers.	13	20	1	34
8	I often buy pastries or cakes.	11	20	3	34
9	I try to keep my overall sugar intake down.	12	12	10	34
10	I make sure I eat at least one serving of vegetables or salad a day.	12	20	2	34
11	If I am having a dessert at home, I try to have something low in fat.	10	18	6	34
12	I rarely takeaway meals.	6	28	0	34
13	I try to ensure that I eat plenty of fruit and vegetables.	9	25	0	34
14	I often eat sweet snacks between meals.	17	17	0	34
15	I usually eat at least one serving of vegetables or salad	13	21	0	34
16	When I am buying a soft drink, I usually choose a diet drink.	26	8	0	34
17	When I put butter or margarine on bread, I usually spread it on thinly.	15	13	6	34
18	If I have a packed lunch, I usually include some chocolate and/or biscuits.	13	17	4	34
19	When I have a snack between meals, I often choose fruit.	9	20	5	34
20	If I am having a dessert or pudding in a restaurant, I usually choose the healthiest one.	12	18	4	34
21	I often have cream on desserts.	12	16	6	34
22	I eat at least servings of fruits most days.	23	11	0	34
23	I generally try to have a healthy diet.	7	27	0	34

Table-2.3.2.1 showed that the highest frequency in the scale most like =26, like=29 and netral=11. Reliability of the questionnaire was checked by Cronbach's Alpha test which is 0.8230 that means there is consistency of the items or variables. The percentage distribution of frequency is given below

sl.no of Items or variables	Most like=2	like=1	Neutral=0	Total	Average f	igure		
1	32%	59%	9%	100	C00/			
2	56%	44%	0%	100	60%		55%	1
3	15%	85%	0%	100				ı
4	44%	56%	0%	100	50% -			
5	32%	68%	0%	100] 30%]			
6	35%	32%	31%	100				
7	38%	59%	3%	100	40% -	37%		
8	32%	59%	9%	100] 40%			
9	35%	35%	29%	100				
10	35%	59%	6%	100	30% -			
11	29%	53%	17%	100				
12	18%	82%	0%	100				
13	26%	74%	0%	100	20% -			
14	50%	50%	0%	100]			
15	38%	62%	0%	100]			
16	76%	24%	0%	100	10% -			8%
17	44%	38%	17%	100				
18	38%	50%	11%	100				
19	26%	59%	14%	100	0% -			
20	35%	53%	11%	100]	1	2	3
21	35%	47%	17%	100		1	_	3
22	68%	32%	0%	100				
23	21%	79%	0%	100				
Average	37%	55%	8%	100				


The item-wise point is presented in table 2.3.2.2.

Table-2.3.2.2: Points of food habits of adolescent based on table 2.3.2.1


Sl. no. of variables/items	Most like=2	like=1	Neutral=0	Total point	position
1	22	20	0	42	
2	38	15	0	53	Third
3	10	29	0	39	
4	30	19	0	49	Fifth
5	22	23	0	45	
6	24	11	0	35	
7	26	20	0	46	Sixth
8	22	20	0	42	
9	12	12	0	24	
10	24	20	0	44	
11	20	18	0	38	
12	12	28	0	40	
13	18	25	0	43	
14	34	17	0	51	Fourth
15	26	21	0	47	
16	52	8	0	60	First
17	30	13	0	43	Seventh
18	26	17	0	43	Seventh
19	18	20	0	38	
20	24	18	0	42	
21	24	16	0	40	
22	46	11	0	57	second
23	14	27	0	41	
	Average of iter	ns		43.56	
	Average of respor			1.89	

The top seven total scoring items are: (i)When I am buying a soft drink, I usually choose a diet drink (60), (ii)I eat at least servings of fruits most days(57),(iii)I usually avoid eating fried foods(53) (iv)I often eat sweet snacks between meals(51),(v) I make sure I eat at least one serving of fruit a day (49)(vi) I avoid eating lots of sausages and burgers.(46) and (vii) When I put butter or margarine on bread and I usually spread it on thinly If I have a packed lunch, I usually include some chocolate and/or biscuits (43). Average of score of the items is 43.56 and the average score of respondents is 1.89. The data of table 2.3.2.2 is presented in figure 2.3.3.2

Figure 2.3.3.2: Points of food habits of adolescent



2.3.3.3 Factor Analysis: The Table of Scale Statistics shows that there are 23 items or variables of adolescent 'eating behavior and to reduce the variables factor analysis is done with Principal Component Analysis with Varimax rotation. The Bartlett's Test of Sphericity by Approx. Chi-Square 295.205 and significant level is 0.00. It means the item size is sufficient to explore the factors. The communalities of the variables varied between 0.57 and 0.903 (Appendix -6). The total variance explained by the PCA is 79.278% that means the PCA is accepted and based on Eigen values eight items are considered important(Appendix-7). The Scree plot based on Eigen values is given in figure 2.3.3.3.



The component transformation matrix is given in the following table

Component Transformation Matrix									
Component	1	2	3	4	5	6	7	8	
1	.510	.325	.488	.428	.272	.358	098	.025	
2	.486	788	.014	007	.177	043	.310	.116	
3	547	303	099	.217	.522	.512	097	088	
4	025	.150	.280	644	.620	219	072	.218	
5	.116	.347	623	.223	.313	058	.391	.417	
6	213	.145	.329	153	115	.235	.842	159	
7	327	130	.371	.238	217	101	014	.790	
8	.195	.015	198	475	283	.701	134	.328	

Therefore the PCA reduces 21 items to eight factors of food habits of adolescent.

2.4. Parental Feeding Style to his Children up to 9 years: The parental feeding style to his children up to 9 years is analyzed below:

2.4.1. Demographic profile and descriptive statistics of the respondents and the babies:

Table-2.4.1: Demographic profile and descriptive statistics of the respondents and the babies

Variable	Number	Variable	Minimum in	Maximum	Mean	Std. Deviation
Sex of responden	t	Age of respondents	years 28	years 59	45.32	8.562
Father	17	Descriptive Statistic	cs of number of Baby		•	
Mother	17	Total no. children	1	5	1.82	1.141
Total	34	9 years	1.0	3.0	1.60	.731
		Descript	tive statistics of parents	s occupation		
	Occupation		Frequency	Percent	Valid Percent	Cumulative Percent
		Business	4	12	12	12
		Service	16	47	47	59
	both	service and business	2	5	5	64
	Others inc	cluding housekeeping	12	36	36	100
		Total	40	100.0	100.0	
Descriptive Stat	istics of Family's an	nual income and expen	ıditure			
			Minimum	Maximum	Mean	Std. Dev.
annual income(in thousand taka)		280	1200	589.00	277.831
annual income(in thousand taka)			246	980	594.75	261.959
Descriptive Stat	istics of Education	of Parents				
Fathers' Schooli	ing year		10	19	14.50	2.631
Mothers School	ing year		8	18	12	2.773

Table 2.4.1 showed that out of 34 respondents 17 mother and 17 fathers. The age of the children varied between 4 year and 9.00 and on average 6.98 with standard deviation 1.57. The number of total children is fluctuated between 1 and 5 while number of children up to 9 years is ranged from 1 to 3. The maximum occupation of the parents are service (16), following others including housekeeping (12), business (4) and both service and business (2). The minimum, maximum and average of annual income of the responding household are 280 thousands, 1200 thousands and 589 thousands respectively and the minimum, maximum and average of annual expenditure are 246, 980 and 594.75 thousands. The minimum, maximum and average education in terms of schooling year of the responding father are 10, 19 and 14.5 respectively while mother's are 8, 18 and 12 respectively.


2.4.2. Descriptive statistics of parental feeding style to his/her children up to 9 years

2.4.2.1: Frequency distribution of feeding style: The respondents were given a scaling questionnaire on five point scale and the result is given in table 4.2.1

Table-2.4.2.1: Frequency Distribution of parental feeding style to his/her children up to 9 years

Sl.no.	Items /variable	1.Never	2.Rarely	3.Sometime	4.often	5.Always 13 15 14 16 4 17 9 8 8 16 4 19 6 7 12 10 9 6 19 8	Total
				S			
1	I allow child to choose which foods to have for meals	3	1	15	2		34
2	I give child something to eat to make him/her feel better when s/he	5	2	9	6	15	34
	is feeling upset						
3	I encourage my child to look forward to the meal	1	1	11	7	1	34
4	I praise my child if s/he eats what I give him/her	3	1	7	7	16	34
5	I decide how many snacks my child should have	6	4	12	8		34
6	I encourage my child to eat a wide variety of foods	1	2	6	8	17	34
7	In order to get my child to behave him/herself I promise him/her something to eat	6	4	13	2	9	34
8	I present food in an attractive way to my child	7	4	11	4	8	34
9	If child misbehaves I withhold his/her favorite food	7	7	10	2	8	34
10	I encourage my child to taste each of the foods I serve at mealtimes	1	3	11	3	16	34
11	I allow my child to wander around during a meal	14	4	9	3	4	34
12	I encourage my child to try foods that s/he hasn't tasted before	2	0	10	3	19	34
13	I give my child something to eat to make him/her feel better when s/he has been hurt	5	3	11	5	6	34
14	I let my child decide when s/he would like to have her meal	8	1	13	5	7	34
15	I give my child something to eat if s/he is feeling bored	4	3	9	6	12	34
16	I allow my child to decide when s/he has had enough snacks to eat	3	4	13	4	10	34
17	I decide when it is time for my child to have a snack	7	1	14	3	9	34
18	I use puddings as a bribe to get my child to eat his/her main course	10	17	10	1	6	34
19	I encourage my child to enjoy his/her food	1	3	8	3	19	34
20	I decide the times when my child eats his/her meals	6	3	12	5	8	34
21	I give my child something to eat if s/he is feeling worried	7	3	12	5	7	34
22	I reward my child with something to eat when s/he is well behaved	6	2	16	4	6	34
23	I let my child eat between meals whenever s/he wants	3	3	10	6	12	34
24	I insist my child eats meals at the table	7	2	10	5	10	34
25	I give my child something to eat to make him/her feel better when	7	4	13	1	9	34
	s/he is feeling angry						
26	I decide what my child eats between meals	9	2	12	7	4	34
27	I praise my child if s/he eats a new food	1	4	12	6	11	34

Table-2.4.2.1 depicted that in the five point scale the highest frequency in the never (14), rarely (17) sometimes (16) ,often (8) and always(19). Reliability of the questionnaire was checked by Cronbach's Alpha test which is 0.931 that means there is consistency of the items or variables. The percentage of frequency distribution is given below:




The total point corresponding to the frequency table 2.4.2.1 is given in table 2.4.2.2.

Table-2.4.2.2: Total Scaling Point of parental feeding style to his/her children up to 9 years based on table-2.4.2.1

Variables no.	1.Never	2.Rarely	3.Sometimes	4.Often	5.Always	Total	position
1	3	2	45	8	65	123	Eight
2	5	4	27	24	75	135	Fourth
3	1	2	33	28	70	134	Fifth
4	3	2	21	28	80	134	Fifth
5	6	8	36	32	20	102	
6	1	4	18	32	85	140	First
7	6	8	39	8	45	106	
8	7	8	33	16	40	104	
9	7	14	30	8	40	99	
10	1	6	33	12	80	132	sixth
11	14	8	27	12	20	81	
12	2	0	30	12	95	139	Second
13	5	6	33	20	30	94	
14	8	2	39	20	35	104	
15	4	6	27	24	60	121	
16	3	8	39	16	50	116	
17	7	2	42	12	45	108	
18	10	34	30	4	30	108	
19	1	6	24	12	95	138	Second
20	6	6	36	20	40	108	
21	7	6	36	20	35	104	
22	6	4	48	16	30	104	
23	3	6	30	24	60	123	Eight
24	7	2	10	5	10	111	
25	7	4	13	1	9	103	
26	9	2	12	7	4	97	
27	1	4	12	6	11	124	Seventh
		Av	erage of items		•	114.51	
		Avera	ge of respondents			4.24	

The total point score of table 2.4.2.2. is given in figure 2. 4.2.2 for easy understanding

Figure-2. 4.2.2: The total point score of table 2.4.2.2.



2.4.3. Factor Analysis: The Table of Scale Statistics shows that there are 27 items or variables of parent's 'eating behavior to their children and to reduce the variables factor analysis is done with Principal Component Analysis with Varimax rotation. Kaiser-Meyer-Olkin Measure of Sampling Adequacy is .670. The Bartlett's Test of Sphericity by Approx. Chi-Square 806.205 and significant level is 0.00. It means the item size is sufficient to explore the factors. The communalities of the variables varied between 0.51 and 0.893 (Appendix -9). The total variance explained by the PCA is 78.96% that means the PCA is accepted and based on Eigen values seven items are considered important(Appendix-10). The Scree plot based on Eigen values is given in figure-2. 4.3.1.




Figure-2.4.3.1

The component matrix is given in appendix-11 and 12. The component score Coefficient Matrix is given in appendix 13. The Component Score Covariance Matrix is given in table 2. 4.3.3

Table - 2. 4.3.3: Component Score Covariance Matrix

Component	1	2	3	4	5	6	7			
1	1.000	.000	.000	.000	.000	.000	.000			
2	.000	1.000	.000	.000	.000	.000	.000			
3	.000	.000	1.000	.000	.000	.000	.000			
4	.000	.000	.000	1.000	.000	.000	.000			
5	.000	.000	.000	.000	1.000	.000	.000			
6	.000	.000	.000	.000	.000	1.000	.000			
7	.000	.000	.000	.000	.000	.000	1.000			
Extraction Metho	Extraction Method: Principal Component Analysis. Rotation Method: Varimax with Kaiser Normalization.									

Therefore PCA produced seven factors from 27 through extraction of parent's eating behavior to their children .

2.5. Food Preference For Adults (20-29) Unmarried Male And Female:

2.5.1: Socioeconomic status of the respondents: The socioeconomic status of the respondents food preferences for adults (20-29) unmarried male and female are discussed through table 2.5.1 to 2.5.6 as given below:

Table-2.5.1: Demographic profile and descriptive statistics of the respondents and the babies

Variable	Number
Sex of respondent	
Male	20
Female	20
Total	40
Marital status	
Unmarried	40
Total	40

Table-2.5.1 depicted that there were 20 male and 20 female respondents and all them were unmarried of 20-29 aged.

Table -2. 5.2: Age Structure of the respondents:

Age	Frequency	Valid Percent	Cumulative Percent
21	13	31.7	31.7
22	12	29.3	61.0
23	7	17.1	78.0
24	4	9.8	87.8
25	3	7.3	95.1
26	1	2.4	97.6
Total	40	100.0	100.00

Table-2.5.2 showed that age structure of the respondents varied between 21 and 26 and 13(31.7%) of respondents have 21 age following 12 (29.3%), 7(17.1%), 4(9.8%) and 3(7.3%) and only 1(2.4) .Table-2.5.3: Occupation of the respondents

occupation	Frequency	Percent	Valid Percent	Cumulative Percent
Service	3	7.3	7.3	7.3
Business	1	2.4	2.4	9.8
Unemployed	4	9.8	9.8	19.5
Others(including Students)	32	78.0	78.0	97.6
Total	40	100.0	100.0	100.00

Table-2.5.3 showed that occupation of most of the respondents is others including students32(78%) following unemployed 4(9.8%) service3(7.3%) and business 1(2.4%).

Table-2.5.4. Education status of the respondents

Schooling year	Frequency	Valid Percent	Cumulative Percent
12	1	2.4	2.4
14	2	4.9	7.3
15	22	53.7	61.0
16	4	9.8	70.7
17	4	9.8	80.5
18	1	2.4	82.9
19	2	4.9	87.8
20	3	7.3	95.1
9	1	2.4	97.6
Total	40	100.0	

Table-2.5.4 showed that occupation of the respondents varied between 9 and 20 and 22(53.7%) of respondents have 15 years education following 16 years of 4(9.8%), 17 years of 4(9.8%) and 20 years 3(7.3%).

Table 2.5.5.Annual income of respondents

income (in thousand)	Frequency	Valid Percent	Cumulative Percent
100	34	82.9	82.9
200	1	2.4	85.4
250	1	2.4	87.8
36	1	2.4	90.2
48	1	2.4	92.7
60	2	4.9	97.6
Total	40	100.0	

Table-2.5.5 showed that annual income of most of the respondents 34(82.9%) is 100 thousand following 60 thousands 2(4.9%) and 200 thousand, 250 thousand 36 thousand and 48 thousand 1(2.4%) each. In table 2.5.6 annual expenditure of respondents is presented and it showed that annual expenditure of the respondents varied between 96 thousands and 200 thousands.

Table-2.5.6: Annual expenditure of respondents

income (in thousand)	Frequency	Valid Percent	Cumulative Percent
100	3	7.3	7.3
120	1	2.4	9.8
150	1	2.4	12.2
200	1	2.4	14.6
48	1	2.4	17.1
60	3	7.3	24.4
70	2	4.9	29.3
72	6	14.6	43.9
78	2	4.9	48.8
80	6	14.6	63.4
82	2	4.9	68.3
84	5	12.2	80.5
85	2	4.9	85.4
90	1	2.4	87.8
94	2	4.9	92.7
95	1	2.4	95.1
96	1	2.4	97.6
Total	40	100.0	100

2.5.2. Descriptive Statistics of food preference: Table-2.5.2.1 depicted that in the five point scale the highest frequency in the dislike a lot (39), dislike a little (12) neither like or dislike (26), like a little (31) and like a lot (34). Reliability of the questionnaire was checked by Cronbach's Alpha test—which is 0.851 that means there is consistency of the items or variables. The total point corresponding to the frequency table 2.5.2.1 is given in table 2.5.2.2

Table-2.5.2.1: Frequency distribution in percentage of food preference of unmarried

Sl.No.	Food Item	1- Dislike	2- Dislike a	3- Neither like or	4- Like a little	5- Like a lot	Total
31.110.	rood item	a lot	little	dislike	4- Like a little	J- LIKE a lot	Total
1	Beef	0%	0%	10%	78%	13%	100
2	Beef burgers	3%	0%	18%	58%	23%	100
3	Lamb	8%	8%	10%	5%	70%	100
4	Chicken	3%	0%	10%	20%	68%	100
5	Bacon	15%	20%	65%	0%	0%	100
6	Ham	70%	25%	5%	0%	0%	100
7	Sausage(Meat)	33%	5%	5%	23%	35%	100
8	White fish	40%	5%	10%	23%	23%	100
9	Oily fish	35%	8%	3%	23%	33%	100
10	Smoked Salmon	70%	8%	23%	0%	0%	100
11	Tinned Tuna	33%	8%	18%	43%	0%	100
12	Eggs	5%	8%	28%	60%	0%	100
13	Baked beans	5%	18%	13%	63%	5%	100
14	Bread or Bread rolls	3%	5%	33%	60%	0%	100
15	Bran cereal	25%	10%	23%	20%	23%	100
16	Porridge	30%	10%	15%	43%	3%	100
17	Plain boiled rice	3%	5%	13%	78%	3%	100
18	Sugared cereal	40%	30%	20%	10%	0%	100
19	Hummus	98%	0%	3%	0%	0%	100
20	Wheat cereal	13%	8%	35%	20%	25%	100
21	Potatoes (boiled or mashed)	0%	8%	93%	0%	0%	100
22	Chips	3%	0%	3%	15%	80%	100
23	Rice or corn cereal	8%	0%	20%	30%	43%	100
24	Soft cheese	10%	8%	35%	48%	0%	100
25	Hard cheese	18%	5%	35%	43%	0%	100
26	Cottage Cheese	13%	3%	8%	30%	40%	100
27	Plain, Low-fat yoghurt	15%	3%	13%	70%	0%	100
28	Oranges	0%	3%	3%	3%	93%	100
29	Grapes	0%	0%	10%	5%	85%	100

					Table cont	inued	
30	Apples	0%	13%	8%	20%	60%	100
31	Melon	8%	3%	13%	78%	0%	100
32	Peaches	70%	13%	8%	8%	0%	100
33	Apricots	78%	3%	3%	5%	13%	100
34	Strawberries	28%	3%	30%	10%	30%	100
35	Avocadoes	70%	3%	13%	15%	0%	100
36	Spinach	20%	3%	15%	8%	55%	100
37	Carrots	3%	13%	15%	70%	0%	100
38	Green beans	25%	30%	30%	15%	0%	100
39	Cucumber	5%	8%	10%	13%	65%	100
40	Celery	3%	5%	8%	10%	75%	100
41	Mushrooms	18%	3%	8%	23%	50%	100
42	Parsnips	30%	8%	15%	20%	28%	100
43	Peas	23%	18%	15%	25%	20%	100
44	Sweet corn	3%	15%	15%	68%	0%	100
45	Broccoli	30%	23%	43%	5%	0%	100
46	Salad leaves	38%	3%	25%	18%	18%	100
47	Red peppers	8%	10%	20%	63%	0%	100
48	Raw tomatoes	15%	10%	13%	23%	53%	100
49	Beetroot	20%	18%	10%	53%	0%	100
50	Brussels sprouts	58%	5%	23%	15%	0%	100
51	Butter	45%	15%	13%	28%	0%	100
52	Butter-like spreads	8%	3%	13%	78%	0%	100
53	Cream	70%	5%	13%	13%	0%	100
54	Mayonnaise	25%	5%	5%	65%	0%	100
55	Plain biscuits	3%	5%	35%	58%	0%	100
56	Chocolate biscuits	5%	5%	8%	83%	0%	100
57	Cake	0%	0%	8%	20%	73%	100
58	Ice cream	0%	3%	3%	10%	85%	100
59	Custard	13%	3%	3%	83%	0%	100
60	Chocolate	0%	0%	3%	10%	88%	100
61	Crisps	0%	0%	0%	5%	95%	100
62	Chewy gummy sweets	10%	23%	20%	48%	0%	100
	Average	21%	8%	17%	31%	24%	100





Table-2.5.2.2: Total Scaling Point of food preference un married between 20-29 years based on table-2.5.2.2

Variable No.	1- Dislike a lot	2- Dislike a little	3- Neither like or dislike	4- Like a little	5- Like a lot	Total	Position
1	0	0	12	124	25	161	
2	1	0	21	92	45	159	
3	3	6	12	8	140	169	
4	1	0	12	32	135	180	seventh
5	6	16	78	0	0	100	
6	28	20	6	0	0	54	
7	13	4	6	36	70	129	
8	16	4	12	36	45	113	
9	14	6	3	36	65	124	
10	28	6	27	0	0	61	
11	13	6	21	68	0	108	
12	2	6	33	96	0	137	
13	2	14	15	100	10	141	
14	1	4	39	96	0	140	
15	10	8	27	32	45	122	
16	12	8	18	68	5	111	
17	1	4	15	124	5	149	
18	16	24	24	16	0	80	
19	39	0	3	0	0	42	
20	5	6	42	32	50	135	
21	0	6	111	0	0	117	
22	1	0	3	24	160	188	Fifth
23	3	0	24	48	85	160	
24	4	6	42	76	0	128	
25	7	4	42	68	0	121	
26	5	2	9	48	80	144	
27	6	2	15	112	0	135	
28	0	2	3	4	185	194	second
29	0	0	12	8	170	190	fourth
30	0	10	9	32	120	171	Eighth
31	3	2	15	124	0	144	
32	28	10	9	12	0	59	
33	31	2	3	8	25	69	
34	11	2	36	16	60	125	
35	28	2	15	24	0	69	
36	8	2	18	12	110	150	
37	1	10	18	112	0	141	
38	10	24	36	24	0	94	
39	2	6	12	20	130	170	
40	1	4	9	16	150	180	seventh
41	7	2	9	36	100	154	
42	12	6	18	32	55	123	
43	9	14	18	40	40	121	
44	1	12	18	108	0	139	
45	12	18	51	8	0	89	
46	15	2	30	28	35	110	
47	3	8	24	100	0	135	
48	6	8	15	36	105	170	
49	8	14	12	84	0	118	
50	23	4	27	24	0	78	
51	18	12	15	44	0	89	
52	3	2	15	124	0	144	
53	28	4	15	20	0	67	
54	10	4	6	104	0	124	
55	1	4	42	92	0	139	
56	2	4	9	132	0	147	
57	0	0	9	32	145	186	sixtl
58	0	2	3	16	170	191	Third
59	5	2	3	132	0	142	
60	0	0	3	16	175	194	Second
61	0	0	0	8	190	198	Firs
62	4	18	24	76	0	122	
			ore of items		•	131.35	
			of respondents			2.11	

The total point of each variable or item of food preference is given in figure -2.5.2 .1 Figure-2.5.2.1: total point of each variable or item of food preference



From the above analysis we can finalize the top ten preferred food items as given in table 2.5.2.3

Table-2.5.2.3: top ten preferred food items

Serial No.	Food items	Total point	Position
1	Crisps	198	First
2	Chocolate	194	Second
3	Oranges	194	second
4	Ice cream	191	Third
5	Grapes	190	fourth
6	Chips	188	Fifth
7	cake	186	sixth
8	Celery	180	seventh
9	Chicken	180	seventh
10	Apples	171	Eighth

2.6. Food Preference for Adults (29-64) Married Male and Female:

2.6.1: Socioeconomic Profile of Respondents: Socioeconomic profiles of respondents are discussed through table 2.6.1.1 to 2.6.1.5 below:

Table-2.6.1.1: Respondents by sex

Variable	Number
Sex of respondent	
Male	20
Female	20
Total	40
	Marital status
married	40
Total	40

Table-2.6.1.1 depicted that there were 20 male and 20 female respondents and all them were married of 29 -64aged.

Table: 2.6.1.2: age, annual income and expenditure of the respondents

Variables	Minimum	Maximum	Mean	Std. Deviation
age	24.0	59.0	42.425	9.2733
annual income(000)	120	6000	574.15	902.991
annual expenditure(000)	120	1200	419.20	221.168

Table-2.6.1.2 showed that age ranged from 24 to 59 with an average 42 and standard deviation 9. The annual income varied between 120 thousands and 6000 thousand with an average 574 and standard deviation 902 while annual expenditure fluctuated between 120 thousands and 1200 thousands with an average 419 and standard deviation 221.

Table-2.6.1.3: Occupation of respondents

Occupation	Frequency	Valid Percent	Cumulative Percent
Service	8	20.0	20.0
Business	26	65.0	85.0
Both of them	4	10.0	95.0
Others	2	5.0	100.0
Total	40	100.0	

Table -2.6.1.3 showed that the occupation of lion portion of respondents is business 26(65%) following service 8(20%) and both business and service 4(10%).

Table-2.6.1.4: Occupation of spouse

Occupation	Frequency	Valid Percent	Cumulative Percent
Household	23	57.5	57.5
Service	12	30.0	87.5
Both service and Household	4	10.0	97.5
Business	1	2.5	100.0
Total	40	100.0	

Table-2. 6.1.4 showed that the occupation of lion portion of spouse is household 23(57.5%) following service 12(30%) and both business and service 4(10%) and business 1(2.5%).

Year of Schooling	Frequency	Valid Percent	Cumulative Percent
5.0	4	10.0	10.0
6.0	1	2.5	12.5
8.0	2	5.0	17.5
9.0	2	5.0	22.5
10.0	8	20.0	42.5
11.0	2	5.0	47.5
12.0	7	17.5	65.0
14.0	3	7.5	72.5
16.0	5	12.5	85.0
17.0	1	2.5	87.5
18.0	5	12.5	100.0
Total	40	100.0	

Table-2.6.1.5.Education status of the respondents

Table-2.6.1.5 showed that education of the respondents in terms of schooling year varied between 5 and 18 years and 10 years schooling is highest number of respondents 8(20%) following 12 years 7(17.5%).

2.6.2. Descriptive Statistics of food preference married adults : Table-2.6.2.1 depicted that in the five point scale the highest frequency in the dislike a lot (35), dislike a little (15) neither like or dislike (15), like a little (32) and like a lot (26). Reliability of the questionnaire was checked by Cronbach's Alpha test which is 0.876 that means there is consistency of the items or variables. The total point corresponding to the frequency table 2.6.2.1 is given in table 2.6.2.2.

Table-2.6.2.1: Frequency distribution in percentage of food preference of married adults

Serial No.	Food Item/Variable	1- Dislike a	2- Dislike a	3- Neither like	4- Like a little	5- Like a lot	Total
		lot	little	or dislike			
1	Beef	3%	5%	5%	80%	8%	100
2	Beef burgers	40%	18%	10%	3%	30%	100
3	Lamb	70%	13%	8%	10%	0%	100
4	Chicken	0%	3%	5%	38%	55%	100
5	Bacon	73%	15%	8%	5%	0%	100
6	Ham	70%	15%	10%	5%	0%	100
7	Sausage(Meat)	33%	5%	5%	23%	35%	100
8	White fish	18%	20%	30%	20%	18%	100
9	Oily fish	25%	8%	20%	23%	25%	100
10	Smoked Salmon	30%	20%	18%	18%	15%	100
11	Tinned Tuna	18%	25%	33%	18%	8%	100
12	Eggs	30%	33%	25%	10%	3%	100
13	Baked beans	43%	20%	23%	10%	5%	100
14	Bread or Bread rolls	38%	30%	33%	0%	0%	100
15	Bran cereal	50%	23%	15%	8%	5%	100
16	Porridge	3%	3%	8%	38%	50%	100
17	Plain boiled rice	3%	5%	13%	38%	43%	100
18	Sugared cereal	0%	5%	15%	30%	50%	100
19	Hummus	0%	13%	13%	25%	50%	100
20	Wheat cereal	0%	0%	5%	30%	65%	100
21	Potatoes	58%	25%	18%	0%	0%	100
22	Chips	88%	13%	0%	0%	0%	100

				Ta	able continued		
23	Rice or corn cereal	0%	0%	3%	30%	68%	100
24	Soft cheese	23%	20%	18%	25%	15%	100
25	Hard cheese	43%	30%	20%	8%	0%	100
26	Cottage Cheese	38%	18%	23%	13%	10%	100
27	Plain, Low-fat yoghurt	5%	8%	15%	30%	43%	100
28	Oranges	8%	15%	15%	15%	48%	100
29	Grapes	13%	13%	33%	30%	13%	100
30	-	8%	18%	13%	45%	18%	100
	Apples						
31	Melon	5%	18%	20%	43%	15%	100
32	Peaches	63%	13%	13%	13%	0%	100
33	Apricots	40%	20%	20%	15%	5%	100
34	Strawberries	23%	15%	38%	15%	10%	100
35	Avocadoes	43%	18%	20%	15%	5%	100
36	Spinach	45%	25%	18%	13%	0%	100
37	Carrots	30%	20%	23%	28%	0%	100
38	Green beans	23%	25%	20%	18%	15%	100
39	Cucumber	25%	30%	35%	10%	0%	100
40	Celery	30%	23%	23%	18%	8%	100
41	Mushrooms	3%	8%	10%	20%	60%	100
42	Parsnips	25%	15%	20%	18%	23%	100
43	Peas	23%	18%	40%	13%	8%	100
44	Sweet corn	45%	23%	15%	18%	0%	100
45	Broccoli	20%	20%	15%	30%	20%	100
46	Salad leaves	30%	20%	25%	18%	8%	100
47	Red peppers	25%	20%	25%	20%	10%	100
48	Raw tomatoes	38%	25%	25%	13%	0%	100
49	Beetroot	45%	38%	10%	8%	0%	100
50	Brussels sprouts	58%	5%	23%	15%	0%	100
51	Butter	50%	20%	15%	15%	0%	100
52	Butter-like spreads	73%	15%	8%	5%	0%	100
53	Cream	78%	10%	8%	5%	0%	100
54	Mayonnaise	93%	8%	0%	0%	0%	100
55	Plain biscuits	50%	25%	15%	8%	3%	100
56 57	Chocolate biscuits Cake	70% 80%	10%	10%	10% 3%	0%	100
58	Ice cream	80% 95%	3%	8% 3%	3% 0%		100
58 59	Custard	95% 13%	3%	3%	83%	0% 0%	100
60	Chocolate	75%	13%	10%	3%	0%	100
61	Crisps	63%	18%	15%	5%	0%	100
62	Chewy gummy sweets	75%	23%	3%	0%	0%	100
02	Average	37%	16%	16%	18%	14%	100
	11volage	31/0	10/0	10/0	10/0	17/0	100




Table-2.6.2.2: Total Scaling Point of food preference married between 29-64 years based on table-2.6.2.1

Variable No.	1- Dislike a lot	2- Dislike a little	3- Neither like or dislike	4- Like a little	5- Like a lot	Total	Position
1	1	4	6	128	15	154	Ninth
2	16	14	12	4	60	106	1 111111
3	28	10	9	16	0	63	
4	0	2	6	60	110	178	third
5	29	12	9	8	0	58	
6	28	12	12	8	0	60	
7	13	4	6	36	70	129	
8	7	16	36	32	35	126	
9	10	6	24	36	50	126	
10	12	16	21	28	30	107	
11	7	20	39	28	15	109	
12	12	26	30	16	5	89	
13	17	16	27	16	10	86	
14	15	24	39	0	0	78	
15	20	18	18	12	10	78	
16	1	2	9	60	100	172	Fourth
17	1	4	15	60	85	165	seventh
18	0	4	18	48	100	170	sixth
19	0	10	15	40	100	165	seventh
20	0	0	6	48	130	184	Second
21	23	20	21	0	0	64	
22	35	10	0	0	0	45	
23	0	0	3	48	135	186	First
24	9	16	21	40	30	116	
25	17	24	24	12	0	77	
26	15	14	27	20	20	96	
27	2	6	18	48	85	159	Eight
28	3	12	18	24	95	152	
29	5	10	39	48	25	127	
30	3	14	15	72	35	139	
31	2	14	24	68	30	138	
32	25	10	15	20	0	70	
33	16	16	24	24	10	90	
34	9	12	45	24	20	110	
35	17	14	24	24	10	89	
36	18	20	21	20	0	79	
37	12	16	27	44	0	99	
38	9	20	24	28	30	111	
39 40	10 12	24 18	42 27	16 28	0 15	92 100	
41	1	6	12	32	120	171	Fifth
42	10	12	24	28	45	119	11111
43	9	14	48	20	15	106	
44	18	18	18	28	0	82	
45	8	16	18	48	40	130	
46	12	16	30 30	28	15 20	101	
47 48	10 15	16 20	30	32 20	0	108 85	
49	18	30	12	12	0	72	
50	23	4	27	24	0	78	
51	20	16	18	24	0	78	
52	29	12	9	8	0	58	
53	31	8	9	8	0	56	
54 55	37 20	6 20	0 18	0 12	5	43 75	
56	28	8	18	16	0	64	
57	32	8	9	4	0	53	
58	38	2	3	0	0	43	
59	5	2	3	132	0	142	
60	30	10	12	4	0	56	
61	25	14	18	8	0	65	
62	30	18	3	0	0	51	
			ge score of items			102.87 1.66	
Average score of respondents					1.00		




Figure: 2.6.2.2. The total point od food preference of adults married

From the above analysis the top ten preferred food items of married adults are listed below

Serial No	Food Item	Total	Position
1	Rice or corn cereal	186	First
2	Wheat cereal	184	Second
3	Chicken	178	third
4	Porridge	172	Fourth
5	Mushrooms	171	Fifth
6	Sugared cereal	170	sixth
7	Plain boiled rice	165	seventh
8	Hummus	165	seventh
9	Plain, Low-fat yoghurt	159	Eight
10	Beef	154	Ninth

2.7. A Comparison of food preference between adults of 20-29 unmarried and 29-64

married: Being the two groups were given same checklist of food preference so we can compare the preferences of them.

2.7.1: Comparison of total point score by food items of the groups: From table-2.5.2.2 and table-2.6.2.2 a comparative table is made as below in table 2.7.1

Table-2.7.1: Comparative food preference score of unmarried adults of (20-29) married adults (29-64) based on table 5.2.2.2 and 6.2.2.2

Variable No.	unmarried adults (20-29)	married adults(29-64)
1	161	154
2	159	106
3	169	63
4	180	178
5	100	58
6	54	60
7	129	129
8	113	126
9	124	126
10	61	107
11 12	108 137	109 89
13	137	86
14	140	78
15	122	78
16	111	172
17	149	165
18	80	170
19	42	165
20	135	184
21	117	64
22	188	45
23	160	186
24	128	116
25	121	77
26	144	96
27	135	159
28	194	152
29	190	127
30	171	139
31	144	138
32	59	70 90
33	69 125	
34 35	69	110 89
36	150	79
37	141	99
38	94	111
39	170	92
40	180	100
41	154	171
42 43	123 121	119 106
43	121	82
45	89	130
46	110	101
47	135	108
48	170	85
49 50	118 78	72 78
51	89	78
52	144	58
53	67	56
54	124	43
55 56	139 147	75 64
56 57	147	53
58	191	43
59	142	142
60	194	56
61	198	65
62	122	51




Figure-2.7.1: Comparative food preference of unmarried adults and married adults

Note: X= Food preference of unmarried adults and Y= Food preference of married adults

2.7.2: Two –sample t- test: The food preference of un married adults and food preference of married are compared and test whether there is any significant difference among their choices. And for this two –sample t- test is applied.

Table-2.7.2.1Group Statistics

Group	N	Mean	Std. Deviation	Std. Error Mean
Unmarried	62	191.10	89.873	11.414
Married	62	102.87	39.889	5.066

Table-2.7.2.2:Independent Samples Test

	Levene's Test for Equality of Variances		t-test for Equality of Means						
	Equality 0	1 variances				•		95% Confidence Interval of the	
	F	Sig.	t	df	Sig. (2- tailed)	Mean Difference	Std. Error Difference	Lower	Upper
Equal variances assumed	36.142	.000	7.065	122	.000	88.226	12.488	63.505	112.946
Equal variances not assumed			7.065	84.135	.000	88.226	12.488	63.393	113.058

In the table of group statistics we can find that mean score of unmarried and married are 191.10 and 102.87 correspondingly with Std. deviation 89.873 and 39.889 for unmarried and married respectively. Therefore the average score of unmarried adult is higher than that of married adults. The table of independent Samples Test depicted two test - Levene's Test for Equality of Variances and t-test for Levene's Test for Equality of means. In the first test i.e. Levene's Test for Equality of Variances here our null hypothesis is that the both the variances are equal and alternative is that the variances are different for the two groups. The Levene's Test for Equality of Variances showed the p- value is 0.00 that means there are significant differences of variances of the two groups. The second test is t-test for Equality of Means which is presented in the second row and showed that the observed test statistics is 7.065 with p- value 0 .000. In this case we can also reject the null hypothesis and can conclude that there is significant difference between the means of the two groups.

Finally we can conclude that there is significant difference between the food preference of unmarried adults and married adults and their preferences s are not same.

8. Overall awareness and behavior of food hygiene and nutrition: There are six kinds of respondents and after analysis the overall average score each group is given in table 8.

Table 8: Overall average score of the respondents in different scale

Serial	Respondents	Number of	Scale of	Item-wise	Average
No.	-	respondents	questionnaire	Average	score by
				score	respondents
1.	Parents of baby up to 4 years	40	5.00	100.5	2.51
2.	Parents of Child eating behavior	40	5.00	117.66	3.36
	between 5-9 years				
3.	Adolescent (10-19 years) Food	34	3.00	43.56	1.89
	Habits				
4.	Parental Feeding Style to his	34	5.00	114.52	4.24
	Children up to 9 years				
5.	Food preference of adults unmarried	40	5.00	131.35	2.12
	(20-29)				
6.	Food Preference for Adults (29-64)	40	5.00	102.87	1.66
	Married Male and Female				

9. Conclusion: This chapter has analyzed collected data through different descriptive and inferential statistical techniques and methods. The major findings produced from the analyses are given in the next chapter.

CHAPTER THREE

FINDINGS AND CONCLUSION

Introduction: In the previous chapter the data has been analyzed using different statistical techniques of six groups of our respondents in this chapter the major findings and suggestions are given below:

3.1. Findings on f Baby's Eating Behavior up to 4(Four Year) Responded by Parents: Table-2.1.2: Frequency of distribution baby's eating behavior showed the highest frequency as follows: (i) in the scale never =1 "Even when my baby had just eaten well he/she was happy to feed again if offered" showed highest frequency 50% (20); (ii) in the scale rarely=2 "Baby could easily take a feed within 30 minutes of the last one showed 45% (18), (iii) in the scale sometimes = 3 "Baby seemed contented while feeding 50% (20), in the scale often=4 "Baby became distressed while feeding and Baby got full before taking all the milk I think he/she should have" 30% (12) each and in the scale 5=Always "Always Baby fed slowly" 30%(12). The average frequency in percentage in the different scales are 23%, 30%, 16%,11% and 11% respectively. Table 2.1.3 showed that the top five score items are – my baby loved milk(155) following my baby seemed contented while feeding (154), my baby fed slowly(150), my baby found it difficult to manage a complete feed(131), my baby got full up easily(113) and my baby frequently wanted more milk than I provided (105). The average score of the items is 100.5 and the average score of the respondents is 2.51. It means the eating behavior of baby lies between rarely and sometimes. Reliability of the questionnaire was checked by Cronbach's Alpha test which is 0.710 that means there is consistency of the items or variables. The PCA reduces 21 items into six factors significant for eating behavior of parents.

3.2. Child eating behavior between 5-9 years: The table -2.2.2.1 depicted the highest frequency among 35 items that in the scale never=1 " If allowed to, my child would eat too much item is highest 50% (20), in the scale rarely=2 " my child eats more when anxious item is highest" 47.5%(19) in the scale sometimes = 3, " my child loves food" 65%(26), in the scale often =4, " my child is difficult to please with meals item is highest" 45% 18 and in the scale always=5, "my child gets full up easily item is highest" 55%(22). The average percentage of frequency distribution in the five point scales are 20%,27%, 23%, 16% and 19% respectively.

The table 2.2.2.2 depicted the total point of each variable and the top seven items are as follows-My child gets full up easily(165), Child cannot eat a meal if s/he has had a snack just before(149), My child is interested in tasting food s/he hasn't tasted before(145), My child is always asking for food(142), My child enjoys a wide variety of foods(142), My child leaves food on his/her plate at the end of a meal(137), My child loves food(137), My child is always asking for a drink(136), My child eats less when angry(135). The average of items is 117.66 and the average of respondents is 3.36. That means the eating behavior of children varied between sometimes and often. Through the PCA 35 items are reduced to 10 factors.

- **3.3.** Adolescent (10-19) Food Habits: The food habits of adolescent is measured in terms of given items or variables with three point scale as Most like=2; Neutral=0; and like=1. Table 2.3.2.1 showed that the highest frequency in the scale Most like=2 is "When I am buying a soft drink, I usually choose a diet drink" 76%(26); in the scale like=1 the item "I rarely takeaway meals 82% (28)" and in the scale Neutral = 0,If I am buying crisps, I often choose a low-fat brand 32% (11) among twenty three items. The average percentages in the scales are 37%, 55% and 8% respectively. The top seven total scoring items are: (i)When I am buying a soft drink, I usually choose a diet drink (60), (ii)I eat at least servings of fruits most days(57),(iii)I usually avoid eating fried foods(53) (iv)I often eat sweet snacks between meals(51),(v) I make sure I eat at least one serving of fruit a day (49)(vi) I avoid eating lots of sausages and burgers.(46) and (vii) When I put butter or margarine on bread and I usually spread it on thinly If I have a packed lunch, I usually include some chocolate and/or biscuits (43). Average of score of the items is 43.56 and the average score of respondents is 1.89. The PCA reduces 21 items to eight factors of food habits of adolescent.
- **3.4. Parental Feeding Style to his Children up to 9 years-:** Table-2.4.2.1 provided that in the five point scale the highest frequency showing items are in the scales never=1 is "I allow my child to wander around during a meal(14)", rarely =2 is "I use puddings as a bribe to get my child to eat his/her main course rarely (17)" sometimes=3 is "I reward my child with something to eat when s/he is well behaved (16)" often =4 is "I decide how many snacks my child should have(8) and I encourage my child to eat a wide variety of foods (8)" and always=5 is "I encourage my child to try foods that s/he hasn't tasted before (19). The average percentage of frequency distribution in the five point scales are 15%, 10%, 33%, 13% and 30% respectively.

The top seven score point items in the scales are I encourage my child to eat a wide variety of foods (140), I encourage my child to try foods that s/he hasn't tasted before(139), I encourage my child to enjoy his/her food(138), I give my child something to eat to make him/her feel better when s/he is feeling upset(135), I encourage my child to look forward to the meal and I praise my child if s/he eats what I give him/her(134), I encourage my child to taste each of the foods I serve at mealtimes(132), I praise my child if s/he eats a new food(124)(Table-2.4.2.2). The total variance explained by the PCA is 78.96% that means the PCA is accepted and based on Eigen values seven items are considered important. The Average of items is 114.52 and the average of respondents is 4.24 that means the parental Feeding Style to his Children up to 9 years varies between often and always.

- **3.5. Food preference of adults unmarried (20-29):** From table-2.5.2.1 we can find that in the five point scale the highest frequency in the scales are dislike a lot=1 is hummus 97.5% (39), dislike a little=2 is Green beans 30% (12) ,neither like or dislike=3 is Bacon 65% (26) , like a little=4 is Oranges 92.5% (37). The average percentages of frequency distribution in the different scales are 21%, 8%, 17% 31% and 24% respectively. The top ten preferred food items are Crisps(198), Chocolate(194), Oranges(194), Ice cream(191), Grapes(190), Chips(188) , cake(186), Celery(180), Chicken(180) and apples(171)(Table -2.5.2.3). The Average of items is 131.35 and the average of respondents is 2.12 that mean the food preference of adults (20-29 years) varies between neither like or dislike and like a little.
- **3.6. Food Preference for Adults (29-64) Married Male and Female:** From table-2.6.2.1 we can find that in the five point scale the highest frequency in the scales are dislike a lot =1 is Chips 87.5% (35), dislike a little=2 is Beetroot 37.5% (15) neither like or dislike=3 is Strawberries 37.5% (15), like a little=4 is Custard 82.5% (33) and like a lot =5 is Rice or corn cereal 67.5% (27). The average percentages of frequency distribution in the different scales are 37%, 16%, 16% 18% and 14% respectively. The top ten preferred food items are Rice or corn cereal(186), Wheat cereal(184), Chicken(178), Porridge(172) Mushrooms(171), Sugared cereal(170) Plain boiled rice(165), Hummus(165), Plain, Low-fat yoghurt(159) and Beef(154)(Table--2.6.2.3). The Average of items is 102.87 and the average of respondents is

- 1.66 that means the food preference of adults (29-64 years) varies between dislike a little and neither like nor dislike.
- 3.7. A Comparison of food preference between adults of 20-29 unmarried and 29-64 married: Being the two groups were given same checklist of food preference so we can compare the preferences of them. The average score of adults of 20-29 unmarried is 131.35 while it is 102.87 for married 102.87 so the average score of adults of 20-29 unmarried is higher than that of married 29-64. Two sample t-test was applied to confirm whether there is any similarity between these two groups. In the table 2.7.1 of group statistics we can find that mean score of unmarried and married are 191.10 and 102.87 correspondingly with Std. deviation 89.873 and 39.889 for unmarried and married respectively. Therefore the average score of unmarried adult is higher than that of married adults. The table of independent Samples Test (table-2.7.2) depicted two test - Levene's Test for Equality of Variances and t-test for Levene's Test for Equality of means. In the first test i.e. Levene's Test for Equality of Variances here our null hypothesis is that the both the variances are equal and alternative is that the variances are different for the two groups. The Levene's Test for Equality of Variances showed the p-value is 0.00 that means there are significant differences of variances of the two groups. The second test is t-test for Equality of Means which is presented in the second row and showed that the observed test statistics is 7.065 with p- value 0 .000. In this case we can also reject the null hypothesis and can conclude that there is significant difference between the means of the two groups. Finally we can conclude that there is significant difference between the food preference of unmarried adults and married adults and their preferences s are not same.
- **3.8 Suggestions for policy implications:** Following the analysis and findings the suggestions are prescribed below for better awareness of food safety hygiene and nutrition:
- (i) The average awareness of parents for their baby up to four years is 2.51 out of 5 around 50%. Therefore, it is needed to increase awareness of the parents in feeding the babies.
- (ii) The average awareness of parents to their child's eating behavior between 5-9 years is 3.36 out of 5. The awareness is around 67% so we should be more careful in this regard.
- (iii) The average awareness adolescent (10-19 years) food habit is 1.89 on three point scale which about 63% only. Therefore the consciousness of adolescent (10-19 years) is be to increase in taking food.

(iv) The average feeding styles of parents is 4.24 out of 5 to their children up to 9 years and in percentage it is 84%. Therefore it is about to satisfactory.

REFERNCES

- ALAM, M. M., & HOSSAIN, M. K. (2018). Policy options on sustainable resource utilization and food security in Haor areas of Bangladesh: A theoretical approach. *International Journal of Social, Political and Economic Research*, 5(1), 11-28.
- Alam, M. M., & Zakaria, A. F. M. (2021). A Probit Estimation of Urban Bases of Environmental Awareness: Evidence from Sylhet City, Bangladesh. *arXiv* preprint arXiv:2107.08342.
- Bhuvneswari, R. and Muthukumar Dr. E.(2015), Food Safety Awareness Among Customers To Avoid The Risk Of Food Borne Illness, *Journal of Business Management & Social Sciences Research* (JBM&SSR) ISSN No: 2319-5614, Volume 4, No.8
- Bruhn Christine M. (1997), Consumer Concerns: Motivating to Action, *Emerging Infectious Diseases*, Vol. 3. No. 4
- Bruhn Christine M and Schutz Howard G. (1999), Consumer Food Safety Knowledge And Practices, Journal of Food Safety, Vol. 19,pp.73-87
- Buzby JC, Roberts T 1995), ERS estimates US foodborne disease costs, Food Review, vol 18, pp37–42)
- Cameron S, Walker W, Beers M et al 1995, Enterohaemorrhagic Escherichia coli outbreak in South Australia associated with the consumption of mettwurst, Communicable Diseases Intelligence, vol 19, pp70–100)
- Emmanuel K. Yiridoe, Samuel Bonti-Ankomah and Ralph C. Martin(2005), Comparison of consumer perceptions and preference toward organic versus conventionally produced foods: A review and update of the literature, *Renewable Agriculture and Food Systems*, Vol. 20, Isuue. 4 pp. 193–205 DOI: 10. 1079 / RAF 2005113
- Daily star Newspaper(2105), Daily star, Bangladesh April 8, 2015
- Doaa M., Nasr-Eldein, Sahar M. Soliman, Samar El Hossiny(2017), Effect of Applying Different Educational Strategies on Dietary Habits of Preparatory Schools' Pupils, *IOSR Journal of Nursing and Health Science* (IOSR-JNHS) e-ISSN: 2320–1959.p- ISSN: 2320–1940 Volume 6, Issue 4 Ver. IV. PP. 12-23
- Government of Bangladesh (2018) Second Country Investment Plan (CIP) for Nutrition-sensitive Food Systems.
- Finance Division. "Climate Fiscal Framework." (2014).
- Finance Division. (2014). Bangladesh Climate Fiscal Framework, Ministry of Finance, GPRB Finance Division, Ministry of Finance, Government of the People's Republic of Bangladesh, Dhaka
- FAO (2014) Guidelines for assessing nutrition-related Knowledge, Attitudes and Practices, KAP Manual.
- Field, A. (2009). Discovering Statistics using SPSS. Sage: London
- Fung Fred, Wang Huei Shyong & Menon Suresh(2018), Food Safety in 21st Century, *Biomedical Journal*, Science Direct, Vol. 41, pp-88-95
- Hoddinott Jhon, Ahmed Akhter, Karachiwall Naureen I and Roy Shalini (2017), Nutrition behavior change communication causes sustained effects on IYCN knowledge in two cluster randomized trials in Bangladesh, *Maternal and Child Nutrition*, Wiley.
- Islam, M. R., Alam, M., & Afzal, M. N. İ. (2021). Nighttime Light Intensity and Child Health Outcomes in Bangladesh. *arXiv preprint* arXiv:2108.00926.
- Karmakar P, Jahan N, Banik S, Das A, Rahman KA, Kundu SK and Sattar MM(2016), Food

- Habits, Obesity and Introduction, Journal of Food & Nutritional Disorders , Nutritional Knowledge among the University Students in Noakhali Region of Bangladesh:ACross Sectional Study Journal of Food & Nutritional Disorders, Vol. 5, Issu.4 http://dx.doi.org/10.4172/2324-9323.1000201
- Karmakar P., Jahan N., Banik S., Das A., Rahman K.A., Kundu S.K. and Sattar M.M. (2017) Food Habits, Obesity and Nutritional Knowledge among the University Students in Noakhali Region of Bangladesh: A Cross Sectional Study, Journal of Food and Nutritional Disorders 5(4)
- Nutrition Iqbal Kabir, A.K.M, S. K. Roy, S. Khatoon (2013) Development of a Complementary Feeding Manual for Bangladesh, BBF for NFPCSP
- Institute of Public Health and WHO(1994, 2003
- Manikam Logan , Raghu Lingam , Isabel Lever , Emma C. Alexander , Chidi Amadi , Yasmin Milner , Taimur Shafi , Lucy Stephenson , Sonia Ahmed and Monica Lakhanpaul(2018), Complementary Feeding Practices for South Asian Young Children Living in High-Income Countries: A Systematic Review, *Nutrients* , 10, 1676; doi:10.3390/nu10111676 www.mdpi.com/journal/nutrients
- Mahmud, I. and N. Mbuya (2016) Water, Sanitation, Hygiene, and Nutrition in Bangladesh Can Building Toilets Affect Children's Growth? World Bank Study
- Nevin Sanlier, Ece Konaklioglu, (2012) "Food safety knowledge, attitude and food handling practices of students", British Food Journal, Vol. 114 Issue: 4, pp.469-480, https://doi.org/10.1108/0007070 12 112 19504
- Nesbitt Andrea, Thomas M.Kate, Marshal Barbara, Snedeker Kate, Meleta Kathryn, Watson Brenda and Bienefeld Monika (2014), Baseline for consumer food safety knowledge and behavior in Canada, Food Control, Vol. 38, pp. 157-173.
- Nina Michaelidou and Hassan Louise M(n.d), The Role of Health Consciousness, Food Safety Concern and Ethical Identity on Attitudes and Intentions towards Organic Food, University of Birmingham Birmingham Business School University House Edgbaston and University of Stirling and the Open University Institute for Social Marketing University of Stirling
- Nunnally JC (1978). Psychometric Theory, 2nd ed. New York: McGraw-Hill.
- Nurul Alam, Swapan Kumar Roy, Tahmeed Ahmed, and A.M. Shamsir Ahmed(2010), Nutritional Status, Dietary Intake, and Relevant Knowledge of Adolescent Girls in Rural Bangladesh, J HEALTH POPUL NUTR 2010 Feb;28(1):86-94 ISSN 1606-0997 | \$ 5.00+0.20
- Quinlan Jennifer J. (2013), Foodborne Illness Incidence Rates and Food Safety Risks for Populations of Low Socioeconomic Status and Minority Race/Ethnicity: A Review of the Literature, *Int. J. Environ. Res. Public Health* Vol. 10, 3634-3652; doi:10.3390/ijerph10083634
- Scallan, Elaine & Hoekstra, Mike & J Angulo, Frederick & V Tauxe, Robert & Widdowson, Marc-Alain & Roy, Sharon & L Jones, Jeffery & M Griffin, Patricia. (2011). Foodborne Illness Acquired in the United States—Major Pathogens. Emerging infectious diseases. 17. 7-15. 10.3201/eid1701.091101p1
- Sedef Akgüngör, Bülent Miran , Canan Abay (2007), Consumer Willingness to Pay for Organic Products in Urban Turk, Contributed Paper prepared for presentation at the 105th EAAE Seminar 'International Marketing and International Trade of Quality Food Products', Bologna, Italy, March 8-10, 2007.

- Saha ML, T Bari, MR Khan and S Hoque 2011. Bacteriological and physico-chemical properties of the Gulshan lake, Dhaka. Bangladesh J. Bot. Vol.40,Iss.2, pp.105-111
- Saha, S., M.K. Zahid, S. Rasheed (2011) The Study of the Level of Knowledge, Attitude, Practices (KAP) as well as the Effects of School Environment on the Nutritional Status of Children (7-12) Coming from Affluent Families in the Dhaka City in Bangladesh, Bangladesh Journal of Nutrition Vol. 24-25
- SPRING (2017) Bangladesh: Farmer Nutrition School Cohort Study. Sustainability of Improved Practices Following Graduation. Strengthening Partnerships, Results, and Innovations in Nutrition Globally (SPRING) project
- Sienny, T. and Serli, W.(2010), The concern and awareness of consumers and food service operators towards food safety and food hygiene in small and medium restaurants in Surabaya, Indonesia, *International Food Research Journal*, Vol.17, pp.641-650.
- Subba rao G. M, Sudershan R. V., Rao Pratima , Vishnu Rao M., Polasa Kalpagam, "Food safety knowledge, attitudes and practices of mothers findings from focus group studies in south India", Appetite 49(2007),
- WHO(2000), The world Health Report, 2000. WHO, Geneva.
- WFP (2015) Safety Nets: Cash With Nutrition Education Has Greatest Impact On Child Nutrition, https://www.wfp.org/news/news-release/safety-nets-cash-nutrition-education-has-greatest-impactchild-nutrition.
- Van Dillen Sonja ME, Hiddink Gerrit J, Koelen Maria A, Graaf Cees de, and Woerkum Cees MJ van(2003), Understanding nutrition communication between health professionals and consumers: development of a model for nutrition awareness based on qualitative consumer research, *American Society for Clinical Nutrition*, Vol. 77(suppl), pp.1065S–72s

Appendix -1

~	Appendix -1Communalities of Baby's eating behavior	01	1
S.No.	Variables/ items of behavior	Initial	Extraction
1	My baby seemed contented while feeding	1.000	.769
2	My baby frequently wanted more milk than I provided	1.000	.844
3	My baby loved milk	1.000	.905
4	My baby had a big appetite	1.000	.782
5	My baby finished feeding quickly	1.000	.745
6	My baby became distressed while feeding	1.000	.859
7	13.7 My baby got full up easily	1.000	.654
8	If allowed to, my baby would take too much milk	1.000	.813
9	My baby took more than 30 minutes to finish feeding	1.000	.781
10	My baby got full before taking all the milk I think he/she should have	1.000	.733
11	My baby fed slowly	1.000	.710
12	Even when my baby had just eaten well he/she was happy to feed again if offered	1.000	.734
13	My baby found it difficult to manage a complete feed	1.000	.855
14	My baby was always demanding a feed	1.000	.722
15	My baby sucked more and more slowly during the course of a feed	1.000	.717
16	If given the chance, my baby would always be feeding	1.000	.830
17	My baby enjoyed feeding time	1.000	.634
18	My baby could easily take a feed within 30 minutes of the last one	1.000	.694

Appendix-2:Total value explained by PCA of Baby's Eating Behavior

]	Initial Eigenva	lues	Rotation Sums of Squared Loadings			
		% of					
Component	Total	Variance	Cumulative %	Total	% of Variance	Cumulative %	
1	5.577	30.982	30.982	2.617	14.540	14.540	
2	2.736	15.202	46.184	2.473	13.737	28.277	
3	1.880	10.444	56.628	2.374	13.191	41.469	
4	1.510	8.390	65.018	2.228	12.376	53.845	
5	1.066	5.922	70.940	2.053	11.407	65.252	
6	1.013	5.626	76.566	2.037	11.314	76.566	
7	.866	4.809	81.375				
8	.727	4.038	85.414				
9	.557	3.097	88.511				
10	.518	2.876	91.387				
11	.340	1.889	93.276				
12	.316	1.755	95.031				
13	.257	1.426	96.457				
14	.204	1.132	97.589				
15	.180	1.000	98.589				
16	.116	.644	99.233				
17	.086	.475	99.708				
18	.053	.292	100.000	·			

	Component								
Items/ Variable	1	2	3	4	5	6			
1	103	513	.029	.070	.189	.674			
2	167	.028	.122	.891	082	004			
3	.328	.094	002	.847	.264	040			
4	.280	.768	204	.248	.096	022			
5	.815	.197	027	.023	102	174			
6	354	.177	.059	.117	031	.827			
7	.010	031	.793	040	.126	.084			
8	.527	.263	135	.634	136	.164			
9	404	.164	.010	.127	.730	.202			
10	056	385	.589	.090	.435	.195			
11	725	006	.222	108	.279	.213			
12	.471	.475	.018	.115	410	325			
13	009	.149	.334	301	.418	.675			
14	.120	.159	824	.014	.000	.063			
15	161	208	.300	.012	.746	.035			
16	.021	.893	053	.019	076	.152			
17	.520	.409	142	.131	393	071			
18	.316	117	606	273	160	336			

Extraction Method: Principal Component Analysis.
Rotation Method: Varimax with Kaiser Normalization.
a. Rotation converged in 8 iterations.

Appendix-4 Item-Total Statistics of Childs' eating behavior

Variable or Items of behavior	Scale Mean if Item Deleted	Scale Variance if Item Deleted	Corrected Item-Total Correlation	Cronbach's Alpha if Item Deleted
My child loves food	65.350	153.105	.201	.762
My child eats more when worried	66.700	150.267	.404	.754
My child has big appetite	66.025	154.999	.046	.773
My child finishes his/her meal quickly	66.100	165.836	275	.798
My child is interested in food	65.300	161.395	176	.783
My child is always asking for a drink	65.375	145.266	.328	.751
My child refuses new foods at first	65.950	156.356	014	.780
My child eats slowly	65.650	143.259	.252	.757
My child eats less when angry	65.400	158.810	079	.784
My child enjoys tasting new foods	64.925	150.584	.254	.758
My child eats less when s/he is tired	65.500	156.462	005	.777
My child is always asking for food	66.375	150.292	.234	.759
My child eats more when annoyed	67.000	153.026	.232	.761
If allowed to, my child would eat too much	66.925	150.892	.213	.761
My child eats more when anxious	66.900	155.785	.058	.770
My child enjoys a wide variety of foods	65.225	153.666	.133	.766
My child leaves food on his/her plate at the end of a meal	65.225	144.281	.283	.754
My child takes more than 30 minutes to finish a meal	65.650	146.387	.291	.754
Given the choice, my child would eat most of the time	66.400	137.579	.526	.732
My child looks forward to mealtimes	66.275	149.230	.264	.757
My child gets full before his/her meal is finished	65.350	145.105	.390	.747

65.250	149.269	.388	.753
65.400	148.503	.399	.752
65.700	149.703	.216	.760
65.800	147.446	.239	.758
64.900	163.990	215	.796
66.425	139.533	.514	.735
65.400	158.656	081	.788
66.350	143.362	.548	.740
65.050	140.818	.461	.739
66.250	144.910	.469	.744
65.150	158.900	079	.783
66.150	145.669	.316	.752
67.025	155.307	.061	.770
65.900	141.579	.484	.739
	65.400 65.700 65.800 64.900 66.425 65.400 66.350 65.050 66.250 65.150 66.150	65.400 148.503 65.700 149.703 65.800 147.446 64.900 163.990 66.425 139.533 65.400 158.656 66.350 143.362 65.050 140.818 66.250 144.910 65.150 158.900 66.150 145.669 67.025 155.307	65.400 148.503 .399 65.700 149.703 .216 65.800 147.446 .239 64.900 163.990 215 66.425 139.533 .514 65.400 158.656 081 66.350 143.362 .548 65.050 140.818 .461 66.250 144.910 .469 65.150 158.900 079 66.150 145.669 .316 67.025 155.307 .061

Appendix-5: Communalities of variables of Childs' eating behavior

Communalities					
	Initial	Extraction			
My child loves food	1.000	.798			
My child eats more when worried	1.000	.857			
My child has big appetite	1.000	.807			
My child finishes his/her meal quickly	1.000	.901			
My child is interested in food	1.000	.890			
My child is always asking for a drink	1.000	.945			
My child refuses new foods at first	1.000	.847			
My child eats slowly	1.000	.960			
My child eats less when angry	1.000	.775			
My child enjoys tasting new foods	1.000	.905			
My child eats less when s/he is tired	1.000	.851			
My child is always asking for food	1.000	.868			
My child eats more when annoyed	1.000	.856			
If allowed to, my child would eat too much	1.000	.807			
My child eats more when anxious	1.000	.911			

My child enjoys a wide variety of foods	1.000	.831
My child leaves food on his/her plate at the end of a meal	1.000	.940
My child takes more than 30 minutes to finish a meal	1.000	.906
Given the choice, my child would eat most of the time	1.000	.866
My child looks forward to mealtimes	1.000	.884
My child gets full before his/her meal is finished	1.000	.931
My child enjoys eating	1.000	.925
My child eats more when s/he is happy	1.000	.898
My child is difficult to please with meals	1.000	.900
My child eats less when upset	1.000	.947
My child gets full up easily	1.000	.970
My child eats more when s/he has nothing else to do	1.000	.923
Even if child is full ups/he finds room to eat his/her favorite food	1.000	.912
If given the chance, child drink continuously throughout the day	1.000	.922
Child cannot eat a meal if s/he has had a snack just before	1.000	.791
if given the chance my child would always be having a drink	1.000	.888
My child is interested in tasting food s/he hasn't tasted before	1.000	.842
Child decides that s/he doesn't like a food, even without tasting it	1.000	.931
If given the chance, child would always have food in his/her mouth	1.000	.900
My child eats more and more slowly during the course of a meal	1.000	.851
Extraction Method: Principal Component Analysis.	1	

Appendix-5: Total Variance Explained by PCA of Childs' Eating Behavior

		Initial Eigenvalues Extraction Sums of Squared Load			Loadings	
Component	Total	% of Variance	Cumulative %	Total	% of Variance	Cumulative %
1	8.639	24.683	24.683	8.639	24.683	24.683
2	5.075	14.499	39.183	5.075	14.499	39.183
3	4.781	13.661	52.844	4.781	13.661	52.844
4	2.713	7.752	60.596	2.713	7.752	60.596
5	2.337	6.676	67.272	2.337	6.676	67.272
6	1.927	5.506	72.777	1.927	5.506	72.777
7	1.602	4.577	77.355	1.602	4.577	77.355
8	1.337	3.821	81.176	1.337	3.821	81.176
9	1.286	3.675	84.850	1.286	3.675	84.850
10	1.238	3.538	88.388	1.238	3.538	88.388
11	.779	2.226	90.615			
12	.733	2.094	92.709			
13	.579	1.653	94.362			
14	.479	1.368	95.730			
15	.341	.975	96.706			
16	.269	.768	97.474			
17	.233	.664	98.138			
18	.177	.505	98.643			
19	.138	.395	99.038			
20	.102	.292	99.330			

22 23 24 25 26 27 28 29	.056 .036 .028 .020 .012 .008	.160 .104 .079 .057 .033 .024	99.686 99.790 99.870 99.926 99.959 99.983 99.994			
24 25 26 27 28 29	.028 .020 .012 .008 .004	.079 .057 .033 .024	99.870 99.926 99.959 99.983			
25 26 27 28 29	.020 .012 .008 .004	.057 .033 .024	99.926 99.959 99.983			
26 27 28 29	.012 .008 .004	.033 .024	99.959 99.983			
27 28 29	.008 .004	.024	99.983			
28 29	.004					
29		.011	99 994			
	001		//.// 1			
	.001	.004	99.998			
30	.001	.002	100.000			
31	.000	.000	100.000			
32	5.788E-16	1.654E-15	100.000			
33	2.478E-16	7.081E-16	100.000			
34	1.658E-16	4.737E-16	100.000	•		
35	-3.625E-16	-1.036E-15	100.000			

Appendix 6: Communalities of Variables of Adolescent behavior

Sl. No. of Variable	Initial	Extraction
1	1.000	.864
2	1.000	.574
3	1.000	.903
4	1.000	.797
5	1.000	.808
6	1.000	.736
7	1.000	.838
8	1.000	.815
9	1.000	.800
10	1.000	.859
11	1.000	.854
12	1.000	.631
13	1.000	.787
14	1.000	.780
15	1.000	.747
16	1.000	.811
17	1.000	.880
18	1.000	.701
19	1.000	.778
20	1.000	.802
21	1.000	.760
22	1.000	.897
23	1.000	.812

Extraction Method: Principal Component Analysis.

Appendix-7: Total Variance Explained

	Initial Eigenvalues			Extraction Sums of Squared Loadings			
Component	Total	% of Variance	Cumulative %	Total	% of Variance	Cumulative %	
1	6.350	27.607	27.607	6.350	27.607	27.607	
2	2.770	12.042	39.649	2.770	12.042	39.649	
3	2.265	9.846	49.495	2.265	9.846	49.495	
4	1.871	8.135	57.631	1.871	8.135	57.631	
5	1.538	6.685	64.316	1.538	6.685	64.316	
6	1.255	5.456	69.772	1.255	5.456	69.772	
7	1.112	4.836	74.608	1.112	4.836	74.608	
8	1.074	4.670	79.278	1.074	4.670	79.278	
9	.882	3.836	83.114				
10	.709	3.081	86.195				
11	.622	2.705	88.900				
12	.573	2.492	91.392				
13	.492	2.140	93.532				
14	.357	1.554	95.086				
15	.259	1.127	96.213				
16	.211	.918	97.131				
17	.197	.858	97.989				
18	.137	.596	98.585				
19	.130	.564	99.149				
20	.076	.331	99.480				
21	.065	.281	99.760				
22	.037	.162	99.923				
23	.018	.077	100.000				

 ${\bf Appendix} \ \textbf{-8 - Component Matrix}$

Component Matrix ^a								
		Component						
	1	2	3	4	5	6	7	8
1_lowfat	.672	.091	.136	457	078	.261	.212	.241
2_eating	.507	.101	.079	249	258	355	158	.146
3_dessert	299	.786	.118	.054	333	254	052	.046
4_least	.782	.168	061	.354	107	071	076	079
5_fat	.681	150	166	185	124	129	.264	398
6_crisps	.513	.046	.482	100	.171	.177	137	.387
7_sausages	.534	082	089	.220	262	.432	.476	.086
8_pastries	268	.777	037	222	.008	.093	.279	034
9_sugar	.568	.135	.071	.201	483	.201	.101	360
10_vegetables	.590	279	375	.262	168	049	435	067
11_home	.697	356	054	273	.258	094	.131	266
12_meals	.380	294	.404	.021	263	015	050	.405
13_fruit	.766	015	.006	.129	373	036	030	.206
14_sweet	.012	.419	169	018	.223	.679	231	106

15_salad	.419	.109	577	325	.219	233	.007	.136
16_softdrink	.516	.583	352	.074	.178	073	197	.020
17_butter	.180	.180	053	.445	.391	321	.544	.249
18_chocolate	.412	.177	.293	.581	.258	045	035	081
19_snack	.385	051	.623	286	.352	.074	111	126
20_healthiest	.385	.193	.585	.378	.267	081	061	225
21_cream	.457	458	406	.126	.359	.122	003	.133
22_servings	.671	.528	318	044	.129	.081	124	.160
23_diet	.608	.205	.270	489	.057	188	.018	222
Extraction Method: Principal Component Analysis.								
a. 8 components extracted.								

pendix-9 Communalities of variables of parents eating behavior

Variable No.	Initial	Extraction
1	1.000	.713
2	1.000	.806
3	1.000	.740
4	1.000	.819
5	1.000	.820
6	1.000	.843
7	1.000	.514
8	1.000	.882
9	1.000	.822
10	1.000	.818
11	1.000	.659
12	1.000	.783
13	1.000	.763
14	1.000	.862
15	1.000	.893
16	1.000	.859
17	1.000	.871
18	1.000	.845
19	1.000	.694
20	1.000	.822
21	1.000	.805
22	1.000	.831
23	1.000	.793

24	1.000	.629			
25	1.000	.810			
26	1.000	.747			
27	1.000	.880			
Extraction Method: Principal Component Analysis.					

Appendix-10 : The total variance explained

		Initial Eigenvalue	es .	Ext	raction Sums of Squared	Loadings
Component	Total	% of Variance	Cumulative %	Total	% of Variance	Cumulative %
1	10.247	37.952	37.952	10.247	37.952	37.952
2	2.767	10.248	48.200	2.767	10.248	48.200
3	2.559	9.479	57.679	2.559	9.479	57.679
4	1.856	6.874	64.553	1.856	6.874	64.553
5	1.473	5.456	70.009	1.473	5.456	70.009
6	1.279	4.736	74.745	1.279	4.736	74.745
7	1.143	4.231	78.976	1.143	4.231	78.976
8	.930	3.444	82.420			
9	.873	3.233	85.653			
10	.750	2.778	88.431			
11	.544	2.013	90.444			
12	.436	1.615	92.060			
13	.363	1.344	93.403			
14	.342	1.266	94.669			
15	.324	1.199	95.868			
16	.282	1.045	96.913			
17	.207	.765	97.678			
18	.165	.612	98.290			
19	.123	.457	98.748			
20	.104	.384	99.132			
21	.075	.278	99.410			
22	.054	.202	99.612			
23	.038	.142	99.754			
24	.033	.120	99.874			
25	.021	.079	99.953			
26	.009	.033	99.986			
27	.004	.014	100.000			

Appendix-11: Component Matrix

		C	omponent Mat	rix ^a						
	Component									
Variable number	1	2	3	4	5	6	7			
1	.433	208	412	.197	.498	.062	.146			
2	.636	240	359	038	294	.058	.352			
3	.604	.343	244	.059	373	.055	226			
4	.623	.364	304	.367	192	171	.076			
5	.357	.434	.235	530	.178	.348	.123			
6	.624	.132	367	321	200	394	047			
7	.531	031	.298	058	027	266	260			
8	.770	.016	.286	101	202	.392	.052			
9	.072	.199	.787	.207	166	.252	157			
10	.650	.506	277	.242	020	033	.049			
11	.568	414	.073	.169	.026	.162	.324			

12	.552	.397	075	.453	.126	.254	171
13	.757	203	009	.258	216	.163	096
14	.693	454	152	046	.374	.101	.028
15	.720	332	.109	278	267	.168	273
16	.702	423	307	.047	.079	.237	168
17	.653	.441	.220	213	.320	.223	.069
18	.571	541	.346	067	020	148	.282
19	.627	.335	297	217	208	082	.058
20	.635	.331	.040	456	059	.041	.308
21	.817	246	.169	.185	065	069	074
22	.777	.031	.222	.286	053	257	.166
23	.552	.301	.250	.138	.445	253	234
24	.445	.078	.407	.455	.048	106	.198
25	.573	404	.240	257	057	226	375
26	.530	.130	.375	231	.192	428	.189
27	.678	014	388	178	.380	.035	304
Extraction Method: Principa	l Component Ana	alysis.					

a. 7 components extracted.

Appendix-12: Rotated component matrix

_		Rotat	ed Component				
67 ' 1 1 1	1	2		Component	5		7
Variable number	1	2	3	4	-	6	7
1	.148	.260	113	018	.767	.101	10
2	.405	.652	.164	.115	.184	138	35
3	.776	.030	.322	.173	.010	048	01
4	.844	.193	025	007	.114	.228	07
5	.042	047	.058	.896	.029	.061	.07
6	.516	.059	.373	.195	.091	.194	59
7	.160	.088	.517	.093	.010	.449	.04
8	.314	.499	.421	.526	.042	.028	.28
9	028	.062	.173	.183	409	.209	.73
10	.811	.073	060	.227	.220	.226	00
11	.058	.732	.136	.062	.290	.093	.0.
12	.656	.023	007	.139	.342	.145	.4
13	.465	.509	.441	.005	.233	.042	.19
14	.013	.453	.335	.136	.706	.142	08
15	.170	.371	.802	.252	.129	055	.00
16	.256	.406	.471	.020	.624	128	0
17	.276	.056	.083	.755	.257	.335	.19
18	173	.729	.375	.075	.081	.355	0
19	.619	.102	.190	.382	.064	.071	32
20	.316	.247	.123	.740	015	.185	25
21	.323	.511	.498	.029	.234	.344	.13
22	.446	.499	.202	.071	.073	.571	.0`
23	.261	138	.176	.202	.320	.697	.2
24	.241	.404	041	005	018	.522	.30
25	046	.191	.813	.040	.126	.292	08
26	.033	.188	.180	.370	024	.715	1′
27	.321	057	.380	.256	.722	.109	1
Extraction Method: Pri	ncipal Component	Analysis. Rota	tion Method: V	rimax with Kais	ser Normalizatio	n.	

Appendix 13: Appendix 13: Component Score Coefficient Matrix

				Component			
	1	2	3	4	5	6	7
1	055	.061	201	016	.357	.048	016
2	.078	.298	109	.025	076	150	180
3	.259	100	.150	037	111	141	.031
4	.268	.036	109	130	069	.063	051
5	102	029	055	.435	.027	093	.051
6	.116	090	.123	033	102	.091	340
7	002	131	.227	083	059	.183	008
8	.015	.144	.060	.196	077	190	.176
9	.008	.002	.108	.055	166	013	.374
10	.224	028	127	005	.018	.038	.003

11	068	.317	143	.022	.052	040	.039
12	.182	091	054	023	.147	048	.290
13	.110	.099	.104	091	008	119	.141
14	138	.066	004	.032	.287	.007	003
15	013	018	.334	.040	052	179	.049
16	001	.018	.128	040	.219	174	.076
17	050	052	091	.302	.115	.040	.108
18	158	.278	015	005	067	.133	101

10	136	.276	015	003	007	.133	101	
1. Res. No.				Sex: Father =0	Mother=	:1		
2. Age								
3. Number of total childr	en							
4. Number of baby up to	4 year							
5. No. Son baby up to 4 y	year							
6. No. Daughter baby up	to 4 year							
7. Name								
8. Contact no.								
Res. Occupation				 Business 				
				2. Service				
				3. both B&S				
				4. Others				
10. Annual Income of Fa	ımily							
11. Annual Expenditure	of Family							
12. Education (in years)								
19	.150	016	.008	.088	093	037	177	
20	011	.105	103	.304	106	006	169	
21	.028	.072	.108	106	003	.071	.061	
22	.072	.141	078	099	088	.225	024	
23	013	236	.038	038	.164	.331	.085	
24	.037	.164	159	081	056	.212	.132	
25	086	138	.381	093	019	.097	051	
26	106	.018	051	.077	073	.366	197	
27	014	234	.134	.038	.315	004	021	
Extraction Method: Pr	incipal Component A	nalysis. Rotation	on Method: `	Varimax with Kai	iser Normalizati	on.		

Questionnaire Set: 1:

Baby Eating Behavior up to 4 years

13. How would you describe your baby's feeding style at a day time feed?

Items /variables	1 Never	2 Rarely	3. Sometimes	4. often	5. Always
13.1 My baby seemed contented while feeding					
13.2 My baby frequently wanted more milk than I provided					
13.3 My baby loved milk					
13.4 My baby had a big appetite					
13.5 My baby finished feeding quickly					
13.6 My baby became distressed while feeding					
13.7 My baby got full up easily					
13.8 If allowed to, my baby would take too much milk					
13.9 My baby took more than 30 minutes to finish feeding					
13.10 My baby got full before taking all the milk I think he/she should have					
13.11 My baby fed slowly					
13.12 Even when my baby had just eaten well he/she was happy to feed again if offered					
13.13 My baby found it difficult to manage a complete feed					
13.14 My baby was always demanding a feed					
13.15 My baby sucked more and more slowly during the course of a feed					
13.16 If given the chance, my baby would always be feeding					
13.17 My baby enjoyed feeding time					
13.18 My baby could easily take a feed within 30 minutes of the last one					

Set-2:Child Eating Behavior Questionnaire (5-9 Years)

13. Please read the following statements and tick the boxes most appropriate to your child's eating behavior.

	1. Never	2. Rarely	3. Sometimes	4. often	5. Always
13.1 My child loves food					
13.2 My child eats more when worried					
13.3 My child has big appetite					
13.4 My child finishes his/her meal quickly					
13.5 My child is interested in food					
13.6 My child is always asking for a drink					
13.7 My child refuses new foods at first					
13.8 My child eats slowly					
13.9 My child eats less when angry					
13.10 My child enjoys tasting new foods					
13.11 My child eats less when s/he is tired					
13.12 My child is always asking for food					
13.13 My child eats more when annoyed					
13.14 If allowed to, my child would eat too much					
13.15 My child eats more when anxious					
13.16 My child enjoys a wide variety of foods					
13.17 My child leaves food on his/her plate at the end of a meal					
13.18 My child takes more than 30 minutes to finish a meal					
13.19 Given the choice, my child would eat most of the time					
13.20 My child looks forward to mealtimes					
13.21 My child gets full before his/her meal is finished					
13.22 My child enjoys eating					
13.23 My child eats more when s/he is happy					
13.24 My child is difficult to please with meals					
13.25 My child eats less when upset					
13.26 My child gets full up easily					
13.27 My child eats more when s/he has nothing else to do					
13.28 Even if child is full finds room to eat his/her favorite food					
13.29 If given the chance, drink continuously throughout the day					
13.30 Child cannot eat a meal if s/he has had a snack just before					
13.31 if given the chance my child would always be having a drink					
13.32 My child is interested in tasting food s/he hasn't tasted before					
13.33 Child decides that s/he doesn't like a food, without tasting					
13.34 If given the chance, child always have food in his/her mouth					
13.35 My child eats more and more slowly the course of a meal					

Adolescent Food Habits (10-19 years)

D. M		
Res. No.		
1. Sex	M	
1. Sex	M	0
2 D A	F	1
2. Res. Age		
3. Schooling in years		
4. Father's Schooling in years		
5. Mother's Schooling in years		
5. Women's Schooling in Jeans		
6. Res. Name		
7. Father's Profession	1.Service	
7. Father 8 Froression	2.Business	
	3.Both	
	4.Others	
8. Mother's Profession	1.House	
8. Mother's Profession	2.Service	
	3.Both	
	4.Business	
	5.Others	
9. Annual Income of Family		
10. Annual Expenditure of Family		
11. Res. Contact no.		
11. Res. Contact no.		

		Most	Neutral=0	like=1
		like=2		
12.1_lowfat	If I am having lunch away from home, I often choose a low-fat option.			
12.2_eating	I usually avoid eating fried foods.			
12.3_dessert	I usually eat a dessert or pudding if there is one available.			
12.4_least	I make sure I eat at least one serving of fruit a day.			
12.5_fat	I try to keep my overall fat intake down.			
12.6_crisps	If I am buying crisps, I often choose a low-fat brand.			
12.7_sausages	I avoid eating lots of sausages and burgers.			
12.8_pastries	I often buy pastries or cakes.			
12.9_sugar	I try to keep my overall sugar intake down.			

12.10_vegetables	I make sure I eat at least one serving of vegetables or salad a day.		
12.11_home	If I am having a dessert at home, I try to have something low in fat.		
12.12_meals	I rarely takeaway meals.		
12.13_fruit	I try to ensure that I eat plenty of fruit and vegetables.		
12.14_sweet	I often eat sweet snacks between meals.		

Res. No.	1. Sex	M	0	
		F	1	
2. Occ_sp	1.Household	•	-	
	2. Service			
	3. both service & household	1		
	4. Business			
	5. Others			
3. Res. Age				
4. no_tc				
5. occ. respondance	1. Service			
_	2. Business			
	3. both Service & Business			
	4. Others			
6. No. of total child up to 9				
7. No. of Son child up to 9				
8. No. of daughter child up to 9				
9. Res. Education years				

12.15_salad	I usually eat at least one serving of vegetables (excluding potatoes) or salad with my		
	evening meal.		
12.16_softdrink	When I am buying a soft drink, I usually choose a diet drink.		
12.17_butter	When I put butter or margarine on bread, I usually spread it on thinly.		
12.18_chocolate	If I have a packed lunch, I usually include some chocolate and/or biscuits.		
12.19_snack	When I have a snack between meals, I often choose fruit.		
12.20_healthiest	If I am having a dessert or pudding in a restaurant, I usually choose the healthiest one.		
12.21_cream	I often have cream on desserts.		
12.22_servings	I eat at least 3 servings of fruits most days.		
12.23_diet	I generally try to have a healthy diet.		

Parental Feeding Style to his Children up to 9 years-

10. Annual Income	
11. Annual Expenditure	
12. Education of spouse	
13. Conjugal life	

Items/variable		1.Never	2.Rarely	3.Someti mes	4.Often	5.Always
I allow my child to choose which foods to have for meals	14.1 choose					
I give my child something to eat to make him/her feel better when s/he is feeling upset	14.2 upset					
I encourage my child to look forward to the meal	14.3 encourage					
I praise my child if s/he eats what I give him/her	14.4 praise					
I decide how many snacks my child should have	14.5 decide					
I encourage my child to eat a wide variety of foods	14.6 variety					
In order to get my child to behave him/herself I promise him/her something to eat	14.7 order					
I present food in an attractive way to my child	14.8 present					
If my child misbehaves I withhold his/her favorite food	14.9 misbehaves					
I encourage my child to taste each of the foods I serve at mealtimes	14.10 taste					
I allow my child to wander around during a meal	14.11 wander					
I encourage my child to try foods that s/he hasn't tasted before	14.12 try					
I give my child something to eat to make him/herfeel better when s/he has been hurt	14.13 feel					
I let my child decide when s/he would like to have her meal	14.14 like					
I give my child something to eat if s/he is feeling bored	14.15 bored					
I allow my child to decide when s/he has had enough snacks to eat	14.16 enough					
I decide when it is time for my child to have a snack	14.17 time					
I use puddings as a bribe to get my child to eat his/her main course	14.18 course					
I encourage my child to enjoy his/her food	14.19 enjoy					
I decide the times when my child eats his/her meals	14.20 meals					
I give my child something to eat if s/he is feeling worried	14.21 btr_wry					
I reward my child with something to eat when s/he is well behaved	14.22 reward					
I let my child eat between meals whenever s/he wants	14.23 between					
I insist my child eats meals at the table	14.24 insist					
I give my child something to eat to make him/her feel better when s/he is feeling angry	14.25 angry					
I decide what my child eats between meals	14.26 eats					
I praise my child if s/he eats a new food	14.27 new_food					

Food preference questionnaire for adults (20-29 unmarried) $\,$

9.

Food Item	1 Dislike a	2. Dislike a	3. Neither like	4. Like a little	5. Like a
	lot	little	or dislike		lot
9.1 Beef					
9.2 Beef burgers					
9.3 Lamb					
9.4 Chicken					
9.5 Bacon					
9.6 Ham					

9.7 Sausages			
9.8 White fish (e.g. cod, haddock)			
9.9 Oily fish (e.g. mackerel, kippers)			
9.10 Smoked Salmon			
9.11 Tinned Tuna			
9.12 Eggs (boiled, scrambled or fried)			
9.13 Baked beans			
9.14 Bread or Bread rolls			
9.15 Bran cereal (e.g. All Bran, Bran Flakes)			
9.16 Porridge			
9.17 Plain boiled rice			
9.18 Sugared cereal (e.g. Frosties, Sugar			
Puffs)			
9.19 Hummus			
9.20 Wheat cereal (e.g. Weetbix, Shredded			
Wheat)			
9.21 Potatoes (boiled or mashed)			
9.22 Chips			
9.23 Rice or corn cereal (e.g. corn Flakes, Rice	+		
(e.g. com Plakes, Rice Krispies)			
9.24 Soft cheese (e.g. Camembert, Brie)			
9.25 Hard cheese (e.g. cheddar)	1		
9.25 Hard cheese (e.g. cheddar) 9.26 Cottage Cheese	 		
9.26 Cottage Cheese 9.27 Plain, Low-fat yoghurt	+		
9.28 Oranges			
9.29 Grapes			
9.30 Apples			
9.31 Melon			
9.32 Peaches			
9.33 Apricots			
9.34 Strawberries			
9.35 Avocadoes			
9.36 Spinach			
9.37 Carrots			
9.38 Green beans			
9.39 Cucumber			
9.40 Celery			
9.41 Mushrooms			
9.42 Parsnips			
9.43 Peas			
9.44 Sweetcorn			
9.45 Broccoli			
9.46 Salad leaves (e.g. lettuce)			
9.47 Red peppers			
9.48 Raw tomatoes			
9.49 Beetroot			
9.50 Brussels sprouts		 	
9.51 Butter		 	
9.52 Butter-like spreads (e.g. sunflower		 · ·	
spread, Flora)		 	
9.53 Cream		 	
9.54 Mayonnaise			
9.55 Plain biscuits (e.g. Digestives)			
9.56 Chocolate biscuits			
9.57 Cake			
9.58 Ice cream			
9.59 Custard			
9.60 Chocolate			
9.61 Crisps			
9.62 Chewy gummy sweets (e.g. Haribo-style			
sweets, wine gums)			
/	1		

Food preference question naire for adults (19-64) Married 16.

Food Item	1 Dislike a	Dislike a	Neither like	Like a little	Like a	l
-----------	-------------	-----------------------------	--------------------------------	---------------------------------	--------------------------	---

	Lot	little	or dislike	I	lot
16.1 Beef	lot	nttie	or distike		101
16.2 Beef burgers					
16.3 Lamb					
16.4 Chicken					
16.5 Bacon					
16.6 Ham					
16.7 Sausages					
16.8 White fish (e.g. cod, haddock)					
16.9 Oily fish (e.g. mackerel, kippers)					
16.10 Smoked Salmon					
16.11 Tinned Tuna 16.12 Eggs (boiled, scrambled or fried)					
16.13 Baked beans					
16.14 Bread or Bread rolls					
16.15 Bran cereal (e.g. All Bran, Bran Flakes)					
16.16 Porridge					
16.17 Plain boiled rice					
16.18 Sugared cereal (e.g. Frosties, Sugar					
Puffs)					
16.19 Hummus					
16.20 Wheat cereal (e.g. Weetbix, Shredded					
Wheat)					
16.21 Potatoes (boiled or mashed) 16.22 Chips	-	-			
16.22 Cmps 16.23 Rice or corn cereal (e.g. corn Flakes,					
Rice Krispies)					
16.24 Soft cheese (e.g. Camembert, Brie)					
16.25 Hard cheese (e.g. cheddar)					
16.26 Cottage Cheese					
16.27 Plain, Low-fat yoghurt					
16.28 Oranges					
16.29 Grapes					
16.30 Apples					
16.31 Melon					
16.32 Peaches					
16.33 Apricots 16.34 Strawberries					
16.35 Avocadoes					
16.36 Spinach					
16.37 Carrots					
16.38 Green beans					
16.39 Cucumber					
16.40 Celery					
16.41 Mushrooms					
16.42 Parsnips					
16.43 Peas					
16.44 Sweetcorn					
16.45 Broccoli					
16.46 Salad leaves (e.g. lettuce)					
16.47 Red peppers					
16.48 Raw tomatoes 16.49 Beetroot	1				
16.50 Brussels sprouts	+				
16.51 Butter					
16.52 Butter-like spreads (e.g. sunflower					
spread, Flora)					
16.53 Cream					
16.54 Mayonnaise					
16.55 Plain biscuits (e.g. Digestives)					
16.56 Chocolate biscuits					
16.57 Cake					
16.58 Ice cream	-				
16.59 Custard	-				
16.60 Chocolate	 				
16.61 Crisps		1			

16.62 Chewy gummy sweets (e.g. Haribo-style			
sweets, wine gums)			