

Verdier duality on conically smooth stratified spaces

Marco Volpe*

June 7, 2022

Abstract

In this paper we prove a duality for constructible sheaves on conically smooth stratified spaces. Here we consider sheaves with values in a stable and bicomplete ∞ -category equipped with a closed symmetric monoidal structure, and in this setting constructible means locally constant along strata and with dualizable stalks. The crucial point where we need to employ the geometry of conically smooth structures is in showing that Lurie’s version of Verdier duality restricts to an equivalence between constructible sheaves and cosheaves: this requires a computation of the exit path ∞ -category of a compact stratified space, that we obtain via resolution of singularities.

Contents

1	Introduction	1
1.1	Linear overview	4
1.2	Acknowledgements	4
2	Finite exit paths	5
2.1	Finite ∞ -categories	5
2.2	Finiteness properties of compact conically smooth spaces	7
3	Homotopy invariance and exodromy with general coefficients	10
3.1	Homotopy invariance of constructible sheaves	11
3.2	Exodromy	14
4	The duality	17

1 Introduction

Constructible sheaves are of great interest both in algebraic and differential geometry, as they provide tools to study invariants for singular spaces (such as *intersection cohomology*, following the approach of [BBDG18]) and have surprising and beautiful relations with D-modules (see [Kas84]). A fundamental feature of constructible sheaves is that, assuming a certain kind of finiteness properties on the stalks and considering nicely behaved stratifications, they carry a duality (sometimes referred to as *Verdier duality*) which enables one, through abstract trace methods, to associate to any such sheaf a class in Borel–Moore homology. One of the interests of these classes is that they can be related to Euler characteristics via routine computations

*University of Regensburg, Universitätsstraße 31, Regensburg. email: marco.volpe@ur.de. The author was supported by the SFB 1085 (Higher Invariants) in Regensburg, Germany, funded by the DFG (German Science Foundation).

with the six functor formalism (see [KS90, Chapter 9] for a discussion on classical index formulas and their microlocal enhancements).

As far the author knows, it was an idea of MacPherson that the duality should actually be thought of as a combination of two different equivalences of categories. The former, given by the usual construction of sections with compact support, was expected to identify constructible sheaves with constructible *cosheaves* (without constructibility assumptions, this was proven by Lurie in [Lur17, Theorem 5.5.5.1], and there named after Verdier); the latter maps back contravariantly constructible cosheaves to sheaves, and obtained using a foreseen combinatorial description of constructible (co)sheaves, similar in spirit to the monodromy for local systems, which is nowadays referred to as *exodromy* (see [Lur17, Theorem A.9.3], [BGH18]). In this paper we make use of the language of ∞ -categories to realize the vision of MacPherson and prove the expected duality result in a very general setting.

Let us spend a few words to specify more precisely the framework in which we are working. Relying on the previous paper [Vol21], we will be able to deal with sheaves valued in any stable bicomplete ∞ -category \mathcal{C} , equipped with a closed symmetric monoidal structure. The machinery of six functors developed in [Vol21] supplies us with a *dualizing sheaf* $\omega_X^{\mathcal{C}}$ for any \mathcal{C} as above and X locally compact Hausdorff stratified space. Our duality functor will thus be defined as usual by taking an internal-hom into $\omega_X^{\mathcal{C}}$, and denoted by $D_X^{\mathcal{C}}$.

Following the nomenclature of [BGH18], we will define a sheaf with values in \mathcal{C} to be *formally constructible*¹ if its restriction to each stratum is locally constant, and *constructible* if furthermore all its stalks are dualizable. Similar definitions can be given for \mathcal{C} -valued cosheaves by observing that, up to passing to an opposite category, these are just \mathcal{C}^{op} -valued sheaves. The requirement of dualizability for stalks is really unavoidable mainly because, when X has a unique point, we see that $\omega_X^{\mathcal{C}}$ is the monoidal unit of \mathcal{C} , and thus the duality functor coincides with the one coming from the monoidal structure on \mathcal{C} . Moreover it is also extremely reasonable, since if $\mathcal{C} = D(R)$ where R is a ring (or more generally, modules over any E_∞ -ring spectrum), it is well-known that a complex is dualizable if and only if it is perfect, and hence our assumptions allow us to recover the classical setting as a special case.

For the geometric side of the story, we will consider *conically smooth stratified spaces*. These were introduced by Ayala, Francis and Tanaka in [AFT17], and provide a natural extension of C^∞ -structures in the stratified/singular setting. The definition of conically smooth atlases is rather involved, as it relies on an elaborate inductive construction based on the *depth*² of a stratification, but we suggest the reader to have a look at the introductions of [AFT17] or [NV21] to get an idea of how this works.

The main feature of conically smooth structures we will use in this paper is *unzip* construction, that allows one to functorially resolve any conically smooth stratified space into a manifold with corners. We will give a brief explanation of how this works later in the paper, but for now let us only mention that, if $X_k \hookrightarrow X$ is the inclusion of a stratum of maximal depth, it consists of a square

¹Notice that in this paper we will only deal with sheaves which are constructible with respect to a fixed stratification, as opposed to [KS90, Chapter 8], for example.

²Recall that, for a stratified space $s : X \rightarrow P$, the depth is defined as

$$\text{depth}(X) = \sup_{x \in X} \dim_x(X) - \dim_x(X_{s(x)}),$$

where \dim denotes the covering dimension and $X_{s(x)}$ is the stratum of X corresponding to $s(x) \in P$.

$$(1.1) \quad \begin{array}{ccc} \text{Link}_k(X) & \hookrightarrow & \text{Unzip}_k(X) \\ \downarrow \pi_X & \lrcorner & \downarrow \\ X_k & \hookrightarrow & X \end{array}$$

which is both pushout and pullback, and $\text{Unzip}_k(X)$ is a conically smooth manifold *boundary* given by $\text{Link}_k(X)$ such that both its interior and $\text{Link}_k(X)$ have depth strictly smaller than the one of X . An interesting consequence of the existence of pushout/pullback square (1.1) is that the notion of conically smooth map is completely determined by the one of smooth maps between manifolds with corners.

We are now ready to state our main result.

Theorem 1.2. Let X be a conically smooth stratified space, and let $\text{Shv}^c(X; \mathcal{C})$ be the full subcategory of $\text{Shv}(X; \mathcal{C})$ spanned by constructible sheaves. Then the restriction to $\text{Shv}^c(X; \mathcal{C})^{op}$ of the functor $D_X^{\mathcal{C}}$ factors through an equivalence

$$D_X^{\mathcal{C}} : \text{Shv}^c(X; \mathcal{C})^{op} \xrightarrow{\simeq} \text{Shv}^c(X; \mathcal{C}).$$

To conclude this introduction, let us make a short comment on how our proof strategy goes. Our first observation, as anticipated above, is that the functor $D_X^{\mathcal{C}}$ factors through Lurie's equivalence

$$\mathbb{D}_{\mathcal{C}} : \text{Shv}(X; \mathcal{C}) \rightarrow \text{CoShv}(X; \mathcal{C}),$$

and most of the work then lies in proving that the restriction of $\mathbb{D}_{\mathcal{C}}$ to constructible sheaves factors through constructible cosheaves. We first show in Proposition 4.1 that $\omega_X^{\mathcal{C}}$ is constructible when $\mathcal{C} = \text{Sp}$ (the ∞ -category of spectra), and from the techniques developed in [Vol21] we deduce immediately that

$$a_{\mathcal{C}}^{\dagger} : \mathcal{C} \rightarrow \text{Shv}(X; \mathcal{C})$$

factors through formally constructible sheaves. As a consequence of this and some easy properties of constructible sheaves that follow from *homotopy invariance* (see Theorem 3.3), one gets that $\mathbb{D}_{\mathcal{C}}$ maps formally constructible sheaves into formally constructible cosheaves. We want to stress that the possibility of working with such a general class of coefficients, which is closed under passing to opposite categories, makes this step extremely formal.

The missing piece then consists in showing that $\mathbb{D}_{\mathcal{C}}$ preserves the property of having dualizable stalks, and this is the point where we will actually have to employ the geometry of conically smooth structures. More specifically, we will use the unzip construction to prove in Proposition 2.12, through an inductive argument on the depth, that any compact stratified space equipped with a conically smooth structure has a finite exit path ∞ -category. The key observation then is that, if $X = C(Z)$ with Z compact, $x \in X$ is the cone point and F is any constructible sheaf on X , then there is a fiber sequence

$$(1.3) \quad \Gamma_x(X; F) \rightarrow F_x \rightarrow \Gamma(Z; F),$$

where $\Gamma_x(X; F)$ denotes the sections of F supported at x (i.e. the stalk of the associated cosheaf of compactly supported sections of F) and F_x is the stalk of F at x . By Proposition 2.12 and the exodromy equivalence (which will be show to hold also for our general class of coefficients in Theorem 3.14), one gets that $\Gamma(Z; F)$ must be dualizable, and thus F_x is dualizable if and only if $\Gamma_x(X; F)$ is, which proves our claim.

1.1 Linear overview

We now give a linear overview of the results in our paper.

Section 2 will be mainly devoted to prove Proposition 2.12. In the first part we will recall the definition of a finite ∞ -category, and show how these can be described in the model of quasi-categories. None of these results or definitions are new, but we decided to include a few words on the subject since we could not find any reference dealing with it in our preferred fashion. In the second part we recall Lurie’s definition of the simplicial set of exit-paths of a stratified topological space, and show in Lemma 2.7 how one can conveniently compute the exit-paths of the cone of a proper stratified fiber bundle $L \rightarrow X$ in terms of L and X . The main idea for proving Proposition 2.12 is then to cover a conically smooth stratified space X into the open subset U given by the locus of points of depth zero and a tubular neighbourhood of the complement of U . By induction and Lemma 2.7 one is then left to show that $\text{Exit}(U)$ is finite, but this follows by observing that, through the unzip construction, U can be identified with the interior of compact manifold with corners.

In Section 3 we deal with extending the results of [HPT20] to sheaves values in stable bicomplete ∞ -categories. This is very simple and formal, after [Vol21]. As a consequence, we show in Corollary 3.5 that the stalk at a point x of a constructible sheaf is the same as sections at any conical chart around x , and compute in Corollary 3.8 the restriction of a constructible sheaf along a stratum in a convenient way. These two results are essential and are used very often in what follows (for example, the first implies immediately the existence of (1.3)). In the second part we will then characterize constructible sheaves by the property of being homotopy invariant (see Proposition 3.12), and use this to deduce exodromy for conically smooth spaces with general stable bicomplete coefficients.

Section 4 deals with proving our main result. We start by showing, through an inductive argument on the depth, that $\omega_X^{\mathcal{C}}$ is constructible whenever X is C^0 -stratified. We first reduce to proving the statement in the case $\mathcal{C} = \text{Sp}$ by employing the techniques developed in [Vol21], and then the only non-trivial part consists in showing that, when $X = C(Z)$, the stalk of the dualizing sheaf at the cone-point is a finite spectrum. We then conclude by proving Theorem 4.2. As explained at the beginning of the introduction, our argument starts by observing that the duality functor factors through Lurie’s Verdier duality, and the hard part then consists in showing that the latter restricts to an equivalence between constructible (co)sheaves, for which we have to implement all the results obtained previously in the paper.

1.2 Acknowledgements

I owe my interest and fascination towards the theory of conically smooth stratified spaces to my supervisor Denis-Charles Cisinski: the enthusiasm he showed while explaining to me the main ideas of [AFR18] gave me the motivation to start reading [AFT17], which incidentally led to the birth of this paper and my collaboration with Guglielmo Nocera [NV21].

Of course, the second person I would like to thank is Guglielmo himself. The hours of discussions we’ve had through zoom during early pandemic times, and later in Regensburg while he stayed here for a visit, have helped me to sharpen my intuition on stratified spaces. The importance of our friendship too cannot be underestimated.

I’d also like to thank Benedikt Preis for asking me, two years ago, if it was possible to deduce the duality for constructible sheaves from Lurie’s Verdier duality; George Raptis for his support, especially one year ago during my self-imposed exile in Italy, and for being a constant reminder of the importance of learning also some geometry, every once in a while; Mauro Porta, Jean-Baptiste Teyssier and Peter Haine for sharing earlier drafts of their papers on homotopy invariance and exodromy with me, and for long and stimulating discussions

during my stay in Paris and through zoom; Denis Nardin, Andrea Gagna, Edoardo Lanari, David Ayala, Niklas Kipp, Sebastian Wolf and Han-Ung Kufner for answering (sometimes silly) questions related to the subject of this paper, and for constantly providing stimulating math conversations.

But most of all, I would like to thank the ghost of Pina Tumminello, who quietly slipped through the cracking wooden door of our room, caressed us gently in our tormented sleeps, and slowly pushed inside our ears the dreams of Scott Gilbert and Robert MacPherson; who shouted, through the mouth of a young girl, the corny, dull (but long forgotten) word "love", at the heart of a world that was desperate, wounded and fallen on its knees – ambulance bells sighing all around us and no trace of warmth, no wine sold after 6 pm – and made it shine once again, "but for less than a moment": a precious, indelible, game changing moment.

This paper will be a part of my PhD thesis.

2 Finite exit paths

This first section contains the main geometric input needed to achieve our goal. Namely, we show that a compact stratified space has a finite stratified homotopy type, as soon as it can be equipped with a conically smooth structure.

2.1 Finite ∞ -categories

This short section will be devoted to recalling the definition of finite ∞ -categories, but before going into that we spend a few words on specifying what an ∞ -category is for us. In this paper we work in the model of quasicategories and, following [Cis19, Example 7.10.14], we define Cat_∞ as the localization of the 1-category of simplicial sets at the class of Joyal equivalences. The class of Joyal equivalences and fibrations equip sSet with the structure of a category with weak equivalences and fibrations in the sense of [Cis19, Definition 7.4.12], and so by [Cis19, Theorem 7.5.18] it follows that any object in Cat_∞ is equivalent to the image through the localization functor $\text{sSet} \rightarrow \text{Cat}_\infty$ of a fibrant object in the Joyal model structure. For this reason, objects of Cat_∞ will be called ∞ -categories.

The first definition we propose is expressed internally to the ∞ -category Cat_∞ in terms of pushouts (and so in a kind of model-independent fashion), and later we will prove that this is actually equivalent to a notion of finiteness that one might expect in the simplicial model. All the results appearing here are not at all original, but we still felt the need to write this section as, in the process of completing the paper, we struggled a lot in trying to find a unique reference dealing with the subject.

Definition 2.1. An ∞ -category is said to be *finite* if it belongs to the smallest full subcategory of Cat_∞ which contains \emptyset , Δ^0 and Δ^1 and is closed under pushouts. An ∞ -groupoid is said to be finite if it is so as an ∞ -category. We will denote by Cat_∞^f and \mathcal{S}^f respectively the full subcategories of Cat_∞ and \mathcal{S} spanned by finite objects.

Remark 2.2. Recall that the inclusion $\mathcal{S} \hookrightarrow \text{Cat}_\infty$ admits both a left and a right adjoint, and one may describe the left adjoint on objects by sending an ∞ -category \mathcal{C} to the localization $\mathcal{C}[\mathcal{C}^{-1}]$. Thus, since $\mathcal{S} \hookrightarrow \text{Cat}_\infty$ preserves colimits and $\Delta^1[(\Delta^1)^{-1}] \simeq \Delta^0$, one may identify the class of finite ∞ -groupoids with the objects of the smallest full subcategory of \mathcal{S} which contains \emptyset and Δ^0 and it's closed under pushouts. This implies in particular that, for any finite ∞ -category \mathcal{C} , $\mathcal{C}[\mathcal{C}^{-1}]$ is again finite.

Lemma 2.3. Let \mathcal{C} be a finite ∞ -category, and let W be a finite subcategory of \mathcal{C} . Then the localization $\mathcal{C}[W^{-1}]$ is again finite.

Proof. We have a pushout

$$\begin{array}{ccc} W & \longrightarrow & \mathcal{C} \\ \downarrow & \lrcorner & \downarrow \\ W[W^{-1}] & \longrightarrow & \mathcal{C}[W^{-1}], \end{array}$$

in Cat_∞ , and thus it suffices to show that $W[W^{-1}]$ is finite, which follows immediately by Remark 2.2. \square

Recall that a simplicial set is said to be finite if it has a finite number of non-degenerate simplices. In the next proposition we reconcile this notion of finiteness with one in Definition 2.1.

Proposition 2.4. An ∞ -category \mathcal{C} is finite if and only if there exists a finite simplicial set K and an equivalence $\mathcal{C} \simeq L(K)$, where $L : \text{sSet} \rightarrow \text{Cat}_\infty$ is the localization functor.

Proof. Let sSet^f be the full subcategory of sSet spanned by the finite ones, and denote by \mathcal{F} the essential image of the restriction of L to sSet^f . We need to show that Cat_∞^f coincides with \mathcal{F} . We will show this by identifying both \mathcal{F} and Cat_∞^f with the localization of sSet^f at the class of Joyal equivalences.

As usual, one can make sSet into a category of cofibrant objects ([Cis19, Definition 7.5.7]) by equipping it with the classes of Joyal equivalences and monomorphisms. We verify that this structure restricts to sSet^f . Since sSet^f is closed under taking pushouts in sSet , the only non-trivial thing to check is that any morphism $f : K \rightarrow L$ of finite simplicial sets can be factored into a monomorphism followed by a Joyal equivalence, which follows by the fact that one may choose an interval object J for the Joyal model structure which belongs to Cat_∞^f (as in [Cis19, Definition 3.3.3]) and by forming as usual the mapping cylinder of f . Thus we get that the functor $\text{sSet}^f \hookrightarrow \text{sSet}$ is right exact in the sense of [Cis19, Definition 7.5.2] and so, if we denote by $L(\text{sSet}^f)$ the localization of sSet^f at the class of Joyal equivalences, by [Cis19, Proposition 7.5.6] we see that $L(\text{sSet}^f)$ has finite colimits and we have a commutative square

$$\begin{array}{ccc} \text{sSet}^f & \hookrightarrow & \text{sSet} \\ \downarrow & & \downarrow \\ L(\text{sSet}^f) & \longrightarrow & \text{Cat}_\infty \end{array}$$

where all functors preserve finite colimits. Moreover the existence of a maximal left calculus of fractions for both sSet^f and sSet implies, by [Cis19, Theorem 7.2.8], that the lower horizontal arrow is fully faithful, and hence we have $\mathcal{F} \simeq L(\text{sSet}^f)$, since both vertical arrows are essentially surjective.

By what we have shown above, \mathcal{F} is a full subcategory of Cat_∞ which is stable under pushouts, and since it contains \emptyset , Δ^0 and Δ^1 , we see that $\text{Cat}_\infty^f \subseteq \mathcal{F}$. For the other inclusion, let K be any finite simplicial set, so that in particular there exists some finite n such that $K = sk_n(K)$. By making an inductive argument on n and using the cellular decomposition

$$\begin{array}{ccc} \coprod_{\partial\Delta^n \rightarrow K} \partial\Delta^n & \longrightarrow & sk_{n-1}(K) \\ \downarrow & \lrcorner & \downarrow \\ \coprod_{\Delta^n \rightarrow K} \Delta^n & \longrightarrow & sk_n(K) \end{array}$$

we see that to prove that $L(K)$ belongs to Cat_∞^f it suffices to show that each Δ^n does, but this is clear because n -simplex is Joyal equivalent to the n -spine. \square

2.2 Finiteness properties of compact conically smooth spaces

The main goal of this section will be to show that the exit path ∞ -category of a compact conically smooth stratified space is finite. For this purpose, we will mainly make use of Lurie's model of the exit path ∞ -category, of which we now recall the definition for the reader's convenience. By a slight abuse of notation, for a poset P we will still denote by P the topological space obtained by equipping the poset with the Alexandroff topology. If $X \rightarrow P$ is a stratified topological space, then we define $\text{Exit}_P(X)$ by forming the pullback

$$\begin{array}{ccc} \text{Exit}_P(X) & \longrightarrow & \text{Sing}(X) \\ \downarrow & \lrcorner & \downarrow \\ N(P) & \longrightarrow & \text{Sing}(P) \end{array}$$

in the category of simplicial sets. When $X \rightarrow P$ is conically stratified, then Lurie shows ([Lur17, Theorem A.6.4]) that the simplicial set $\text{Exit}_P(X)$ is an ∞ -category. The reason why we prefer to focus on this perspective is that, as opposed to the model given in [AFT17], it has an evident much richer functoriality: by the functoriality of Sing , Exit is functorial with respect to general stratified maps, while the one in [AFT17] is only functorial with respect to conically smooth open embeddings. We also see immediately that, if we stratify P over itself through the identity, then $\text{Exit}_P(P) = N(P)$. We will also need the following easy lemma.

Lemma 2.5. Let $X \rightarrow P$ and $Y \rightarrow Q$ be stratified spaces such that $\text{Exit}_P(X)$ is an ∞ -category. Assume that we have a stratified embedding $Y \hookrightarrow X$ which is a consecutive inclusion of strata. Then $\text{Exit}_Q(Y)$ is an ∞ -category and the induced functor $\text{Exit}_Q(Y) \rightarrow \text{Exit}_P(X)$ is fully faithful.

Proof. Since the functor Sing from topological spaces to simplicial sets preserves limits and since $Y \hookrightarrow X$ is inclusion of strata, we get a pullback square

$$\begin{array}{ccc} \text{Exit}_Q(Y) & \longrightarrow & \text{Exit}_P(X) \\ \downarrow & \lrcorner & \downarrow \\ N(Q) & \longrightarrow & N(P) \end{array}$$

of simplicial sets, and moreover, since the inclusion is consecutive, we know that the functor $N(Q) \rightarrow N(P)$ is fully faithful. To prove the desired conclusion it thus suffices to show that the right vertical map in the square is a fibration in the Joyal model structure, but this follows immediately by the assumption that $\text{Exit}_P(X)$ is fibrant and P is a poset. \square

Recall that, for a proper conically smooth fiber bundle $\pi : L \rightarrow X$, we define the *fiberwise cone* of π as the pushout

$$(2.6) \quad \begin{array}{ccc} L & \longrightarrow & L \times \mathbb{R}_{\geq 0} \\ \downarrow & \lrcorner & \downarrow \\ X & \longrightarrow & C(\pi) \end{array}$$

taken in the category of conically smooth stratified spaces. By definition, we get a new fiber bundle $C(\pi) \rightarrow X$ whose fibers are isomorphic to basiscs. We now show how to compute the exit paths of $C(\pi)$ in terms of L and X .

Lemma 2.7. The commutative square

$$\begin{array}{ccc} \text{Exit}(L) & \longrightarrow & \text{Exit}(L \times \mathbb{R}_{\geq 0}) \\ \downarrow & & \downarrow \\ \text{Exit}(X) & \longrightarrow & \text{Exit}(C(\pi)) \end{array}$$

in Cat_∞ induced by (2.6) is a pushout.

Proof. By Van Kampen theorem for exit paths, we may assume that X is a basic and thus, by [AFT17, Corollary 7.1.4], we may also assume that π is a trivial bundle. Hence, since Exit commutes with finite products, we may assume that $X = *$, and hence we are only left to prove that the square

$$\begin{array}{ccc} \text{Exit}(L) & \longrightarrow & \text{Exit}(L \times \mathbb{R}_{\geq 0}) \\ \downarrow & & \downarrow \\ \Delta^0 & \longrightarrow & \text{Exit}(C(L)) \end{array}$$

is a pushout in Cat_∞ for any compact conically smooth L . This follows for example by [AFT17, Lemma 6.1.4], or can be easily checked by hand even without any conically smooth assumption on L . \square

We will also need to use the *unzip* and *link* construction, as defined in [AFT17, Definition 7.3.11]: for any proper constructible embedding $X \hookrightarrow Y$ we have a pullback square

$$(2.8) \quad \begin{array}{ccc} \text{Link}_X(Y) & \hookrightarrow & \text{Unzip}_X(Y) \\ \downarrow \pi_X & \lrcorner & \downarrow \\ X & \hookrightarrow & Y \end{array}$$

where $\text{Unzip}_X(Y)$ is a conically smooth manifold with corners whose interior is identified with $Y \setminus X$, and $\text{Unzip}_X(Y) \rightarrow Y$ and $\pi_X : \text{Link}_X(Y) \rightarrow X$ are proper constructible bundles. To get a feeling of how $\text{Unzip}_X(Y)$ works, one may think of it as a generalization of the spherical blow-up (see [AK10]): more precisely, when Y is a smooth manifold stratified with a closed submanifold X and its open complement, then the unzip of $X \hookrightarrow Y$ coincides with the spherical blow-up of X in Y , and the link is diffeomorphic to the boundary of any normalized tubular neighbourhood of X in Y .

The link of a proper constructible embedding is used to provide tubular neighbourhoods in the stratified setting. In [AFT17, Proposition 8.2.5], the authors show that there is a conically smooth embedding

$$(2.9) \quad C(\pi_X) \hookrightarrow Y$$

under X which is a refinement onto its image and whose image is open in Y (here we are using the same notations as in (2.8)). Hence, if we denote by $\widetilde{\text{Link}_X(Y)} \times \mathbb{R}_{>0}$ and $\widetilde{C(\pi_X)}$ the respective refinements of $\text{Link}_X(Y) \times \mathbb{R}_{>0}$ and $C(\pi_X)$ through the embedding (2.9), a straightforward application of Van Kampen theorem gives the following.

Corollary 2.10. The square

$$\begin{array}{ccc} \text{Exit}(\widetilde{\text{Link}_X(Y)} \times \mathbb{R}_{>0}) & \longrightarrow & \text{Exit}(Y \setminus X) \\ \downarrow & & \downarrow \\ \text{Exit}(\widetilde{C(\pi_X)}) & \longrightarrow & \text{Exit}(Y) \end{array}$$

is a pushout in Cat_∞ .

Remark 2.11. Notice that, for the existence of tubular neighbourhoods, one may relax the assumption of properness for a constructible embedding $i : X \hookrightarrow Y$ to just requiring that there is a factorization

$$\begin{array}{ccc} & Y' & \\ i' \nearrow & & \nwarrow j \\ X & \xrightarrow{i} & Y \end{array}$$

where i' is a proper constructible embedding and j is a conically smooth open embedding. For example, if P is the stratifying poset of Y and $X = Y_\alpha$ for some $\alpha \in P$, one may pick $Y' = Y_{\geq \alpha}$ and thus get a tubular neighbourhood of Y_p .

We are now ready to prove the main result of the section.

Proposition 2.12. Let X be any compact conically smooth stratified space. Then $\text{Exit}(X)$ is a finite ∞ -category.

Proof. Since X is compact, it follows that it must be finite dimensional, and hence also has finite depth. We then argue by induction on $\text{depth}(X) = k$.

If $k = 0$, it is well known that that $\text{Exit}(X) = \text{Sing}(X)$ is a finite ∞ -groupoid: for example, this follows by Van Kampen theorem and the existence of finite good covers for X .

Assume now that k is positive. Denote by X_0 the union of strata of minimal depth, and by $X_{>0}$ its complement in X . Clearly $X_{>0} \hookrightarrow X$ is a proper constructible embedding, and hence by Corollary 2.10 we get a pushout

$$\begin{array}{ccc} \widetilde{\text{Exit}(\text{Link}_{>0}(X) \times \mathbb{R}_{>0})} & \longrightarrow & \text{Exit}(X_0) \\ \downarrow & & \downarrow \\ \widetilde{\text{Exit}(C(\pi_{>0}))} & \longrightarrow & \text{Exit}(X). \end{array}$$

Thus, by Lemma 2.7, to conclude the proof it suffices to show that $\widetilde{\text{Exit}(\text{Link}_{>0}(X) \times \mathbb{R}_{>0})}$, $\widetilde{\text{Exit}(C(\pi_{>0}))}$ and $\text{Exit}(X_0)$ are finite.

Being a closed subset of X , $X_{>0}$ must be compact, and by the pullback square (2.8) $\text{Link}_{>0}(X)$ is compact too. Since the depths of both are strictly less than $\text{depth}(X)$, by the inductive hypothesis we get that they both have finite exit path ∞ -categories. Notice that we have a stratified embedding $\widetilde{\text{Link}_{>0}(X) \times \mathbb{R}_{>0}} \hookrightarrow X_0$ and X_0 is a smooth manifold, so in particular we see that the stratification on $\widetilde{\text{Link}_{>0}(X)}$ is trivial. Thus $\widetilde{\text{Exit}(\text{Link}_{>0}(X))}$ is a localization of $\text{Exit}(\text{Link}_{>0}(X))$ at all maps, and by Lemma 2.3 we see that it has to be finite.

By Lemma 2.7 and the inductive hypothesis, we also know that $\text{Exit}(C(\pi_{>0}))$ is finite, and by [AFT17, Proposition 1.2.13] we know that the canonical functor

$$\phi : \text{Exit}(C(\pi_{>0})) \rightarrow \widetilde{\text{Exit}(C(\pi_{>0}))}$$

is a localization at the class of exit paths that are inverted by ϕ . Since the inclusion $C(\pi_{>0}) \hookrightarrow X$ lies under $X_{>0}$, the same argument as above shows that an non-invertible exit path is inverted ϕ is and only if it lies inside $\text{Link}_{>0}(X) \times \mathbb{R}_{>0} \hookrightarrow C(\pi_{>0})$. Furthermore, since $\text{Link}_{>0}(X) \times \mathbb{R}_{>0} \hookrightarrow C(\pi_{>0})$ is a consecutive inclusion of strata, by Lemma 2.5 the induced functor on exit paths is the inclusion of a full subcategory, which implies that one can identify

$\text{Exit}(\widetilde{C(\pi_{>0})})$ with the localization of $\text{Exit}(C(\pi_{>0}))$ at $\text{Link}_{>0}(X) \times \mathbb{R}_{>0}$, which means again by Lemma 2.3 that it must be finite.

We know that X_0 is the interior of the compact manifold with corners $\text{Unzip}_{>0}(X)$. The existence of collaring for corners ([AFT17, Lemma 8.2.1]) implies, with an argument completely analogous to [AMGR19, Lemma 2.1.3], that the inclusion $X_0 \hookrightarrow \text{Unzip}_{>0}(X)$ is an homotopy. Hence to conclude the proof it suffices to show that $\text{Sing}(\text{Unzip}_{>0}(X))$ is finite, but this follows by the existence of good covers for manifolds with corners. \square

Corollary 2.13. Let X be a finitary conically smooth stratified space. Then $\text{Exit}(X)$ is a finite ∞ -category. In particular, if X is the interior of a compact conically smooth manifold with corners, then $\text{Exit}(X)$ is a finite ∞ -category.

Proof. By Proposition 2.12 and Lemma 2.7 we know that the class of conically smooth spaces having finite exit path ∞ -category contains all basics, and thus it suffices to show that it closed under taking collar glueings. But this is clear, since for any collar glueing $f: Y \rightarrow \overline{\mathbb{R}}$ we get an open covering of Y given by $f^{-1}(\mathbb{R}_{\leq \infty})$, $f^{-1}(\mathbb{R}_{\leq \infty})$ and $f^{-1}(0)$ are finitary.

The last part of the statement follows by [AFT17, Theorem 8.3.10 (1)]. \square

Remark 2.14. We strongly believe that one should be able to prove similar finiteness properties for compact C^0 -stratified spaces, in the sense of [AFT17, Definition 2.1.15]. However, in the process of writing this paper, we have decided to ease ourselves (and the reader) from the expected technicalities needed to obtain such results at a higher level of generality, mainly because in future applications we will only be interested in stratified spaces admitting conically smooth structures. Nevertheless, let us indulge ourselves on a very brief and sketchy outline of how we expect one could try to tackle this problem.

Recall that, following [HPT20], for a fixed poset P one may define an ∞ -category of P -stratified homotopy types Str_P , that can be easily described as the full subcategory of the slice $\text{Cat}_{\infty/P}$ spanned by conservative functors. At least if P is finite, one can show that an object of Str_P is compact (or finite) if and only if it is so as an ∞ -category. This observation, as trivial as it may look, is in fact quite important if one notices that it's much more convenient to check compactness (or finiteness) of exit paths as an object of Str_P : namely, a sufficient condition for the compactness (finiteness) of an object of Str_P is the compactness (finiteness) of all its *homotopy links*³. Evidence of the compactness of homotopy links without assuming the presence of any smooth structure can be found in [Qui88, Theorem 4.7]. Let us note to conclude that the explicit models of homotopy links provided in [DW21] might also turn out to be relevant for this discussion.

3 Homotopy invariance and exodromy with general coefficients

In this section we will explain how to use [Vol21] to extend *homotopy invariance* and the *exodromy equivalence* to (formally) constructible sheaves (see Definition 3.2) valued in stable and bicomplete ∞ -categories. With these at hand, we will show in Corollary 3.16 that global sections of constructible sheaves on compact conically smooth stratified spaces are dualizable.

³It's not unreasonable to expect the latter condition to be necessary as well, but by now we are not aware of any proof of this.

3.1 Homotopy invariance of constructible sheaves

From now on, all ∞ -categories appearing as coefficients for sheaves will be assumed to be stable and bicomplete, all topological spaces locally compact Hausdorff and all posets Noetherian.

Remark 3.1. Recall that, by [AFT17, Lemma 2.2.2], any C^0 -stratified space admits a basis given by its open subsets homeomorphic to $\mathbb{R}^n \times C(Z)$, where Z is a compact C^0 -stratified space. A simple inductive argument shows that the stratifying poset of Z , and hence of $\mathbb{R}^n \times C(Z)$, must be finite, and therefore the stratifying poset of any C^0 -stratified space is locally finite, and therefore Noetherian.

Definition 3.2. Let $X \rightarrow P$ be any stratified space. We say that a sheaf $F \in \text{Shv}(X; \mathcal{C})$ is *formally constructible* if for any $\alpha \in P$ the restriction of F to the stratum X_α is locally constant.

Assume now that \mathcal{C} admits a closed symmetric monoidal structure, and denote by \mathcal{C}^c the full subcategory of \mathcal{C} spanned by dualizable objects. We say that F is *constructible* if F is formally constructible and each stalk of F belongs to \mathcal{C}^c .

We will denote by $\text{Shv}^{fc}(X; \mathcal{C})$ and $\text{Shv}^c(X; \mathcal{C})$ the full subcategories of $\text{Shv}(X; \mathcal{C})$ spanned respectively by formally constructible and constructible sheaves. Dually, we define formally constructible and constructible cosheaves on X just as $\text{CoShv}^{fc}(X; \mathcal{C}) := \text{Shv}^{fc}(X; \mathcal{C}^{op})^{op}$ and $\text{CoShv}^c(X; \mathcal{C}) := \text{Shv}^c(X; \mathcal{C}^{op})^{op}$.

In this paper we will only be dealing with constructible sheaves with respect to a specified stratification. Therefore, we will take the liberty of omitting to include the stratified poset into our notation for constructible sheaves, for the sake of making it cleaner.

We first recall the proof of the homotopy invariance of constructible sheaves. Our argument follows precisely the one in [HPT20], nevertheless we will try to quickly outline the main steps just to convince the reader that all the results in [HPT20], after [Vol21], generalize to the setting of non-presentable coefficients, at least if we restrict ourselves to locally compact Hausdorff spaces.

Theorem 3.3 (Homotopy invariance). Let $X \rightarrow P$ be a stratified space. Let $p : X \times I \rightarrow X$ be the canonical projection, where I is the closed interval $[0, 1] \subset \mathbb{R}$. Then $p^* : \text{Shv}(X; \mathcal{C}) \rightarrow \text{Shv}(X \times I; \mathcal{C})$ restricts to an equivalence

$$\text{Shv}^{fc}(X; \mathcal{C}) \simeq \text{Shv}^{fc}(X \times I; \mathcal{C}).$$

As a consequence, if $Y \rightarrow P$ is another stratified space and $f : X \rightarrow Y$ is a stratified homotopy equivalence, then the functor

$$f^* : \text{Shv}^{fc}(Y; \mathcal{C}) \rightarrow \text{Shv}^{fc}(X; \mathcal{C}).$$

is an equivalence.

Proof. We first treat the case of locally constant sheaves, i.e. when $P = *$. By [Lur17, Lemma A.2.9] and [Vol21, Corollary 5.2] we know that p^* is fully faithful. We first show that f_* preserves constant sheaves: if $a : X \rightarrow *$ and $b : X \times I \rightarrow *$ are the unique maps then, for any object $M \in \mathcal{C}$, the fully faithfulness of f^* implies

$$\begin{aligned} p_* b^* M &\simeq p_* p^* a^* M \\ &\simeq a^* M. \end{aligned}$$

Moreover, since I is compact, [Vol21, Proposition 6.1] provides a base change formula for p_* . Using basechange and [HPT20, Lemma 4.9], we then see that p_* preserves locally constant sheaves. Hence to conclude we only need to show that, for any locally constant sheaf F on $X \times I$, the counit map $p^*p_*F \rightarrow F$ is an equivalence. Again by basechange and [HPT20, Lemma 4.9], we may reduce to $F \simeq b^*M$ constant, in which case we have a commuting diagram

$$\begin{array}{ccc}
p^*p_*b^*M & \xrightarrow{\text{counit}_{b^*M}} & b^*M \\
\downarrow \simeq & & \downarrow \simeq \\
p^*p_*p^*a^*M & \xrightarrow{\text{counit}_{p^*a^*M}} & p^*a^*M \\
\uparrow p^*(\text{unit}_{a^*M}) \simeq & \nearrow \text{id} & \\
p^*a^*M & &
\end{array}$$

which implies what we want.

Now assume that P is any Noetherian poset. Using basechange, one sees that p^* preserves constructible sheaves, and thus we are left to show that $p^*p_*F \rightarrow F$ is an equivalence for any F constructible. By [Vol21, Corollary 4.2] any stable and bicomplete ∞ -category respects glueing in the sense of [HPT20, Definition 5.17], and hence [HPT20, Lemma 5.19] implies that the functor given by restricting to the strata of X are jointly surjective. By base change we may thus assume that F is locally constant, in which case the counit is known to be an equivalence by the previous step.

The last part of the statement then follows by a standard argument analogous to the proof of [Vol21, Corollary 3.1]. \square

We now present a couple of useful corollaries of homotopy invariance.

Corollary 3.4. Let $f : X \rightarrow Y$ be a stratified homotopy equivalence, $F \in \text{Shv}^{fc}(Y; \mathcal{C})$. The canonical map

$$\Gamma(Y; F) \rightarrow \Gamma(X; f^*F)$$

is an equivalence.

Proof. The commutative triangle

$$\begin{array}{ccc}
X & \xrightarrow{f} & Y \\
& \searrow a & \swarrow b \\
& & *
\end{array}$$

where a and b are the unique maps, induces an invertible natural transformation $f^*b^* \simeq a^*$. Since both a^* and b^* factor through formally constructible sheaves, we get $b^* \simeq \eta a^*$, where η is any adjoint inverse of the restriction of f^* to $\text{Shv}^{fc}(Y; \mathcal{C})$. Thus, by passing to right adjoints, we get the desired equivalence. \square

Corollary 3.5. Let X be a C^0 -stratified space and $F \in \text{Shv}^{fc}(X; \mathcal{C})$. For any point $x \in X$ and any conical chart $\mathbb{R}^n \times C(Z)$ centered in x , the canonical map

$$\Gamma(\mathbb{R}^n \times C(Z); F) \rightarrow F_x$$

is an equivalence.

Proof. Let $\mathbb{R}_{\geq 0} \times Z \rightarrow C(Z)$ be the quotient map, and denote by $C_\epsilon(Z)$ the image of the open subset $[0, \epsilon) \times Z$. We first show that the family of open subsets

$$\{C_\epsilon(Z) \mid \epsilon \in \mathbb{R}_{>0}\}$$

forms a basis at the cone point. To do this we will prove that for every open subset W of $\mathbb{R}_{\geq 0} \times Z$ containing $\{0\} \times Z$ there exists some $\epsilon > 0$ such that $[0, \epsilon) \times Z \subseteq W$. Since Z is compact, one can obtain a finite covering of $\{0\} \times Z$ with opens of the type $V_i \times [0, \epsilon_i) \subseteq W$, and thus by taking ϵ to be the minimum of the ϵ_i we get the claim.

Hence, by a cofinality argument, we see that

$$\varinjlim_{\epsilon > 0} \Gamma(\mathbb{R}^n \times C_\epsilon(Z); F) \simeq F_x$$

and by Corollary 3.4 we have

$$\Gamma(\mathbb{R}^n \times C(Z); F) \simeq \varinjlim_{\epsilon > 0} \Gamma(\mathbb{R}^n \times C_\epsilon(Z); F).$$

□

Let now $X \rightarrow P$ be a conically smooth stratified space, and let $\alpha \in P$. Then, by Remark 2.11, we get commutative triangle

$$\begin{array}{ccc} & X_\alpha & \\ i \swarrow & & \searrow i_\alpha \\ C(\pi_\alpha) & \xrightarrow{j} & X \end{array}$$

where π_α is the fiber bundle $\text{Link}_{X_\alpha}(X_{\geq \alpha}) \rightarrow X_\alpha$, i is the cone-point section of the fiber bundle $p : C(\pi_\alpha) \rightarrow X_\alpha$ and j is a conically smooth open immersion. For any $F \in \text{Shv}(X; \mathcal{C})$, the unit of the adjunction $i^* \dashv i_*$ gives a natural map

$$(3.6) \quad p_* j^* F \rightarrow p_* i_* i^* j^* F \simeq i_\alpha^* F.$$

Furthermore the map (3.6) can be obtained by applying the sheafification functor to

$$(3.7) \quad p_* j^* F \rightarrow p_* i_* (i^*)^{pre} j^* F \simeq (i_\alpha^*)^{pre} F$$

where $(i^*)^{pre}$ and $(i_\alpha^*)^{pre}$ denote the corresponding presheaf pullback functors.

Corollary 3.8. Let F be any formally constructible sheaf on X . Then the map (3.6) is an equivalence.

Proof. Since F is a sheaf, it suffices to show that (3.7) is an equivalence. As usual, it suffices to prove that it is an equivalence after taking sections on any euclidean chart U of X_α . For any such U , by [AFT17, Corollary 7.1.4] we have $\Gamma(U; p_* j^* F) = \Gamma(U \times C(Z); F)$ for some compact conically smooth stratified space Z . Thus we are left to show that the canonical map

$$(3.9) \quad \Gamma(U \times C(Z); F) \rightarrow \varinjlim_{U \subseteq V} \Gamma(V; F)$$

is an equivalence.

By a cofinality argument, the map (3.9) factors through an equivalence

$$\varinjlim_{\epsilon > 0} \Gamma(C_\epsilon(Z); F) \simeq \varinjlim_{U \subseteq V} \Gamma(V; F),$$

and thus we are left to show that

$$\Gamma(U \times C(Z); F) \rightarrow \varinjlim_{\epsilon > 0} \Gamma(C_\epsilon(Z); F)$$

is an equivalence. This last assertion then follows by Corollary 3.4. □

3.2 Exodromy

This subsection will be devoted to extending the proof of the exodromy equivalence on conically smooth stratified spaces to constructible sheaves valued in stable and bicomplete ∞ -categories. To do this we will use the model of the exit path ∞ -category of a conically smooth stratified space given in [AFT17, Definition 1.1.3], so we briefly recall how this is defined.

One has categories and functors

$$\mathcal{Bsc} \rightarrow \mathcal{Snglr} \rightarrow \mathcal{Strat}$$

where \mathcal{Strat} is the 1-category whose objects are conically smooth stratified spaces and morphisms are conically smooth maps, \mathcal{Snglr} is the subcategory of \mathcal{Strat} with the same objects and with morphisms given by open immersions, \mathcal{Bsc} is the full subcategory of \mathcal{Snglr} spanned by *basic* conically smooth stratified spaces, i.e. those which are isomorphic in \mathcal{Strat} to one of the type $\mathbb{R}^n \times C(Z)$, where Z is compact, and the functors are the respective inclusions.

In [AFT17, Lemma 4.1.4], the authors show that all the categories appearing in (3.2) admit enrichments in Kan-complexes and thus, by passing to homotopy coherent nerves, one gets ∞ -categories that we denote by

$$(3.10) \quad \mathcal{Bsc} \rightarrow \mathcal{Snglr} \rightarrow \mathcal{Strat}.$$

For any conically smooth stratified space, the authors of [AFT17] then define $\text{Exit}(X)$ as the opposite of the slice $\mathcal{Bsc}/_X$. Their proof of the exodromy equivalence for constructible sheaves of spaces, combined with the one in [Lur17], implies that Lurie's version of the exit path ∞ -category has to be equivalent to it (see [AFT17, Corollary 1.2.10]).

One has obvious functors

$$\begin{array}{ccc} \mathcal{Bsc}/_X & \xrightarrow{\gamma} & \mathcal{Bsc}/_X \\ & & \downarrow \iota \\ & & \mathcal{U}(X) \end{array}$$

where ι sends an open immersion to its image in X . [AFT17, Lemma 4.5.1] shows γ is a localization at the class W of open immersions of basics $U \hookrightarrow V$ such that U and V are abstractly isomorphic in \mathcal{Strat} , i.e. precomposing with γ gives an equivalence

$$(3.11) \quad \gamma^* : \text{Fun}(\text{Exit}(X), \mathcal{C}) \rightarrow \text{Fun}_W(\mathcal{Bsc}/_X^{op}, \mathcal{C})$$

where the left-hand side denotes the full subcategory of $\text{Fun}(\mathcal{Bsc}/_X^{op}, \mathcal{C})$ spanned by functors which send all morphisms in W to equivalences. In the next proposition we show that W

coincides with the class of open immersions which are stratified homotopy equivalences, and then characterize the property of being formally constructible through these maps.

Proposition 3.12. Let $X \rightarrow P$ be a conically smooth stratified space, and let $F \in \text{Shv}(X; \mathcal{C})$. Then the following assertions are equivalent:

- (i) F is formally constructible;
- (ii) for any inclusion $V \hookrightarrow U$ of basic open subsets of X which is a stratified homotopy equivalence, then

$$\Gamma(U; F) \rightarrow \Gamma(V; F)$$

is an equivalence;

- (iii) for any inclusion $V \hookrightarrow U$ of basic open subsets of X which are abstractly isomorphic, then

$$\Gamma(U; F) \rightarrow \Gamma(V; F)$$

is an equivalence.

Proof. We first prove that (ii) is equivalent to (iii) by showing that an open immersion $j : V \hookrightarrow U$ of basic open subsets of X is a stratified homotopy equivalence if and only if U and V are abstractly isomorphic.

If j is a stratified homotopy equivalence, it follows that U and V must be stratified over the same subposet of P , and so by [AFT17, Lemma 4.3.7] U and V must be isomorphic. Viceversa, if $j : \mathbb{R}^n \times C(Z) \hookrightarrow \mathbb{R}^n \times C(Z)$ is a conically smooth open embedding, then by [AFT17, Lemma 4.3.6] j is homotopy equivalent to D_0j , and the latter is a stratified homotopy equivalence.

By Theorem 3.3 we have that (i) implies (ii), so we are left to show that (iii) implies (i). Let $i^* : Y \rightarrow X$ be the inclusion of a stratum, and let $V \hookrightarrow U$ be an inclusion of euclidean charts of U . By Corollary 3.8, the horizontal arrows in the commutative square

$$\begin{array}{ccc} \Gamma(U \times C(Z); F) & \longrightarrow & \Gamma(U; i^*F) \\ \downarrow & & \downarrow \\ \Gamma(V \times C(Z); F) & \longrightarrow & \Gamma(V; i^*F) \end{array}$$

are invertible, and thus $\Gamma(U; i^*F) \rightarrow \Gamma(V; i^*F)$ is invertible too. Hence, we may assume that X is a smooth manifold. The result could then be seen as a very special case of [HPT20, Proposition 3.1], but we will now review and adapt the proof to our setting for the reader's convenience.

The question being local, we may assume that $X = \mathbb{R}^n$, in which case we will prove that condition (iii) implies that F is constant. More precisely, we will show that the counit map

$$(3.13) \quad a^* a_* F \rightarrow F$$

is an equivalence, where $a : X \rightarrow *$ is the unique map. Since X is hypercomplete and admits a basis given by those open subsets diffeomorphic to itself, it then suffices to check that for any such open $j : U \hookrightarrow X$, the map $a_* j_* j^* a^* a_* F \rightarrow a_* j_* j^* F$ obtained by applying to (3.13) the functor of sections at U is invertible. Notice that we have a commutative triangle

$$\begin{array}{ccc} a_* j_* j^* a^* a_* F & \xrightarrow{a_* j_* j^* \text{counit}_F} & a_* j_* j^* F \\ \text{counit}_{a_* F} \downarrow & \nearrow \simeq & \\ a_* F & \xrightarrow{a_*(\text{unit}_F)} & \end{array}$$

where the diagonal map is invertible by the assumption in (iii), and thus to conclude the proof it suffices to show that a^* is fully faithful, which follows by the homotopy invariance of the shape (see [Vol21, Corollary 3.1]). \square

Theorem 3.14 (Exodromy). The composition

$$\mathrm{Fun}(\mathrm{Exit}(X), \mathcal{C}) \xrightarrow{\gamma^*} \mathrm{Fun}_W(\mathrm{Bsc}/_X, \mathcal{C}) \xrightarrow{\iota_*} \mathrm{Fun}(\mathcal{U}(X)^{op}, \mathcal{C})$$

is fully faithful with essential image given by $\mathrm{Shv}^{fc}(X; \mathcal{C})$. Moreover, if we assume \mathcal{C} to have a closed symmetric monoidal structure, the statement remains true if we replace \mathcal{C} by \mathcal{C}^c and $\mathrm{Shv}^{fc}(X; \mathcal{C})$ by $\mathrm{Shv}^c(X; \mathcal{C})$.

Proof. By Proposition 3.12, it suffices to show that the restriction of ι_* to $\mathrm{Fun}_W(\mathrm{Bsc}/_X, \mathcal{C})$ factors through $\mathrm{Shv}^{fc}(X; \mathcal{C})$.

Let U be basic open subset of X and let $\kappa : T \hookrightarrow \mathrm{Bsc}/_U$ be a covering sieve. Clearly there must be at least one $V \in T$ whose image in U intersects the locus of maximal depth, and thus by [AFT17, Lemma 4.3.7] U and V must be abstractly isomorphic. Then for any $F \in \mathrm{Fun}_W(\mathrm{Bsc}/_X, \mathcal{C})$ we have a commutative triangle

$$\begin{array}{ccc} \Gamma(U; F) & \longrightarrow & \varinjlim_{O \in T} \Gamma(O; F) \\ & \searrow \simeq & \downarrow \\ & & \Gamma(V; F) \end{array}$$

where the diagonal map is invertible by assumption. By [AFT17, Proposition 3.2.23] and [Aok20, Theorem A.6], to conclude the proof it suffices to show that the vertical map is invertible. Let $\delta : T \rightarrow \mathcal{T}$ be the localization of T at W . Since δ is final and κ^* sends maps in X to equivalences, the result then follows by observing that V is a terminal object in \mathcal{T} .

The second part of the statement is obvious after Corollary 3.5. \square

Remark 3.15. Notice that even though we assumed from the beginning the coefficients to be stable and bicomplete, all the arguments we have discussed work whenever $\mathrm{Shv}(X; \mathcal{C}) \hookrightarrow \mathrm{Fun}(\mathcal{U}(X)^{op}, \mathcal{C})$ admits a left adjoint and \mathcal{C} respects glueings in the sense of [HPT20, Definition 5.17]. In particular, our proof recovers also the case $\mathcal{C} = \mathcal{S}$. A proof of the exodromy equivalence with presentable coefficients but on a much bigger class of stratified spaces will appear soon in [PT22].

Corollary 3.16. Let Z be any compact conically smooth stratified space, and let $F \in \mathrm{Shv}^c(X; \mathcal{C})$. Then $\Gamma(Z; F)$ is dualizable.

Proof. Since γ is final, by Theorem 3.14 we know that there exists a functor $G : \mathrm{Exit}(Z) \rightarrow \mathcal{C}^c$ such that global sections of F are equivalent to the limit of G . Therefore the proof is concluded by applying Proposition 2.12 and observing that, since \mathcal{C} is stable and its monoidal structure is closed, \mathcal{C}^c is itself stable. \square

Remark 3.17. Let Z be a stratified space such that $\mathrm{Exit}(Z)$ is a retract in Cat_∞ of a finite ∞ -category, and assume that the exodromy equivalence holds for constructible sheaves on Z . Then, since \mathcal{C}^c is idempotent complete, the same argument of Corollary 3.16 shows that for any $F \in \mathrm{Shv}^c(X; \mathcal{C})$, $\Gamma(Z; F)$ is dualizable.

4 The duality

This final section is devoted to proving Verdier duality on conically smooth spaces. For this reason, from now on our ∞ -categories of coefficients will be assumed to be equipped with a closed symmetric monoidal structure.

We first introduce the usual duality functor. For any stratified space X , we will denote by $\omega_X^{\mathcal{C}}$ the sheaf of $a^!(\mathbb{1}_{\mathcal{C}})$, where $a : X \rightarrow *$ is the unique map and $\mathbb{1}_{\mathcal{C}}$ is the monoidal unit in \mathcal{C} . $\omega_X^{\mathcal{C}}$ will be called the \mathcal{C} -valued *dualizing sheaf* of X . We will denote the functor

$$\underline{\mathrm{Hom}}_X(-, \omega_X^{\mathcal{C}}) : \mathrm{Shv}(X; \mathcal{C})^{op} \rightarrow \mathrm{Shv}(X; \mathcal{C})$$

simply by $D_X^{\mathcal{C}}$ and, when $X = *$, we will only write $D^{\mathcal{C}} : \mathcal{C}^{op} \rightarrow \mathcal{C}$.

Proposition 4.1. For any C^0 -stratified space X , the dualizing sheaf $\omega_X^{\mathcal{C}}$ is constructible.

Proof. First of all, notice that, by [Vol21, Proposition 6.4], we have an equivalence $\omega_X^{\mathcal{C}} \simeq \omega_X^{\mathrm{Sp}} \otimes \mathbb{1}_{\mathcal{C}}$, where the right-hand side denotes the image of the couple $(\omega_X^{\mathrm{Sp}}, \mathbb{1}_{\mathcal{C}})$ through the canonical variablewise cocontinuous functor

$$\mathrm{Shv}(X; \mathrm{Sp}) \times \mathcal{C} \rightarrow \mathrm{Shv}(X; \mathrm{Sp}) \otimes \mathcal{C} \simeq \mathrm{Shv}(X; \mathcal{C})$$

which corresponds with the usual tensoring over spectra given by cocompleteness and stability of \mathcal{C} when $X = *$. Moreover, by [Vol21, Corollary 5.3], for each point $x : * \rightarrow X$ we have commutative square

$$\begin{array}{ccc} \mathrm{Shv}(X; \mathrm{Sp}) \times \mathcal{C} & \longrightarrow & \mathrm{Shv}(X; \mathcal{C}) \\ \downarrow x_{\mathrm{Sp} \times \mathcal{C}}^* & & \downarrow x_{\mathcal{C}}^* \\ \mathrm{Sp} \times \mathcal{C} & \longrightarrow & \mathcal{C}. \end{array}$$

Since a spectrum is dualizable if and only if it is finite, and since dualizable objects in \mathcal{C} are closed under finite colimits, we see that that it suffices to prove the proposition in the case in which $\mathcal{C} = \mathrm{Sp}$.

We will proceed by induction on the depth of X . If X has depth 0, then it is a topological manifold and hence ω_X is locally constant of rank the dimension of X . Now assume that X has finite non zero depth. Since the question is local on X , by [AFT17, Lemma 2.2.2] we may assume that $X = \mathbb{R}^n \times C(Z)$, where Z is a compact C^0 -stratified space with $\mathrm{depth}(Z) < \mathrm{depth}(X)$.

Let $p : \mathbb{R}^n \times C(Z) \rightarrow C(Z)$ be the canonical projection, $b : C(Z) \rightarrow *$ the unique map. Since for any sheaf F on $C(Z)$ we have a functorial equivalence $p^!F \simeq \Sigma^n p^*F$, it suffices to show that $b^!\mathbb{S} = \omega_{C(Z)}$ is constructible, and hence we may assume that $X = C(Z)$.

Let $i : \{x\} \hookrightarrow X$ be the inclusion of the cone point and $j : U \hookrightarrow X$ its open complement, which is evidently homeomorphic to $\mathbb{R} \times Z$. We have a localization sequence

$$j_{\#}j^*\omega_X \rightarrow \omega_X \rightarrow i_*i^*\omega_X.$$

By the inductive hypothesis $j^*\omega_X$ is constructible, and thus for every stratum $T \subseteq X$ which does not contain the cone point, the restriction of ω_X is locally constant. Hence it remains to prove that the stalk of ω_X at the cone point is a finite spectrum. But by a routine cofinality argument we get

$$\begin{aligned} (\omega_X)_x &= \varinjlim_{x \in U} \Gamma(U; \omega_X) \\ &\simeq \varinjlim_{\epsilon > 0} \Gamma(C_{\epsilon}(Z); \omega_X) \\ &\simeq \varinjlim_{\epsilon > 0} \underline{\mathrm{Hom}}(\Gamma_c(C_{\epsilon}(Z); \mathbb{S}_X), \mathbb{S}). \end{aligned}$$

To conclude the proof, we will now show that the canonical extension map $\Gamma_c(C_\epsilon(Z); \mathbb{S}_X) \rightarrow \Gamma_c(X; \mathbb{S}_X)$ is an equivalence for each ϵ and that $\Gamma_c(X; \mathbb{S}_X)$ is a finite spectrum.

First of all, notice that for any $K \subseteq C_\epsilon(Z)$ compact containing the cone point, there exists a $T \geq 0$ such that $K \subseteq \overline{C_T(Z)}$ (namely, take T to be the maximum in the image of K through the projection $C_\epsilon(Z) \rightarrow \mathbb{R}_{\geq 0}$). Hence we have a commutative square

$$\begin{array}{ccc} \Gamma_c(C_\epsilon(Z); \mathbb{S}_X) & \xrightarrow{\simeq} & \varinjlim_{0 \leq T < \epsilon} \Gamma_{\overline{C_T(Z)}}(X; \mathbb{S}_X) \\ \downarrow & & \downarrow \\ \Gamma_c(X; \mathbb{S}_X) & \xrightarrow{\simeq} & \varinjlim_{T \geq 0} \Gamma_{\overline{C_T(Z)}}(X; \mathbb{S}_X). \end{array}$$

Since both colimits on the right-hand side of the square are indexed by filtered posets, to prove that the horizontal maps are invertible it will suffice to show that the functors of which we compute both colimits are constant. By definition, there is a fiber sequence

$$\Gamma_{\overline{C_T(Z)}}(X; \mathbb{S}_X) \rightarrow \Gamma(X; \mathbb{S}_X) \rightarrow \Gamma(X \setminus \overline{C_T(Z)}; \mathbb{S}_X).$$

By homotopy invariance of the shape, since X is contractible and $X \setminus \overline{C_T(Z)} \simeq Z$ there is a commutative diagram

$$\begin{array}{ccc} \Gamma(X; \mathbb{S}_X) & \longrightarrow & \Gamma(X \setminus \overline{C_T(Z)}; \mathbb{S}_X) \\ \downarrow \simeq & & \downarrow \simeq \\ \underline{\mathrm{Hom}}_{\mathrm{Sp}}(\Sigma_+^\infty X, \mathbb{S}) & \longrightarrow & \underline{\mathrm{Hom}}_{\mathrm{Sp}}(\Sigma_+^\infty (X \setminus \overline{C_T(Z)}), \mathbb{S}) \\ \downarrow \simeq & & \downarrow \simeq \\ \mathbb{S} & \longrightarrow & \underline{\mathrm{Hom}}_{\mathrm{Sp}}(\Sigma_+^\infty Z, \mathbb{S}) \end{array}$$

where the lower horizontal arrow is induced by applying the functor $\underline{\mathrm{Hom}}_{\mathrm{Sp}}(\Sigma_+^\infty(-), \mathbb{S})$ to the unique arrow $Z \rightarrow *$. Thus we get

$$\Gamma_{\overline{C_T(Z)}}(X; \mathbb{S}_X) \simeq \mathrm{fib}(\mathbb{S} \rightarrow \underline{\mathrm{Hom}}_{\mathrm{Sp}}(\Sigma_+^\infty Z, \mathbb{S}))$$

for any $T \geq 0$. Hence to conclude the proof it suffices to show that $\Sigma_+^\infty Z$ is a finite spectrum. This is clear since, for any spectrum M , we have a functorial equivalence

$$\underline{\mathrm{Hom}}(\Sigma_+^\infty Z, M) \simeq a_1 a^* M$$

and $a_1 a^*$ preserves (filtered) colimits: since the class of compact objects in Sp coincides with the one of finite spectra, we may conclude. Otherwise, one may see this as a consequence of Lemma 2.3, Proposition 2.12 and the fact that Σ_+^∞ preserves finite objects. \square

Theorem 4.2. Let X be a conically smooth stratified space. Then the restriction to $\mathrm{Shv}^c(X; \mathcal{C})^{op}$ of the functor D_X^c factors through an equivalence

$$D_X^c : \mathrm{Shv}^c(X; \mathcal{C})^{op} \xrightarrow{\simeq} \mathrm{Shv}^c(X; \mathcal{C}).$$

Proof. Let $j : U \hookrightarrow X$ be any open subset of X . Then, for any $F \in \mathrm{Shv}(X; \mathcal{C})$, by applying [Vol21, Corollary 3.4], [Vol21, Lemma 6.2] and [Vol21, Proposition 6.2] we get functorial

equivalences

$$\begin{aligned}\Gamma(U; \underline{\mathbf{Hom}}_X(F, \omega_X)) &\simeq \Gamma(U; \underline{\mathbf{Hom}}_U(j^*F, j^*\omega_X^{\mathcal{C}})) \\ &\simeq \Gamma(U; \underline{\mathbf{Hom}}_U(j^*F, \omega_U^{\mathcal{C}})) \\ &\simeq \underline{\mathbf{Hom}}_{\mathcal{C}}(\Gamma_c(U; F), \mathbb{1}_{\mathcal{C}})\end{aligned}$$

and thus there is a factorization

$$\begin{array}{ccc}\mathrm{Shv}(X; \mathcal{C})^{op} & \xrightarrow{\underline{\mathbf{Hom}}_X(-, \omega_X)} & \mathrm{Shv}(X; \mathcal{C}). \\ & \searrow \cong \mathbb{D} & \nearrow D_{\bullet}^{\mathcal{C}} \\ & \mathrm{CoShv}(X; \mathcal{C})^{op} & \end{array}$$

where $D_{\bullet}^{\mathcal{C}}$ denotes the functor obtained by postcomposing with $D^{\mathcal{C}} : \mathcal{C}^{op} \rightarrow \mathcal{C}$.

We start by showing that \mathbb{D} restricts to an equivalence

$$(4.3) \quad \mathbb{D} : \mathrm{Shv}^{fc}(X; \mathcal{C}) \simeq \mathrm{CoShv}^{fc}(X; \mathcal{C}).$$

First of all, we prove that if $F \in \mathrm{Shv}(X; \mathcal{C})$ is locally constant, then $\mathbb{D}F$ is a formally constructible cosheaf. Since both properties are local, it suffices to show that \mathbb{D} sends constant sheaves to formally constructible sheaves, so assume $F \simeq a^*M$ where $a : X \rightarrow *$ is the unique map. In this case we have that $\mathbb{D}F \simeq a_{\mathcal{C}^{op}}^!M$, and so by Proposition 4.1 we get what we want.

Assume now that F is any formally constructible sheaf, and let $i : X_{\alpha} \hookrightarrow X$ by the inclusion of a stratum of X , with complement $j : U \hookrightarrow X$. Thus, we need to show that $i_{\mathcal{C}^{op}}^*\mathbb{D}F \simeq \mathbb{D}i_{\mathcal{C}}^!F$ is locally constant. Since the question is local, by restricting to $X_{\geq \alpha}$ one may assume that X_{α} is a closed stratum, so that we have a fiber sequence

$$(4.4) \quad i_{\mathcal{C}}^!F \rightarrow i_{\mathcal{C}}^*F \rightarrow i_{\mathcal{C}}^*j_*^{\mathcal{C}}j_{\mathcal{C}}^*F.$$

Hence to conclude, since we have already shown that \mathbb{D} preserves locally constant sheaves on smooth manifolds, it suffices to show that $j_*^{\mathcal{C}}j_{\mathcal{C}}^*F$ is formally constructible. For any inclusion $V \hookrightarrow W$ of basic open subsets of X , we have a commutative square

$$\begin{array}{ccc}\Gamma(V; j_*^{\mathcal{C}}j_{\mathcal{C}}^*F) & \xrightarrow{\cong} & \Gamma(V \cap U; F) \\ \downarrow & & \downarrow \\ \Gamma(W; j_*^{\mathcal{C}}j_{\mathcal{C}}^*F) & \xrightarrow{\cong} & \Gamma(W \cap U; F),\end{array}$$

where the right vertical map is an equivalence by Corollary 3.4, since $V \cap U \hookrightarrow W \cap U$ is still a stratified homotopy equivalence. Hence, by Proposition 3.12 we conclude that $j_*^{\mathcal{C}}j_{\mathcal{C}}^*F$ is formally constructible.

We now prove that \mathbb{D} restricts to an equivalence

$$(4.5) \quad \mathbb{D} : \mathrm{Shv}^c(X; \mathcal{C}) \simeq \mathrm{CoShv}^c(X; \mathcal{C}).$$

For this it suffices to show that, for any $x \in X$ and $F \in \mathrm{Shv}^{fc}(X; \mathcal{C})$, $F_x \in \mathcal{C}$ is dualizable if and only if $(\mathbb{D}F)_x$ is dualizable, where the latter denotes the costalk of $\mathbb{D}F$ at x . Let $x : * \hookrightarrow X$ be the inclusion of a point $x \in X$. By definition, there are equivalences $(\mathbb{D}F)_x \simeq x_{\mathcal{C}^{op}}^*(\mathbb{D}F) \simeq x_{\mathcal{C}}^!F$. Thus, by applying global sections to the localization sequence associated to i and j , we obtain a fiber sequence

$$x^!F \simeq \Gamma(X; i_*i^!F) \rightarrow \Gamma(X; F) \rightarrow \Gamma(U; F)$$

and hence an equivalence

$$\Gamma_{\{x\}}(X; F) \simeq i^!F.$$

Thus, by choosing a conical chart $\mathbb{R}^n \times C(Z)$ around x and by applying Corollary 3.5, we get a fiber sequence

$$(\mathbb{D}F)_x \rightarrow F_x \rightarrow \Gamma((\mathbb{R}^n \times C(Z)) \setminus (0, *); F)$$

where $*$ $\in C(Z)$ denotes the cone point. Therefore by Corollary 3.16, to conclude it will suffice to show that $\text{Exit}((\mathbb{R}^n \times C(Z)) \setminus (0, *))$ is finite. But by Van Kampen for exit paths, one has a pushout

$$\begin{array}{ccc} \text{Exit}(\mathbb{R}^n \setminus \{0\} \times \mathbb{R}_{>0} \times Z) & \longrightarrow & \text{Exit}(\mathbb{R}^n \times \mathbb{R}_{>0} \times Z) \\ \downarrow & & \downarrow \\ \text{Exit}(\mathbb{R}^n \setminus \{0\} \times C(Z)) & \longrightarrow & \text{Exit}(\mathbb{R}^n \times C(Z)) \setminus (0, *) \end{array}$$

and so the result follows from Proposition 2.12 and Lemma 2.7.

We finally show that

$$D_{\bullet}^{\mathcal{C}} : \text{CoShv}^c(X; \mathcal{C})^{op} \rightarrow \text{Shv}^c(X; \mathcal{C})$$

is an equivalence. The diagram

$$\begin{array}{ccc} \text{Fun}(\text{Exit}(X), (\mathcal{C}^c)^{op}) & \xrightarrow{D_{\bullet}^{\mathcal{C}}} & \text{Fun}(\text{Exit}(X), \mathcal{C}^c) \\ \simeq \downarrow \gamma^* & & \simeq \downarrow \gamma^* \\ \text{Fun}_W(\text{Bsc}/X, (\mathcal{C}^c)^{op}) & \xrightarrow{D_{\bullet}^{\mathcal{C}}} & \text{Fun}_W(\text{Bsc}/X, \mathcal{C}^c) \\ \simeq \uparrow \iota^* & & \simeq \uparrow \iota^* \\ \text{CoShv}^c(X; \mathcal{C}) & \xrightarrow{D_{\bullet}^{\mathcal{C}}} & \text{Shv}^c(X; \mathcal{C}) \end{array}$$

commutes since the horizontal arrows are given by postcompositions and the vertical arrows by precompositions. Moreover, since the restriction of $D^{\mathcal{C}}$ induces a duality on \mathcal{C}^c , the upper horizontal arrows are equivalences, and thus we get the desired conclusion. \square

Remark 4.6. Since the functor (4.3) can be promoted to a natural transformation of sheaves of ∞ -categories on X , to prove that it is actually an equivalence one could have assumed from the beginning that X is a basic. In this case the stratifying poset of X is finite, and hence one could have deduced that (4.3) is an equivalence by [AMGR19, Corollary 7.4.25] (see [AMGR19, Example 1.10.8]). However, we preferred to give a more concrete and independent proof to make our discussion as self contained as possible.

Remark 4.7. Notice that the equivalences (4.3) and (4.5) are already interesting on their own, because they imply that for any stratified map $f : X \rightarrow Y$, $f_*^{\mathcal{C}}$ or $f_{\mathcal{C}}^*$ preserves (formal) constructibility if and only if $f_!^{\mathcal{C}}$ or $f_{\mathcal{C}}^!$ does. In particular, we see that $f_{\mathcal{C}}^!$ always preserves (formally) constructible sheaves.

Remark 4.8. Any μ -stratification of an analytic manifold in the sense of [KS90] satisfies the Whitney conditions, and hence by [NV21] defines a conically smooth structure. Thus, Theorem 4.2 recovers and generalizes the duality on constructible sheaves on analytic manifolds as defined in [KS90] (i.e. sheaves which are constructible in our sense with respect to *some* μ -stratification).

Remark 4.9. Theorem 4.2 implies that one may equip $\text{Shv}^c(X; \mathcal{C})$ with (at least) two structures of a Poincaré ∞ -category in the sense of [CDH⁺20]. This could be relevant for example to generalize the results in [SW20].

References

- [AFR18] David Ayala, John Francis, and Nick Rozenblyum. A stratified homotopy hypothesis. *Journal of the European Mathematical Society*, 21(4):1071–1178, 2018.
- [AFT17] David Ayala, John Francis, and Hiro Lee Tanaka. Local structures on stratified spaces. *Advances in Mathematics*, 307:903–1028, 2017.
- [AK10] Gregory Arone and Marja Kankaanrinta. On the functoriality of the blow-up construction. *Bulletin of the Belgian Mathematical Society-Simon Stevin*, 17(5):821–832, 2010.
- [AMGR19] David Ayala, Aaron Mazel-Gee, and Nick Rozenblyum. Stratified noncommutative geometry. *arXiv preprint arXiv:1910.14602*, 2019.
- [Aok20] Ko Aoki. Tensor triangular geometry of filtered objects and sheaves. *arXiv preprint arXiv:2001.00319*, 2020.
- [BBDG18] Alexander Beilinson, Joseph Bernstein, Pierre Deligne, and Ofer Gabber. *Faisceaux pervers*. Société mathématique de France, 2018.
- [BGH18] Clark Barwick, Saul Glasman, and Peter Haine. Exodromy. *arXiv preprint arXiv:1807.03281*, 2018.
- [CDH⁺20] Baptiste Calmés, Emanuele Dotto, Yonatan Harpaz, Fabian Hebestreit, Markus Land, Kristian Moi, Denis Nardin, Thomas Nikolaus, and Wolfgang Steimle. Hermitian K-theory for stable ∞ -categories I: Foundations. *preprint*, 2020.
- [Cis19] Denis-Charles Cisinski. *Higher categories and homotopical algebra*, volume 180. Cambridge University Press, 2019.
- [DW21] Sylvain Douteau and Lukas Waas. From homotopy links to stratified homotopy theories. *arXiv preprint arXiv:2112.02394*, 2021.
- [Gor21] Mark Goresky. Lecture notes on sheaves and perverse sheaves. *arXiv preprint arXiv:2105.12045*, 2021.
- [Hai18] Peter J Haine. On the homotopy theory of stratified spaces. *arXiv preprint arXiv:1811.01119*, 2018.
- [Hai21] Peter J. Haine. From nonabelian basechange to basechange with coefficients. *arXiv preprint arXiv:2108.03545*, 2021.
- [Hen] Simon Henry. Why are finite cell complexes also finite as infinity-categories? MathOverflow answer. <https://mathoverflow.net/q/314697>.
- [HPT20] Peter J. Haine, Mauro Porta, and Jean-Baptiste Teyssier. The homotopy-invariance of constructible sheaves of spaces. *arXiv: Algebraic Topology*, 2020.
- [Kas84] Masaki Kashiwara. The riemann-hilbert problem for holonomic systems. *Publications of the Research Institute for Mathematical Sciences*, 20(2):319–365, 1984.
- [KS90] Masaki Kashiwara and Pierre Schapira. *Sheaves on Manifolds: With a Short History. Les débuts de la théorie des faisceaux. By Christian Houzel*, volume 292. Springer Science & Business Media, 1990.

- [Lur09] Jacob Lurie. *Higher topos theory*. Princeton University Press, 2009. <https://www.math.ias.edu/~lurie/papers/HTT.pdf>.
- [Lur17] Jacob Lurie. *Higher algebra*. 2017. <https://www.math.ias.edu/~lurie/papers/HA.pdf>.
- [NV21] Guglielmo Nocera and Marco Volpe. Whitney stratifications are conically smooth. *arXiv preprint arXiv:2105.09243*, 2021.
- [PT22] Mauro Porta and Jean-Baptiste Teyssier. Topological exodromy with coefficients. *To appear*, 2022.
- [Qui88] Frank Quinn. Homotopically stratified sets. *Journal of the American Mathematical Society*, pages 441–499, 1988.
- [SW20] Jörg Schürmann and Jonathan Woolf. Witt groups of abelian categories and perverse sheaves. *Annals of K-Theory*, 4(4):621–670, 2020.
- [Vol21] Marco Volpe. Six functor formalism for sheaves with non-presentable coefficients. *arXiv preprint arXiv:2110.10212*, 2021.
- [Wal65] C. T. C. Wall. Finiteness conditions for CW-complexes. *Annals of Mathematics*, 81(1):56–69, 1965.