ON AN ANALOGUE OF THE HUREWICZ THEOREM FOR MEAN DIMENSION

MASAKI TSUKAMOTO

ABSTRACT. The Hurewicz theorem is a fundamental result in classical dimension theory concerning continuous maps which lower topological dimension. We study whether or not its analogue holds for mean dimension of dynamical systems. Our first main result shows that an analogue of the Hurewicz theorem does not hold for mean dimension in general. Our second main result shows that it holds true if a base system has zero mean dimension.

1. INTRODUCTION

1.1. Background and basic definitions. The Hurewicz theorem is one of the fundamental results in the classical topological dimension theory. For a compact metrizable space X we denote its topological dimension (Lebesgue covering dimension) by dim X. Let $f : X \to Y$ be a continuous map between compact metrizable spaces. Then the Hurewicz theorem [HW41, p. 91, Theorem VI 7] states that

$$\dim X \le \dim Y + \sup_{y \in Y} \dim f^{-1}(y).$$

The purpose of this paper is to study an analogue of this theorem for mean dimension of dynamical systems.

Mean dimension is a dynamical version of topological dimension introduced by Gromov [Gro99]. It quantifies how many parameters per iterate we need for describing orbits of a dynamical system. It has applications to several problems in topological dynamics [LW00, Lin99, MT19, GT20].

We need to prepare some definitions before rigorously stating our problem. Throughout of this paper we assume that "simplicial complex" means a *finite* simplicial complex (namely, the number of its simplices is finite). Let (X, d) be a metric space and Y a topological space. A continuous map $f : X \to Y$ is called an ε -embedding if we have $\operatorname{Diam} f^{-1}(y) < \varepsilon$ for all $y \in Y$. We define the ε -width dimension $\operatorname{Widim}_{\varepsilon}(X, d)$ as the minimum integer $n \geq 0$ such that there exist an n-dimensional simplicial complex P and

²⁰²⁰ Mathematics Subject Classification. 37B99, 54F45.

Key words and phrases. Dynamical system, factor map, mean dimension, Hurewicz theorem.

M.T. was supported by JSPS KAKENHI JP21K03227.

an ε -embedding $f: X \to P$. If X is compact then its topological dimension is defined by

$$\dim X = \lim_{\varepsilon \to 0} \operatorname{Widim}_{\varepsilon}(X, d).$$

A pair (X, T) is called a **dynamical system** if X is a compact metrizable space and $T: X \to X$ is a homeomorphism. Let (X, T) be a dynamical system with a metric d on X. For each natural number N we define a metric d_N on X by

$$d_N(x,y) = \max_{0 \le n < N} d\left(T^n x, T^n y\right).$$

We define the **mean dimension** of (X, T) by

$$\operatorname{mdim}(X,T) = \lim_{\varepsilon \to 0} \left(\lim_{N \to \infty} \frac{\operatorname{Widim}_{\varepsilon}(X,d_N)}{N} \right)$$

The value of $\operatorname{mdim}(X,T)$ is independent of the choice of a metric d, and it provides a topological invariant of (X,T).

Let A be a (not necessarily invariant) closed subset of X. We define the **upper and** lower mean dimensions of A by

$$\overline{\mathrm{mdim}}(A,T) = \lim_{\varepsilon \to 0} \left(\limsup_{N \to \infty} \frac{\mathrm{Widim}_{\varepsilon}(A,d_N)}{N} \right),$$
$$\underline{\mathrm{mdim}}(A,T) = \lim_{\varepsilon \to 0} \left(\liminf_{N \to \infty} \frac{\mathrm{Widim}_{\varepsilon}(A,d_N)}{N} \right).$$

These are also independent of the choice of d.

Let (X, T) and (Y, S) be dynamical systems. A map $\pi : X \to Y$ is called a **factor map** between dynamical systems if π is a continuous surjection satisfying $\pi \circ T = S \circ \pi$. We often denote it by $\pi : (X, T) \to (Y, S)$ for clarifying the underlying dynamics.

Let $\pi : (X, T) \to (Y, S)$ be a factor map between dynamical systems. We would like to study the mean dimension of fibers of π . For this purpose we define the **relative mean dimension** of $\pi : (X, T) \to (Y, S)$ by

$$\operatorname{mdim}(\pi, T) = \lim_{\varepsilon \to 0} \left(\lim_{N \to \infty} \frac{\sup_{y \in Y} \operatorname{Widim}_{\varepsilon} \left(\pi^{-1}(y), d_N \right)}{N} \right).$$

It is easy to check that the quantity $\sup_{y \in Y} \operatorname{Widim}_{\varepsilon}(\pi^{-1}(y), d_N)$ is sub-additive in N and monotone in ε . So the above limits exist.

The next proposition clarifies the meaning of this definition.

Proposition 1.1. Let $\pi : (X,T) \to (Y,S)$ be a factor map between dynamical systems. Then

$$\operatorname{mdim}(\pi, T) = \sup_{y \in Y} \operatorname{\overline{mdim}} \left(\pi^{-1}(y), T \right) = \sup_{y \in Y} \operatorname{\underline{mdim}} \left(\pi^{-1}(y), T \right).$$

So we can say that the relative mean dimension $\operatorname{mdim}(\pi, T)$ properly measures the mean dimension of fibers of π .

Now we formally state the main problem we study in the paper:

Problem 1.2. Let $\pi : (X,T) \to (Y,S)$ be a factor map between dynamical systems. Does the following inequality hold true?

(1.1)
$$\operatorname{mdim}(X,T) \le \operatorname{mdim}(Y,S) + \operatorname{mdim}(\pi,T).$$

This problem was originally posed by the author in [Tsu08, Problem 4.8] more than ten years ago. He encountered it when he studied mean dimension of certain dynamical systems coming from geometric analysis. At that time he spent a lot of time trying to prove the inequality (1.1), but he did not succeed. He could only prove an inequality much weaker than (1.1) under a rather artificial assumption [Tsu08, Theorem 4.6].

Recently Liang [Lia21] revisited this problem from a new angle. He proved, among other things, that the inequality $(1\cdot 1)$ holds true in the category of *algebraic actions*¹ [Lia21, Corollary 2.18]. See also [LL18, Corollary 6.1] for a closely related result.

We will give both (partially) positive and negative answers to Problem 1.2 below.

1.2. Main results. Our first main result shows a negative answer to Problem 1.2 in a rather strong sense:

Theorem 1.3. For any positive number δ there exists a factor map between dynamical systems

$$\pi: (X,T) \to (Y,S)$$

satisfying

 $\operatorname{mdim}(X,T) = 1$, $\operatorname{mdim}(Y,S) < \delta$, $\operatorname{mdim}(\pi,T) = 0$.

In particular, letting $\delta < 1$, this shows that the inequality

$$\operatorname{mdim}(X,T) \le \operatorname{mdim}(Y,S) + \operatorname{mdim}(\pi,T)$$

does not hold in general.

Remark 1.4. In the above statement, we consider the condition $\operatorname{mdim}(X, T) = 1$. This is just for simplicity, and indeed we can make $\operatorname{mdim}(X, T)$ arbitrary large as follows: Let n be a natural number and δ a positive number. By Theorem 1.3 there exists a factor map $\pi : (X, T) \to (Y, S)$ satisfying

$$\operatorname{mdim}(X,T) = 1$$
, $\operatorname{mdim}(Y,S) < \frac{\delta}{n}$, $\operatorname{mdim}(\pi,T) = 0$.

Then it is easy to check that the factor map $\pi: (X, T^n) \to (Y, S^n)$ satisfies

 $\mathrm{mdim}(X,T^n)=n,\quad \mathrm{mdim}(Y,S^n)<\delta,\quad \mathrm{mdim}(\pi,T^n)=0.$

Furthermore we can even prove that for any positive number δ there exists a factor map $\pi': (X', T') \to (Y', S')$ satisfying

 $\mathrm{mdim}(X',T')=\infty,\quad \mathrm{mdim}(Y',S')<\delta,\quad \mathrm{mdim}(\pi',T')=0.$

¹Indeed, in the category of algebraic actions, the inequality $(1\cdot 1)$ becomes an equality. This is also proved in [Lia21, Corollary 2.18].

See Remarks 3.10 and 3.11 in §3.2 for further discussions.

Some readers might wonder whether one can even require $\operatorname{mdim}(Y, S) = 0$ instead of $\operatorname{mdim}(Y, S) < \delta$ in the statement of Theorem 1.3. However this turns out to be impossible. This is our second main result:

Theorem 1.5. Let $\pi : (X,T) \to (Y,S)$ be a factor map between dynamical systems. If $\operatorname{mdim}(Y,S) = 0$ then

 $\operatorname{mdim}(X, T) = \operatorname{mdim}(\pi, T).$

Namely, an analogue of the Hurewicz theorem holds true if the base system (Y, S) has zero mean dimension. This provides a partially positive answer to Problem 1.2.

The proof of Theorem 1.3 is based on a result of Gromov [Gro88] and its variations developed in §3.1 below. A main ingredient of the proof of Theorem 1.5 is Lindenstrauss–Weiss' theory of small boundary property.

Acknowledgment. I would like to deeply thank Tom Meyerovitch. The conversation with him revived my interest in Problem 1.2. We discussed the problem together. He generously declined to be a coauthor of this paper. But his influence is visible in several places of the paper. In particular I came up with Remark 3.11 thanks to his suggestion.

2. PROOF OF PROPOSITION 1.1

2.1. Preparations on ε -width dimension. Here we prepare some simple results on ε -width dimension. Let (X, d) and (Y, d') be metric spaces. We consider its product

 $(X, d) \times (Y, d') = (X \times Y, d \times d'),$

where $d \times d'$ is a metric on $X \times Y$ defined by

$$d \times d'((x_1, y_1), (x_2, y_2)) = \max(d(x_1, x_2), d'(y_1, y_2)).$$

Lemma 2.1. For any $\varepsilon > 0$ we have

$$\operatorname{Widim}_{\varepsilon} (X \times Y, d \times d') \leq \operatorname{Widim}_{\varepsilon}(X, d) + \operatorname{Widim}_{\varepsilon}(Y, d').$$

Proof. If $f : (X, d) \to K$ and $g : (Y, d') \to L$ are both ε -embeddings then $f \times g : (X \times Y, d \times d') \to K \times L$ is also an ε -embedding.

We say that a map $f: X \to Y$ is **distance non-decreasing** if for every $x_1, x_2 \in X$ we have

$$d(x_1, x_2) \le d'(f(x_1), f(x_2)) .$$

Lemma 2.2. If there exists a distance non-decreasing continuous map $f : X \to Y$ then for any $\varepsilon > 0$

$$\operatorname{Widim}_{\varepsilon}(X, d) \leq \operatorname{Widim}_{\varepsilon}(Y, d').$$

Proof. If $g: Y \to K$ is an ε -embedding then $g \circ f: X \to K$ is also.

2.2. Proof of Proposition 1.1. Let $\pi : (X,T) \to (Y,S)$ be a factor map between dynamical systems. Here we prove Proposition 1.1. It is obvious from the definitions that

$$\sup_{y \in Y} \underline{\mathrm{mdim}}\left(\pi^{-1}(y), T\right) \le \sup_{y \in Y} \overline{\mathrm{mdim}}\left(\pi^{-1}(y), T\right) \le \mathrm{mdim}(\pi, T).$$

So it is enough to prove

$$\operatorname{mdim}(\pi, T) \le \sup_{y \in Y} \operatorname{mdim}\left(\pi^{-1}(y), T\right)$$

Take any positive number a with

$$\sup_{y \in Y} \underline{\mathrm{mdim}}\left(\pi^{-1}(y), T\right) < a.$$

We will show $\operatorname{mdim}(\pi, T) \leq a$.

Let ε be any positive number. For every $y \in Y$ there exists $N_y > 0$ satisfying $\operatorname{Widim}_{\varepsilon} (\pi^{-1}(y), d_{N_y}) < aN_y$.

Claim 2.3. We can find an open neighborhood U_y of $\pi^{-1}(y)$ satisfying Widim_{ε} $(U_y, d_{N_y}) < aN_y$.

Proof. Take an ε -embedding $f : \pi^{-1}(y) \to K$ with a simplicial complex K of dimension smaller than aN_y . Since a simplicial complex is ANR (absolute neighborhood retract), we can find an open neighborhood U_y of $\pi^{-1}(y)$ and extend f to a continuous map $f : U_y \to K$. If we choose U_y sufficiently small, then $f : U_y \to K$ is also an ε -embedding and we have $\operatorname{Widim}_{\varepsilon}(U_y, d_{N_y}) < aN_y$.

There exists an open neighborhood V_y of y satisfying $\pi^{-1}(V_y) \subset U_y$. Then we have $\operatorname{Widim}_{\varepsilon}(\pi^{-1}(V_y), d_{N_y}) < aN_y$.

Since Y is compact, we can find an open cover $Y = V_1 \cup V_2 \cup \cdots \cup V_m$ and natural numbers N_1, N_2, \ldots, N_m satisfying $\operatorname{Widim}_{\varepsilon}(\pi^{-1}(V_i), d_{N_i}) < aN_i$ for all $1 \leq i \leq m$. Set

$$\bar{N} := \max_{1 \le i \le m} N_i$$

Claim 2.4. For every $y \in Y$ and natural number N we have

Widim_{$$\varepsilon$$} $(\pi^{-1}(y), d_N) < a(N + \overline{N}).$

Proof. We choose a sequence i_1, i_2, \ldots, i_k such that

$$y \in V_{i_1}, \quad S^{N_{i_1}} y \in V_{i_2}, \quad S^{N_{i_1}+N_{i_2}} y \in V_{i_3}, \quad \dots \quad , S^{N_{i_1}+N_{i_2}+\dots+N_{i_{k-1}}} y \in V_{i_k},$$
$$N_{i_1} + N_{i_2} + \dots + N_{i_{k-1}} < N \le N_{i_1} + N_{i_2} + \dots + N_{i_k}.$$

Consider a map

$$f: (\pi^{-1}(y), d_N) \to (\pi^{-1}(V_{i_1}), d_{N_{i_1}}) \times (\pi^{-1}(V_{i_2}), d_{N_{i_2}}) \times \dots \times (\pi^{-1}(V_{i_k}), d_{N_{i_k}}),$$

defined by $f(x) = (x, T^{N_{i_1}}x, T^{N_{i_1}+N_{i_2}}x, \dots, T^{N_{i_1}+N_{i_2}+\dots+N_{i_{k-1}}}x)$. This is a distance nondecreasing continuous map. Hence by Lemmas 2.1 and 2.2 we have

Widim_{$$\varepsilon$$} $\left(\pi^{-1}(y), d_N\right) \leq \sum_{j=1}^k \text{Widim}_{\varepsilon} \left(\pi^{-1}(V_{i_j}), d_{N_{i_j}}\right)$
 $< a \sum_{j=1}^k N_{i_j} < a(N + \bar{N}).$

Thus we have

$$\sup_{y \in Y} \operatorname{Widim}_{\varepsilon} \left(\pi^{-1}(y), d_N \right) \le a(N + \bar{N}).$$

Hence

$$\lim_{N \to \infty} \frac{\sup_{y \in Y} \operatorname{Widim}_{\varepsilon} \left(\pi^{-1}(y), d_N \right)}{N} \le a.$$

Since $\varepsilon > 0$ is arbitrary, we have

$$\operatorname{mdim}\left(\pi,T\right) = \lim_{\varepsilon \to 0} \left(\lim_{N \to \infty} \frac{\sup_{y \in Y} \operatorname{Widim}_{\varepsilon}\left(\pi^{-1}(y), d_{N}\right)}{N}\right) \le a$$

This proves Proposition 1.1.

3. Proof of Theorem 1.3

3.1. Variations of Gromov's lemma. Gromov proved the following statement in [Gro88, p.107, (H''_1) Example]. This shows that a direct analogue of the Hurewicz theorem does not hold for ε -width dimension.

Lemma 3.1 (Gromov 1988). Let (X, d) be a (2n + 1)-dimensional compact Riemannian manifold. For any positive number ε there exists a smooth map $f : X \to [0, 1]$ such that for every $t \in [0, 1]$ we have

Widim_{$$\varepsilon$$} $(f^{-1}(t), d) \le n.$

The purpose of this subsection is to develop some variations of this lemma. The main results are Lemma 3.4 and Corollary 3.5 below. We do not explain the proof of Lemma 3.1 itself because we will provide a full proof of a more detailed version below. Our argument follows Gromov's idea.

We need to prepare some basic terminologies of simplicial complex. (Recall that we always assume that a simplicial complex has only finitely many simplices.) For a simplicial complex K we denote by V(K) the set of vertices of K.

Definition 3.2. Let K be a simplicial complex and $L \subset K$ a subcomplex. L is said to be a **full subcomplex** of K if for every simplex $\Delta \subset K$ with $V(\Delta) \subset L$ we have $\Delta \subset L$.

There is a one-to-one correspondence between full subcomplexes of K and a subset of V(K). For a subset $A \subset V(K)$ we denote by K(A) the (unique) full subcomplex of K satisfying V(K(A)) = A. If $A = \emptyset$ (empty set), then $K(A) = \emptyset$.

For a vertex $v \in K$ we define the **star** St(v) by

 $\operatorname{St}(v) = \bigcup \left\{ \operatorname{Int}(\Delta) \, | \, \Delta \text{ is a simplex of } K \text{ with } v \in \Delta \right\},$

where $Int(\Delta)$ is the interior of Δ defined by (letting $V(\Delta) = \{p_0, p_1, \dots, p_n\}$)

$$\operatorname{Int}(\Delta) = \left\{ \sum_{i=0}^{n} t_i p_i \, \middle| \, t_0 + t_1 + \dots + t_n = 1, \, t_i > 0 \; (\forall 0 \le i \le n) \right\}.$$

When $\Delta = \{v\}$ (just one point) then its interior is $\{v\}$. So in particular $\operatorname{St}(v)$ contains v. Indeed it is easy to check that $\operatorname{St}(v)$ is an open neighborhood of v. So the stars $\operatorname{St}(v)$ $(v \in V(K))$ form an open covering of K.

Let *m* be a natural number. We denote the standard basis of \mathbb{R}^m by e_1, e_2, \ldots, e_m . We define the standard (m-1)-dimensional simplex $\Delta^{m-1} \subset \mathbb{R}^m$ by

$$\Delta^{m-1} = \left\{ \sum_{i=1}^{m} t_i e_i \, \middle| \, t_1 + t_2 + \dots + t_m = 1, \, t_i \ge 0 \; (\forall 1 \le i \le m) \right\}.$$

The following lemma is our first variation of Gromov's lemma.

Lemma 3.3. Let K be a simplicial complex with a metric d. Let ε be a positive number satisfying

$$\max_{v \in V(K)} \operatorname{Diam} \operatorname{St}(v) < \varepsilon.$$

Let m be a natural number. Suppose we are given a partition

$$V(K) = A_1 \cup A_2 \cup \dots \cup A_m \quad (disjoint \ union).$$

Then there exists a simplicial map $f: K \to \Delta^{m-1}$ such that for every $p \in \Delta^{m-1}$ we have

Widim_{$$\varepsilon$$} $(f^{-1}(p), d) \leq \max_{1 \leq i \leq m} \dim K(A_i).$

Here $K(A_i)$ is the full subcomplex of K corresponding to A_i .

Proof. We define a simplicial map $f : K \to \Delta^{m-1}$ by the condition $f(A_i) = \{e_i\}$ and extending it linearly. Namely, if $x \in K$ has the form

$$x = \sum_{i=1}^{m} \left(\sum_{u \in A_i} x_u u \right),$$

where x_u are nonnegative numbers with $\sum_{i=1}^m \sum_{u \in A_i} x_u = 1$, we define

$$f(x) = \sum_{i=1}^{m} \left(\sum_{u \in A_i} x_u \right) e_i.$$

Let $p = \sum_{i=1}^{m} t_i e_i \in \Delta^{m-1}$. We assume, say, $t_1 > 0$. (Other cases can be treated similarly.) If $x = \sum_{i=1}^{m} (\sum_{u \in A_i} x_u u) \in f^{-1}(p)$ then

$$f(x) = \sum_{i=1}^{m} \left(\sum_{u \in A_i} x_u \right) e_i = \sum_{i=1}^{m} t_i e_i,$$

and hence

$$\sum_{u \in A_1} x_u = t_1 > 0.$$

We define a map $g: f^{-1}(p) \to K(A_1)$ by

$$g(x) = \frac{\sum_{u \in A_1} x_u u}{\sum_{u \in A_1} x_u}$$

Every fiber of g is contained in some star St(v) of $v \in A_1$. So its diameter is smaller than ε . Hence g is an ε -embedding. Therefore

Widim_{$$\varepsilon$$} $(f^{-1}(p), d) \leq \dim K(A_1).$

We need to recall the terminologies on barycentric subdivision. Let K be a simplicial complex. For each simplex $\Delta \subset K$ we denote the barycenter of Δ by $bc(\Delta)$. (Namely, if $V(\Delta) = \{p_0, p_1, \ldots, p_n\}$ then $bc(\Delta) = (p_0 + p_1 + \cdots + p_n)/(n+1)$.)

We define the **barycentric subdivision** K' of K by the following two conditions.

- $V(K') := \{ \operatorname{bc}(\Delta) | \Delta \subset K : \operatorname{simplex} \}.$
- If $\Delta_0 \subset \Delta_1 \subset \cdots \subset \Delta_n$ is a flag of mutually distinct simplices of K then $bc(\Delta_0), bc(\Delta_1), \ldots, bc(\Delta_n)$ form an *n*-simplex in K'.

As a topological space, K' is naturally identified with K. See Figure 1.




FIGURE 1. The 2-simplex (left) and its barycentric subdivision (right).

The next lemma is our second variation of Gromov's lemma.

Lemma 3.4. Let K be a simplicial complex with a metric d. Let ε be a positive number and m a natural number. After subdividing K sufficiently fine, we can find a simplicial map $f: K \to \Delta^{m-1}$ such that for every point $p \in \Delta^{m-1}$

Widim_{$$\varepsilon$$} $\left(f^{-1}(p), d\right) \leq \frac{\dim K}{m}.$

Proof. By subdividing K sufficiently fine, we can assume that

$$\max_{v \in V(K)} \operatorname{Diam} \operatorname{St}(v) < \varepsilon.$$

Let K' be the barycentric subdivision of K. Then for each simplex $\Delta \subset K$ we have a vertex $bc(\Delta)$ of K'. We define a partition $V(K') = A_1 \cup A_2 \cup \cdots \cup A_m$ (disjoint union) by

$$A_{1} = \left\{ \operatorname{bc}(\Delta) \middle| \Delta \text{ is a simplex of } K \text{ with } \dim \Delta \leq \frac{\dim K}{m} \right\},$$
$$A_{i} = \left\{ \operatorname{bc}(\Delta) \middle| \Delta \text{ is a simplex of } K \text{ with } \frac{(i-1)\dim K}{m} < \dim \Delta \leq \frac{i\dim K}{m} \right\}, \quad (2 \leq i \leq m).$$

Consider the full subcomplexes $K'(A_i)$ $(1 \le i \le m)$ corresponding to A_i . Every simplex of $K'(A_1)$ corresponds to a flag $\Delta_0 \subset \Delta_1 \subset \cdots \subset \Delta_n$ of distinct simplices of K of dimension $\le \frac{\dim K}{m}$. The length of such a flag is at most $\frac{\dim K}{m} + 1$. So we have

$$\dim K'(A_1) \le \frac{\dim K}{m}$$

Similarly

$$\dim K'(A_i) < \frac{\dim K}{m}, \quad (2 \le i \le m).$$

Figure 2 shows the case that K is two dimensional and m = 2.




FIGURE 2. This shows the construction in the case that K is two dimensional and m = 2. The left is a two-dimensional simplicial complex K. The right is $K'(A_1)$ and $K'(A_2)$. $K'(A_1)$ is the one-dimensional skeleton of K (more precisely, $K'(A_1)$ is the barycentric subdivision of the onedimensional skeleton of K). The vertices of $K'(A_1)$ is depicted as dots. $K'(A_2)$ consists of five points depicted as *, which are the barycenters of two-dimensional simplices of K. So $K'(A_1)$ is one dimensional and $K'(A_2)$ is zero dimensional.

Then by Lemma 3.3 we can find a simplicial map $f: K' \to \Delta^{m-1}$ such that for every point $p \in \Delta^{m-1}$ we have

Widim_{$$\varepsilon$$} $(f^{-1}(p), d) \le \max_{1 \le i \le m} \dim K'(A_i) \le \frac{\dim K}{m}.$

The following corollary is a crucial ingredient of the proof of Theorem 1.3 in the next subsection.

Corollary 3.5. Let K be a simplicial complex with a metric d. For any positive number ε and any natural number m there exists a continuous map $F: K \to [0,1]^{m-1}$ such that for every point $p \in [0,1]^{m-1}$ we have

Widim_{$$\varepsilon$$} $\left(F^{-1}(p), d\right) \leq \frac{\dim K}{m}.$

Proof. The (m-1) dimensional cube $[0, 1]^{m-1}$ is homeomorphic to the (m-1) dimensional simplex Δ^{m-1} . So the statement follows from Lemma 3.4.

Remark 3.6. The factor of $\frac{1}{m}$ in the statements of Lemma 3.4 and Corollary 3.5 is optimal, according to [Gro88, p.107, Corollaries (H'_1)]. It says that if $f: X \to [0, 1]^{m-1}$ is a continuous map from a compact metric space (X, d) then for any $\varepsilon > 0$

$$\sup_{p \in [0,1]^{m-1}} \operatorname{Widim}_{\varepsilon} \left(f^{-1}(p), d \right) \ge \frac{\operatorname{Widim}_{\varepsilon}(X, d) - m + 1}{m}.$$

Remark 3.7. Lemma 3.4 and Corollary 3.5 show a rather counter-intuitive phenomena. The following example illustrates its significance: Let K be a simplicial complex of

dim K = 1000,000 (one million).

Let m = 1001. From Corollary 3.5, for any positive number ε , there exists a continuous map $F: K \to [0, 1]^{1000}$ such that for all $p \in [0, 1]^{1000}$

Widim_{$$\varepsilon$$} $(F^{-1}(p), d) < 1000.$

Hence every fiber of F looks like a space whose dimension is smaller than 1000 (up to distortion bounded by ε). The range of F is also 1000 dimensional. However the total space K has dimension one million!

3.2. **Proof of Theorem 1.3.** The purpose of this subsection is to prove Theorem 1.3. Consider the two-sided infinite product of copies of the unit interval [0, 1]:

$$[0,1]^{\mathbb{Z}} = \cdots \times [0,1] \times [0,1] \times [0,1] \times \cdots$$

We define a metric ρ on $[0,1]^{\mathbb{Z}}$ by

$$\rho\left((x_n)_{n\in\mathbb{Z}},(y_n)_{n\in\mathbb{Z}}\right) = \sum_{n\in\mathbb{Z}} 2^{-|n|} |x_n - y_n|.$$

We define the shift map $\sigma:[0,1]^{\mathbb{Z}}\rightarrow [0,1]^{\mathbb{Z}}$ by

$$\sigma\left((x_n)_{n\in\mathbb{Z}}\right) = (x_{n+1})_{n\in\mathbb{Z}}.$$

The pair $([0,1]^{\mathbb{Z}},\sigma)$ is a dynamical system.

Let (Z, R) be a zero dimensional free minimal dynamical system. Here "zero dimensional" means that Z is totally disconnected, "free" means that it has no periodic point, and "minimal" means that every orbit is dense in Z. (Indeed, the following argument works well without the minimality assumption. But we assume it for simplicity of the explanation.) It is a standard fact that such a dynamical system exists. For example, we can construct it as a subshift of $(\{0,1\}^{\mathbb{Z}}, \text{shift})$. We take a metric ρ' on Z.

We define a dynamical system (X, T) as the product of $([0, 1]^{\mathbb{Z}}, \sigma)$ and (Z, R):

$$(X,T) := ([0,1]^{\mathbb{Z}} \times Z, \sigma \times R).$$

We define a metric d on $X = [0, 1]^{\mathbb{Z}} \times Z$ by

$$d((x, z), (x', z')) = \max(\rho(x, x'), \rho'(z, z')).$$

The mean dimension of (X, T) is one:

$$\mathrm{mdim}(X,T) = 1.$$

Throughout this subsection we fix $(X,T) = ([0,1]^{\mathbb{Z}} \times Z, \sigma \times R)$ and construct factor maps from this (X,T).

The next proposition is a preliminary version of Theorem 1.3.

Proposition 3.8. For any positive numbers ε and δ there exist a dynamical system (Y, S) and a factor map $\pi : (X, T) \to (Y, S)$ such that

$$\operatorname{mdim}(Y,S) < \delta, \quad \lim_{N \to \infty} \frac{\sup_{y \in Y} \operatorname{Widim}_{\varepsilon} (\pi^{-1}(y), d_N)}{N} < \delta.$$

Proof. We prepare ingredients of the construction:

- Fix a natural number m with $\frac{1}{m} < \delta$.
- Fix a natural number L > m with $\frac{m}{L} < \frac{\delta}{2}$.
- Since (Z, R) is free and zero dimensional, we can take a non-empty clopen set $U \subset Z$ satisfying $U \cap R^{-n}U = \emptyset$ for all $1 \leq n \leq L$. Here "clopen" means that U is both closed and open. Since (Z, R) is minimal, there exists a natural number L' > L satisfying

$$Z = \bigcup_{n=1}^{L'-1} R^{-n} U.$$

• For $z \in Z$ we set $E(z) = \{n \in \mathbb{Z} \mid R^n z \in U\}$. For any two distinct points $a, b \in E(z)$ we have |b - a| > L. Moreover for any $a \in E(z)$ there exists $b \in E(z)$ satisfying a + L < b < a + L'.

• For each natural number n, we apply Corollary 3.5 to the n-dimensional cube $[0,1]^n$. Then we find a continuous map $F_n: [0,1]^n \to [0,1]^{m-1}$ such that for every point $p \in [0,1]^{m-1}$ we have

Widim_{$$\varepsilon/4$$} $\left(F_n^{-1}(p), \left\|\cdot\right\|_{\infty}\right) \le \frac{n}{m} \left(<\delta n\right).$

Here a metric on $[0,1]^n$ is given by the ℓ^{∞} -norm $||x||_{\infty} = \max_{1 \le i \le n} |x_i|$.

• For $n \ge m$ we define $G_n : [0,1]^n \to [0,1]^n$ by

$$G_n(x) = \left(F_n(x), \underbrace{0, 0, \dots, 0, 0}_{n-m+1}\right).$$

Notice that if n is large then the proportion of non-zero entries of $G_n(x)$ is very small. For every point $p \in [0, 1]^n$ we have

Widim_{$$\varepsilon/4$$} $\left(G_n^{-1}(p), \|\cdot\|_{\infty}\right) \leq \frac{n}{m} (<\delta n).$

We will construct an equivariant continuous map $f: (X,T) \to ([0,1]^{\mathbb{Z}},\sigma)$ by using the above data. For a point $x = (x_n)_{n \in \mathbb{Z}} \in [0,1]^{\mathbb{Z}}$ and integers a < b, we denote

$$x|_{[a,b)} := (x_a, x_{a+1}, x_{a+2}, \dots, x_{b-1}).$$

Let $(x, z) \in [0, 1]^{\mathbb{Z}} \times Z = X$. Take any point $a \in E(z)$ and set $b := \min(E(z) \cap (a, \infty))$. We have L < b - a < L'. We define

$$f(x,z)|_{[a,b)} := G_{b-a}(x|_{[a,b)}) \in [0,1]^{b-a}.$$

We consider this for every $a \in E(z)$. Then we have defined $f(x, z) \in [0, 1]^{\mathbb{Z}}$. The map f is equivariant and continuous. (The continuity follows from the clopenness of U.) We define $\pi : X \to [0, 1]^{\mathbb{Z}} \times Z$ by

$$\pi(x,z) = (f(x,z),z).$$

We set $Y = \pi(X)$, which becomes a dynamical system under the map $S := \sigma \times R$. We will show that a factor map $\pi : (X, T) \to (Y, S)$ satisfies the statement.

First we estimate the mean dimension of (Y, S). Let $(x, z) \in [0, 1]^{\mathbb{Z}} \times Z$ and let N be a natural number. Denote

$$[0, N) \cap E(z) = \{a_1 < a_2 < a_3 < \dots < a_k\}.$$

Since $a_{i+1} - a_i > L$, we have $k - 1 < \frac{N}{L}$. Set

$$a_0 := \max (E(z) \cap (-\infty, 0)), \quad a_{k+1} := \min (E(z) \cap [N, \infty)).$$

Then we have

$$[0,N) \subset \bigcup_{i=0}^{k} [a_i, a_{i+1}) = [a_0, a_{k+1}).$$

For $0 \le i \le k$

$$f(x,z)|_{[a_i,a_{i+1})} = G_{a_{i+1}-a_i}\left(x|_{[a_i,a_{i+1})}\right).$$

The number of its non-zero entries is at most m-1. Therefore

The number of non-zero entries of $f(x,z)|_{[0,N)} \le (k+1)(m-1)$

$$< \left(\frac{N}{L} + 2\right)(m-1)$$
$$< \frac{\delta N}{2} + 2m \quad \text{by } \frac{m}{L} < \frac{\delta}{2}.$$

Denote by $\Pi_N : [0,1]^{\mathbb{Z}} \to [0,1]^N$ the projection to the $0, 1, 2, \ldots, (N-1)$ -th coordinates. Then $\Pi_N(f(X))$ is contained in

$$\left\{ (y_0, y_1, \dots, y_{N-1}) \, \middle| \, y_n = 0 \text{ except for at most } \frac{\delta N}{2} + 2m \text{ entries} \right\}.$$

whose dimension is at most $\frac{\delta N}{2} + 2m$. This implies that

$$\operatorname{mdim}\left(f(X),\sigma\right) \leq \lim_{N \to \infty} \left(\frac{\delta N}{2} + 2m\right)/N = \frac{\delta}{2} < \delta.$$

Since Z is zero dimensional, we have

$$\operatorname{mdim}(Y, S) = \operatorname{mdim}(f(X), \sigma) < \delta.$$

Next we study the fibers of π . Fix M > 0 with $\sum_{|n| \ge M} 2^{-|n|} < \frac{\varepsilon}{2}$. Let $y = (p, z) \in Y$ with $p \in [0, 1]^{\mathbb{Z}}$ and $z \in Z$. Let N be a natural number. Denote

$$(-M, N+M) \cap E(z) = \{a_1 < a_2 < \dots < a_k\},\$$

and set

$$a_0 := \max (E(z) \cap (-\infty, -M]), \quad a_{k+1} := \min (E(z) \cap [N+M, \infty)).$$

Then we have

$$(-M, N+M) \subset \bigcup_{i=0}^{k} [a_i, a_{i+1}) = [a_0, a_{k+1}).$$

Since $a_{i+1} - a_i < L'$, we have $a_{k+1} - a_0 < N + 2M + 2L'$. For any $0 \le i \le k$ and $(x, z) \in \pi^{-1}(y)$ we have

$$f(x,z)|_{[a_i,a_{i+1})} = G_{a_{i+1}-a_i}\left(x|_{[a_i,a_{i+1})}\right) = p|_{[a_i,a_{i+1})}.$$

We define a projection $\Pi_{[a_0,a_{k+1})}: [0,1]^{\mathbb{Z}} \times Z \to [0,1]^{a_{k+1}-a_0}$ by

$$\Pi_{[a_0,a_{k+1})}\left((x_n)_{n\in\mathbb{Z}},z\right) = (x_n)_{n\in[a_0,a_{k+1})}$$

Then

$$\Pi_{[a_0,a_{k+1})}\left(\pi^{-1}(y)\right) \subset G_{a_1-a_0}^{-1}\left(p|_{[a_0,a_1)}\right) \times G_{a_2-a_1}^{-1}\left(p|_{[a_1,a_2)}\right) \times \cdots \times G_{a_{k+1}-a_k}^{-1}\left(p|_{[a_k,a_{k+1})}\right).$$

It follows that

Widim_{ε} $\left(\pi^{-1}(y), d_N\right) \leq$ Widim_{$\varepsilon/4$} $\left(\Pi_{[a_0, a_{k+1})} \left(\pi^{-1}(y)\right), \|\cdot\|_{\infty}\right)$ by the definition of the metric d $\leq \sum_{i=0}^{k} \text{Widim}_{\varepsilon/2} \left(G_{a_{i+1}-a_{i}}^{-1} \left(p|_{[a_{i}, a_{i+1})}\right), \|\cdot\|_{\infty}\right)$ $\leq \sum_{i=0}^{k} \frac{a_{i+1}-a_{i}}{m}$ $= \frac{a_{k+1}-a_{0}}{m} < \frac{N+2M+2L'}{m}.$

This holds for every $y \in Y$. So we get

 $\sup_{y \in Y} \operatorname{Widim}_{\varepsilon} \left(\pi^{-1}(y), d_N \right) < \frac{N + 2M + 2L'}{m}.$

Notice that M and L' are independent of N. Thus we conclude

$$\lim_{N \to \infty} \frac{\sup_{y \in Y} \operatorname{Widim}_{\varepsilon} (\pi^{-1}(y), d_N)}{N} \le \frac{1}{m} < \delta.$$

Now we are ready to prove Theorem 1.3.

Theorem 3.9. For any positive number δ there exists a dynamical system (Y, S) and a factor map $\pi : (X, T) \to (Y, S)$ such that

$$\operatorname{mdim}(Y, S) < \delta, \quad \operatorname{mdim}(\pi, T) = 0.$$

Recall that the dynamical system $(X,T) = ([0,1]^{\mathbb{Z}} \times Z, \sigma \times R)$ has mean dimension one. So Theorem 1.3 follows from this theorem.

Proof. For each natural number n we apply Proposition 3.8 to $\varepsilon = \frac{1}{n}$ and $\frac{\delta}{2^n}$. Then we find a factor map $\pi_n : (X, T) \to (Y_n, S_n)$ such that

$$\operatorname{mdim}(Y_n, S_n) < \frac{\delta}{2^n}, \quad \lim_{N \to \infty} \frac{\sup_{y \in Y_n} \operatorname{Widim}_{1/n} (\pi_n^{-1}(y), d_N)}{N} < \frac{\delta}{2^n}$$

Define

$$\pi := \pi_1 \times \pi_2 \times \pi_3 \times \cdots : X \to Y_1 \times Y_2 \times Y_3 \times \cdots, \quad x \mapsto (\pi_1(x), \pi_2(x), \pi_3(x), \ldots).$$

Set $Y = \pi(X) \subset Y_1 \times Y_2 \times Y_3 \times \cdots$ with a map $S := S_1 \times S_2 \times S_3 \times \cdots$. The pair (Y, S) is a dynamical system. We show that the factor map $\pi : (X, T) \to (Y, S)$ satisfies the statement.

The mean dimension of (Y, S) is bounded by

$$\operatorname{mdim}(Y,S) \le \sum_{n=1}^{\infty} \operatorname{mdim}(Y_n,S_n) < \sum_{n=1}^{\infty} \frac{\delta}{2^n} = \delta.$$

Let $y = (y_1, y_2, y_3, ...) \in Y (y_n \in Y_n)$. We have

$$\pi^{-1}(y) = \bigcap_{n=1}^{\infty} \pi_n^{-1}(y_n).$$

Then for any natural numbers n and N

Widim_{1/n}
$$(\pi^{-1}(y), d_N) \leq \text{Widim}_{1/n} (\pi_n^{-1}(y_n), d_N).$$

Hence

$$\lim_{N \to \infty} \frac{\sup_{y \in Y} \operatorname{Widim}_{1/n} \left(\pi^{-1}(y), d_N \right)}{N} \leq \lim_{N \to \infty} \frac{\sup_{y_n \in Y_n} \operatorname{Widim}_{1/n} \left(\pi^{-1}_n(y_n), d_N \right)}{N} < \frac{\delta}{2^n}.$$

Letting $n \to \infty$, we get $\operatorname{mdim}(\pi, T) = 0.$

Remark 3.10. Let δ be a positive number. For each natural number n we apply Theorem 3.9 to $\frac{\delta}{2^n}$. Then there exists a factor map $\pi_n : (X,T) \to (Y_n,S_n)$ such that

$$\operatorname{mdim}(Y_n, S_n) < \frac{\delta}{2^n}, \quad \operatorname{mdim}(\pi_n, T) = 0.$$

We set

$$(X',T') := (X,T) \times (X,T) \times (X,T) \times \cdots, \quad (Y',S') := (Y_1,S_1) \times (Y_2,S_2) \times (Y_3,S_3) \times \cdots.$$

We define a factor map $\pi' : (X',T') \to (Y',S')$ by

$$\pi'(x_1, x_2, x_3, \dots) = (\pi_1(x_1), \pi_2(x_2), \pi_3(x_3), \dots).$$

Then it is easy to check that

$$\operatorname{mdim}(X',T') = \infty, \quad \operatorname{mdim}(Y',S') < \delta, \quad \operatorname{mdim}(\pi',T') = 0.$$

See Remark 1.4 in $\S1.2$.

Remark 3.11. The dynamical system (X', T') constructed in the above remark is a rather "universal" one. It has the form

$$(X',T') = \left(\left([0,1]^{\mathbb{N}} \right)^{\mathbb{Z}}, \text{shift} \right) \times (Z',S')$$

where (Z', S') is a zero dimensional dynamical system given by

$$(Z',S') = (Z,S) \times (Z,S) \times (Z,S) \times \cdots$$

It is easy to see that *every* dynamical system embeds in $(([0,1]^{\mathbb{N}})^{\mathbb{Z}}, \text{shift})$. Hence, given an arbitrary dynamical system $(\mathcal{X}, \mathcal{T})$, we can embed $(\mathcal{X}, \mathcal{T}) \times (Z', S')$ in (X', T'). Therefore, for any positive number δ , there exists a factor map

$$\Pi: (\mathcal{X} \times Z', \mathcal{T} \times S') \to (\mathcal{Y}, \mathcal{S})$$

satisfying

$$\operatorname{mdim}\left(\mathcal{Y},\mathcal{S}\right) < \delta, \quad \operatorname{mdim}\left(\Pi,\mathcal{T}\right) = 0.$$

Notice that

$$\operatorname{mdim}\left(\mathcal{X}\times Z', \mathcal{T}\times S'\right) = \operatorname{mdim}\left(\mathcal{X}, \mathcal{T}\right)$$

can be an arbitrary nonnegative number. Thus we can say that the construction of this section shows an universal phenomena.

4. Proof of Theorem 1.5

4.1. **Preliminaries on relative mean dimension.** Here we prepare some simple facts on the relative mean dimension.

Lemma 4.1. Let $\pi_i : (X_i, T_i) \to (Y_i, S_i)$ (i = 1, 2) be two factor maps between dynamical systems. We consider their product:

$$\pi_1 \times \pi_2 : (X_1 \times X_2, T_1 \times T_2) \to (Y_1 \times Y_2, S_1 \times S_2).$$

For this factor map we have

$$\text{mdim} (\pi_1 \times \pi_2, T_1 \times T_2) \leq \text{mdim}(\pi_1, T_1) + \text{mdim}(\pi_2, T_2), \\ \text{mdim} (\pi_1 \times \pi_2, T_1 \times T_2) \geq \max (\text{mdim}(\pi_1, T_1), \text{mdim}(\pi_2, T_2)).$$

Proof. Let d and d' be metrics on X_1 and X_2 respectively. The product space $X \times X'$ has a metric $d \times d'$. (See §2.1.)

Let $(y_1, y_2) \in Y_1 \times Y_2$. For every natural number N we have

$$\left((\pi_1 \times \pi_2)^{-1}(y_1, y_2), (d \times d')_N\right) = \left(\pi_1^{-1}(y_1), d_N\right) \times \left(\pi_2^{-1}(y_2), d'_N\right).$$

By Lemma 2.1, for $\varepsilon > 0$

$$\operatorname{Widim}_{\varepsilon}\left((\pi_1 \times \pi_2)^{-1}(y_1, y_2), (d \times d')_N\right) \leq \operatorname{Widim}_{\varepsilon}\left(\pi_1^{-1}(y_1), d_N\right) + \operatorname{Widim}_{\varepsilon}\left(\pi_2^{-1}(y_2), d'_N\right).$$

Thus we have the first inequality.

Fix $p \in \pi_2^{-1}(y_2)$. The map

$$(\pi_1^{-1}(y_1), d_N) \to ((\pi_1 \times \pi_2)^{-1}(y_1, y_2), (d \times d')_N), \quad x \mapsto (x, p),$$

is an isometric embedding. Hence

Widim_{$$\varepsilon$$} $\left(\pi_1^{-1}(y_1), d_N\right) \leq \text{Widim}_{\varepsilon} \left((\pi_1 \times \pi_2)^{-1}(y_1, y_2), (d \times d')_N\right).$

Similarly

Widim_{$$\varepsilon$$} $\left(\pi_2^{-1}(y_2), d_N\right) \leq \text{Widim}_{\varepsilon} \left((\pi_1 \times \pi_2)^{-1}(y_1, y_2), (d \times d')_N\right).$

Then we get the second inequality.

Corollary 4.2. Let $\pi : (X,T) \to (Y,S)$ be a factor map between dynamical systems, and let (Z,R) a dynamical system. We consider

$$\pi \times \mathrm{Id} : (X \times Z, T \times R) \to (Y \times Z, S \times R), \quad (x, z) \mapsto (\pi(x), z).$$

Then we have

$$m\dim(\pi \times \mathrm{Id}, T \times R) = m\dim(\pi, T).$$

Proof. The trivial factor map

$$\mathrm{Id}: (Z,R) \to (Z,R)$$

has zero relative mean dimension. So the corollary follows from Lemma 4.1.

4.2. Small boundary property. Here we review the theory of *small boundary property* introduced by Lindenstrauss–Weiss [LW00]. Let (X,T) be a dynamical system. For a subset $A \subset X$ we define the orbit capacity of A by

$$ocap(A) = \lim_{N \to \infty} \frac{1}{N} \sup_{x \in X} \sum_{n=0}^{N-1} 1_A(T^n x).$$

The quantity $\sup_{x \in X} \sum_{n=0}^{N-1} 1_A(T^n x)$ is sub-additive in N. So we have

(4.1)
$$\operatorname{ocap}(A) = \inf_{N \ge 1} \frac{1}{N} \sup_{x \in X} \sum_{n=0}^{N-1} 1_A(T^n x).$$

It is easy to see that $ocap(A \cup B) \leq ocap(A) + ocap(B)$. In particular, if ocap(A) = ocap(B) = 0 then $ocap(A \cup B) = 0$.

The next lemma was proved in [Lin99, Lemma 6.3].

Lemma 4.3. Let $E \subset X$ be a closed subset. For any positive number δ there exists an open neighborhood U of E satisfying

$$\operatorname{ocap}(U) < \operatorname{ocap}(E) + \varepsilon.$$

Proof. This follows from the formula $(4 \cdot 1)$.

We say that a dynamical system (X, T) has the **small boundary property** if for every point $x \in X$ and for every open neighborhood U of x there exists an open set V such that

$$x \in V \subset U$$
, $\operatorname{ocap}(\partial V) = 0$.

Here ∂V is the boundary of V, namely $\partial V := \overline{V} \setminus V$. The small boundary property is a dynamical version of totally disconnectedness.

Lindenstrauss–Weiss [LW00, Theorem 5.4] proved that if a dynamical system has the small boundary property then its mean dimension is zero. Lindenstrauss [Lin99, Theorem 6.2] proved a partial converse of this statement as follows. This will be crucial in the next subsection.

Theorem 4.4 (Lindenstrauss 1999). If (X,T) is an extension of a free minimal system and mdim(X,T) = 0 then (X,T) has the small boundary property.

Here the assumption that "(X, T) is an extension of a free minimal system" means that there exists a factor map $\pi : (X, T) \to (Y, S)$ such that (Y, S) is a free minimal dynamical system.

Lemma 4.5. Let (Y, S) be a dynamical system having the small boundary property. Let δ be a positive number. For any open covering $Y = V_1 \cup V_2 \cup \cdots \cup V_m$ there exist compact subsets $E_i \subset V_i$ $(1 \le i \le m)$ such that

- $E_i \cap E_j = \emptyset$ for $i \neq j$.
- $\operatorname{ocap}(Y \setminus (E_1 \cup \cdots \cup E_m)) < \delta.$

Proof. By the small boundary property there exist open sets W_i $(1 \le i \le m)$ such that $\overline{W_i} \subset V_i$, $\operatorname{ocap}(\partial W_i) = 0$ and $Y = W_1 \cup W_2 \cup \cdots \cup W_m$. By Lemma 4.3 there exist open sets $U_i \supset \partial W_i$ with $\operatorname{ocap}(U_i) < \frac{\delta}{m}$ for $1 \le i \le m$. Then $\overline{W_i} \cup U_i = W_i \cup U_i$ is an open set. We set

$$E_1 := \overline{W_1}, \quad E_2 := \overline{W_2} \setminus (\overline{W_1} \cup U_1), \quad E_3 := \overline{W_3} \setminus (\overline{W_1} \cup U_1 \cup \overline{W_2} \cup U_2), \quad \dots,$$
$$E_m := \overline{W_m} \setminus (\overline{W_1} \cup U_1 \cup \overline{W_2} \cup U_2 \cup \dots \cup \overline{W_{m-1}} \cup U_{m-1}).$$

 E_i are compact and

$$Y \setminus (E_1 \cup \cdots \cup E_m) \subset U_1 \cup U_2 \cup \cdots \cup U_m.$$

Then

$$\operatorname{ocap}\left(Y\setminus\left(E_{1}\cup\cdots\cup E_{m}\right)\right)\leq\sum_{i=1}^{m}\operatorname{ocap}(U_{i})<\delta.$$

4.3. **Proof of Theorem 1.5.** In this subsection we first prove an analogue of the Hurewicz theorem for mean dimension in the case that the base system has the small boundary property. Next we prove it in the case that the base system has zero mean dimension.

Before stating the proposition, we prepare terminologies on *cone*. This will be used in the proof. Let K be a topological space. We define the **cone** C(K) by

$$C(K) = [0,1] \times K/\sim,$$

where $(0, x) \sim (0, y)$ for any $x, y \in K$. For $0 \le t \le 1$ and $x \in X$, the equivalence class of (t, x) is denoted by tx. The point 0x is called the **vertex** of the cone C(K) and often denoted by *.

When K is a simplicial complex, the cone C(K) naturally admits a structure of a simplicial complex. Its dimension is dim K + 1.

Let K_1, \ldots, K_m be topological spaces and let $C(K_1), \ldots, C(K_m)$ their cones. We denote by $*_i$ the vertex of the cone $C(K_i)$. Let $C(K_1) \cup C(K_2) \cup \cdots \cup C(K_m)$ be the disjoint union of $C(K_1), \ldots, C(K_m)$. We define

$$C(K_1) \cup_* C(K_2) \cup_* \cdots \cup_* C(K_m) = C(K_1) \cup C(K_2) \cup \cdots \cup C(K_m) / \sim$$

where $*_i \sim *_j$ for all i, j. Namely we glue $C(K_1), \ldots, C(K_m)$ at their vertices, and the resulting space is denoted by $C(K_1) \cup_* C(K_2) \cup_* \cdots \cup_* C(K_m)$. The shared vertex (i.e. the equivalence class of $*_i$) is denoted by *. See Figure 3.

When K_1, \ldots, K_m are simplicial complexes then $C(K_1) \cup_* \cdots \cup_* C(K_m)$ is also a simplicial complex and its dimension is the maximum of dim $K_i + 1$ $(1 \le i \le m)$.




FIGURE 3. The left is the cone C(K) with the vertex *. The right is $C(K_1) \cup_* C(K_2) \cup_* C(K_3)$ with the vertex *. Three cones are glued at their vertices.

Proposition 4.6. Let $\pi : (X,T) \to (Y,S)$ be a factor map between dynamical systems. If (Y,S) has the small boundary property then

$$\operatorname{mdim}(X, T) = \operatorname{mdim}(\pi, T).$$

Proof. $\operatorname{mdim}(\pi, T) \leq \operatorname{mdim}(X, T)$ is obvious from the definition. Let *a* be any positive number with $\operatorname{mdim}(\pi, T) < a$. We will prove $\operatorname{mdim}(X, T) \leq a$.

We take a metric d on X. Let ε be any positive number. We can take a natural number N such that

$$\sup_{y \in Y} \operatorname{Widim}_{\varepsilon} \left(\pi^{-1}(y), d_N \right) + 1 < aN.$$

As in Claim 2.3 in §2.2, for each $y \in Y$ there exists an open neighborhood U_y of $\pi^{-1}(y)$ satisfying

Widim_{$$\varepsilon$$} $(U_y, d_N) + 1 < aN.$

There exists an open neighborhood V_y of y with $\pi^{-1}(V_y) \subset U_y$.

Since Y is compact, we can find an open covering $Y = V_1 \cup V_2 \cup \cdots \cup V_m$ such that for all $1 \le i \le m$

Widim_{$$\varepsilon$$} $\left(\pi^{-1}(V_i), d_N\right) + 1 < aN.$

Let δ be any positive number. By applying Lemma 4.5 to an open cover $Y = V_1 \cup \cdots \cup V_m$, we find compact subsets $E_i \subset V_i$ $(1 \le i \le m)$ such that

$$E_i \cap E_j = \emptyset \quad (i \neq j), \quad \operatorname{ocap}\left(Y \setminus (E_1 \cup \dots \cup E_m)\right) < \delta.$$

We take open sets W_i $(1 \le i \le m)$ satisfying

$$E_i \subset W_i \subset V_i, \quad W_i \cap W_j = \emptyset \quad (i \neq j).$$

We also take a continuous function $\rho: Y \to [0,1]$ such that $\rho = 1$ on $E_1 \cup E_2 \cup \cdots \cup E_m$ and $\operatorname{supp} \rho \subset W_1 \cup W_2 \cup \cdots \cup W_m$. See Figure 4.




FIGURE 4. The continuous function ρ .

For each $1 \leq i \leq m$ we take an ε -embedding $f_i : (\pi^{-1}(V_i), d_N) \to K_i$ such that K_i is a simplicial complex of dimension smaller than aN - 1. Let $C(K_i)$ be the cone over K_i . We set

$$K' := C(K_1) \cup_* C(K_2) \cup_* \cdots \cup_* C(K_m).$$

This is a simplicial complex of dimension smaller than aN. We define a continuous map $f': X \to K'$ as follows: If $x \in \pi^{-1}(W_i)$ $(1 \le i \le m)$ then we set

$$f'(x) = \rho(\pi(x)) f_i(x) \in C(K_i) \subset K'.$$

If $x \notin \pi^{-1}(W_1) \cup \pi^{-1}(W_2) \cup \cdots \cup \pi^{-1}(W_m)$ then we set f'(x) = *.

We take an ε -embedding $g : (X, d_N) \to L$ such that L is a simplicial complex of dimension $\operatorname{Widim}_{\varepsilon}(X, d_N) < \infty$. Let L' := C(L) be the cone over L. We have $\dim L' = \operatorname{Widim}_{\varepsilon}(X, d_N) + 1$. We define a continuous map $g' : X \to L'$ by

$$g'(x) = (1 - \rho(\pi(x))) g(x).$$

For $n \ge 1$ we define $F_n : X \to (K' \times L')^n$ by

$$F_n(x) = \left(f'(x), g'(x), f'(T^N x), g'(T^N x), \dots, f'(T^{(n-1)N} x), g'(T^{(n-1)N} x)\right)$$

Claim 4.7. F_n is an ε -embedding with respect to the metric d_{nN} .

Proof. It is enough to prove that

$$f' \times g' : X \to K' \times L', \quad x \mapsto (f'(x), g'(x))$$

is an ε -embedding with respect to the metric d_N .

Suppose (f'(x), g'(x)) = (f'(x'), g'(x')) for $x, x' \in X$. Then we have $\rho(\pi(x)) = \rho(\pi(x'))$. If this common value is zero then g(x) = g(x') and hence we have $d_N(x, x') < \varepsilon$. If the value is positive then there exists $1 \le i \le m$ such that $x, x' \in \pi^{-1}(W_i)$ and $f_i(x) = f_i(x')$. Then $d_N(x, x') < \varepsilon$. **Claim 4.8.** If n is sufficiently large then for all $x \in X$

$$\left| \{ 0 \le k < n \mid g'(T^{kN}x) \neq * \} \right| < \delta n N.$$

Therefore the image of F_n is contained in a simplicial complex of dimension smaller than $n \dim K' + \delta n N \dim L'$.

Proof. Let $x \in X$ and set $y = \pi(x)$. The condition $g'(T^{kN}x) \neq *$ is equivalent to $\rho(S^{kN}y) < 1$. The latter condition implies $S^{kN}y \in Y \setminus (E_1 \cup \cdots \cup E_m)$. Then

$$\left| \{ 0 \le k < n \mid g'(T^{kN}x) \neq * \} \right| \le \sum_{k=0}^{n-1} \mathbb{1}_{Y \setminus (E_1 \cup \dots \cup E_m)}(S^{kN}y)$$
$$\le \sum_{k=0}^{nN-1} \mathbb{1}_{Y \setminus (E_1 \cup \dots \cup E_m)}(S^ky)$$
$$< \delta nN \quad \text{by ocap}\left(Y \setminus (E_1 \cup \dots \cup E_m)\right) < \delta$$

for all sufficiently large n (uniformly in $x \in X$).

By Claims 4.7 and 4.8, for sufficiently large n

Widim_{$$\varepsilon$$} $(X, d_{nN}) < n \dim K' + \delta nN \dim L'$
 $< anN + \delta nN (Widim_{\varepsilon}(X, d_N) + 1).$

Then

$$\lim_{n \to \infty} \frac{\operatorname{Widim}_{\varepsilon} (X, d_{nN})}{nN} \le a + \delta \left(\operatorname{Widim}_{\varepsilon} (X, d_{N}) + 1 \right).$$

Here δ is independent of ε, N . So we can let $\delta \to 0$ and get

$$\lim_{n \to \infty} \frac{\operatorname{Widim}_{\varepsilon}(X, d_n)}{n} \le a.$$

Letting $\varepsilon \to 0$, we conclude

$$\operatorname{mdim}\left(X,T\right) \leq a.$$

Now we are ready to prove Theorem 1.5. We write the statement again.

Theorem 4.9 (= Theorem 1.5). Let $\pi : (X,T) \to (Y,S)$ be a factor map between dynamical systems. If mdim(Y,S) = 0 then

$$\operatorname{mdim}(X,T) = \operatorname{mdim}(\pi,T).$$

Proof. Let (Z, R) be a zero dimensional free minimal dynamical system. We consider

(4.2)
$$\pi \times \mathrm{Id} : (X \times Z, T \times R) \to (Y \times Z, S \times R)$$

The system $(Y \times Z, S \times R)$ has a free minimal factor (Z, R) and its mean dimension is zero:

$$\operatorname{mdim}(Y \times Z, S \times R) = \operatorname{mdim}(Y, S) = 0$$
 since $\operatorname{dim} Z = 0$.

By the Lindenstrauss Theorem (Theorem 4.4), the system $(Y \times Z, S \times R)$ has the small boundary property.

Now we can apply Proposition 4.6 to the above factor map $(4\cdot 2)$ and get

$$\operatorname{mdim}(X \times Z, T \times R) = \operatorname{mdim}(\pi \times \operatorname{Id}, T \times R).$$

We have

$$\operatorname{mdim}(X \times Z, T \times R) = \operatorname{mdim}(X, T) \quad \text{since } \dim Z = 0,$$

$$\operatorname{mdim}(\pi \times \operatorname{Id}, T \times R) = \operatorname{mdim}(\pi, T)$$
 by Corollary 4.2.

Therefore we conclude $\operatorname{mdim}(X, T) = \operatorname{mdim}(\pi, T)$.

References

- [Gro88] M. Gromov, Width and related invariants of Riemannian manifolds, Astérisque 163-164 (1988) 93-109.
- [Gro99] M. Gromov, Topological invariants of dynamical systems and spaces of holomorphic maps: I, Math. Phys. Anal. Geom. 2 (1999) 323-415.
- [GT20] Y. Gutman, M. Tsukamoto, Embedding minimal dynamical systems into Hilbert cubes, Invent. math. 221 (2020) 113-166.
- [HW41] W. Hurewicz, H. Wallman, Dimension theory, Princeton University Press, 1941.
- [LL18] H. Li, B. Liang, Mean dimension, mean rank, and von Neumann-Lück rank, J. Reine Angew. Math. 739 (2018) 207–240.
- [Lia21] B. Liang, Conditional mean dimension, Ergodic Theory Dynam. Systems, First Views, DOI: https://doi.org/10.1017/etds.2021.85, arXiv:2001.02603.
- [Lin99] E. Lindenstrauss, Mean dimension, small entropy factors and an embedding theorem, Inst. Hautes Études Sci. Publ. Math. 89 (1999) 227-262.
- [LW00] E. Lindenstrauss, B. Weiss, Mean topological dimension, Israel J. Math. 115 (2000) 1-24.
- [MT19] T. Meyerovitch, M. Tsukamoto, Expansive multiparameter actions and mean dimension, Trans. Amer. Math. Soc. 371 (2019), 7275-7299.
- [Tsu08] M. Tsukamoto, Moduli space of Brody curves, energy and mean dimension, Nagoya Math. J. 192 (2008) 27-58.

Department of Mathematics, Kyoto University, Kitashirakawa Oiwake-cho, Sakyo-ku, Kyoto 606-8502, Japan

Email address: tsukamoto@math.kyoto-u.ac.jp