

Comment on the Analytical Bounds in the Rezzolla-Zhidenko Parametrization

Prashant Kocherlakota¹ and Luciano Rezzolla^{1,2,3}

¹*Institut für Theoretische Physik, Goethe-Universität, Max-von-Laue-Str. 1, 60438 Frankfurt, Germany*

²*Frankfurt Institute for Advanced Studies, Ruth-Moufang-Str. 1, 60438 Frankfurt, Germany and*

³*School of Mathematics, Trinity College, Dublin 2, Ireland*

In this short note, we briefly comment on the analytical bounds that must be imposed on the parameter space of the Rezzolla-Zhidenko (RZ) metric-parametrization approach introduced in Ref. [1]. We hope this will clarify some of the confusion recently emerged on this issue [2].

Introduction. The Rezzolla-Zhidenko (RZ) metric parametrization approach was introduced in Ref. [1] to characterize efficiently and accurately the exterior horizon geometry of asymptotically-flat, spherically-symmetric and static black-hole (BH) spacetimes. This approach has been employed successfully to model a large number of generic BH spacetimes¹ in terms of a series of coefficients $\{\epsilon, a_i, b_i\}$ and by Padé approximants in the form of continued fractions (see Ref. [3] and references therein); the approach has also been extended to stationary BH spacetimes in Ref. [4].

Comment. In the RZ-parametrization approach, the radial coordinate $r = r_0$ must, *by construction*, correspond to the location of the outermost Killing horizon. We recall that spherically symmetric BH spacetimes admit a time-translation Killing vector field \mathbf{K} which, in the static ($\mathbf{K} \wedge d\mathbf{K} = 0$) exterior horizon geometry, can be identified to be $\mathbf{K} = \partial_t$. \mathbf{K} is asymptotically timelike, i.e., $g_{tt} < 0$, and becomes null on Killing horizons located at $r = r_{\text{kh};k}$, i.e., $g_{tt}(r_{\text{kh};k}) = 0$ (with k indexing the roots of g_{tt}). As per the definition above, if r_0 is the largest, real, positive root, i.e., $r_0 = \max\{r_{\text{kh};k}, \forall k\} \in \mathbb{R}^+$, then clearly for \mathbf{K} to remain timelike on the exterior horizon geometry, $N^2(r) := -g_{tt}$ must be positive-definite on $r > r_0$. Therefore, and as already noted in Ref. [1, 3, 5, 6], this requires that the following condition must be met

$$N^2 =: xA(x, \epsilon, a_i) > 0, \quad \text{for } 0 \leq x \leq 1, \quad (1)$$

where $x := 1 - r_0/r$ is a compactified coordinate such that $x = 0$ corresponds to the location of the event horizon ($N^2(x) = 0$) and $x = 1$ to spatial infinity (we recall that $r_0 := 2M/(1 + \epsilon)$, where M is the BH mass). As pointed out already in 2014 (see Eq. (5) of Ref. [1]), Eq. (1) partitions the relevant parameter space into mathematically “allowed” and “disallowed” regions, and its impact on the relevant parameter spaces for various slices of the general RZ metric has *already* been reported in Table 2 of Ref. [5], but also in Figs. 3-5 of Ref. [5], where the regions in which $r = r_0$ does not locate the outermost Killing horizon are shown as gray regions.

Recently, Ref. [2] has considered a two-parameter realization of the RZ approach, that is, it has employed the space of parameters (r_0, a_1) , or equivalently (ϵ, a_1) , to express the g_{tt} metric function of a generic BH spacetime and solved for all possible roots of $N^2(r, r_0, a_1) = 0$, which corresponds to a quartic equation in this specific case. Finding that $r = r_0$ does not locate the outermost Killing horizon for arbitrary choices of the RZ parameters, Ref. [2] partitions the relevant (r_0, a_1) RZ parameter space into regions where $r = r_0$ does in fact locate the outermost Killing horizon and where it does not, essentially re-iterating the computation reported in Ref. [5]. Indeed, Eqs. (25) and (26) of Ref. [2] are not new and have already been reported respectively in the seventh and eight rows of Table 2 of Ref. [5] (i.e., the case $\mathcal{M}(\epsilon, a_1)$); the same table also provides the ranges for other choices of the parametrization.

Conclusion. Contrary to the recent claim in Ref. [2], the horizon radius in the RZ parameterization does not change discontinuously. This can happen only if the parameterization is performed incorrectly outside of the region of validity required by Eq. (1). Such conditions were *already* discussed and published in Refs. [1, 3, 5, 6]. Finally, we note that to ensure that the Ricci and Kretschmann scalars corresponding to the RZ metric do not diverge outside or on the horizon, the metric function $B^2(r) := g_{rr}(r)N^2(r)$ must generically be nonvanishing for $r \geq r_0$. This represents another nontrivial constraint on the b_i coefficients of the RZ parameter space.

We thank Alexander Zhidenko for useful comments. Support comes from the ERC Advanced Grant “JETSET: Launching, propagation and emission of relativistic jets from binary mergers and across mass scales” (Grant No. 884631).

¹ As noted in Ref. [3], the RZ parametrization can trivially be employed also for spherically symmetric static spacetimes that do not contain BHs but other compact objects, including nonsingular and singular black holes, boson stars, and naked singularities.

-
- [1] L. Rezzolla and A. Zhidenko, *Phys. Rev. D* **90**, 084009 (2014), arXiv:1407.3086 [gr-qc].
 - [2] D. Heumann and D. Psaltis, arXiv e-prints , arXiv:2205.12994 (2022), arXiv:2205.12994 [gr-qc].
 - [3] P. Kocherlakota and L. Rezzolla, *Phys. Rev. D* **102**, 064058 (2020), arXiv:2007.15593 [gr-qc].
 - [4] R. Konoplya, L. Rezzolla, and A. Zhidenko, *Phys. Rev. D* **93**, 064015 (2016), arXiv:1602.02378 [gr-qc].
 - [5] P. Kocherlakota and L. Rezzolla, *MNRAS* **513**, 1229 (2022).
 - [6] Event Horizon Telescope Collaboration *et al.*, *ApJL* **930**, L17 (2022).