

Decentralized Online Regularized Learning Over Random Time-Varying Graphs

Xiwei Zhang, Tao Li, *Senior Member, IEEE*, and Xiaozheng Fu

Abstract

We study the decentralized online regularized linear regression algorithm over random time-varying graphs. At each time step, every node runs an online estimation algorithm consisting of an innovation term processing its own new measurement, a consensus term taking a weighted sum of estimations of its own and its neighbors with additive and multiplicative communication noises and a regularization term preventing over-fitting. It is not required that the regression matrices and graphs satisfy special statistical assumptions such as mutual independence, spatio-temporal independence or stationarity. We develop the nonnegative supermartingale inequality of the estimation error, and prove that the estimations of all nodes converge to the unknown true parameter vector almost surely if the algorithm gains, graphs and regression matrices jointly satisfy the *sample path spatio-temporal persistence of excitation* condition. Especially, this condition holds by choosing appropriate algorithm gains if the graphs are *uniformly conditionally jointly connected* and *conditionally balanced*, and the regression models of all nodes are *uniformly conditionally spatio-temporally jointly observable*, under which the algorithm converges in mean square and almost surely. In addition, we prove that the regret upper bound is $\mathcal{O}(T^{1-\tau}\ln T)$, where $\tau \in (0.5, 1)$ is a constant depending on the algorithm gains.

Index Terms

Decentralized online linear regression, regularization, random time-varying graph, persistence of excitation, regret analysis.

This work was supported by the National Natural Science Foundation of China under Grant 62261136550. (*Corresponding author: Tao Li.*)

Xiwei Zhang is with the School of Mathematical Sciences, East China Normal University, Shanghai 200241, China (e-mail: xwzhangmath@sina.com).

Tao Li is with the School of Mathematical Sciences, East China Normal University, Shanghai 200241, China (e-mail: tli@math.ecnu.edu.cn).

Xiaozheng Fu is with the School of Mathematical Sciences, East China Normal University, Shanghai 200241, China (e-mail: fxz4926@163.com).

I. INTRODUCTION

Empirical risk minimization is an important criterion to judge the predictive ability of models in statistical learning. Problems in machine learning are usually ill-posed, and the solutions obtained by using the empirical risk minimization principle are unstable, easily overfitted and usually have large norms. To solve this problem, Poggio, Girosi *et al.* [1]-[3] introduced regularization methods in inverse problems. Regularization is an effective tool for dealing with the complexity of a model, and its role is to choose a model with less empirical risk and complexity at the same time. In addition, this method can transform the original ill-posed problem into a well-posed one, reduce the norm of estimations of the unknown true parameter vector, and keep the sum of the squared errors small [4]. Regularization methods are widely used in various research areas, such as classification in machine learning, online state estimation for power systems and image reconstruction [5]-[9]. There are two classic regularization methods: Ivanov regularization [10], which restricts the hypothesis space, and Tikhonov regularization, which restricts certain parameters in the loss function, i.e., adds regularization terms which are also called penalty terms into the algorithm. In learning theory, ridge regression adopts Tikhonov regularization [11].

An offline algorithm usually needs to acquire a finite amount of data generated by an unknown, stationary probability distribution in advance, which heavily relies on the memory of the system especially when the dataset is far too large, while an online algorithm processes infinite data streams that are continuously generated at rapid rates, where the data is discarded after it has been processed, and the unknown data generation process is possibly non-stationary. So far, the centralized online regularized algorithms have been widely studied in [12]-[22]. In a centralized algorithm, there is an information fusion center collecting the measurements of all nodes and giving the global estimation.

In reality, many learning tasks process very large datasets, and thus decentralized parallel processing of data by communicating and computing units in the network is necessary, see e.g. [23]-[24] and references therein. Besides, if the data contains sensitive private information (e.g. medical and social network data, etc.), they may come from different units in the network, and transmitting all these data subsets to the fusion center may lead to potential privacy risks [25]-[26]. Therefore, decentralized learning is needed, which can improve the efficiency of communication and protect the privacies of users.

At present, the non-regularized decentralized linear regression problems have been widely

studied in [27]-[38]. Xie and Guo [32]-[33] considered the time-varying linear regression with measurement noises, where the cooperative information condition on the conditional expectations of the regression matrices was proposed over a deterministic, undirected and strongly connected graph. Chen *et al.* [35] proposed a saturated innovation update algorithm for the decentralized estimation under sensor attacks, where the interagent communication is noiseless. They proved that if the communication graph is undirected and fixed, the nodes are locally observable, and the number of attacked nodes is less than half of the total, then all nodes' estimations converge to the unknown true parameter with a polynomial rate. Yuan *et al.* [36] investigated the decentralized online linear regression problem, where the graph is assumed to be strongly connected and balanced. They gave the regret upper bounds. Wang *et al.* [38] investigated a consensus plus innovation based decentralized linear regression algorithm over random networks with random regression matrices. They proved that if the regression matrices and communication graphs satisfy the stochastic spatio-temporal persistence of excitation condition, properly choosing the algorithm gains guarantees the convergence of the estimations of all nodes to the unknown true parameter. Some scholars have also considered both measurement and communication noises among nodes, e.g. [28]-[29] and [31]. Kar *et al.* [28]-[29] studied the decentralized parameter estimation algorithms based on consensus plus innovation with measurement and communication noises. It is required that the expectations of the regression matrices be known, and the graphs, regression matrices, measurement and communication noises be spatially i.i.d. Moreover, Kar *et al.* [28] and Sahu *et al.* [31] investigated the decentralized estimation algorithms with nonlinear observation models.

Based on the advantages of the decentralized information structure, the online algorithm and the regularization method, we propose a decentralized online regularized algorithm for the linear regression problem over random time-varying graphs. The algorithm of each node contains an innovation term, a consensus term and a regularization term. In each iteration, the innovation term is used to update the node's own estimation, the consensus term is the weighted sum of estimations of its own and its neighbors with additive and multiplicative communication noises, and the regularization term is helpful for constraining the norm of the estimation in the algorithm and preventing the unknown true parameter vector from being overfitted. Although regularization is an effective method to deal with linear regression problems, it brings essential difficulties to the convergence analysis of the algorithm. Compared with the non-regularized decentralized linear regression algorithm, the estimation error equation of this algorithm contains a non-martingale

difference term with the regularization parameter, which cannot be directly analyzed by using martingale convergence theorem as [39]. We no longer require that the sequences of regression matrices and graphs satisfy special statistical properties, such as mutual independence, spatio-temporal independence and stationarity. Compared with the case with i.i.d. data, dependent observations and data contain less information and therefore lead to more unstable learning errors as well as the performance degradation [40]. Besides, we consider both additive and multiplicative communication noises in the process of the information exchange among nodes. All these challenges make it difficult to analyze the convergence and performance of the algorithm, and the methods in the existing literature are no longer applicable. For example, the methods in [27]-[30] and [34] are applicable for the case that the graphs, regression matrices and noises are i.i.d. and mutually independent and it is required that the expectations of the regression matrices be known in [28]-[29]. Liu *et al.* [25] studied the decentralized regularized gossip gradient descent algorithm for linear regression models, where the method is applicable for the case that only two nodes exchange information at each instant. In addition, they require that the graphs be strongly connected and the observation vectors and the noises be i.i.d. and bounded. Wang *et al.* [38] studied the non-regularized decentralized online algorithm, where the communication noises were not considered, and the method therein is only applicable for the case that the inhomogeneous part of the estimation error equation is a martingale difference sequence.

To overcome the difficulties mentioned above, we develop the nonnegative supermartingale inequality of the estimation error, and further establish the *sample path spatio-temporal persistence of excitation* condition by combining information of the regression matrices, graphs and algorithm gains, under which sufficient conditions for the convergence of the algorithm are obtained. We prove that if the algorithm gains and the sequences of regression matrices and Laplacian matrices of the graphs jointly satisfy the *sample path spatio-temporal persistence of excitation* condition, then the estimations of all nodes converge to the unknown true parameter vector almost surely. Furthermore, we give an intuitive sufficient condition of the *sample path spatio-temporal persistence of excitation* condition, i.e., the graphs are *conditionally balanced* and *uniformly conditionally jointly connected*, and the regression models of all nodes are *uniformly conditionally spatio-temporally jointly observable*, which drives the algorithm to converge in mean square and almost surely by the proper choice of the algorithm gains. Especially, for the case with Markovian switching graphs and regression matrices, we prove that the *sample path spatio-temporal persistence of excitation* condition holds if the stationary graph is balanced

with a spanning tree and regression model is *spatio-temporally jointly observable*, implying that neither *local observability* of each node nor *instantaneous global observability* of all the regression models is necessary.

Historically, Guo [41] first proposed the stochastic persistence of excitation condition for analyzing the centralized Kalman filtering algorithm, which was then refined in [42]. Whereafter, the cooperative information condition on the conditional expectations of the regression matrices over the deterministic connected graph for the decentralized adaptive filtering algorithms was proposed in [32]-[33]. The stochastic spatio-temporal persistence of excitation condition on the conditional expectations of the regression matrices and graphs was proposed for the non-regularized decentralized online estimation algorithms over the random time-varying graphs in [37]-[38]. Specifically, the stochastic spatio-temporal persistence of excitation condition in [37]-[38] requires that the minimum eigenvalues of the random information matrices have a positive lower bound independent of the sample paths. In this paper, this excitation condition is further weakened to the *sample path spatio-temporal persistence of excitation* condition, which only requires that the infinite series composed of the minimum eigenvalues of the information matrices diverge for almost all sample paths. Meanwhile, we extend the work of Li and Wang [39] from distributed averaging to decentralized online learning and generalize the network model of Xie and Guo [32]-[33] from deterministic, time-invariant and undirected graphs to random and time-varying digraphs. Xie and Guo [32]-[33] considered tracking time-varying parameters by using constant algorithm gains and obtained the L_p -boundedness of the tracking errors. In [32]-[33], the homogeneous part of the estimation error equation is L_p -exponentially stable. Different from [32]-[33], we consider estimating time-invariant parameters. To ensure the strong consistency of the algorithm, we use decaying algorithm gains, which result in that the homogeneous part of the estimation error equation is not L_p -exponentially stable. To ensure not only boundedness but also strong consistency of the algorithm, the decaying rates of the algorithm gains and regularization parameter need to be designed precisely.

We use the regret to evaluate the performance of the decentralized optimization algorithm, which has been investigated in [36] and [43]-[44]. Yuan *et al.* [36] studied the non-regularized decentralized online linear regression problem over the fixed graph. Bedi *et al.* [43] considered the multi-agent stochastic optimization problems with constraints in heterogeneous networks. Dixit *et al.* [44] studied the decentralized online dynamic optimization problems over deterministic and time-varying graphs. Compared with [36] and [43]-[44], which assumed the graphs to be

deterministic, strongly connected and balanced, we consider the decentralized online regularized linear regression algorithm over the random time-varying graphs, and our techniques are helpful to further study the optimization problems over random time-varying graphs. We prove that the upper bound of the regret is $\mathcal{O}(T^{1-\tau}\ln T)$, where $\tau \in (0.5, 1)$ is a constant depending on the decaying algorithm gains.

The rest of this paper is organized as follows. Section II proposes the decentralized online regularized linear regression algorithm. Section III gives the main results, including the convergence analysis of the algorithm and the analysis of the regret upper bound. Section IV gives the numerical examples. Section V concludes the full paper and gives some future topics.

Notation and symbols: \mathbb{R}^n : the n dimensional real vector space; $\mathbb{R}^{m \times n}$: the $m \times n$ dimensional real matrix space; \otimes : the Kronecker product; $\mathbf{diag}(A_1, \dots, A_n)$: the block diagonal matrix with entries being A_1, \dots, A_n ; $\|A\|$: the 2-norm of matrix A ; $\|A\|_F$: the Frobenius-norm of matrix A ; $\mathbb{P}\{A\}$: the probability of event A ; $\mathbb{E}[\xi]$: the mathematical expectation of random variable ξ ; A^T : the transpose of matrix A ; $\rho(A)$: the spectral radius of matrix A ; $\lambda_{\min}(A)$: the minimum eigenvalue of real symmetric matrix A ; $\lambda_2(A)$: the second smallest eigenvalue of real symmetric matrix A ; $A \geq B$: the matrix $A - B$ is positive semi-definite; $A \succeq B$: the matrix $A - B$ is nonnegative; I_n : the n dimensional identity matrix; $\mathbf{O}_{n \times m}$: the $n \times m$ dimensional zero matrix; $\mathbf{1}_N$: the N dimensional vector with all ones; $\lfloor x \rfloor$: the largest integer less than or equal to x ; $\lceil x \rceil$: the smallest integer greater than or equal to x ; $a_n = \mathcal{O}(b_n)$: $\lim_{n \rightarrow \infty} \sup \frac{|a_n|}{b_n} < \infty$, where $\{a_n, n \geq 0\}$ is a real sequence and $\{b_n, n \geq 0\}$ is a positive real sequence; $a_n = o(b_n)$: $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 0$; For a sequence of $n \times n$ dimensional matrices $\{Z(k), k \geq 0\}$, denote

$$\Phi_Z(j, i) = \begin{cases} Z(j) \cdots Z(i), & j \geq i \\ I_n, & j < i. \end{cases}, \quad \prod_{k=i}^j Z(k) = \Phi_Z(j, i).$$

II. DECENTRALIZED REGULARIZED LINEAR REGRESSION

Suppose that $x_0 \in \mathbb{R}^n$ is the unknown true parameter vector. We consider a network modeled by a sequence of random digraphs with N nodes $\{\mathcal{G}(k) = \{\mathcal{V}, \mathcal{E}_{\mathcal{G}(k)}, \mathcal{A}_{\mathcal{G}(k)}\}, k \geq 0\}$, where \mathcal{V} is the node set, $\mathcal{E}_{\mathcal{G}(k)}$ is the edge set at instant k , and $\mathcal{A}_{\mathcal{G}(k)}$ is the weighted adjacency matrix. The regression model of node i at instant k is given by

$$y_i(k) = H_i(k)x_0 + v_i(k), \quad k \geq 0, \quad i \in \mathcal{V}, \quad (1)$$

where $H_i(k) \in \mathbb{R}^{n_i \times n}$ is the random regression matrix of node i at instant k , $v_i(k) \in \mathbb{R}^{n_i}$ is the additive measurement noise, and $y_i(k) \in \mathbb{R}^{n_i}$ is the observation data.

Remark 1. The regression model (1) with random regression matrices over random graphs have been widely studied in [25], [27]-[31], [34], [37]-[38] and [45]-[46]. There are various uncertainties in real-world networks, where intermittent sensing failures and packet dropouts may occur at random times [29]. (i) Node/link failures can be modeled by a sequence of random communication graphs [45]. (ii) Node sensing failures or measurement losses can be modeled by a sequence of random observation/regression matrices, for example, $H_i(k) = \frac{1}{p}\mu_i(k)C_i$, where $\{\mu_i(k), k \geq 0\}$ is a sequence of zero-one i.i.d. Bernoulli variables account for sensor failures, $p > 0$ is the sensing probability, and C_i models the normal operation of the sensor [28]-[29]. Besides, in decentralized parameter identification [38], all the nodes over graphs cooperatively identify the parameters of an auto-regressive (AR) model from noisy data, and each node's measurement equation is given by

$$y_i(k) = \sum_{j=1}^d c_j y_i(k-j) + v_i(k), \quad k \geq 0, \quad i \in \mathcal{V}, \quad (2)$$

where $\{c_j \in \mathbb{R}, 1 \leq j \leq d\}$ are the model parameters to be identified, and $v_i(k) \in \mathbb{R}$ is the additive zero-mean white noise. Here, the unknown parameter $x_0 = [c_1, \dots, c_d]^T \in \mathbb{R}^d$ and the random regression matrix $H_i(k) = [y_i(k-1), \dots, y_i(k-d)] \in \mathbb{R}^{1 \times d}$.

Remark 2. In this paper, the network structure is modeled by a sequence of random digraphs $\mathcal{G}(k, \omega) = \{\mathcal{V}, \mathcal{E}_{\mathcal{G}(k, \omega)}, \mathcal{A}_{\mathcal{G}(k, \omega)}\}$, $k \geq 0$, where ω is the sample path, $\mathcal{A}_{\mathcal{G}(k, \omega)} = [w_{ij}(k, \omega)]_{N \times N}$ is an N -dimensional random weighted adjacency matrix with zero diagonal elements, and $\mathcal{E}_{\mathcal{G}(k, \omega)}$ is the edge set, where each edge represents a communication link. Denote the neighborhood of node i at instant k by $\mathcal{N}_i(k, \omega) = \{j \in \mathcal{V} | (j, i) \in \mathcal{E}_{\mathcal{G}(k, \omega)}\}$. There is a one-to-one correspondence between $\mathcal{G}(k, \omega)$ and $\mathcal{A}_{\mathcal{G}(k, \omega)}$. Here, the sample path ω is omitted. The in-degree of node i at instant k $\deg_{\text{in}}^i(k) = \sum_{j \in \mathcal{V}} w_{ij}(k)$ and the out-degree of node i at instant k $\deg_{\text{out}}^i(k) = \sum_{j \in \mathcal{V}} w_{ji}(k)$. If $\deg_{\text{in}}^i(k) = \deg_{\text{out}}^i(k)$, $\forall i \in \mathcal{V}$, then $\mathcal{G}(k)$ is balanced. We call $\mathcal{L}_{\mathcal{G}(k)} = \mathcal{D}_{\mathcal{G}(k)} - \mathcal{A}_{\mathcal{G}(k)}$ the Laplacian matrix of $\mathcal{G}(k)$, where $\mathcal{D}_{\mathcal{G}(k)} = \mathbf{diag}(\deg_{\text{in}}^1(k), \dots, \deg_{\text{in}}^N(k))$.

We now provide the following real-world example of decentralized linear regression with random regression matrices over random graphs.

Example 1. In the decentralized multi-area state estimation in power systems [38], the power grid is partitioned into multiple geographically non-overlapping areas, where the communication topology is modeled by random graphs. The state of the grid x_0 to be estimated consists of the amplitude of the voltages and the phase angles of all the buses. The measurement $y_i(k)$ in each region consists of the active power flow and the reactive power flow. After a DC power flow approximation [47], the grid state to be estimated degenerates into a vector of phase angles of all buses. The relationship between the measurement in the i -th region and the grid state is represented by

$$y_i(k) = s_i(k)\overline{H}_i(k)x_0 + v_i(k), \quad k \geq 0,$$

where $\{v_i(k), k \geq 0\}$ is the sensing noise sequence, $\{s_i(k) \in \mathbb{R}, k \geq 0\}$ is a sequence of i.i.d. Bernoulli variables, which represents the intermittent sensor failures, and $\{\overline{H}_i(k), k \geq 0\}$ is the sequence of observation matrices without sensing failures.

Denote $y(k) = [y_1^T(k), \dots, y_N^T(k)]^T$, $H(k) = [H_1^T(k), \dots, H_N^T(k)]^T$, $\mathcal{H}(k) = \mathbf{diag}(H_1(k), \dots, H_N(k))$, and $v(k) = [v_1^T(k), \dots, v_N^T(k)]^T$. Rewrite (1) by the compact form

$$y(k) = H(k)x_0 + v(k), \quad k \geq 0. \quad (3)$$

The estimation of $\mathbf{1}_N \otimes x_0$ can be obtained by minimizing the loss function $\Psi(\cdot)$ as

$$\hat{x}_0 = \arg \min_{x \in \mathbb{R}^{Nn}} \Psi(x) \triangleq \frac{1}{2} (\mathbb{E} [\|y(k) - \mathcal{H}(k)x\|^2] + \mathbb{E} [\langle (\mathcal{L}_{\mathcal{G}(k)} \otimes I_n) x, x \rangle] + \lambda \|x\|^2), \quad \lambda > 0,$$

if the sequences of the graphs, regression matrices and noises are identically distributed, respectively, the mean graph is undirected, and the sequences of regression matrices and noises are mutually independent. The loss function $\Psi(\cdot)$ consists of three parts: the mean square error of the estimation $\mathbb{E}[\|y(k) - \mathcal{H}(k)x\|^2]$, the consensus error $\mathbb{E}[\langle (\mathcal{L}_{\mathcal{G}(k)} \otimes I_n) x, x \rangle]$ and the regularization term $\lambda \|x\|^2$. To solve the above optimization problem, we consider the stochastic gradient descent (SGD) algorithm:

$$x(k+1) = x(k) + a(k)\mathcal{H}^T(k)(y(k) - \mathcal{H}(k)x(k)) - b(k)(\mathcal{L}_{\mathcal{G}(k)} \otimes I_n)x(k) - \lambda(k)x(k), \quad (4)$$

where $\lambda(k) \triangleq \lambda c(k)$. Let $x(k) = [x_1^T(k), \dots, x_N^T(k)]^T$, then $x_i(k) \in \mathbb{R}^n$ is the estimation of the node i at instant k . From (4), it follows that

$$x_i(k+1) = x_i(k) + a(k)H_i^T(k)(y_i(k) - H_i(k)x_i(k)) + b(k) \sum_{j \in \mathcal{N}_i(k)} w_{ij}(k)(x_j(k) - x_i(k)) - \lambda(k)x_i(k), \quad k \geq 0, \quad i \in \mathcal{V},$$

where $a(k)H_i^T(k)(y_i(k) - H_i(k)x_i(k))$ is the innovation term updating the estimation $x_i(k)$ with the innovation gain $a(k)$, $b(k)\sum_{j \in \mathcal{N}_i(k)} w_{ij}(k)(x_j(k) - x_i(k))$ is the consensus term taking a weighted sum of estimations of its own and its neighbors with the consensus gain $b(k)$, and $\lambda(k)x_i(k)$ is the regularization term constraining the estimation $x_i(k)$ with the regularization gain $\lambda(k)$, which avoids overfitting the estimation of the unknown true parameter vector x_0 . In the practical algorithm, there are communication noises among nodes. Specifically, node j acquires the estimations of its neighbors by

$$\mu_{ji}(k) = x_j(k) + f_{ji}(x_j(k) - x_i(k))\xi_{ji}(k), \quad j \in \mathcal{N}_i(k), \quad (5)$$

where $\xi_{ji}(k)$ is the communication noise, and $f_{ji}(x_j(k) - x_i(k))$ is the noise intensity function.

Therefore, the decentralized online regularization algorithm is given by

$$\begin{aligned} x_i(k+1) = & x_i(k) + a(k)H_i^T(k)(y_i(k) - H_i(k)x_i(k)) + b(k) \sum_{j \in \mathcal{N}_i(k)} w_{ij}(k)(\mu_{ji}(k) - x_i(k)) \\ & - \lambda(k)x_i(k), \quad k \geq 0, \quad i \in \mathcal{V}. \end{aligned} \quad (6)$$

Here, by assuming that (i) the sequences of the graphs, regression matrices and noises are identically distributed, respectively; (ii) the mean graph is undirected; (iii) the sequences of regression matrices and noises are mutually independent, we obtain the above algorithm. In fact, even if the mean graphs are directed, and the graphs and regression matrices do not satisfy special statistical properties such as independence and stationarity, the algorithm (6) will still be proved to converge.

Remark 3. The communication model (5) with relative-state-dependent communication noises $f_{ji}(x_j(k) - x_i(k))\xi_{ji}(k)$ are reasonable for many realistic applications. (i) In some multiple robots or UAV systems, due to the non-reliability of the communications, the transmission states are more prone to noises whose density functions depend on the distances between the transmitter and receiver, i.e., the relative distances between agents [48]. (ii) In consensus problems with quantized measurements of relative states [49], the logarithmic quantized measurement by agent i of $x_j(k) - x_i(k)$ is given by $x_j(k) - x_i(k) + (x_j(k) - x_i(k))\Delta_{ji}(k)$, which is a special case of (5), where $\Delta_{ji}(k)$ is the quantization uncertainty. (iii) In distributed averaging with Gaussian fading channels [50], the measurement of $x_j(k) - x_i(k)$ is given by $z_{ji}(k) = \xi_{ij}(k)(x_j(k) - x_i(k))$, where $\{\xi_{ij}(k), k \geq 0\}$ are independent Gaussian noises with mean value γ_{ij} . In [50], the above equation is transformed into $z_{ji}(k) = \gamma_{ij}(x_j(k) - x_i(k)) + \Delta_{ij}(k)(x_j(k) - x_i(k))$ with $\Delta_{ij}(k) = \xi(k) - \gamma_{ij}$ being independent zero-mean Gaussian noises, which is a special case of (5).

Denote $\mathcal{F}(k) = \sigma(\mathcal{A}_{\mathcal{G}(s)}, H_i(s), v_i(s), \xi_{ji}(s), j, i \in \mathcal{V}, 0 \leq s \leq k)$ with $\mathcal{F}(-1) = \{\Omega, \emptyset\}$, and $\xi(k) = [\xi_{11}(k), \dots, \xi_{N1}(k), \dots, \xi_{1N}(k), \dots, \xi_{NN}(k)]^T$. For the algorithm (6), we need the following assumptions.

(A1) For the noise intensity function $f_{ji}(\cdot) : \mathbb{R}^n \rightarrow \mathbb{R}$, there exist constants σ and b , such that $|f_{ji}(x)| \leq \sigma\|x\| + b, \forall x \in \mathbb{R}^n$.

Remark 4. Assumption **(A1)** indicates that the communication noises in (5) cover both additive and multiplicative noises, where σ and b are multiplicative and additive noise intensity coefficients, respectively.

(A2) The noises $\{v(k), \mathcal{F}(k), k \geq 0\}$ and $\{\xi(k), \mathcal{F}(k), k \geq 0\}$ are both martingale difference sequences and independent of $\{\mathcal{H}(k), \mathcal{A}_{\mathcal{G}(k)}, k \geq 0\}$. There exists a constant β_v , such that $\sup_{k \geq 0} \mathbb{E}[\|v(k)\|^2 + \|\xi(k)\|^2 | \mathcal{F}(k-1)] \leq \beta_v$ a.s.

Remark 5. Different from [27]-[30], the measurement noises are only assumed to be a martingale difference sequence and independent of the graphs and regression matrices in Assumption **(A2)**. In this paper, neither mutual independence nor spatio-temporal independence is assumed on the regression matrices and graphs. This is applicable to complex scenarios where regression matrices and graphs are spatio-temporal dependent.

The problems of decentralized online regression over graphs have been investigated in most of the literature, including regression with time-varying unknown parameters (R.T.V.P.) (e.g. [32]-[33]) and regression with time-invariant unknown parameters (R.T.I.P.) (e.g. [27]-[30], [34], [38], [45] and [51]-[52]), where different assumptions of models and excitation conditions have been required. Here, we sum up in Table I the assumptions, excitation conditions that they required and their main results.

III. MAIN RESULTS

The convergence and performance analysis of the algorithm (6) are presented in this section. First, Lemma 1 gives a nonnegative supermartingale type inequality of the squared estimation error. Based on which, Theorem 1 proves the almost sure convergence of the algorithm. Then, Theorem 2 gives intuitive convergence conditions for the case with balanced conditional digraphs by Lemma 2. Whereafter, Corollary 2 gives more intuitive convergence conditions for the case

TABLE I: Summary of assumptions, excitation conditions and main results in relevant works.

		Regression matrices	Graphs	Excitation conditions	Results
R.T.V.P.	[32]-[33]	random and time-varying	deterministic, undirected and time-invariant	connected graph and cooperative information condition	L_p -stable tracking errors
R.T.I.P.	[51]	spatio-temporally independent	deterministic and time-invariant	stability of the information matrix	mean and mean square convergence
	[45]	i.i.d. over time	balanced digraphs and homogeneous ergodic Markov chain	global observability and a spanning tree contained in the union of the graphs	almost sure and mean square convergence
	[29]	i.i.d. over time	undirected and i.i.d.	global observability and connected mean graph	almost sure convergence
	[52]	temporally strictly stationary and temporally correlated	deterministic and time-invariant	spatially joint observability	mean square convergence
	[34]	time-invariant and deterministic	undirected and i.i.d.	global observability and connected mean graph	almost sure convergence
	[38]	random and time-varying	random and time-varying digraphs	stochastic spatio-temporal persistence of excitation condition	almost sure and mean square convergence
	This paper	random and time-varying	random and time-varying digraphs	sample path spatio-temporal persistence of excitation condition	almost sure and mean square convergence

with Markovian switching graphs and regression matrices. Finally, Theorem 3 establishes an upper bound for the regret of the algorithm by Lemma 3, and Theorem 4 gives a non-asymptotic rate for the algorithm. The proofs of theorems, Proposition 1 and Corollary 2 are in Appendix A, and those of the lemmas in this section are in Appendix B.

Denote $f_i(k) = \mathbf{diag}(f_{1i}(x_1(k) - x_i(k)), \dots, f_{Ni}(x_N(k) - x_i(k)))$; $Y(k) = \mathbf{diag}(f_1(k), \dots, f_N(k))$; $M(k) = Y(k) \otimes I_n$; $W(k) = \mathbf{diag}(\alpha_1^T(k) \otimes I_n, \dots, \alpha_N^T(k) \otimes I_n)$, where $\alpha_i^T(k)$ is the i -th row of $\mathcal{A}_{\mathcal{G}(k)}$. A compact form equivalent to (6) is given by

$$x(k+1) = [(1 - \lambda(k))I_{Nn} - b(k)\mathcal{L}_{\mathcal{G}(k)} \otimes I_n - a(k)\mathcal{H}^T(k)\mathcal{H}(k)]x(k) + a(k)\mathcal{H}^T(k)y(k) + b(k)W(k)M(k)\xi(k). \quad (7)$$

Denote $\delta(k) = x(k) - \mathbf{1}_N \otimes x_0$ as the global estimation error. Noting that $(\mathcal{L}_{\mathcal{G}(k)} \otimes I_n)(\mathbf{1}_N \otimes x_0) = (\mathcal{L}_{\mathcal{G}(k)}\mathbf{1}_N) \otimes x_0 = 0$ and $\mathcal{H}(k)(\mathbf{1}_N \otimes x_0) = H(k)x_0$, subtracting $\mathbf{1}_N \otimes x_0$ from both sides of (7) at the same time gives

$$\begin{aligned} & \delta(k+1) \\ &= ((1 - \lambda(k))I_{Nn} - b(k)\mathcal{L}_{\mathcal{G}(k)} \otimes I_n - a(k)\mathcal{H}^T(k)\mathcal{H}(k))x(k) + a(k)\mathcal{H}^T(k)y(k) \\ & \quad + b(k)W(k)M(k)\xi(k) - \mathbf{1}_N \otimes x_0 \\ &= ((1 - \lambda(k))I_{Nn} - b(k)\mathcal{L}_{\mathcal{G}(k)} \otimes I_n - a(k)\mathcal{H}^T(k)\mathcal{H}(k))(x(k) - \mathbf{1}_N \otimes x_0 + \mathbf{1}_N \otimes x_0) \end{aligned}$$

$$\begin{aligned}
& +a(k)\mathcal{H}^T(k)y(k) + b(k)W(k)M(k)\xi(k) - \mathbf{1}_N \otimes x_0 \\
& = ((1 - \lambda(k))I_{Nn} - b(k)\mathcal{L}_{\mathcal{G}(k)} \otimes I_n - a(k)\mathcal{H}^T(k)\mathcal{H}(k)) (\delta(k) + \mathbf{1}_N \otimes x_0) \\
& \quad +a(k)\mathcal{H}^T(k)y(k) + b(k)W(k)M(k)\xi(k) - \mathbf{1}_N \otimes x_0 \\
& = ((1 - \lambda(k))I_{Nn} - b(k)\mathcal{L}_{\mathcal{G}(k)} \otimes I_n - a(k)\mathcal{H}^T(k)\mathcal{H}(k)) \delta(k) + a(k)\mathcal{H}^T(k)(y(k) \\
& \quad - H(k)x_0) + b(k)W(k)M(k)\xi(k) - \lambda(k)(\mathbf{1}_N \otimes x_0) \\
& = ((1 - \lambda(k))I_{Nn} - b(k)\mathcal{L}_{\mathcal{G}(k)} \otimes I_n - a(k)\mathcal{H}^T(k)\mathcal{H}(k)) \delta(k) + a(k)\mathcal{H}^T(k)v(k) \\
& \quad +b(k)W(k)M(k)\xi(k) - \lambda(k)(\mathbf{1}_N \otimes x_0) \\
& = \Phi_P(k, 0)\delta(0) + \sum_{i=0}^k a(i)\Phi_P(k, i+1)\mathcal{H}^T(i)v(i) + \sum_{i=0}^k b(i)\Phi_P(k, i+1)W(i)M(i)\xi(i) \\
& \quad - \sum_{i=0}^k \lambda(i)\Phi_P(k, i+1)(\mathbf{1}_N \otimes x_0), \tag{8}
\end{aligned}$$

where $P(k) = (1 - \lambda(k))I_{Nn} - b(k)\mathcal{L}_{\mathcal{G}(k)} \otimes I_n - a(k)\mathcal{H}^T(k)\mathcal{H}(k)$.

Denote $\widehat{\mathcal{L}}_{\mathcal{G}(k)} = \frac{\mathcal{L}_{\mathcal{G}(k)} + \mathcal{L}_{\mathcal{G}(k)}^T}{2}$. For any given positive integers h and k , denote

$$\Lambda_k^h = \lambda_{\min} \left[\sum_{i=kh}^{(k+1)h-1} \left(b(i)\mathbb{E} \left[\widehat{\mathcal{L}}_{\mathcal{G}(i)} | \mathcal{F}(kh-1) \right] \otimes I_n + a(i)\mathbb{E} \left[\mathcal{H}^T(i)\mathcal{H}(i) | \mathcal{F}(kh-1) \right] \right) \right].$$

Denote $V(k) = \|\delta(k)\|^2$. A nonnegative supermartingale type inequality of the squared estimation error $V(k)$ is obtained in the following lemma, which plays a key role in the convergence and performance analyses of the algorithm.

Lemma 1. For the algorithm (6), if Assumptions (A1)-(A2) hold, the algorithm gains $a(k)$, $b(k)$ and $\lambda(k)$ monotonically decrease to zero, and there exists a positive integer h and a positive constant ρ_0 , such that $\sup_{k \geq 0} (\|\mathcal{L}_{\mathcal{G}(k)}\| + (\mathbb{E}[\|\mathcal{H}^T(k)\mathcal{H}(k)\|^{2\max\{h,2\}} | \mathcal{F}(k-1)])^{\frac{1}{2\max\{h,2\}}}) \leq \rho_0$ a.s., then there exists a positive integer k_0 , such that

$$\begin{aligned}
\mathbb{E}[V((k+1)h) | \mathcal{F}(kh-1)] & \leq (1 + \Omega(k))V(kh) - 2 \left(\Lambda_k^h + \sum_{i=kh}^{(k+1)h-1} \lambda(i) \right) V(kh) \\
& \quad + \Gamma(k) \text{ a.s., } k \geq k_0,
\end{aligned}$$

where $\{\Omega(k), k \geq 0\}$ and $\{\Gamma(k), k \geq 0\}$ are nonnegative deterministic real sequences satisfying $\Omega(k) + \Gamma(k) = \mathcal{O}(a^2(kh) + b^2(kh) + \lambda(kh))$.

Remark 6. The entries of $\mathcal{A}_{\mathcal{G}(k)}$ represent the weights of all edges of the graph. It is reasonable to assume that the weights are uniformly bounded with respect to the sample paths.

It is realistic to assume that the regression matrices are conditionally bounded. For example, consider the AR model (2) in Example 1 with i.i.d. Gaussian white noises. For this case, we

can see that $\mathbb{E}[\|H_i^T(k)H_i(k)\||\mathcal{F}(k-1)]$ is bounded but $H_i(k)$ is not uniformly bounded with respect to the sample paths due to the effects of the Gaussian white noises.

The main results are as follows. Firstly, we analyze the convergence of the algorithm.

Theorem 1. For the algorithm (6), if Assumptions (A1)-(A2) hold, the algorithm gains $a(k)$, $b(k)$ and $\lambda(k)$ decrease monotonically, and there exists a positive integer h and a positive constant ρ_0 , such that (i) $\sum_{k=0}^{\infty} \Lambda_k^h = \infty$ a.s. with $\inf_{k \geq 0} (\Lambda_k^h + \lambda(k)) \geq 0$ a.s.; (ii) $\sum_{k=0}^{\infty} (a^2(k) + b^2(k) + \lambda(k)) < \infty$; (iii) $\sup_{k \geq 0} (\|\mathcal{L}_{\mathcal{G}(k)}\| + (\mathbb{E}[\|\mathcal{H}^T(k)\mathcal{H}(k)\|^{2^{\max\{h,2\}}}]|\mathcal{F}(k-1)])^{\frac{1}{2^{\max\{h,2\}}}} \leq \rho_0$ a.s., then $\lim_{k \rightarrow \infty} x_i(k) = x_0$, $i \in \mathcal{V}$ a.s.

The following proof sketch provides the main steps in the proof of Theorem 1, from which we have solved the limitations of prior works.

Proof sketch. The non-commutative, non-independent and non-stationary random matrices pose intrinsic difficulties for analyzing the convergence of the online algorithm (6), for which our proof is mainly consisted of the following three steps. (I) We start by introducing the tools of conditional mathematical expectations in probability theory, and then derive the upper bound of the state transition matrix, i.e., $\|\mathbb{E}[\Phi_P^T((k+1)h-1, kh)\Phi_P((k+1)h-1, kh)|\mathcal{F}(kh-1)]\| \leq 1 - 2(\Lambda_k^h + \sum_{i=kh}^{(k+1)h-1} \lambda(i)) + \mathcal{O}(a^2(kh) + b^2(kh) + \lambda^2(kh))$ a.s. through the binomial expansion techniques, which helps us to transform the analysis of random matrix products into that of the minimum eigenvalue of the information matrix Λ_k^h and we no longer need to separate the product of the random matrices by assuming independence as [28]-[29]. (II) Different from the existing techniques in [28]-[29] and [37]-[38], which directly analyzed the mean squared error $\mathbb{E}[V(k)]$, we next turn to consider the squared estimation error $V(kh)$. By applying the inequality techniques in probability theory, we reveal the relation between $\mathbb{E}[V((k+1)h)|\mathcal{F}(kh-1)]$ and $V(kh)$ and derive the nonnegative supermartingale type inequality of the squared estimation error in Lemma 1. (III) Then, by the probabilistic structure of the estimation error obtained in Lemma 1 and the convergence theorem for nonnegative supermartingales, we conclude that $V(kh)$ and the infinite series $\sum_{k=0}^{\infty} (\Lambda_k^h + \sum_{i=kh}^{(k+1)h-1} \lambda(i))V(kh)$ almost surely converge by choosing appropriate algorithm gains and the regularization parameter. (IV) Finally, we prove that $V(kh) \rightarrow 0$ a.s. as $k \rightarrow \infty$ by the condition (i) in Theorem 1, where we no longer require the minimum eigenvalue of the information matrix Λ_k^h has a positive and deterministic lower bound as in [37]-[38]. As a consequence, our results not only remove the reliance on the special statistical properties of

regression matrices and graphs, but also weaken the conditions in our previous works.

Remark 7. The choices of the algorithm gains $a(k)$, $b(k)$ and the regularization parameter $\lambda(k)$ is crucial for the nodes to successfully estimate the unknown true parameter vector x_0 . The conditions (i) and (iii) in Theorem 1 imply $\sum_{k=0}^{\infty} (a(k) + b(k)) = \infty$, which means that the algorithm gains can not be too small. Besides, to reduce the effects of perturbing the innovations and consensus by the measurement and communication noises, the monotonically decreasing condition for the algorithm gains and the condition (ii) in Theorem 1 ensure that all nodes avoid making excessive changes to the current estimate when acquiring new noisy data, which guarantees that the impact of each successive new noisy data decays with the algorithm gains $a(k)$, $b(k)$ and $\lambda(k)$. The conditions (i) and (ii) give what need to be satisfied for the algorithm gains.

It is worth noting that the information matrix Λ_k^h in the condition (i) is coupled with the gains, graphs and regression matrices. For some special cases, Theorem 2 and Corollary 2 later provide acceptable ranges on the values of these gains.

Remark 8. Most existing literature on decentralized online regression suppose that the mean graphs are balanced and strongly connected (e.g. [28]-[29] and [36]). Here, the condition (i) in Theorem 1 may still hold even if the mean graphs are unbalanced and not strongly connected. For example, consider a fixed digraph $\mathcal{G} = \{\mathcal{V} = \{1, 2\}, \mathcal{A}_{\mathcal{G}} = [w_{ij}]_{2 \times 2}\}$ with $w_{12} = 1$ and $w_{21} = 0$. Obviously, \mathcal{G} is unbalanced and not strongly connected. Suppose $H_1 = 0, H_2 = 1$. Choose $a(k) = b(k) = \frac{1}{k+1}$. We have $\lambda_{\min}(b(k)\hat{\mathcal{L}}_{\mathcal{G}} + a(k)\mathcal{H}^T\mathcal{H}) = \frac{1}{k+1}\lambda_{\min}(\hat{\mathcal{L}}_{\mathcal{G}} + \mathcal{H}^T\mathcal{H}) = \frac{1}{2k+2}$. Then, the condition (i) holds with $h = 1$ and $\sum_{k=0}^{\infty} \Lambda_k^h = \sum_{k=0}^{\infty} \frac{1}{2k+2} = \infty$.

For the special case without regularization, we directly obtain the following corollary by Theorem 1.

Corollary 1. For the algorithm (6) with $\lambda(k) \equiv 0$, if Assumptions **(A1)**-**(A2)** hold, and there exists a positive integer h and a positive constant ρ_0 , such that (i) $\sum_{k=0}^{\infty} \Lambda_k^h = \infty$ a.s. with $\inf_{k \geq 0} \Lambda_k^h \geq 0$ a.s.; (ii) $\sum_{k=0}^{\infty} (a^2(k) + b^2(k)) < \infty$; (iii) $\sup_{k \geq 0} (\|\mathcal{L}_{\mathcal{G}(k)}\| + (\mathbb{E}[\|\mathcal{H}^T(k)\mathcal{H}(k)\|^{2\max\{h, 2\}}] | \mathcal{F}(k-1)])^{\frac{1}{2\max\{h, 2\}}} \leq \rho_0$ a.s., then $\lim_{k \rightarrow \infty} x_i(k) = x_0$, $i \in \mathcal{V}$ a.s.

Remark 9. The conditions for the regularized algorithm in Theorem 1 are less conservative

than those for the non-regularized algorithm in Corollary 1. For example, consider the digraph sequence $\{\mathcal{G}(k) = \{\mathcal{V} = \{1, 2\}, \mathcal{A}_{\mathcal{G}(k)} = [w_{ij}(k)]_{2 \times 2}\}, k \geq 0\}$ with $w_{12}(k) = \begin{cases} 1, & k = 2m; \\ \frac{1}{k}, & k = 2m + 1 \end{cases}$ and $w_{21}(k) = 0$, and the regression matrices $H_1(k) = 0, H_2(k) = \begin{cases} 1, & k = 2m; \\ \frac{1}{\sqrt{5k}}, & k = 2m + 1 \end{cases}, m \geq 0$. Choose $a(k) = b(k) = \frac{1}{k+1}$. Then, we have $\sum_{k=0}^{\infty} \Lambda_k^1 = \infty$. Obviously, the condition (i) in Corollary 1 does not hold since $\inf_{k \geq 0} \Lambda_k^1 \leq \inf_{k \geq 0} \Lambda_{2k+1}^1 = \inf_{k \geq 0} (-\frac{\sqrt{41}-6}{10(2k+1)^2}) < 0$. Choose $\lambda(k) = \frac{1}{10(k+1)^2}$. Then, the condition (i) in Theorem 1 holds with $\inf_{k \geq 0} (\Lambda_k^1 + \lambda(k)) = 0$.

Subsequently, we list the conditions on the algorithm gains that may be needed later.

(C1) The algorithm gains $\{a(k), k \geq 0\}, \{b(k), k \geq 0\}$ and $\{\lambda(k), k \geq 0\}$ are all nonnegative sequences monotonically decreasing, and satisfy $\sum_{k=0}^{\infty} \min\{a(k), b(k)\} = \infty, \sum_{k=0}^{\infty} (a^2(k) + b^2(k) + \lambda(k)) < \infty, a(k) = \mathcal{O}(a(k+1)), b(k) = \mathcal{O}(b(k+1))$ and $\lambda(k) = o(\min\{a(k), b(k)\})$.

(C2) The algorithm gains $\{a(k), k \geq 0\}, \{b(k), k \geq 0\}$ and $\{\lambda(k), k \geq 0\}$ are all nonnegative sequences monotonically decreasing to zero, and satisfy $\sum_{k=0}^{\infty} \min\{a(k), b(k)\} = \infty, a(k) = \mathcal{O}(a(k+1)), b(k) = \mathcal{O}(b(k+1))$ and $a^2(k) + b^2(k) + \lambda(k) = o(\min\{a(k), b(k)\})$.

Remark 10. As mentioned previously, the regularization gain $\lambda(k)$ has the function of constraining the estimation $x_i(k+1)$ in the algorithm (6). To ensure that the algorithm can cooperatively estimate the unknown true parameter vector x_0 by means of the innovation and consensus terms, Conditions **(C1)**-**(C2)** require the regularization gain $\lambda(k)$ to decay faster than the innovation gain $a(k)$ and the consensus gain $b(k)$.

Next, we consider the sequence of balanced conditional digraphs

$$\Gamma_1 = \{\{\mathcal{G}(k), k \geq 0\} | \mathbb{E}[\mathcal{A}_{\mathcal{G}(k)} | \mathcal{F}(k-1)] \succeq \mathbf{0}_{N \times N} \text{ a.s., } \mathcal{G}(k|k-1) \text{ is balanced a.s., } k \geq 0\}.$$

Here, $\mathbb{E}[\mathcal{A}_{\mathcal{G}(k)} | \mathcal{F}(m)], m < k$, is called the *conditional generalized weighted adjacency matrix* of $\mathcal{A}_{\mathcal{G}(k)}$ w.r.t. $\mathcal{F}(m)$, and its associated random graph is called the *conditional digraph* of $\mathcal{G}(k)$ w.r.t. $\mathcal{F}(m)$, denoted by $\mathcal{G}(k|m)$, i.e., $\mathcal{G}(k|m) = \{\mathcal{V}, \mathbb{E}[\mathcal{A}_{\mathcal{G}(k)} | \mathcal{F}(m)]\}$.

For any given positive integers h and k , we denote

$$\tilde{\Lambda}_k^h = \lambda_{\min} \left[\sum_{i=kh}^{(k+1)h-1} \mathbb{E} \left[\hat{\mathcal{L}}_{\mathcal{G}(i)} \otimes I_n + \mathcal{H}^T(i) \mathcal{H}(i) | \mathcal{F}(kh-1) \right] \right].$$

Then $\tilde{\Lambda}_k^h$ contains information of both the Laplacian matrices of the graphs and regression matrices. As shown in Remark 7, the information matrix Λ_k^h in the condition (i) in Theorem 1 is coupled with the gains, graphs and regression matrices, it is generally difficult to decouple the algorithm gains from Λ_k^h for general graphs and regression matrices since $\hat{\mathcal{L}}_{\mathcal{G}(k)}$ is indefinite. For the case with the conditionally balanced digraphs $\mathcal{G}(k) \in \Gamma_1$, we have $\Lambda_k^h \geq \min\{a(i), b(i), kh \leq i < (k+1)h\} \tilde{\Lambda}_k^h$ a.s. The following lemma gives a lower bound of $\tilde{\Lambda}_k^h$, where one can see how the conditionally balanced graphs and regression matrices of all nodes affect the lower bound, respectively.

Lemma 2. For the algorithm (6), suppose that $\{\mathcal{G}(k), k \geq 0\} \in \Gamma_1$, and there exists a positive constant ρ_0 , such that

$$\sup_{k \geq 0} \mathbb{E} [\|\mathcal{H}^T(k)\mathcal{H}(k)\| | \mathcal{F}(k-1)] \leq \rho_0 \text{ a.s.}$$

Then, for any given positive integer h ,

$$\begin{aligned} \tilde{\Lambda}_k^h \geq & \frac{\lambda_2 \left[\sum_{i=kh}^{(k+1)h-1} \mathbb{E} [\hat{\mathcal{L}}_{\mathcal{G}(i)} | \mathcal{F}(kh-1)] \right]}{2Nh\rho_0 + N\lambda_2 \left[\sum_{i=kh}^{(k+1)h-1} \mathbb{E} [\hat{\mathcal{L}}_{\mathcal{G}(i)} | \mathcal{F}(kh-1)] \right]} \\ & \times \lambda_{\min} \left(\sum_{i=1}^N \sum_{j=kh}^{(k+1)h-1} \mathbb{E} [H_i^T(j)H_i(j) | \mathcal{F}(kh-1)] \right) \text{ a.s.} \end{aligned}$$

Remark 11. If the network structure degenerates into a deterministic, undirected and connected graph \mathcal{G} , and $H_i^T(k)H_i(k) \leq \rho_0 I_n$, $i \in \mathcal{V}$ a.s., then the inequality in Lemma 2 degenerates to

$$\tilde{\Lambda}_k^h \geq \frac{\lambda_2(\mathcal{L}_{\mathcal{G}})}{2Nh\rho_0 + N\lambda_2(\mathcal{L}_{\mathcal{G}})} \lambda_{\min} \left(\sum_{i=1}^N \sum_{j=kh}^{(k+1)h-1} \mathbb{E} [H_i^T(j)H_i(j) | \mathcal{F}(kh-1)] \right) \text{ a.s.,}$$

which is given in [32]-[33].

Then, we give intuitive convergence conditions for the case with balanced conditional digraphs. We first introduce the following definitions.

Definition 1. For the random undirected graph sequence $\{\mathcal{G}(k), k \geq 0\}$, if there exists a positive integer h and a positive constant θ , such that

$$\inf_{k \geq 0} \lambda_2 \left(\sum_{i=kh}^{(k+1)h-1} \mathbb{E} [\mathcal{L}_{\mathcal{G}(i)} | \mathcal{F}(kh-1)] \right) \geq \theta \text{ a.s.,}$$

then $\{\mathcal{G}(k), k \geq 0\}$ is said to be *uniformly conditionally jointly connected*.

Remark 12. Definition 1 means that the conditional graphs $\mathcal{G}(i|kh-1), i = kh, \dots, (k+1)h-1$, are jointly connected a.s., and the average algebraic connectivity is uniformly positive bounded away from zero.

Definition 2. For the sequence of regression matrices $\{H_i(k), i = 1, \dots, N, k \geq 0\}$, if there exists a positive integer h and a positive constant θ , such that

$$\inf_{k \geq 0} \lambda_{\min} \left(\sum_{i=1}^N \sum_{j=kh}^{(k+1)h-1} \mathbb{E} [H_i^T(j) H_i(j) | \mathcal{F}(kh-1)] \right) \geq \theta \text{ a.s.},$$

then $\{H_i(k), i = 1, \dots, N, k \geq 0\}$ is said to be *uniformly conditionally spatio-temporally jointly observable*.

Denote the symmetrized graph of $\mathcal{G}(k)$ by $\widehat{\mathcal{G}}(k) = \{\mathcal{V}, \mathcal{E}_{\mathcal{G}(k)} \cup \mathcal{E}_{\widetilde{\mathcal{G}}(k)}, \frac{\mathcal{A}_{\mathcal{G}(k)} + \mathcal{A}_{\widetilde{\mathcal{G}}(k)}^T}{2}\}$, where $\widetilde{\mathcal{G}}(k)$ is the reversed digraph of $\mathcal{G}(k)$ [39]. For the case with ϕ -mixing random process [32], we have the following proposition.

Proposition 1. Suppose that $\mathcal{G}(k) \in \Gamma_1$ and $\{\mathcal{A}_{\mathcal{G}(k)}, H_i(k), v_i(k), \xi_i(k), i = 1, \dots, N, k \geq 0\}$ is a ϕ -mixing sequence.

- (i) If there exists a positive integer h_1 and a positive constant α_1 , such that $\inf_{k \geq 0} \lambda_2(\sum_{i=kh_1}^{(k+1)h_1-1} \mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(i)}]) \geq \alpha_1$, then $\{\widehat{\mathcal{G}}(k), k \geq 0\}$ is *uniformly conditionally jointly connected*.
- (ii) If there exists a positive integer h_2 and a positive constant α_2 , such that $\inf_{k \geq 0} \lambda_{\min}(\sum_{i=1}^N \sum_{j=kh_2}^{(k+1)h_2-1} \mathbb{E}[H_i^T(j) H_i(j)]) \geq \alpha_2$, then $\{H_i(k), i = 1, \dots, N, k \geq 0\}$ is *uniformly conditionally spatio-temporally jointly observable*.

Theorem 2. For the algorithm (6), suppose that $\{\mathcal{G}(k), k \geq 0\} \in \Gamma_1$, Assumptions **(A1)**-(**A2**) hold, and there exists a positive integer h and a positive constant ρ_0 , such that (i) $\{\widehat{\mathcal{G}}(k), k \geq 0\}$ is *uniformly conditionally jointly connected*; (ii) $\{H_i(k), i \in \mathcal{V}, k \geq 0\}$ is *uniformly conditionally spatio-temporally jointly observable*; (iii) $\sup_{k \geq 0} (\|\mathcal{L}_{\mathcal{G}(k)}\| + (\mathbb{E}[\|\mathcal{H}^T(k) \mathcal{H}(k)\|^{2^{\max\{h, 2\}}}] | \mathcal{F}(k-1)])^{\frac{1}{2^{\max\{h, 2\}}}}) \leq \rho_0$ a.s.

(I). If Condition **(C1)** holds, then $\lim_{k \rightarrow \infty} x_i(k) = x_0, i \in \mathcal{V}$ a.s.

(II). If Condition **(C2)** holds, then $\lim_{k \rightarrow \infty} \mathbb{E}[\|x_i(k) - x_0\|^2] = 0, i \in \mathcal{V}$.

The combination of the conditions (i)-(ii) in Theorem 2 with Condition **(C1)** or Condition **(C2)** gives an intuitive sufficient condition for the condition (i) in Theorem 1 to hold.

The condition (i) in Theorem 1 is called the *sample path spatio-temporal persistence of excitation* condition, which is an indispensable part of the convergence and performance analysis of the algorithm. Specifically, *spatio-temporal persistence of excitation* means that the infinite series of the minimum eigenvalues of the information matrices composed of the graphs and regression matrices in fixed-length time intervals diverges for almost all sample paths, i.e., $\sum_{k=0}^{\infty} \Lambda_k^h = \infty$ a.s., which is to avoid the failure of estimating the unknown true parameter vector due to lack of effective measurement information or sufficient information exchange among nodes. To illustrate this, let us consider the extreme case that the regression matrix of each node at each instant is always zero, then $\Lambda_k^h = 0$ for any given positive integer h . In this case, no matter how to design the algorithm gains, one can not obtain any information about the unknown true parameter vector since there is no measurement information and no information interflow. Here, *spatio-temporality* focuses on the state of the information matrices consisting of the graphs and regression matrices of all nodes over a fixed-length time period rather than the state at each instant, where the temporality is captured by h . From Theorem 2, we know that neither the *locally temporally joint observability* of each node, i.e., $\inf_{k \geq 0} \lambda_{\min}(\mathbb{E}[\sum_{j=kh}^{(k+1)h-1} H_i^T(j)H_i(j)|\mathcal{F}(kh-1)])$ being uniformly bounded away from zero for each node i , nor the *instantaneously globally spatially joint observability* of all the regression models, i.e., $\inf_{k \geq 0} \lambda_{\min}(\mathbb{E}[\sum_{i=1}^N H_i^T(j)H_i(j)|\mathcal{F}(kh-1)])$ being uniformly bounded away from zero for each instant j , is necessary.

At present, most results on decentralized online linear regression algorithms (e.g. [28]-[29] and [45]-[46]) all require that the regression matrices and graphs satisfy some special statistical properties, such as i.i.d., spatio-temporal independence or stationary, etc. However, these special statistical assumptions are difficult to be satisfied if the regression matrices are generated by the auto-regressive models. In order to solve this problem, in the past several decades, many scholars have proposed the persistence of excitation condition based on the conditional expectations of the regression matrices. The stochastic persistence of excitation condition was first proposed in the analysis of the centralized Kalman filter algorithm in [41] and then refined in [42]. For the decentralized adaptive filtering algorithms in [32]-[33], the cooperative information condition on the conditional expectations of the regression matrices was proposed for the case with deterministic connected graphs. For the decentralized online estimation algorithms over random time-varying graphs in [37]-[38], the stochastic spatio-temporal persistence of excitation

condition was proposed. The stochastic spatio-temporal persistence of excitation condition in [38] means that the minimum eigenvalue of the matrix consisting of the spatial-temporal observation matrices and Laplacian matrices in each time period $[kh, (k+1)h-1]$ has a positive lower bound $c(k)$ independent of the sample paths, i.e., $\Lambda_k^h \geq c(k)$ a.s. with $\sum_{k=0}^{\infty} c(k) = \infty$. Here, in Corollary 1, we show that this is actually not needed. Therefore, the *sample path spatio-temporal persistence of excitation* condition in this paper is more general than the stochastic spatio-temporal persistence of excitation condition in [38].

We give an example for which the stochastic spatio-temporal persistence of excitation condition does not hold but the *sample path spatio-temporal persistence of excitation* condition hold.

Example 2. Consider a fixed digraph $\mathcal{G} = \{\mathcal{V} = \{1, 2\}, \mathcal{A}_{\mathcal{G}} = [w_{ij}]_{2 \times 2}\}$ with $w_{12} = 1$ and $w_{21} = 0$. Let $H_1 = 0$ and $H_2 = \sqrt{x}$, where the random variable x is uniformly distributed in $(0.25, 1.25)$. Choose $a(k) = b(k) = \frac{1}{k+1}$. For any given positive integer h , if Assumption (A2) holds, then the *sample path spatio-temporal persistence of excitation* condition holds and there does not exist a positive real sequence $\{c(k), k \geq 0\}$ satisfying $\Lambda_k^h \geq c(k)$ a.s., i.e., the stochastic spatio-temporal persistence of excitation condition in [38] does not hold.

Therefore, the *sample path spatio-temporal persistence of excitation* condition weakens the stochastic spatio-temporal persistence of excitation condition in [38]. To our best knowledge, we have obtained the most general persistence of excitation condition ever.

Whereafter, we give more intuitive convergence conditions for the case with Markovian switching graphs and regression matrices. We first make the following assumption.

(A3) $\{\langle \mathcal{H}(k), \mathcal{A}_{\mathcal{G}(k)} \rangle, k \geq 0\} \subseteq \mathcal{S}$ is a homogeneous and uniform ergodic Markov chain with a unique stationary distribution π , $\sup_{l \geq 1} \|\mathcal{A}_l\| < \infty$ and $\sup_{l \geq 1} \|\mathcal{H}_l\| < \infty$.

Here, $\mathcal{H}_l = \text{diag}(H_{1,l}, \dots, H_{N,l})$ and $\mathcal{S} = \{\langle \mathcal{H}_l, \mathcal{A}_l \rangle, l \geq 1\}$, where $\{H_{i,l} \in \mathbb{R}^{n_i \times n}, l \geq 1\}$ is the state space of the regression matrices of node i and $\{\mathcal{A}_l, l \geq 1\}$ is the state space of the weighted adjacency matrices, $\pi = [\pi_1, \pi_2, \dots]^T$, $\pi_l \geq 0$, $l \geq 1$, and $\sum_{l=1}^{\infty} \pi_l = 1$ with π_l representing $\pi(\langle \mathcal{H}_l, \mathcal{A}_l \rangle)$. The readers may refer to [54] for the definition of uniform ergodic Markov chain.

Corollary 2. For the algorithm (6), suppose that $\{\mathcal{G}(k), k \geq 0\} \in \Gamma_1$, Assumptions (A1)-(A3) hold, the stationary weighted adjacency matrix $\sum_{l=1}^{\infty} \pi_l \mathcal{A}_l$ is nonnegative and its associated graph is balanced with a spanning tree, and the regression model (1) is *spatio-temporally jointly*

observable, i.e.,

$$\lambda_{\min} \left(\sum_{i=1}^N \left(\sum_{l=1}^{\infty} \pi_l H_{i,l}^T H_{i,l} \right) \right) > 0. \quad (9)$$

(I). If Condition **(C1)** holds, then $\lim_{k \rightarrow \infty} x_i(k) = x_0$, $i \in \mathcal{V}$ a.s.

(II). If Condition **(C2)** holds, then $\lim_{k \rightarrow \infty} \mathbb{E}[\|x_i(k) - x_0\|^2] = 0$, $i \in \mathcal{V}$.

Remark 13. Corollary 2 shows that for homogeneous and uniform ergodic Markovian switching graphs and regression matrices, the *sample path spatio-temporal persistence of excitation* condition holds if the stationary graph is balanced with a spanning tree and the regression model is *spatio-temporally jointly observable*, that is, (9) holds, implying that neither *local observability* of each node, i.e., $\lambda_{\min}(\sum_{l=1}^{\infty} \pi_l H_{i,l}^T H_{i,l}) > 0$, $i \in \mathcal{V}$, nor *instantaneous global observability* of all the regression models, i.e., $\lambda_{\min}(\sum_{i=1}^N H_{i,l}^T H_{i,l}) > 0$, $l = 1, 2, \dots$, is needed.

To appraise the performance of the algorithm (6), the regret upper bound of the decentralized online regularized algorithm will be taken into account. The loss function of node j at instant t is defined by

$$l_{j,t}(x) \triangleq \frac{1}{2} \|H_j(t)x - y_j(t)\|^2, \quad x \in \mathbb{R}^n.$$

The performance of the algorithm (6) is appraised, at node i , through its regret defined by

$$\text{Regret}_{\text{LMS}}(i, T) \triangleq \sum_{t=0}^T \sum_{j=1}^N \mathbb{E}[l_{j,t}(x_i(t))] - \sum_{t=0}^T \sum_{j=1}^N \mathbb{E}[l_{j,t}(x_{\text{LMS}}^*)],$$

where

$$x_{\text{LMS}}^* \triangleq \arg \min_{x \in \mathbb{R}^n} \sum_{t=0}^T \sum_{j=1}^N \frac{1}{2} \mathbb{E}[\|H_j(t)x - y_j(t)\|^2]$$

is the linear optimal estimated parameter. Before giving the upper bound of $\text{Regret}_{\text{LMS}}(i, T)$, we need the following lemma.

Lemma 3. For the algorithm (6), suppose that Assumption **(A2)** holds and there exists a positive constant ρ_0 , such that

$$\sup_{k \geq 0} (\mathbb{E}[\|\mathcal{H}^T(k)\mathcal{H}(k)\| | \mathcal{F}(k-1)]) \leq \rho_0 \text{ a.s.}$$

Then

$$\text{Regret}_{\text{LMS}}(i, T) \leq \frac{1}{2} N \rho_0 \sum_{t=0}^T \mathbb{E}[V(t)], \quad i \in \mathcal{V}.$$

Remark 14. Lemma 3 indicates that the regret of each node at time instant T can be bounded by the accumulation of the mean square estimation errors during the period $[0, T]$, which directs the way for obtaining the regret upper bound.

For the *uniformly conditionally jointly connected* graphs and *uniformly conditionally spatio-temporally jointly observable* regression models, the following theorem gives an upper bound of $\text{Regret}_{\text{LMS}}(i, T)$, $i \in \mathcal{V}$.

Theorem 3. For the algorithm (6), suppose that $\{\mathcal{G}(k), k \geq 0\} \in \Gamma_1$ and Assumptions (A1)-(A2) hold. If $a(k) = b(k) = \frac{c_1}{(k+1)^\tau}$ and $\lambda(k) = \frac{c_2}{(k+1)^{2\tau}}$, $c_1 > 0$, $c_2 > 0$, $0.5 < \tau < 1$, and there exists a positive integer h and a positive constant ρ_0 , such that (i) $\{\hat{\mathcal{G}}(k), k \geq 0\}$ is *uniformly conditionally jointly connected*; (ii) $\{H_i(k), i \in \mathcal{V}, k \geq 0\}$ is *uniformly conditionally spatio-temporally jointly observable*; (iii) $\sup_{k \geq 0} (\|\mathcal{L}_{\mathcal{G}(k)}\| + (\mathbb{E}[\|\mathcal{H}^T(k)\mathcal{H}(k)\|^{2^{\max\{h, 2\}}} | \mathcal{F}(k-1)]))^{\frac{1}{2^{\max\{h, 2\}}}} \leq \rho_0$ a.s., then the regret of node i satisfies

$$\text{Regret}_{\text{LMS}}(i, T) = \mathcal{O}(T^{1-\tau} \ln T), \quad i \in \mathcal{V}. \quad (10)$$

Remark 15. Yuan *et al.* [36] studied the non-regularized decentralized online linear regression algorithm over the fixed graph. In this paper, we consider the decentralized online regularized linear regression algorithm over the random time-varying graphs. Theorem 3 shows that the upper bound of the regret is $\mathcal{O}(T^{1-\tau} \ln T)$, where $\tau \in (0.5, 1)$ is a constant depending on the decaying algorithm gains, which improves the regret upper bound $\mathcal{O}(\sqrt{T})$ with the fixed gain in [36].

For the *uniformly conditionally jointly connected* graphs and *uniformly conditionally spatio-temporally jointly observable* regression models, the following theorem further gives a non-asymptotic rate for the algorithm.

Theorem 4. For the algorithm (6), suppose that $\{\mathcal{G}(k), k \geq 0\} \in \Gamma_1$ and Assumptions (A1)-(A2) hold. If $a(k) = b(k) = \frac{1}{(k+1)^\tau}$ and $\lambda(k) = \frac{1}{(k+1)^{2\tau}}$, $0.5 < \tau < 1$, and there exists a positive integer h and a positive constant ρ_0 , such that (i) $\{\hat{\mathcal{G}}(k), k \geq 0\}$ is *uniformly conditionally jointly connected*; (ii) $\{H_i(k), i \in \mathcal{V}, k \geq 0\}$ is *uniformly conditionally spatio-temporally jointly observable*; (iii) $\sup_{k \geq 0} (\|\mathcal{L}_{\mathcal{G}(k)}\| + (\mathbb{E}[\|\mathcal{H}^T(k)\mathcal{H}(k)\|^{2^{\max\{h, 2\}}} | \mathcal{F}(k-1)]))^{\frac{1}{2^{\max\{h, 2\}}}} \leq \rho_0$ a.s., then

there exist constants c_i , $i = 1, \dots, 6$, such that

$$\mathbb{E}[V(k)] \leq 2^h \left(\frac{25c_4(2h)^{1-\tau} \ln k}{c_5 k^{1-\tau}} + N \|x_0\|^2 e^{-\frac{c_5 \left(\left(\frac{k}{2h} \right)^{1-\tau} - 2 \right)}{1-\tau}} \right) + \frac{c_6}{(k+2)^2},$$

when

$$k \geq \left(\frac{12}{c_5(1-\tau)} \ln \left(\frac{4}{c_5(1-\tau)} \right) \right)^{\frac{1}{1-\tau}} + \left\lfloor \frac{2c_1}{hc_2} + c_3^2 \right\rfloor + 16h^2(\rho_0 + 1)^2 + 1.$$

IV. NUMERICAL EXAMPLE

We consider the graphs composed of 10 nodes, and the states are $x_i(k)$, $i = 1, \dots, 10$, $k \geq 0$, whose initial states are given by $x_1(0) = [12, 11, 6]^T$, $x_2(0) = [10, 16, 8]^T$, $x_3(0) = [14, 16, 13]^T$, $x_4(0) = [15, 12, 9]^T$, $x_5(0) = [10, 14, 8]^T$, $x_6(0) = [9, 16, 7]^T$, $x_7(0) = [8, 13, 11]^T$, $x_8(0) = [12, 10, 9]^T$, $x_9(0) = [13, 10, 9]^T$, $x_{10}(0) = [12, 10, 8]^T$. Each node estimates the unknown true parameter vector $x_0 = [5, 4, 3]^T$.

Model settings. The local observation data $y_i(k)$ of node i is given by $y_i(k) = H_i(k)x_0 + v_i(k)$, $i = 1, \dots, 10$. Here, the regression matrices are taken as

$$\begin{pmatrix} 0 & 0 & 0 \\ \bar{h}_{s,t,k} & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & \bar{h}_{s+2,t,k} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & \bar{h}_{5,t,k} & 0 \\ 0 & 0 & 0 \\ \bar{h}_{6,t,k} & 0 & 0 \end{pmatrix}, \quad s = 1, 2, \quad t = 1, 2,$$


where $\bar{h}_{s,t,k} = (-1)^s h_{s,t}(k) + (-1)^t 0.5$, $\bar{h}_{s+2,t,k} = (-1)^s h_{s+2,t}(k) + (-1)^t 0.5$, $\bar{h}_{5,t,k} = (-1)^t h_{5,t}(k) + (-1)^t 0.5$, $\bar{h}_{6,t,k} = (-1)^{t+1} h_{6,t}(k) + (-1)^{t+1} 0.5$, and $h_{i,j}(k)$, $i = 1, 2, 3, 4, 5, 6$, $j = 1, 2$, are independent random variables with uniform distribution on $(0, 1)$. The signal at the receiver side of a communication channel is described by (5), where the noise intensity functions are given by $f_{ji}(x_j(k) - x_i(k)) = 0.1 \|x_j(k) - x_i(k)\| + 0.1$, $i, j = 1, \dots, 10$. The updating rules of the estimation states follow algorithm (6), where the random weights $\{w_{ij}(k), i, j = 1, \dots, 10, k \geq 0\}$ are selected by the following rules. When $k = 2m$, $m \geq 0$, the random weights are uniformly distributed in the interval $[0, 1]$; when $k \neq 2m$, $m \geq 0$, the random weights are uniformly distributed in $[-0.5, 0.5]$. So the random weights may be negative at some time instants. Here, $\{w_{ij}(k), i, j = 1, \dots, 10, k \geq 0\}$ are spatio-temporally independent. Thus, when $k = 2m$, $m \geq 0$, the average graph is balanced and connected, and when $k \neq 2m$, $m \geq 0$, the average graph is empty. Suppose that the measurement noises $\{v_i(k), i = 1, \dots, 10, k \geq 0\}$ and the communication noises $\{\xi_{ji}(k), i, j = 1, \dots, 10, k \geq 0\}$ are random variables obeying the

TABLE II: Settings of the algorithm gains

	$a(k)$	$b(k)$	$\lambda(k)$
Setting I	$(k+1)^{-0.6}$	$(k+1)^{-0.6}$	$(k+1)^{-2}$
Setting II	$(k+1)^{-0.6}$	$(k+1)^{-0.6}$	$(k+1)^{-3}$
Setting III	$(k+1)^{-0.8}$	$(k+1)^{-0.8}$	$(k+1)^{-2}$
Setting IV	$(k+1)^{-0.8}$	$(k+1)^{-0.8}$	$(k+1)^{-3}$

normal distribution and independent of the random graphs. Different settings of the algorithm gains are shown in Table II, where we take $a(k) = b(k)$ for simplicity.

To verify the *sample path spatio-temporal persistence of excitation* condition, Fig. 1 presents the curves of the quantity $R(k) \triangleq (\sum_{i=0}^k \Lambda_i^2)^{-1} = (\sum_{i=0}^k \lambda_{\min}(\sum_{j=2i}^{2i+1} (b(j) \mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(j)} | \mathcal{F}(2i-1)] \otimes I_2 + a(j) \mathbb{E}[\mathcal{H}^T(j) \mathcal{H}(j) | \mathcal{F}(2i-1)]))^{-1}$ over time k under different settings in Table II.

Fig. 1: The sample paths of $R(k)$ with different settings.

Here, all the conditions of Theorem 1 hold with $h = 2$ and $\rho_0 = 5$. The trajectories of the estimation errors are shown in Fig. 2. It can be seen that as time goes on, the estimations of each node's algorithms with different algorithm gains in Table II converge to the true value of x_0 almost surely. We can see from Fig. 1 and Fig. 2 that larger $\sum_{i=0}^k \Lambda_i^2$ leads to faster convergence of the algorithm. It shows in Fig. 2 that (i) for the same regularization parameter $\lambda(k)$, larger gains $a(k)$ and $b(k)$ accelerate the convergence of the algorithm; (ii) for the same gains, it is possible that the algorithm with larger regularization parameter converges faster than that with smaller one.





Fig. 2: The sample paths of errors with different settings.

Meanwhile, the trajectories of the mean value of the norms of agents' states are plotted in Fig. 3, where setting V and setting VI are taken as $a(k) = b(k) = (k + 1)^{-0.6}$, $\lambda(k) = 0$ and $a(k) = b(k) = (k + 1)^{-0.8}$, $\lambda(k) = 0$, respectively. Fig. 3 shows that compared with the non-regularized algorithms, the regularization is effective for reducing the magnitudes of estimations of the unknown true vector, which helps to reduce the model complexity and to avoid overfitting.


Subsequently, we take $a(k) = b(k) = 0.1(k + 1)^{-0.6}$ and $\lambda(k) = 0.2(k + 1)^{-1.2}$. A direct calculation gives

$$\inf_{k \geq 0} \lambda_{\min} \left(\sum_{i=1}^{10} \sum_{j=2k}^{2k+1} \mathbb{E} [H_i^T(j) H_i(j) | \mathcal{F}(2k-1)] \right) \geq 0.2,$$


which shows that the regression matrices are *uniformly conditionally spatio-temporally jointly*




(a) Setting I and Setting V



(b) Setting II and Setting V



(c) Setting III and Setting VI



(d) Setting IV and Setting VI

Fig. 3: Trajectories of the mean value of norms of agents' states.

observable. $\{\hat{\mathcal{G}}(k), k \geq 0\}$ is *uniformly conditionally spatio-temporally jointly connected* with

$$\inf_{k \geq 0} \lambda_2 \left(\sum_{i=2k}^{2k+1} \mathbb{E} \left[\hat{\mathcal{L}}_{\mathcal{G}(i)} | \mathcal{F}(2k-1) \right] \right) \geq 0.5.$$

Then, all the conditions of Theorem 3 hold. Fig. 4(a) shows that the estimation state of each node converges to x_0 in mean square. The performance of the algorithm is evaluated by the following maximum average regret

$$\text{MAR} = \frac{\max_{i \in V} \text{Regret}_{\text{LMS}}(i, T)}{T^{1-\tau} \ln T},$$

which is a function of T in Fig. 4(b). We can see from Fig. 4(b) that the maximum average regret tends to be a constant, which is consistent with the result of Theorem 3 with $\tau = 0.6$.




Fig. 4: Mean square errors and maximum average regret.

V. CONCLUSIONS

We study the decentralized online regularized linear regression algorithm over random time-varying graphs. At each time step, every node runs an online estimation algorithm consisting of an innovation term processing its own new measurement, a consensus term taking a weighted sum of estimations of its own and its neighbors with additive and multiplicative communication noises and a regularization term preventing over-fitting. For the generalized regression models and graphs, we obtain the sufficient conditions for the almost sure and mean square convergences of the decentralized online regularization algorithm. We prove that if the algorithm gains, the sequences of regression matrices and Laplacian matrices of the graphs jointly satisfy the *sample path spatio-temporal persistence of excitation* condition, the estimations of all nodes then converge to the unknown true parameter vector almost surely. Especially, if the graphs are *conditionally balanced* and *uniformly conditionally jointly connected*, and the regression models of all nodes are *uniformly conditionally spatio-temporally jointly observable*, then the *sample path spatio-temporal persistence of excitation* condition holds and the estimations of all nodes converge to the unknown true parameter vector almost surely and in mean square by properly choosing the algorithm gains. For the special case with Markovian switching graphs and regression matrices, we show that if the stationary graph is balanced and contains a spanning tree, and the regression model is *spatio-temporally jointly observable*, then appropriate algorithm gains can be designed such that the *sample path spatio-temporal persistence of excitation* condition

holds. To appraise the performance of the decentralized online regularized algorithm, we prove that the regret upper bound is $\mathcal{O}(T^{1-\tau}\ln T)$, where $\tau \in (0.5, 1)$ is a constant depending on the algorithm gains.

There are a number of interesting problems for further investigation. For examples, (i) how to design the appropriate regularization parameter to achieve the optimal convergence rate for decentralized online regularized algorithm; (ii) for the case that the relative-state difference $x_j(k) - x_i(k)$ goes to zero but the multiplicative term does not go to zero in communication model (5), the method in this paper may not be applicable. It is interesting to develop effective tools to analyze the decentralized online learning algorithm with more general state-dependent communication noises; (iii) it is also interesting to study the decentralized online regression problems over random time-varying graphs with time-varying unknown parameters by combining our analytical techniques with those of Xie and Guo [32]-[33]; (iv) it is possible to study the asynchronous regularized algorithm with time delays by applying the techniques of Wang *et al.* [38] and this paper.

APPENDIX A

PROOFS OF THEOREMS 1-4, PROPOSITION 1, COROLLARY 2 AND VERIFICATION OF EXAMPLE 2

Proof of Theorem 1. It follows from the condition (iii) and Lemma 1 that there exists a positive integer k_0 , such that

$$\begin{aligned} \mathbb{E}[V((k+1)h)|\mathcal{F}(kh-1)] &\leq (1 + \Omega(k))V(kh) - 2 \left(\Lambda_k^h + \sum_{i=kh}^{(k+1)h-1} \lambda(i) \right) V(kh) \\ &\quad + \Gamma(k) \text{ a.s., } k \geq k_0, \end{aligned}$$

where $\Omega(k) + \Gamma(k) = \mathcal{O}(a^2(kh) + b^2(kh) + \lambda(kh))$ is nonnegative. It follows from the condition (i) and $V(k) = \|\delta(k)\|^2$ that $\Lambda_k^h + \sum_{i=kh}^{(k+1)h-1} \lambda(i)$ and $V(kh)$ are both nonnegative and adapted to $\mathcal{F}(kh-1)$ for $k \geq k_0$. By the condition (ii), we have

$$\sum_{k=0}^{\infty} \Omega(k) + \sum_{k=0}^{\infty} \Gamma(k) < \infty,$$

which together with above and Theorem 1 in [55] leads to

$$\sum_{k=0}^{\infty} \left(\Lambda_k^h + \sum_{i=kh}^{(k+1)h-1} \lambda(i) \right) V(kh) < \infty \text{ a.s.,}$$

and $V(kh)$ converges almost surely, which together with condition (i) further leads to

$$\lim_{k \rightarrow \infty} \delta(kh) = 0 \text{ a.s.} \quad (\text{A.1})$$

Denote $m_k = \lfloor \frac{k}{h} \rfloor$. For any $\varepsilon > 0$, by Markov inequality and the conditions (ii)-(iii), we have

$$\sum_{k=0}^{\infty} \mathbb{P}\{\|b(k)\mathcal{L}_{\mathcal{G}(k)} \otimes I_n + a(k)\mathcal{H}^T(k)\mathcal{H}(k)\| \geq \varepsilon\} \leq \varepsilon^{-2} \rho_0^2 \sum_{k=0}^{\infty} (a^2(k) + b^2(k)) < \infty \text{ a.s.,}$$

which together with Borel-Cantelli Lemma gives

$$\mathbb{P}\{\|b(k)\mathcal{L}_{\mathcal{G}(k)} \otimes I_n + a(k)\mathcal{H}^T(k)\mathcal{H}(k)\| \geq \varepsilon \text{ i.o.}\} = 0,$$

i.e., $\|b(k)\mathcal{L}_{\mathcal{G}(k)} \otimes I_n + a(k)\mathcal{H}^T(k)\mathcal{H}(k)\|$ converges to zero almost surely, which gives

$$\sup_{k \geq 0} \|b(k)\mathcal{L}_{\mathcal{G}(k)} \otimes I_n + a(k)\mathcal{H}^T(k)\mathcal{H}(k)\| < \infty \text{ a.s.}$$

Noting that $0 \leq k - m_k h < h$, we have

$$\Phi_0 \triangleq \sup_{k \geq 0} \sup_{m_k h \leq i \leq k} \|\Phi_P(k-1, i)\| < \infty \text{ a.s.} \quad (\text{A.2})$$

By Lemma A.1, we know that $\|W(k)\| \leq \sqrt{N}\|\mathcal{A}_{\mathcal{G}(k)}\|$ and $\|M(k)\| \leq \sqrt{4\sigma^2 V(k) + 2b^2} \leq 2\sigma\delta(k) + \sqrt{2}b$, it follows from the estimation error equation (8) and (A.2) that

$$\|\delta(k)\| \leq f(k) + \sum_{i=m_k h}^{k-1} g(i)\|\delta(i)\|,$$

where

$$f(k) = \Phi_0(\|\delta(m_k h)\|) + \sum_{i=m_k h}^{k-1} (a(i)\|v(i)\|\|\mathcal{H}^T(i)\| + b\sqrt{2N}b(i)\|\xi(i)\|\|\mathcal{A}_{\mathcal{G}(i)}\| + \sqrt{N}\|x_0\|\lambda(i))$$

and

$$g(i) = 2\sigma\sqrt{N}\Phi_0 b(i)\|\xi(i)\|\|\mathcal{A}_{\mathcal{G}(i)}\|.$$

By Gronwall inequality, we get

$$\|\delta(k)\| \leq f(k) + \sum_{i=m_k h}^{k-1} f(i)g(i) \prod_{j=i+1}^{k-1} (1 + g(j)). \quad (\text{A.3})$$

For any $\varepsilon > 0$, it follows from Assumption (A2), the conditions (ii)-(iii), Lemma A.1 and Markov inequality that

$$\sum_{k=0}^{\infty} \mathbb{P}\{b(k)\|\xi(k)\|\|\mathcal{A}_{\mathcal{G}(k)}\| \geq \varepsilon\} \leq \varepsilon^{-2} N \beta_v \rho_0^2 \sum_{k=0}^{\infty} b^2(k) < \infty \text{ a.s.,}$$

which shows that $b(k)\|\xi(k)\|\|\mathcal{A}_{\mathcal{G}(k)}\|$ converges to zero almost surely by Borel-Cantelli Lemma. Following the same way, we know that $a(k)\|v(k)\|\|\mathcal{H}^T(k)\|$ also converges to zero almost surely, which together with (A.1)-(A.2) gives $\lim_{k \rightarrow \infty} f(k) = 0$ a.s. By (A.2), we also have $\lim_{k \rightarrow \infty} g(k) = 0$ a.s. It follows from the above and (A.3) that $\lim_{k \rightarrow \infty} \delta(k) = 0$ a.s.

Proof of Proposition 1. Let the mixing rate of $\{\mathcal{A}_{\mathcal{G}(k)}, H_i(k), v_i(k), \xi_i(k), i \in \mathcal{V}, k \geq 0\}$ be $\phi(k)$. Since $\phi(k) \rightarrow 0$ as $k \rightarrow \infty$, there exists a positive integer T such that

$$\phi(k) \leq \frac{\alpha_1}{4h_0 + 2}, \quad \forall k \geq T. \quad (\text{A.4})$$

Let $h = 2h_1 + T + 1$ and $\theta = 0.5\alpha_1$. Noting that $\mathcal{G}(k) \in \Gamma_1$ implies $\mathcal{G}(k|k-1)$ is balanced a.s. and $\mathbb{E}[\mathcal{A}_{\mathcal{G}(k)}|\mathcal{F}(k-1)] \succeq \mathbf{O}_{n \times n}$ a.s., we know that $\mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(k)}|\mathcal{F}(k-1)]$ is the Laplacian matrix of $\frac{\mathbb{E}[\mathcal{A}_{\mathcal{G}(k)} + \mathcal{A}_{\mathcal{G}(k)}^T|\mathcal{F}(k-1)]}{2}$. By the definition of Laplacian matrix, we have $x^T \mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(k)}|\mathcal{F}(k-1)]x = \frac{1}{4} \sum_{i=1}^N \sum_{j=1}^N \mathbb{E}[w_{ij}(k) + w_{ji}(k)|\mathcal{F}(k-1)](x_i - x_j)^2 \geq 0$, $\forall x = [x_1, \dots, x_N]^T \in \mathbb{R}^N$, i.e., $\mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(k)}|\mathcal{F}(k-1)] \geq \mathbf{O}_{N \times N}$, which gives

$$\lambda_2 \left(\sum_{i=kh+T}^{(k+1)h-1} \mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(i)}] \right) \geq \alpha_1, \quad \forall k \geq 0. \quad (\text{A.5})$$

By the property of ϕ -mixing process and (A.4), we get

$$\left\| \mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(kh+i)}|\mathcal{F}(kh-1)] - \mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(kh+i)}] \right\| \leq \frac{\alpha_1}{4h_1 + 2}, \quad (\text{A.6})$$

where $i \geq T$. Combining (A.4)-(A.6) gives

$$\begin{aligned} & \lambda_2 \left(\sum_{i=kh}^{(k+1)h-1} \mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(i)}|\mathcal{F}(kh-1)] \right) \\ & \geq \lambda_2 \left(\sum_{i=kh+T}^{(k+1)h-1} \mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(i)}|\mathcal{F}(kh-1)] \right) \\ & \geq \lambda_2 \left(\sum_{i=kh+T}^{(k+1)h-1} \mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(i)}] \right) \\ & \quad - \left\| \sum_{i=kh+T}^{(k+1)h-1} \mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(i)}|\mathcal{F}(kh-1)] - \sum_{i=kh+T}^{(k+1)h-1} \mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(i)}] \right\| \\ & \geq \alpha_1 - (h-T) \frac{\alpha_1}{4h_1 + 2} \\ & = \theta_1 \text{ a.s., } \forall k \geq 0. \end{aligned} \quad (\text{A.7})$$

Noting that $H_i^T(j)H_i(j) \geq \mathbf{O}_{N \times N}$ a.s., by following the same way with (A.5)-(A.7), we can similarly prove Theorem 1. (ii).

Proof of Theorem 2. It follows from the conditions (i)-(ii) and Definitions 1-2 that there exist positive constants θ_1 and θ_2 , such that

$$\inf_{k \geq 0} \lambda_2 \left(\sum_{j=kh}^{(k+1)h-1} \mathbb{E} \left[\widehat{\mathcal{L}}_{\mathcal{G}(j)} | \mathcal{F}(kh-1) \right] \right) \geq \theta_1 \text{ a.s.}$$

and

$$\lambda_{\min} \left(\sum_{i=1}^N \sum_{j=kh}^{(k+1)h-1} \mathbb{E} \left[H_i^T(j) H_i(j) | \mathcal{F}(kh-1) \right] \right) \geq \theta_2 \text{ a.s.,}$$

respectively. Denote

$$T_k = (2Nh\rho_0 + N\theta_1)^{-1} \theta_1 \theta_2 \min\{a((k+1)h), b((k+1)h)\} + \sum_{i=kh}^{(k+1)h-1} \lambda(i).$$

By Condition **(C1)**, we have

$$\sum_{k=0}^{\infty} T_k = \infty.$$

Hence, by Condition **(C1)** and (I) of Lemma A.4, we have proved (I) of Theorem 2.

Next, we will prove (II) of Theorem 2. It follows from Condition **(C2)** that

$$\sum_{k=0}^{\infty} T_k = \infty.$$

Noting that $a(k) = \mathcal{O}(a(k+1))$ and $b(k) = \mathcal{O}(b(k+1))$, we know that there exist positive constants C_1 and C_2 , such that $a(k) \leq C_1 a(k+1)$ and $b(k) \leq C_2 b(k+1)$. Denote $C = \max\{C_1, C_2\}$, we have

$$\min\{a(k), b(k)\} \leq C \min\{a(k+1), b(k+1)\},$$

which leads to

$$\begin{aligned} & \frac{\min\{a(kh), b(kh)\}}{\min\{a((k+1)h), b((k+1)h)\}} \\ &= \prod_{i=0}^{h-1} \left[\frac{\min\{a(kh+i), b(kh+i)\}}{\min\{a(kh+i+1), b(kh+i+1)\}} \right] \\ &\leq C^h, \end{aligned}$$

thus, we obtain

$$\min\{a(kh), b(kh)\} \leq C^h \min\{a((k+1)h), b((k+1)h)\},$$

which together with Condition **(C2)** gives $a^2(kh) + b^2(kh) + \lambda(kh) = o(T_k)$. It follows from (II) of Lemma A.4 that $\lim_{k \rightarrow \infty} \mathbb{E}[V(k)] = 0$.

Verification of Example 2. By Assumption (2), we have $\mathbb{E}[\mathcal{H}^T \mathcal{H} | \mathcal{F}(kh-1)] = \mathbb{E}[\mathcal{H}^T \mathcal{H} | \sigma(\mathcal{H})] = \mathcal{H}^T \mathcal{H}$ a.s. For any given positive integer h , by the definition of Laplacian matrix, we get

$$\Lambda_k^h = \sum_{i=kh}^{(k+1)h-1} \frac{x+1-\sqrt{x^2-2x+2}}{2(i+1)} \text{ a.s., } \forall k \geq 0. \quad (\text{A.8})$$

Noting that x is uniformly distributed in $(0.25, 1.25)$, we obtain $x+1-\sqrt{x^2-2x+2} > 0$ a.s., which together with (A.8) gives

$$\sum_{k=0}^{\infty} \Lambda_k^h = \left(x+1-\sqrt{x^2-2x+2} \right) \sum_{k=0}^{\infty} \sum_{i=kh}^{(k+1)h-1} \frac{1}{2(i+1)} = \infty \text{ a.s.}$$

Suppose that there exists a positive real sequence $\{c(k), k \geq 0\}$ satisfying $\Lambda_k^h \geq c(k)$ a.s. For any given integer $k_0 > 0$, denote $\mu = \frac{(k_0 h + 1)c(k_0)}{h}$. It follows from $\Lambda_k^h \geq c(k)$ a.s. that

$$0 < \mu \leq \frac{x+1-\sqrt{x^2-2x+2}}{2} < 1 \text{ a.s.,} \quad (\text{A.9})$$

which leads to $x \geq \mu + \frac{1}{4(1-\mu)}$ a.s. Hence, by (A.9), we have

$$\begin{aligned} \mathbb{P} \{ \Lambda_{k_0}^h \geq c(k_0) \} &\leq \mathbb{P} \left\{ \mu + \frac{1}{4(1-\mu)} \leq x \leq \frac{5}{4} \right\} \\ &= \frac{5}{4} - \mu - \frac{1}{4(1-\mu)} < 1, \end{aligned}$$

which is contradictory to $\Lambda_k^h \geq c(k)$ a.s.

Proof of Corollary 2. Due to the one-to-one correspondence between the weighted adjacency matrix $\mathcal{A}_{\mathcal{G}(k)}$ and the Laplacian matrix $\mathcal{L}_{\mathcal{G}(k)}$, by Assumption (A3) and the definition of uniform ergodic Markov chain, we know that $\mathcal{L}_{\mathcal{G}(k)}$ is a homogeneous and uniform ergodic Markov chain with the unique stationary distribution π . Denote the associated Laplacian matrix of \mathcal{A}_l by \mathcal{L}_l and $\hat{\mathcal{L}}_l = \frac{1}{2}(\mathcal{L}_l + \mathcal{L}_l^T)$, by the definition of $\tilde{\Lambda}_k^h$, we have

$$\begin{aligned} &\tilde{\Lambda}_k^h \\ &= \lambda_{\min} \left(\sum_{i=kh}^{(k+1)h-1} \mathbb{E} \left[\hat{\mathcal{L}}_{\mathcal{G}(i)} \otimes I_n + \mathcal{H}^T(i) \mathcal{H}(i) | \mathcal{F}(kh-1) \right] \right) \\ &= \lambda_{\min} \left(\sum_{i=kh}^{(k+1)h-1} \mathbb{E} \left[\hat{\mathcal{L}}_{\mathcal{G}(i)} \otimes I_n \right. \right. \\ &\quad \left. \left. + \mathcal{H}^T(i) \mathcal{H}(i) | \left\langle \hat{\mathcal{L}}_{\mathcal{G}(kh-1)}, \mathcal{H}(kh-1) \right\rangle = S_0 \right] \right) \\ &= \lambda_{\min} \left(\sum_{i=1}^h \sum_{l=1}^{\infty} \left(\hat{\mathcal{L}}_{\mathcal{G}_l} \otimes I_n + \mathcal{H}_l^T \mathcal{H}_l \right) \mathbb{P}^i \left(S_0, \left\langle \hat{\mathcal{L}}_{\mathcal{G}_l}, \mathcal{H}_l \right\rangle \right) \right), \\ &\quad S_0 \in \mathcal{S}, \quad k \geq 0, \quad h \geq 1. \end{aligned}$$

Noting the uniform ergodicity of $\{\hat{\mathcal{L}}_{\mathcal{G}(k)}, k \geq 0\}$ and $\{\mathcal{H}(k), k \geq 0\}$ and the uniqueness of the stationary distribution π , it follows from $\sup_{l \geq 1} \|\mathcal{L}_l\| < \infty$ and $\sup_{l \geq 1} \|\mathcal{H}(l)\| < \infty$ that

$$\begin{aligned}
& \left\| \frac{1}{h} \sum_{i=1}^h \sum_{l=1}^{\infty} \left(\hat{\mathcal{L}}_{\mathcal{G}_l} \otimes I_n + \mathcal{H}_l^T \mathcal{H}_l \right) \mathbb{P}^i \left(S_0, \langle \hat{\mathcal{L}}_{\mathcal{G}_l}, \mathcal{H}_l \rangle \right) - \sum_{l=1}^{\infty} \pi_l \left(\hat{\mathcal{L}}_{\mathcal{G}_l} \otimes I_n + \mathcal{H}_l^T \mathcal{H}_l \right) \right\| \\
&= \left\| \frac{1}{h} \sum_{i=1}^h \sum_{l=1}^{\infty} \left(\hat{\mathcal{L}}_{\mathcal{G}_l} \otimes I_n + \mathcal{H}_l^T \mathcal{H}_l \right) \mathbb{P}^i \left(S_0, \langle \hat{\mathcal{L}}_{\mathcal{G}_l}, \mathcal{H}_l \rangle \right) - \frac{1}{h} \sum_{i=1}^h \sum_{l=1}^{\infty} \pi_l \left(\hat{\mathcal{L}}_{\mathcal{G}_l} \otimes I_n + \mathcal{H}_l^T \mathcal{H}_l \right) \right\| \\
&= \left\| \frac{1}{h} \sum_{i=1}^h \sum_{l=1}^{\infty} \left(\hat{\mathcal{L}}_{\mathcal{G}_l} \otimes I_n + \mathcal{H}_l^T \mathcal{H}_l \right) \left(\mathbb{P}^i \left(S_0, \langle \hat{\mathcal{L}}_{\mathcal{G}_l}, \mathcal{H}_l \rangle \right) - \pi_l \right) \right\| \\
&\leq \left(\sup_{l \geq 1} \|\mathcal{L}_l\| + \sup_{l \geq 1} \|\mathcal{H}_l\|^2 \right) h^{-1} \sum_{i=1}^h Rr^{-i} \rightarrow 0, \quad h \rightarrow \infty,
\end{aligned}$$

where $R > 0$ and $r > 1$ are both constants. By the definition of uniform convergence, we know that

$$\begin{aligned}
& \frac{1}{h} \sum_{i=kh}^{(k+1)h-1} \mathbb{E} \left[\hat{\mathcal{L}}_{\mathcal{G}(i)} \otimes I_n + \mathcal{H}^T(i) \mathcal{H}(i) | \mathcal{F}(kh-1) \right] \\
& \rightarrow \sum_{l=1}^{\infty} \pi_l \left(\hat{\mathcal{L}}_{\mathcal{G}_l} \otimes I_n + \mathcal{H}_l^T \mathcal{H}_l \right) \text{ a.s., } h \rightarrow \infty,
\end{aligned} \tag{A.10}$$

which is uniform w.r.t. k and the sample paths a.s. For arbitrary non-zero vector $x = [x_1^T, \dots, x_N^T] \in \mathbb{R}^{Nn}$, where $x_i \in \mathbb{R}^n, i = 1, \dots, N$, if there exists a non-zero vector $a \in \mathbb{R}^n$ such that $x_1 = x_2 = \dots = x_N = a$. Noting that the regression model (1) is spatio-temporally jointly observable, we have

$$x^T \left(\sum_{l=1}^{\infty} \pi_l \left(\hat{\mathcal{L}}_{\mathcal{G}_l} \otimes I_n + \mathcal{H}_l^T \mathcal{H}_l \right) \right) x \geq a^T \left(\sum_{i=1}^N \sum_{l=1}^{\infty} \pi_l H_{i,l}^T H_{i,l} \right) a > 0, \tag{A.11}$$

otherwise, there exists $i \neq j$ such that $x_i \neq x_j$. Here, we conclude that $\sum_{l=1}^{\infty} \pi_l \hat{\mathcal{L}}_l$ is the Laplacian matrix of a connected graph since the stationary weighted adjacency matrix $\sum_{l=1}^{\infty} \pi_l \mathcal{A}_l$ is nonnegative and its associated graph is balanced with a spanning tree. Then, by Lemma A.7 in [38], we have

$$x^T \left(\sum_{l=1}^{\infty} \pi_l \left(\hat{\mathcal{L}}_{\mathcal{G}_l} \otimes I_n + \mathcal{H}_l^T \mathcal{H}_l \right) \right) x \geq x^T \left(\sum_{l=1}^{\infty} \pi_l \left(\hat{\mathcal{L}}_{\mathcal{G}_l} \otimes I_n \right) \right) x > 0. \tag{A.12}$$

Hence, Combining (A.11) and (A.12), we get

$$\lambda_{\min} \left(\sum_{l=1}^{\infty} \pi_l \left(\hat{\mathcal{L}}_{\mathcal{G}_l} \otimes I_n + \mathcal{H}_l^T \mathcal{H}_l \right) \right) > 0.$$

One one hand, since the function $\lambda_{\min}(\cdot)$, whose arguments are matrices, is continuous, there exists a constant $\delta > 0$ such that for any matrix L satisfying $\|L - (\sum_{l=1}^{\infty} \pi_l (\hat{\mathcal{L}}_{\mathcal{G}_l} \otimes I_n + \mathcal{H}_l^T \mathcal{H}_l))\| \leq \delta$,

we have $\|\lambda_{\min}(L) - \lambda_{\min}(\sum_{l=1}^{\infty} \pi_l(\widehat{\mathcal{L}}_{\mathcal{G}_l} \otimes I_n + \mathcal{H}_l^T \mathcal{H}_l))\| \leq \frac{1}{2} \lambda_{\min}(\sum_{l=1}^{\infty} \pi_l(\widehat{\mathcal{L}}_{\mathcal{G}_l} \otimes I_n + \mathcal{H}_l^T \mathcal{H}_l))$.

One the other hand, by (A.10), there exists a positive integer h_0 , such that

$$\sup_{k \geq 0} \left\| \frac{1}{h} \sum_{i=kh}^{(k+1)h-1} \mathbb{E} \left[\widehat{\mathcal{L}}_{\mathcal{G}(i)} \otimes I_n + \mathcal{H}^T(i) \mathcal{H}(i) | \mathcal{F}(kh-1) \right] - \sum_{l=1}^{\infty} \pi_l \left(\widehat{\mathcal{L}}_{\mathcal{G}_l} \otimes I_n + \mathcal{H}_l^T \mathcal{H}_l \right) \right\| \leq \delta, \quad h \geq h_0,$$

which gives

$$\begin{aligned} & \sup_{k \geq 0} \left\| \frac{1}{h} \widetilde{\Lambda}_k^h - \lambda_{\min} \left(\sum_{l=1}^{\infty} \pi_l \left(\widehat{\mathcal{L}}_{\mathcal{G}_l} \otimes I_n + \mathcal{H}_l^T \mathcal{H}_l \right) \right) \right\| \\ & \leq \frac{1}{2} \lambda_{\min} \left(\sum_{l=1}^{\infty} \pi_l \left(\widehat{\mathcal{L}}_{\mathcal{G}_l} \otimes I_n + \mathcal{H}_l^T \mathcal{H}_l \right) \right), \quad h \geq h_0. \end{aligned}$$

Then we arrive at

$$\widetilde{\Lambda}_k^h \geq \frac{1}{2} \lambda_{\min} \left(\sum_{l=1}^{\infty} \pi_l \left(\widehat{\mathcal{L}}_{\mathcal{G}_l} \otimes I_n + \mathcal{H}_l^T \mathcal{H}_l \right) \right) \text{ a.s., } h \geq h_0,$$

thus, the proof of Corollary 2 is completed due to Lemma A.5.

Proof of Theorem 3. It follows from $\{\mathcal{G}(k), k \geq 0\} \in \Gamma_1$ that $\mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(k)} | \mathcal{F}(k-1)]$ is positive semi-definite. Noting that

$$\mathbb{E} \left[\widehat{\mathcal{L}}_{\mathcal{G}(k)} | \mathcal{F}(mh-1) \right] = \mathbb{E} \left[\mathbb{E} \left[\widehat{\mathcal{L}}_{\mathcal{G}(k)} | \mathcal{F}(k-1) \right] | \mathcal{F}(mh-1) \right], \quad k \geq mh,$$

we know that $\mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(k)} | \mathcal{F}(mh-1)]$ is also positive semi-definite, which shows that

$$\begin{aligned} & \sum_{i=kh}^{(k+1)h-1} \mathbb{E} \left[b(i) \widehat{\mathcal{L}}_{\mathcal{G}(i)} \otimes I_n + a(i) \mathcal{H}^T(i) \mathcal{H}(i) | \mathcal{F}(kh-1) \right] \\ & \geq \min\{a((k+1)h), b((k+1)h)\} \sum_{i=kh}^{(k+1)h-1} \mathbb{E} \left[\widehat{\mathcal{L}}_{\mathcal{G}(i)} \otimes I_n + \mathcal{H}^T(i) \mathcal{H}(i) | \mathcal{F}(kh-1) \right] \text{ a.s.} \end{aligned}$$

It follows from the conditions (i)-(ii) and Definitions 1-2 that there exist positive constants θ_1 and θ_2 , such that

$$\inf_{k \geq 0} \lambda_2 \left(\sum_{j=kh}^{(k+1)h-1} \mathbb{E} \left[\widehat{\mathcal{L}}_{\mathcal{G}(j)} | \mathcal{F}(kh-1) \right] \right) \geq \theta_1 \text{ a.s.}$$

and

$$\lambda_{\min} \left(\sum_{i=1}^N \sum_{j=kh}^{(k+1)h-1} \mathbb{E} \left[H_i^T(j) H_i(j) | \mathcal{F}(kh-1) \right] \right) \geq \theta_2 \text{ a.s.,}$$

respectively, which together with condition (iii) and Lemma 2 gives

$$\Lambda_k^h \geq (2Nh\rho_0 + N\theta_1)^{-1} \theta_1 \theta_2 \min\{a((k+1)h), b((k+1)h)\} \text{ a.s.}$$

Denote

$$L(k) = (2Nh\rho_0 + N\theta_1)^{-1}\theta_1\theta_2 \min\{a((k+1)h), b((k+1)h)\} + \sum_{i=kh}^{(k+1)h-1} \lambda(i),$$

we have

$$\Lambda_k^h + \sum_{i=kh}^{(k+1)h-1} \lambda(i) \geq L(k) \text{ a.s.,}$$

which together with Lemma 1 shows that there exists a positive integer k_0 , such that

$$\mathbb{E}[V((k+1)h)|\mathcal{F}(kh-1)] \leq (1 + \Omega(k))V(kh) - 2L(k)V(kh) + \Gamma(k) \text{ a.s., } k \geq k_0, \text{ (A.13)}$$

where $\Omega(k) + \Gamma(k) = \mathcal{O}(a^2(kh) + b^2(kh) + \lambda(kh))$, which together with the choice of the algorithm gains shows that there exists a positive constant u such that

$$\Gamma(k) \leq u(kh+1)^{-2\tau}, \quad k \geq 0.$$

Noting that $\Omega(k) = o(L(k))$ and $L(k) = o(1)$, without loss of generality, we suppose that $0 < \Omega(k) \leq L(k) < 1$, $k \geq k_0$. Denote

$$v = (4Nh\rho_0 + 2N\theta_1)^{-1}c_1\theta_1\theta_2,$$

it follows that

$$L(k) \geq v(kh+h+1)^{-\tau}, \quad k \geq k_0.$$

By taking mathematical expectation on both sides of (A.13), we get

$$\begin{aligned} & \mathbb{E}[V((k+1)h)] \\ & \leq \prod_{i=k_0+1}^k \left(1 - \frac{v}{(ih+h+1)^\tau}\right) \sup_{k \geq 0} \mathbb{E}[V(k)] \\ & + \sum_{i=k_0}^k \frac{u}{(ih+1)^{2\tau}} \prod_{j=i+1}^k \left(1 - \frac{v}{(jh+h+1)^\tau}\right), \quad k > k_0. \end{aligned} \quad \text{(A.14)}$$

Noting that

$$\begin{aligned} & \sum_{j=i+1}^k \frac{v}{(jh+h+1)^\tau} \\ & \geq \int_{i+1}^k \frac{v}{(xh+h+1)^\tau} dx \\ & = \frac{v}{h(1-\tau)} ((kh+h+1)^{1-\tau} - (ih+2h+1)^{1-\tau}), \quad i \geq 0, \end{aligned}$$

we have

$$\prod_{j=i+1}^k \left(1 - \frac{v}{(jh+h+1)^\tau}\right)$$

$$\leq \exp \left(-\frac{v}{h(1-\tau)} ((kh+h+1)^{1-\tau} - (ih+2h+1)^{1-\tau}) \right), \quad k_0 \leq i \leq k. \quad (\text{A.15})$$

By Theorem 2, we have $\lim_{k \rightarrow \infty} \mathbb{E}[V(k)] = 0$, which shows that

$$\sup_{k \geq 0} \mathbb{E}[V(k)] < \infty.$$

It follows from (A.15) that

$$\exp \left(-\frac{v}{h(1-\tau)} (kh+h+1)^{1-\tau} \right) = o \left((kh+h)^{-\tau} \ln(kh+h) \right),$$

thus, we have

$$\prod_{i=k_0+1}^k \left(1 - \frac{v}{(ih+h+1)^\tau} \right) \sup_{k \geq 0} \mathbb{E}[V(k)] = o \left((kh+h)^{-\tau} \ln(kh+h) \right). \quad (\text{A.16})$$

Denote

$$\epsilon(k) = \left\lceil \frac{2}{v} (kh+h+1)^\tau \ln(kh+h+1) \right\rceil,$$

then we have $\epsilon(k) = o(k)$. Noting that $\epsilon(k) \rightarrow \infty$ as $k \rightarrow \infty$, without loss of generality, we suppose that $k_0 < \epsilon(k) \leq 2\epsilon(k) \leq k$. On one hand, for the case with $k_0 \leq i \leq k-1-\epsilon(k)$, we know that

$$ih+2h+1 \leq kh+h+1-\epsilon_1(k), \quad (\text{A.17})$$

where $\epsilon_1(k) = \lceil \frac{2h}{v} (kh+h+1)^\tau \ln(kh+h+1) \rceil$. Noting that

$$(1-x)^\alpha \leq 1 - \alpha x, \quad 0 \leq x, \alpha \leq 1,$$

we obtain

$$\left(\frac{kh+h+1-\epsilon_1(k)}{kh+h+1} \right)^{1-\tau} \leq 1 - \frac{\epsilon_1(k)(1-\tau)}{kh+h+1},$$

which directly gives

$$(kh+h+1)^{1-\tau} - (kh+h+1-\epsilon_1(k))^{1-\tau} \geq (kh+h+1)^{-\tau} \epsilon_1(k)(1-\tau) \geq v^{-1} 2h(1-\tau) \ln(kh+h+1).$$

This together with (A.17) gives

$$(h(1-\tau))^{-1} v \left((kh+h+1)^{1-\tau} - (ih+2h+1)^{1-\tau} \right) \geq 2 \ln(kh+h+1).$$

Then, it follows from (A.15) that

$$\prod_{j=i+1}^k \left(1 - \frac{v}{(jh+h+1)^\tau} \right) \leq \frac{1}{(kh+h+1)^2}, \quad k_0 \leq i \leq k-1-\epsilon(k),$$

from which we know that

$$\sum_{i=k_0}^{k-1-\epsilon(k)} \frac{u}{(ih+1)^{2\tau}} \prod_{j=i+1}^k \left(1 - \frac{v}{(jh+h+1)^\tau}\right) = \mathcal{O}(k^{-1}). \quad (\text{A.18})$$

On the other hand, for the case with $k - \epsilon(k) \leq i \leq k$, we have $k \leq 2k - 2\epsilon(k) \leq 2i$, which shows that

$$\frac{u}{(ih+1)^{2\tau}} \leq \frac{4^\tau u}{(kh+2)^{2\tau}}, \quad k - \epsilon(k) \leq i \leq k.$$

Then it follows from

$$\prod_{j=i+1}^k \left(1 - \frac{v}{(jh+h+1)^\tau}\right) \leq 1$$

that

$$\sum_{i=k-\epsilon(k)}^k \frac{u}{(ih+1)^{2\tau}} \prod_{j=i+1}^k \left(1 - \frac{v}{(jh+h+1)^\tau}\right) = \mathcal{O}((kh+h)^{-\tau} \ln(kh+h)). \quad (\text{A.19})$$

Thus, by (A.18)-(A.19), we obtain

$$\sum_{i=k_0}^k \frac{u}{(ih+1)^{2\tau}} \prod_{j=i+1}^k \left(1 - \frac{v}{(jh+h+1)^\tau}\right) = \mathcal{O}((kh+h)^{-\tau} \ln(kh+h)). \quad (\text{A.20})$$

Combining (A.14), (A.16) and (A.20) directly shows that

$$\mathbb{E}[V(kh)] = \mathcal{O}((kh+h)^{-\tau} \ln(kh+h)).$$

By Lemma A.2, there exists a positive integer k_1 such that

$$\mathbb{E}[V(k)] \leq 2^h \mathbb{E}[V(m_k h)] + h 2^h \gamma(m_k h), \quad k \geq k_1,$$

where $\gamma(k) = \mathcal{O}((k+1)^{-2\tau})$ and $m_k = \lfloor \frac{k}{h} \rfloor$. Thus, we get

$$\sum_{t=0}^T \mathbb{E}[V(t)] = \mathcal{O} \left(\sum_{t=0}^T (t+1)^{-\tau} \ln(t+1) \right).$$

Noting that

$$\sum_{t=0}^T (t+1)^{-\tau} \ln(t+1) = \mathcal{O} \left(\int_0^T (x+1)^{-\tau} \ln(x+1) dx \right),$$

by Lemma 3, we get (10).

Proof of Theorem 4. It follows from $\{\mathcal{G}(k), k \geq 0\} \in \Gamma_1$ that $\mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(k)} | \mathcal{F}(k-1)]$ is positive semi-definite. Noting that

$$\mathbb{E} \left[\widehat{\mathcal{L}}_{\mathcal{G}(k)} | \mathcal{F}(mh-1) \right] = \mathbb{E} \left[\mathbb{E} \left[\widehat{\mathcal{L}}_{\mathcal{G}(k)} | \mathcal{F}(k-1) \right] | \mathcal{F}(mh-1) \right], \quad k \geq mh,$$

we know that $\mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(k)}|\mathcal{F}(mh-1)]$ is also positive semi-definite, which shows that

$$\begin{aligned} & \sum_{i=kh}^{(k+1)h-1} \mathbb{E} \left[b(i) \widehat{\mathcal{L}}_{\mathcal{G}(i)} \otimes I_n + a(i) \mathcal{H}^T(i) \mathcal{H}(i) | \mathcal{F}(kh-1) \right] \\ & \geq \min\{a((k+1)h), b((k+1)h)\} \sum_{i=kh}^{(k+1)h-1} \mathbb{E} \left[\widehat{\mathcal{L}}_{\mathcal{G}(i)} \otimes I_n + \mathcal{H}^T(i) \mathcal{H}(i) | \mathcal{F}(kh-1) \right] \text{ a.s.} \end{aligned}$$

It follows from the conditions (i)-(ii) and Definitions 1-2 that there exist positive constants θ_1 and θ_2 , such that

$$\inf_{k \geq 0} \lambda_2 \left(\sum_{j=kh}^{(k+1)h-1} \mathbb{E} \left[\widehat{\mathcal{L}}_{\mathcal{G}(j)} | \mathcal{F}(kh-1) \right] \right) \geq \theta_1 \text{ a.s.}$$

and

$$\lambda_{\min} \left(\sum_{i=1}^N \sum_{j=kh}^{(k+1)h-1} \mathbb{E} \left[H_i^T(j) H_i(j) | \mathcal{F}(kh-1) \right] \right) \geq \theta_2 \text{ a.s.,}$$

respectively, which together with condition (iii) and Lemma 2 gives

$$\Lambda_k^h \geq (2Nh\rho_0 + N\theta_1)^{-1} \theta_1 \theta_2 \min\{a((k+1)h), b((k+1)h)\} \text{ a.s.}$$

Denote

$$L(k) = (2Nh\rho_0 + N\theta_1)^{-1} \theta_1 \theta_2 \min\{a((k+1)h), b((k+1)h)\} + \sum_{i=kh}^{(k+1)h-1} \lambda(i),$$

we have

$$\Lambda_k^h + \sum_{i=kh}^{(k+1)h-1} \lambda(i) \geq L(k) \text{ a.s.,} \quad (\text{A.21})$$

By the proof of Lemma 1, we have

$$\begin{aligned} \Omega(k) &= (h+1)(4^h - 1 - 2h)(\rho_0 a(kh) + \lambda(kh))^2 + 2^{h+3} h N^2 \beta_v \sigma^2 \rho_0^2 a^2(kh) \\ &\quad + 2^{h+5} h N^2 \beta_v \sigma^2 \rho_0^2 a^2(kh) + \sum_{i=kh}^{(k+1)h-1} \lambda(i) \\ &\leq \frac{(4^h - 2h - 1)(h+1)(\rho_0 + 1)^2 + 2^{h+3} h N^2 \beta_v \sigma^2 \rho_0^2 + 2^{h+5} h N^2 \beta_v \sigma^2 \rho_0^2 + h}{(kh+1)^2} \\ &\triangleq \frac{c_1}{(kh+1)^2}, \quad \forall k \geq 0, \end{aligned}$$

Noting that

$$L(k) \geq \frac{\theta_1 \theta_2}{(2Nh\rho_0 + N\theta_1)((k+1)h+1)^\tau} \triangleq \frac{c_2}{((k+1)h+1)^\tau}, \quad \forall k \geq 0,$$

and

$$L(k) \leq \frac{\theta_1 \theta_2}{(2Nh\rho_0 + N\theta_1)(kh+1)^\tau} + \frac{h}{(kh+1)^\tau} \triangleq \frac{c_3}{(kh+1)^\tau}, \quad \forall k \geq 0,$$

Denote

$$k_0 = \left\lfloor \frac{2c_1}{hc_2} + c_3^2 \right\rfloor + 16h^2(\rho_0 + 1)^2 + 1.$$

We can verify that $0 < \Omega(k) \leq L(k) < 1$, $k \geq k_0$. By (A.21) and the proof of Lemma 1, we obtain

$$\mathbb{E}[V((k+1)h) | \mathcal{F}(kh-1)] \leq (1 + \Omega(k))V(kh) - 2L(k)V(kh) + \Gamma(k) \text{ a.s., } k \geq k_0, \quad (\text{A.22})$$

where

$$\begin{aligned} \Gamma(k) &= 2h\rho_0\beta_v a^2(kh) + (2^{h+3}h\sigma^2 + 4b^2)hN^2\beta_v\rho_0^2b^2(kh) + 8ha(kh)b(kh)(\beta_v\rho_0 \\ &\quad + N^2\beta_v\rho_0^2(2^{h+2}h\sigma^2 + 2b^2)) + 2hN\|x_0\|^2\lambda(kh) + 2Nh^2\|x_0\|^2\lambda^2(kh) \\ &\leq 2h\rho_0\beta_v a^2(kh) + (2^{h+3}h\sigma^2 + 4b^2)hN^2\beta_v\rho_0^2a^2(kh) + 8ha^2(kh)(\beta_v\rho_0 \\ &\quad + N^2\beta_v\rho_0^2(2^{h+2}h\sigma^2 + 2b^2)) + 2hN\|x_0\|^2a^2(kh) + 2Nh^2\|x_0\|^2a^2(kh) \\ &\triangleq \frac{c_4}{(kh+1)^{2\tau}}, \quad \forall k \geq 0. \end{aligned}$$

Denote

$$c_5 = (4Nh\rho_0 + 2N\theta_1)^{-1}c_1\theta_1\theta_2,$$

it follows that

$$L(k) \geq c_5(kh + h + 1)^{-\tau}, \quad k \geq k_0.$$

By taking mathematical expectation on both sides of (A.22), we get

$$\mathbb{E}[V((k+1)h)] \leq \left(1 - \frac{c_5}{(kh + h + 1)^\tau}\right) \mathbb{E}[V(kh)] + \frac{c_4}{(kh + 1)^{2\tau}}, \quad \forall k > k_0, \quad (\text{A.23})$$

which together with Lemma 2.9 in [53] gives

$$\mathbb{E}[V(kh)] \leq \frac{25c_4 \ln k}{c_5 k^{1-\tau}} + N\|x_0\|^2 e^{-\frac{c_5(k^{1-\tau}-2)}{1-\tau}}, \quad \forall k \geq \left(\frac{12}{c_5(1-\tau)} \ln \left(\frac{4}{c_5(1-\tau)}\right)\right)^{\frac{1}{1-\tau}} + k_0.$$

Denote $m_k = \lfloor \frac{k}{h} \rfloor$, $\forall k \geq 0$. Noting that $m_k h \leq k < (m_k + 1)h$, then for

$$k \geq \left(\frac{12}{c_5(1-\tau)} \ln \left(\frac{4}{c_5(1-\tau)}\right)\right)^{\frac{1}{1-\tau}} + \left\lfloor \frac{2c_1}{hc_2} + c_3^2 \right\rfloor + 16h^2(\rho_0 + 1)^2 + 1,$$

it follows from (A.23) and Lemma A.2 that

$$\begin{aligned} \mathbb{E}[V(k)] &\leq 2^h \mathbb{E}[V(m_k h)] + h2^h \gamma(m_k h) \\ &\leq 2^h \left(\frac{25c_4 \ln m_k}{c_5 m_k^{1-\tau}} + N\|x_0\|^2 e^{-\frac{c_5(m_k^{1-\tau}-2)}{1-\tau}} \right) + h2^h \gamma(m_k h) \\ &= 2^h \left(\frac{25c_4 \ln m_k}{c_5 m_k^{1-\tau}} + N\|x_0\|^2 e^{-\frac{c_5(m_k^{1-\tau}-2)}{1-\tau}} \right) + h2^h (4b^2 a^2(m_k h) N^2 \rho_0^2 \beta_v \end{aligned}$$

$$\begin{aligned}
& +2a^2(m_k h)\rho_0\beta_v + (N\|x_0\|^2 + a^2(m_k h)\rho_0^2)\lambda(m_k h) + N\|x_0\|^2\lambda^2(m_k h)) \\
\leq & 2^h \left(\frac{25c_4 \ln m_k}{c_5 m_k^{1-\tau}} + N\|x_0\|^2 e^{-\frac{c_5(m_k^{1-\tau}-2)}{1-\tau}} \right) + h2^h (4b^2 a^2(m_k h) N^2 \rho_0^2 \beta_v \\
& + 2a^2(m_k h)\rho_0\beta_v + (N\|x_0\|^2 + \rho_0^2)a^2(m_k h) + N\|x_0\|^2 a^2(m_k h)) \\
\leq & 2^h \left(\frac{25c_4 \ln k}{c_5 \left(\frac{k}{2h}\right)^{1-\tau}} + N\|x_0\|^2 e^{-\frac{c_5\left(\left(\frac{k}{2h}\right)^{1-\tau}-2\right)}{1-\tau}} \right) + h2^h \left(4b^2 a^2 \left(\frac{k}{2}\right) N^2 \rho_0^2 \beta_v \right. \\
& \left. + 2a^2 \left(\frac{k}{2}\right) \rho_0 \beta_v + (N\|x_0\|^2 + \rho_0^2)a^2 \left(\frac{k}{2}\right) + N\|x_0\|^2 a^2 \left(\frac{k}{2}\right) \right) \\
\triangleq & 2^h \left(\frac{25c_4 (2h)^{1-\tau} \ln k}{c_5 k^{1-\tau}} + N\|x_0\|^2 e^{-\frac{c_5\left(\left(\frac{k}{2h}\right)^{1-\tau}-2\right)}{1-\tau}} \right) + \frac{c_6}{(k+2)^2}.
\end{aligned}$$

APPENDIX B

PROOFS OF LEMMAS 1-3

Proof of Lemma 1. By the estimation error equation (8), we have

$$\mathbb{E}[V((k+1)h)|\mathcal{F}(kh-1)] = \sum_{i=1}^4 \mathbb{E} \left[A_i^T(k)A_i(k) + 2 \sum_{1 \leq i < j \leq 4} A_i^T(k)A_j(k) | \mathcal{F}(kh-1) \right], \quad (\text{B.1})$$

where $A_1(k) = \Phi_P((k+1)h-1, kh)\delta(kh)$, $A_2(k) = \sum_{i=kh}^{(k+1)h-1} a(i)\Phi_P((k+1)h-1, i+1)\mathcal{H}^T(i)v(i)$, $A_3(k) = \sum_{i=kh}^{(k+1)h-1} b(i)\Phi_P((k+1)h-1, i+1)W(i)M(i)\xi(i)$, and $A_4(k) = \sum_{i=kh}^{(k+1)h-1} \lambda(i)\Phi_P((k+1)h-1, i+1)(\mathbf{1}_N \otimes x_0)$.

We now consider the RHS of (B.1) term by term. It follows from Assumption (A2) that $\Phi_P^T((k+1)h-1, kh)\Phi_P((k+1)h-1, i+1)\mathcal{H}^T(i)$ and $v(i)$ are independent, $kh \leq i \leq (k+1)h-1$, which further shows that $\Phi_P^T((k+1)h-1, kh)\Phi_P((k+1)h-1, i+1)\mathcal{H}^T(i)$ and $v(i)$ are conditionally independent w.r.t. $\mathcal{F}(kh-1)$ by Lemma A.1 in [39]. Noting that $\delta(kh) \in \mathcal{F}(kh-1)$, we have

$$\mathbb{E} [A_1^T(k)A_2(k)|\mathcal{F}(kh-1)] = 0. \quad (\text{B.2})$$

Similarly, we also have

$$\mathbb{E} [A_1^T(k)A_3(k) + A_2^T(k)A_4(k) + A_3^T(k)A_4(k)|\mathcal{F}(kh-1)] = 0. \quad (\text{B.3})$$

By Lemma A.3, there exist positive integers k_2 and k_3 , such that

$$\begin{aligned}
& \mathbb{E} [A_1^T(k)A_4(k)|\mathcal{F}(kh-1)] \\
\leq & \frac{1}{2} \left(\sum_{i=kh}^{(k+1)h-1} \lambda(i) \right) (1 + p(k))V(kh) + Nh\|x_0\|^2\lambda(kh) \text{ a.s., } k \geq \max\{k_2, k_3\}, \quad (\text{B.4})
\end{aligned}$$

where $p(k) = (4^h - 1 - 2h)(\rho_0 \max\{a(kh), b(kh)\} + \lambda(kh))^2$. By Assumption **(A2)** and Lemma A.1 in [39], we know that

$$\begin{aligned} & \mathbb{E}[v^T(i)\mathcal{H}(i)\Phi_P^T((k+1)h-1, i+1) \\ & \times \Phi_P((k+1)h-1, j+1)W(j)M(j)\xi(j)|\mathcal{F}(kh-1)] \\ & = 0, \quad kh \leq j < i \leq (k+1)h-1, \end{aligned} \quad (\text{B.5})$$

and

$$\begin{aligned} & \mathbb{E}[v^T(i)\mathcal{H}(i)\Phi_P^T((k+1)h-1, i+1) \\ & \times \Phi_P((k+1)h-1, j+1)W(j)M(j)\xi(j)|\mathcal{F}(kh-1)] \\ & = 0, \quad kh \leq i < j \leq (k+1)h-1. \end{aligned} \quad (\text{B.6})$$

Therefore, by (B.5)-(B.6), Assumptions **(A1)**-**(A2)**, Lemmas A.1-A.3 and Lemma A.1 in [39], there exists a positive integer k_1 , such that

$$\begin{aligned} & \mathbb{E}[A_2^T(k)A_3(k)|\mathcal{F}(kh-1)] \\ & \leq 4 \sum_{i=kh}^{(k+1)h-1} a(i)b(i)\mathbb{E}[\|\mathcal{H}(i)\|^2\|v(i)\|^2 + \|W(i)\|^2\|M(i)\|^2\|\xi(i)\|^2|\mathcal{F}(kh-1)] \\ & \leq 4h(\beta_v\rho_0 + N^2\beta_v\rho_0^2(2^{h+2}\sigma^2V(kh) + 2^{h+2}h\sigma^2 \\ & + 2b^2))a(kh)b(kh) \text{ a.s., } k \geq \max\{k_1, k_3\}. \end{aligned} \quad (\text{B.7})$$

Similar to (B.5)-(B.6), we get

$$\begin{aligned} & \mathbb{E}[v^T(i)\mathcal{H}(i)\Phi_P^T((k+1)h-1, i+1) \\ & \times \Phi_P((k+1)h-1, j+1)\mathcal{H}^T(j)v(j)|\mathcal{F}(kh-1)] \\ & = 0, \quad kh \leq j \neq i \leq (k+1)h-1, \end{aligned}$$

which together with Assumption **(A2)**, Lemma A.3 and Lemma A.1 in [39] gives

$$\begin{aligned} & \mathbb{E}[A_2^T(k)A_2(k)|\mathcal{F}(kh-1)] \\ & \leq 2 \sum_{i=kh}^{(k+1)h-1} a^2(i)\mathbb{E}[\|\mathcal{H}^T(i)\|^2\|v(i)\|^2|\mathcal{F}(kh-1)] \\ & \leq 2h\beta_v\rho_0a^2(kh) \text{ a.s., } k \geq k_3. \end{aligned} \quad (\text{B.8})$$

Following the same way as (B.5)-(B.6), we have

$$\begin{aligned} & \mathbb{E}[\xi^T(i)M^T(i)W^T(i)\Phi_P^T((k+1)h-1, i+1) \\ & \times \Phi_P((k+1)h-1, j+1)W(j)M(j)\xi(j)|\mathcal{F}(kh-1)] \\ & = 0, \quad kh \leq j \neq i \leq (k+1)h-1, \end{aligned}$$

which along with Assumption (A2), Lemmas A.1-A.3 and Lemma A.1 in [39] leads to

$$\begin{aligned} & \mathbb{E}[A_3^T(k)A_3(k)|\mathcal{F}(kh-1)] \\ & \leq 2 \sum_{i=kh}^{(k+1)h-1} b^2(i) \mathbb{E}[\|W(i)\|^2 \|M(i)\|^2 \|\xi(i)\|^2 | \mathcal{F}(kh-1)] \\ & \leq 2hN^2\beta_v\rho_0^2(2^{h+2}\sigma^2V(kh) + 2^{h+2}h\sigma^2 + 2b^2) b^2(kh) \text{ a.s., } k \geq \max\{k_1, k_3\}. \end{aligned} \quad (\text{B.9})$$

By Lemma A.3, we know that

$$\mathbb{E}[A_4^T(k)A_4(k)|\mathcal{F}(kh-1)] \leq 2Nh^2\|x_0\|^2\lambda^2(kh) \text{ a.s.,} \quad (\text{B.10})$$

for $k \geq k_3$. By Lemma A.3, we have

$$\mathbb{E}[A_1^T(k)A_1(k)|\mathcal{F}(kh-1)] \leq \left(1 - 2\Lambda_k^h - 2 \sum_{i=kh}^{(k+1)h-1} \lambda(i) + p(k)\right) V(kh) \text{ a.s.,} \quad (\text{B.11})$$

for $k \geq k_2$. Noting that $\lambda(k)$ converges to zero, we know that there exists a positive integer k_4 , such that $\lambda(k) \leq 1$, $k \geq k_4$. Substituting (B.2)-(B.4) and (B.7)-(B.11) into (B.1) leads to

$$\begin{aligned} & \mathbb{E}[V((k+1)h)|\mathcal{F}(kh-1)] \\ & \leq (1 + \Omega(k)) V(kh) - \left(2\Lambda_k^h + 2 \sum_{i=kh}^{(k+1)h-1} \lambda(i)\right) V(kh) + \Gamma(k) \text{ a.s., } k \geq k_0, \end{aligned}$$

where $k_0 = \max\{k_1, k_2, k_3, k_4\}$, $\Omega(k) = p(k) + 2^{h+3}hN^2\beta_v\sigma^2\rho_0^2b^2(kh) + 2^{h+5}hN^2\beta_v\sigma^2\rho_0^2a(kh)b(kh) + hp(k) + \sum_{i=kh}^{(k+1)h-1} \lambda(i)$ and $\Gamma(k) = 2h\rho_0\beta_v a^2(kh) + (2^{h+3}h\sigma^2 + 4b^2)hN^2\beta_v\rho_0^2b^2(kh) + 8ha(kh)b(kh)(\beta_v\rho_0 + N^2\beta_v\rho_0^2(2^{h+2}h\sigma^2 + 2b^2)) + 2hN\|x_0\|^2\lambda(kh) + 2Nh^2\|x_0\|^2\lambda^2(kh)$. Noting that $p(k) = \mathcal{O}(a^2(kh) + b^2(kh) + \lambda^2(kh))$, we get $\Omega(k) + \Gamma(k) = \mathcal{O}(a^2(kh) + b^2(kh) + \lambda(kh))$.

Proof of Lemma 2. It follows from $\mathcal{L}_{\mathcal{G}(k)} = \mathcal{D}_{\mathcal{G}(k)} - \mathcal{A}_{\mathcal{G}(k)}$ that $\mathbf{1}_N$ is the eigenvector corresponding to the zero eigenvalue of $\mathcal{L}_{\mathcal{G}(k)}$. $\mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(k)}|\mathcal{F}(k-1)]$ has a unique zero eigenvalue with eigenvector $\mathbf{1}_N$ since $\mathcal{G}(k)$ is conditionally balanced. Therefore, $\mathbb{E}[\sum_{i=kh}^{(k+1)h-1} \widehat{\mathcal{L}}_{\mathcal{G}(i)} \otimes I_n | \mathcal{F}(kh-1)]$ has only n zero eigenvalues, whose corresponding eigenvectors consist of $s_1(k) = \frac{1}{\sqrt{N}}\mathbf{1}_N \otimes e_1, \dots, s_n(k) = \frac{1}{\sqrt{N}}\mathbf{1}_N \otimes e_n$, where e_i is the standard orthonormal basis in \mathbb{R}^n . Suppose that the remaining eigenvalues of $\mathbb{E}[\sum_{i=kh}^{(k+1)h-1} \widehat{\mathcal{L}}_{\mathcal{G}(i)} \otimes I_n | \mathcal{F}(kh-1)]$ are $\gamma_{n+1}(k), \dots, \gamma_{Nn}(k)$, and the corresponding unit orthogonal eigenvectors are $s_{n+1}(k), \dots, s_{Nn}(k)$. Then, given a positive integer k , for any unit vector $\eta \in \mathbb{R}^{Nn}$, there exist $r_i(k) \in \mathbb{R}, 1 \leq i \leq Nn$, such that

$$\eta = \eta_1(k) + \eta_2(k),$$

where $\eta_1(k) = \sum_{i=1}^n r_i(k)s_i(k)$, $\eta_2(k) = \sum_{i=n+1}^{Nn} r_i(k)s_i(k)$, and $\sum_{i=1}^{Nn} r_i^2(k) = 1$. Denote

$$H_k = \mathbf{diag}(H_{k,1}, \dots, H_{k,N}),$$

where $H_{k,i} = \sum_{j=kh}^{(k+1)h-1} \mathbb{E}[H_i^T(j)H_i(j)|\mathcal{F}(kh-1)]$, $\mathcal{L}_k = \sum_{i=kh}^{(k+1)h-1} \mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(i)}|\mathcal{F}(kh-1)]$, $u_{st}(k) = \eta_s^T(k)H_k\eta_t(k)$ and $\tilde{w}_{st}(k) = \eta_s^T(k)(\mathcal{L}_k \otimes I_n)\eta_t(k)$, $s, t = 1, 2$. It follows that

$$\begin{aligned} & \eta^T \left(\sum_{i=kh}^{(k+1)h-1} \mathbb{E} \left[\widehat{\mathcal{L}}_{\mathcal{G}(i)} \otimes I_n + \mathcal{H}^T(i)\mathcal{H}(i)|\mathcal{F}(kh-1) \right] \right) \eta \\ &= (\eta_1^T(k) + \eta_2^T(k)) (H_k + \mathcal{L}_k \otimes I_n) (\eta_1(k) + \eta_2(k)) \\ &= u_{11}(k) + u_{22}(k) + 2u_{12}(k) + \tilde{w}_{11}(k) + \tilde{w}_{22}(k) + 2\tilde{w}_{12}(k). \end{aligned} \quad (\text{B.12})$$

Denote

$$\zeta(k) = \frac{2h\rho_0}{2h\rho_0 + \lambda_2(\mathcal{L}_k)}.$$

Noting that H_k is positive semi-definite, by Cauchy inequality, we have

$$2 \left| \eta_1^T(k)H_k\eta_2(k) \right| = 2 \left| \eta_1^T(k)H_k^{\frac{1}{2}}H_k^{\frac{1}{2}}\eta_2(k) \right| \leq \zeta(k)\eta_1^T(k)H_k\eta_1(k) + \frac{1}{\zeta(k)}\eta_2^T(k)H_k\eta_2(k). \quad (\text{B.13})$$

Substituting (B.13) into (B.12) leads to

$$\begin{aligned} & \eta^T \left(\sum_{i=kh}^{(k+1)h-1} \mathbb{E} \left[\widehat{\mathcal{L}}_{\mathcal{G}(i)} \otimes I_n + \mathcal{H}^T(i)\mathcal{H}(i)|\mathcal{F}(kh-1) \right] \right) \eta \\ & \geq (1 - \zeta(k))u_{11}(k) + \left(1 - \frac{1}{\zeta(k)} \right) u_{22}(k) + \tilde{w}_{11}(k) + \tilde{w}_{22}(k) + 2\tilde{w}_{12}(k). \end{aligned} \quad (\text{B.14})$$

We now consider the RHS of (B.14) item by item. Denote $A_k = [r_1(k), \dots, r_n(k)]^T$ and $B_k = [s_1(k), \dots, s_n(k)]$, we have

$$u_{11}(k) = \eta_1^T(k)H_k\eta_1(k) = \left(\sum_{i=1}^n r_i(k)s_i(k) \right)^T H_k \left(\sum_{i=1}^n r_i(k)s_i(k) \right) = A_k^T B_k^T H_k B_k A_k. \quad (\text{B.15})$$

Noting that $H_k = \mathbf{diag}(H_{k,1}, \dots, H_{k,N})$ and $s_i(k) = \frac{1}{\sqrt{N}}\mathbf{1}_N \otimes e_i$, $1 \leq i \leq n$, it follows that

$$B_k^T H_k B_k = \frac{1}{N} \sum_{i=1}^N \sum_{j=kh}^{(k+1)h-1} \mathbb{E} [H_i^T(j)H_i(j)|\mathcal{F}(kh-1)].$$

By (B.15), we get

$$u_{11}(k) \geq \frac{1}{N} \lambda_{\min} \left(\sum_{i=1}^N \sum_{j=kh}^{(k+1)h-1} \mathbb{E} [H_i^T(j)H_i(j)|\mathcal{F}(kh-1)] \right) \left(\sum_{i=1}^n r_i^2(k) \right). \quad (\text{B.16})$$

It can be verified that

$$u_{22}(k) \leq \|H_k\| \|\eta_2(k)\|^2 \leq h\rho_0 \sum_{i=n+1}^{Nn} r_j^2(k) \text{ a.s.} \quad (\text{B.17})$$

Noting that $1 - \frac{1}{\zeta(k)} < 0$, by (B.17), we have

$$\left(1 - \frac{1}{\zeta(k)} \right) u_{22}(k) \geq h\rho_0 \left(1 - \frac{1}{\zeta(k)} \right) \left(1 - \sum_{i=1}^n r_j^2(k) \right). \quad (\text{B.18})$$

It follows from

$$\left(\mathbb{E} \left[\widehat{\mathcal{L}}_{\mathcal{G}(i)} | \mathcal{F}(m_i h - 1) \right] \otimes I_n \right) (\mathbf{1}_N \otimes e_j) = \left(\mathbb{E} \left[\widehat{\mathcal{L}}_{\mathcal{G}(i)} | \mathcal{F}(m_i h - 1) \right] \mathbf{1}_N \right) \otimes e_j = 0$$

that

$$\tilde{w}_{11}(k) = \tilde{w}_{12}(k) = 0. \quad (\text{B.19})$$

Since $\mathcal{G}(k|k-1)$ is balanced, then

$$\lambda_{n+1}(\mathcal{L}_k \otimes I_n) = \lambda_2(\mathcal{L}_k) > 0,$$

where $\lambda_{n+1}(\cdot)$ denotes the $(n+1)$ -th smallest eigenvalue. Noting that $\{s_i(k), n+1 \leq i \leq Nn\}$ is an orthonormal system, we know that

$$\tilde{w}_{22}(k) \geq \lambda_2(\mathcal{L}_k) \left(1 - \sum_{i=1}^n r_i^2(k) \right) \text{ a.s.} \quad (\text{B.20})$$

Denote

$$\begin{aligned} & F_k(x) \\ &= \left(\frac{1 - \zeta(k)}{N} \lambda_{\min} \left(\sum_{i=1}^N \sum_{j=kh}^{(k+1)h-1} \mathbb{E} [H_i^T(j) H_i(j) | \mathcal{F}(kh-1)] \right) - \lambda_2(\mathcal{L}_k) - h\rho_0 + \frac{h\rho_0}{\zeta(k)} \right) x \\ & \quad + \lambda_2(\mathcal{L}_k) + h\rho_0 - \frac{h\rho_0}{\zeta(k)}, \quad x \in \mathbb{R}. \end{aligned}$$

Then, by (B.14) and (B.16)-(B.20), we have

$$\eta^T \left(\sum_{i=kh}^{(k+1)h-1} \mathbb{E} [\widehat{\mathcal{L}}_{\mathcal{G}(i)} \otimes I_n + \mathcal{H}^T(i) \mathcal{H}(i) | \mathcal{F}(kh-1)] \right) \eta \geq F_k \left(\sum_{i=1}^n r_i^2(k) \right) \text{ a.s.}, \quad (\text{B.21})$$

which along with $\zeta(k) = \frac{2h\rho_0}{2h\rho_0 + \lambda_2(\mathcal{L}_k)}$ gives

$$\frac{dF_k(x)}{dx} \leq 0 \text{ a.s.}$$

Hence, the function $F_k(x)$ is monotonically decreasing. It follows from (B.21) and

$$0 \leq \sum_{i=1}^n r_i^2(k) \leq 1$$

that

$$\eta^T \left(\sum_{i=kh}^{(k+1)h-1} \mathbb{E} [\widehat{\mathcal{L}}_{\mathcal{G}(i)} \otimes I_n + \mathcal{H}^T(i) \mathcal{H}(i) | \mathcal{F}(kh-1)] \right) \eta \geq F_k(1) \text{ a.s.},$$

which completes the proof.

Proof of Lemma 3. By the definition of the linear optimal estimated parameter x_{LMS}^* , the regression model (1), Assumption **(A2)** and Lemma A.1 in [39], we have

$$x_{\text{LMS}}^* = \arg \min_{x \in \mathbb{R}^n} \sum_{t=0}^T \sum_{j=1}^N \frac{1}{2} [\mathbb{E}[\|H_j(t)(x - x_0)\|^2] + \mathbb{E}[\|v_j(t)\|^2]].$$

Thus, we have

$$\mathbb{E}[l_{j,t}(x_{\text{LMS}}^*)] = \frac{1}{2} \mathbb{E}[\|v_j(t)\|^2].$$

By the definition of the loss function $l_{j,t}(\cdot)$ and regression model (1), we get

$$\mathbb{E}[l_{j,t}(x_0)] = \frac{1}{2} \mathbb{E}[\|H_j(t)x_0 - y_j(t)\|^2] = \frac{1}{2} \mathbb{E}[\|v_j(t)\|^2] = \mathbb{E}[l_{j,t}(x_{\text{LMS}}^*)],$$

and

$$\mathbb{E}[l_{j,t}(x_i(t))] = \frac{1}{2} \mathbb{E}[\mathbb{E}[\|H_j(t)(x_i(t) - x_0) - v_j(t)\|^2 | \mathcal{F}(t-1)]] .$$

Noting that $x_i(t)$ is adapted to $\mathcal{F}(t-1)$, by Assumption **(A2)** and Lemma A.1 in [39], we obtain

$$\mathbb{E}[l_{j,t}(x_i(t))] = \frac{1}{2} \mathbb{E}[\|H_j(t)(x_i(t) - x_0)\|^2] + \frac{1}{2} \mathbb{E}[\|v_j(t)\|^2].$$

Then it follows that

$$\text{Regret}_{\text{LMS}}(i, T) \leq \frac{\rho_0 N}{2} \sum_{t=0}^T \mathbb{E}[V(t)].$$

APPENDIX C

KEY LEMMAS

Lemma A.1. For the algorithm (6), if Assumption **(A1)** holds and there exists a positive constant ρ_0 , such that $\sup_{k \geq 0} \|\mathcal{L}_{\mathcal{G}(k)}\| \leq \rho_0$ a.s., then $\sup_{k \geq 0} \mathbb{E}[\|\mathcal{A}_{\mathcal{G}(k)}\|^2 | \mathcal{F}(k-1)] \leq N\rho_0^2$ a.s., $\|W(k)\| \leq \sqrt{N}\|\mathcal{A}_{\mathcal{G}(k)}\|$, $\sup_{k \geq 0} \mathbb{E}[\|W(k)\|^2 | \mathcal{F}(k-1)] \leq N^2\rho_0^2$ a.s. and $\|M(k)\|^2 \leq 4\sigma^2 V(k) + 2b^2$.

Proof. It follows from the definition of $W(k)$ that

$$\|W(k)\|^2 = \lambda_{\max}(W^T(k)W(k)) = \max_{i \in \mathcal{V}} \alpha_i^T(k) \alpha_i(k).$$

Noting that $\alpha_i^T(k)$ is the i -th row of the weighted adjacency matrix $\mathcal{A}_{\mathcal{G}(k)}$, we have $\alpha_i^T(k) \alpha_i(k) = \sum_{j=1}^N w_{ij}^2(k)$, which gives

$$\|W(k)\|^2 = \max_{i \in \mathcal{V}} \sum_{j=1}^N w_{ij}^2(k) \leq \|\mathcal{A}_{\mathcal{G}(k)}\|_F^2. \quad (\text{C.1})$$

Meanwhile, noting that $\|\cdot\| \leq \|\cdot\|_F \leq \sqrt{N}\|\cdot\|$, by (C.1), we get $\|W(k)\| \leq \sqrt{N}\|\mathcal{A}_{\mathcal{G}(k)}\|$, which gives

$$\mathbb{E}[\|W(k)\|^2 | \mathcal{F}(k-1)] \leq N\mathbb{E}[\|\mathcal{A}_{\mathcal{G}(k)}\|^2 | \mathcal{F}(k-1)]. \quad (\text{C.2})$$

Furthermore, it follows from the definition of the Laplacian matrix $\mathcal{L}_{\mathcal{G}(k)} = \mathcal{D}_{\mathcal{G}(k)} - \mathcal{A}_{\mathcal{G}(k)}$ that

$$\begin{aligned} & \mathbb{E}[\|\mathcal{A}_{\mathcal{G}(k)}\|_F^2 | \mathcal{F}(k-1)] \\ &= \mathbb{E}\left[\sum_{i=1}^N \sum_{j=1}^N w_{ij}^2(k) | \mathcal{F}(k-1)\right] \\ &\leq \mathbb{E}\left[\sum_{1 \leq i \neq j \leq N} w_{ij}^2(k) + \sum_{i=1}^N \left(\sum_{j \neq i} w_{ij}^2(k)\right) | \mathcal{F}(k-1)\right] \\ &= \mathbb{E}[\|\mathcal{L}_{\mathcal{G}(k)}\|_F^2 | \mathcal{F}(k-1)] \\ &\leq N\mathbb{E}[\|\mathcal{L}_{\mathcal{G}(k)}\|^2 | \mathcal{F}(k-1)] \\ &\leq N\rho_0^2 \text{ a.s.}, \end{aligned}$$

which gives

$$\mathbb{E}[\|\mathcal{A}_{\mathcal{G}(k)}\|^2 | \mathcal{F}(k-1)] \leq N\rho_0^2 \text{ a.s.} \quad (\text{C.3})$$

Substituting (C.3) into (C.2) leads to $\mathbb{E}[\|W(k)\|^2 | \mathcal{F}(k-1)] \leq N^2\rho_0^2$ a.s. By Assumption (A1), we have

$$\begin{aligned} & \|M(k)\|^2 \\ &= \max_{1 \leq i, j \leq N} (f_{ji}(x_j(k) - x_i(k)))^2 \\ &\leq \max_{1 \leq i, j \leq N} 2\sigma^2(x_j(k) - x_i(k)) + 2b^2 \\ &\leq 4\sigma^2 \max_{1 \leq i, j \leq N} \{(x_j(k) - x_0)^2 + (x_i(k) - x_0)^2\} + 2b^2 \\ &\leq 4\sigma^2 \sum_{i=1}^N (x_i(k) - x_0)^2 + 2b^2 \\ &= 4\sigma^2 V(k) + 2b^2. \end{aligned}$$

□

Lemma A.2. For the algorithm (6), if Assumptions (A1)-(A2) hold, the algorithm gains $a(k)$, $b(k)$ and $\lambda(k)$ monotonically decrease to zero, and there exists a positive constant ρ_0 , such that $\sup_{k \geq 0} (\|\mathcal{L}_{\mathcal{G}(k)}\| + (\mathbb{E}[\|\mathcal{H}^T(k)\mathcal{H}(k)\|^2 | \mathcal{F}(k-1)]^{\frac{1}{2}}) \leq \rho_0$ a.s., then there exist nonnegative deterministic sequences $\{\alpha(k), k \geq 0\}$ and $\{\gamma(k), k \geq 0\}$ satisfying $\alpha(k) = o(1)$ and $\gamma(k) = \mathcal{O}(a^2(k) + b^2(k) + \lambda(k))$, such that $\mathbb{E}[V(k) | \mathcal{F}(m_k h - 1)] \leq \prod_{i=m_k h}^{k-1} (1 + \alpha(i)) V(m_k h) +$

$\sum_{i=m_k h}^{k-1} \gamma(i) \prod_{j=i+1}^{k-1} (1 + \alpha(j))$ a.s., $h \geq 1$, where $m_k = \lfloor \frac{k}{h} \rfloor$, especially, there exists a positive integer k_1 , such that $\mathbb{E}[V(k)|\mathcal{F}(m_k h - 1)] \leq 2^h V(m_k h) + h 2^h \gamma(m_k h)$ a.s., $k \geq k_1$.

Proof. By the estimation error equation (8), we have

$$\begin{aligned}
& V(k+1) \\
&= (1 - \lambda(k))^2 V(k) - (1 - \lambda(k)) \delta^T(k) (D(k) + D^T(k)) \delta(k) + \delta^T(k) D^T(k) D(k) \delta(k) \\
&\quad + S^T(k) S(k) + 2S^T(k) ((1 - \lambda(k)) I_{Nn} - D(k)) \delta(k) + \lambda^2(k) \|\mathbf{1}_N \otimes x_0\|^2 \\
&\quad - 2\lambda(k) \delta^T(k) ((1 - \lambda(k)) I_{Nn} - D^T(k)) (\mathbf{1}_N \otimes x_0) - 2\lambda(k) S^T(k) (\mathbf{1}_N \otimes x_0) \\
&\leq (1 - \lambda(k))^2 V(k) + |1 - \lambda(k)| |\delta^T(k) (D(k) + D^T(k)) \delta(k)| + \|D(k)\|^2 \|\delta(k)\|^2 + \|S(k)\|^2 \\
&\quad - 2\lambda(k) S^T(k) (\mathbf{1}_N \otimes x_0) + 2S^T(k) ((1 - \lambda(k)) I_{Nn} - D(k)) \delta(k) + N\lambda^2(k) \|x_0\|^2 \\
&\quad + 2\lambda(k) |\delta^T(k) ((1 - \lambda(k)) I_{Nn} - D^T(k)) (\mathbf{1}_N \otimes x_0)|, \tag{C.4}
\end{aligned}$$

where $S(k) = a(k)\mathcal{H}^T(k)v(k) + b(k)W(k)M(k)\xi(k)$, $D(k) = b(k)\mathcal{L}_G(k) \otimes I_n + a(k)\mathcal{H}^T(k)\mathcal{H}(k)$.

We now consider the conditional mathematical expectation of each term on the RHS of (C.4)

w.r.t. $\mathcal{F}(m_k h - 1)$. Noting that $k - 1 \geq m_k h - 1$, by Assumption (A2), we get

$$\begin{aligned}
& \mathbb{E} [S^T(k) ((1 - \lambda(k)) I_{Nn} - D(k)) \delta(k) | \mathcal{F}(m_k h - 1)] \\
&= \mathbb{E} [a(k) v^T(k) \mathcal{H}(k) ((1 - \lambda(k)) I_{Nn} - D(k)) \delta(k) | \mathcal{F}(m_k h - 1)] \\
&\quad + \mathbb{E} [b(k) \xi^T(k) M^T(k) W^T(k) ((1 - \lambda(k)) I_{Nn} - D(k)) \delta(k) | \mathcal{F}(m_k h - 1)] \\
&= \mathbb{E} [\mathbb{E} [a(k) v^T(k) \mathcal{H}(k) ((1 - \lambda(k)) I_{Nn} - D(k)) \delta(k) | \mathcal{F}(k - 1)] | \mathcal{F}(m_k h - 1)] \\
&\quad + \mathbb{E} [\mathbb{E} [b(k) \xi^T(k) M^T(k) W^T(k) ((1 - \lambda(k)) I_{Nn} - D(k)) \delta(k) | \mathcal{F}(k - 1)] | \mathcal{F}(m_k h - 1)] \\
&= a(k) \mathbb{E} [\mathbb{E} [v^T(k) | \mathcal{F}(k - 1)] \mathbb{E} [\mathcal{H}(k) ((1 - \lambda(k)) I_{Nn} - D(k)) | \mathcal{F}(k - 1)] \delta(k) | \mathcal{F}(m_k h - 1)] \\
&\quad + b(k) \mathbb{E} [\mathbb{E} [\xi^T(k) | \mathcal{F}(k - 1)] M^T(k) \\
&\quad \times \mathbb{E} [W^T(k) ((1 - \lambda(k)) I_{Nn} - D(k)) | \mathcal{F}(k - 1)] \delta(k) | \mathcal{F}(m_k h - 1)] \\
&= 0, \tag{C.5}
\end{aligned}$$

where the penultimate equality is due to $\delta(k) \in \mathcal{F}(k - 1)$, $M(k) \in \mathcal{F}(k - 1)$ and Lemma A.1 in [39]. Following the same way as above, we have

$$\begin{aligned}
& \mathbb{E} [S^T(k) (\mathbf{1}_N \otimes x_0) | \mathcal{F}(m_k h - 1)] \\
&= \mathbb{E} [a(k) v^T(k) \mathcal{H}(k) + b(k) \xi^T(k) M^T(k) W^T(k) | \mathcal{F}(m_k h - 1)] (\mathbf{1}_N \otimes x_0) \\
&= a(k) \mathbb{E} [\mathbb{E} [v^T(k) | \mathcal{F}(k - 1)] \mathbb{E} [\mathcal{H}(k) | \mathcal{F}(k - 1)] | \mathcal{F}(m_k h - 1)] (\mathbf{1}_N \otimes x_0) \\
&\quad + b(k) \mathbb{E} [\mathbb{E} [\xi^T(k) | \mathcal{F}(k - 1)] M^T(k) \mathbb{E} [W^T(k) | \mathcal{F}(k - 1)] | \mathcal{F}(m_k h - 1)] (\mathbf{1}_N \otimes x_0) \\
&= 0. \tag{C.6}
\end{aligned}$$

Denote $q(k) = \max\{a(k), b(k)\}$. Noting that $V(k) \in \mathcal{F}(k-1)$, we obtain

$$\begin{aligned}
& \mathbb{E} [\|D(k)\|^2 \|\delta(k)\|^2 | \mathcal{F}(m_k h - 1)] \\
& \leq \mathbb{E} \left[(b(k) \|\mathcal{L}_{\mathcal{G}(k)}\| + a(k) \|\mathcal{H}^T(k) \mathcal{H}(k)\|)^2 \|\delta(k)\|^2 | \mathcal{F}(m_k h - 1) \right] \\
& \leq q^2(k) \mathbb{E} \left[\mathbb{E} \left[(\|\mathcal{L}_{\mathcal{G}(k)}\| + \|\mathcal{H}^T(k) \mathcal{H}(k)\|)^2 V(k) | \mathcal{F}(k-1) \right] | \mathcal{F}(m_k h - 1) \right] \\
& = q^2(k) \mathbb{E} \left[\mathbb{E} \left[(\|\mathcal{L}_{\mathcal{G}(k)}\| + \|\mathcal{H}^T(k) \mathcal{H}(k)\|)^2 | \mathcal{F}(k-1) \right] V(k) | \mathcal{F}(m_k h - 1) \right] \\
& \leq q^2(k) \rho_0^2 \mathbb{E}[V(k) | \mathcal{F}(m_k h - 1)] \text{ a.s.}
\end{aligned} \tag{C.7}$$

It follows from Lemma A.1 that $\mathbb{E}[\|M(k)\|^2 | \mathcal{F}(m_k h - 1)] \leq 4\sigma^2 \mathbb{E}[V(k) | \mathcal{F}(m_k h - 1)] + 2b^2$, which together with Assumption (A2), Lemma A.1 and Lemma A.1 in [39] gives

$$\begin{aligned}
& \mathbb{E} [\|S(k)\|^2 | \mathcal{F}(m_k h - 1)] \\
& \leq 2q^2(k) \mathbb{E} \left[\|\mathcal{H}^T(k)\|^2 \|v(k)\|^2 + \|W(k)\|^2 \|M(k)\|^2 \|\xi(k)\|^2 | \mathcal{F}(m_k h - 1) \right] \\
& \leq 2q^2(k) \mathbb{E} [\rho_0 \beta_v + \mathbb{E} [\|W(k)\|^2 \|M(k)\|^2 \|\xi(k)\|^2 | \mathcal{F}(k-1)] | \mathcal{F}(m_k h - 1)] \\
& = 2q^2(k) \mathbb{E} [\rho_0 \beta_v + \mathbb{E} [\|W(k)\|^2 \|\xi(k)\|^2 | \mathcal{F}(k-1)] \|M(k)\|^2 | \mathcal{F}(m_k h - 1)] \\
& = 2q^2(k) \mathbb{E} [\rho_0 \beta_v + \mathbb{E} [\|W(k)\|^2 | \mathcal{F}(k-1)] \mathbb{E} [\|\xi(k)\|^2 | \mathcal{F}(k-1)] \|M(k)\|^2 | \mathcal{F}(m_k h - 1)] \\
& \leq 2q^2(k) \rho_0 \beta_v + 2q^2(k) N^2 \rho_0^2 \beta_v (4\sigma^2 \mathbb{E}[V(k) | \mathcal{F}(m_k h - 1)] + 2b^2) \text{ a.s.}
\end{aligned} \tag{C.8}$$

Noting that

$$\begin{aligned}
& \mathbb{E} [\|\delta^T(k)(D(k) + D^T(k))\delta(k)\| | \mathcal{F}(m_k h - 1)] \\
& \leq 2\mathbb{E} [V(k) (b(k) \|\mathcal{L}_{\mathcal{G}(k)}\| + a(k) \|\mathcal{H}^T(k) \mathcal{H}(k)\|) | \mathcal{F}(m_k h - 1)] \\
& = 2\mathbb{E} [\mathbb{E} [V(k) (b(k) \|\mathcal{L}_{\mathcal{G}(k)}\| + a(k) \|\mathcal{H}^T(k) \mathcal{H}(k)\|) | \mathcal{F}(k-1)] | \mathcal{F}(m_k h - 1)] \\
& \leq 2q(k) \mathbb{E} [\mathbb{E} [\|\mathcal{L}_{\mathcal{G}(k)}\| + \|\mathcal{H}^T(k) \mathcal{H}(k)\| | \mathcal{F}(k-1)] V(k) | \mathcal{F}(m_k h - 1)] \\
& \leq 2q(k) \rho_0 \mathbb{E}[V(k) | \mathcal{F}(m_k h - 1)] \text{ a.s.},
\end{aligned} \tag{C.9}$$

by mean value inequality, we have

$$\begin{aligned}
& \mathbb{E} [\delta^T(k) ((1 - \lambda(k))I_{Nn} - D^T(k)) (\mathbf{1}_N \otimes x_0) | \mathcal{F}(m_k h - 1)] \\
& \leq \mathbb{E} [\|\delta(k)\| \|(1 - \lambda(k))I_{Nn} - D^T(k)\| \|\mathbf{1}_N \otimes x_0\| | \mathcal{F}(m_k h - 1)] \\
& \leq \mathbb{E} [\|\delta(k)\| (|1 - \lambda(k)| + \|D(k)\|) \|\mathbf{1}_N \otimes x_0\| | \mathcal{F}(m_k h - 1)] \\
& \leq \mathbb{E} [\|\delta(k)\| (|1 - \lambda(k)| + b(k) \|\mathcal{L}_{\mathcal{G}(k)}\| + a(k) \|\mathcal{H}^T(k) \mathcal{H}(k)\|) \|\mathbf{1}_N \otimes x_0\| | \mathcal{F}(m_k h - 1)] \\
& \leq \mathbb{E} [\|\delta(k)\| \|\mathbf{1}_N \otimes x_0\| + \|\delta(k)\| \|\mathbf{1}_N \otimes x_0\| q(k) (\|\mathcal{L}_{\mathcal{G}(k)}\| + \|\mathcal{H}^T(k) \mathcal{H}(k)\|) | \mathcal{F}(m_k h - 1)] \\
& \leq \frac{1}{2} (\mathbb{E}[V(k) | \mathcal{F}(m_k h - 1)] + N\|x_0\|^2) \\
& \quad + \frac{1}{2} (N\|x_0\|^2 \mathbb{E}[V(k) | \mathcal{F}(m_k h - 1)] + q^2(k) \rho_0^2) \text{ a.s.},
\end{aligned} \tag{C.10}$$

which together with (C.4)-(C.10) leads to

$$\mathbb{E}[V(k+1)|\mathcal{F}(m_k h - 1)] \leq (1 + \alpha(k))\mathbb{E}[V(k)|\mathcal{F}(m_k h - 1)] + \gamma(k) \text{ a.s.}, \quad (\text{C.11})$$

where $\alpha(k) = \lambda^2(k) + \lambda(k)(1 + N\|x_0\|^2) + q^2(k)\rho_0^2 + 8q^2(k)N^2\rho_0^2\beta_v\sigma^2 + 2\rho_0|1 - \lambda(k)|q(k)$ and $\gamma(k) = 4b^2q^2(k)N^2\rho_0^2\beta_v + 2q^2(k)\rho_0\beta_v + (N\|x_0\|^2 + q^2(k)\rho_0^2)\lambda(k) + N\|x_0\|^2\lambda^2(k)$. Thus, by (C.11), we get

$$\mathbb{E}[V(k)|\mathcal{F}(m_k h - 1)] \leq \prod_{i=m_k h}^{k-1} (1 + \alpha(i))V(m_k h) + \sum_{i=m_k h}^{k-1} \gamma(i) \prod_{j=i+1}^{k-1} (1 + \alpha(j)) \text{ a.s.} \quad (\text{C.12})$$

Since $a(k)$, $b(k)$ and $\lambda(k)$ converge to 0, then $\alpha(k) \rightarrow 0$ and $\gamma(k) \rightarrow 0$ as $k \rightarrow \infty$, from which we know that there exists a positive integer k_1 such that $\alpha(i) \leq 1$, $i \geq k_1$. Noting that $0 \leq k - m_k h < h$, the lemma is proved by (C.12). \square

The following lemma proves that the norm of the conditional expectation of $\Phi_P^T((k+1)h - 1, kh)\Phi_P((k+1)h - 1, kh)$ w.r.t. $\mathcal{F}(kh - 1)$ has an upper bound strictly less than 1 in the time interval $[kh, (k+1)h - 1]$ by performing binomial expansion and inequality estimation on the homogeneous part of the estimation error equation (8), which plays an important role in the analysis of the convergence of the algorithm.

Lemma A.3. For the algorithm (6), if the algorithm gains $a(k)$, $b(k)$ and $\lambda(k)$ monotonically decrease satisfying $\sum_{k=0}^{\infty} (a^2(k) + b^2(k) + \lambda^2(k)) < \infty$, and there exists a positive integer h and a positive constant ρ_0 , such that $\sup_{k \geq 0} (\|\mathcal{L}_{\mathcal{G}(k)}\| + (\mathbb{E}[\|\mathcal{H}^T(k)\mathcal{H}(k)\|^{2^{\max\{h, 2\}}}] | \mathcal{F}(k-1)])^{\frac{1}{2^{\max\{h, 2\}}}}) \leq \rho_0$ a.s., then there exists a positive integer k_2 , such that $\|\mathbb{E}[\Phi_P^T((k+1)h - 1, kh)\Phi_P((k+1)h - 1, kh)|\mathcal{F}(kh - 1)]\| \leq 1 - 2\Lambda_k^h - 2\sum_{i=k h}^{(k+1)h-1} \lambda(i) + p(k)$ a.s., $k \geq k_2$, where $p(k) = (4^h - 1 - 2h)(\rho_0 \max\{a(kh), b(kh)\} + \lambda(kh))^2$, especially, there exists a positive integer k_3 , such that $\|\mathbb{E}[\Phi_P^T((k+1)h - 1, i+1)\Phi_P((k+1)h - 1, i+1)|\mathcal{F}(kh - 1)]\| \leq 2$ a.s., $k \geq k_3$, $\forall i \in [kh, (k+1)h - 1]$.

Proof. Denote $D(k) = I_{Nn} - P(k)$. It follows that

$$\begin{aligned} & \left\| \mathbb{E}[\Phi_P^T((k+1)h - 1, kh)\Phi_P((k+1)h - 1, kh)|\mathcal{F}(kh - 1)] \right\| \\ &= \left\| \mathbb{E}[(1 - \lambda(kh))I_{Nn} - D^T(kh)] \times \cdots \times [(1 - \lambda((k+1)h - 1))I_{Nn} - D^T((k+1)h - 1)] \right. \\ & \quad \times [(1 - \lambda((k+1)h - 1))I_{Nn} - D((k+1)h - 1)] \times \cdots \\ & \quad \left. \times [(1 - \lambda(kh))I_{Nn} - D(kh)] | \mathcal{F}(kh - 1) \right\| \end{aligned}$$

$$\begin{aligned}
&= \left\| I_{Nn} - \sum_{i=kh}^{(k+1)h-1} \mathbb{E} [D^T(i) + D(i) + 2\lambda(i)I_{Nn} | \mathcal{F}(kh-1)] + \mathbb{E} \left[\sum_{i=2}^{2h} M_i(k) | \mathcal{F}(kh-1) \right] \right\| \\
&\leq \left\| I_{Nn} - \sum_{i=kh}^{(k+1)h-1} \mathbb{E} [D^T(i) + D(i) + 2\lambda(i)I_{Nn} | \mathcal{F}(kh-1)] \right\| + \left\| \mathbb{E} \left[\sum_{i=2}^{2h} M_i(k) | \mathcal{F}(kh-1) \right] \right\|,
\end{aligned} \tag{C.13}$$

where $M_i(k), i = 2, \dots, 2h$ represent the i -th order terms in the binomial expansion of $\Phi_P^T((k+1)h-1, kh)\Phi_P((k+1)h-1, kh)$. By the definition of spectral radius, we have

$$\begin{aligned}
&\max_{1 \leq i \leq Nn} \lambda_i \left(\sum_{j=kh}^{(k+1)h-1} \mathbb{E} [D^T(j) + D(j) + 2\lambda(j)I_{Nn} | \mathcal{F}(kh-1)] \right) \\
&\leq \left\| \sum_{j=kh}^{(k+1)h-1} \mathbb{E} [D^T(j) + D(j) + 2\lambda(j)I_{Nn} | \mathcal{F}(kh-1)] \right\| \\
&\leq 2 \sum_{j=kh}^{(k+1)h-1} \mathbb{E} [b(j) \|\mathcal{L}_{\mathcal{G}(j)}\| + a(j) \|\mathcal{H}^T(j)\mathcal{H}(j)\| + \lambda(j) | \mathcal{F}(kh-1)] \\
&\leq 2 \sum_{j=kh}^{(k+1)h-1} \max\{a(j), b(j)\} \rho_0 + 2 \sum_{j=kh}^{(k+1)h-1} \lambda(j) \\
&\leq 2 \sum_{j=kh}^{(k+1)h-1} (\rho_0 a(j) + \rho_0 b(j) + \lambda(j)) \text{ a.s.}
\end{aligned} \tag{C.14}$$

Noting that algorithm gains decrease to zero, we know that the RHS of the last inequality of (C.14) converges to zero as $k \rightarrow \infty$, which is independent of the sample paths. Hence, there exists a positive integer l_1 , such that

$$\lambda_i \left(\sum_{j=kh}^{(k+1)h-1} \mathbb{E} [D^T(j) + D(j) + 2\lambda(j)I_{Nn} | \mathcal{F}(kh-1)] \right) \leq 1, \quad i = 1, \dots, Nn, \quad k \geq l_1, \text{ a.s.},$$

from which we get

$$\begin{aligned}
&\left\| I_{Nn} - \sum_{j=kh}^{(k+1)h-1} \mathbb{E} [D^T(j) + D(j) + 2\lambda(j)I_{Nn} | \mathcal{F}(kh-1)] \right\| \\
&= \rho \left(I_{Nn} - \sum_{j=kh}^{(k+1)h-1} \mathbb{E} [D^T(j) + D(j) + 2\lambda(j)I_{Nn} | \mathcal{F}(kh-1)] \right) \\
&= \max_{1 \leq i \leq Nn} \left| \lambda_i \left(I_{Nn} - \sum_{j=kh}^{(k+1)h-1} \mathbb{E} [D^T(j) + D(j) + 2\lambda(j)I_{Nn} | \mathcal{F}(kh-1)] \right) \right| \\
&= 1 - \lambda_{\min} \left(\sum_{j=kh}^{(k+1)h-1} \mathbb{E} [D^T(j) + D(j) + 2\lambda(j)I_{Nn} | \mathcal{F}(kh-1)] \right)
\end{aligned}$$

$$\begin{aligned}
&= 1 - 2\lambda_{\min} \left(\sum_{j=kh}^{(k+1)h-1} \mathbb{E} \left[b(j) \widehat{\mathcal{L}}_{\mathcal{G}(j)} \otimes I_n + a(j) \mathcal{H}^T(j) \mathcal{H}(j) + \lambda(j) I_{Nn} \mid \mathcal{F}(kh-1) \right] \right) \\
&= 1 - 2\Lambda_k^h - 2 \sum_{j=kh}^{(k+1)h-1} \lambda(j), \quad k \geq l_1.
\end{aligned} \tag{C.15}$$

By Cr-inequality, we have

$$\begin{aligned}
&\left[\mathbb{E} \left[\|D(j) + \lambda(j) I_{Nn}\|^{2^r} \mid \mathcal{F}(kh-1) \right] \right]^{\frac{1}{2^r}} \\
&\leq \left[\mathbb{E} \left[\left(\|b(j) \mathcal{L}_{\mathcal{G}(j)} \otimes I_{Nn} + a(j) \mathcal{H}^T(j) \mathcal{H}(j)\| + \lambda(j) \right)^{2^r} \mid \mathcal{F}(kh-1) \right] \right]^{\frac{1}{2^r}} \\
&\leq \left[\mathbb{E} \left[2^{2^r-1} \left(\|b(j) \mathcal{L}_{\mathcal{G}(j)} \otimes I_{Nn} + a(j) \mathcal{H}^T(j) \mathcal{H}(j)\|^{2^r} + \lambda^{2^r}(j) \right) \mid \mathcal{F}(kh-1) \right] \right]^{\frac{1}{2^r}} \\
&\leq 2 \left(\rho_0^{2^r} q^{2^r}(j) + \lambda^{2^r}(j) \right)^{\frac{1}{2^r}} \\
&\leq 2(\rho_0 q(j) + \lambda(j)), \quad 1 \leq r \leq h, \quad kh \leq j \leq (k+1)h-1,
\end{aligned}$$

which together with conditional Hölder inequality and conditional Lyapunov inequality leads to

$$\begin{aligned}
&\mathbb{E} \left[\left\| \prod_{j=1}^r (D(n_j) + \lambda(n_j) I_{Nn}) \right\| \mid \mathcal{F}(kh-1) \right] \\
&\leq \left[\mathbb{E} \left[\left\| \prod_{j=1}^{r-1} (D(n_j) + \lambda(n_j) I_{Nn}) \right\|^2 \mid \mathcal{F}(kh-1) \right] \right]^{\frac{1}{2}} \left[\mathbb{E} \left[\|D(n_r) + \lambda(n_r) I_{Nn}\|^2 \mid \mathcal{F}(kh-1) \right] \right]^{\frac{1}{2}} \\
&\leq 2(\rho_0 q(n_r) + \lambda(n_r)) \left[\mathbb{E} \left[\left\| \prod_{j=1}^{r-1} (D(n_j) + \lambda(n_j) I_{Nn}) \right\|^2 \mid \mathcal{F}(kh-1) \right] \right]^{\frac{1}{2}} \\
&\leq 2^r \prod_{j=1}^r (\rho_0 q(n_j) + \lambda(n_j)) \text{ a.s.}, \quad 1 \leq r \leq h,
\end{aligned} \tag{C.16}$$

where $kh \leq n_1 \leq \dots \leq n_r \leq (k+1)h-1$. On one hand, we have

$$\begin{aligned}
&\mathbb{E}[\|M_i(k)\| \mid \mathcal{F}(kh-1)] \\
&= \mathbb{E} \left[\left\| \sum_{s+t=i} \prod_{l=1}^s (D^T(n_l) + \lambda(n_l) I_{Nn}) \prod_{w=1}^t (D(v_{t+1-w}) + \lambda(v_{t+1-w}) I_{Nn}) \right\| \mid \mathcal{F}(kh-1) \right] \\
&\leq \sum_{s+t=i} \mathbb{E} \left[\left\| \prod_{l=1}^s (D^T(n_l) + \lambda(n_l) I_{Nn}) \right\| \left\| \prod_{w=1}^t (D(v_{t+1-w}) + \lambda(v_{t+1-w}) I_{Nn}) \right\| \mid \mathcal{F}(kh-1) \right] \\
&\leq \sum_{s+t=i} 2^{s+t} \prod_{l=1}^s (\rho_0 q(n_l) + \lambda(n_l)) \prod_{w=1}^t (\rho_0 q(v_w) + \lambda(v_w)) \text{ a.s.}, \quad 2 \leq i \leq h,
\end{aligned} \tag{C.17}$$

where $kh \leq n_1 \leq \dots \leq n_s \leq (k+1)h-1$, $kh \leq v_1 \leq \dots \leq v_t \leq (k+1)h-1$. Noting that there exists a positive integer l_2 , such that $\rho_0 q(k) + \lambda(k) \leq 1, k \geq l_2$. By (C.17), we get

$$\mathbb{E}[\|M_i(k)\| \mid \mathcal{F}(kh-1)] \leq C_{2h}^i 2^i (\rho_0 q(kh) + \lambda(kh))^2 \text{ a.s.}, \quad 2 \leq i \leq h, \quad k \geq l_2. \tag{C.18}$$

On the other hand, for the case with $h < i \leq 2h$, each term of $M_i(k)$ is multiplied by at most i ($2 \leq i \leq h$) different elements, where the matrix and its transpose are regarded as the same element. Noting that $\|A^T A\| = \|A\|^2$ for any matrix A . By (C.16), we get

$$\mathbb{E}[\|M_i(k)\| | \mathcal{F}(kh-1)] \leq C_{2h}^i 2^i (\rho_0 q(kh) + \lambda(kh))^2 \text{ a.s., } h < i \leq 2h, k \geq l_2. \quad (\text{C.19})$$

By (C.18) and (C.19), we have

$$\begin{aligned} & \left\| \mathbb{E} \left[\sum_{i=2}^{2h} M_i(k) | \mathcal{F}(kh-1) \right] \right\| \\ & \leq \sum_{i=2}^{2h} C_{2h}^i 2^i (\rho_0 q(kh) + \lambda(kh))^2 \\ & = (4^h - 1 - 2h) (\rho_0 q(kh) + \lambda(kh))^2 \text{ a.s., } k \geq l_2. \end{aligned} \quad (\text{C.20})$$

Denote the RHS of (C.20) by $p(k)$, we know that $\sum_{k=0}^{\infty} p(k) < \infty$. Denote $k_2 = \max\{l_1, l_2\}$.

By (C.13), (C.15) and (C.20), we get

$$\begin{aligned} & \left\| \mathbb{E} \left[\Phi_P^T((k+1)h-1, kh) \Phi_P((k+1)h-1, kh) | \mathcal{F}(kh-1) \right] \right\| \\ & \leq 1 - 2\Lambda_k^h - 2 \sum_{i=kh}^{(k+1)h-1} \lambda(i) + p(k) \text{ a.s., } k \geq k_2. \end{aligned}$$

Furthermore, denote $G_j(k, i)$ the j -th order terms in the binomial expansion of $\Phi_P^T((k+1)h-1, i+1) \Phi_P((k+1)h-1, i+1)$. Similar to (C.19), we get $\mathbb{E}[\|G_j(k, i)\| | \mathcal{F}(i)] \leq C_{2((k+1)h-i-1)}^j 2^j (\rho_0 q(kh) + \lambda(kh))^2 \text{ a.s., } k \geq k_2, kh \leq i \leq (k+1)h-1, 2 \leq j \leq 2((k+1)h-i-1)$, which along with (C.14) gives

$$\begin{aligned} & \left\| \mathbb{E} \left[\Phi_P^T((k+1)h-1, i+1) \Phi_P((k+1)h-1, i+1) | \mathcal{F}(i) \right] \right\| \\ & = \left\| I_{Nn} - \sum_{j=i+1}^{(k+1)h-1} \mathbb{E} [D^T(j) + D(j) + 2\lambda(j) I_{Nn} | \mathcal{F}(i)] + \mathbb{E} \left[\sum_{j=2}^{2((k+1)h-i-1)} G_j(k, i) | \mathcal{F}(i) \right] \right\| \\ & \leq \left\| I_{Nn} - \sum_{j=i+1}^{(k+1)h-1} \mathbb{E} [D^T(j) + D(j) + 2\lambda(j) I_{Nn} | \mathcal{F}(i)] \right\| + \left\| \mathbb{E} \left[\sum_{j=2}^{2((k+1)h-i-1)} G_j(k, i) | \mathcal{F}(i) \right] \right\| \\ & \leq 1 + \left\| \sum_{j=i+1}^{(k+1)h-1} \mathbb{E} [D^T(j) + D(j) + 2\lambda(j) I_{Nn} | \mathcal{F}(i)] \right\| + \sum_{j=2}^{2((k+1)h-i-1)} \mathbb{E} [\|G_j(k, i)\| | \mathcal{F}(i)] \\ & \leq 1 + \sum_{\substack{j=kh \\ (k+1)h-1}}^{(k+1)h-1} \left\| \mathbb{E} [D^T(j) + D(j) + 2\lambda(j) I_{Nn} | \mathcal{F}(i)] \right\| + \sum_{j=2}^{2((k+1)h-i-1)} \mathbb{E} [\|G_j(k, i)\| | \mathcal{F}(i)] \\ & \leq 1 + 2 \sum_{j=kh}^{(k+1)h-1} (\rho_0 a(j) + \rho_0 b(j) + \lambda(j)) + \sum_{j=2}^{2((k+1)h-i-1)} C_{2((k+1)h-i-1)}^j 2^j (\rho_0 q(kh) + \lambda(kh))^2 \end{aligned}$$

$$\begin{aligned}
&\leq 1 + 2 \sum_{j=kh}^{(k+1)h-1} (\rho_0 a(j) + \rho_0 b(j) + \lambda(j)) + \sum_{j=2}^{2h} C_{2h}^j 2^j (\rho_0 q(kh) + \lambda(kh))^2 \\
&= 1 + 2 \sum_{j=kh}^{(k+1)h-1} (\rho_0 a(j) + \rho_0 b(j) + \lambda(j)) + p(k) \text{ a.s., } k \geq k_2,
\end{aligned}$$

which further shows that there exists a positive integer k_3 , such that $\|\mathbb{E}[\Phi_P^T((k+1)h-1, i+1)\Phi_P((k+1)h-1, i+1)|\mathcal{F}(i)]\| \leq 2$ a.s., $k \geq k_3$. \square

Lemma A.4. For the algorithm (6), suppose that $\{\mathcal{G}(k), k \geq 0\} \in \Gamma_1$, Assumptions **(A1)**–**(A2)** hold, the algorithm gains $a(k)$, $b(k)$ and $\lambda(k)$ monotonically decrease to zero, and there exist positive constants ρ_0 , θ_1 and θ_2 and a positive integer h , such that (i) $\inf_{k \geq 0} \lambda_2(\sum_{j=kh}^{(k+1)h-1} \mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(j)} | \mathcal{F}(kh-1)]) \geq \theta_1$ a.s.; (ii) $\lambda_{\min}(\sum_{i=1}^N \sum_{j=kh}^{(k+1)h-1} \mathbb{E}[H_i^T(j)H_i(j) | \mathcal{F}(kh-1)]) \geq \theta_2$ a.s.; (iii) $\sup_{k \geq 0} (\|\mathcal{L}_{\mathcal{G}(k)}\| + (\mathbb{E}[\|\mathcal{H}^T(k)\mathcal{H}(k)\|^2]^{\frac{1}{2\max\{h,2\}}}) |\mathcal{F}(k-1)]) \leq \rho_0$ a.s. Denote

$$G(k) = \frac{\theta_1 \theta_2}{2Nh\rho_0 + N\theta_1} \min\{a((k+1)h), b((k+1)h)\} + \sum_{i=kh}^{(k+1)h-1} \lambda(i).$$

- (I). If $\sum_{k=0}^{\infty} G(k) = \infty$ and $\sum_{k=0}^{\infty} (a^2(k) + b^2(k) + \lambda(k)) < \infty$, then $\lim_{k \rightarrow \infty} x_i(k) = x_0$, $i \in \mathcal{V}$ a.s.
- (II). If $\sum_{k=0}^{\infty} G(k) = \infty$ and $a^2(kh) + b^2(kh) + \lambda(kh) = o(G(k))$, then $\lim_{k \rightarrow \infty} \mathbb{E}[\|x_i(k) - x_0\|^2] = 0$, $i \in \mathcal{V}$.

Proof. Noting that $\{\mathcal{G}(k), k \geq 0\} \in \Gamma_1$, it follows that $\mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(k)} | \mathcal{F}(k-1)]$ is positive semi-definite. Noting that $\mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(k)} | \mathcal{F}(mh-1)] = \mathbb{E}[\mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(k)} | \mathcal{F}(k-1)] | \mathcal{F}(mh-1)]$, $k \geq mh$, then $\mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(k)} | \mathcal{F}(mh-1)]$ is also positive semi-definite, which shows

$$\begin{aligned}
&\sum_{i=kh}^{(k+1)h-1} \mathbb{E} \left[b(i) \widehat{\mathcal{L}}_{\mathcal{G}(i)} \otimes I_n + a(i) \mathcal{H}^T(i) \mathcal{H}(i) | \mathcal{F}(kh-1) \right] \\
&\geq \min\{a((k+1)h), b((k+1)h)\} \sum_{i=kh}^{(k+1)h-1} \mathbb{E} \left[\widehat{\mathcal{L}}_{\mathcal{G}(i)} \otimes I_n + \mathcal{H}^T(i) \mathcal{H}(i) | \mathcal{F}(kh-1) \right]. \quad (\text{C.21})
\end{aligned}$$

By Lemma 2, condition (ii) and (C.21), we get

$$\Lambda_k^h \geq \frac{\theta_1 \theta_2}{2Nh\rho_0 + N\theta_1} \min\{a((k+1)h), b((k+1)h)\} \text{ a.s.,}$$

which leads to

$$\Lambda_k^h + \sum_{i=kh}^{(k+1)h-1} \lambda(i) \geq G(k). \quad (\text{C.22})$$

We first prove (I) of Lemma A.4. Here, the algorithm gains guarantee that the sample path spatio-temporal persistence of excitation condition holds, then the algorithm (6) converges almost surely by Theorem 1.

Next, we will prove (II) of Lemma A.4. By (C.22) and Lemma 1, there exists a positive integer k_0 , such that

$$\mathbb{E}[V((k+1)h)|\mathcal{F}(kh-1)] \leq (1 + \Omega(k))V(kh) - 2G(k)V(kh) + \Gamma(k) \text{ a.s., } k \geq k_0, \quad (\text{C.23})$$

where $\Omega(k) + \Gamma(k) = \mathcal{O}(a^2(kh) + b^2(kh) + \lambda(kh))$. Noting that $G(k) = o(1)$ and $\Omega(k) = o(G(k))$, without loss of generality, we suppose that $0 < \Omega(k) \leq G(k) < 1$, $k \geq k_0$. Taking mathematical expectation on both sides of (C.23), we get $\mathbb{E}[V((k+1)h)] \leq (1 - G(k))\mathbb{E}[V(kh)] + \Gamma(k)$, $k \geq k_0$. On one hand, we know that $\Gamma(k) = o(G(k))$ and $\sum_{k=0}^{\infty} G(k) = \infty$, which together with Lemma A.3 in [39] gives

$$\lim_{k \rightarrow \infty} \mathbb{E}[V(kh)] = 0. \quad (\text{C.24})$$

On the other hand, by Lemma A.2, we know that there exists a positive integer k_1 , such that

$$\mathbb{E}[V(k)] \leq 2^h \mathbb{E}[V(m_k h)] + h 2^h \gamma(m_k h), \quad k \geq k_1,$$

where $\gamma(k) = o(1)$ and $m_k = \lfloor \frac{k}{h} \rfloor$. Noting that $0 \leq k - m_k h < h$, by (C.24), we get $\lim_{k \rightarrow \infty} \mathbb{E}[V(k)] = 0$. \square

Especially, for the case with the graphs and regression matrices, which jointly satisfies the stochastic spatio-temporal persistence of excitation condition in [38], we have the following lemma.

Lemma A.5. For the algorithm (6), suppose that $\{\mathcal{G}(k), k \geq 0\} \in \Gamma_1$, Assumptions (A1)-(A2) hold, the algorithm gains $a(k)$, $b(k)$ and $\lambda(k)$ monotonically decrease to zero, and there exists a positive integer h and positive constants θ and ρ_0 , such that (i) $\inf_{k \geq 0} \tilde{\Lambda}_k^h \geq \theta$ a.s.; (ii) $\sup_{k \geq 0} (\|\mathcal{L}_{\mathcal{G}(k)}\| + (\mathbb{E}[\|\mathcal{H}^T(k)\mathcal{H}(k)\|^{2^{\max\{h,2\}}}]|\mathcal{F}(k-1)])^{\frac{1}{2^{\max\{h,2\}}}}) \leq \rho_0$ a.s. Denote

$$J(k) = \min\{a((k+1)h), b((k+1)h)\}\theta + \sum_{i=kh}^{(k+1)h-1} \lambda(i).$$

(I). If $\sum_{k=0}^{\infty} J(k) = \infty$, and $\sum_{k=0}^{\infty} (a^2(k) + b^2(k) + \lambda(k)) < \infty$, then $\lim_{k \rightarrow \infty} x_i(k) = x_0$, $i \in \mathcal{V}$ a.s.

(II). If $\sum_{k=0}^{\infty} J(k) = \infty$, and $a^2(kh) + b^2(kh) + \lambda(kh) = o(J(k))$, then $\lim_{k \rightarrow \infty} \mathbb{E}[\|x_i(k) - x_0\|^2] = 0$, $i \in \mathcal{V}$.

Proof. It follows from $\{\mathcal{G}(k), k \geq 0\} \in \Gamma_1$ that $\mathbb{E}[\widehat{\mathcal{L}}_{\mathcal{G}(k)} | \mathcal{F}(k-1)]$ is positive semi-definite, which shows that

$$\Lambda_k^h \geq \min\{a((k+1)h), b((k+1)h)\} \widetilde{\Lambda}_k^h \text{ a.s.,}$$

thus, by condition (i), we get

$$\Lambda_k^h + \sum_{i=kh}^{(k+1)h-1} \lambda(i) \geq J(k) \text{ a.s.,} \quad (\text{C.25})$$

which together with $\sum_{k=0}^{\infty} J(k) = \infty$ gives $\sum_{k=0}^{\infty} (\Lambda_k^h + \sum_{i=kh}^{(k+1)h-1} \lambda(i)) = \infty$ a.s. Noting that $\sum_{k=0}^{\infty} (a^2(k) + b^2(k) + \lambda(k)) < \infty$, by Theorem 1, the algorithm converges almost surely.

Next, we will prove (II) of Lemma A.5. By Lemma 1 and (C.25), there exists a positive integer k_0 such that

$$\mathbb{E}[V((k+1)h) | \mathcal{F}(kh-1)] \leq (1 + \Omega(k))V(kh) - 2J(k)V(kh) + \Gamma(k) \text{ a.s., } k \geq k_0, \quad (\text{C.26})$$

where $\Omega(k) + \Gamma(k) = \mathcal{O}(a^2(kh) + b^2(kh) + \lambda(kh))$. Noting that $a^2(kh) + b^2(kh) + \lambda(kh) = o(J(k))$, we know that $\Omega(k) = o(J(k))$. Since $J(k) = o(1)$, then, without loss of generality, we suppose that $0 < \Omega(k) \leq J(k) < 1$, $k \geq k_0$. Taking mathematical expectation on both sides of (C.26) leads to

$$\mathbb{E}[V((k+1)h)] \leq (1 - J(k))V(kh) + \Gamma(k) \text{ a.s., } k \geq k_0.$$

On one hand, noting that $\Gamma(k) = o(J(k))$, it follows from $\sum_{k=0}^{\infty} J(k) = \infty$ and Lemma A.3 in [39] that

$$\lim_{k \rightarrow \infty} \mathbb{E}[V(kh)] = 0. \quad (\text{C.27})$$

On the other hand, by Lemma A.2, there exists a positive integer k_1 , such that

$$\mathbb{E}[V(k)] \leq 2^h \mathbb{E}[V(m_k h)] + h 2^h \gamma(m_k h), \quad k \geq k_1, \quad (\text{C.28})$$

where $\gamma(k) = o(1)$ and $m_k = \lfloor \frac{k}{h} \rfloor$. Noting that $0 \leq k - m_k h < h$, by (C.27) and (C.28), we have $\lim_{k \rightarrow \infty} \mathbb{E}[V(k)] = 0$. \square

REFERENCES

- [1] T. Evgeniou, M. Pontil, and T. Poggio, “Regularization networks and support vector machines,” *Advances in Computational Mathematics*, vol. 13, no. 1, pp. 1-50, Apr. 2000.
- [2] F. Girosi, “An equivalence between sparse approximation and support vector machines neural computation,” *Neural Computation*, vol. 10, no. 6, pp. 1455-1480, Aug. 1998.
- [3] T. Poggio and S. Smale, “The Mathematics of learning: dealing with data,” *Notices of the American Mathematical Society*, vol. 50, no. 5, pp. 537-544, May 2003.
- [4] S. Theodoridis, *Machine learning: a Bayesian and optimization perspective*. London, UK: Elsevier, 2015.
- [5] H. Xue and Z. Ren, “Sketch discriminatively regularized online gradient descent classification,” *Applied Intelligence*, vol. 50, pp. 1367-1378, Jan. 2020.
- [6] N. Zhou, D. J. Trudnowski, J. W. Pierre, and W. A. Mittelstadt, “Electromechanical mode online estimation using regularized robust RLS methods,” *IEEE Trans. Power Systems*, vol. 23, no. 4, pp. 1670-1680, Nov. 2008.
- [7] Y. Sun, B. Wohlberg, and U. S. Kamilov, “An online plug-and-play algorithm for regularized image reconstruction,” *IEEE Trans. Computational Imaging*, vol. 5, no. 3, pp. 395-408, Sep. 2019.
- [8] T. Matsushita, “Algorithm for atomic resolution holography using modified L_1 -regularized linear regression and steepest descent method,” *Physica Status Solidi (b)*, vol. 255, no. 11, pp. 1800091, Jul. 2018.
- [9] K. H. Ng, S. Tatinati, and A. W. Khong, “Grade prediction from multi-valued click-stream traces via bayesian-regularized deep neural networks,” *IEEE Trans. Signal Processing*, vol. 69, pp. 1477-1491, Feb. 2021.
- [10] V. K. Ivanov, V. V. Vasin, and V. P. Tanana, *Theory of linear ill-posed problems and its applications*, Berlin, Germany: Walter de Gruyter, 2002.
- [11] E. D. Vito, L. Rosasco, A. Caponnetto, U. D. Giovannini, F. Odone, and P. Bartlett, “Learning from examples as an inverse problem,” *J. Machine Learning Research*, vol. 6, no. 5, pp. 883-904, May 2005.
- [12] N. Cesa-bianchi, P. M. Long, and M. K. Warmuth, “Worst-case quadratic loss bounds for prediction using linear functions and gradient descent,” *IEEE Trans. Neural Networks*, vol. 7, no. 3, pp. 604-619, May 1996.
- [13] J. Kivinen and M. K. Warmuth, “Exponentiated gradient versus gradient descent for linear predictors,” *Information and Computation*, vol. 132, no. 1, pp. 1-63, Jan. 1997.
- [14] V. Vovk, “Competitive on-line statistics,” *International Statistical Review*, vol. 69, no. 2, pp. 213-248, Aug. 2001.
- [15] P. Gaillard, S. Gerchinovitz, M. Huard, and G. Stoltz, “Uniform regret bounds over \mathbb{R}^d for the sequential linear regression problem with the square loss,” in *Proc. 30th Int. Conf. Algorithmic Learning Theory*, Chicago, USA, Mar. 2019, pp. 404-432.
- [16] S. Gerchinovitz, “Sparsity regret bounds for individual sequences in online linear regression,” *J. Machine Learning Research*, vol. 14, no. 1, pp. 729-769, Mar. 2013.
- [17] W. Jamil and A. Bouchachia, “Competitive regularised regression,” *Neurocomputing*, vol. 390, pp. 374-383, May 2020.
- [18] C. Thrampoulidis, S. Oymak, and B. Hassibi, “Regularized linear regression: a precise analysis of the estimation error,” in *Proc. 28th Conf. Learning Theory*, Jun. 2015, pp. 1683-1709.
- [19] H. Zou, “The adaptive lasso and its oracle properties,” *J. Amer. Stat. Assoc.*, vol. 101, no. 476, pp. 1418-1429, Dec. 2006.
- [20] J. Fan and R. Li, “Variable selection via nonconcave penalized likelihood and its oracle properties,” *J. Amer. Stat. Assoc.*, vol. 96, no. 456, pp. 1348-1360, Dec. 2001.
- [21] S. Shalev-Shwartz, “Online learning and online convex optimization,” *Foundations and Trends in Machine Learning*, vol. 4, no. 2, pp. 107-194, Mar. 2012.

- [22] E. Hazan, “Introduction to online convex optimization,” *Foundations and Trends in Optimization*, vol. 2, no. 3-4, pp. 157-325, Aug. 2016.
- [23] K. Scaman, F. Bach, S. Bubeck, L. Massoulié, and Y. T. Lee, “Optimal algorithms for non-smooth distributed optimization in networks,” in *Proc. 32nd Conf. Neural Information Processing Systems*, Montréal, Canada, Dec. 2018, pp. 2745-2754.
- [24] J. B. Predd, S. R. Kulkarni, and H. V. Poor, “A collaborative training algorithm for distributed learning,” *IEEE Trans. Information Theory*, vol. 55, no. 4, pp. 1856-1871, Mar. 2009.
- [25] Y. Liu, J. Liu, and T. Basar, “Differentially private gossip gradient descent,” in *Proc. 57th IEEE Conf. Decision and Control*, Miami, USA, Dec. 2018, pp. 2777-2782.
- [26] F. Yan, S. Sundaram, S. Vishwanathan, and Y. Qi, “Distributed autonomous online learning: Regrets and intrinsic privacy-preserving properties,” *IEEE Trans. Knowledge and Data Engineering*, vol. 25, no. 11, pp. 2483-2493, Nov. 2013.
- [27] S. Kar and J. M. F. Moura, “Convergence rate analysis of distributed gossip (linear parameter) estimation: fundamental limits and tradeoff,” *IEEE J. Sel. Topics Signal Processing*, vol. 5, no. 4, pp. 674-690, Aug. 2011.
- [28] S. Kar, J. M. F. Moura, and K. Ramanan, “Distributed parameter estimation in sensor networks: Nonlinear observation models and imperfect communication,” *IEEE Trans. Information Theory*, vol. 58, no. 6, pp. 3575-3605, Jun. 2012.
- [29] S. Kar and J. M. F. Moura, “Consensus+innovations distributed inference over networks: Cooperation and sensing in networked systems,” *IEEE Signal Processing Magazine*, vol. 30, no. 3, pp. 99-109, May 2013.
- [30] S. Kar, J. M. F. Moura, and H. V. Poor, “Distributed linear parameter estimation: asymptotically efficient adaptive strategies,” *SIAM Journal on Control and Optimization*, vol. 51, no. 3, pp. 2200-2229, May 2013.
- [31] A. K. Sahu, S. Kar, J. M. F. Moura, and H. V. Poor, “Distributed constrained recursive nonlinear least-squares estimation: algorithms and asymptotics,” *IEEE Trans. Signal and Information Processing over Networks*, vol. 2, no. 4, pp. 426-441, Dec. 2016.
- [32] S. Xie and L. Guo, “Analysis of normalized least mean squares-based consensus adaptive filters under a general information condition,” *SIAM J. Control and Optimization*, vol. 56, no. 5, pp. 3404-3431, Sep. 2018.
- [33] S. Xie and L. Guo, “Analysis of distributed adaptive filters based on diffusion strategies over sensor networks,” *IEEE Trans. Automatic Control*, vol. 63, no. 11, pp. 3643-3658, Nov. 2018.
- [34] A. K. Sahu, D. Jakovetic, and S. Kar, “ $CTRF_{\varepsilon}$: A distributed random fields estimator,” *IEEE Trans. Signal Processing*, vol. 66, no. 18, pp. 4980-4995, Sep. 2018.
- [35] Y. Chen, S. Kar, and J. M. F. Moura, “Resilient distributed estimation: Sensor attacks,” *IEEE Trans. Automatic Control*, vol. 64, no. 9, pp. 3772-3779, Sep. 2019.
- [36] D. Yuan, A. Proutiere, and G. Shi, “Distributed online linear regressions,” *IEEE Trans. Information Theory*, vol. 67, no. 1, pp. 616-639, Oct. 2020.
- [37] X. Zhang, T. Li, and Y. Gu, “Consensus+innovations distributed estimation with random network graphs, observation matrices and noises,” in *Proc. 59th IEEE Conf. Decision and Control*, Jeju Island, South Korea, Dec. 2020, pp. 4318-4323.
- [38] J. Wang, T. Li, and X. Zhang, “Decentralized cooperative online estimation with random observation matrices, communication graphs and time delays,” *IEEE Trans. Information Theory*, vol. 67, no. 6, pp. 4035-4059, Jun. 2021.
- [39] T. Li and J. Wang, “Distributed averaging with random network graphs and noises,” *IEEE Trans. Information Theory*, vol. 64, no. 11, pp. 7063-7080, Nov. 2018.
- [40] Z. Zhang, Y. Zhang, D. Guo, S. Zhao, and X. L. Zhu, “Communication-efficient federated continual learning for distributed learning system with Non-IID data,” *SCIENCE CHINA Information Sciences*, vol. 66, no. 2, pp. 122102, Dec. 2022.
- [41] L. Guo, “Estimating time-varying parameters by Kalman filter based algorithm: Stability and convergence,” *IEEE Trans. Automatic Control*, vol. 35, no. 2, pp. 141-147, Feb. 1990.

- [42] J. F. Zhang, L. Guo, and H. F. Chen, " L_p -stability of estimation errors of Kalman filter for tracking time-varying parameters", *Int. J. Adaptive Control and Signal Processing*, vol. 5, no. 3, pp. 155-174, May 1991.
- [43] A. S. Bedi, A. Koppel, and K. Rajawat, "Asynchronous saddle point algorithm for stochastic optimization in heterogeneous networks," *IEEE Trans. Signal Processing*, vol. 67, no. 7, pp. 1742-1757, Apr. 2019.
- [44] R. Dixit, A. S. Bedi, and K. Rajawat, "Online learning over dynamic graphs via distributed proximal gradient algorithm," *IEEE Trans. Automatic Control*, vol. 66, no. 11, pp. 5065-5079, Nov. 2021.
- [45] Q. Zhang and J. F. Zhang, "Distributed parameter estimation over unreliable networks with markovian switching topologies," *IEEE Trans. Automatic Control*, vol. 57, no. 10, pp. 2545-2560, Oct. 2012.
- [46] J. Zhang, X. He, and D. Zhou, "Distributed filtering over wireless sensor networks with parameter and topology uncertainties," *Int. J. Control*, vol. 93, no. 4, pp. 910-921, Apr. 2020.
- [47] A. J. Wood and B. F. Wollenberg, *Power Generation, Operation, and Control*. New York, NY, USA: Wiley, 2012.
- [48] S. Djaidja and Q. Wu, "An overview of distributed consensus of multi-agent systems with measurement/communication noises," in *Proc. 34th Chin. Control Conf.*, Hangzhou, China, Jul. 2015, pp. 7285-7290.
- [49] D. V. Dimarogonas and K. H. Johansson, "Stability analysis for multi-agent systems using the incidence matrix: Quantized communication and formation control," *Automatica*, vol. 46, no. 4, pp. 695-700, Apr. 2010.
- [50] J. Wang and N. Elia, "Mitigation of complex behavior over networked systems: Analysis of spatially invariant structures," *Automatica*, vol. 49, no. 6, pp. 1626-1638, Jun. 2013.
- [51] F. S. Cattivelli and A. H. Sayed, "Diffusion LMS strategies for distributed estimation," *IEEE Trans. Signal Processing*, vol. 58, no. 3, pp. 1035-1048, Mar. 2009.
- [52] M. J. Piggott and V. Solo, "Diffusion LMS with correlated regressors II: Performance," *IEEE Trans. Signal Processing*, vol. 65, no. 15, pp. 3934-3947, Aug. 2017.
- [53] Leonard, N. Ehrich, and A. Olshevsky, "Cooperative learning in multiagent systems from intermittent measurements," *SIAM J. Control and Optimization*, vol. 53, no. 1, pp. 1-29, Jan. 2015.
- [54] S. P. Meyn and R. L. Tweedie, *Markov Chains and Stochastic Stability*. London, UK: Springer-Verlag, 1993.
- [55] H. Robbins and D. Siegmund, "A convergence theorem for non negative almost supermartingales and some applications," in *Selected Papers*, T. L. Lai, and D. Siegmund, Eds., New York, USA: Springer-Verlag, 1985.