

Normalized solutions for Schrödinger-Bopp-Podolsky system

Chuan-Min He, Lin Li* and Shang-Jie Chen[†]

School of Mathematics and Statistics & Chongqing Key Laboratory of Economic and Social Application Statistics,
Chongqing Technology and Business University,
Chongqing 400067, China

Abstract

In this paper, we study the following energy functional originates from the Schrödinger-Bopp-Podolsky system

$$I(u) = \frac{1}{2} \int_{\mathbb{R}^3} |\nabla u|^2 dx + \frac{1}{4} \int_{\mathbb{R}^3} \phi_u u^2 dx - \frac{1}{p} \int_{\mathbb{R}^3} |u|^p dx$$

constrained on $B_\rho = \{u \in H^1(\mathbb{R}^3, C) : \|u\|_2 = \rho\}$, where $\rho > 0$. As such constrained problem $I(u)$ is bounded from below on B_ρ when $p \in (2, \frac{10}{3})$. We use minimizing method to get a normalized solution.

Key words: Normalized solutions, Schrödinger-Bopp-Podolsky system, Constrained minimization

2010 Mathematics Subject Classification: 35Q55, 35J30, 35J50

1 Introduction

In this paper, we consider the following Schrödinger-Bopp-Podolsky system:

$$\begin{cases} -\Delta u + \omega u + \phi u = |u|^{p-2}u & \text{in } \mathbb{R}^3, \\ -\Delta \phi + a^2 \Delta^2 \phi = 4\pi u^2 & \text{in } \mathbb{R}^3. \end{cases} \quad (1.1)$$

where $u, \phi : \mathbb{R}^3 \rightarrow \mathbb{R}$, $\omega > 0$ and let $a = 1$. Such a system was first proposed by d'Avenia and Siciliano[12] and can be used to describe solitary waves for nonlinear stationary equations of Schrödinger type interacting with an electrostatic field in the Bopp-Podolsky electromagnetic theory. The Bopp-Podolsky theory, developed by Bopp[6] and Podolsky[17] independently, can be interpreted as an effective theory for short distances and for large distance it is experimentally

*Corresponding author. E-mail address: linli@ctbu.edu.cn & lilin420@gmail.com (L. Li).

[†]This work is supported by Research Fund of National Natural Science Foundation of China (No. 11861046), National Natural Science Foundation of Chongqing(No. 2019jcyj-msxmX0115), Chongqing Municipal Education Commission (No. KJQN20190081), Chongqing Technology and Business University(No. CTBUZDPTTD201909).

indistinguishable from Maxwell one. For more physical aspects of this system, please refer to [7, 8, 14] and the references therein. Next we focus on the mathematical aspects of the problem.

In recent years, many people consider this system through using variational methods. Next we recall some previous results. At first, d'Avenia and Siciliano[12] have been devoted to the following autonomous system with subcritical growth:

$$\begin{cases} -\Delta u + \omega u + q^2 \phi u = |u|^{p-2}u & \text{in } \mathbb{R}^3, \\ -\Delta \phi + a^2 \Delta^2 \phi = 4\pi u^2 & \text{in } \mathbb{R}^3. \end{cases} \quad (1.2)$$

where $\omega, a > 0, q \neq 0$ and $p > 2$. They obtain that problem (1.2) has a nontrivial solution when $p \in (3, 6)$ and $q > 0$ or $p \in (2, 3]$ and $q > 0$ small enough. In the radial case, they get the solutions tend to solutions of classical Schrödinger-Poisson system as $a \rightarrow 0$. Furthermore, they prove that problem (1.2) does not have a nontrivial solution when $p \geq 6$ through using a Pohožaev identity. Siciliano and Silva[19], by means of the fibering approach, prove the system (1.2) has no solutions at all for large enough of q and has two radical solutions for small enough of q when $p \in (2, 3]$, which improve some results in [12]. Furthermore, under the assumption that ω is replaced by a coercive potential $V(x)$ and $|u|^{p-2}u$ is replaced with $f(u)$ in [24], they proved that problem (1.2) has a ground state when $f(u) = |u|^{p-2}u + h(x)$ with $p \in [4, 6)$ and at least one positive solution for $f(u) = P(x)u^5 + \mu|u|^{p-2}u$ with $p \in (2, 6)$. They also proved that the problem (1.2) possesses infinitely many nontrivial solutions when $f(u)$ satisfies the following conditions:

$$(f_1) \quad f(t) = -f(-t).$$

$$(f_2) \quad \text{There exist } t \in (1, 5) \text{ such that } \lim_{t \rightarrow 0} \frac{f(t)}{t} = \lim_{t \rightarrow +\infty} \frac{f(t)}{|t|^t} = 0.$$

$$(f_3) \quad \lim_{|t| \rightarrow +\infty} \frac{F(t)}{|t|^4} = \infty, \text{ and there exists } \mu \geq 4, \kappa > 0 \text{ such that } \mu F(t) \leq tf(t) + \kappa t^2, \text{ where } F(t) = \int_0^t f(r)dr.$$

Afterwards, Chen and Tang[10] study the following critical Schrödinger-Bopp-Podolsky with subcritical perturbations of general function:

$$\begin{cases} -\Delta u + V(x)u + \phi u = \mu f(u) + u^5 & \text{in } \mathbb{R}^3, \\ -\Delta \phi + a^2 \Delta^2 \phi = 4\pi u^2 & \text{in } \mathbb{R}^3, \end{cases} \quad (1.3)$$

get a nontrivial solution and ground state solution. Yang, Chen and Liu[21] study the existence of nontrivial solution when $f(u)$ in (1.3) without any growth and Ambrosetti-Rabinowitz conditions. Later, Li, Pucci and Tang[15], by using the method of the Pohožaev-Nehari manifold, get a nontrivial ground state solution for (1.3) when $f(u) = |u|^{p-1}u$.

Observe all the above articles, it is not difficult to find that they all regard ω as a fixed frequency parameter to seek nontrivial solutions. Hence, nothing can be given a priori on the L^2 -norm of the solutions. However, many physicists are very interested in normalized solutions, i.e. solutions with prescribed L^2 -norm. To my best knowledge, for solving the normalized solution of the Schrödinger-Bopp-Podolsky system, only paper [1] consider it under Neumann boundary conditions. In the present paper, we study whether (1.1) has normalized solutions without Neumann boundary

conditions. A normalized solution of (1.1) can be obtained as a constrained critical point of the functional

$$I(u) = \frac{1}{2} \int_{\mathbb{R}^3} |\nabla u|^2 dx + \frac{1}{4} \int_{\mathbb{R}^3} \phi_u u^2 dx - \frac{1}{p} \int_{\mathbb{R}^3} |u|^p dx \quad (1.4)$$

on the L^2 -sphere in $H^1(\mathbb{R}^3, C)$

$$B_\rho = \{u \in H^1(\mathbb{R}^3, C) : \|u\|_2 = \rho\},$$

where ϕ_u is defined in Section 2. It should be noted that in this case the frequency ω is no longer imposed but instead appears as a Lagrange parameter.

Consider the following minimization problem

$$I_{\rho^2} = \inf_{B_\rho} I(u),$$

which makes sense for $p \in (2, \frac{10}{3})$. Moreover, in this case, we know that the C^1 functional I is bounded from below and coercive on B_ρ . The detailed proof procedure can be seen in Lemma 2.7. Hence, the most dramatic difficult problem is the minimizing sequences $\{u_n\} \subset B_\rho$ the lack of compactness. In fact, there will be two bad possibilities, namely

- (i) $u_n \rightharpoonup 0$.
- (ii) $u_n \rightharpoonup \hat{u} \neq 0$ and $0 < \|\hat{u}\|_2 < \rho$.

We can rule out both cases using the general method. i.e. first prove any minimizing sequence weakly converges, up to translation, to a function \hat{u} which is different from zero, excluding the vanishing case and then we have to show $\|\hat{u}\|_2 = \rho$. In order to prove $\|\hat{u}\|_2 = \rho$, we need to know that function I satisfies the strong subadditivity inequality, namely

$$I_{\rho^2} < I_{\mu^2} + I_{\rho^2 - \mu^2} \quad \text{for all } 0 < \mu < \rho. \quad (1.5)$$

When $p \in (3, \frac{10}{3})$, we adapt a similar argument as in [4]. By standard scaling arguments and some basic properties of functions we can get (1.5). However, when $p \in (2, 3)$, it will be difficult to prove (1.5) using the above method. Therefore, we use the method of [3]. We must show that

(HD) the function $s \mapsto \frac{I_{s^2}}{s^2}$ is monotone decreasing.

In fact, if **(HD)** hold when $\mu \in (0, \rho)$, from direct calculation we obtain

$$\frac{\mu^2}{\rho^2} I_{\rho^2} < I_{\mu^2} \quad \frac{\rho^2 - \mu^2}{\rho^2} < I_{\rho^2 - \mu^2}.$$

Therefore,

$$I_{\rho^2} = \frac{\mu^2}{\rho^2} I_{\rho^2} + \frac{\rho^2 - \mu^2}{\rho^2} I_{\rho^2} < I_{\mu^2} + I_{\rho^2 - \mu^2}.$$

Proving that satisfying **(HD)** is not an easy task. The function $s \mapsto \frac{I_{s^2}}{s^2}$ have a fast oscillating behavior, even in a neighborhood of the origin. We use a large class of functionals provided that they satisfy some good scaling properties and give sufficient condition to guarantee **(HD)** hold.

Our results are as follows.

Theorem 1.1. *Let $p \in (3, \frac{10}{3})$ and $\rho > 0$, there exists $\rho_1 > 0$ such that for any $\rho \in (\rho_1, +\infty)$ there exists a couple $(u_\rho, \lambda_\rho) \in H^1(\mathbb{R}^3) \times \mathbb{R}^+$ solution of (1.1) with $\|u_\rho\|_2 = \rho$.*

Theorem 1.2. *Let $p \in (2, 3)$ and $\rho > 0$, there exists $\rho_2 > 0$ such that for any $\rho \in (0, \rho_2)$ there exists a couple $(u_\rho, \lambda_\rho) \in H^1(\mathbb{R}^3) \times \mathbb{R}^+$ solution of (1.1) with $\|u_\rho\|_2 = \rho$.*

Finally, we prove the orbital stability of standing waves for Schrödinger-Bopp-Podolsky system. It is well-known that the original approach of proving orbital stability is very complicated. Therefore, we draw on the method of [9] to get the following theorem.

Theorem 1.3. *Let $p \in (2, \frac{10}{3})$. Then the set*

$$S_\rho = \left\{ e^{i\theta} u(x) : \theta \in [0, 2\pi), \|u\|_2 = \rho, I(u) = I_{\rho^2} \right\}$$

is orbitally stable.

Remark 1.4. Notice that if $p \geq \frac{10}{3}$, by observing Lemma 2.7, we know the functional I is unbounded from below on B_ρ . Therefore, the minimizing method will no longer apply. At this point we know that if $a \rightarrow 0$ in problem (1.1) the Schrödinger-Bopp-Podolsky system becomes the classical Schrödinger-Poisson system, namely

$$\begin{cases} -\Delta u + \omega u + \phi u = |u|^{p-2}u & \text{in } \mathbb{R}^3, \\ -\Delta \phi = 4\pi u^2 & \text{in } \mathbb{R}^3. \end{cases} \quad (1.6)$$

Regarding the normalized solution of the mass sup-criticality of problem (1.6), the paper [2] investigates the case $p \in (\frac{10}{3}, 6)$. They proved that problem (1.6) has a mountain-pass geometry and similar to [5] develop a deformation argument to get a localization of Palais-Smale sequence. However, the mountain-pass geometry can not guarantee the existence of a bounded Palais-Smale sequence, so a set of constraints about Pohožaev equality was constructed to solve this problem. Moreover, concerning the lack of compactness for Palais-Smale sequence, it does not to reduce the problem to the classical vanishing-dichotomy-compactness scenario or like this paper method to check of the strong subadditivity inequalities. They studied some properties about the mountain-pass level function to overcome this difficulty. Afterwards, using the above geometric structure and adding a new constraint $\int_{\mathbb{R}^3} |\nabla u|^2 dx < \frac{3}{5} \int_{\mathbb{R}^3} |u|^{\frac{10}{3}} dx$, Ye [22] studied the existence and the concentration behavior for problem (1.6) when $p = \frac{10}{3}$. In addition, those papers [11, 13, 20, 23] has also studied the normalized solution of problem (1.6) under different suitable assumptions. Unfortunately, None of the methods in the above papers can be used in the Schrödinger-Bopp-Podolsky system. Because if we do the scaling $u_\lambda(\cdot) = \lambda^\alpha u(\lambda^\beta(\cdot))$, $\alpha, \beta \in \mathbb{R}$, $\lambda > 0$ of the nonlocal term ϕ_u , form (2.2) we know it is impossible to get $\phi_{u_{\lambda(\cdot)}} = \lambda^\xi \phi_u$ where $\xi = \xi(\alpha, \beta)$. Moreover, $\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} e^{-|x-y|} u^2(x) u^2(y) dx dy$ in Pohožaev equality is also difficult to handle. Therefore, we can not study the normalized solution for problem (1.1) when $p \geq \frac{10}{3}$.

2 Preliminaries

In this section we explain the notations and some auxiliary lemmas which are useful later. We cite the book [16] for the standard reference book.

Let $H^1(\mathbb{R}^3)$ denote the usual Sobolev space with respect to the norm

$$\|u\|_{H^1(\mathbb{R}^3)}^2 = \int_{\mathbb{R}^3} (|\nabla u|^2 + u^2) dx.$$

$L^p(\mathbb{R}^3)$ denote the usual Lebesgue space with respect to the norm

$$\|u\|_p^p = \int_{\mathbb{R}^3} |u|^p dx.$$

$D^{1,2}(\mathbb{R}^3)$ is the completion of C_0^∞ with respect with

$$\|u\|_{D^{1,2}(\mathbb{R}^3)}^2 = \int_{\mathbb{R}^3} |\nabla u|^2 dx.$$

The \mathcal{D} is defined by the completion of $C_0^\infty(\mathbb{R}^3)$ equipped with the scalar product

$$\langle u, v \rangle_{\mathcal{D}} := \int_{\mathbb{R}^3} \nabla u \nabla v dx + \int_{\mathbb{R}^3} \Delta u \Delta v dx.$$

It is easy to know that $\mathcal{D}(\mathbb{R}^3)$ is continuously embedded in $D^{1,2}(\mathbb{R}^3)$ and consequently in $L^\infty(\mathbb{R}^3)$.

C_1, C_2, \dots denote positive constant possibly different in different places.

For the sake of brevity, from now on we define the following quantities:

$$\begin{aligned} A(u) &:= \frac{1}{2} \int_{\mathbb{R}^3} |\nabla u|^2 dx & B(u) &:= \frac{1}{4} \int_{\mathbb{R}^3} \phi_u |u|^2 dx \\ C(u) &:= -\frac{1}{p} \int_{\mathbb{R}^3} |u|^p dx & T(u) &:= B(u) + C(u) \end{aligned}$$

so that

$$I(u) = A(u) + T(u).$$

Definition 2.1. Let $u \in H^1(\mathbb{R}^3)$, $u \neq 0$. A continuous path $g_u : \theta \in \mathbb{R}^+ \mapsto g_u(\theta) \in H^1(\mathbb{R}^3)$ such that $g_u(1) = u$ is said to be a scaling path of u if $\Theta_{g_u}(\theta) := \frac{\|g_u(\theta)\|_2^2}{\|u\|_2^2}$ is differentiable and $\Theta'_{g_u}(1) \neq 0$. We denote with G_u the set of the scaling paths of u .

Moreover, we define a real valued function when $u \neq 0$, which will be useful in subsequent proofs

$$h_{g_u}(\theta) := I(g_u(\theta)) - \Theta_{g_u}(\theta)I(u), \quad \theta \geq 0.$$

Definition 2.2. Let $u \neq 0$ be fixed and $g_u \in G_u$. We say that the scaling path g_u is admissible for the functional I if h_{g_u} is a differentiable function.

Proposition 2.3. ([4]) Let T be a C^1 functional on $H^1(\mathbb{R}^3)$ and $\{u_n\} \subset B_\rho$ be a minimizing sequence for I_ρ such that $u_n \rightharpoonup u \neq 0$; Assume

$$(i) \quad \langle T'(u_n), u_n \rangle = O(1);$$

$$(ii) \quad T(u_n - u) + T(u) = T(u_n) + o(1);$$

$$(iii) \quad T(\alpha_n(u_n - u)) - T(u_n - u) = o(1), \text{ where } \alpha_n = \frac{\rho^2 - \mu^2}{\|u_n - u\|_2};$$

$$(iv) \quad \langle T'(u_n) - T'(u_m), u_n - u_m \rangle = o(1);$$

$$(v) \quad I_{\rho^2} < I_{\mu^2} + I_{\rho^2 - \mu^2} \text{ for any } 0 < \mu < \rho.$$

Then $\|u_n - u\|_{H^1(\mathbb{R}^3)} \rightarrow 0$.

Lemma 2.4. ([12, Lemma 3.2]) *The space $C_0^\infty(\mathbb{R}^3)$ is dense in $\mathcal{A} := \{\phi \in D^{1,2}(\mathbb{R}^3) : \Delta\phi \in L^2(\mathbb{R}^3)\}$ normed by $\sqrt{\langle \phi, \phi \rangle_{\mathcal{D}}}$. Hence, $\mathcal{D} = \mathcal{A}$.*

From [12] we obtain that there exists a unique solution $\phi_u \in \mathcal{D}$ to the second equation in (1.1). Its expression is

$$\phi_u(x) := \int_{\mathbb{R}^3} \frac{1 - e^{-|x-y|}}{|x-y|} u^2(y) dy. \quad (2.1)$$

Notice that if we assume $u_\lambda(\cdot) = \lambda^\alpha u(\lambda^\beta(\cdot))$, $\alpha, \beta \in \mathbb{R}$, $\lambda > 0$, then

$$\begin{aligned} \phi_{u_\lambda}(x) &= \int_{\mathbb{R}^3} \lambda^{2\alpha+\beta} (1 - e^{-\frac{1}{\lambda^\beta}|x^\beta - \lambda^\beta y|}) \frac{|u(\lambda^\beta y)|^2}{|\lambda^\beta x - \lambda^\beta y|} dy \\ &= \lambda^{2(\alpha-\beta)} \int_{\mathbb{R}^3} (1 - e^{-\frac{1}{\lambda^\beta}|x^\beta - y|}) \frac{|u(y)|^2}{|\lambda^\beta x - y|} dy. \end{aligned} \quad (2.2)$$

Moreover ϕ_u has the following properties.

Lemma 2.5. ([12]) *For every $u \in H^1(\mathbb{R}^3)$, we have:*

$$(i) \quad \phi_u \geq 0 \text{ for all } u \in H^1(\mathbb{R}^3).$$

$$(ii) \quad \|\phi_u\|_{\mathcal{D}} \leq C \|u\|^2 \text{ and } \int_{\mathbb{R}^3} \phi_u u^2 dx \leq C \|u\|_{\frac{12}{5}}^4.$$

$$(iii) \quad \text{if } u_n \rightharpoonup u \text{ in } H^1(\mathbb{R}^3), \text{ then } \phi_{u_n} \rightharpoonup \phi_u \text{ in } \mathcal{D}.$$

Lemma 2.6. *Assume I satisfied the following conditions*

(i) *I satisfied the weak subadditivity inequality, namely:*

$$I_{\rho^2} \leq I_{\mu^2} + I_{\rho^2 - \mu^2} \quad \text{for all } 0 < \mu < \rho. \quad (2.3)$$

$$(ii) \quad -\infty < I_{s^2} < 0 \text{ for all } s > 0.$$

$$(iii) \quad s \mapsto I_{s^2} \text{ is continuous.}$$

$$(iv) \quad \lim_{s \rightarrow 0} \frac{I_{s^2}}{s^2} = 0.$$

(v) $\forall u \in M(\rho) := \bigcup_{\mu \in (0, \rho]} \{u \in B_\mu : I(u) = I_{\mu^2}\}$, $\exists g_u \in G(u)$ admissible, such that $\frac{d}{d\theta} h_{g_u}(\theta) \big|_{\theta=1} \neq 0$.

Then **(HD)** holds. Hence, we know $I_{\rho^2} < I_{\mu^2} + I_{\rho^2 - \mu^2}$ for any $0 < \mu < \rho$.

Proof. The detailed proof process can be found in Theorem 2.1 of references [3]. \square

Lemma 2.7. *If $p \in (2, \frac{10}{3})$, then for every $\rho > 0$ the functional I is bounded from below and coercive on B_ρ .*

Proof. Due to Gagliardo–Nirenberg inequality, we can get

$$\|u\|_p^p \leq C_p \|u\|_2^{\frac{6-p}{2}} \|\nabla u\|_2^{\frac{3p}{2}-3}.$$

From (1.4) it is easily to know

$$\begin{aligned} I(u) &\geq \frac{1}{2} \int_{\mathbb{R}^3} |\nabla u|^2 dx - \frac{1}{p} \int_{\mathbb{R}^3} |u|^p dx \\ &\geq \frac{1}{2} \|\nabla u\|_2^2 - \frac{1}{p} C_p \|u\|_2^{\frac{6-p}{2}} \|\nabla u\|_2^{\frac{3p}{2}-3}. \end{aligned}$$

Because of $p < \frac{10}{3}$, hence $\frac{3p}{2} - 3 < 2$, we can obtain

$$I(u) \geq \frac{1}{2} \|\nabla u\|_2^2 + O(\|\nabla u\|_2^2).$$

\square

Lemma 2.8. *If $p \in (2, \frac{10}{3})$, let $u_n \subset B_\rho$ be a minimizing sequence for I_{ρ^2} such that $u_n \rightharpoonup u \neq 0$, then T satisfies the following properties*

$$(i) \quad \langle T'(u_n), u_n \rangle = O(1);$$

$$(ii) \quad T(u_n - u) + T(u) = T(u_n) + o(1);$$

$$(iii) \quad T(\alpha_n(u_n - u)) - T(u_n - u) = o(1), \text{ where } \alpha_n = \frac{\rho^2 - \mu^2}{\|u_n - u\|_2};$$

$$(iv) \quad \langle T'(u_n) - T'(u_m), u_n - u_m \rangle = o(1).$$

Proof. Due to u_n be a minimizing sequence, we can easily to deduce that $\langle T'(u_n), u_n \rangle = O(1)$. From Lemma 2.7 we know that u_n is bounded in $H^1(\mathbb{R}^3)$. Therefore, u_n is bounded in L^s for $s \in [2, 2^*]$ and there exists $u \in H^1(\mathbb{R}^3)$ such that $u_n \rightharpoonup u$ in $H^1(\mathbb{R}^3)$ and $u_n \rightarrow u$ a.e. in \mathbb{R}^3 . We give these notations

$$G(x, y) = \frac{1 - e^{-|x-y|}}{|x-y|}, \quad A := \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} u^2(y) u^2(x) G(x, y) dx dy,$$

$$I_n^{(1)} := \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} u_n^2(y) u^2(x) G(x, y) dx dy,$$

$$I_n^{(2)} := \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} u_n(y) u(y) u_n(x) u(x) G(x, y) dx dy,$$

$$I_n^{(3)} := \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} u_n^2(y) u_n(x) u(x) G(x, y) dx dy,$$

$$I_n^{(3)} := \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} u_n(y) u(y) u^2(x) G(x, y) dx dy.$$

From direct calculation we can deduce

$$B(u_n - u) - (B(u_n) - B(u)) = 2I_n^{(1)} + 4I_n^{(2)} - 4I_n^{(3)} - 4I_n^{(4)} + 2A.$$

Next we will show that $\lim_{n \rightarrow \infty} I_n^{(i)} = A$, $i = 1, 2, 3, 4$. Let

$$v_n(x) := \int_{\mathbb{R}^3} \frac{1 - e^{-|x-y|}}{|x-y|} u_n^2(y) dy, \quad v(x) = \int_{\mathbb{R}^3} \frac{1 - e^{-|x-y|}}{|x-y|} u^2(y) dy.$$

$$\begin{aligned} |v_n(x) - v(x)| &\leq \int_{\mathbb{R}^3} |u_n^2(y) - u^2(y)| \frac{1 - e^{-|x-y|}}{|x-y|} dy \\ &\leq \int_{\mathbb{R}^3} |u_n^2(y) - u^2(y)| \frac{1}{|x-y|} dy \\ &\leq C \|u_n^2 - u^2\|_{L^2(B_R(x))} \left(\int_{|y-x| \leq R} \frac{1}{|x-y|^2} dy \right)^{\frac{1}{2}} \\ &\quad + C \|u_n^2 - u^2\|_{L^{\frac{4}{3}}(B_R^c(x))} \left(\int_{|y-x| \geq R} \frac{1}{|x-y|^2} dy \right)^{\frac{1}{2}}. \end{aligned}$$

Letting $n \rightarrow \infty$ and $R \rightarrow \infty$, we know $v_n \rightarrow v$ a.e. on \mathbb{R}^3 . Due to $v_n = \phi_{u_n} \in \mathcal{D}(\mathbb{R}^3)$ and Lemma 2.5, we know

$$\|v_n\|_6 = C \|v_n\|_D \leq C \|u_n\|_{\frac{12}{5}}^2 \leq C.$$

Hence we can assume $v_n \rightharpoonup v$ in $L^6(\mathbb{R}^3)$ and due to $u \in H^1(\mathbb{R}^3)$ then $u^2 \in L^{\frac{6}{5}}(\mathbb{R}^3)$, we can get

$$\int_{\mathbb{R}^3} v_n u^2 dx \rightarrow \int_{\mathbb{R}^3} v u^2 dx.$$

We prove that $\lim_{n \rightarrow \infty} I_n^{(1)} = A$. Next we prove $\lim_{n \rightarrow \infty} I_n^{(2)} = A$. we set

$$\hat{v}_n(x) := \int_{\mathbb{R}^3} \frac{1 - e^{-|x-y|}}{|x-y|} u_n(y) u(y) dy.$$

A similar proof process to the previous one can easily obtain $\hat{v}_n \rightarrow v$ a.e. in \mathbb{R}^3 . From Hardy-Littlewood-Sobolev inequality, $\hat{v}_n \in L^6(\mathbb{R}^3)$ and

$$\|\hat{v}_n\|_6 \leq \int_{\mathbb{R}^3} \frac{u_n(y) u(y)}{|x-y|} dy \leq C \|u_n u\|_{\frac{6}{5}} \leq C \|u_n\|_{\frac{12}{5}} \|u\|_{\frac{12}{5}}.$$

So we get

$$\|\hat{v}_n u_n\|_2 \leq \|\hat{v}_n\|_6 \|u_n\|_3 \leq C \|u_n\|^2 \|u\| \leq C.$$

Therefore up to a subsequence $\hat{v}_n u_n \rightharpoonup v u$ in $L^2(\mathbb{R}^3)$. Due to $u \in L^2(\mathbb{R}^3)$ we get

$$\int_{\mathbb{R}^3} \hat{v}_n u_n u dx \rightarrow \int_{\mathbb{R}^3} v u^2 dx.$$

Therefore, we prove that $\lim_{n \rightarrow \infty} I_n^{(2)} = A$. A similar proof process can be deduced $\lim_{n \rightarrow \infty} I_n^{(3)} = A$ and $\lim_{n \rightarrow \infty} I_n^{(4)} = A$. Hence we get

$$B(u_n - u) + B(u) = B(u_n) + o(1).$$

From Brezis-Leib lemma can deduce

$$C(u_n - u) + C(u) = C(u_n) + o(1).$$

(ii) certified.

From Sobolev inequality and Gagliardo–Nirenberg inequality, we know

$$B(u_n) = \int_{\mathbb{R}^3} \phi_{u_n} u_n^2 dx \leq C \|u_n\|_{\frac{12}{5}}^4 \leq C \|u_n\|_2^3 \|\nabla u_n\|_2^3.$$

Due to Brezis-Leib lemma, we know

$$\|u_n - u\|_2^2 + \|u\|_2^2 = \|u_n\|_2^2 + o(1),$$

hence

$$\alpha_n = \frac{\rho^2 - \mu^2}{\|u_n - u\|_2} \rightarrow 1.$$

We can know

$$B(\alpha_n(u_n - u)) - B(u_n - u) = (\alpha_n^4 - 1)B(u_n - u) = o(1)$$

$$C(\alpha_n(u_n - u)) - C(u_n - u) = (\alpha_n^p - 1)C(u_n - u) = o(1).$$

(iii) certified. Due to u_n, u_m are minimizing sequence and Gagliardo–Nirenberg inequality we know

$$\|u_n - u_m\|_p \leq \|u_n - u_m\|_2^{\frac{6-p}{2p}} \|u_n - u_m\|_2^{\frac{3}{2}-3p} = o(1).$$

From Hölder inequality we deduce

$$\int_{\mathbb{R}^3} |u_n|^{p-1} |u_n - u| dx \leq \left(\int_{\mathbb{R}^3} |u_n|^p dx \right)^{\frac{1}{q}} \left(\int_{\mathbb{R}^3} |u_n - u|^p dx \right)^{\frac{1}{p}} = o(1),$$

where $q = \frac{p}{p-1}$. Hence, we know

$$\left| \int_{\mathbb{R}^3} (|u_n|^{p-1} - |u_m|^{p-1}) (u_n - u_m) dx \right| \leq C \|u_n - u_m\|_p = o(1).$$

We can obtain $\langle C'(u_n) - C'(u_m), u_n - u_m \rangle = o(1)$, then we need to verification B . From Hölder inequality we deduce

$$\begin{aligned} \int_{\mathbb{R}^3} \phi_{u_n} u_n (u_n - u_m) dx &\leq C \|\phi_{u_n}\|_6 \|u_n\|_2 \|u_n - u_m\|_3 \\ &\leq C \|u_n\|^2 \|u_n\|_2 \|u_n - u_m\|_3 \\ &= o(1). \end{aligned} \tag{2.4}$$

Hence, we get

$$\langle T'(u_n) - T'(u_m), u_n - u_m \rangle = o(1).$$

□

Lemma 2.9. *For every $\rho > 0$, $\{u_n\}$ be a minimizing sequence in B_ρ with $I_{\rho^2} < 0$, then $u_n \rightharpoonup u \neq 0$.*

Proof. Due to $\{u_n\}$ be a minimizing sequence in B_ρ for I_{ρ^2} and translation does not deform, we know for any sequence $\{y_n\} \subset \mathbb{R}^3$, $u(\cdot + y_n)$ is still a minimizing sequence for I_{ρ^2} . From Lions' lemma, if

$$\lim_{n \rightarrow \infty} \left(\sup_{y \in \mathbb{R}^3} \int_{B(y,1)} |u_n|^2 dx \right) = 0,$$

where $B(a, r) = \{x \in \mathbb{R}^3, |x - a| \leq r\}$. Then $u_n \rightarrow 0$ in $L^q(\mathbb{R}^3)$ for $q \in (2, 2^*)$, therefore $C(u_n) \rightarrow 0$. Moreover, From $I_{\rho^2} < 0$ we obtained

$$\sup_{y \in \mathbb{R}^3} \int_{B(y,1)} |u_n|^2 dx \geq \mu > 0,$$

hence we can choose $y_n \in \mathbb{R}^3$ such that

$$\int_{B(0,1)} |u_n(\cdot + y_n)|^2 dx \geq \mu > 0.$$

Since $H^1(B(0,1)) \hookrightarrow L^2(B(0,1))$ is a compact embedding, the weak limit of $u_n(\cdot + y_n)$ is nonzero. So it can be deduced that $u_n \rightharpoonup u \neq 0$. \square

3 Proof of the Theorem 1.1

Proof of Theorem 1.1. Now, proof Theorems 1.1 only need to prove that function I satisfies the strong subadditivity inequality, namely:

There exists $\rho_1 > 0$ such that $I_{\mu^2} < 0$ for all $\mu \in (\rho_1, +\infty)$ and

$$I_{\rho^2} < I_{\mu^2} + I_{\rho^2 - \mu^2},$$

for all $\rho > \rho_1$ and $0 < \mu < \rho$. We define $u_\theta(x) = \theta^{1-\frac{3}{2}\beta} u(\frac{x}{\theta^\beta})$, then from direct calculations we can get

$$\begin{aligned} \|u_\theta\|_2 &= \theta \|u\|_2, \\ A(u_\theta) &= \frac{1}{2} \int_{\mathbb{R}^3} |\nabla u_\theta|^2 dx = \theta^{2-2\beta} A(u), \\ B(u_\theta) &= \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{1 - e^{-|x-y|}}{|x-y|} u_\theta^2(x) u_\theta^2(y) dx dy, \\ &\leq \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{1}{|x-y|} u_\theta^2(x) u_\theta^2(y) dx dy = \theta^{4-\beta} H(u), \\ C(u_\theta) &= \theta^{(1-\frac{3}{2}\beta)p+3\beta} C(u), \end{aligned}$$

where $H(u) = \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{1}{|x-y|} u^2(x) u^2(y) dx dy$. We can also know

$$\begin{aligned} I(u_\theta) &= A(u_\theta) + B(u_\theta) + C(u_\theta) \\ &\leq \theta^{2-2\beta} A(u) + \theta^{4-\beta} H(u) + \theta^{(1-\frac{3}{2}\beta)p+3\beta} C(u) \\ &= \theta^2 \left(I(u) + (\theta^{-2\beta} - 1) A(u) + (\theta^{(1-\frac{3}{2}\beta)p+3\beta-2} - 1) C(u) + \theta^{2-\beta} H(u) - B(u) \right) \\ &= \theta^2 (I(u) + f(\theta, u)), \end{aligned}$$

where $f(\theta, u) = (\theta^{-2\beta} - 1)A(u) + (\theta^{(1-\frac{3}{2}\beta)p+3\beta-2} - 1)C(u) + \theta^{2-\beta}H(u) - B(u)$.

When $\beta = -2$, we get

$$I(u_\theta) \leq \theta^6 A(u) + \theta^6 H(u) + \theta^{4p-6} C(u),$$

and $4p - 6 > 6$ for $3 < p < \frac{10}{3}$. Hence, for θ sufficiently large we have $I(u_\theta) \rightarrow 0^-$. Let $\{u_n\}$ be a minimizing sequence in B_ρ with $I_{\rho^2} < 0$, then

$$0 < k_1 < A(u_n) < k_2,$$

$$0 < \eta_1 < |C(u_n)| < \eta_2.$$

Indeed, if $A(u_n) = o(1)$ we have $|C(u_n)| = o(1)$ and $I_{\rho^2} = 0$. For $\beta = -2$ we get

$$f(\theta, u) = (\theta^4 - 1)A(u) + (\theta^{4p-8} - 1)C(u) + \theta^4 H(u) - B(u),$$

with $4p - 8 > 4$ and

$$f'(\theta, u) |_{\theta=1} = 4A(u) + 4H(u) + (4p - 8)C(u) < k < 0,$$

$$f''(\theta, u) = 12\theta^2 A(u) + (4p - 8)(4p - 9)\theta^{4p-10}C(u) + 12\theta^2 H(u) < k < 0.$$

we know $f(1, u) = H(u) - B(u) > 0$ and $f(\theta, u_n) < k(\theta) < 0$ for any $\theta > 1$. From the mean value theorem, we can get there exists θ_0 such that if $\theta > \theta_0$ then $f(\theta, u) < 0$. hence

$$I_{\theta^2 \rho^2} < \theta^2 I(u_n) = \theta^2 I_{\rho^2}.$$

Let us suppose that $\mu^2 < \rho^2 - \mu^2$. We distinguish three cases

- $\mu^2 < \rho^2 - \mu^2 < \rho_1^2$
- $\mu^2 < \rho_1^2 < \rho^2 - \mu^2$
- $\rho_1^2 < \mu^2 < \rho^2 - \mu^2$.

The first case is trivial. For the second one, we know $I_{\rho^2 - \mu^2} > I_{\rho^2}$ and we conclude. For the third case

$$\begin{aligned} I_{\rho^2} &= I_{\frac{\rho^2}{\mu^2} \mu^2} < \frac{\rho^2}{\mu^2} I_{\mu^2} \\ &= \frac{\rho^2 - \mu^2 + \mu^2}{\mu^2} I_{\mu^2} \\ &= \frac{\rho^2 - \mu^2}{\mu^2} I_{\frac{\mu^2}{\rho^2 - \mu^2} \rho^2 - \mu^2} + I_{\mu^2} \leq I_{\mu^2} + I_{\rho^2 - \mu^2}. \end{aligned} \tag{3.1}$$

Since the strong subadditivity inequality condition holds, then combine Lemma 2.8 with Lemma 2.9, we can apply Proposition 2.3 and conclude the proof of Theorem 1.1. \square

4 Proof of the Theorem 1.2

When $p \in (2, 3)$ we apply Lemma 2.6 to proof strong subadditivity inequality. Hence the next step will be divided into several lemmas to verify all the assumptions in Lemma 2.6 in turn.

Lemma 4.1. $-\infty < I_{s^2} < 0$ for all $s > 0$ and condition (2.3) holds.

Proof. The detailed proof process for the establishment of condition (2.3) can be proved using the same way in Proposition 2.3 in [18]. From Lemma (2.7) we can get $I_{s^2} > -\infty$. Hence, we just need to proof $I_{s^2} < 0$. Let $g_u(\theta) = \theta^{1-\frac{3}{2}\beta}u(\frac{x}{\theta^\beta})$. From calculation, we obtained $\Theta_{g_u}(\theta) = \theta^2$ and $\|g_u(\theta)\|_2 = \theta$. By variable substitution

$$\begin{aligned} A(g_u(\theta)) &= \theta^{2-2\beta}A(u), \\ C(g_u(\theta)) &= \theta^{(1-\frac{3}{2}\beta)p+3\beta}C(u), \\ B(g_u(\theta)) &= \theta^{4-\beta} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{1 - e^{-\theta^\beta|x-y|}}{|x-y|} u(x)^2 u(y)^2 dx dy. \end{aligned}$$

Assume $\beta = -2$ we have

$$I(g_u(\theta)) = \frac{\theta^6}{2}A(u) + \frac{\theta^6}{4} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{1 - e^{-\theta^{-2}|x-y|}}{|x-y|} u(x)^2 u(y)^2 dx dy + \frac{\theta^{4p-6}}{p}C(u).$$

Due to $p \in (2, 3)$, we know $4p - 6 < 6$. Hence, $I(u_\theta) \rightarrow 0^-$ as $\theta \rightarrow 0$. Then there exists a small θ_0 such that

$$I_{s^2} < 0 \quad \forall s \in (0, \theta_0].$$

Let $\rho \in (\theta_0, \sqrt{2}\theta_0)$, for every $s \in (\theta_0, \rho)$ from adding conditions to get

$$I_{s^2} \leq I_{\theta_0^2} + I_{s^2 - \theta_0^2} < 0.$$

Due to $s^2 - \theta_0^2 < \theta_0^2$ we know $I_{s^2} < 0$ for s in the large interval $(0, \rho]$. Repeat the above process to get $I_{s^2} < 0$ for every $s > 0$. \square

Lemma 4.2. The function $s \mapsto I_{s^2}$ is continuous.

Proof. To prove this lemma just prove if $\rho_n \rightarrow \rho$ then $\lim_{n \rightarrow \infty} I_{\rho_n^2} = I_{\rho^2}$. Let $w_n \in B_{\rho_n}$ such that $I(w_n) < I_{\rho_n^2} + \frac{1}{n} < \frac{1}{n}$. Hence, from Gagliardo–Nirenberg inequality, Sobolev inequality and Lemma 2.5, we can deduce

$$\begin{aligned} \frac{1}{n} &> I(w_n) \\ &\geq \frac{1}{2} \|\nabla w_n\|_2^2 - \frac{1}{p} \|w_n\|_p^p \\ &\geq \frac{1}{2} \|\nabla w_n\|_2^2 - C \rho_n^{\frac{6-p}{2}} \|\nabla w_n\|_2^{\frac{3(p-2)}{2}}. \end{aligned}$$

Due to $\frac{3(p-2)}{2} < 2$ and $\{\rho_n\}$ is bounded sequence, Hence we know that $\{w_n\}$ is bounded in $H^1(\mathbb{R}^3)$. Moreover, $\{A(w_n)\}$ and $\{C(w_n)\}$ are bounded sequence. We also know

$$B(w_n) = \int_{\mathbb{R}^3} \phi_{w_n} |w_n|^2 dx \leq C \|w_n\|^4.$$

So $\{B(w_n)\}$ bounded in $H^1(\mathbb{R}^3)$. From simple calculation we can obtain

$$\begin{aligned}
I_{\rho^2} &\leq I\left(\frac{\rho}{\rho_n}w_n\right) \\
&= \frac{1}{2}A\left(\frac{\rho}{\rho_n}w_n\right) + \frac{1}{4}B\left(\frac{\rho}{\rho_n}w_n\right) + \frac{1}{p}C\left(\frac{\rho}{\rho_n}w_n\right) \\
&= \frac{1}{2}\left(\frac{\rho}{\rho_n}\right)^2 A(w_n) + \frac{1}{p}\left(\frac{\rho}{\rho_n}\right)^p + \frac{1}{4}\left(\frac{\rho}{\rho_n}\right)^5 \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{1 - e^{-\frac{\rho_n}{\rho}|x-y|}}{|x-y|} w_n^2(x) w_n^2(y) dx dy \\
&\leq I(w_n) + o(1) \\
&\leq I_{\rho_n^2} + o(1).
\end{aligned} \tag{4.1}$$

On the other hand, given a minimizing sequence $\{v_n\} \subset B_\rho$ for I_{ρ^2} we have

$$\begin{aligned}
I_{\rho_n^2} &\leq I\left(\frac{\rho_n}{\rho}v_n\right) \\
&= I(v_n) + o(1) \\
&= I_{\rho^2} + o(1).
\end{aligned} \tag{4.2}$$

According (4.1) and (4.2), we get $\lim_{n \rightarrow \infty} I_{\rho_n^2} = I_{\rho^2}$. \square

Lemma 4.3. $\lim_{s \rightarrow 0} \frac{I_{s^2}}{s^2} = 0$.

Proof. We let $G_{\rho^2} = \inf \left\{ \frac{1}{2} \|u\|_{D^{1,2}}^2 - \frac{1}{p} \int_{\mathbb{R}^3} |u|^p dx \right\}$. We know that $G_{\rho^2}/\rho^2 \leq I_{\rho^2}/\rho^2 < 0$. Since $G_{\rho^2}/\rho^2 \rightarrow 0$ (see Appendix A in [3]) we can easily conclude. \square

Lemma 4.4. For small ρ the function I satisfies (v) of proposition 2.3.

Proof. From Theorem 2.1 of [3], we know that under the conditional assumptions of proposition 2.6 we can get $M(\rho)$ is nonempty, where $M(\rho)$ is defined in Lemma 2.6. Moreover, since $0 \notin M(\rho)$, $A(u)$, $B(u)$ and $C(u)$ are different from zero whenever $u \in M(\rho)$. As we all know that the solution of problem (1.1) must satisfied the following Pohožev equality which was proved in [15],

$$0 = A(u) + 5B(u) + 3C(u) + \frac{1}{4} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} e^{-|x-y|} u^2(x) u^2(y) dx dy + \frac{3\omega}{2} \int_{\mathbb{R}^3} u^2 dx. \tag{4.3}$$

Moreover, when without the L^2 -norm constraint, the energy functional of problem 1.1 is

$$J(u) = A(u) + B(u) + C(u) + \frac{\omega}{2} \int_{\mathbb{R}^2} u^2 dx.$$

If u is the solution of problem 1.1 then $\langle J'(u), u \rangle = 0$ namely,

$$2A(u) + 4B(u) + pC(u) + \omega \int_{\mathbb{R}^3} |u|^2 dx = 0. \tag{4.4}$$

From (4.3) and (4.4), we claim that for any $u \in M(\rho)$ we get

$$0 = 2A(u) + B(u) - \frac{6-3p}{2}C(u) - \frac{1}{4} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} e^{-|x-y|} u^2(x) u^2(y) dx dy. \tag{4.5}$$

Now, for $u \neq 0$ we compute explicitly $h_{g_u}(\theta)$ by choosing the family of scaling paths of u with $\beta \in \mathbb{R}$ given by

$$G_u^\beta = \left\{ g_u(\theta) = \theta^{1-\frac{3}{2}\beta} u(x/\theta^\beta) \right\}.$$

All the paths of this family have as associated function $\Theta(\theta) = \theta^2$. We get

$$\begin{aligned} h_{g_u}(\theta) &= I(g_u(\theta)) - \theta^2 I(u) \\ &= (\theta^{2-2\beta} - \theta^2) A(u) + (\theta^{(1-\frac{3}{2}\beta)p+3\beta} - \theta^2) C(u) - \theta^2 B(u) \\ &\quad + \frac{1}{4} \theta^{4-\beta} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{1 - e^{-\theta^\beta |x-y|}}{|x-y|} u(x)^2 u(y)^2 dx dy, \end{aligned} \quad (4.6)$$

which show that h_{g_u} is differentiable for every $g_u \in G_u^\beta$. We have also, for $g_u \in G_u^\beta$:

$$\begin{aligned} h'_{g_u}(\theta) &= \left((2-2\beta)\theta^{2-2\beta-1} - 2\theta \right) A(u) - 2\theta B(u) + \left(\left[\left(1 - \frac{3}{2}\beta\right)p + 3\beta \right] \theta^{(1-\frac{3}{2}\beta)p+3\beta-1} - 2\theta \right) C(u) \\ &\quad + \frac{1}{4} (4-\beta) \theta^{4-\beta-1} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{1 - e^{-\theta^\beta |x-y|}}{|x-y|} u^2(x) u^2(y) dx dy \\ &\quad + \frac{1}{4} \theta^{4-\beta} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} e^{-\theta^\beta |x-y|} \beta \theta^{\beta-1} u^2(x) u^2(y) dx dy. \end{aligned} \quad (4.7)$$

Let $\theta = 1$, we get

$$\begin{aligned} h'_{g_u}(1) &= -2\beta A(u) + \left(\left(1 - \frac{3}{2}\beta\right)p + 3\beta - 2 \right) C(u) + (2-\beta) B(u) \\ &\quad + \frac{\beta}{4} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} e^{-|x-y|} u^2(x) u^2(y) dx dy. \end{aligned} \quad (4.8)$$

Assume there exists a sequence $\{u_n\} \subset M(\rho)$ with $\rho \geq \|u_n\|_2 = \rho_n \rightarrow 0$ such that for all $\beta \in \mathbb{R}$.

$$\begin{aligned} 0 &= h'_{g_{u_n}}(1) \\ &= -2\beta A(u_n) + \left(\left(1 - \frac{3}{2}\beta\right)p + 3\beta - 2 \right) C(u_n) + (2-\beta) B(u_n) \\ &\quad + \frac{\beta}{4} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} e^{-|x-y|} u_n^2(x) u_n^2(y) dx dy. \end{aligned} \quad (4.9)$$

From (4.5) and (4.9), we can get

$$2B(u_n) + (p-2)C(u_n) = 0. \quad (4.10)$$

and hence

$$\begin{aligned} B(u_n) &= A(u_n) - \frac{1}{2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} e^{-|x-y|} u_n^2(x) u_n^2(y) dx dy, \\ C(u_n) &= \frac{2}{2-p} A(u_n) - \frac{1}{2-p} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} e^{-|x-y|} u_n^2(x) u_n^2(y) dx dy, \\ I(u_n) &= \frac{6-2p}{2-p} A(u_n) - \frac{4-p}{2(2-p)} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} e^{-|x-y|} u_n^2(x) u_n^2(y) dx dy. \end{aligned} \quad (4.11)$$

Through direct calculation, we can know that $e^{-|x-y|}$ is a bounded function, then

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} e^{-|x-y|} u_n^2(x) u_n^2(y) dx dy \rightarrow 0,$$

so

$$\begin{cases} I(u_n) = I_{\rho_n^2} \rightarrow 0, \\ A(u_n), B(u_n), C(u_n) \rightarrow 0. \end{cases} \quad (4.12)$$

Notice the following Hardy-Littlewood-Sobolev inequality

$$B(u_n) \leq C \|u_n\|_{\frac{12}{5}}^4,$$

that we will frequently use.

Case 1: When $2 < p < \frac{12}{5}$.

Then

$$B(u_n) \leq C \|u_n\|_{\frac{12}{5}}^4 \leq C \|u_n\|_p^{4\alpha} \|u_n\|_6^{4(1-\alpha)}, \quad \alpha = \frac{3p}{2(6-p)}.$$

Thanks to (4.11) and Sobolev inequality $\|u_n\|_6^2 \leq 2SA(u_n)$, where S is the best constant for Sobolev embedding $D^{1,2}(\mathbb{R}^3) \hookrightarrow L^6(\mathbb{R}^3)$, namely $S := \inf_{u \in D^{1,2}(\mathbb{R}^3) \setminus \{0\}} \frac{\int_{\mathbb{R}^3} |\nabla u|^2 dx}{(\int_{\mathbb{R}^3} |u|^6 dx)^{\frac{1}{3}}}$. We get

$$B(u_n) \leq CB(u_n)^{\frac{4\alpha}{p}} \left(B(u_n) + \frac{1}{2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} e^{-|x-y|} u_n^2(x) u_n^2(y) dx dy \right)^{\frac{4(1-\alpha)}{2}}.$$

Due to $\frac{4\alpha}{p} + \frac{4(1-\alpha)}{2} = 3 - \frac{2p}{6-p} > 1$ for $2 < p < \frac{12}{5}$, hence this is a contradiction with (4.12).

Case 2: When $p = \frac{12}{5}$.

Due to (4.10) we obtain

$$\|u_n\|_{\frac{12}{5}}^{\frac{12}{5}} = CB(u_n) \leq C \|u_n\|_{\frac{12}{5}}^4,$$

which contradicts (4.12).

Case 3: When $\frac{12}{5} < p < \frac{8}{3}$.

Interpolating $L^{\frac{12}{5}}$ between L^2 and L^p we get

$$\|u_n\|_p^p = CB(u_n) \leq C \|u_n\|_{\frac{12}{5}}^4 \leq C \|u_n\|_2^{4\alpha} \|u_n\|_p^{4(1-\alpha)}, \quad \alpha = \frac{5p-12}{6(p-2)}.$$

Since $p < 4(1-\alpha)$, i.e. $p < \frac{8}{3}$, we get a contradiction with (4.12).

Case 4: When $p = \frac{8}{3}$.

Again by interpolation we get

$$B(u_n) \leq C \|u_n\|_{\frac{12}{5}}^4 \leq C \rho_n^{\frac{4}{3}} \|u_n\|_{\frac{8}{3}}^{\frac{8}{3}},$$

and again, using that $B(u_n) = C \|u_n\|_{\frac{8}{3}}^{\frac{8}{3}}$ we get a contradiction.

Case 5: When $\frac{8}{3} < p < 3$.

We first compute

$$\begin{aligned}
\frac{I(g_u(\theta))}{\theta^2 \|u\|_2^2} &= \frac{h_{g_u}(\theta)}{\theta^2 \|u\|_2^2} + \frac{I(u)}{\|u\|_2^2} \\
&= \frac{1}{\|u\|_2^2} \left\{ \theta^{-2\beta} A(u) + \frac{1}{4} \theta^{2-\beta} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{1 - e^{-\theta^\beta |x-y|}}{|x-y|} u^2(x) u^2(y) dx dy \right. \\
&\quad \left. + \theta^{(1-\frac{3}{2}\beta)p+3\beta-2} C(u) \right\}.
\end{aligned} \tag{4.13}$$

In this case for u_0 satisfying (4.11) with $\|u_0\|_2 = \rho_0$, hence we can get

$$\begin{aligned}
\frac{I_{\theta^2 \rho^2}}{\theta^2 \rho^2} &\leq \frac{I(g_u(\theta))}{\theta^2 \rho^2} \\
&\leq \frac{1}{\rho_0^2} \left(\theta^{-2\beta} A(u_0) + \frac{2}{2-p} \theta^{(1-\frac{3}{2}\beta)p+3\beta-2} A(u_0) + \frac{1}{4} \theta^{2-\beta} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{1}{|x-y|} u(x)^2 u(y)^2 dx dy \right) \\
&\leq \frac{1}{\rho_0^2} \left(\theta^{-2\beta} A(u_0) + \theta^{2-\beta} A(u_0) + \frac{2}{2-p} \theta^{(1-\frac{3}{2}\beta)p+3\beta-2} A(u_0) \right).
\end{aligned} \tag{4.14}$$

Now let us choose $\beta = \frac{2(2-p)}{10-3p}$ so that $0 < -2\beta = (1 - \frac{3}{2}\beta)p + \beta - 2 < 2 - \beta$. Hence we can obtain

$$\begin{aligned}
\frac{I_{\theta^2 \rho_0^2}}{\theta^2 \rho_0^2} &\leq \frac{I(g_{u_0}(\theta))}{\theta^2 \rho_0^2} = \frac{A(u_0)}{\rho_0^2} \left[\frac{4-p}{(2-p)} \theta^{\frac{4(p-2)}{10-3p}} + \theta^{\frac{4(4-p)}{10-3p}} \right] \\
&= \frac{\frac{4-p}{8(3-p)} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} e^{-|x-y|} u_0(x)^2 u_0(y)^2 dx dy}{\rho_0^2} \left[\frac{4-p}{2-p} \theta^{\frac{4(p-2)}{10-3p}} + \theta^{\frac{4(4-p)}{10-3p}} \right] \\
&\quad + \frac{\frac{2-p}{3-p} I(u_0)}{\rho_0^2} \left[\frac{4-p}{2-p} \theta^{\frac{4(p-2)}{10-3p}} + \theta^{\frac{4(4-p)}{10-3p}} \right].
\end{aligned} \tag{4.15}$$

Due to $\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} e^{-|x-y|} u_0(x)^2 u_0(y)^2 dx dy \rightarrow 0$ as s is enough small and $\theta^2 \rho_0^2 = s^2$, we get

$$\frac{I_{s^2}}{s^2} \leq -cs^{\frac{4(p-2)}{10-3p}} + o(s^{\frac{4(p-2)}{10-3p}}). \tag{4.16}$$

On the other hand for u_n satisfying (4.11)

$$\|u_n\|_p^p = CB(u_n) \leq C \|u_n\|_{\frac{12}{5}}^4 \leq C \|u_n\|_2^{4\alpha} \|u_n\|_p^{4(1-\alpha)}, \quad \alpha = \frac{5p-12}{6(p-2)},$$

that is

$$\|u_n\|_p^p \leq C \rho_n^{4\alpha} \|u_n\|_p^{4(1-\alpha)}. \tag{4.17}$$

we know that $\frac{8}{3} < p$ from (4.17) we get $\|u_n\|_p^p \leq C \rho_n^{\frac{4(p-2)}{3p-8}}$. Combine with (4.11) we obtain

$$\frac{I_{\rho_n^2}}{\rho_n^2} \geq -C \rho_n^{\frac{4(p-2)}{3p-8}}. \tag{4.18}$$

Combine (4.16) with (4.18) we obtain

$$-C \rho_n^{\frac{4(p-2)}{3p-8}} \leq \frac{I_{\rho_n^2}}{\rho_n^2} \leq -cs^{\frac{4(p-2)}{10-3p}} + o(s^{\frac{4(p-2)}{10-3p}}).$$

Since $\frac{4(p-2)}{3p-8} > \frac{4(p-2)}{10-3p}$, for $\rho_n \rightarrow 0$ we get a contradiction. \square

5 The orbital stability

Before proving Theorem 1.3, the definition of orbital stability is introduced. Let

$$S_\rho = \left\{ e^{i\theta} u(x) : \theta \in [0, 2\pi), \|u\|_2 = \rho, I(u) = I_{\rho^2} \right\}.$$

We call S_ρ is orbitally stable if for every $\varepsilon > 0$ there exists $\delta > 0$ such that for any $\psi_0 \in H^1(\mathbb{R}^3)$ with $\inf_{v \in S_\rho} \|v - \psi_0\|_{H^1(\mathbb{R}^3; \mathbb{C})} < \delta$ we have

$$\forall t > 0 \quad \inf_{v \in S_\rho} \|\psi(t, \cdot) - v\|_{H^1(\mathbb{R}^3; \mathbb{C})} < \varepsilon,$$

where $\psi(t, \cdot)$ is the solution of problem (1.1) with initial datum ψ_0 . It is worth noting that S_ρ is translational invariant, that is, when $v \in S_\rho$ then $v(\cdot - y) \in S_\rho$ for every $y \in \mathbb{R}^3$.

Proof of Theorem 1.3. Using the method of contradiction, it is assumed that there exists a $\rho > 0$ such that S_ρ is not orbitally stable. Therefore, there exist $\varepsilon > 0$ and a sequence of initial data $\{\psi_n, 0\} \subset H^1(\mathbb{R}^3)$ and $\{t_n\} \subset \mathbb{R}$, such that the maximal solution ψ_n , which is global and $\psi_n(0, \cdot) = \psi_{n,0}$, satisfies

$$\lim_{n \rightarrow +\infty} \inf_{v \in S_\rho} \|\psi_{n,0} - v\|_{H^1(\mathbb{R}^3)} = 0,$$

and

$$\inf_{v \in S_\rho} \|\psi_n(t_n, \cdot) - v\|_{H^1(\mathbb{R}^3)} \geq \varepsilon.$$

Therefore, there exists $u_\rho \in H^1(\mathbb{R}^3)$ which is a minimizer of I_{ρ^2} and $\theta \in \mathbb{R}$ such that $v = e^{i\theta} u_\rho$ and

$$\|\psi_{n,0}\|_2 \rightarrow \|v\|_2 = \rho \quad I(\psi_{n,0}) \rightarrow I(v) = I_{\rho^2}.$$

On the other hand, we can assume that $\psi_{n,0} \in B_\rho$. Then $\{\psi_{n,0}\}$ is a minimizing sequence for I_{ρ^2} , and due to

$$I(\psi_n(\cdot, t_n)) = I(\psi_{n,0}).$$

So $\{\psi_n(\cdot, t_n)\}$ is a minimizing sequence for I_{ρ^2} . Since we have that every minimizing sequence has a subsequence converging in $H^1(\mathbb{R}^3)$ to a minimum on the sphere B_ρ , we have a contradiction. \square

References

- [1] Danilo G Afonso and Gaetano Siciliano. Normalized solutions to a schrödinger–bopp–podolsky system under neumann boundary conditions. *Communications in Contemporary Mathematics*, page 2150100, 2021.
- [2] Jacopo Bellazzini, Louis Jeanjean, and Tingjian Luo. Existence and instability of standing waves with prescribed norm for a class of Schrödinger-Poisson equations. *Proc. Lond. Math. Soc. (3)*, 107(2):303–339, 2013.
- [3] Jacopo Bellazzini and Gaetano Siciliano. Scaling properties of functionals and existence of constrained minimizers. *J. Funct. Anal.*, 261(9):2486–2507, 2011.

- [4] Jacopo Bellazzini and Gaetano Siciliano. Stable standing waves for a class of nonlinear Schrödinger-Poisson equations. *Z. Angew. Math. Phys.*, 62(2):267–280, 2011.
- [5] H. Berestycki and P.-L. Lions. Nonlinear scalar field equations. II. Existence of infinitely many solutions. *Arch. Rational Mech. Anal.*, 82(4):347–375, 1983.
- [6] Fritz Bopp. Eine lineare Theorie des Elektrons. *Ann. Physik (5)*, 38:345–384, 1940.
- [7] Max Born. Modified field equations with a finite radius of the electron. *Nature*, 132(3329):282–282, 1933.
- [8] Max Born and Leopold Infeld. Foundations of the new field theory. *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character*, 144(852):425–451, 1934.
- [9] T. Cazenave and P.-L. Lions. Orbital stability of standing waves for some nonlinear Schrödinger equations. *Comm. Math. Phys.*, 85(4):549–561, 1982.
- [10] Sitong Chen and Xianhua Tang. On the critical Schrödinger-Bopp-Podolsky system with general nonlinearities. *Nonlinear Anal.*, 195:111734, 25, 2020.
- [11] Sitong Chen, Xianhua Tang, and Shuai Yuan. Normalized solutions for Schrödinger-Poisson equations with general nonlinearities. *J. Math. Anal. Appl.*, 481(1):123447, 24, 2020.
- [12] Pietro d’Avenia and Gaetano Siciliano. Nonlinear Schrödinger equation in the Bopp-Podolsky electrodynamics: solutions in the electrostatic case. *J. Differential Equations*, 267(2):1025–1065, 2019.
- [13] Louis Jeanjean and Thanh Trung Le. Multiple normalized solutions for a Sobolev critical Schrödinger-Poisson-Slater equation. *J. Differential Equations*, 303:277–325, 2021.
- [14] Suteo Kanesawa and Sin-itiro Tomonaga. On a relativistically invariant formulation of the quantum theory of wave fields. V. Case of interacting electromagnetic and meson fields. *Progr. Theoret. Phys.*, 3:101–113, 1948.
- [15] Lin Li, Patrizia Pucci, and Xianhua Tang. Ground state solutions for the nonlinear Schrödinger-Bopp-Podolsky system with critical Sobolev exponent. *Adv. Nonlinear Stud.*, 20(3):511–538, 2020.
- [16] Nikolaos S. Papageorgiou, Vicențiu D. Rădulescu, and Dušan D. Repovš. *Nonlinear analysis—theory and methods*. Springer Monographs in Mathematics. Springer, Cham, 2019.
- [17] Boris Podolsky. A generalized electrodynamics. I. Nonquantum. *Phys. Rev. (2)*, 62:68–71, 1942.
- [18] Óscar Sánchez and Juan Soler. Long-time dynamics of the Schrödinger-Poisson-Slater system. *J. Statist. Phys.*, 114(1-2):179–204, 2004.

- [19] Gaetano Siciliano and Kaye Silva. The fibering method approach for a non-linear Schrödinger equation coupled with the electromagnetic field. *Publ. Mat.*, 64(2):373–390, 2020.
- [20] Weihong Xie, Haibo Chen, and Hongxia Shi. Existence and multiplicity of normalized solutions for a class of Schrödinger-Poisson equations with general nonlinearities. *Math. Methods Appl. Sci.*, 43(6):3602–3616, 2020.
- [21] Jie Yang, Haibo Chen, and Senli Liu. The existence of nontrivial solution of a class of Schrödinger-Bopp-Podolsky system with critical growth. *Bound. Value Probl.*, pages Paper No. 144, 16, 2020.
- [22] Hongyu Ye. The existence and the concentration behavior of normalized solutions for the L^2 -critical Schrödinger-Poisson system. *Comput. Math. Appl.*, 74(2):266–280, 2017.
- [23] Hongyu Ye and Tingjian Luo. On the mass concentration of L^2 -constrained minimizers for a class of Schrödinger-Poisson equations. *Z. Angew. Math. Phys.*, 69(3):Paper No. 66, 13, 2018.
- [24] Yuting Zhu, Chunfang Chen, and Jianhua Chen. The Schrödinger-Bopp-Podolsky equation under the effect of nonlinearities. *Bull. Malays. Math. Sci. Soc.*, 44(2):953–980, 2021.