The TCFHs of D=11 AdS backgrounds and hidden symmetries

G. Papadopoulos and E. Pérez-Bolaños

Department of Mathematics King's College London Strand London WC2R 2LS, UK

george.papadopoulos@kcl.ac.uk
edgar.perez_bolanos@kcl.ac.uk

Abstract

We present the TCFHs of all AdS backgrounds of 11-dimensional supergravity and explore some of the properties of the associated connections. Therefore we demonstrate that all Killing spinor bilinears satisfy a generalisation of the conformal Killing-Yano equations with respect to the TCFH connection. In addition we explore the hidden symmetries of spinning particle probes propagating on these backgrounds. We give several examples of hidden symmetries for probes on the maximal supersymmetric AdS backgrounds as well as on some AdS backgrounds that arise as near horizon geometries of intersecting M-branes.

1 Introduction

There are two methods to solve the Killing spinor equations (KSEs) of a supergravity theory, the bilinears method [1] and the spinorial geometry method [2]; for a review see [3]. In the latter method the KSEs are solved directly using the (gauge) symmetries of supergravity theories and a realisation of spinors in terms of forms. While in the former method one solves the equations that the KSEs impose on the spacetime form bilinears (Dirac currents) of the Killing spinors. Originally these equations on the form bilinears were not thought to have a geometric interpretation but more recently it was realised that they can be arranged as a twisted covariant form hierarchy (TCFH) [4, 5]. This means that there is a connection $\mathcal{D}^{\mathcal{F}}$ on the space of spacetime forms which depends on the form fluxes \mathcal{F} of the supergravity theory under consideration such that schematically

$$\mathcal{D}_X^F \Omega = i_X \mathcal{P} + X \wedge \mathcal{Q} , \qquad (1)$$

for every spacetime vector field¹ X, where Ω is a multi-form spanned by the form bilinears, and \mathcal{P} and \mathcal{Q} are also multi-forms that depend on the supergravity fields as well as the form bilinears. The connection $\mathcal{D}^{\mathcal{F}}$ is not necessarily form degree preserving. One can also demonstrate that all theories with local supersymmetry, which may include higher order curvature corrections, on spacetimes of any signature exhibit a TCFH. The presence of a TCFH in a gravity theory with supersymmetry depends on the structure of the supercovariant connection, and so on the form of gravitino KSE, instead on the field dependence and the couplings of the theory.

A consequence of the TCFH, and equation (1) in particular, is that the form bilinears satisfy a generalisation of the conformal Killing-Yano (CKY) equation² with respect to the connection $\mathcal{D}^{\mathcal{F}}$. It is well known for sometime that Killing-Yano (KY) forms are associated with constants of motion for relativistic particles propagating on a spacetime [6, 7]. Key applications include the integrability of the geodesic motion as well as that of some classic field equations in a variety of black holes, see e.g. [8]-[14] and also reviews [15, 16]. For some other applications, see also [17, 18, 19, 20, 21]. In addition, KY forms generate symmetries [22] in spinning particle actions [23]. As a consequence it is natural to ask whether (1) gives rise to KY forms which in turn will generate symmetries in various spinning particle probes propagating in supersymmetric backgrounds. This question has been investigated in several supergravity theories in [24, 25] and several examples have been presented.

The purpose of this paper is to give the TCFHs on the internal space of all warped AdS backgrounds of 11-dimensional supergravity theory. This will put the conditions on the form bilinears implied by the KSEs in a firm geometric basis. For this we shall use the solution of the (gravitino) KSE of the theory along the AdS subspace of a background presented in [26]. Then we shall explore some of the properties of the TCFH connections which include their (reduced) holonomy for generic internal spaces. In addition we shall

¹We have also denoted with X the 1-form dual of X constructed as X(Y) = g(X, Y), where g is the spacetime metric and Y a vector field on the spacetime.

²A k-form ω on a n-dimensional manifold is CKY, iff $\nabla_X \omega = i_X d\omega - (n-k+1)^{-1} X \wedge \delta \omega$, where ∇ is the Levi-Civita connection. In addition, if the form ω is co-closed, $\delta \omega = 0$, then ω is Killing-Yano. Whereas if $d\omega = 0$, then ω is closed CKY (CCKY).

examine the conditions under which the form bilinears give rise to KY or CCKY forms³. The existence of such forms will imply in turn the presence of constants of motion in the propagation of spinning particles on the internal spaces of such backgrounds. We shall present several backgrounds with KY forms arising from the TCFH of their internal space. These include the maximally supersymmetric AdS solutions of the theory as well as the near horizon geometries of some intersecting M-brane configurations.

This paper is organised as follows. In sections 2, 3 and 4, we present the TCFH of warped AdS_2 , AdS_3 and AdS_4 backgrounds of 11-dimensional supergravity, respectively, and investigate some of the properties of the TCFH connections. In section 5, we give the TCFHs and investigate their properties of the remaining AdS backgrounds. In section 6, we explore the hidden symmetries of probes that arise from the TCFH of some solutions that include the maximally supersymmetric AdS backgrounds as well as some AdS backgrounds which are the near horizon geometries of intersecting M-branes, and in section 7 we give our conclusions.

2 The TCFH of warped AdS_2 backgrounds

2.1 Fields and Killing spinors

The bosonic fields of 11-dimensional supergravity for warped AdS_2 backgrounds, $AdS_2 \times_w M^9$, can be written as

$$g = 2\mathbf{e}^+\mathbf{e}^- + \delta_{ij}\mathbf{e}^i\mathbf{e}^j , \quad F = \mathbf{e}^+ \wedge \mathbf{e}^- \wedge Y + X , \qquad (2)$$

where Y and X are a 2-form and 4-form on the internal space M^9 with metric $g(M^9) = \delta_{ij} \mathbf{e}^i \mathbf{e}^j$, respectively,

$$\mathbf{e}^{+} = du$$
, $\mathbf{e}^{-} = dr + rh - \frac{1}{2}r^{2}\ell^{-2}A^{-2}du$, $\mathbf{e}^{i} = e^{i}{}_{J}dy^{J}$, (3)

is a null pseudo-orthonormal frame on the spacetime with \mathbf{e}^i an orthonormal frame on the internal space M^9 , $h = -d \log A^2$ and A is the warp factor which is a function on M^9 . The y^I are coordinates on M^9 and (u, r) are the remaining coordinates on the spacetime. It can be seen after a change of coordinates that the above metric can be rewritten in the standard warped product form, $g = A^2 g_\ell (AdS_2) + g(M^9)$, where $g_\ell (AdS_2)$ is the standard metric on AdS₂ of radius ℓ .

The KSE of 11-dimensional supergravity can be solved along the AdS₂ subspace of AdS₂ ×_w M^9 [26] and the Killing spinors ϵ can expressed as $\epsilon = \epsilon_1 + \epsilon_2$ with

$$\epsilon_1 = \phi_- + u\Gamma_+\Theta_-\phi_- + ru\Gamma_-\Theta_+\Gamma_+\Theta_-\phi_- , \quad \epsilon_2 = \phi_+ + r\Gamma_-\Theta_+\phi_+ . \tag{4}$$

where ϕ_{\pm} are spinors that depend only on the coordinates of M^9 and satisfy the light-cone projections $\Gamma_{\pm}\phi_{\pm} = 0$, where Γ_{\pm} have been adapted to the frame⁴ (3) and Θ_{\pm} are Clifford algebra elements that depend on the fields. For the explicit expressions of Θ_{\pm} as well

³The Hodge dual of a CCKY form is a KY form.

⁴From here on, the gamma matrices are always adapted to a (pseudo-)orthonormal frame.

as for the spinor notation we use below, see [26]. The dependence of Killing spinors in (4) on the (u, r) coordinates is explicit as it is that of the fields in (2). In addition, the (spacetime) gravitino KSE implies that ϕ_{\pm} satisfy the KSEs

$$\mathcal{D}_i^{\pm}\phi_{\pm} = 0 , \quad \mathcal{D}_i^{\pm} \equiv \nabla_i + \Psi_i^{\pm} , \qquad (5)$$

on the internal space M^9 , where ∇ is the Levi-Civita connection of the metric $g(M^9)$ induced on the spin bundle and

$$\Psi_i^{\pm} = \mp \frac{1}{4} h_i - \frac{1}{288} \Gamma X_i + \frac{1}{36} X_i \pm \frac{1}{24} \Gamma Y_i \mp \frac{1}{6} Y_i .$$
(6)

The TCFHs on M^9 that we shall explore below are associated with the supercovariant connections \mathcal{D}^{\pm} .

2.2 The TCFH on the internal space

A basis in the space of form bilinears on the internal space M^9 is

$$f^{\pm r,s} = \langle \phi^r_{\pm}, \phi^s_{\pm} \rangle , \quad k^{\pm r,s} = \langle \phi^r_{\pm}, \Gamma_i \phi^s_{\pm} \rangle \mathbf{e}^i ,$$

$$\theta^{\pm r,s} = \frac{1}{4!} \langle \phi^r_{\pm}, \Gamma_{ijk\ell} \phi^s_{\pm} \rangle \mathbf{e}^i \wedge \mathbf{e}^j \wedge \mathbf{e}^k \wedge \mathbf{e}^\ell , \quad \omega^{\pm r,s} = \frac{1}{2} \langle \phi^r_{\pm}, \Gamma_{ij} \phi^s_{\pm} \rangle \mathbf{e}^i \wedge \mathbf{e}^j ,$$

$$\varphi^{\pm r,s} = \frac{1}{3!} \langle \phi^r_{\pm}, \Gamma_{ijk} \phi^s_{\pm} \rangle \mathbf{e}^i \wedge \mathbf{e}^j \wedge \mathbf{e}^k , \qquad (7)$$

where all (time-like) space-like gamma matrices are (anti-)Hermitian with respect to the inner product $\langle \cdot, \cdot \rangle$. As the Killing spinors of 11-dimensional supergravity are real note that the bilinears f, k and θ are symmetric in the exchange of spinors ϕ_{\pm}^{r} and ϕ_{\pm}^{s} and the rest are skew-symmetric.

The TCFH associated with the supercovariant connections \mathcal{D}^{\pm} on the bilinears that are symmetric in the exchange of ϕ^r_{\pm} with ϕ^s_{\pm} is

$$\mathcal{D}_{i}^{(\pm)\mathcal{F}}f^{\pm} \equiv \nabla_{i}f^{\pm} \pm \partial_{i}\log A f^{\pm} = \pm \frac{1}{144} * X_{i\ell_{1}\ell_{2}\ell_{3}\ell_{4}}\theta^{\pm\ell_{1}\ell_{2}\ell_{3}\ell_{4}} \pm \frac{1}{3}Y_{i\ell}k^{\pm\ell} ,$$

$$\mathcal{D}_{i}^{(\pm)\mathcal{F}}k_{j}^{\pm} \equiv \nabla_{i}k_{j}^{\pm} \pm \partial_{i}\log A k_{j}^{\pm} - \frac{1}{12}X_{i\ell_{1}\ell_{2}\ell_{3}}\theta^{\pm\ell_{1}\ell_{2}\ell_{3}}j$$

$$= \frac{1}{18}X_{\ell_{1}\ell_{2}\ell_{3}[i}\theta^{\pm\ell_{1}\ell_{2}\ell_{3}}_{j]} + \frac{1}{144}\delta_{ij}X_{\ell_{1}\ell_{2}\ell_{3}\ell_{4}}\theta^{\pm\ell_{1}\ell_{2}\ell_{3}\ell_{4}} \pm \frac{1}{3}Y_{ij}f^{\pm} \pm \frac{1}{12}Y_{\ell_{1}\ell_{2}}\theta^{\pm\ell_{1}\ell_{2}}_{j]},$$

$$\mathcal{D}_{i}^{(\pm)\mathcal{F}}\theta_{j_{1}j_{2}j_{3}j_{4}}^{\pm} \equiv \nabla_{i}\theta_{j_{1}j_{2}j_{3}j_{4}}^{\pm} \pm \partial_{i}\log A \theta_{j_{1}j_{2}j_{3}j_{4}}^{\pm} \pm \frac{3}{2} * X_{\ell_{1}\ell_{2}i[j_{1}j_{2}}\theta^{\pm\ell_{1}\ell_{2}}_{j]}$$

$$+2X_{i[j_{1}j_{2}j_{3}}k_{j_{4}}^{\pm}] + \frac{1}{3} * Y_{\ell_{1}\ell_{2}\ell_{3}i[j_{1}j_{2}j_{3}}\theta^{\pm\ell_{1}\ell_{2}\ell_{3}}_{j_{4}}]$$

$$= \pm \frac{5}{3} * X_{\ell_{1}\ell_{2}[i_{1}j_{2}}\theta^{\pm\ell_{1}\ell_{2}}_{j_{3}j_{4}}] + \frac{5}{6} X_{[ij_{1}j_{2}j_{3}}k_{j_{4}}^{\pm}] \pm \frac{1}{6} * X_{ij_{1}j_{2}j_{3}j_{4}}f^{\pm}$$

$$\mp \frac{2}{3}\delta_{i[j_{1}} * X_{j_{2}j_{3}|k_{1}k_{2}k_{3}}\theta^{\pm\ell_{1}\ell_{2}\ell_{3}}_{j_{4}}] + \frac{2}{3}\delta_{i[j_{1}}X_{j_{2}j_{3}j_{4}]}\ell^{k_{2}\ell_{3}}\ell^{k_{2}\ell_{4}} - \frac{1}{18}\delta_{i[j_{1}} * Y_{j_{2}j_{3}j_{4}]\ell_{1}\ell_{2}\ell_{3}\ell_{4}}\theta^{\pm\ell_{1}\ell_{2}\ell_{3}\ell_{4}}$$

$$+ \frac{5}{18} * Y_{[ij_{1}j_{2}j_{3}|\ell_{1}\ell_{2}\ell_{3}]}\theta^{\pm\ell_{1}\ell_{2}\ell_{3}}_{j_{4}}] \mp 2\delta_{i[j_{1}}Y_{j_{2}j_{3}}k_{j_{4}}^{\pm}] .$$
(8)

While the TCFH on the bilinears which are skew-symmetric in the exchange of ϕ^r_\pm with ϕ^s_\pm is

$$\mathcal{D}_{i}^{(\pm)\mathcal{F}}\omega_{j_{1}j_{2}}^{\pm} \equiv \nabla_{i}\omega_{j_{1}j_{2}}^{\pm} \pm \partial_{i}\log A\,\omega_{j_{1}j_{2}}^{\pm} - \frac{1}{2}X_{\ell_{1}\ell_{2}i[j_{1}}\varphi^{\pm\ell_{1}\ell_{2}}{}_{j_{2}]} \mp \frac{1}{2}Y_{i\ell}\varphi^{\pm\ell_{1}}{}_{j_{1}j_{2}}$$

$$= -\frac{1}{4} X_{\ell_{1}\ell_{2}[ij_{1}} \varphi^{\pm \ell_{1}\ell_{2}}{}_{j_{2}]} \mp \frac{1}{12} * X_{ij_{1}j_{2}\ell_{1}\ell_{2}} \omega^{\pm \ell_{1}\ell_{2}} - \frac{1}{18} \delta_{i[j_{1}} X_{j_{2}]\ell_{1}\ell_{2}\ell_{3}} \varphi^{\pm \ell_{1}\ell_{2}\ell_{3}}$$

$$\pm \frac{1}{6} \delta_{i[j_{1}} \varphi^{\pm \ell_{1}\ell_{2}}{}_{j_{2}]} Y_{\ell_{1}\ell_{2}} \pm \frac{1}{2} Y_{\ell[i} \varphi^{\pm \ell}{}_{j_{1}j_{2}]} ,$$

$$\mathcal{D}_{i}^{(\pm)\mathcal{F}} \varphi_{j_{1}j_{2}j_{3}}^{\pm} \equiv \nabla_{i} \varphi_{j_{1}j_{2}j_{3}}^{\pm} \pm \partial_{i} \log A \varphi_{j_{1}j_{2}j_{3}}^{\pm} \pm \frac{3}{4} * X_{\ell_{1}\ell_{2}i[j_{1}j_{2}} \varphi^{\pm \ell_{1}\ell_{2}}{}_{j_{3}]}$$

$$- \frac{3}{2} X_{\ell i[j_{1}j_{2}} \omega^{\pm \ell}{}_{j_{3}]} \mp \frac{3}{2} Y_{i[j_{1}} \omega_{j_{2}j_{3}]}^{\pm}$$

$$= \pm \frac{2}{3} * X_{\ell_{1}\ell_{2}[ij_{1}j_{2}} \varphi^{\pm \ell_{1}\ell_{2}}{}_{j_{3}]} - \frac{2}{3} X_{\ell[ij_{1}j_{2}} \omega^{\pm \ell}{}_{j_{3}]} \pm \frac{1}{6} \delta_{i[j_{1}}} * X_{j_{2}j_{3}]\ell_{1}\ell_{2}\ell_{3}} \varphi^{\pm \ell_{1}\ell_{2}\ell_{3}}$$

$$- \frac{1}{4} \delta_{i[j_{1}} X_{j_{2}j_{3}]\ell_{1}\ell_{2}} \omega^{\pm \ell_{1}\ell_{2}} - \frac{1}{36} * Y_{ij_{1}j_{2}j_{3}\ell_{1}\ell_{2}\ell_{3}} \varphi^{\pm \ell_{1}\ell_{2}\ell_{3}} \mp Y_{[ij_{1}} \omega_{j_{2}j_{3}]}^{\pm} \mp \delta_{i[j_{1}} Y_{j_{2}|\ell|} \omega^{\pm \ell}{}_{j_{3}]} , \quad (9)$$

where X is the Hodge dual⁵ of X and similarly for the other fields. In the TCFH above we have suppressed the r, s indices on the bilinears that count the number of Killing spinors.

To compute (8) and (9), one utilises the identity

$$\nabla_i \langle \phi^r_{\pm}, \Gamma_{i_1 \dots i_k} \phi^s_{\pm} \rangle = \langle \nabla_i \phi^r_{\pm}, \Gamma_{i_1 \dots i_k} \phi^s_{\pm} \rangle + \langle \phi^r_{\pm}, \Gamma_{i_1 \dots i_k} \nabla_i \phi^s_{\pm} \rangle
= -\langle \Psi^{\pm}_i \phi^r_{\pm}, \Gamma_{i_1 \dots i_k} \phi^s_{\pm} \rangle - \langle \phi^r_{\pm}, \Gamma_{i_1 \dots i_k} \Psi^{\pm}_i \phi^s_{\pm} \rangle ,$$
(10)

where the last equality follows after a use the KSEs (5). Then one proceed with some extensive Clifford algebra computation.

The TCFH above has been expressed in terms of the minimal connection $\mathcal{D}^{(\pm)\mathcal{F}}$, see [5] for the definition. As expected $\mathcal{D}^{(\pm)\mathcal{F}}$ is form degree non-preserving connection. As the action of $\mathcal{D}_i^{(\pm)\mathcal{F}}$ on the space of forms preserves the subspaces of symmetric and skew-symmetric bilinears in the exchange of ϕ^r and ϕ^s Killing spinors and acts trivially on the scalars f, the reduced holonomy⁶ of the connection is included in (the connected to identity component of) $GL(135) \times GL(120)$. Note that the holonomy of the maximal TCFH connection, see again [5], is contained in $GL(136) \times GL(120)$ as it acts non-trivially on the scalars.

An alternative way to see that the holonomy of $\mathcal{D}^{(\pm)\mathcal{F}}$ is included in $GL(135) \times GL(120)$ is to observe that ϕ_{\pm} can be thought of as Majorana spinors of $\mathfrak{spin}(9)$. It is well known that the tensor product of two $\mathfrak{spin}(9)$ Majorana representations, Δ_{16} , decomposes as

$$\Delta_{\mathbf{16}} \otimes \Delta_{\mathbf{16}} = \bigoplus_{k=0}^{4} \Lambda^k(\mathbb{R}^9) , \qquad (11)$$

where $\Lambda^k(\mathbb{R}^9)$ is the irreducible representation of $\mathfrak{spin}(9)$ on the space of k-degree forms on \mathbb{R}^9 . The action of the supercovariant connection on the tensor product of two spin bundles, i.e. on a bispinor, preserves the symmetric and skew symmetric subspaces. As the rank of the spin bundle is 16, these sub-bundles have rank 136 and 120, respectively. So the holonomy of all connections of the TCFH is included in $GL(136) \times GL(120)$.

Although the form bilinears are CKY forms with respect to the TCFH connection $\mathcal{D}^{(\pm)\mathcal{F}}$ as expected, it is clear from the TCFH (8) and (9) that they are neither KY nor CCKY forms for generic supersymmetric backgrounds. However we shall demonstrate that for special solutions several terms in the TCFH vanish and as a result some bilinears become either KY or CCKY forms.

⁵In our conventions $\Gamma_{i_1...i_9}\phi_{\pm} = \pm \epsilon_{i_1...i_9}\phi_{\pm}$ with $\epsilon^{123456789} = 1$.

⁶From here on with the term holonomy we shall always refer to the reduced holonomy of the TCFH connection, i.e. the connected to the identity part of the holonomy group.

3 The TCFH of warped AdS_3 backgrounds

3.1 Fields and Killing spinors

The bosonic fields of 11-dimensional supergravity for a warped AdS_3 background, $AdS_3 \times_w M^8$, can be written as

$$g = 2\mathbf{e}^{+}\mathbf{e}^{-} + (\mathbf{e}^{z})^{2} + \delta_{ij}\mathbf{e}^{i}\mathbf{e}^{j} , \quad F = \mathbf{e}^{+} \wedge \mathbf{e}^{-} \wedge dz \wedge Q + X , \qquad (12)$$

where the metric, $g(M^8)$, on the internal space M^8 is $g(M^8) = \delta_{ij} \mathbf{e}^i \mathbf{e}^j$ and Q and X are a 1-form and a 4-form on M^8 , respectively. Moreover

$$\mathbf{e}^{+} = du$$
, $\mathbf{e}^{-} = dr - 2\ell^{-1}rdz - 2rd\ln A$, $\mathbf{e}^{z} = Adz$, $\mathbf{e}^{i} = e_{J}^{i}dy^{J}$, (13)

is a null pseudo-orthonormal frame on the spacetime with \mathbf{e}^i an orthonormal frame on M^8 , y are coordinates on M^8 and (u, r, z) are the rest of coordinates of spacetime and A is the warp factor. As for the AdS₂ backgrounds in the previous section, there is a coordinate transformation such that the spacetime metric g can be put into the standard warped form, $g = A^2 g_{\ell}(AdS_3) + g(M^8)$, where $g_{\ell}(AdS_3)$ is the standard metric on AdS₃ with radius ℓ .

The gravitino KSE of 11-dimensional supergravity can be solved [26] along the AdS₃ subspace of a AdS₃ ×_w M^8 background with fields (12). The Killing spinors ϵ can be expressed as, $\epsilon = \epsilon(\sigma_{\pm}) + \epsilon(\tau_{\pm})$, with

$$\begin{aligned} \epsilon(\sigma_{\pm}) &= \sigma_{+} + \sigma_{-} - \ell^{-1} A^{-1} u \Gamma_{+z} \sigma_{-} , \\ \epsilon(\tau_{\pm}) &= e^{-\frac{z}{\ell}} \tau_{+} - \ell^{-1} A^{-1} r e^{-\frac{z}{\ell}} \Gamma_{-z} \tau_{+} + e^{\frac{z}{\ell}} \tau_{-} , \end{aligned} \tag{14}$$

where σ_{\pm} and τ_{\pm} spinors satisfy the lightcone projections $\Gamma_{\pm}\sigma_{\pm} = \Gamma_{\pm}\tau_{\pm} = 0$, and depend only on the coordinates of M^8 . In addition, they satisfy the KSEs

$$\mathcal{D}_i^{\pm} \sigma_{\pm} = \mathcal{D}_i^{\pm} \tau_{\pm} = 0 , \qquad (15)$$

on the internal space M^8 , where

$$\mathcal{D}_{i}^{\pm} \equiv \nabla_{i} \pm \frac{1}{2} \partial_{i} \log A - \frac{1}{288} \Pi X_{i} + \frac{1}{36} X_{i} \mp \frac{1}{12} A^{-1} \Gamma_{z} \Pi Q_{i} \pm \frac{1}{6} A^{-1} \Gamma_{z} Q_{i} , \qquad (16)$$

are the supercovariant connections on M^8 and ∇ is the connection induced on the spin bundle from the Levi-Civita connection of the metric $g(M^8)$. Furthermore σ_{\pm} and τ_{\pm} satisfy an additional algebraic KSE on M^8 arising from the integration of the gravitino KSE of 11-dimensional supergravity along the z coordinate. These algebraic KSEs have been explained in detail in [26] and they will be used in the examples below to produce the right counting for the number of Killing spinors of the AdS backgrounds but they do not contribute in the TCFH below.

3.2 The TCFH on the internal space

A basis in the space of form bilinear on the internal space M^8 is

$$f^{\pm r,s} = \langle \phi^r_{\pm}, \phi^s_{\pm} \rangle , \quad \tilde{f}^{\pm r,s} = \langle \phi^r_{\pm}, \Gamma_z \phi^s_{\pm} \rangle , \quad k^{\pm r,s} = \langle \phi^r_{\pm}, \Gamma_i \phi^s_{\pm} \rangle \mathbf{e}^i ,$$

$$\theta^{\pm r,s} = \frac{1}{4!} \langle \phi^r_{\pm}, \Gamma_{ijk\ell} \phi^s_{\pm} \rangle \mathbf{e}^i \wedge \mathbf{e}^j \wedge \mathbf{e}^k \wedge \mathbf{e}^\ell , \quad \tilde{\varphi}^{\pm r,s} = \frac{1}{3!} \langle \phi^r_{\pm}, \Gamma_{ijk} \Gamma_z \phi^s_{\pm} \rangle \mathbf{e}^i \wedge \mathbf{e}^j \wedge \mathbf{e}^k ,$$

$$\tilde{k}^{\pm r,s} = \langle \phi^r_{\pm}, \Gamma_i \Gamma_z \phi^s_{\pm} \rangle \mathbf{e}^i , \quad \omega^{\pm r,s} = \frac{1}{2} \langle \phi^r_{\pm}, \Gamma_{ij} \phi^s_{\pm} \rangle \mathbf{e}^i \wedge \mathbf{e}^j ,$$

$$\tilde{\omega}^{\pm r,s} = \frac{1}{2} \langle \phi^r_{\pm}, \Gamma_{ij} \Gamma_z \phi^s_{\pm} \rangle \mathbf{e}^i \wedge \mathbf{e}^j , \quad \varphi^{\pm r,s} = \frac{1}{3!} \langle \phi^r_{\pm}, \Gamma_{ijk} \phi^s_{\pm} \rangle \mathbf{e}^i \wedge \mathbf{e}^j \wedge \mathbf{e}^k , \quad (17)$$

where ϕ_{\pm} stands⁷ for either σ_{\pm} or τ_{\pm} . The first five form bilinears are symmetric in the exchange of ϕ_{\pm}^r with ϕ_{\pm}^s while the rest are skew-symmetric. The TCFH expressed⁸ in terms of the minimal connection is

$$\begin{split} \mathcal{D}_{i}^{(\pm)\mathcal{F}} f^{\pm} &\equiv \nabla_{i} f^{\pm} \pm \partial_{i} \log A f^{\pm} = \pm \frac{1}{36} * X_{i}^{\ell_{1}\ell_{2}\ell_{3}} \tilde{\varphi}_{\ell_{1}\ell_{2}\ell_{3}}^{\pm} \mp \frac{1}{3} A^{-1} Q_{i} \tilde{f}^{\pm} , \\ \mathcal{D}_{i}^{(\pm)\mathcal{F}} k_{j}^{\pm} &\equiv \nabla_{i} k_{j}^{\pm} \pm \partial_{i} \log A k_{j}^{\pm} - \frac{1}{12} X_{i\ell_{1}\ell_{2}\ell_{3}} \theta^{\pm\ell_{1}\ell_{2}\ell_{3}} g^{\pm\ell_{1}\ell_{2}\ell_{3}} \\ &= -\frac{1}{18} X_{[i|\ell_{1}\ell_{2}\ell_{3}]} \theta^{\pm\ell_{1}\ell_{2}\ell_{3}} g^{\pm\ell_{1}\ell_{2}\ell_{3}} d^{\pm\ell_{1}\ell_{2}\ell_{3}} q^{\pm\ell_{1}\ell_{2}\ell_{3}} g^{\pm\ell_{1}\ell_{2}\ell_{3}} g^{\pm\ell_{1}\ell_{2}\ell_{3}} g^{\pm\ell_{1}\ell_{2}\ell_{3}} g^{\pm\ell_{1}\ell_{2}\ell_{3}} g^{\pm\ell_{1}\ell_{2}\ell_{3}} k_{j_{1}}^{\pm} \\ &+ \frac{1}{3} A^{-1} * Q_{\ell_{1}\ell_{2}\ell_{3}\ell_{1}(j_{1}j_{2}j_{3}} \theta^{\pm\ell_{1}\ell_{2}\ell_{3}} g^{\pm\ell_{1}\ell_{2}\ell_{3}} g^{\pm\ell$$

⁷One can also consider mixed σ_{\pm} and τ_{\pm} form bilinears. The TCFH is the same as the one stated below for the form bilinear basis above.

⁸In our conventions $\Gamma_z \Gamma_{i_1...i_8} \phi_{\pm} = \pm \epsilon_{i_1...i_8} \phi_{\pm}$ with $\epsilon^{12345678} = 1$.

$$= \mp \frac{1}{2} * X_{\ell_1 \ell_2 [ij_1} \varphi^{\pm \ell_1 \ell_2}{}_{j_2]} - \frac{1}{3} X_{ij_1 j_2 \ell} \tilde{k}^{\pm \ell} \mp \frac{1}{9} \delta_{i[j_1} * X_{j_2] \ell_1 \ell_2 \ell_3} \varphi^{\pm \ell_1 \ell_2 \ell_3} \pm \frac{1}{2} A^{-1} Q_{[i} \omega_{j_1 j_2]}^{\pm} \\ \pm \frac{1}{3} A^{-1} \delta_{i[j_1} \omega^{\pm \ell}{}_{j_2]} Q_{\ell} , \qquad (18)$$

where *X is the Hodge dual of X on the internal space M^8 and similarly for the other fields. We have also suppressed the r, s indices on the form bilinears that count the number of Killing spinors.

The action of the minimal TCFH connection on the space of forms preserves the subspaces of forms with are symmetric and skew-symmetric in the exchange of the Killing spinors ϕ_{\pm}^r and ϕ_{\pm}^s . Furthermore, it preserves the subspaces of 1-forms \tilde{k}^{\pm} , where it acts as the Levi-Civita connection up to a rescalling with the warp factor A, and acts trivially on the scalars f^{\pm} and \tilde{f}^{\pm} . Therefore the reduced holonomy is included in $GL(134) \times SO(8) \times GL(112)$. The reduced holonomy of the maximal TCFH connection instead is included in $GL(136) \times GL(120)$ because it preserves only the subspaces of forms which are symmetric and skew-symmetric in the exchange of the Killing spinors ϕ_{\pm}^r and ϕ_{\pm}^s .

The holonomy of the TCFH connection can be understood in a way similar to that of AdS_2 backgrounds. As ϕ_{\pm} can be viewed as Majorana $\mathfrak{spin}(8)$ spinors, it is known that the product of two such Majorana representations, Δ_{16} , can be decomposed in terms of form representations, $\Lambda^k(\mathbb{R}^8)$, of $\mathfrak{spin}(8)$ as

$$\Delta_{\mathbf{16}} \otimes \Delta_{\mathbf{16}} = \bigoplus_{k=0}^{8} \Lambda^{k}(\mathbb{R}^{8}) .$$
⁽¹⁹⁾

As the supercovariant derivative preserves the space of symmetric and skew-symmetric bi-spinors, it is clear that the holonomy of all TCFH connections will be included in $GL(136) \times GL(120)$, where 136 is the rank of the sub-bundle of symmetric bi-spinors while 120 is the rank of the sub-bundle of skew-symmetric bi-spinors.

As expected all form bilinears are CKY forms with respect to the TCFH connections $\mathcal{D}^{(\pm)\mathcal{F}}$ in agreement with the general result in [5]. Apart from $A^{\pm 1}\tilde{k}^{\pm}$ which is a Killing 1-form, the TCFH does not imply that the remaining form bilinears are KY forms for generic supersymmetric backgrounds. However, we shall demonstrate that many of them are either KY or CCKY forms for some AdS₃ solutions of 11-dimensional supergravity.

4 The TCFH of warped AdS_4 backgrounds

4.1 Fields and Killing spinors

The bosonic fields of 11-dimensional supergravity of warped AdS_4 backgrounds, $AdS_4 \times_w M^7$, can be written as

$$g = 2\mathbf{e}^{+}\mathbf{e}^{-} + (\mathbf{e}^{z})^{2} + (\mathbf{e}^{x})^{2} + g(M^{7}) , \quad F = S \,\mathbf{e}^{+} \wedge \mathbf{e}^{-} \wedge \mathbf{e}^{z} \wedge \mathbf{e}^{x} + X , \quad (20)$$

with $g(M^7) = \delta_{ij} \mathbf{e}^i \mathbf{e}^j$, where S and X are a 0-form and 4-form on the internal space M^7 , respectively. In addition,

$$\mathbf{e}^{+} = du , \quad \mathbf{e}^{-} = dr - 2\ell^{-1}rdz - 2rd\ln A , \quad \mathbf{e}^{z} = Adz , \quad \mathbf{e}^{x} = Ae^{\frac{z}{\ell}}dx ,$$
$$\mathbf{e}^{i} = e^{i}{}_{J}dy^{J} , \qquad (21)$$

is a null pseudo-orthonormal frame on the spacetime with \mathbf{e}^i an orthonormal frame on M^7 , y are coordinates on M^7 and (u, r, z, x) are the remaining coordinates of the spacetime, and ℓ is the radius of AdS₄. As in previous cases, after a coordinate transformation the spacetime metric can be written in the standard warped form with warp factor A^2 .

The gravitino KSE of 11-dimensional supergravity can be explicitly integrated along the (u, r, z, x) coordinates and the Killing spinors can be written as, $\epsilon = \epsilon(\sigma_{\pm}) + \epsilon(\tau_{\pm})$, see [26], with

$$\epsilon(\sigma_{\pm}) = \sigma_{+} + \sigma_{-} - \ell^{-1} e^{\frac{z}{\ell}} x \Gamma_{xz} \sigma_{-} - \ell^{-1} A^{-1} u \Gamma_{+z} \sigma_{-} , \epsilon(\tau_{\pm}) = e^{-\frac{z}{\ell}} \tau_{+} - \ell^{-1} A^{-1} r e^{\frac{z}{\ell}} \Gamma_{-z} \tau_{+} - \ell^{-1} x \Gamma_{xz} \tau_{+} + e^{\frac{z}{\ell}} \tau_{-} ,$$
(22)

where the spinors σ_{\pm} and τ_{\pm} depend only on the coordinates of M^7 and satisfy the lightcone projections $\Gamma_{\pm}\sigma_{\pm} = 0$ and $\Gamma_{\pm}\tau_{\pm} = 0$. Furthermore, these spinors satisfy the KSEs

$$\mathcal{D}_i^{\pm}\sigma_{\pm} , \quad \mathcal{D}_i^{\pm}\tau_{\pm} = 0 , \qquad (23)$$

on the internal space M^7 , where the supercovariant connection is

$$\mathcal{D}_{i}^{\pm} \equiv \nabla_{i} \pm \frac{1}{2} \partial_{i} \log A - \frac{1}{288} \Gamma X_{i} + \frac{1}{36} X_{i} \pm \frac{1}{12} S \Gamma_{izx} , \qquad (24)$$

with ∇ induced by the Levi-Civita connection of the metric $g(M^7)$ on M^7 . The spinors σ_{\pm} and τ_{\pm} satisfy an additional algebraic KSE which arises from the integration of the gravitino KSE of 11-dimensional supergravity along the z coordinate. These algebraic KSEs can be found in [26] and they are essential for the correct counting of Killing spinors for warped AdS backgrounds.

4.2 The TCFH on the internal manifold

A basis in the space of form bilinears on the internal space M^7 is

$$f^{\pm r,s} = \langle \phi^r_{\pm}, \phi^s_{\pm} \rangle , \quad k^{\pm r,s} = \langle \phi^r_{\pm}, \Gamma_i \phi^s_{\pm} \rangle \mathbf{e}^i , \quad \tilde{\omega}^{\pm r,s} = \frac{1}{2} \langle \phi^r_{\pm}, \Gamma_{ij} \Gamma_{zx} \phi^s_{\pm} \rangle \mathbf{e}^i \wedge \mathbf{e}^j ,$$
$$\tilde{\varphi}^{\pm r,s} = \frac{1}{3!} \langle \phi^r_{\pm}, \Gamma_{ijk} \Gamma_{zx} \phi^s_{\pm} \rangle \mathbf{e}^i \wedge \mathbf{e}^j \wedge \mathbf{e}^k , \quad \tilde{k}^{\pm r,s} = \langle \phi^r_{\pm}, \Gamma_i \Gamma_{zx} \phi^s_{\pm} \rangle \mathbf{e}^i ,$$
$$\omega^{\pm r,s} = \frac{1}{2} \langle \phi^r_{\pm}, \Gamma_{ij} \phi^s_{\pm} \rangle \mathbf{e}^i \wedge \mathbf{e}^j , \quad \varphi^{\pm r,s} = \frac{1}{3!} \langle \phi^r_{\pm}, \Gamma_{ijk} \phi^s_{\pm} \rangle \mathbf{e}^i \wedge \mathbf{e}^k ,$$
$$\tilde{f}^{\pm r,s} = \langle \phi^r_{\pm}, \Gamma_{zx} \phi^s_{\pm} \rangle . \tag{25}$$

where ϕ_{\pm} stands⁹ for either σ_{\pm} or τ_{\pm} . Note that the bilinears $f, k \tilde{\omega}$ and $\tilde{\varphi}$ are symmetric in the exchange of spinors ϕ_{\pm}^r and ϕ_{\pm}^s while the rest are skew-symmetric.

The TCFH expressed in terms of the minimal connection can be written as¹⁰

$$\mathcal{D}_i^{(\pm)\mathcal{F}} f^{\pm} \equiv \nabla_i f^{\pm} \pm \partial_i \log A f^{\pm} = \pm \frac{1}{12} {}^* X_{i\ell_1\ell_2} \tilde{\omega}^{\pm \ell_1\ell_2} ,$$

⁹Unlike the case of warped AdS₃ backgrounds, the σ_{\pm} and τ_{\pm} Killing spinors of all warped AdS_k, k > 3, backgrounds are related with Clifford algebra operations.

¹⁰In our conventions $\Gamma_{zx}\Gamma_{i_1i_2i_3i_4i_5i_6i_7}\phi_{\pm} = \pm \epsilon_{i_1i_2i_3i_4i_5i_6i_7}\phi_{\pm}$ with $\epsilon^{1234567} = 1$

$$\begin{split} \mathcal{D}_{i}^{(\pm)F}k_{j}^{\pm} &\equiv \nabla_{i}k_{j}^{\pm} \pm \partial_{i}\log Ak_{j}^{\pm} \mp \frac{1}{4}*X_{i\ell_{\ell}2}\tilde{\varphi}^{\pm\ell_{\ell}2_{\ell}}_{j} \\ &= \pm \frac{1}{6}S\tilde{\omega}_{ij}^{\pm} \mp \frac{1}{3}*X_{\ell_{\ell}2[i}\tilde{\varphi}^{\pm\ell_{\ell}\ell_{2}}_{j}] \mp \frac{1}{18}\delta_{ij}*X_{\ell_{\ell}2\ell_{3}}\tilde{\varphi}^{\pm\ell_{\ell}\ell_{2}\ell_{3}} , \\ \mathcal{D}_{i}^{(\pm)F}\tilde{\omega}_{j_{1}j_{2}}^{\pm} &\equiv \nabla_{i}\tilde{\omega}_{j_{1}j_{2}}^{\pm} \pm \partial_{i}\log A\tilde{\omega}_{j_{1}j_{2}}^{\pm} - \frac{1}{2}X_{\ell_{\ell}\ell_{2}[i]j}\tilde{\varphi}^{\pm\ell_{\ell}\ell_{2}}_{j_{2}]} \\ &= -\frac{1}{4}X_{\ell_{1}\ell_{2}[ij)}\tilde{\varphi}^{\pm\ell_{\ell}\ell_{2}}_{j_{2}]} - \frac{1}{18}\delta_{i[j_{1}}X_{j_{2}]\ell_{\ell}\ell_{2}}\tilde{\varphi}^{\pm\ell_{\ell}\ell_{2}}_{j_{1}\ell_{2}j_{3}} \pm \frac{1}{6}*X_{ij_{1}j_{2}}f^{\pm} \mp \frac{1}{3}S\delta_{i[j_{1}}k_{j_{2}}^{\pm}] , \\ \mathcal{D}_{i}^{(\pm)F}\tilde{\varphi}_{j_{1}j_{2}j_{3}}^{\pm} \equiv \nabla_{i}\tilde{\varphi}_{j_{1}j_{2}j_{3}}^{\pm} \pm \partial_{i}\log A\tilde{\varphi}_{j_{1}j_{2}j_{3}}^{\pm} - \frac{3}{2}X_{\ell_{\ell}[j_{1}j_{2}}\tilde{\omega}^{\pm\ell}_{j_{3}}] \mp \frac{3}{2}*X_{i[j_{1}j_{2}}k_{j_{3}^{\pm}}^{\pm}] \\ &= \mp \frac{4}{3}*X_{[ij_{1}j_{2}}k_{j_{3}^{\pm}}^{\pm} - \frac{2}{3}X_{\ell_{l}[j_{1}j_{2}}\tilde{\omega}^{\pm\ell}_{j_{3}}] \mp \delta_{i[j_{1}}*X_{j_{2}j_{3}]}\ell^{k^{\pm\ell}} \\ - \frac{1}{4}\delta_{i[j_{1}}X_{j_{2}j_{3}]\ell_{\ell^{2}}}\tilde{\omega}^{\pm\ell_{\ell^{2}}}_{j_{1}j_{2}} = \nabla_{i}\tilde{\omega}_{j_{1}j_{2}}^{\pm} \pm \partial_{i}\log A\omega_{j_{1}j_{2}}^{\pm} - \frac{1}{2}X_{\ell_{\ell}\ell_{2}[j_{1}}\varphi^{\pm\ell_{\ell^{2}}}_{j_{2}}] \\ &= -\frac{1}{4}X_{\ell_{1}\ell_{2}[ij_{1}}\varphi^{\pm\ell_{\ell^{2}}}_{j_{2}}] - \frac{1}{18}\delta_{i[j_{1}}X_{j_{2}]\ell_{\ell^{2}}\delta\varphi}^{\pm\ell_{\ell^{2}}}\tilde{\varphi}^{\pm\ell_{\ell^{2}}}_{j_{2}} \\ &= -\frac{1}{4}X_{\ell_{1}\ell_{2}[ij_{1}}\varphi^{\pm\ell_{\ell^{2}}}_{j_{2}}] - \frac{1}{18}\delta_{i[j_{1}}X_{j_{2}]\ell_{\ell^{2}}\delta\varphi}^{\pm\ell_{\ell^{2}}}_{j_{2}} \mp \frac{1}{6}*X_{ij_{1}j_{2}}\tilde{f}^{\pm} \pm \frac{1}{3}S\delta_{i[j_{1}}\tilde{k}_{j_{2}}] , \\ \mathcal{D}_{i}^{(\pm)F}\varphi_{j_{1}j_{2}j_{3}}^{\pm} \equiv \nabla_{i}\varphi_{j_{1}j_{2}j_{3}}^{\pm} \pm \partial_{i}\log A\varphi_{j_{1}j_{2}j_{3}}^{\pm} - \frac{3}{2}X_{\ell_{\ell}[j_{1}j_{2}}\omega^{\pm\ell_{2}}] \pm \frac{1}{3}S\delta_{i[j_{1}}\tilde{k}_{j_{3}}] \\ &= \pm \frac{4}{3}*X_{[i_{1}j_{2}}\tilde{k}_{j_{3}}^{\pm}] - \frac{2}{3}X_{\ell_{\ell}[i_{1}j_{2}}\omega^{\pm\ell_{\ell^{2}}}_{j_{3}}] \pm \delta_{i}[j_{1}}*X_{j_{2}j_{3}]\ell\tilde{k}^{\pm\ell}} \\ - \frac{1}{4}\delta_{i[j_{1}}X_{j_{2}j_{3}]\ell_{\ell^{2}}\omega^{\pm\ell_{\ell^{2}}}_{j_{3}}] \pm \delta_{i}\log A\tilde{f}^{\pm} \pm \mp \frac{1}{12}*X_{\ell^{2}}\omega^{\pm\ell_{\ell^{2}}}_{j_{4}} , \\ \mathcal{D}_{i}^{(\pm)F}\tilde{f}^{\pm} \equiv \nabla_{i}\tilde{f}^{\pm} \pm \partial_{i}\log A\tilde{f}^{\pm} \pm \mp \frac{1}{12}}*X_{i\ell_{\ell^{2}}}\varphi^{\pm\ell_{\ell^{2}}}_{j_{4}} \\ \\$$

where *X and *S are the Hodge duals of X and S on the internal space M^7 , respectively, and we have suppress the r, s indices on the form bilinears that label the number of Killing spinors.

The action of the minimal TCFH connection on the space of forms preserves the subspaces of symmetric and skew-symmetric bilinears in the exchange of ϕ_{\pm}^{r} and ϕ_{\pm}^{s} Killing spinors and acts trivially on the scalars f and \tilde{f} . As a consequence, the holonomy of the connection is included in $GL(63) \times GL(63)$. Note that the TCFH on the 0-, 1- and 3-form bilinears which are symmetric in the exchange of ϕ_{\pm}^{r} and ϕ_{\pm}^{s} is almost identical to that of the corresponding form bilinears which are skew-symmetric in the exchange of the same spinors. The difference is a sign in the terms containing the fluxes S and *F. The holonomy of the maximal TCFH connection is included in $GL(64) \times GL(64)$ since it acts non-trivially on the scalars.

The spinors σ_{\pm} and τ_{\pm} are associated with the (reducible) Majorana representation Δ_{16} of $\mathfrak{spin}(8)$. This decomposes under $\mathfrak{spin}(7)$ as $\Delta_{16} = \Delta_8 \oplus \Delta_8$, where Δ_8 is the (irreducible) Majorana representation of $\mathfrak{spin}(7)$. Moreover the tensor product of two such representations, Δ_8 , decomposes in terms of form representations as

$$\Delta_{\mathbf{8}} \otimes \Delta_{\mathbf{8}} = \sum_{k=0}^{3} \Lambda^{k}(\mathbb{R}^{7}) \ . \tag{27}$$

Clearly the TCFH includes of two copies of forms that appear in the above decomposition which lead for the holonomy of the maximal connection of TCFH to be included in $GL(64) \times GL(64)$.

5 The TCFH of warped AdS_k , k = 5, 6, 7, backgrounds

5.1 Fields and Killing spinors

The fields of 11-dimensional supergravity for warped $\operatorname{AdS}_k \times_w M^{11-k}$, k = 5, 6, 7, backgrounds can be written as

$$g = 2\mathbf{e}^{+}\mathbf{e}^{-} + (\mathbf{e}^{z})^{2} + \sum_{a} (\mathbf{e}^{a})^{2} + g(M^{11-k}) , \quad F = X ,$$
 (28)

with $g(M^{11-k}) = \delta_{ij} \mathbf{e}^i \mathbf{e}^j$, where X is a 4-form on the internal space M^{11-k} . The null pseudo-orthonormal frame $(\mathbf{e}^+, \mathbf{e}^-, \mathbf{e}^z, \mathbf{e}^a, \mathbf{e}^i)$ is expressed as

$$\mathbf{e}^{+} = du$$
, $\mathbf{e}^{-} = dr - 2\ell^{-1}rdz - 2rd\ln A$, $\mathbf{e}^{z} = Adz$, $\mathbf{e}^{a} = Ae^{\frac{z}{\ell}}dx^{a}$, $\mathbf{e}^{i} = e^{i}_{J}dy^{j}$,(29)

where y are coordinates of the internal space M^{11-k} and (u, r, z, x^a) are the rest of the coordinates of the spacetime, ℓ is the radius of AdS subspace and A is the warp factor.

As in all previous cases, the KSEs of 11-dimensional supergravity can be integrated over the AdS subspace and the Killing spinors can be expressed as, $\epsilon = \epsilon(\sigma_{\pm}) + \epsilon(\tau_{\pm})$, see [26], with

$$\begin{aligned} \epsilon(\sigma_{\pm}) &= \sigma_{+} + \sigma_{-} - \ell^{-1} e^{\frac{z}{\ell}} x^{a} \Gamma_{az} \sigma_{-} - \ell^{-1} A^{-1} u \Gamma_{+z} \sigma_{-} ,\\ \epsilon(\tau_{\pm}) &= e^{-\frac{z}{\ell}} \tau_{+} - \ell^{-1} A^{-1} r e^{-\frac{z}{\ell}} \Gamma_{-z} \tau_{+} - \ell^{-1} x^{a} \Gamma_{az} \tau_{+} + e^{\frac{z}{\ell}} \tau_{-} , \end{aligned}$$
(30)

where the σ_{\pm} and τ_{\pm} spinors satisfy the lightcone projections $\Gamma_{\pm}\sigma_{\pm} = \Gamma_{\pm}\tau_{\pm} = 0$, and depend only on the coordinates of M^{11-k} . In addition, they satisfy the KSEs

$$\mathcal{D}_i^{\pm}\phi_{\pm} = 0 , \qquad (31)$$

along the internal space M^{11-k} , where the supercovariant connection is

$$\mathcal{D}_{i}^{\pm} \equiv \nabla_{i} \pm \frac{1}{2} \partial_{i} \log A - \frac{1}{288} \Pi X_{i} + \frac{1}{36} X_{i} , \qquad (32)$$

 ∇ is the connection on the spin bundle of M^{11-k} induced from the metric $g(M^{11-k})$ and ϕ_{\pm} stands from either σ_{\pm} or τ_{\pm} . Note that for warped AdS₇ backgrounds the term ΓX in the supercovariant connection vanishes. As in previous cases σ_{\pm} and τ_{\pm} satisfy an additional algebraic KSE which arises from the integration of the gravitino KSE of 11-dimensional supergravity along the z coordinate and can be found in [26]. It will be used to determine the number of Killing spinors in some examples below.

5.2 The TCFH of warped AdS_5 backgrounds

A basis in the space of form bilinears on the internal space M^6 is

$$f^{\pm r,s} = \langle \phi^r_{\pm}, \phi^s_{\pm} \rangle , \quad k^{\pm r,s} = \langle \phi^r_{\pm}, \Gamma_i \phi^s_{\pm} \rangle \mathbf{e}^i , \quad \tilde{k}^{\pm r,s} = \langle \phi^r_{\pm}, \Gamma_i \Gamma_{(3)} \phi^s_{\pm} \rangle \mathbf{e}^i ,$$
$$\tilde{\omega}^{\pm r,s} = \frac{1}{2} \langle \phi^r_{\pm}, \Gamma_{ij} \Gamma_{(3)} \phi^s_{\pm} \rangle \mathbf{e}^i \wedge \mathbf{e}^j , \quad \omega^{\pm r,s} = \frac{1}{2} \langle \phi^r_{\pm}, \Gamma_{ij} \phi^s_{\pm} \rangle \mathbf{e}^i \wedge \mathbf{e}^j ,$$
$$\varphi^{\pm r,s}_{ijk} = \frac{1}{3!} \langle \phi^r_{\pm}, \Gamma_{ijk} \phi^s_{\pm} \rangle \mathbf{e}^i \wedge \mathbf{e}^j \wedge \mathbf{e}^k , \quad \tilde{f}^{\pm r,s} = \langle \phi^r_{\pm}, \Gamma_{(3)} \phi^s_{\pm} \rangle , \qquad (33)$$

where $\Gamma_{(3)} = \Gamma_{zx^1x^2}$, i.e. it is the product of gamma matrices along the directions \mathbf{e}^z and \mathbf{e}^a for a = 1, 2. The first four bilinears are symmetric in the exchange of ϕ^r_{\pm} and ϕ^s_{\pm} spinors while the rest are skew-symmetric.

The TCFH expressed¹¹ in terms of the minimal connection is

$$\begin{aligned} \mathcal{D}_{i}^{(\pm)\mathcal{F}}f^{\pm} &\equiv \nabla_{i}f^{\pm} \pm \partial_{i}\log Af^{\pm} = \pm \frac{1}{6}^{*}X_{i\ell}\tilde{k}^{\pm\ell} ,\\ \mathcal{D}_{i}^{(\pm)\mathcal{F}}k_{j}^{\pm} &\equiv \nabla_{i}k_{j}^{\pm} \pm \partial_{i}\log Ak_{j}^{\pm} \pm \frac{1}{2}^{*}X_{i\ell}\tilde{\omega}^{\pm\ell}{}_{j} \\ &= \pm \frac{2}{3}^{*}X_{[i|\ell|}\tilde{\omega}^{\pm\ell}{}_{j]} \mp \frac{1}{6}\delta_{ij}^{*}X_{\ell_{1}\ell_{2}}\tilde{\omega}^{\pm\ell_{1}\ell_{2}} ,\\ \mathcal{D}_{i}^{(\pm)\mathcal{F}}\tilde{k}_{j}^{\pm} &\equiv \nabla_{i}\tilde{k}_{j}^{\pm} \pm \partial_{i}\log A\tilde{k}_{j}^{\pm} = \frac{1}{6}X_{ij\ell_{1}\ell_{2}}\tilde{\omega}^{\pm\ell_{1}\ell_{2}} \pm \frac{1}{6}^{*}X_{ij}f^{\pm} ,\\ \mathcal{D}_{i}^{(\pm)\mathcal{F}}\tilde{\omega}_{j_{1}j_{2}}^{\pm} &\equiv \nabla_{i}\tilde{\omega}_{j_{1}j_{2}}^{\pm} \pm \partial_{i}\log A\tilde{\omega}_{j_{1}j_{2}}^{\pm} \pm ^{*}X_{i[j_{1}}k_{j_{2}]} \\ &= -\frac{1}{3}X_{ij_{1}j_{2}\ell}\tilde{k}^{\pm\ell} \pm \frac{2}{3}\delta_{i[j_{1}}^{*}X_{j_{2}]\ell}k^{\pm\ell} \pm ^{*}X_{[ij_{1}}k_{j_{2}]} ,\\ \mathcal{D}_{i}^{(\pm)\mathcal{F}}\omega_{j_{1}j_{2}}^{\pm} &\equiv \nabla_{i}\omega_{j_{1}j_{2}}^{\pm} \pm \partial_{i}\log A\omega_{j_{1}j_{2}}^{\pm} - \frac{1}{2}X_{\ell_{1}\ell_{2}i[j_{1}}\varphi^{\pm\ell_{1}\ell_{2}} ,\\ &= -\frac{1}{4}X_{\ell_{1}\ell_{2}[ij_{1}}\varphi^{\pm\ell_{1}\ell_{2}} ,\\ \mathcal{D}_{i}^{(\pm)\mathcal{F}}\varphi_{\ell_{1}\ell_{2}\ell_{3}}^{\pm} &\equiv \nabla_{i}\varphi_{\ell_{1}\ell_{2}\ell_{3}}^{\pm} \pm \partial_{i}\log A\varphi_{\ell_{1}\ell_{2}\ell_{3}}^{\pm} - \frac{3}{2}X_{\ell_{i}[j_{1}j_{2}}\omega^{\pm\ell} ,\\ &= -\frac{1}{4}\delta_{i[j_{1}}X_{j_{2}j_{3}]\ell_{1}\ell_{2}}\omega^{\pm\ell_{1}\ell_{2}} - \frac{2}{3}X_{\ell_{i}[j_{1}j_{2}}\omega^{\pm\ell} ,\\ &= -\frac{1}{4}\delta_{i[j_{1}}X_{j_{2}j_{3}]\ell_{1}\ell_{2}}\omega^{\pm\ell_{1}\ell_{2}} - \frac{2}{3}X_{\ell_{i}[j_{1}j_{2}}\omega^{\pm\ell} ,\\ &= -\frac{1}{6}^{*}X_{\ell_{1}\ell_{2}}\varphi^{\pm\ell_{1}\ell_{2}} , \end{aligned}$$

where *X is the Hodge dual of X on M^6 and we have suppressed the indices r, s of the form bilinears.

As expected the minimal connection of the TCFH $\mathcal{D}^{(\pm)\mathcal{F}}$ is not form degree preserving. On the other hand its action closes on the form bilinears which are either symmetric or skew-symmetric in the interchange of spinors ϕ_{\pm}^r and ϕ_{\pm}^s . The holonomy of $\mathcal{D}^{(\pm)\mathcal{F}}$ is contained in $SO(6) \times GL(21) \times GL(35)$ as in addition it acts with the Levi-Civita connection on the 1-form bilinear $A \tilde{k}$ and trivially on the scalar bilinears f and \tilde{f} . Note that the holonomy of the maximal TCFH connection is contained in $GL(28) \times GL(35)$ as it only closes on the symmetric and skew-symmetric form bilinears under the exchange of spinors ϕ_{\pm}^r and ϕ_{\pm}^s .

¹¹In our conventions $\Gamma_{(3)}\Gamma_{i_1...i_6}\phi_{\pm} = \pm \epsilon_{i_1...i_6}\phi_{\pm}$ and $\epsilon^{123456} = 1$.

5.3 The TCFH of warped AdS_6 backgrounds

A basis in the space of form bilinears on the internal space M^5 is

$$f^{\pm r,s} = \langle \phi^r_{\pm}, \phi^s_{\pm} \rangle , \quad k^{\pm r,s} = \langle \phi^r_{\pm}, \Gamma_i \phi^s_{\pm} \rangle \mathbf{e}^i , \quad \tilde{f}^{\pm r,s} = \langle \phi^r_{\pm}, \Gamma_{(4)} \phi^s_{\pm} \rangle ,$$

$$\tilde{k}^{\pm r,s} = \langle \phi^r_{\pm}, \Gamma_i \Gamma_{(4)} \phi^s_{\pm} \rangle \mathbf{e}^i ,$$

$$\omega^{\pm r,s} = \frac{1}{2} \langle \phi^r_{\pm}, \Gamma_{ij} \phi^s_{\pm} \rangle \mathbf{e}^i \wedge \mathbf{e}^j , \quad \tilde{\omega}^{\pm r,s} = \frac{1}{2} \langle \phi^r_{\pm}, \Gamma_{ij} \Gamma_{(4)} \phi^s_{\pm} \rangle \mathbf{e}^i \wedge \mathbf{e}^j , \quad (35)$$

where the first four form bilinears are symmetric in the exchange of ϕ_{\pm}^r and ϕ_{\pm}^s spinors while the rest are skew-symmetric and $\Gamma_{(4)} = \Gamma_{zx^1x^2x^3}$.

The TCFH expressed¹² in terms of the minimal connection, $\mathcal{D}^{(\pm)\mathcal{F}}$, is

$$\mathcal{D}_{i}^{(\pm)\mathcal{F}}f^{\pm} \equiv \nabla_{i}f^{\pm} \pm \partial_{i}\log Af^{\pm} = \pm \frac{1}{6} {}^{*}X_{i}\tilde{f}^{\pm} ,$$

$$\mathcal{D}_{i}^{(\pm)\mathcal{F}}k_{j}^{\pm} \equiv \nabla_{i}k_{j}^{\pm} \pm \partial_{i}\log Ak_{j}^{\pm} \mp \frac{1}{2} {}^{*}X_{i}\tilde{k}_{j}^{\pm} = \mp \frac{2}{3} {}^{*}X_{[i}\tilde{k}_{j]}^{\pm} \mp \frac{1}{3}\delta_{ij} {}^{*}X_{\ell}\tilde{k}^{\pm\ell} ,$$

$$\mathcal{D}_{i}^{(\pm)\mathcal{F}}\tilde{f}^{\pm} \equiv \nabla_{i}\tilde{f}^{\pm} \pm \partial_{i}\log A\tilde{f}^{\pm} = \pm \frac{1}{6} {}^{*}X_{i}f^{\pm} ,$$

$$\mathcal{D}_{i}^{(\pm)\mathcal{F}}\tilde{k}_{j}^{\pm} \equiv \nabla_{i}\tilde{k}_{j}^{\pm} \pm \partial_{i}\log A\tilde{k}_{j}^{\pm} \mp \frac{1}{2} {}^{*}X_{i}k_{j}^{\pm} = \mp \frac{2}{3} {}^{*}X_{[i}k_{j]}^{\pm} \mp \frac{1}{3}\delta_{ij} {}^{*}X_{\ell}k^{\pm\ell} ,$$

$$\mathcal{D}_{i}^{(\pm)\mathcal{F}}\omega_{j_{1}j_{2}}^{\pm} \equiv \nabla_{i}\omega_{j_{1}j_{2}}^{\pm} \pm \partial_{i}\log A\omega_{j_{1}j_{2}}^{\pm} \mp \frac{1}{2} {}^{*}X_{i}\tilde{\omega}_{j_{1}j_{2}}^{\pm} = \mp {}^{*}X_{[i}\tilde{\omega}_{j_{1}j_{2}}^{\pm}] \mp \frac{2}{3}\delta_{i[j_{1}}\tilde{\omega}^{\pm\ell}{}_{j_{2}]} {}^{*}X_{\ell} ,$$

$$\mathcal{D}_{i}^{(\pm)\mathcal{F}}\tilde{\omega}_{j_{1}j_{2}}^{\pm} \equiv \nabla_{i}\tilde{\omega}_{j_{1}j_{2}}^{\pm} \pm \partial_{i}\log A\tilde{\omega}_{j_{1}j_{2}}^{\pm} \mp \frac{1}{2} {}^{*}X_{i}\omega_{j_{1}j_{2}}^{\pm} = \mp {}^{*}X_{[i}\omega_{j_{1}j_{2}}^{\pm}] \mp \frac{2}{3}\delta_{i[j_{1}}\omega^{\pm\ell}{}_{j_{2}]} {}^{*}X_{\ell} (36)$$

where *X is the Hodge dual of X and we have suppressed the r, s indices on the form bilinears as in previous cases.

Unlike previous cases, the minimal TCFH connection $\mathcal{D}^{(\pm)\mathcal{F}}$ for AdS₆ backgrounds is form degree preserving. Furthermore its action can be diagonalised on the forms

$$\zeta_{(+)}^{\pm} = \zeta^{\pm} + \tilde{\zeta}^{\pm} , \quad \zeta_{(-)}^{\pm} = \zeta^{\pm} - \tilde{\zeta}^{\pm} , \qquad (37)$$

where ζ^{\pm} stands for either k^{\pm} or ω^{\pm} , i.e. one has that

$$\mathcal{D}_{i}^{(\pm)\mathcal{F}}\zeta_{(+)}^{\pm} = \nabla_{i}\zeta_{(+)}^{\pm} \pm \partial_{i}\log A\zeta_{(+)}^{\pm} \mp \frac{1}{2} * X_{i}\zeta_{(+)}^{\pm} ,$$
$$\mathcal{D}_{i}^{(\pm)\mathcal{F}}\zeta_{(-)}^{\pm} = \nabla_{i}\zeta_{(-)}^{\pm} \pm \partial_{i}\log A\zeta_{(-)}^{\pm} \pm \frac{1}{2} * X_{i}\zeta_{(-)}^{\pm} .$$
(38)

Such a connection arises provided one gauges the scale transformation $\zeta \to s \zeta$ accompanied with $*X \to *X \pm 2s^{-1}ds$, where the sign is plus for $\zeta^+_{(+)}$ and $\zeta^-_{(-)}$ while it is minus for the rest of the form bilinears. Clearly, there are two sectors and the holonomy of the connection in each sector is $SO(5) \times (\mathbb{R} - \{0\})$.

5.4 The TCFH of AdS₇ backgrounds

A basis in the space of spinor bilinears on the internal space M^4 is

$$f^{\pm r,s} = \langle \phi^r_{\pm}, \phi^s_{\pm} \rangle , \quad k^{\pm r,s} = \langle \phi^r_{\pm}, \Gamma_i \phi^s_{\pm} \rangle \mathbf{e}^i , \quad \tilde{f}^{\pm r,s} = \langle \phi^r_{\pm}, \Gamma_{(5)} \phi^s_{\pm} \rangle ,$$

¹²In our conventions $\Gamma_{(4)}\Gamma_{i_1...i_5}\phi_{\pm} = \pm \epsilon_{i_1...i_5}\phi_{\pm}$ with $\epsilon^{12345} = 1$.

$$\tilde{k}^{\pm r,s} = \langle \phi_{\pm}^r, \Gamma_i \Gamma_{(5)} \phi_{\pm}^s \rangle \mathbf{e}^i , \quad \omega^{\pm r,s} = \frac{1}{2} \langle \phi_{\pm}^r, \Gamma_{ij} \phi_{\pm}^s \rangle \mathbf{e}^i \wedge \mathbf{e}^j , \qquad (39)$$

where the first three are symmetric in the exchange of spinors ϕ_{\pm}^r and ϕ_{\pm}^s while the rest are skew-symmetric and $\Gamma_{(5)} = \Gamma_{zx^1x^2x^3x^4}$.

The TCFH expressed¹³ in terms of the minimal connection $\mathcal{D}^{(\pm)\mathcal{F}}$ is

$$\mathcal{D}_{i}^{(\pm)\mathcal{F}}f^{\pm} \equiv \nabla_{i}f^{\pm} \pm \partial_{i}\log Af^{\pm} = 0 ,$$

$$\mathcal{D}_{i}^{(\pm)\mathcal{F}}k_{j}^{\pm} \equiv \nabla_{i}k_{j}^{\pm} \pm \partial_{i}\log Ak_{j}^{\pm} = \mp \frac{1}{3}\delta_{ij} * X\tilde{f}^{\pm} ,$$

$$\mathcal{D}_{i}^{(\pm)\mathcal{F}}\tilde{f}^{\pm} \equiv \nabla_{i}\tilde{f}^{\pm} \pm \partial_{i}\log A\bar{f}^{\pm} = \pm \frac{1}{3} * Xk_{i}^{\pm} ,$$

$$\mathcal{D}_{i}^{(\pm)\mathcal{F}}\omega_{j_{1}j_{2}}^{\pm} \equiv \nabla_{i}\omega_{j_{1}j_{2}}^{\pm} \pm \partial_{i}\log A\omega_{j_{1}j_{2}}^{\pm} = \pm \frac{2}{3}\delta_{i[j_{1}}\bar{k}_{j_{2}]}^{\pm} * X ,$$

$$\mathcal{D}_{i}^{(\pm)\mathcal{F}}\tilde{k}_{j}^{\pm} \equiv \nabla_{i}\tilde{k}_{j}^{\pm} \pm \partial_{i}\log A\tilde{k}_{j}^{\pm} = \mp \frac{1}{3} * X\omega_{ij}^{\pm} ,$$

(40)

where X is the Hodge dual of the 4-form X on the internal space M^4 .

It is clear that the (reduced) holonomy of $\mathcal{D}^{(\pm)\mathcal{F}}$ is contained in SO(4). Furthermore $A^{\pm 1}k^{\pm}$ and $A^{\pm 1}\omega^{\pm}$ are CCKY forms. Therefore their dual in M^4 are KY. In addition $A^{\pm 1}\tilde{k}^{\pm}$ is a KY tenors. It is well-known KY tensors generate symmetries in spinning particle actions.

6 Probes and symmetries

6.1 Relativistic and spinning particles

We have integrated the KSE of 11-dimensional supergravity along the AdS_k subspace of a warped spacetime, $AdS_k \times_w M^{11-k}$, and found the TCFHs on the internal space M^{11-k} . To investigate whether the form bilinears of the TCFHs on the internal space generate symmetries for spinning particle actions, we have to integrate the dynamics of the spinning particle along the AdS_k subspace and describe the effective dynamics of the system on the internal space M^{11-k} .

For this consider first the dynamics of a relativistic particle on a warped spacetime, $N \times_w M$, with metric $g = A^2h + \gamma$, where h is a metric on N and γ is a metric on M and A is the warped factor. Varying the action

$$A = \frac{1}{2} \int dt \, g_{\mu\nu} \dot{x}^{\mu} \dot{x}^{\nu} \,, \tag{41}$$

one finds that the equations of motion are

$$\nabla_t^h (A^2 \dot{\rho}^a) = 0 , \quad \nabla_t^\gamma \dot{y}^I - \frac{1}{2} \gamma^{IJ} \partial_J A^2 h_{ab} \dot{\rho}^a \dot{\rho}^b = 0 , \qquad (42)$$

¹³In our conventions $\Gamma_{(5)}\Gamma_{i_1...i_4}\phi_{\pm} = \pm \epsilon_{i_1...i_4}\phi_{\pm}$ and $\epsilon^{1234} = 1$.

where ∇^h and ∇^γ denote the Levi-Civita connections of h and γ , respectively, ρ^a are coordinates on N and y^I are coordinates on M. It is clear that

$$Q^{2} = \frac{1}{2} A^{4} h_{ab} \dot{\rho}^{a} \dot{\rho}^{b} , \qquad (43)$$

is conserved as a consequence of the field equation on N. Then notice that the dynamics of the relativistic particle on M can be described by the effective action

$$A_M = \frac{1}{2} \int dt \left(\frac{1}{2} \gamma_{IJ} \dot{y}^I \dot{y}^J - Q^2 A^{-2} \right) \,. \tag{44}$$

The action apart from the usual kinetic term exhibits a potential depending on the warped factor. There are various sectors to consider parameterised by the value of Q^2 . If either $Q^2 = 0$, which is the case for ρ constant, or A^2 is constant, A_M becomes the standard action for geodesic motion on M possibly shifted by an ignorable constant.

A similar analysis can be performed for a spinning particle probe [23] propagating on a spacetime with metric g described by the action

$$A = -\frac{i}{2} \int dt \, d\theta \, g_{\mu\nu} \, Dx^{\mu} \, \dot{x}^{\nu} \,, \qquad (45)$$

where (t, θ) are worldline superspace coordinates, x are worldline superfields $x = x(t, \theta)$ and D is a worldline superspace derivative with $D^2 = i\partial_t$.

The equations of motion of the spinning particle (45) propagating on a warped spacetime $N \times_w M$, as for the relativistic particle above, are

$$\nabla^h (A^2 \dot{\rho}^a) + \nabla^h_t (A^2 D \rho^a) = 0 , \quad \nabla^\gamma \dot{y}^I = \frac{1}{2} \gamma^{IJ} \partial_J A^2 h_{ab} D \rho^a \dot{\rho}^b .$$

$$\tag{46}$$

In this case, there is not a simple description of the effective dynamics on M as for the relativistic particle described by the action (44). However note that ρ equals to a constant is a solution of the equations of motion above. So if either ρ is constant or the warp factor A is constant, the effective dynamics of the spinning particle on M is described by the action

$$A_M = -\frac{i}{2} \int dt \, d\theta \, \gamma_{IJ} \, Dy^I \, \dot{y}^J \,. \tag{47}$$

It is well known that the action above is invariant under an infinitesimal transformation

$$\delta y^I = \epsilon \, \alpha^I{}_{J_1 \cdots J_{m-1}} D y^{J_1} \cdots D y^{J_{m-1}} , \qquad (48)$$

provided that α is a KY form [22], where ϵ is the infinitesimal parameter.

There is an extensive list of 11-dimensional supersymmetric AdS solutions, see e.g. [27, 28, 29, 30, 31, 32]. The purpose here is to give some examples of TCFHs and investigate their properties instead of being comprehensive. So we shall focus below on the TCFH of the maximally supersymmetric AdS solutions and some AdS solutions that arise as near horizon geometries of intersecting M-branes.

6.2 Maximally supersymmetric AdS backgrounds

6.2.1 $AdS_4 \times S^7$

The TCFH of warped AdS_4 backgrounds with only electric flux, i.e. X = 0 in (20), can be written as

$$\begin{aligned} \mathcal{D}_{i}^{(\pm)\mathcal{F}} f^{\pm} &\equiv \nabla_{i} f^{\pm} \pm \partial_{i} \log A f^{\pm} = 0 , \\ \mathcal{D}_{i}^{(\pm)\mathcal{F}} k_{j}^{\pm} &\equiv \nabla_{i} k_{j}^{\pm} \pm \partial_{i} \log A k_{j}^{\pm} = \pm \frac{1}{6} S \tilde{\omega}_{ij}^{\pm} , \\ \mathcal{D}_{i}^{(\pm)\mathcal{F}} \tilde{\omega}_{j_{1}j_{2}}^{\pm} &\equiv \nabla_{i} \tilde{\omega}_{j_{1}j_{2}}^{\pm} \pm \partial_{i} \log A \tilde{\omega}_{j_{1}j_{2}}^{\pm} = \mp \frac{1}{3} S \delta_{i[j_{1}} k_{j_{2}}^{\pm}] , \\ \mathcal{D}_{i}^{(\pm)\mathcal{F}} \tilde{\varphi}_{j_{1}j_{2}j_{3}}^{\pm} &\equiv \nabla_{i} \tilde{\varphi}_{j_{1}j_{2}j_{3}}^{\pm} \pm \partial_{i} \log A \tilde{\varphi}_{j_{1}j_{2}j_{3}}^{\pm} = -\frac{1}{36} * S_{ij_{1}j_{2}j_{3}\ell_{1}\ell_{2}\ell_{3}} \tilde{\varphi}^{\pm\ell_{1}\ell_{2}\ell_{3}} \\ \mathcal{D}_{i}^{(\pm)\mathcal{F}} \omega_{j_{1}j_{2}}^{\pm} &\equiv \nabla_{i} \omega_{j_{1}j_{2}}^{\pm} \pm \partial_{i} \log A \omega_{j_{1}j_{2}}^{\pm} = \pm \frac{1}{3} S \delta_{i[j_{1}} \tilde{k}_{j_{2}}^{\pm}] , \\ \mathcal{D}_{i}^{(\pm)\mathcal{F}} \varphi_{j_{1}j_{2}j_{3}}^{\pm} &\equiv \nabla_{i} \varphi_{j_{1}j_{2}j_{3}}^{\pm} \pm \partial_{i} \log A \varphi_{j_{1}j_{2}j_{3}}^{\pm} = -\frac{1}{36} * S_{ij_{1}j_{2}j_{3}\ell_{1}\ell_{2}\ell_{3}} \varphi^{\pm\ell_{1}\ell_{2}\ell_{3}} , \\ \mathcal{D}_{i}^{(\pm)\mathcal{F}} \tilde{f}^{\pm} &\equiv \nabla_{i} \tilde{f} \pm \partial_{i} \log A \tilde{f}^{\pm} = 0 , \\ \mathcal{D}_{i}^{(\pm)\mathcal{F}} \tilde{k}_{j}^{\pm} &\equiv \nabla_{i} \tilde{k}_{j}^{\pm} \pm \partial_{i} \log A \tilde{k}_{j}^{\pm} = \mp \frac{1}{6} S \omega_{ij}^{\pm} . \end{aligned}$$

It is clear from the TCFH above that $A^{\pm 1}k$, $A^{\pm 1}\tilde{k}$, $A^{\pm 1}\varphi$ and $A^{\pm 1}\tilde{\varphi}$ are KY forms, and $A^{\pm 1}\omega$ and $A^{\pm 1}\tilde{\omega}$ are CCKY forms. As $A^{\pm 1}\omega$ and $A^{\pm 1}\tilde{\omega}$ are CCKY forms their duals in the internal space are KY forms. The holonomy of the TCFH connection is included in SO(7).

The maximally supersymmetric AdS_4 solution is a Freund-Rubin type of background with internal space M^7 the round 7-sphere, $M^7 = S^7$, and the warp factor A constant. All the forms bilinears above generate symmetries for the spinning particle probe action (47). Note that the form bilinears on S^7 are not necessarily invariant forms under the SO(8) isometry group of S^7 .

6.2.2 $\operatorname{AdS}_7 \times S^4$

The maximally supersymmetric AdS_7 solution is again a Freund-Rubin type of background with internal space M^4 the round 4-sphere, $M^4 = S^4$, and the warp factor A constant. An inspection of the TCFH of warped AdS_7 backgrounds in (40) reveals that the bilinear \tilde{k}^{\pm} is a KY form, and k^{\pm} and ω^{\pm} are CCKY forms. Again the duals $*k^{\pm}$ and $*\omega^{\pm}$ of k^{\pm} and ω^{\pm} in S^4 , respectively, are KY forms and so \tilde{k}^{\pm} , $*k^{\pm}$ and $*\omega^{\pm}$ generate symmetries for the spinning particle probe action (47).

6.3 AdS backgrounds from intersecting branes

More examples of AdS backgrounds emerge as near horizon geometries of intersecting Mbranes [33]. We shall not explore all the possibilities, see [34] for more examples. Instead we shall focus on the AdS₂ and AdS₃ solutions that arise as near horizon geometries of the intersection of three M2-branes on an 0-brane, the intersection of an M2-brane and M5brane on a 1-brane and the intersection of three M5-branes on a 1-brane configurations.

6.3.1 AdS₂ solution from intersecting M2-branes

One could take the near horizon geometry of the three intersecting M2-brane solution on a 0-brane and proceed to examine the associated TCFH. Instead, we shall write an ansatz for the fields which includes the solution. In particular, we set

$$g = g_{\ell}(AdS_2) + g(S^3) + g(\mathbb{R}^6) , \quad F = d\operatorname{vol}_{\ell}(AdS_2) \wedge Y , \qquad (50)$$

where $g_{\ell}(AdS_2)$ and $dvol_{\ell}(AdS_2)$ is the standard metric and volume 2-form on AdS₂ with radius ℓ , respectively, $g(S^3)$ is the round metric on S^3 of unit radius, $g(\mathbb{R}^6)$ is the Euclidean metric on \mathbb{R}^6 and Y is a constant 2-form on \mathbb{R}^6 . Using the scale symmetry $g \to \Omega^2 g$ and $F \to \Omega^3 F$ of 11-dimensional supergravity as well as some coordinate transformations, one can show that the near horizon geometry of three M2-branes, with arbitrary charge densities, intersecting on a 0-brane solution can be cast into the above form. Clearly for this ansatz X = 0 in (2) and we have set A = 1.

Focusing on the ϕ_+ Killing spinors, the gravitino KSE along the directions of \mathbb{R}^6 implies that Y is a non-degenerate 2-form and proportional to a Kähler form λ on \mathbb{R}^6 associated with the Euclidean metric, i.e. $Y = \gamma \lambda$, $\gamma \in \mathbb{R}$. Furthermore ϕ_+ has to satisfy the conditions $\Gamma_{1234}\phi_+ = \eta\phi_+$ and $\Gamma_{1256}\phi_+ = \zeta\phi_+$, where $\eta, \zeta = \pm 1$ and we have arranged such that \mathbb{R}^6 lies in the directions $1, \ldots, 6$. Then the warp factor field equation in [26] implies that $\gamma^2 = \ell^{-2}$. Next the field equation along S^3 , which is the round unit sphere, gives $\ell^{-2} = 4$. All the remaining KSEs and field equations are satisfied without further conditions. Therefore the background (50) with the above choice of parameters admits 4 ϕ_+ Killing spinors. The solution also admits 4 more ϕ_- Killing spinor and so it preserves 1/4 of the supersymmetry.

Next notice that the supercovariant derivative along S^3 can be written as

$$\mathcal{D}^{\pm}_{\alpha} = \nabla^{S^3}_{\alpha} \pm \frac{1}{24} \Gamma_{\alpha} \not\!\!\! / \, , \qquad (51)$$

where α here labels the three orthonormal directions tangential to S^3 . Note that $\Gamma_{\alpha} \not Y = \not Y \Gamma_{\alpha}$. Moreover considering only those components of the form bilinears that lie on S^3 , i.e.

$$k_{\alpha}^{\pm rs} = \langle \phi_{\pm}^{r}, \Gamma_{\alpha} \phi_{\pm}^{s} \rangle , \quad \omega_{\alpha\beta}^{\pm rs} = \langle \phi_{\pm}^{r}, \Gamma_{\alpha\beta} \phi_{\pm}^{s} \rangle , \quad \varphi_{\alpha\beta\delta}^{\pm rs} = \langle \phi_{\pm}^{r}, \Gamma_{\alpha\beta\delta} \phi_{\pm}^{s} \rangle , \tag{52}$$

one can demonstrate that k and φ are KY forms while ω is CCKY form. Therefore all of them or their Hodge duals on S^3 generate symmetries for the probe action (47).

6.3.2 AdS₃ solution from M2- and M5-branes

An ansatz which includes the near horizon geometry of an M2-brane intersecting an M5brane on a 1-brane is

$$g = g_{\ell}(AdS_3) + g(S^3) + g(\mathbb{R}^5) , \quad F = d\operatorname{vol}_{\ell}(AdS_3) \wedge Q + d\operatorname{vol}(S^3) \wedge P , \quad (53)$$

i.e. $Q, X \neq 0$ in (12), where $g_{\ell}(AdS_3)$ and $dvol_{\ell}(AdS_3)$ are the standard metric and volume 3-form of AdS₃ with radius ℓ , respectively, and $g(\mathbb{R}^5)$ is the Euclidean metric on

 \mathbb{R}^5 . Similarly, $g(S^3)$ and $d\text{vol}(S^3)$ are the metric and volume 3-form of unit round 3-sphere, respectively, and the 1-forms P and Q are constant and lie along the same direction in \mathbb{R}^5 , e.g. $P = p \, dw$ and $Q = q \, dw$ with p, q constants.

Focusing on the KSEs on σ_+ and setting without loss of generality A = 1, the integrability of the gravitino KSE along the \mathbb{R}^5 directions implies that $p^2 = q^2$. Moreover, one has to also consider the algebraic KSE $\Xi^+\sigma_+ = 0$ which arises from the integration of the KSE of 11-dimensional supergravity along the z direction of AdS₃. As $\Xi^+ = -(2\ell)^{-1} + \frac{1}{288}\Gamma_z X + \frac{1}{6}Q$, see [26], the algebraic KSE can be arranged as

$$\left(-\frac{1}{\ell}\Gamma_w + \frac{p}{6}\Gamma_z\Gamma_{(3)} + \frac{1}{3}q\right)\sigma_+ = 0 , \qquad (54)$$

where $\Gamma_{(3)}$ is the product of the three gamma matrices along orthonormal directions tangential to S^3 .

As $p^2 = q^2$ to solve (54) let us set p = q. The other case p = -q can be treated in a similar way. Then decompose (54) into eigenspaces of Γ_w and $\Gamma_z\Gamma_{(3)}$. Using $\Gamma_w^2 = 1$ and $(\Gamma_z\Gamma_{(3)})^2 = 1$ and $\Gamma_w\Gamma_z\Gamma_{(3)} = \Gamma_z\Gamma_{(3)}\Gamma_w$, we have that

$$-\eta \frac{1}{\ell} + \zeta \frac{q}{6} + \frac{1}{3}q = 0 , \qquad (55)$$

where $\Gamma_w \sigma_+ = \eta \sigma_+$ and $\Gamma_z \Gamma_{(3)} \sigma_+ = \zeta \sigma_+$ with $\eta, \zeta = \pm 1$. There are four cases to consider leading to $q = \pm 2\ell^{-1}$ and $q = \pm 6\ell^{-1}$. Two of these solutions are related to the other two by a change of the overall sign of the 4-form field strength F. So there are only two remaining independent solutions. Furthermore the $q = \pm 6\ell^{-1}$ solution is ruled out by the warp factor field equation [26]. In addition, the field equation along S^3 implies that $p^2 = 4$. As $q = \pm 2\ell^{-1}$, one finds that $\ell^2 = 1$, which is the near horizon geometry of the M2- and M5-brane intersection on a 1-brane solution. This solution preserves 1/2 of supersymmetry as each of the KSEs on σ_{\pm} and τ_{\pm} give 4 independent solutions.

As in the previous AdS_2 case, we next consider the KSE along the S^3 directions whose supercovariant derivative can be put in the form

$$\mathcal{D}_{\alpha}^{+} = \nabla_{\alpha}^{S^{3}} + \Gamma_{\alpha}\Gamma_{z}(\frac{1}{6}p\zeta + \frac{1}{12}q)\eta , \qquad (56)$$

where α labels the three othonormal directions tangential to S^3 . Considering the form bilinears (52) with $\phi_+ = \sigma_+$, it is easy to show that k and ω are CCKY forms on S^3 . As a result their Hodge duals on S^3 are KY forms and generate symmetries for the probe action (47). The bilinear φ is also a CCKY form but its dual is a scalar.

6.4 AdS_3 solution for intersecting M5-branes

An ansatz which includes the near horizon geometry of three M5-branes intersecting on a 1-brane is

$$g = g_{\ell}(AdS_3) + g(S^2) + g(\mathbb{R}^6) , \quad F = dvol(S^2) \wedge W ,$$
 (57)

i.e. Q = 0 in (12), where $g_{\ell}(AdS_3)$ is the metric of AdS₃ with radius ℓ , $g(S^2)$ and $dvol(S^2)$ are the metric and volume 2-form of round 2-sphere with unit radius, respectively, $g(\mathbb{R}^6)$ is the Euclidean metric on \mathbb{R}^6 and W is a constant non-degenerate 2-form on \mathbb{R}^6 .

To continue let us focus on the gravitino KSE on σ_+ . The integrability condition of this equation along the \mathbb{R}^6 directions implies that $W = \gamma \lambda$ and that $\Gamma_{1234}\sigma_+ = \eta \sigma_+$ and $\Gamma_{1256}\sigma_+ = \zeta \sigma_+$, where λ is a Kähler form of the Euclidean metric on \mathbb{R}^6 and we have chosen \mathbb{R}^6 along the 123456 directions. Without loss of generality one can always choose $\lambda = \lambda_1 dx^1 \wedge dx^2 + \lambda_2 dx^3 \wedge dx^4 + \lambda_3 dx^5 \wedge dx^6$ with $\lambda_1, \lambda_2, \lambda_3 = \pm 1$. Then the algebraic KSE, $\Xi^+\sigma_+ = (-(2\ell)^{-1} + (288)^{-1}\Gamma_z \not{X})\sigma_+ = 0$, implies, after imposing $\Gamma_z\Gamma_{12}\Gamma_{(2)}\sigma_+ = \theta\sigma_+$, $\theta = \pm 1$, that either $\gamma = \pm 6\ell^{-1}$ or $\gamma = \pm 2\ell^{-1}$, where $\Gamma_{(2)}$ is the product of two gamma matrices along two orthonormal directions tangential to S^2 . The warp factor field equation is not satisfied for $\gamma = \pm 6\ell^{-1}$. While for $\gamma = \pm 2\ell^{-1}$, the Einstein equation along S^2 gives $\ell = 2$. This is the solution that describes the near horizon geometry of three intersecting M5-branes and preserves 1/4 of supersymmetry.

After imposing all the conditions above on σ_+ appropriate for this solution, the gravitino KSE along S^2 can be written as

$$\mathcal{D}_{\alpha}^{+} = \nabla_{\alpha} \pm \frac{1}{2} \epsilon_{\alpha\beta} \Gamma^{\beta} \Gamma_{12} , \qquad (58)$$

where a, b are restricted along two orthonormal tangential directions of S^2 . Next it is straightforward to show that the 1-form bilinears restricted on S^2 are KY forms the 2-form bilinears restricted on S^2 are CCKY forms. So the 1-form bilinears generate symmetries for the probe action (47).

7 Concluding remarks

We have presented the TCFH of all supersymetric AdS backgrounds of 11-dimensional supergravity. Therefore we have demonstrated that all the form bilinears on the internal space of these backgrounds are CKY forms with respect to the TCFH connection. This provides a geometric interpretation for all the conditions on these form bilinears implied by the KSE of the theory. We have also given the reduced holonomy of the TCFH connections for generic supersymmetric backgrounds and we have found that it factorises on the space symmetric and skew-symmetric form bilinears under the exchange of the two Killing spinors. We have illustrated our results with some examples that include the maximally supersymmetric AdS backgrounds of 11-dimensional supergravity as well as some other AdS backgrounds that arise as near horizon geometries of intersecting M-branes. We have found that some of the form bilinears on these backgrounds are KY forms and so generate symmetries for spinning particle probes propagating on the internal spaces.

It is clear from the results of [5] that TCFHs can be constructed for all supersymmetric theories that exhibit a gravitino KSE. The generality of this result allows for the exploration of many more possibilities. As a consequence, all form bilinears are CKY forms with respect to some connection. This indicates they may be related to symmetries for particle and string probes propagating on supersymmetric backgrounds. Of course for special backgrounds this can be confirmed with explicit calculations like those presented in section 6 of this paper. However given a TCFH of a supersymmetric theory, there is not a systematic way to construct a probe action which exhibits symmetries generated by the form bilinears. Although there are many partial results that corroborate such constructions, the general question remains open.

Acknowledgments

EPB is supported by the CONACYT, the Mexican Council of Science.

References

- J. P. Gauntlett, J. B. Gutowski, C. M. Hull, S. Pakis and H. S. Reall, "All supersymmetric solutions of minimal supergravity in five- dimensions," Class. Quant. Grav. 20 (2003), 4587-4634 doi:10.1088/0264-9381/20/21/005 [arXiv:hep-th/0209114 [hep-th]].
- [2] J. Gillard, U. Gran and G. Papadopoulos, "The Spinorial geometry of supersymmetric backgrounds," Class. Quant. Grav. 22 (2005), 1033-1076 doi:10.1088/0264-9381/22/6/009 [arXiv:hep-th/0410155 [hep-th]].
- [3] U. Gran, J. Gutowski and G. Papadopoulos, "Classification, geometry and applications of supersymmetric backgrounds," Phys. Rept. **794** (2019), 1-87 doi:10.1016/j.physrep.2018.11.005 [arXiv:1808.07879 [hep-th]].
- [4] J. Gutowski and G. Papadopoulos, "Eigenvalue estimates for multiform modified Dirac operators," J. Geom. Phys. 160 (2021), 103954 doi:10.1016/j.geomphys.2020.103954 [arXiv:1911.02281 [math.DG]].
- G. Papadopoulos, "Twisted form hierarchies, Killing-Yano equations and supersymmetric backgrounds," JHEP 07 (2020), 025 doi:10.1007/JHEP07(2020)025 [arXiv:2001.07423 [hep-th]].
- [6] R. Penrose, Ann. N.Y. Acad. Sci. 224 (1973) 125.
- [7] R. Floyd, The dynamics of Kerr fields, Ph. D. Thesis, London (1973).
- [8] B. Carter, "Global structure of the Kerr family of gravitational fields," Phys. Rev. 174 (1968) 1559.
- [9] S. Chandrasekhar, "The Solution Of Dirac's Equation In Kerr Geometry," Proc. Roy. Soc. Lond. A 349 (1976) 571.
- [10] B. Carter, "Killing Tensor Quantum Numbers And Conserved Currents In Curved Space," Phys. Rev. D 16 (1977) 3395.
- [11] B. Carter and R. G. Mclenaghan, "Generalized Total Angular Momentum Operator For The Dirac Equation In Curved Space-Time," Phys. Rev. D 19 (1979) 1093.

- [12] P. Krtous, D. Kubiznak, D. N. Page and V. P. Frolov, "Killing-Yano tensors, rank-2 Killing tensors, and conserved quantities in higher dimensions," JHEP 0702 (2007) 004 [arXiv:hep-th/0612029].
- [13] F. De Jonghe, K. Peeters and K. Sfetsos, "Killing-Yano supersymmetry in string theory," Class. Quant. Grav. 14 (1997) 35 [arXiv:hep-th/9607203].
- [14] Y. Chervonyi and O. Lunin, "Killing(-Yano) Tensors in String Theory," JHEP 1509 (2015) 182 doi:10.1007/JHEP09(2015)182 [arXiv:1505.06154 [hep-th]].
- [15] M. Cariglia, "Hidden Symmetries of Dynamics in Classical and Quantum Physics," Rev. Mod. Phys. 86 (2014) 1283 doi:10.1103/RevModPhys.86.1283 [arXiv:1411.1262 [math-ph]].
- [16] V. Frolov, P. Krtous and D. Kubiznak, "Black holes, hidden symmetries, and complete integrability," Living Rev. Rel. 20 (2017) no.1, 6 doi:10.1007/s41114-017-0009-9 [arXiv:1705.05482 [gr-qc]].
- [17] D. Kastor and J. Traschen, "Conserved gravitational charges from Yano tensors," JHEP 08 (2004), 045 doi:10.1088/1126-6708/2004/08/045 [arXiv:hep-th/0406052 [hep-th]].
- [18] G. Papadopoulos, "Killing-Yano equations and G-structures," Class. Quant. Grav.
 25 (2008), 105016 doi:10.1088/0264-9381/25/10/105016 [arXiv:0712.0542 [hep-th]].
- [19] P. S. Howe and U. Lindström, "Some remarks on (super)-conformal Killing-Yano tensors," JHEP 11 (2018), 049 doi:10.1007/JHEP11(2018)049 [arXiv:1808.00583 [hepth]].
- [20] U. Lindström and O. Sarıoğlu, "New currents with Killing–Yano tensors," Class. Quant. Grav. 38 (2021) no.19, 195011 doi:10.1088/1361-6382/ac1871
 [arXiv:2104.12451 [hep-th]].
- [21] U. Lindström and O. Sarıoğlu, "Tensionless strings and Killing(-Yano) tensors," Phys. Lett. B 829 (2022), 137088 doi:10.1016/j.physletb.2022.137088 [arXiv:2202.06542 [hep-th]].
- [22] G. W. Gibbons, R. H. Rietdijk and J. W. van Holten, "SUSY in the sky," Nucl. Phys. B 404 (1993) 42 [arXiv:hep-th/9303112].
- [23] L. Brink, P. Di Vecchia and P. S. Howe, "A Lagrangian Formulation of the Classical and Quantum Dynamics of Spinning Particles," Nucl. Phys. B 118 (1977), 76-94 doi:10.1016/0550-3213(77)90364-9
- [24] G. Papadopoulos and E. Pérez-Bolaños, "Symmetries, spinning particles and the TCFH of D=4,5 minimal supergravities," Phys. Lett. B 819 (2021), 136441 doi:10.1016/j.physletb.2021.136441 [arXiv:2101.10709 [hep-th]].

"TCFHs, hidden symmetries and M-theory backgrounds," [arXiv:2201.11563 [hep-th]].

- [25] L. Grimanellis, G. Papadopoulos and J. Phillips, "TCFHs, hidden symmetries and type II theories," [arXiv:2111.15405 [hep-th]].
- [26] J. B. Gutowski and G. Papadopoulos, "Supersymmetry of AdS and flat backgrounds in M-theory," JHEP 02 (2015), 145 doi:10.1007/JHEP02(2015)145 [arXiv:1407.5652 [hep-th]].
- [27] N. Kim and J. D. Park, "Comments on AdS(2) solutions of D=11 supergravity," JHEP 09 (2006), 041 doi:10.1088/1126-6708/2006/09/041 [arXiv:hep-th/0607093 [hep-th]].
- [28] J. P. Gauntlett, N. Kim and D. Waldram, "Supersymmetric AdS(3), AdS(2) and Bubble Solutions," JHEP 04 (2007), 005 doi:10.1088/1126-6708/2007/04/005 [arXiv:hepth/0612253 [hep-th]].
- [29] M. Gabella, D. Martelli, A. Passias and J. Sparks, " $\mathcal{N} = 2$ supersymmetric AdS₄ solutions of M-theory," Commun. Math. Phys. **325** (2014), 487-525 doi:10.1007/s00220-013-1865-0 [arXiv:1207.3082 [hep-th]].
- [30] J. P. Gauntlett, D. Martelli, J. Sparks and D. Waldram, "Supersymmetric AdS(5) solutions of M theory," Class. Quant. Grav. 21 (2004), 4335-4366 doi:10.1088/0264-9381/21/18/005 [arXiv:hep-th/0402153 [hep-th]].
- [31] F. Apruzzi, M. Fazzi, A. Passias, D. Rosa and A. Tomasiello, "AdS₆ solutions of type II supergravity," JHEP **11** (2014), 099 [erratum: JHEP **05** (2015), 012] doi:10.1007/JHEP11(2014)099 [arXiv:1406.0852 [hep-th]].
- [32] J. Gutowski and G. Papadopoulos, "On supersymmetric AdS₆ solutions in 10 and 11 dimensions," JHEP **12** (2017), 009 doi:10.1007/JHEP12(2017)009 [arXiv:1702.06048 [hep-th]].
- [33] G. Papadopoulos and P. K. Townsend, "Intersecting M-branes," Phys. Lett. B 380 (1996), 273-279 doi:10.1016/0370-2693(96)00506-0 [arXiv:hep-th/9603087 [hep-th]].
- [34] H. J. Boonstra, B. Peeters and K. Skenderis, "Brane intersections, anti-de Sitter space-times and dual superconformal theories," Nucl. Phys. B 533 (1998), 127-162 doi:10.1016/S0550-3213(98)00512-4 [arXiv:hep-th/9803231 [hep-th]].