DIFFERENTIAL CALCULUS FOR GENERALIZED GEOMETRY AND GEOMETRIC LAX FLOWS

SHENGDA HU

ABSTRACT. Employing a class of generalized connections, we describe certain differential complices $\left(\widetilde{\Omega}^*_{\mathbb{T}}(M), \widetilde{\mathfrak{d}}^{\mathbb{T}}\right)$ constructed from $\wedge^* \mathbb{T} M$ and study some of their basic properties, where $\mathbb{T} M = TM \oplus T^*M$ is the generalized tangent bundle on M. A number of classical geometric notions are extended to $\mathbb{T} M$, such as the curvature tensor for a generalized connection. In particular, we describe an analogue to the Levi-Civita connection when $\mathbb{T} M$ is endowed with a generalized metric and a structure of exact Courant algebroid. We further describe in generalized geometry the analogues to the Chern-Weil homomorphism, a Weitzenböck identity, the Ricci flow and Ricci soliton, the Hermitian-Einstein equation and the degree of a holomorphic vector bundle. Furthermore, the Ricci flows are put into the context of geometric Lax flows, which may be of independent interest.

1. INTRODUCTION

In generalized geometry à la Hitchin [19], over a smooth manifold M of real dimension n, the bundle $\mathbb{T}M := TM \oplus T^*M$ is considered the analogue of the classical tangent bundle TM. It fits into the natural exact sequence

$$0 \to T^*M \hookrightarrow \mathbb{T}M \xrightarrow{\pi} TM \to 0$$

and is endowed with the natural pairing

$$\langle x, y \rangle = \langle X + \xi, Y + \eta \rangle := \frac{1}{2} \left(\iota_X \eta + \iota_Y \xi \right)$$

where $x, y \in C^{\infty}(\mathbb{T}M)$ and $X, Y \in C^{\infty}(TM)$, $\xi, \eta \in C^{\infty}(T^*M)$ are their respective components. The dual of $\mathbb{T}M$ can be identified with itself under the pairing $2\langle,\rangle$. Wellknown geometric structures on $\mathbb{T}M$ such as generalized complex, Riemannian, Hermitian, Kähler structures and generalized connections are natural extensions of the corresponding classical notions on TM. There are by now many references in the literature, including the pioneering works by Gualtieri [14, 15, 16, 17] on the subjects.

We show that the analogue can be pushed further, leading to coherent extensions of known geometric notions, as well as Lax forms for some geometric flows. For simplicity, we will restrict our considerations to compact connected orientable smooth manifolds without boundary, while aside from cohomology computations, most descriptions are of a local nature. It starts with a differential complex constructed with $\mathbb{T}M$ in place of TM, as a quotient of $\Omega^*_{\mathbb{T}}(M) := C^{\infty}(\wedge^*\mathbb{T}M)$, whose differential is defined from a generalized

²⁰²⁰ Mathematics Subject Classification. 53D18, 53E20, 53B15, 53B20.

Key words and phrases. Generalized connections; de Rham cohomology; generalized curvature; geometric Lax flows; generalized Kähler geometry; generalized holomorphic bundles.

connection on $\mathbb{T}M$. Let V be a vector bundle, as in [15], a generalized connection on V is a derivation

 $\mathcal{D}: C^{\infty}(V) \to C^{\infty}(\mathbb{T}M \otimes V)$

such that for $f \in C^{\infty}(M)$, $v \in C^{\infty}(V)$, $x \in C^{\infty}(\mathbb{T}M)$ and $X = \pi(X)$

 $\mathcal{D}_{x}(fv) = X(f)v + f\mathcal{D}_{x}v$

Given a generalized connection $\mathcal{D}^{\mathbb{T}}$ on $\mathbb{T}M$, it is *TM*-torsion free if for all $x, y \in C^{\infty}(\mathbb{T}M)$

(1.1)
$$\pi \left(x \diamond_{\mathbb{T}} y \right) = \left[\pi \left(x \right), \pi \left(y \right) \right]$$

where the $\mathcal{D}^{\mathbb{T}}$ -diamond bracket $\diamond_{\mathbb{T}}$ is the skew symmetrization of $\mathcal{D}^{\mathbb{T}}$:

(1.2)
$$x \diamond_{\mathbb{T}} y := \mathcal{D}_x^{\mathbb{T}} y - \mathcal{D}_y^{\mathbb{T}} x$$

The first observation is a differential complex obtained from such a generalized connection.

Theorem 1.1 (§2.1). *Consider the derivation* $\mathbb{d}^{\mathbb{T}}$: $\Omega^*_{\mathbb{T}}(M) \to \Omega^{*+1}_{\mathbb{T}}(M)$ defined by

$$\left(\mathbb{d}^{\mathbb{T}}\theta\right)(x_0, x_1, \dots, x_k) := \sum_i (-1)^i \left(\mathcal{D}_{x_i}^{\mathbb{T}}\theta\right)\left(x_0, x_1, \dots, \widehat{x}_i, \dots, x_k\right)$$

where $\theta \in \Omega^k_{\mathbb{T}}(M)$ and $x_j \in C^{\infty}(\mathbb{T}M)$. Then $\mathbb{d}^{\mathbb{T}} \circ \mathbb{d}^{\mathbb{T}}$ is tensorial iff $\mathcal{D}^{\mathbb{T}}$ is T *M*-torsion free, in which case, the quotient $\widetilde{\Omega}^*_{\mathbb{T}}(M)$ of $\Omega^*_{\mathbb{T}}(M)$ by the image of $\mathbb{d}^{\mathbb{T}} \circ \mathbb{d}^{\mathbb{T}}$ is a differential complex with the induced derivation $\widetilde{\mathbb{d}}^{\mathbb{T}}$.

The cohomology of the resulting complex $\left(\widetilde{\Omega}^*_{\mathbb{T}}(M), \widetilde{\mathbb{d}}^{\mathbb{T}}\right)$ is denoted $\widetilde{H}^*_{\mathbb{T}}(M)$, and the projection π induces a natural map $\widetilde{\pi}^*$: $H^*(M) \to \widetilde{H}^*_{\mathbb{T}}(M)$, where $H^*(M)$ is the classical de Rham cohomology.

Suppose that $\mathbb{T}M$ is endowed with a generalized Riemannian structure \mathbb{G}^{b} [14, 16], defined by a Riemannian metric g and a 2-form $b \in \Omega^{2}(M)$

(1.3)
$$\mathbb{G}^{b}(x, y) := \frac{1}{2} \left[g(X, Y) + g^{-1} \left(\xi - \iota_{X} b, \eta - \iota_{Y} b \right) \right]$$

The \mathbb{G}^{b} -metric connections are generalized connections on $\mathbb{T}M$ that preserve \langle, \rangle and \mathbb{G}^{b} , among which the *TM*-torsion free ones are classified by $\phi \in \Omega^{3}(M)$ and denoted $\mathcal{D}^{\phi,b}$.

The (ϕ, b) -connections $\mathcal{D}^{\phi,b}$ can be seen as analogues of the Levi-Civita connections in this context, since they are also compatible (Definition 2.7) with the corresponding almost Dorfman brackets $*_{\gamma}$ defined by $\gamma = \phi - db$:

(1.4)
$$(X+\xi) *_{\gamma} (Y+\eta) = [X,Y] + \mathcal{L}_X \eta - d\iota_Y \xi + \iota_Y \iota_X \gamma$$

When $d\gamma = 0$, the bracket $*_{\gamma}$ is a *Dorfman bracket*.

Theorem 1.2 (Theorem 2.8, §2.3). Let $\gamma \in \Omega^3(M)$ be a closed 3-form and $\phi = \gamma + db$. The (ϕ, b) -connection $\mathcal{D}^{\phi,b}$ associated with \mathbb{G}^b is the unique T M-torsion free \mathbb{G}^b -metric connection on $\mathbb{T}M$ that is metric compatible with a Dorfman bracket $*_{\gamma}$. The natural map $\widetilde{\pi}^*$: $H^*(M) \to \widetilde{H}^*_{\phi,b}(M)$ is injective. Moreover, $\widetilde{H}^{2n}_{\phi,b}(M) \cong \mathbb{R}$. The differential calculus thus established on $\mathbb{T}M$ leads to a natural definition of curvature for any generalized connection \mathcal{D} on any vector bundle V:

$$\mathcal{F}^{\mathbb{T}}(\mathcal{D}) := \mathbb{d}_{\mathcal{D}}^{\mathbb{T}} \circ \mathcal{D} \in \Omega^{2}_{\mathbb{T}} \left(\operatorname{End}\left(V \right) \right)$$

where $\mathbb{d}_{D}^{\mathbb{T}}$ is the extension of $\mathbb{d}^{\mathbb{T}}$ to $C^{\infty}(\mathbb{T}M \otimes V)$ as usual. In terms of covariant derivatives, $\mathcal{F}^{\mathbb{T}}(D)$ is given by

(1.5)
$$\mathcal{F}_{x,y}^{\mathbb{T}}(\mathcal{D}) v := \left(\mathcal{D}_x \mathcal{D}_y - \mathcal{D}_y \mathcal{D}_x - \mathcal{D}_{x \diamond_{\mathbb{T}} y} \right) v$$

where $x, y \in C^{\infty}(\mathbb{T}M)$ and $v \in C^{\infty}(V)$. A side-effect of this is that the resulting curvature tensor $\mathcal{F}^{\mathbb{T}}$ now depends on the generalized connection $\mathcal{D}^{\mathbb{T}}$ on $\mathbb{T}M$. Nonetheless, passing to the quotient $\widetilde{\Omega}^*_{\mathbb{T}}$ (End (*V*)), the Chern-Weil homomorphism naturally extends.

Theorem 1.3 (§3.1). Any invariant polynomial of the $\mathcal{D}^{\mathbb{T}}$ -curvature $\mathcal{F}^{\mathbb{T}}(\mathcal{D})$ defines a class in $\widetilde{H}^*_{\mathbb{T}}(M)$, which coincides with the image under $\widetilde{\pi}^*$ of the corresponding classical characteristic class in $H^*(M)$.

The (ϕ, b) -curvature for $\mathcal{D}^{\phi,b}$ itself, denoted $\mathcal{R}^{\phi,b}$, can be regarded as the analogue to the Riemannian curvature in this context. The (ϕ, b) -Ricci curvature $\mathcal{R}ic^{\phi,b}$, as well as the corresponding scalar curvature are defined via the usual contractions of $\mathcal{R}^{\phi,b}$. Associated to the connection $\mathcal{D}^{\phi,b}$, further operators, including the analogues to the Bochner and Hodge Laplacians, can be defined on the tensor bundles. An analogue to the Weitzenböck identity holds, which states that the Laplacians differ by the (ϕ, b) -Ricci curvature.

Theorem 1.4 (Theorem 4.7). On $\Omega^*_{\mathbb{T}}(M)$, let $\Delta^{\phi,b}$ be the $\mathcal{D}^{\phi,b}$ -Hodge Laplacian and $\Delta_{\mathcal{D}^{\phi,b}}$ be the $\mathcal{D}^{\phi,b}$ -Bochner Laplacian, then

$$\Delta^{\phi,b} = \Delta_{\mathcal{D}^{\phi,b}} + \mathbb{G}^b \mathcal{R}ic^{\phi,b}\mathbb{G}^b$$

Consider next a generalized complex manifold $(M, \gamma; \mathbb{J})$. A generalized connection $\mathcal{D}^{\mathbb{T}}$ is \mathbb{J} -compatible with $*_{\gamma}$ if the diamond bracket $\diamond_{\mathbb{T}}$ coincides with $*_{\gamma}$ when restricted to sections of the same eigenbundle of \mathbb{J} . For such $\mathcal{D}^{\mathbb{T}}$, $\mathbb{d}^{\mathbb{T}}$ admits the natural decomposition

$$\mathbf{d}^{\mathrm{T}} = \partial_{\mathbf{J}}^{\mathrm{T}} + \overline{\partial}_{\mathbf{J}}^{\mathrm{T}}$$

according to the types with respect to J. Together with a generalized metric \mathbb{G}^b commuting with J, we have a generalized Hermitian manifold $(M, \gamma; \mathbb{G}^b, \mathbb{J})$, which corresponds classically to an almost bi-Hermitian structure $(M, \gamma; g, I_{\pm}; b)$, where g is Hermitian with respect to both almost complex structures I_{\pm} . We show that the J-compatibility of $\mathcal{D}^{\phi,b}$ with $*_{\gamma}$ is a generalized Kähler condition, equivalent to the condition given in [15].

Theorem 1.5 (Theorem 5.7). On a generalized Hermitian manifold $(M, \gamma; \mathbb{G}^b, \mathbb{J})$, let $\phi = \gamma + db$. Then it is a generalized Kähler manifold iff $\mathcal{D}^{\phi,b}$ is \mathbb{J} -compatible with $*_{\gamma}$.

In terms of I_{\pm} , the J-compatibility is equivalent to $\nabla^{\pm\phi}I_{\pm} = 0$. On a generalized Kähler manifold, I_{\pm} are integrable. Working with $\mathcal{D}^{\phi,b}$, we recover a well-known result obtained via holomorphic reduction [17], that the I_{\pm} -(anti)holomorphic tangent bundles on a generalized Kähler manifold carry natural I_{\pm} -holomorphic structures respectively.

For a \mathbb{J} -holomorphic Hermitian vector bundle $(V, \overline{\partial}_{\mathbb{J}}, h)$, the notion of Chern connection extends naturally. Over a generalized Hermitian manifold $(M, \gamma; \mathbb{G}^b, \mathbb{J})$, there is a natural contraction $\Lambda_{\mathbb{J}}$ on $\Omega^2_{\mathbb{T}}(M)$, where for $x, y \in C^{\infty}(\mathbb{T}M)$

$$\Lambda_{\mathbb{J}} (x \wedge y) := \mathbb{G}^{b} (\mathbb{J}x, y)$$

Given a γ - \mathbb{J} -connection $\mathcal{D}^{\mathbb{T}}$ (Definition 5.1) on $\mathbb{T}M$, analogous to the classical case (Lübke-Teleman [25]), we propose the following $\mathcal{D}^{\mathbb{T}}$ - \mathbb{J} -*Hermitian-Einstein equation* for the Hermitian metric *h*:

$$\sqrt{-1}\Lambda_{\mathbb{J}_{-}}\left(\mathcal{F}^{\mathbb{T},C}(V)\right) = 2c\mathrm{Id}_{V}$$

where $c \in \mathbb{R}$ and $\mathcal{F}^{\mathbb{T},C}$ denotes the $\mathcal{D}^{\mathbb{T}}$ -curvature of the generalized Chern connection. The contraction also leads to the notion of degree for a J-holomorphic Hermitian vector bundle. The degree is independent to the choice of Hermitian metric on *V* if the generalized Hermitian manifold is $\mathcal{D}^{\mathbb{T}}$ -J-*Gauduchon*, i.e. for all $f \in C^{\infty}(M)$

$$\int_{M} \Lambda_{\mathbb{J}_{-}} \left(\partial_{\mathbb{J}}^{\mathbb{T}} \overline{\partial}_{\mathbb{J}}^{\mathbb{T}} f \right) d \operatorname{vol}_{g} = 0$$

For such manifolds, the notions of slope and stability naturally extend and one should expect a version of Kobayashi-Hitchin correspondence to hold. On a generalized Kähler manifold, these notions relate to their classical counterparts, in particular, the J-Hermitian-Einstein equation is equivalent to an equation proposed by Hitchin [20] (Remark in §3.3 *ibid.*).

Theorem 1.6 (§6.3). Let $(M, \gamma; \mathbb{G}^b, \mathbb{J})$ be a \mathbb{J} -Gauduchon generalized Kähler manifold and $\omega_{\pm} \in \Omega^2(M)$ be the Kähler forms for I_{\pm} respectively. Let $(V, \partial_{\mathbb{J}})$ be a \mathbb{J} -holomorphic vector bundle, then the \mathbb{J} -Hermitian-Einstein equation is equivalent to

$$\frac{\sqrt{-1}}{2} \left(F_{+}^{C}(V) \wedge \omega_{+}^{m-1} + (-1)^{\varepsilon} F_{-}^{C}(V) \wedge \omega_{-}^{m-1} \right) = c (m-1)! \mathrm{Id}_{V} d \mathrm{vol}_{g}$$

where F_{\pm}^{C} are the classical Chern curvatures with respect to I_{\pm} , $\varepsilon = 0$ if I_{\pm} induce the same orientation on T M and $\varepsilon = 1$ otherwise. The \mathbb{G}^{b} -degree of (V, h) is computed by

$$\deg_{\mathbb{G}^{b}}(V,h) = \frac{1}{2} \left(\deg_{+}(V,h) + \deg_{-}(V,h) \right)$$

where \deg_{\pm} are the degrees with respect to the I_{\pm} -holomorphic structure.

Geometric flows such as the mean curvature flow (Brakke [5]) and the Ricci flow (Hamilton [18]) are very important in understanding smooth manifolds and structures associated to them. In generalized geometry, it is natural to consider flows involving the structures on $\mathbb{T}M$ such as the generalized metrics or generalized complex structures. A generalized metric \mathbb{G}^b defines a bundle automorphism of $\mathbb{T}M$ via the pairing \langle, \rangle :

(1.6)
$$\langle \mathbb{G}^b x, y \rangle := \mathbb{G}^b \langle x, y \rangle$$

which enjoys the following properties

(1.7)
$$(\mathbb{G}^b)^2 = 1 \text{ and } \langle \mathbb{G}^b x, y \rangle = \langle x, \mathbb{G}^b y \rangle$$

For a smooth family of generalized metrics $\mathbb{G}_{t}^{b_{t}}$, differentiating the above identites shows that $\mathbb{K}_{t} = \frac{d}{dt}\mathbb{G}_{t}^{b_{t}}$ is skew with respect to $\mathbb{G}_{t}^{b_{t}}$ and symmetric with respect to \langle, \rangle , i.e. it satisfies

(1.8)
$$\mathbb{G}_{t}^{b_{t}}\mathbb{K}_{t} + \mathbb{K}_{t}\mathbb{G}_{t}^{b_{t}} = 0 \text{ and } \langle \mathbb{K}_{t}x, y \rangle = \langle x, \mathbb{K}_{t}y \rangle$$

Hence, a flow of generalized metrics can be defined from geometric quantities satisfying the above identities. Similar identities can be derived for a smooth family of generalized complex structures. For any smooth family $\mathbb{L}_t : \mathbb{T}M \to \mathbb{T}M$, the commutator $[\mathbb{L}_t, \mathbb{G}_t^{b_t}]$ satisfies the first identity in (1.8), which lends to the consideration of Lax pairs (Lax [24]).

In this context, we generally assume that there is a fixed closed 3-form γ such that $\phi = \gamma + db$ holds for all (ϕ, b) under consideration. Hence the flow generated can be regarded as over a fixed Courant algebroid, defined by the Dorfman bracket $*_{\gamma}$. For instance, any $\mathbb{T}M$ -form $\theta \in \Omega^2_{\mathbb{T}}(M)$ defines a 2-form Lax flow via the induced map $\theta_t : \mathbb{T}M \to \mathbb{T}M$. The action of generalized symmetries of $\mathbb{T}M$ on a generalized metric can be represented as Lax flows generated by the (ϕ, b) -curvatures of a Hermitian line bundle.

Theorem 1.7 (Theorem 7.10). Consider a family of unitary generalized connections $\{D_t\}$ on a Hermitian line bundle V. Let $\phi_t = \gamma + db_t$, then the Lax flow

$$\frac{d}{dt}\mathbb{G}_{t}^{b_{t}}=\left[\mathcal{F}^{\phi_{t},b_{t}}\left(\mathcal{D}_{t}\right),\mathbb{G}_{t}^{b_{t}}\right]$$

corresponds to the push-forward of an initial generalized metric \mathbb{G}^b by a family of generalized diffeomorphisms of the exact Courant algebroid defined by $*_{\gamma}$.

Here we recall ([14], Hu-Uribe [23]) that a generalized diffeomorphism for the Courant algebroid structure on $\mathbb{T}M$ defined by $*_{\gamma}$ is a pair (λ, B) , where $\lambda \in \text{Diff}(M)$ is a diffeomorphism and $B \in \Omega^2(M)$, such that

$$\lambda^* \gamma = \gamma - dB$$

whose action on $\mathbb{T}M$ by push-forward is given by

$$(\lambda, B)_* (X + \xi) := \lambda_* (X + \iota_X B + \xi)$$

The action of $B \in \Omega^2(M)$ is also denoted e^B :

$$e^B \left(X + \xi \right) := X + \iota_X B + \xi$$

We comment that the diffeomorphism components of the generalized diffeomorphisms in Theorem 1.7 arise from the vector components of the generalized connections on V, which are 0 when D_t are liftings of classical connections on V.

Even though the Bianchi identites do not hold for $\mathcal{R}^{\phi,b}$ in general, it turns out that the (ϕ, b) -Ricci tensor $\mathcal{R}c^{\phi,b}$ is symmetric. The corresponding Lax flow is the *Ricci Lax flow*

(1.9)
$$\frac{d}{dt} \mathbb{G}_t^{b_t} = \left[\mathcal{R}ic^{\phi_t, b_t}, \mathbb{G}_t^{b_t} \right]$$

which defines the Ricci flow in this context. We note that in (1.9), only the components of $\mathcal{R}c^{\phi,b}$ that involve sections from different eigenbundles of \mathbb{G}^{b} contribute, similar to the Ricci flows for Courant algebroids considered in Ševera-Valach [26, 27] and Streets [29].

Theorem 1.8 (Theorem 7.13). *The Ricci Lax flow equation above is equivalent to the generalized Ricci flow equation*

$$\begin{cases} \frac{d}{dt}g_t = -2Rc_t + \frac{1}{2}\left(\gamma + db_t\right)^2\\ \frac{d}{dt}b_t = -d^*\left(\gamma + db_t\right) \end{cases}$$

The above system has been studied in mathematics and physics literatures, for instance, Garcia-Fernandez-Streets [9], and references therein. We note that an equation similar to (1.9) appeared in [29] citing communication with Gualtieri (reference [11] *ibid.*) and the equivalence with the generalized Ricci flow appeared in [9] (Remark 4.8 *ibid.*), via a somewhat different approach to the notion of generalized Ricci curvature.

A few more examples of geometric Lax flows are given. Conformal deformations of the Riemannian metric *g* can be represented as a Lax flow, of the form

$$\frac{d}{dt}\mathbb{G}_t^{b_t} = \left[\theta_t, \mathbb{G}_t^{b_t}\right]$$

where $\theta_t \in \Omega^2_{\mathbb{T}}(M)$ are $\mathbb{G}_t^{b_t}$ -conformal $\mathbb{T}M$ -forms (Definition 7.7). The *Ricci soliton equation* (Streets [28] and references therein) takes the following form (Definition 7.15)

$$\left[\mathcal{R}ic^{\phi_t,b_t} - \mathbb{d}^{\phi_t,b_t}u_t - \theta_t, \mathbb{G}_t^{b_t}\right] = 0$$

where the $\mathbb{G}_{l}^{b_{t}}$ -conformal $\mathbb{T}M$ -forms θ_{l} have constant conformal weights. We also see that the classical Kähler-Ricci flow can be recast as a geometric Lax flow (§7.4).

We expect that many classical constructions should admit natural extensions to $\mathbb{T}M$ via the differential calculus developed here. For instance, the notions of second fundamental form and mean curvature for submanifolds extend naturally. Spinors, which are behind the notion of J_-contraction in Definition 6.7, relate the geometry on $\mathbb{T}M$ back to $\Omega^*(M)$, which, in particular, lead to the canonical line of a generalized (almost) complex structure [14, 16] as well as the notion of scalar curvature in generalized Kähler geometry (Goto [11, 12], Wang [31]). Functionals involving curvatures, such as the Yang-Mills functional, can be extended (§3.3) and lead to natural questions on extremal / critical (generalized) connections / metrics with respect to them. Explicit examples such as compact Lie groups ([12], Hu [21]) could provide further insights into understanding these extensions. It should be worth exploring the interaction of the Riemannian, the complex and the Poisson geometric interpretations (Griffiths [13]), it would be interesting to understand if it provides new perspective for the related geometric flows.

We briefly give some further descriptions of the structure of the paper. In §2, we set up the differential calculus on $\mathbb{T}M$ and compute in §2.5 the group $\widetilde{H}^*_{\gamma,0}(G)$ for a semi-simple real Lie group G, with the bi-invariant metric and the Cartan 3-form γ . The generalized curvature tensors are introduced in §3. The rest of the article applies the constructions in various contexts. The analogue to the Riemann curvature is discussed in §4, together with the associated Ricci and scalar curvatures, as well as the generalized Bismut connections [15]. In §5, we apply the differential calculus to generalized complex and Hermitian manifolds. The degree, stability and Hermitian-Einstein equation for a generalized holomorphic bundle over a generalized Hermitian manifold are discussed in §6. In the last section §7, we discuss the notion of geometric Lax flows.

Acknowledgement. I would like to thank Spiro Karigiannis for helpful discussions and enlightening questions, and Yucong Jiang for pointing out the references [26, 27]. This research was partially supported by the NSERC Discovery Grant RGPIN-2019-05899.

2. DIFFERENTIAL CALCULUS ON $\mathbb{T}M$

Let $V \rightarrow M$ be a vector bundle. Recall that a generalized connection on V is a derivation:

$$D: C^{\infty}(V) \to C^{\infty}(\mathbb{T}M \otimes V): D(fv) = df \otimes v + fDv$$

where $\mathbb{T}M = TM \oplus T^*M$, $f \in C^{\infty}(M)$ and $v \in C^{\infty}(V)$. It is the *lift* of a classical connection ∇_0 on V if

$$(2.1) D_x v = \nabla_{0,\pi(x)} v$$

for all $x \in C^{\infty}(\mathbb{T}M)$ and $v \in C^{\infty}(V)$. The generalized connections naturally extend to tensor bundles in the standard fashion.

2.1. $\mathbb{T}M$ -forms. Under the pairing $2\langle , \rangle$, sections of $\wedge^*\mathbb{T}M$ can be seen as $\mathbb{T}M$ -forms and the space of such forms will be suggestively denoted by

$$\Omega^*_{\scriptscriptstyle \mathbb{T}}(M) := C^{\infty}(\wedge^* \mathbb{T}M)$$

Let $\mathcal{D}^{\mathbb{T}}$ be a generalized connection on $\mathbb{T}M$. The skew-symmetrization of the covariant derivative by $\mathcal{D}^{\mathbb{T}}$ induces the $\mathcal{D}^{\mathbb{T}}$ -derivation $\mathbb{d}^{\mathbb{T}}$. Namely, for $\theta \in \Omega^k_{\mathbb{T}}(M)$:

(2.2)
$$\left(\mathbb{d}^{\mathbb{T}} \theta \right) (x_0, x_1, \dots, x_k) := \sum_i (-1)^i \left(\mathcal{D}_{x_i}^{\mathbb{T}} \theta \right) \left(x_0, x_1, \dots, \hat{x_i}, \dots, x_k \right)$$

where $x_i \in C^{\infty}(\mathbb{T}M)$. For $f \in C^{\infty}(M)$, $\mathbb{d}^{\mathbb{T}}$ coincides with the usual differential:

(2.3)
$$\mathbb{d}^{\mathbb{T}}f := df \in \Omega^{1}_{\mathbb{T}}(M)$$

Moverover, $\mathbb{d}^{\mathbb{T}}$ is a graded derivation on $\Omega^*_{\mathbb{T}}(M)$, i.e. for $\theta_1 \in \Omega^k(M)$ and $\theta_2 \in \Omega^*(M)$:

(2.4)
$$\mathbf{d}^{\mathbb{T}}\left(\theta_{1} \wedge \theta_{2}\right) = \left(\mathbf{d}^{\mathbb{T}}\theta_{1}\right) \wedge \theta_{2} + (-1)^{k} \theta_{1} \wedge \left(\mathbf{d}^{\mathbb{T}}\theta_{2}\right)$$

Standard computations lead to

(2.5)
$$\begin{pmatrix} \left(\mathbb{d}^{\mathbb{T}} \circ \mathbb{d}^{\mathbb{T}} \theta \right) \left(x_{0}, x_{1}, \dots, x_{k+1} \right) \\ = \sum_{i < j} \left(-1 \right)^{i+j} \left[\tau_{T} \left(x_{i}, x_{j} \right) \right] \theta \left(x_{0}, \dots, \hat{x}_{i}, \dots, \hat{x}_{j}, \dots, x_{k+1} \right) \\ - \sum_{i < j < \ell} \left(-1 \right)^{i+j+\ell} \theta \left(\left[x_{i} \diamond_{\mathbb{T}} x_{j} \diamond_{\mathbb{T}} x_{\ell} \right], x_{0}, \dots, \hat{x}_{i}, \dots, \hat{x}_{j}, \dots, \hat{x}_{\ell}, \dots, x_{k+1} \right)$$

where τ_T is the *TM*-torsion of $\mathcal{D}^{\mathbb{T}}$ given in (2.7) and $[x \diamond_{\mathbb{T}} y \diamond_{\mathbb{T}} z]$ is the *Jacobiator* of the diamond bracket $\diamond_{\mathbb{T}}$ (1.2):

(2.6)
$$[x \diamond_{\mathbb{T}} y \diamond_{\mathbb{T}} z] := (x \diamond_{\mathbb{T}} y) \diamond_{\mathbb{T}} z + c.p.$$

Definition 2.1. Let $\mathcal{D}^{\mathbb{T}}$ be a generalized connection on $\mathbb{T}M$. Its *TM*-torsion is

(2.7)
$$\tau_T(x, y) := \pi \left(x \diamond_{\mathbb{T}} y \right) - \left[\pi \left(x \right), \pi \left(y \right) \right]$$

where $x, y \in C^{\infty}(\mathbb{T}M)$. Then $\mathcal{D}^{\mathbb{T}}$ is *TM*-torsion free if its *TM*-torsion vanishes.

Let $\mathcal{R}^{\mathbb{T}}$ be the (not necessarily tensorial) curvature operator for $\mathcal{D}^{\mathbb{T}}$:

$$\mathcal{R}_{x,y}^{\mathbb{T}}z := \mathcal{D}_{x}^{\mathbb{T}}\mathcal{D}_{y}^{\mathbb{T}}z - \mathcal{D}_{y}^{\mathbb{T}}\mathcal{D}_{x}^{\mathbb{T}}z - \mathcal{D}_{\mathcal{D}_{x}^{\mathbb{T}}y}^{\mathbb{T}}z + \mathcal{D}_{\mathcal{D}_{y}^{\mathbb{T}}x}^{\mathbb{T}}z$$

where $x, y, z \in C^{\infty}(\mathbb{T}M)$, then straightforward rearrangements lead to

(2.8)
$$\left[x \diamond_{\mathbb{T}} y \diamond_{\mathbb{T}} z \right] = -\mathcal{R}_{x,y}^{\mathbb{T}} z - c.p.$$

It follows that $d^{\mathbb{T}} \circ d^{\mathbb{T}}$ is tensorial when $\mathcal{D}^{\mathbb{T}}$ is TM-torsion free, since in this case $\mathcal{R}^{\mathbb{T}}$ is tensorial by (3.3). Furthermore, $\mathcal{D}^{\mathbb{T}}$ being TM-torsion free also implies its Jacobiator has values in T^*M :

(2.9)
$$\pi \left[x \diamond_{\mathbb{T}} y \diamond_{\mathbb{T}} z \right] = \left[\left[\pi \left(x \right), \pi \left(y \right) \right], \pi \left(z \right) \right] + c.p. = 0$$

Any torsion free affine connection ∇^T on TM lifts to a TM-torsion free generalized connection on $\mathbb{T}M$:

$$\mathcal{D}_{X}^{\mathbb{T}} y := \nabla_{X}^{T} y = \nabla_{X}^{T} Y + \nabla_{X}^{T} \eta$$

for $x, y \in C^{\infty}(\mathbb{T}M)$ with $x = X + \xi$ and $y = Y + \eta$. The affine space $\mathcal{D}(\mathbb{T}M)$ of generalized connections on $\mathbb{T}M$ is modeled on the space of bundle homomorphisms:

 $\mathcal{D}(\mathbb{T}M) \cong \{A : \mathbb{T}M \otimes \mathbb{T}M \to \mathbb{T}M\}$

in which, the subspace $\mathcal{D}_{\tau}(\mathbb{T}M)$ of TM-torsion free ones is modeled on the subspace of the right hand side consisting of the symmetric ones valued in T^*M :

(2.10)
$$\mathscr{D}(\mathbb{T}M) \supseteq \mathscr{D}_{\tau}(\mathbb{T}M) \cong \{A : \mathbb{T}M \odot \mathbb{T}M \to T^*M\}$$

The *contraction* by $x \in C^{\infty}(\mathbb{T}M)$ is a graded derivation on $\Omega^*_{\pi}(M)$ defined by

where $y \in \Omega^1_{\mathbb{T}}(M)$. The *Lie derivative* along $x \in C^{\infty}(\mathbb{T}M)$ is given by

(2.12)
$$\mathcal{L}_{x}^{\mathsf{T}}\theta := \iota_{x} \mathrm{d}^{\mathsf{T}}\theta + \mathrm{d}^{\mathsf{T}}\iota_{x}\theta$$

where $\theta \in \Omega^*_{\mathbb{T}}(M)$. In particular, for $f \in C^{\infty}(M)$ and $X = \pi(x) \in C^{\infty}(TM)$:

$$\mathcal{L}_{x}^{\mathbb{T}}f = Xf$$

Suppose that $\mathcal{D}^{\mathbb{T}}$ is TM-torsion free, then the familiar relations among the operators $\mathbb{d}^{\mathbb{T}}$, ι_x and $\mathcal{L}_x^{\mathbb{T}}$ almost hold, up to possible terms involving the Jacobiator, similar to (2.5).

Proposition 2.2. Let $x, y, z, w \in C^{\infty}(\mathbb{T}M), \theta \in \Omega^k_{\mathbb{T}}(M), \alpha \in \Omega^1(M)$ and $X = \pi(x)$. Suppose that $\mathcal{D}^{\mathbb{T}}$ is T M-torsion free, then

- (1) $d\alpha = 0 \Longrightarrow d^{\mathsf{T}} \alpha = 0;$
- (2) $[x \diamond_{\mathbb{T}} y \diamond_{\mathbb{T}} z] \in C^{\infty}(T^*M);$
- (3) $X \langle y, z \rangle = \langle x \diamond_{\mathbb{T}} y, z \rangle + \langle y, \mathcal{L}_{x}^{\mathbb{T}} z \rangle;$ (4) $\mathcal{L}_{x}^{\mathbb{T}} \iota_{y} \theta \iota_{y} \mathcal{L}_{x}^{\mathbb{T}} \theta = \iota_{x \diamond_{\mathbb{T}} y} \theta;$

(5)
$$\left\langle \mathcal{L}_{x}^{\mathbb{T}}y - x \diamond_{\mathbb{T}} y, z \right\rangle = \left\langle \mathcal{D}_{y}^{\mathbb{T}}x, z \right\rangle + \left\langle \mathcal{D}_{z}^{\mathbb{T}}x, y \right\rangle;$$

(6) $\left\langle \left[\mathcal{L}_{x}^{\mathbb{T}}, \mathcal{L}_{y}^{\mathbb{T}} \right] z, w \right\rangle = \left\langle \mathcal{L}_{x \diamond_{\mathbb{T}} y}^{\mathbb{T}} z, w \right\rangle + \iota_{\left[x \diamond_{\mathbb{T}} y \diamond_{\mathbb{T}} w\right]} z.$
(7) For $x_{1}, \dots, x_{k+1} \in C^{\infty} (\mathbb{T}M):$
 $\left(\left[d^{\mathbb{T}}, \mathcal{L}_{x}^{\mathbb{T}} \right] \theta \right) (x_{1}, \dots, x_{k+1})$
 $= \sum_{i < j} (-1)^{i+j+1} \theta \left(\left[x \diamond_{\mathbb{T}} x_{i} \diamond_{\mathbb{T}} x_{j} \right], x_{1}, \dots, \hat{x}_{i}, \dots, \hat{x}_{j}, \dots, x_{k+1} \right)$

Proof. The verification follows from standard computations and is left for the reader. \Box

If $\mathcal{D}^{\mathbb{T}}$ is TM-torsion free, by item (1) of Proposition 2.2, the cohomology of $(\Omega^*_{\mathbb{T}}(M), \mathbb{d}^{\mathbb{T}})$ is well-defined for degrees k < 2. Furthermore, the Jacobiator (2.6) defines a degree 2 map $\mathscr{J}^{\mathbb{T}}$ on $\Omega^*_{\mathbb{T}}(M)$, which commutes with $\mathbb{d}^{\mathbb{T}}$

$$(2.13) \qquad \begin{pmatrix} \mathscr{J}^{\mathbb{T}}\theta \end{pmatrix} \begin{pmatrix} x_0, \dots, x_{k+1} \end{pmatrix} := \left(\mathbb{d}^{\mathbb{T}} \circ \mathbb{d}^{\mathbb{T}}\theta \right) \begin{pmatrix} x_0, \dots, x_{k+1} \end{pmatrix} \\ = \sum_{i < j < \ell} (-1)^{i+j+\ell+1} \theta \left(\begin{bmatrix} x_i \diamond_{\mathbb{T}} x_j \diamond_{\mathbb{T}} x_\ell \end{bmatrix}, x_0, \dots, \hat{x}_i, \dots, \hat{x}_j, \dots, \hat{x}_\ell, \dots, x_{k+1} \end{pmatrix}$$

where $\theta \in \Omega^k_{\mathbb{T}}(M)$ and $x_j \in C^{\infty}(\mathbb{T}M)$. It follows that $\mathbb{d}^{\mathbb{T}}$ induces a differential $\widetilde{\mathbb{d}}^{\mathbb{T}}$ on the quotient space of the $\mathcal{D}^{\mathbb{T}}$ -reduced $\mathbb{T}M$ -forms:

(2.14)
$$0 \to \operatorname{img} \mathscr{J}^{\mathbb{T}} \to \Omega^*_{\mathbb{T}}(M) \xrightarrow{\mathscr{R}^{\mathbb{T}}} \widetilde{\Omega}^*_{\mathbb{T}}(M) \to 0$$

where $\mathscr{R}^{\mathbb{T}}$ denotes the quotient map. In particular

(2.15)
$$\widetilde{\Omega}^k_{\mathbb{T}}(M) = \Omega^k_{\mathbb{T}}(M) \text{ for } k \leq 2$$

Definition 2.3. Let $\mathcal{D}^{\mathbb{T}}$ be a *T M*-torsion free generalized connection on $\mathbb{T}M$. The complex $\left(\widetilde{\Omega}^*_{\mathbb{T}}(M), \widetilde{\mathbb{d}}^{\mathbb{T}}\right)$ is the $\mathcal{D}^{\mathbb{T}}$ -de Rham complex and its k-th cohomology is the k-th $\mathcal{D}^{\mathbb{T}}$ -de Rham cohomology of M:

(2.16)
$$\widetilde{H}^{k}_{\mathbb{T}}(M) := \frac{\ker\left(\widetilde{\mathbb{d}}^{\mathbb{T}} : \widetilde{\Omega}^{k}_{\mathbb{T}}(M) \to \widetilde{\Omega}^{k+1}_{\mathbb{T}}(M)\right)}{\operatorname{img}\left(\widetilde{\mathbb{d}}^{\mathbb{T}} : \widetilde{\Omega}^{k-1}_{\mathbb{T}}(M) \to \widetilde{\Omega}^{k}_{\mathbb{T}}(M)\right)}$$

Because $\mathscr{J}^{\mathbb{T}}$ is of degree 2, when k < 2, $\widetilde{H}^{k}_{\mathbb{T}}(M)$ compute the corresponding cohomology groups of $\Omega^{*}_{\mathbb{T}}(M)$. By (2.3), it is evident that $\widetilde{H}^{0}_{\mathbb{T}}(M) = \mathbb{R} = H^{0}(M)$, which consists of the constant functions. The item (1) in Proposition 2.2 then gives a natural inclusion

(2.17)
$$H^1(M) \subseteq \widetilde{H}^1_{\pi}(M)$$

in which the equality may not hold in general (c.f. Proposition 2.11).

In general, the map π induces a natural injection π^* : $\Omega^k(M) \hookrightarrow \Omega^k_{\pi}(M)$ for all k:

(2.18)
$$(\pi^*\alpha)(x_1,\ldots,x_k) := \alpha(\pi(x_1),\ldots,\pi(x_k))$$

where $\alpha \in \Omega^k(M)$ and $x_j \in C^{\infty}(\mathbb{T}M)$. Alternatively, π^* is induced from the inclusion $T^*M \hookrightarrow \mathbb{T}M$. When $\mathcal{D}^{\mathbb{T}}$ is TM-torsion free, π^* commutes with the derivations

(2.19)
$$\pi^*(d\alpha) = \mathbb{d}^{\mathbb{I}}(\pi^*\alpha)$$

Thus π^* defines a morphism of cochain complices after passing to the quotient

$$\widetilde{\pi}^*$$
: $\Omega^k(M) \to \widetilde{\Omega}^k_{\mathbb{T}}(M)$

which induces the corresponding maps on the cohomology groups:

(2.20)
$$\widetilde{\pi}^* : H^k(M) \to \widetilde{H}^k_{\pi}(M)$$

2.2. (ϕ, b) -connections. To restrict $\mathcal{D}^{\mathbb{T}}$ further, consider a generalized metric \mathbb{G}^{b} (1.3), corresponding to the pair (g, b) of Riemannian metric g on M and 2-form $b \in \Omega^{2}(M)$.

Definition 2.4. A generalized metric connection with respect to \mathbb{G}^b , or a \mathbb{G}^b -metric connection, is a generalized connection $\mathcal{D}^{\mathbb{T}}$ on $\mathbb{T}M$ that preserves both \mathbb{G}^b and \langle, \rangle :

(2.21)
$$X \langle y, z \rangle = \langle D_x^{\mathbb{T}} y, z \rangle + \langle y, D_x^{\mathbb{T}} z \rangle$$
 and $X \mathbb{G}^b (y, z) = \mathbb{G}^b \left(D_x^{\mathbb{T}} y, z \right) + \mathbb{G}^b \left(y, D_x^{\mathbb{T}} z \right)$

where $x, y, z \in C^{\infty}(\mathbb{T}M)$ and $X = \pi(x)$.

A \mathbb{G}^b -metric connection $\mathcal{D}^{\mathbb{T}}$ preserves the ± 1 -eigenbundles C^b_+ of \mathbb{G}^b :

(2.22)
$$C_{\pm}^{b} := \left\{ x^{b\pm} := X + (b\pm g) X : X \in TM \right\}$$

which induces a \mathbb{G}^{b} -eigendecomposition into a quadruple of metric connections ∇_{\star}^{\bullet} on TM:

(2.23)
$$\nabla^{\bullet}_{\star,X}Y := \pi \left(\mathcal{D}^{\mathbb{T}}_{x^{b^{\star}}} y^{b^{\bullet}} \right)$$

where \star and \bullet respectively stand for + or -. Furthermore, the corresponding $\mathbb{d}^{\mathbb{T}}$ admits the induced \mathbb{G}^{b} -eigendecomposition.

Lemma 2.5. Let $\mathcal{D}^{\mathbb{T}}$ be a \mathbb{G}^{b} -metric connection, then the operator $\mathbb{d}^{\mathbb{T}}$ decomposes into components as follows:

(2.24)
$$\mathbb{d}^{\mathbb{T}} = \mathbb{d}^{\mathbb{T}}_{+} + \mathbb{d}^{\mathbb{T}}_{-} : \Omega^{p,q}_{\mathbb{G}}(M) \to \Omega^{p+1,q}_{\mathbb{G}}(M) \oplus \Omega^{p,q+1}_{\mathbb{G}}(M)$$

where

(2.25)
$$\Omega^{p,q}_{\mathbb{G}}(M) := C^{\infty} \left(\wedge^{p} C^{b}_{+} \otimes \wedge^{q} C^{b}_{-} \right) \cong \Omega^{p}(M) \otimes \Omega^{q}(M)$$

Proof. Consider $\theta \in \Omega^{p,0}_{\mathbb{G}}(M)$. For k > 1, since $\mathcal{D}^{\mathbb{T}}$ preserves C^b_{\pm} , it is straightforward to verify that for $x_j^{b\pm} \in C^{\infty}(C^b_{\pm})$

$$\left(\mathbb{d}^{\mathbb{T}}\theta\right)\left(x_{0}^{b-}, x_{1}^{b-}, \dots, x_{k-1}^{b-}, x_{k}^{b+}, \dots, x_{q}^{b+}\right) = 0$$

Thus $\mathbb{d}^{\mathbb{T}}\theta$ cannot contain any components in $\Omega_{\mathbb{G}}^{p-k+1,k}(M)$ for k > 1. The general situation follows from the analogue for $\theta' \in \Omega_{\mathbb{G}}^{0,q}(M)$ then noticing that $\mathbb{d}^{\mathbb{T}}$ is a derivation (2.4). \square

10

In its \mathbb{G}^{b} -eigendecomposition, a \mathbb{G}^{b} -metric connection $\mathcal{D}^{\mathbb{T}}$ is TM-torsion free iff $\nabla^{+}_{+} = \nabla^{-}_{-} = \nabla$ is the Levi-Civita connection for g and for all $X, Y \in C^{\infty}(TM)$:

$$\nabla_{+,X}^{-}Y - \nabla_{-,Y}^{+}X - [X,Y] = 0$$

It follows that a 3-form $\phi \in \Omega^3(M)$, which may not be closed, can be defined by

(2.26)
$$\phi(X, Y, Z) := 2g((\nabla_{-}^{+} - \nabla)_{X}Y, Z) = 2g((\nabla_{+}^{-} - \nabla)_{Y}X, Z)$$

The mixed components in the \mathbb{G}^{b} -eigendecomposition can be expressed in terms of ϕ , e.g.

(2.27)
$$\nabla_{-,X}^{+}Y = \nabla_{X}^{+\phi}Y := \nabla_{X}Y + \frac{1}{2}g^{-1}\iota_{Y}\iota_{X}\phi$$

which is a metric connection on TM with totally skew torsion ϕ . The computations are summarized in the following Theorem / Definition.

Theorem 2.6 ((ϕ , b)-connections). *The* TM*-torsion free* \mathbb{G}^{b} *-metric connections on* $\mathbb{T}M$ are classified by 3-forms $\phi \in \Omega^{3}(M)$, which is denoted $D^{\phi,b}$, and referred to as the (ϕ , b)-connection. The \mathbb{G}^{b} -eigendecomposition of $D^{\phi,b}$ is explicitly presented below:

(2.28)
$$\mathcal{D}_{x^{b+}}^{\phi,b} y^{b+} = \nabla_X Y + (b+g) \left(\nabla_X Y \right) \text{ and } \mathcal{D}_{x^{b+}}^{\phi,b} y^{b-} = \nabla_X^{-\phi} Y + (b-g) \left(\nabla_X^{-\phi} Y \right) \\ \mathcal{D}_{x^{b-}}^{\phi,b} y^{b+} = \nabla_X^{+\phi} Y + (b+g) \left(\nabla_X^{+\phi} Y \right) \text{ and } \mathcal{D}_{x^{b-}}^{\phi,b} y^{b-} = \nabla_X Y + (b-g) \left(\nabla_X Y \right)$$

where $X, Y \in C^{\infty}(TM)$, $x^{b\pm} = X + (b \pm g) X \in C^{\infty}(C^{b}_{\pm})$ and so on. In the splitting $\mathbb{T}M = TM \oplus T^*M$, the (ϕ, b) -connection takes the form:

(2.29)
$$\mathcal{D}_{e^{b_{X}}}^{\phi,b}e^{b_{Y}} = e^{b} \left[\nabla_{X}y + \frac{1}{4}g^{-1} \left(\iota_{g^{-1}\eta}\iota_{X}\phi - \iota_{Y}\iota_{g^{-1}\xi}\phi \right) + \frac{1}{4} \left(\iota_{Y}\iota_{X}\phi - \iota_{g^{-1}\eta}\iota_{g^{-1}\xi}\phi \right) \right]$$

where $x = X + \xi$, $y = Y + \eta \in C^{\infty}(\mathbb{T}M)$.

Proof. Under the pairing $2\langle,\rangle$, the space $\mathcal{D}(\mathbb{T}M)$ of generalized connections on $\mathbb{T}M$ is modeled on

$$\mathcal{D}\left(\mathbb{T}M\right) \cong \{A : \mathbb{T}M \to \mathbb{T}M \otimes \mathbb{T}M\}$$

The subspace $\mathscr{D}_{\mathbb{G}^b}(\mathbb{T}M)$ of \mathbb{G}^b -metric connections is then modeled on

(2.30)
$$\mathscr{D}_{\mathbb{G}^{b}}(\mathbb{T}M) \cong \left\{ A : \mathbb{T}M \to \left(\wedge^{2}C_{+}^{b}\right) \oplus \left(\wedge^{2}C_{-}^{b}\right) \cong \operatorname{End}\left(TM,g\right)^{\oplus 2} \right\}$$

From (2.10) and (2.30), it follows that space of TM-torsion free \mathbb{G}^b -metric connections on $\mathbb{T}M$ is modeled on the intersection. The identity (2.26) can be seen also as

$$\phi(X, Y, Z) = 2 \left\langle A_{x^{b-}} y^{b+}, z^{b+} \right\rangle = -2 \left\langle A_{y^{b+}} x^{b-}, z^{b-} \right\rangle$$

which shows that the intersection is isomorphic to $\Omega^3(M)$. The rest follows from straightforward computations.

Using the explicit formulae in (2.28), it is straightforward to verify that $\mathcal{D}^{\phi,b}$ is metric compatible with the (almost) Dorfman bracket $*_{\phi-db}$, in the sense of Definition 2.7 below.

Definition 2.7. A \mathbb{G}^{b} -metric connection $\mathcal{D}^{\mathbb{T}}$ is *metric compatible with* $*_{\gamma}$ (1.4) if the diamond bracket $\diamond_{\mathbb{T}}$ coincides with $*_{\gamma}$ on mixed \mathbb{G}^{b} -eigensections, i.e.

(2.31)
$$x^{b+} \diamond_{\mathbb{T}} y^{b-} = x^{b+} *_{\gamma} y^{b-}$$

where $x^{b+} \in C^{\infty}(C^b_+)$ and $y^{b-} \in C^{\infty}(C^b_-)$.

The space of \mathbb{G}^{b} -metric connections that are metric compatible with $*_{\gamma}$ but not necessarily TM-torsion free is modeled on the following subspace of (2.30):

$$\mathscr{D}_{\mathbb{G}^{b},\gamma}(\mathbb{T}M) \cong \left\{ A : C^{b}_{+} \to \wedge^{2}C^{b}_{+} \right\} \oplus \left\{ A : C^{b}_{-} \to \wedge^{2}C^{b}_{-} \right\} \cong \Omega^{1}\left(\operatorname{End}\left(TM,g\right)\right)^{\oplus 2}$$

Theorem 2.8. For a closed form $\gamma \in \Omega^3(M)$, let $\phi = \gamma + db$. Then $\mathcal{D}^{\phi,b}$ is the unique *TM*-torsion free \mathbb{G}^b -metric connection on $\mathbb{T}M$ that is metric compatible with $*_{\gamma}$.

2.3. $\mathcal{D}^{\phi,b}$ -de Rham cohomology. The \mathbb{G}^{b} -eigendecomposition of $\mathbb{T}M$ induces two left inverses of the natural injection π^{*} in (2.18). There are two obvious projections p^{\pm} : $\Omega^{k}_{\mathbb{T}}(M) \to C^{\infty}(\wedge^{k}C^{b}_{+})$ for each k:

$$(p^{\pm}\theta)(x_1^{b\pm},\ldots,x_k^{b\pm}) := \theta(x_1^{b\pm},\ldots,x_k^{b\pm})$$

where $\theta \in \Omega^k_{\mathbb{T}}(M)$, $X_j \in C^{\infty}(TM)$ and $x_j^{b\pm} = X_j + (b \pm g) X_j$ for all *j*. The projection π induces the natural isomorphisms $\pi_* : C^{\infty}(\wedge^* C^b_{\pm}) \cong \Omega^*(M)$:

 $(\pi_*\theta_{\pm})(X_1,\ldots,X_k) := \theta_{\pm}(x_1^{b\pm},\ldots,x_k^{b\pm})$

where $\theta_{\pm} \in C^{\infty}\left(\wedge^{*}C_{\pm}^{b}\right)$. Then $\pi_{*}^{\pm} = \pi_{*}\circ p^{\pm}$: $\Omega_{\mathbb{T}}^{k}(M) \to \Omega^{k}(M)$ is given by

$$\left(\pi_{*}^{\pm}\theta\right)\left(X_{1},\ldots,X_{k}\right) := \theta\left(x_{1}^{b\pm},\ldots,x_{k}^{b\pm}\right)$$

It is now straightforward to see that

$$\pi^{\pm}_{*} \circ \pi^{*} = \mathrm{id} : \Omega^{*}(M) \to \Omega^{*}(M)$$

Let $\mathbb{d}^{\phi,b}$ denote the $\mathcal{D}^{\phi,b}$ -derivation (2.2) induced by $\mathcal{D}^{\phi,b}$. The decomposition (2.24) can be explicitly described by the classical de Rham differential and covariant derivatives. For instance, for $\alpha \in \Omega^k(M)$, let $\alpha_+ \in \Omega^{p,0}_{\mathbb{G}}(M)$ such that $\alpha = \pi^+_*(\alpha_+)$. Then $\mathbb{d}^{\phi,b}_+ \alpha_+$ is essentially the de Rham differential:

$$(\mathbb{d}^{\phi,b}\alpha_{+}) \left(x_{0}^{b+}, x_{1}^{b+}, \dots, x_{k}^{b+}\right) = \sum_{i} (-1)^{i} \left(\mathcal{D}_{x_{i}^{b+}}^{\phi,b}\alpha_{+}\right) \left(x_{0}^{b+}, x_{1}^{b+}, \dots, \widehat{x_{i}^{b+}}, \dots, x_{k}^{b+}\right)$$
$$= \sum_{i} (-1)^{i} \left(\nabla_{x_{i}}\alpha\right)_{+} \left(x_{0}^{b+}, x_{1}^{b+}, \dots, \widehat{x_{i}^{b+}}, \dots, x_{k}^{b+}\right) = (d\alpha)_{+} \left(x_{0}^{b+}, x_{1}^{b+}, \dots, x_{k}^{b+}\right)$$

while the component $\mathbb{d}_{-}^{\phi,b}\alpha_{+} \in \Omega_{\mathbb{G}}^{p,1}(M)$ is essentially given by $\nabla^{+\phi}$:

(2.33)
$$\left(\mathbb{d}_{-}^{\phi,b}\alpha_{+}\right)\left(x_{0}^{b-},x_{1}^{b+},\ldots,x_{q}^{b+}\right) = \left(\nabla_{X_{0}}^{+\phi}\alpha\right)_{+}\left(x_{1}^{b+},\ldots,x_{q}^{b+}\right)$$

where $x_j^{b\pm} \in C^{\infty}(C_{\pm}^b)$, and $\pi(x_j^{b\pm}) = X_j$.

Lemma 2.9. The Jacobiator for $\diamond_{\phi,b}$ is given by

(2.34)
$$\left[x^{b\pm} \diamond_{\phi,b} y^{b\pm} \diamond_{\phi,b} z^{b\mp}\right] = \pm 2g\left(R_{X,Y}^{\mp\phi}Z\right)$$

where $x^{b\pm}$, $y^{b\pm}$, $z^{b\pm} \in C^{\infty}(C^{b}_{\pm})$ with $\pi(x^{b\pm}) = X$ and so on, and $R^{\pm\phi}$ are respectively the classical curvature for $\nabla^{\pm\phi}$. All other components in the \mathbb{G}^{b} -eigendecomposition vanish.

Proof. Follows by relatively lengthy but standard computations from the definitions. \Box

Proposition 2.10. The natural map $\widetilde{\pi}^*$: $H^*(M) \to \widetilde{H}^*_{\phi,b}(M)$ in (2.20) is injective.

Proof. For $\alpha, \beta \in \Omega^*(M)$, suppose that $\pi^* \alpha = \pi^*(d\beta)$, then the injectivity of π^* implies that $\alpha = d\beta$. The statement then follows from $\operatorname{img} \mathcal{J}^{\phi,b} \cap \operatorname{img} \pi^* = \{0\}$. In fact, suppose $\mathcal{J}^{\phi,b}\theta = \pi^*\alpha \in \operatorname{img} \mathcal{J}^{\phi,b} \cap \operatorname{img} \pi^*$, then

$$\alpha = \pi^+_*(\pi^*\alpha) = \pi^+_*\left(\mathcal{J}^{\phi,b}\theta\right) = 0$$

The last equality is due to $[x^{b+} \diamond_{\phi,b} y^{b+} \diamond_{\phi,b} z^{b+}] = 0$, which follows from Lemma 2.9.

Since $\widetilde{\Omega}^k_{\mathbb{T}}(M) \cong \Omega^k_{\mathbb{T}}(M)$ for $k \leq 2$ and the derivations coincide for k < 2, the groups $\widetilde{H}^k_{\phi,b}(M)$ for k < 2 can be determined. When k = 0, it is straightforward to see that $\widetilde{H}^0_{\phi,b}(M) \cong \mathbb{R}$ consists of the constant functions.

Proposition 2.11. Let $P_{\phi}^{1}(M)$ be the space of ∇ -parallel 1-forms on M, which also annilates ϕ , i.e.

$$\xi \in P^1_{\phi}(M) \iff \nabla \xi = 0 \text{ and } \iota_{g^{-1}\xi} \phi = 0$$

then

(2.35)
$$\widetilde{H}^{1}_{\phi,b}(M) \cong H^{1}(M) \oplus P^{1}_{\phi}(M)$$

Proof. Let $\theta \in \Omega^1_{\mathbb{T}}(M)$. For $X, Y \in C^{\infty}(TM)$, let $x^{b\pm} = X + (b \pm g) X$ and so on, define

(2.36)
$$\alpha(X) := \theta(x^{b+}) \text{ and } \beta(X) := \theta(x^{b-})$$

It then gives the identification

(2.37)
$$Q: \Omega^{1}_{\mathbb{T}}(M) \cong \Omega^{1}(M) \oplus \Omega^{1}(M): \theta \mapsto \frac{1}{2}(\alpha + \beta, \alpha - \beta)$$

Suppose that $d^{\phi,b}\theta = 0$, then (2.32) implies that

$$d\alpha = d\beta = 0$$

Hence

$$0 = \left(\mathbb{d}_{-}^{\phi,b}\theta\right)\left(x^{b+}, y^{b-}\right) = -\left[\nabla_{Y}^{+\phi}\left(\alpha - \beta\right)\right](X)$$

i.e. $\nabla^{+\phi} (\alpha - \beta) = 0$. Set $\xi = \alpha - \beta$, then $d\xi = 0$ implies that

$$0 = \left(\nabla_X^{+\varphi}\xi\right)(Y) - \left(\nabla_Y^{+\varphi}\xi\right)(X) = -\phi\left(X, Y, g^{-1}\xi\right)$$

for all $X, Y \in C^{\infty}(TM)$, from which $\xi \in P_{\phi}^{1}(M)$ follows. Thus on $\Omega_{\mathbb{T}}^{1}(M)$

$$Q\left(\ker d^{\phi,b}\right) = \ker d \oplus P^{1}_{\phi}(M)$$

Then (2.35) follows from Q(df) = (df, 0) for $f \in C^{\infty}(M)$.

Corollary 2.12. $\widetilde{H}^1_{\phi,h}(M) \cong H^1(M)$ if one of the following holds

(1) ϕ is non-degenerate, i.e. the following map is injective

$$\iota_{\bullet}\phi: C^{\infty}(TM) \to \Omega^{2}(M): X \mapsto \iota_{X}\phi$$

(2) *M* admits no non-trivial ∇ -parallel vector fields.

Example 2.13. Let $M = \mathbb{R}^n / \mathbb{Z}^n$, with the induced flat metric. Suppose that $\phi = 0$, then the ∇ -parallel forms on M are the constant forms. Thus

$$\widetilde{H}^1_{0,b}(M) \cong H^1(M) \oplus T_0 M \cong \mathbb{R}^{2\prime}$$

where T_0M is the tangent space at $0 \in M$. On the other hand, for n = 3, let $\phi = \operatorname{vol}_g$, the volume form of the flat metric on M, then it is non-degenerate. By Corollary 2.12

$$\widetilde{H}^{1}_{\mathrm{vol}_{g},b}(M) = H^{1}(M) \cong \mathbb{R}^{3}$$

Let $n = \dim_{\mathbb{R}} M$. The group $H^{2n}_{\mathbb{T}}(M)$ is always well-defined for a *T M*-torsion free $\mathcal{D}^{\mathbb{T}}$

$$H^{2n}_{\mathbb{T}}(M) := \frac{\Omega^{2n}_{\mathbb{T}}(M)}{\operatorname{img}\left(\mathbb{d}^{\mathbb{T}} : \Omega^{2n-1}_{\mathbb{T}}(M) \to \Omega^{2n}_{\mathbb{T}}(M)\right)}$$

When $\mathcal{D}^{\mathbb{T}} = \mathcal{D}^{\phi,b}$, the groups $\widetilde{H}^{2n}_{\phi,b}(M)$ and $H^{2n}_{\phi,b}(M)$ can be determined.

Proposition 2.14. For any $(\phi, b) \in \Omega^3(M) \oplus \Omega^2(M)$, $\widetilde{H}^{2n}_{\phi,b}(M) \cong H^{2n}_{\phi,b}(M) \cong \mathbb{R}$.

Proof. The derivation $\mathbb{Q}^{\phi,b}$ on $\Omega^{2n-1}_{\mathbb{T}}(M)$ splits in the decomposition (2.25) as

$$\mathbb{d}^{\phi,b} = \mathbb{d}^{\phi,b}_{+} \oplus \mathbb{d}^{\phi,b}_{-} : \Omega^{n-1,n}_{\mathbb{T}}(M) \oplus \Omega^{n,n-1}_{\mathbb{T}}(M) \to \Omega^{n,n}_{\mathbb{T}}(M) = \Omega^{2n}_{\mathbb{T}}(M)$$

For instance, $\mathbb{d}^{\phi,b}_+$: $\Omega^{n-1,n}_{\mathbb{T}}(M) \to \Omega^{n,n}_{\mathbb{T}}(M)$ is given by the lifting of d: $\Omega^{n-1}(M) \to \Omega^n(M)$ as in (2.32). It follows that

$$\widetilde{H}^{2n}_{\phi,b}(M)\cong H^{2n}_{\phi,b}(M)\cong \mathbb{R}$$

since $\operatorname{img} \mathcal{J}^{\phi,b} \cap \Omega^{2n}(M) = \{0\}$ by Proposition 4.5, which gives $\widetilde{\Omega}^{2n}_{\mathbb{T}}(M) = \Omega^{2n}_{\mathbb{T}}(M)$.

Let *M* and *M'* be two smooth manifolds. It is straightforward to see that if a diffeomorphism $\lambda : M \to M'$ relates the corresponding data on both manifolds, i.e.

$$(g, b, \phi) = (\lambda^* g', \lambda^* b', \lambda^* \phi')$$

it induces isomorphisms throughout the constructions. In particular, it induces the natural isomorphism of cohomology groups $\lambda^* : \widetilde{H}^*_{\phi',b'}(M') \xrightarrow{\cong} \widetilde{H}^*_{\phi,b}(M)$.

2.4. **Laplacians.** Analogous to the classical case, a generalized connection \mathcal{D} on V defines the corresponding \mathcal{D} -Bochner Laplacian on $C^{\infty}(V)$. Let $\{X_i\}$ be a local orthonormal frame on TM for the Riemannian metric g and let $\{e_i^{b\pm}\}$ be the corresponding local \mathbb{G}^{b} -orthonormal frame on $\mathbb{T}M$:

$$e_i^{b\pm} := e^b \left[X_i \pm g \left(X_i \right) \right]$$

Definition 2.15. Let \mathcal{D} be a generalized connection on V. The \mathcal{D} -Bochner Laplacian (with respect to $\mathcal{D}^{\phi,b}$) on V is given for $v \in C^{\infty}(V)$ by:

(2.38)
$$\Delta_D v := -\sum_i \left[\left(\mathcal{D}_{e_i^{b+}} \mathcal{D}_{e_i^{b+}} - \mathcal{D}_{\mathcal{D}_{e_i^{b+}}^{\phi,b}} e_i^{b+} \right) v + \left(\mathcal{D}_{e_i^{b-}} \mathcal{D}_{e_i^{b-}} - \mathcal{D}_{\mathcal{D}_{e_i^{b-}}^{\phi,b}} e_i^{b-} \right) v \right]$$

It is straightforward to verify that (2.38) is independent of the choice of $\{X_i\}$. Because $\mathcal{D}_{e_i^{b_{\pm}}}^{\phi,b}e_i^{b_{\pm}}$ involve only the Levi-Civita connection for g, it is evident that

$$\Delta_{\mathcal{D}} v = \left(\Delta_{+}^{b} + \Delta_{-}^{b}\right) v$$

where Δ_{\pm}^{b} are the classical Bochner Laplacians for ∇_{\pm}^{b} respectively. Thus Δ_{D} is a second order elliptic operator, which in general depends on both *g* and *b*, i.e. on \mathbb{G}^{b} , but not on ϕ . When D is a lift of a classical connection, Δ_{D} reduces to (a constant multiple of) the corresponding classical Bochner Laplacian.

The operator $d^{\phi,b}$ can alternatively be written as

(2.39)
$$\mathbb{d}^{\phi,b}\theta = \sum_{j} \left(e_{j}^{b+} \wedge \mathcal{D}_{e_{j}^{b+}}^{\phi,b}\theta - e_{j}^{b-} \wedge \mathcal{D}_{e_{j}^{b-}}^{\phi,b}\theta \right)$$

where $\theta \in \Omega^*_{\pi}(M)$. Thus the symbol of $\mathbb{d}^{\phi,b}$ is

$$\sigma\left(\mathbb{d}^{\phi,b}\right) = 2\sqrt{-1}\xi\wedge : \wedge^{k}\mathbb{T}M \to \wedge^{k+1}\mathbb{T}M$$

where $\xi \in T^*M$. It follows that whenever $\mathbb{d}^{\phi,b}$ squares to 0 it defines an elliptic complex. Analogous to the classical case, for $\theta \in \Omega^*_{\mathbb{T}}(M)$, define

(2.40)
$$\mathbb{d}^{\phi,b*}\theta := -\frac{1}{2}\sum_{j} \left(\iota_{e_{j}^{b+}} \mathcal{D}_{e_{j}^{b+}}^{\phi,b}\theta + \iota_{e_{j}^{b-}} \mathcal{D}_{e_{j}^{b-}}^{\phi,b}\theta \right)$$

where the $\frac{1}{2}$ is due to the convention (2.11). The symbol of $d^{\phi,b*}$ is then

$$\sigma\left(\mathbb{d}^{\phi,b*}\right) = -\sqrt{-1}\iota_{g^{-1}\xi + bg^{-1}\xi} : \wedge^{k}\mathbb{T}M \to \wedge^{k-1}\mathbb{T}M$$

where $\xi \in T^*M$.

Definition 2.16. The $\mathcal{D}^{\phi,b}$ -Hodge Laplacian is the operator on $\Omega^*_{\mathbb{T}}(M)$ given by

(2.41)
$$\Delta^{\phi,b} := \mathrm{d}^{\phi,b} \mathrm{d}^{\phi,b*} + \mathrm{d}^{\phi,b*} \mathrm{d}^{\phi,b}$$

Proposition 2.17. The $\mathcal{D}^{\phi,b}$ -Hodge Laplacian $\Delta^{\phi,b}$ is a second order elliptic operator.

Proof. It is clear from the discussions above that the symbol of $\Delta^{\phi,b}$ is

$$\sigma\left(\Delta^{\phi,b}\right) = 2\|\xi\|_g^2 : \wedge^k \mathbb{T}M \to \wedge^k \mathbb{T}M$$

where $\xi \in T^*M$, from which the statement follows.

Theorem 2.18. Let M be a closed manifold. Then the following holds:

(2.42)
$$\widetilde{H}^{1}_{\phi,b}(M) \cong \ker \mathbb{d}^{\phi,b} \big|_{\Omega^{1}_{\mathbb{T}}(M)} \cap \ker \mathbb{d}^{\phi,b*} \big|_{\Omega^{1}_{\mathbb{T}}(M)} \subseteq \ker \Delta^{\phi,b} \big|_{\Omega^{1}_{\mathbb{T}}(M)}$$

Proof. Let $\theta \in \Omega^1_{\mathbb{T}}(M)$ and consider $\alpha, \beta \in \Omega^1(M)$ as in (2.36), then it can be shown that

$$\mathbb{d}^{\phi,b*}\theta = d^* \left(\alpha + \beta\right)$$

Combining the identity above with the proof of Proposition 2.11, the first isomorphism follows from the classical Hodge theorem. The last inclusion is obvious. \Box

Even though $(\Omega^*_{\mathbb{T}}(M), \mathbb{d}^{\phi,b})$ is generally not a chain complex, Theorem 2.18 nonetheless hints at the analogue of its "cohomology groups".

Definition 2.19. Let \mathbb{G}^b be a generalized metric on M. The (ϕ, b) -pseudo-cohomology groups $\check{H}^*_{\phi,b}(M)$ of M consist of the intersection of the kernels of $\mathbb{d}^{\phi,b}$ and $\mathbb{d}^{\phi,b*}$:

$$\check{H}^{k}_{\phi,b}(M) := \ker d^{\phi,b} \big|_{\Omega^{k}_{\mathbb{T}}(M)} \cap \ker d^{\phi,b*} \big|_{\Omega^{k}_{\mathbb{T}}(M)}$$

The (ϕ, b) -Laplacian kernels $\widehat{H}^k_{\phi, b}(M)$ are the subspaces

$$\widehat{H}^{k}_{\phi,b}(M) := \ker \Delta^{\phi,b} \big|_{\Omega^{k}_{\pi}(M)}$$

Corollary 2.20. For a closed manifold M, both $\hat{H}^*_{\phi,b}(M)$ and $\check{H}^*_{\phi,b}(M)$ are finite dimensional.

Proof. It follows from $\check{H}^k_{\phi,b}(M) \subseteq \widehat{H}^k_{\phi,b}(M)$ and the ellipticity of $\Delta^{\phi,b}$.

Remark 2.21. The inclusions $\check{H}_{\phi,b}^{k}(M) \subseteq \widehat{H}_{\phi,b}^{k}(M)$ may not be equalities. For k = 1, direct computation using the \mathbb{G}^{b} -eigendecomposition shows that

$$\theta = \hat{H}^{1}_{\phi,b}(M) \iff \begin{cases} \Delta(\alpha + \beta) + \Delta_{+\phi}(\alpha - \beta) - \frac{1}{2} \sum_{j,k} (d\beta) \left(X_{j}, X_{k}\right) \iota_{X_{k}} \iota_{X_{j}} \phi = 0\\ \Delta(\alpha + \beta) - \Delta_{-\phi}(\alpha - \beta) + \frac{1}{2} \sum_{j,k} (d\alpha) \left(X_{j}, X_{k}\right) \iota_{X_{k}} \iota_{X_{j}} \phi = 0 \end{cases}$$

where α , β are as in (2.36) and $\{X_j\}$ is a local *g*-orthonormal frame of *TM*. In the case that $d\alpha = d\beta = 0$, it can be shown that the right hand side is exactly equivalent to $\theta \in \check{H}^1_{\phi,b}(M)$. Namely, under the identification (2.37)

$$\check{H}^{1}_{\phi,b}(M) = \widehat{H}^{1}_{\phi,b}(M) \cap (\ker d \oplus \ker d)$$

П

2.5. Compact Lie groups. Manifolds admitting flat metric connections with non-trivial completely skew torsions are known by Cartan and Schouten ([6], [7]), which are essentially compact Lie groups and S^7 (see also Agricola-Friedrich [1]).

Suppose that *G* is a real semi-simple Lie group, endowed with the bi-invariant Killing metric *g* and the corresponding bi-invariant Cartan 3-form $\gamma \in \Omega^3(G)$. In this case, as will become clear below, the computations are very much parallel to those for the classical de Rham cohomology of the doubled group $G \times G$.

The metric connections $\nabla^{\pm \gamma}$, with torsions $\pm \gamma$ respectively, are flat. Let $(\phi, b) = (\gamma, 0)$ and consider the $(\gamma, 0)$ -connection $\mathcal{D}^{\gamma, 0}$. By Lemma 2.9, for all $x, y, z \in C^{\infty}(\mathbb{T}G)$

$$\left[x \diamond_{\gamma,0} y \diamond_{\gamma,0} z\right] = 0$$

which implies that $\mathscr{J}^{\gamma,0} = \mathbb{d}^{\gamma,0} \circ \mathbb{d}^{\gamma,0} = 0$. Hence $\left(\Omega^*_{\mathbb{T}}(G), \mathbb{d}^{\gamma,0}\right)$ is the $\mathcal{D}^{\gamma,0}$ -de Rham complex, whose cohomology is the $\mathcal{D}^{\gamma,0}$ -de Rham cohomology $\widetilde{H}^*_{\gamma,0}(G)$.

Let X_u^l denote the left invariant vector field on G such that $X_u^l(e) = u \in \mathfrak{g} := T_e G$. The corresponding right invariant vector fields are denoted X_u^r . The Lie algebra structure on \mathfrak{g} is identified with the Lie algebra of left invariant vector fields:

$$\left[X_{u}^{l}, X_{v}^{l}\right] = X_{\left[u, v\right]}^{l}$$

For $u, v \in \mathfrak{g}$, set $\theta_u^r := g(X_u^r)$ and $\theta_v^l := g(X_v^l)$, then $x^+ = X^r + \theta^r \subset C^{\infty}(C)$ and $x^- = X^l - \theta^l \in C^{\infty}(C)$

$$x_u^+ = X_u^r + \theta_u^r \in C^\infty(C_+) \text{ and } x_v^- = X_v^t - \theta_v^t \in C^\infty(C_-)$$

It is straightforward to see that

$$\mathcal{D}_{x_{u}^{+}}^{\gamma,0}x_{v}^{+} = -\frac{1}{2}x_{[u,v]}^{+}, \mathcal{D}_{x_{u}^{-}}^{\gamma,0}x_{v}^{-} = \frac{1}{2}x_{[u,v]}^{-}, \text{ and } \mathcal{D}_{x_{u}^{+}}^{\gamma,0}x_{v}^{-} = \mathcal{D}_{x_{u}^{-}}^{\gamma,0}x_{v}^{+} = 0$$

Let $\mathbf{u} = (u, u'), \mathbf{v} = (v, v') \in \mathbf{g} \oplus \mathbf{g}$ and

$$x_{u} = -x_{u}^{+} + x_{u'}^{-}, x_{v} = -x_{v}^{+} + x_{v'}^{-} \in C^{\infty} (\mathbb{T}G)$$

then direct computation leads to

(2.43)
$$\mathcal{D}_{x_{\mathfrak{u}}}^{\gamma,0} x_{\mathfrak{v}} = \frac{1}{2} x_{[\mathfrak{u},\mathfrak{v}]} \Longrightarrow x_{\mathfrak{u}} \diamond_{\gamma,0} x_{\mathfrak{v}} = x_{\mathfrak{u}} *_{\gamma} x_{\mathfrak{v}} = x_{[\mathfrak{u},\mathfrak{v}]}$$

where the Lie bracket on $\mathfrak{g} \oplus \mathfrak{g}$ is the direct sum of those on each factor. The last equality in (2.43) can be seen also from the *Courant trivialization* of Alekseev-Bursztyn-Meinrenken [2]. The (γ , 0)-curvature can then be computed as

$$\mathcal{R}_{x_{\mathrm{u}},x_{\mathrm{v}}}^{\gamma,0}x_{\mathrm{w}} = -\frac{1}{4}\left[\left[x_{\mathrm{u}},x_{\mathrm{v}}\right],x_{\mathrm{w}}\right] = -\frac{1}{4}x_{\left[\left[u,v\right],w\right]}$$

which gives the $(\gamma, 0)$ -Ricci tensor

$$\mathcal{R}c^{\gamma,0}\left(x_{\mathrm{u}},x_{\mathrm{v}}\right) = \frac{1}{4}\mathbb{G}\left(x_{\mathrm{u}},x_{\mathrm{v}}\right)$$

In particular, $(G, \gamma; \mathbb{G})$ may be seen as an example of a $(\gamma, 0)$ -*Einstein manifold*, where the $(\gamma, 0)$ -Ricci curvature is proportional to the generalized metric.

To compute $\widetilde{H}_{\gamma,0}(G)$, recall that $\mathbb{T}G$ is the dual of itself via $2\langle , \rangle$, which leads to

$$\left(\mathbb{d}^{\gamma,0}x_{\mathrm{w}}\right)\left(x_{\mathrm{v}},x_{\mathrm{w}}\right) = -x_{\mathrm{w}}\left(x_{[\mathrm{v},\mathrm{w}]}\right)$$

Let $\theta \in \Omega^*_{\mathbb{T}}(G)$ be decomposable as the product of k sections of the form $x_{\mathbb{V}}$, then

$$\left(\mathbb{d}^{\gamma,0}\theta\right)\left(x_{\mathfrak{u}_{0}},\ldots,x_{\mathfrak{u}_{k}}\right)=\sum_{i< j}\left(-1\right)^{i+j}\theta\left(x_{\left[\mathfrak{u}_{i},\mathfrak{u}_{j}\right]},x_{\mathfrak{u}_{0}},\ldots,\widehat{x}_{\mathfrak{u}_{i}},\ldots,\widehat{x}_{\mathfrak{u}_{j}},\ldots,x_{\mathfrak{u}_{k}}\right)$$

Let $f_{\mathfrak{u}} \in \mathfrak{g}^* \oplus \mathfrak{g}^*$ be defined by

$$f_{\mathsf{u}}(\mathsf{v}) := x_{\mathsf{u}}\left(x_{\mathsf{v}}\right)$$

It induces an inclusion of the Chevalley-Eilenberg complex of $\mathfrak{g} \oplus \mathfrak{g}$ for the trivial module

$$(\wedge^{*}\left(\mathfrak{g}^{*}\oplus\mathfrak{g}^{*}\right),\delta)\hookrightarrow\left(\Omega_{\mathbb{T}}^{*}\left(G\right),\mathbb{d}^{\gamma,0}\right)\,\colon\,f_{\mathbb{U}}\mapsto x_{\mathbb{U}}$$

Similar to the classical case, this induces an isomorphism on the cohomology:

$$\widetilde{H}^*_{\gamma,0}(G) \cong H^*(\mathfrak{g} \oplus \mathfrak{g}) \cong H^*(G \times G)$$

The isomorphism above in fact is an isomorphism of rings, where on $\widetilde{H}^*_{\gamma,0}(G)$ the product is induced by the wedge product in $\Omega^*_{\pi}(M)$.

3. CURVATURE TENSORS

Consider a generalized connection \mathcal{D} on V. Let $\mathcal{D}^{\mathbb{T}}$ be any generalized connection on $\mathbb{T}M$. The $\mathcal{D}^{\mathbb{T}}$ -derivation $\mathbb{d}^{\mathbb{T}}$ (2.2) extends to $\Omega^k_{\mathbb{T}}(V) := C^{\infty} (\wedge^k \mathbb{T}M \otimes V)$:

(3.1)
$$\mathbf{d}_{D}^{\mathbb{T}}(\theta \otimes v) := \left(\mathbf{d}^{\mathbb{T}}\theta\right) \otimes v + (-1)^{k}\theta \wedge Dv$$

where $v \in C^{\infty}(V)$. The $\mathcal{D}^{\mathbb{T}}$ -curvature operator $\mathcal{F}^{\mathbb{T}}(\mathcal{D})$ of \mathcal{D} is then given by

(3.2)
$$\mathcal{F}^{\mathbb{T}}(\mathcal{D}) := \mathbb{d}_{\mathcal{D}}^{\mathbb{T}} \circ \mathcal{L}$$

which generally is not tensorial in v if $\mathcal{D}^{\mathbb{T}}$ is not TM-torsion free. When \mathcal{D} is understood, it will be dropped from the notations $\mathcal{F}^{\mathbb{T}}(\mathcal{D})$ and $\mathbb{d}_{\mathcal{D}}^{\mathbb{T}}$.

In terms of covariant derivatives, the $\mathcal{D}^{\mathbb{T}}$ -curvature operator $\mathcal{F}^{\mathbb{T}}$ is given by (1.5). It is tensorial iff $\mathcal{D}^{\mathbb{T}}$ is TM-torsion free, in which case, for any $f \in C^{\infty}(M)$

(3.3)
$$\mathcal{F}_{x,y}^{\mathbb{T}}(fv) - f\mathcal{F}_{x,y}^{\mathbb{T}}v = \left([\pi(x), \pi(y)] - \pi \left(x \diamond_{\mathbb{T}} y \right) \right)(f)v = 0$$

The resulting tensor $\mathcal{F}^{\mathbb{T}} \in \Omega^2_{\mathbb{T}}$ (End (*V*)) is the $\mathcal{D}^{\mathbb{T}}$ -curvature of \mathcal{D} . Similar to (3.1), let $\widetilde{\mathbb{d}}_{\mathcal{D}}^{\mathbb{T}}$ be the extension of $\widetilde{\mathbb{d}}^{\mathbb{T}}$ to $\widetilde{\Omega}^*_{\mathbb{T}}$ (*V*) := $\widetilde{\Omega}^*_{\mathbb{T}}$ (*M*) $\otimes C^{\infty}$ (*V*). By (2.15), $\mathcal{F}^{\mathbb{T}}$ (*D*) can be seen as an element in $\widetilde{\Omega}^2_{\mathbb{T}}$ (End (*V*)) and (3.2) can also be rewritten as

(3.4)
$$\mathcal{F}^{\mathbb{T}}(\mathcal{D}) = \widetilde{d}_{\mathcal{D}}^{\mathbb{T}} \circ \mathcal{D}$$

Example 3.1. Let (V, h) be a Hermitian vector bundle on M. A generalized connection D on V preserves h, or is (h-)unitary, if for $v_i \in C^{\infty}(V)$ and $x \in \mathbb{T}M$ with $X = \pi(x)$:

$$Xh(v_1, v_2) = h\left(\mathcal{D}_x v_1, v_2\right) + h\left(v_1, \mathcal{D}_x v_2\right)$$

Suppose now that V is a Hermitian line bundle and s is a local section of V such that h(s, s) = 1. Since D is unitary, it is determined by a local section $u \in C^{\infty}(\mathbb{T}M)$, such that for $x \in \mathbb{T}M$

$$\sqrt{-1}\mathcal{D}_x s = 2\left\langle x, u \right\rangle s$$

Analogous to the classical computation, the $\mathcal{D}^{\mathbb{T}}$ -curvature for the line bundle V is then

(3.5)
$$\sqrt{-1}\mathcal{F}_{x,y}^{\mathbb{T}}(\mathcal{D}) = 2\left(\left\langle y, \mathcal{D}_{x}^{\mathbb{T}}u\right\rangle - \left\langle x, \mathcal{D}_{y}^{\mathbb{T}}u\right\rangle\right) = \left(\mathbb{d}^{\mathbb{T}}u\right)(x, y)$$

3.1. **Chern-Weil homomorphism.** Let $\mathcal{D}^{\mathbb{T}}$ be a TM-torsion free generalized connection on $\mathbb{T}M$. Since $\mathbb{d}^{\mathbb{T}}$ generally does not square to 0 (2.5), the Bianchi identity generally does not hold for $\mathcal{F}^{\mathbb{T}}(\mathcal{D})$. In terms of covariant derivatives, $\mathbb{d}_{\mathcal{D}}^{\mathbb{T}}\mathcal{F}^{\mathbb{T}}(\mathcal{D})$ can be expanded into

(3.6)
$$\begin{pmatrix} \left(\mathbb{d}_{D}^{\mathbb{T}} \mathcal{F}^{\mathbb{T}} \right)_{x,y,z} = \mathcal{D}_{x} \mathcal{F}_{y,z}^{\mathbb{T}} - \mathcal{F}_{D_{x}^{\mathbb{T}}y,z}^{\mathbb{T}} - \mathcal{F}_{y,D_{x}^{\mathbb{T}}z}^{\mathbb{T}} - \mathcal{F}_{y,z}^{\mathbb{T}} \mathcal{D}_{x} + c.p. \text{ in } x, y, z \\ = -\mathcal{D}_{\mathcal{R}_{x,y}^{\mathbb{T}}z}^{\mathbb{T}} - c.p. \text{ in } x, y, z$$

where $x, y, z \in C^{\infty}(\mathbb{T}M)$. By (2.8) and (2.9), it gives

(3.7)
$$\left(\mathbb{d}_{D}^{\mathbb{T}} \mathcal{F}^{\mathbb{T}} \right)_{x,y,z} = \mathcal{D}_{\left[x \diamond_{\mathbb{T}} y \diamond_{\mathbb{T}} z \right]} = \psi_{\left[x \diamond_{\mathbb{T}} y \diamond_{\mathbb{T}} z \right]}$$

which leads to the Bianchi identity over $\widetilde{\Omega}^*_{\mathbb{T}}(M)$.

Lemma 3.2. Let $\mathcal{F}^{\mathbb{T}} \in \Omega^2_{\mathbb{T}}(M) \otimes \operatorname{End}(V)$ be the $\mathcal{D}^{\mathbb{T}}$ -curvature of a generalized connection \mathcal{D} on V, then

(3.8)
$$\widetilde{\mathbf{d}}_D^{\mathbb{T}} \mathcal{F}^{\mathbb{T}} = 0$$

Proof. It follows from (3.7) that

$$\mathbb{d}_{D}^{\mathbb{T}}\mathcal{F}^{\mathbb{T}} \in \operatorname{img} \mathscr{J}^{\mathbb{T}} \otimes \operatorname{End} (V)$$
$$\mathcal{F}^{\mathbb{T}} = 0.$$

Thus $\widetilde{\mathbf{d}}_D^{\mathbb{T}} \mathcal{F}^{\mathbb{T}} = \mathscr{R}^{\mathbb{T}} \left(\mathbf{d}_D^{\mathbb{T}} \mathcal{F}^{\mathbb{T}} \right) = 0.$

The space $\mathscr{D}(V)$ of generalized connections on V is an affine space modeled on $\Omega^1_{\mathbb{T}}$ (End (V)), which coincides with $\widetilde{\Omega}^1_{\mathbb{T}}$ (End (V)). For $A \in \widetilde{\Omega}^1_{\mathbb{T}}$ (End (V)), standard computation gives

$$\mathcal{F}^{\mathbb{T}}\left(\mathcal{D}+A\right)-\mathcal{F}^{\mathbb{T}}\left(\mathcal{D}\right)=\mathbb{d}_{\mathcal{D}}^{\mathbb{T}}A+A\wedge A$$

It follows that

(3.9)
$$\operatorname{tr}_{V}\left(\mathcal{F}^{\mathbb{T}}\left(\mathcal{D}+A\right)\right) - \operatorname{tr}_{V}\left(\mathcal{F}^{\mathbb{T}}\left(\mathcal{D}\right)\right) = \operatorname{tr}_{V}\left(\widetilde{\operatorname{d}}_{D}^{\mathbb{T}}A\right) = \widetilde{\operatorname{d}}^{\mathbb{T}}\operatorname{tr}_{V}\left(A\right)$$

As in the classical case, (3.8) implies that

$$\widetilde{d}^{\mathbb{T}} \operatorname{tr}_{V} \mathcal{F}^{\mathbb{T}}(\mathcal{D}) = \operatorname{tr}_{V} \widetilde{d}_{\mathcal{D}}^{\mathbb{T}} \mathcal{F}^{\mathbb{T}}(\mathcal{D}) = 0$$

The gauge group Aut (V) acts on $\mathcal{D}(V)$ by push-forward

$$(\lambda D)_x v := \lambda^{-1} [D_x (\lambda v)]$$

where $\lambda \in Aut(V)$, $x \in TM$ and $v \in C^{\infty}(V)$. It induces the action on the curvature by conjugation

$$\mathcal{F}^{\mathbb{T}}(\lambda \mathcal{D}) = \lambda^{-1} \mathcal{F}^{\mathbb{T}}(\mathcal{D}) \lambda$$

The Chern-Weil homomorphism extends to define characteristic classes for V in $\widetilde{H}^*_{\pi}(M)$.

Definition 3.3. For a Hermitian vector bundle (V, h) over M, its k-th $\mathcal{D}^{\mathbb{T}}$ -Chern class is

(3.10)
$$c_{k}^{\mathbb{T}}(V) := \left[\operatorname{tr}_{V} \left(\sqrt{-1} \mathcal{F}^{\mathbb{T}}(\mathcal{D}) \right)^{k} \right] \in \widetilde{H}_{\mathbb{T}}^{2k}(M)$$

where \mathcal{D} is a generalized connection on V. For real vector bundles, their $\mathcal{D}^{\mathbb{T}}$ -Euler and $\mathcal{D}^{\mathbb{T}}$ -*Pontrjagin* classes can similarly be defined, as elements of $\widetilde{H}^*_{\mathbb{T}}(M)$ of appropriate degrees.

By (3.9), the $\mathcal{D}^{\mathbb{T}}$ -Chern classes do not depend on the choice of \mathcal{D} on V. Let \mathcal{D} be the lift of a classical connection ∇_0 on V and let F_0 be the classical curvature of ∇_0 , then

$$\mathcal{F}_{x,y}^{\mathbb{T}} = F_{0;\pi(x),\pi(y)}$$

for $x, y \in C^{\infty}(\mathbb{T}M)$. This relates $c_*^{\mathbb{T}}(V)$ to the classical Chern classes $c_*(V)$.

Proposition 3.4. Let (V, h) be a Hermitian vector bundle, then

$$c_k^{\mathbb{I}}(V) = \widetilde{\pi}^* c_k(V)$$

for all k. In particular, $c_k^{\mathbb{T}}(V) = 0$ for all k such that 2k > n. Similarly, the $\mathcal{D}^{\mathbb{T}}$ -Euler and Pontrjagin classes are the images of the respective classical classes under $\tilde{\pi}^*$.

3.2. (ϕ, b) -curvatures. A generalized Riemannian metric \mathbb{G}^b induces an eigendecomposition of a generalized connection D on V. For $X \in C^{\infty}(TM)$ and $v \in C^{\infty}(V)$, let $x^{b\pm} = X + (b \pm g) X$, then:

(3.11)
$$\nabla^b_{\pm,X} v := \mathcal{D}_{x^{b\pm}} v$$

The connections ∇^b_+ depend on *b*, while their difference does not (c.f. [15]):

(3.12)
$$\psi_X := \frac{1}{2} \left(\nabla^b_{+,X} - \nabla^b_{-,X} \right) = \frac{1}{2} \left(\mathcal{D}_{x^{b+}} - \mathcal{D}_{x^{b-}} \right) = \mathcal{D}_{g(X)}$$

The average of ∇^b_+ gives the *b*-neutral connection of \mathcal{D} :

(3.13)
$$\nabla_{0,X}^{b} := \frac{1}{2} \left(\nabla_{+,X}^{b} + \nabla_{-,X}^{b} \right) = \mathcal{D}_{X+i_{X}b}$$

which leads to

(3.14)
$$\nabla^b_{\pm} = \nabla^b_0 \pm \psi$$

When $\psi = 0$, the generalized connection \mathcal{D} is the lift of a classical connection ∇_0 on V, in which case $\nabla^b_{\pm} = \nabla_0$ are independent of b as well. The dependence on b of the \mathbb{G}^b -eigendecomposition of \mathcal{D} can be described in terms of ψ as follows.

Proposition 3.5. For $b \in \Omega^2(M)$, let

$$j_b := g^{-1}b : TM \to TM : X \mapsto g^{-1}(\iota_X b)$$

and (∇^b_+, ∇^b_-) be the \mathbb{G}^b -eigendecomposition of \mathcal{D} on V. For $a \in \Omega^2(M)$, then

$$\nabla^{b+a}_{\pm} = \nabla^b_{\pm} + \psi_j$$

gives the \mathbb{G}^{b+a} -eigendecomposition of \mathcal{D} .

 \square

Definition 3.6. Let \mathcal{D} be a generalized connection on a vector bundle V over M. The (ϕ, b) -curvature $\mathcal{F}^{\phi, b}(\mathcal{D})$ of \mathcal{D} is its $\mathcal{D}^{\phi, b}$ -curvature (3.2). When \mathcal{D} is understood it is also simply denoted $\mathcal{F}^{\phi, b}$.

Given the pair of classical connections (∇^b_+, ∇^b_-) , besides the curvature F^b_{\pm} of each of them, there is also a *mixed curvature* $F^b_{\pm-}$ ([31]):

(3.15)
$$F_{+,-;X,Y}^{\phi,b} v := \left(\nabla_{+,X}^{b} \nabla_{-,Y}^{b} - \nabla_{-,\nabla_{X}^{-\phi}Y}^{b}\right) v - \left(\nabla_{-,Y}^{b} \nabla_{+,X}^{b} - \nabla_{+,\nabla_{Y}^{+\phi}X}^{b}\right) v$$

where $X, Y \in C^{\infty}(TM)$ and $v \in C^{\infty}(V)$. It can also be expressed using the tensor ψ

(3.16)
$$F_{+,-;X,Y}^{\phi,b} = F_{+,X,Y}^{b} - 2\left(\nabla_{+,X}^{b}\psi\right)_{Y} - \left(\iota_{g^{-1}\psi}\phi\right)(X,Y)$$

Let F_0^b be the classical curvature for ∇_0^b (3.13), it gives another decomposition for the mixed curvature (3.15):

(3.17)
$$F_{+,-;X,Y}^{\phi,b} = F_{0;X,Y}^{b} + \left(\iota_{g^{-1}\psi}\phi\right)(X,Y) - \left[\psi_{X},\psi_{Y}\right] - \left[\left(\nabla_{0,X}^{b}\psi\right)_{Y} + \left(\nabla_{0,Y}^{b}\psi\right)_{X}\right]$$

Theorem 3.7. The (ϕ, b) -curvature admits \mathbb{G}^{b} -eigendecomposition in terms of the (mixed) curvatures of the pair $(\nabla^{b}_{+}, \nabla^{b}_{-})$ of classical connections as follows:

(3.18)
$$\mathcal{F}_{x^{b\pm},y^{b\pm}}^{\phi,b} v = F_{\pm,X,Y}^{b} v \text{ and } \mathcal{F}_{x^{b+},y^{b-}}^{\phi,b} v = F_{+,-;X,Y}^{\phi,b} v$$

Proof. Straightforward from the definition, via the \mathbb{G}^{b} -eigendecomposition.

Example 3.8. Continue with Example 3.1 for $\mathcal{D}^{\mathbb{T}} = \mathcal{D}^{\phi,b}$. In this case, the local section $u \in C^{\infty}(\mathbb{T}M)$ decomposes into

$$u = \frac{1}{2} \left[\left(g^{-1} v_{+} + b g^{-1} v_{+} + v_{+} \right) - \left(g^{-1} v_{-} + b g^{-1} v_{-} - v_{-} \right) \right]$$

where $\sqrt{-1}v_{\pm} \in \Omega^1(M)$ are the local 1-forms defining the connections ∇^b_{\pm} respectively. It follows that

$$\sqrt{-1}F^b_{\pm} = dv_{\pm}$$

and the mixed component in $\mathcal{F}^{\phi,b}$ is given by

$$\sqrt{-1}F^{b}_{+,-;X,Y} = \left[\nabla^{-\phi}_{X}\nu_{-}\right](Y) - \left[\nabla^{+\phi}_{Y}\nu_{+}\right](X)$$

Note that $F_{+,-}^b$ is neither symmetric nor skew-symmetric in X and Y, and decomposes into symmetric and skew-symmetric parts as

$$\sqrt{-1}F_{+,-}^{b} = -\mathcal{L}_{g^{-1}\psi}g + \left(\sqrt{-1}F_{0}^{b} - \iota_{g^{-1}\psi}\phi\right)$$

where F_0^b is the curvature of the *b*-neutral connection ∇_0^b and $\psi = \frac{1}{2} (v_+ - v_-)$, in the decomposition (3.14) of \mathcal{D} .

Corollary 3.9. In the decomposition $\mathbb{T}M = e^bTM \oplus T^*M$ the (ϕ, b) -curvature is

(3.19)
$$\mathcal{F}_{x,y}^{\phi,b} = F_{0;X,Y}^{b} + \left(\nabla_{0,X}^{b}\psi\right)_{g^{-1}\eta} - \left(\nabla_{0,Y}^{b}\psi\right)_{g^{-1}\xi} + \left[\psi_{g^{-1}\xi},\psi_{g^{-1}\eta}\right] \\ + \frac{1}{2}\left[\left(\iota_{g^{-1}\psi}\phi\right)(X,Y) - \left(\iota_{g^{-1}\psi}\phi\right)\left(g^{-1}\xi,g^{-1}\eta\right)\right]$$

where $x = X + \iota_X b + \xi$ and $y = Y + \iota_Y b + \eta$.

Remark 3.10. The dependence of $\mathcal{F}^{\phi,b}$ on ϕ is completely contained in the last term of (3.16), e.g. for $\phi \in \Omega^3(M)$:

$$F_{+,-;X,Y}^{\phi+\phi,b} - F_{+,-;X,Y}^{\phi,b} = -\left(\iota_{g^{-1}\phi}\phi\right)(X,Y)$$

The dependence of $\mathcal{F}^{\phi,b}$ on *b* is more complicated, which can be derived from (3.19) by relatively lengthy computations, noting that ξ and η in the expression depend on *b* as well.

3.3. **Yang-Mills functional.** It is straightforward to extend the Yang-Mills functional to this context. The generalized metric \mathbb{G}^b induces natural inner product on $\wedge^* \mathbb{T}M$. For a Hermitian bundle (V, h), it induces a natural norm on $\wedge^* \mathbb{T}M \otimes \text{End}(V)$, denoted $\|\bullet\|_{\mathbb{G}^b,h}$. The $\mathcal{D}^{\mathbb{T}}$ -*Yang-Mills functional* on $\mathcal{D}(V)$ is given by

(3.20)
$$\mathcal{YM}_{\mathbb{T}}(\mathcal{D}) := \int_{M} \left\| \mathcal{F}^{\mathbb{T}}(\mathcal{D}) \right\|_{\mathbb{G}^{b},h}^{2} d\operatorname{vol}_{g}$$

It is evidently invariant under the gauge action on $\mathcal{D}(V)$. When restricted to the subspace of the lifts of classical connections on V, $\mathcal{YM}_{\mathbb{T}}(D)$ reduces to (a constant multiple of) the classical Yang-Mills functional. It can also be regarded as a functional of the pair $(\mathcal{D}, \mathcal{D}^{\mathbb{T}})$ of generalized connections on V and $\mathbb{T}M$ respectively.

When $\mathcal{D}^{\mathbb{T}} = \mathcal{D}^{\phi,b}$, it can be represented as

(3.21)
$$\mathcal{YM}_{\phi,b}(\mathcal{D}) = YM\left(\nabla^{b}_{+}\right) + YM\left(\nabla^{b}_{-}\right) + 2\int_{M} \left\|F^{\phi,b}_{+,-}\right\|_{g,h}^{2} d\operatorname{vol}_{g}$$

where $YM(\bullet)$ denotes the classical Yang-Mills functional. The 2-form *b* affects only the \mathbb{G}^{b} -eigendecomposition of \mathcal{D} . The right hand side can be seen as a functional for a pair of classical connections (∇_{+}, ∇_{-}) , where the last term encodes the dependence on $\phi \in \Omega^{3}(M)$ (Remark 3.10), as well as the interaction within the pair.

4. CURVATURES ON $\mathbb{T}M$

For a \mathbb{G}^{b} -metric connection $\mathcal{D}^{\mathbb{T}}$ on $\mathbb{T}M$, its (ϕ, b) -curvature is denoted $\mathcal{R}^{\mathbb{T},\phi,b}$ and the associated *curvature tensor* is given by

(4.1)
$$\mathcal{R}^{\mathbb{T},\phi,b}(x,y,z,w) := \mathbb{G}^b\left(\mathcal{R}^{\mathbb{T},\phi,b}_{x,y}z,w\right)$$

where $x, y, z, w \in C^{\infty}(\mathbb{T}M)$. Similar to the classical situation, it is skew in the first two and the last two entries respectively:

$$\mathcal{R}^{\mathbb{T},\phi,b}(x,y,z,w) = -\mathcal{R}^{\mathbb{T},\phi,b}(y,x,z,w) = \mathcal{R}^{\mathbb{T},\phi,b}(y,x,w,z)$$

Definition 4.1. The (ϕ, b) -*Riemannian curvature* $\mathcal{R}^{\phi,b}$ for (M, g) is the (ϕ, b) -curvature for $\mathcal{D}^{\phi,b}$, and the corresponding curvature tensor is the (ϕ, b) -*Riemann tensor*, which is also denoted by $\mathcal{R}^{\phi,b}$.

4.1. **Bianchi identities.** Since $\mathcal{D}^{\phi,b}$ preserves C^b_{\pm} , $\mathcal{R}^{\phi,b}(x, y, z, w)$ vanishes when the last two entries are sections of different \mathbb{G}^b -eigenbundles. The non-vanishing components in the \mathbb{G}^b -eigendecomposition of $\mathcal{R}^{\phi,b}$ are given by the classical Riemann tensor R of g as well as the curvature tensors $R^{\pm\phi}$ of $\nabla^{\pm\phi}$.

Proposition 4.2. Let $X, Y, Z, W \in TM$ and $x^{b\pm} \in C^b_{\pm}$ with $X = \pi(x^{b\pm})$ etc, then

 $\begin{array}{l} (1) \ \mathcal{R}^{\phi,b}\left(x^{b\pm}, y^{b\pm}, z^{b\pm}, w^{b\pm}\right) = R(X, Y, Z, W) \\ (2) \ \mathcal{R}^{\phi,b}\left(x^{b\mp}, y^{b\pm}, z^{b\pm}, w^{b\pm}\right) = R^{\pm\phi}(X, Y, Z, W) \mp \frac{1}{2}\left(\nabla_{X}^{\pm\phi}\phi\right)(Y, Z, W) \\ (3) \ \mathcal{R}^{\phi,b}\left(x^{b\mp}, y^{b\mp}, z^{b\pm}, w^{b\pm}\right) = R^{\pm\phi}(X, Y, Z, W) \end{array}$

All other components of $\mathcal{R}^{\phi,b}$ vanish.

Proof. Standard computations from the definitions, which is left for the reader.

By (1) above, the algebraic Bianchi identity for $\mathcal{R}^{\phi,b}$ holds when all entries involved are from the same \mathbb{G}^{b} -eigenbundle. The analogues to the algebraic and differential Bianchi identities follow from previous discussions.

Lemma 4.3. In their respective \mathbb{G}^{b} -eigendecompositions:

(1) For
$$\mathcal{R}_{x,y}^{\phi,b}z + c.p.$$
:
(4.2) $\mathcal{R}_{x^{b\mp},y^{b\mp}}^{\phi,b}z^{b\pm} + \mathcal{R}_{y^{b\mp},z^{b\pm}}^{\phi,b}x^{b\mp} + \mathcal{R}_{z^{b\pm},x^{b\mp}}^{\phi,b}y^{b\mp} = \pm 2g\left(\mathcal{R}_{X,Y}^{\pm\phi}Z\right)$

(2) For $\mathbb{d}_{D}^{\phi,b}\mathcal{F}^{\phi,b}$:

(4.3)
$$\left(\mathbb{d}_{D}^{\phi,b} \mathcal{F}^{\phi,b} \right)_{x^{b\mp}, y^{b\mp}, z^{b\pm}} = \mp 2\mathcal{D}_{g\left(R_{X,Y}^{\pm\phi}Z\right)} = \mp 2\psi_{R_{X,Y}^{\pm\phi}Z}$$

All other components vanish.

Proof. The identity (4.2) follows from (2.8) and (2.34). If $d\phi = 0$, (4.2) can also be obtained from the explicit expressions in Proposition 4.2 and the identity [4] below

)

(4.4)
$$R^{+\phi}(X,Y,Z,W) = R^{-\phi}(Z,W,X,Y)$$

Then (4.3) follows from (3.7) and (4.2).

In particular, the differential Bianchi identity holds for the (ϕ, b) -curvature when $\psi = 0$, i.e. \mathcal{D} is the lifting of a classical connection on V. Another special case is when $\nabla^{\pm \phi}$ are flat ([1, 6, 7]), i.e. $R^{\pm \phi} = 0$, then both Bianchi identities hold. In this second special case, $\mathcal{R}^{\phi,b}$ enjoys all the symmetries of a classical Riemann curvature.

Theorem 4.4. Suppose that the connections $\nabla^{\pm \phi}$ are flat on T M. Then

(1) $\mathcal{R}^{\phi,b}(x, y, z, w) = -\mathcal{R}^{\phi,b}(y, x, z, w) = \mathcal{R}^{\phi,b}(y, x, w, z)$ (2) $\mathcal{R}^{\phi,b}_{x,y}z + \mathcal{R}^{\phi,b}_{y,z}x + \mathcal{R}^{\phi,b}_{z,x}y = 0$ (3) $\mathbb{d}^{\phi,b}_{D}\mathcal{R}^{\phi,b} = 0$

(4) $\mathcal{R}^{\phi,b}(x, y, z, w) = \mathcal{R}^{\phi,b}(z, w, x, y)$

In this case, $\mathcal{R}^{\phi,b}$ defines a symmetric pairing on $\wedge^2 \mathbb{T} M$:

$$\mathcal{R}^{\phi,b}:\wedge^{2}\mathbb{T}M\otimes\wedge^{2}\mathbb{T}M\to\mathbb{R}:\mathcal{R}^{\phi,b}(x\wedge y,w\wedge z):=\mathcal{R}^{\phi,b}(x,y,z,w)$$

which defines the corresponding operator on $\wedge^2 \mathbb{T}M$ via \mathbb{G}^b .

Proof. Items (1) - (3) follow from preceeding discussions and the flatness assumption, while (4) follows from (1) - (3) as in the classical situation. The last statement is a consequence of (1) and (4).

The following consequence of Lemma 4.3 was used in the proof of Proposition 2.14.

Proposition 4.5. $\mathbb{d}^{\phi,b}$ is a differential at $\Omega^{2n-1}_{\pi}(M)$.

Proof. Let $\{X_j\}$ be a local *g*-orthonormal frame of TM and $\{e_j^{b\pm}\}$ the induced \mathbb{G}^b -orthonormal frame of $\mathbb{T}M$. Let $\theta \in \Omega_{\mathbb{T}}^{2n-2}(M)$, then by (2.8), (2.13) and (4.2):

$$\begin{split} \left(\mathbb{d}^{\phi,b} \circ \mathbb{d}^{\phi,b} \Theta \right) \left(e_{1}^{b+}, \dots, e_{n}^{b+}, e_{1}^{b-}, \dots, e_{n}^{b-} \right) \\ &= -\sum_{i < j,k} (-1)^{i+j+k+n} \theta \left(-g \left(R_{X_{i},X_{j}}^{-\phi} X_{k} \right), \dots, \hat{e}_{i}^{b+}, \dots, \hat{e}_{j}^{b+}, \dots, \hat{e}_{k}^{b-}, \dots \right) \\ &- \sum_{i,j < k} (-1)^{i+j+k} \theta \left(g \left(R_{X_{j},X_{k}}^{+\phi} X_{i} \right), \dots, \hat{e}_{i}^{b+}, \dots, \hat{e}_{j}^{b-}, \dots, \hat{e}_{k}^{b-}, \dots \right) \\ &= \sum_{i,j,k} (-1)^{j+k+n} \left[-R^{-\phi} \left(X_{i}, X_{j}, X_{k}, X_{i} \right) + R^{+\phi} \left(X_{i}, X_{k}, X_{j}, X_{i} \right) \right] \theta \left(\dots, \hat{e}_{j}^{b+}, \dots, \hat{e}_{k}^{b-}, \dots \right) \\ &= 0 \end{split}$$

The last step above follows from (4.4).

4.2. **Ricci curvature.** The trace of the (ϕ, b) -curvature on $\mathbb{T}M$ defines the corresponding (ϕ, b) -Ricci curvature.

Definition 4.6. For a \mathbb{G}^b -metric connection $\mathcal{D}^{\mathbb{T}}$, the (ϕ, b) -*Ricci curvature* $\mathcal{R}ic^{\mathbb{T}} : \mathbb{T}M \to \mathbb{T}M$ for (M, \mathbb{G}^b) is the trace of the (ϕ, b) -curvature $\mathcal{R}^{\mathbb{T}}$. For $x, y \in C^{\infty}(\mathbb{T}M)$, in the local orthonormal frame $\left\{e_i^{b+}, e_j^{b-}\right\}$ of $\mathbb{T}M$ induced from a local *g*-orthonormal frame $\{X_i\}$

(4.5)
$$\mathcal{R}ic^{\mathbb{T}}(x) := \sum_{i} \left[\mathcal{R}_{x,e_{i}^{b+}}^{\mathbb{T}} e_{i}^{b+} + \mathcal{R}_{x,e_{i}^{b-}}^{\mathbb{T}} e_{i}^{b-} \right]$$

The (ϕ, b) -Ricci tensor $\mathcal{R}c^{\mathbb{T}} \in C^{\infty} (\mathbb{T}M \otimes \mathbb{T}M)$ is

(4.6)
$$\mathcal{R}c^{\mathbb{T}}(x,y) := \mathbb{G}^{b}\left(\mathcal{R}ic^{\mathbb{T}}(x),y\right)$$

Specialized to the (ϕ, b) -connection $\mathcal{D}^{\phi,b}$, the \mathbb{G}^{b} -eigendecomposition of $\mathcal{R}c^{\phi,b}$ can be determined from that of $\mathcal{R}^{\phi,b}$ as follows:

(4.7)
$$\mathcal{R}c^{\phi,b}\left(x^{b\pm}, y^{b\pm}\right) = \mathcal{R}c\left(X, Y\right) \text{ and } \mathcal{R}c^{\phi,b}\left(x^{b\pm}, y^{b\mp}\right) = \mathcal{R}c^{\mp\phi}\left(X, Y\right)$$

where Rc is the Ricci tensor for ∇ , $Rc^{\pm\phi}$ are the Ricci tensors for $\nabla^{\pm\phi}$ respectively:

(4.8)
$$Rc^{\pm\phi} = Rc \mp \frac{1}{2}d^*\phi - \frac{1}{4}\phi^2$$

and

$$\phi^2(X,Y) := \sum_{i,j} \phi(X,X_i,X_j)\phi(Y,X_i,X_j)$$

It follows that $\mathcal{R}c^{\phi,b}$ is symmetric, in other words

(4.9)
$$\langle \mathcal{R}ic^{\phi,b}(x), \mathbb{G}^{b}y \rangle = \langle \mathbb{G}^{b}\mathcal{R}ic^{\phi,b}(x), y \rangle = \langle \mathbb{G}^{b}\mathcal{R}ic^{\phi,b}(y), x \rangle = \langle \mathbb{G}^{b}x, \mathcal{R}ic^{\phi,b}(y) \rangle$$

Similar to the classical case, the (ϕ, b) -Ricci curvature appears in a Weitzenböck identity relating two natural Laplacians on $\wedge^* \mathbb{T}M$ described in §2.4.

Theorem 4.7. On $\Omega^*_{\pi}(M)$, the following Weitzenböck identity holds:

(4.10)
$$\Delta^{\phi,b} = \Delta_{D^{\phi,b}} + \mathbb{G}^b \mathcal{R} i c^{\phi,b} \mathbb{G}^b$$

where $\Delta^{\phi,b}$ is the $\mathcal{D}^{\phi,b}$ -Hodge Laplacian while $\Delta_{\mathcal{D}^{\phi,b}}$ is the $\mathcal{D}^{\phi,b}$ -Bochner Laplacian.

Proof. It can be shown following standard computation that

$$\Delta^{\phi,b}\theta = \Delta_{D^{\phi,b}}\theta - \frac{1}{2}\sum_{\alpha,\beta} \mathbb{G}^{b}\left(e^{b}_{\alpha}\right) \wedge \iota_{e^{b}_{\beta}}\left(\mathcal{R}^{\phi,b}_{e^{b}_{\alpha},e^{b}_{\beta}}\theta\right)$$

where $e_{\alpha}^{b}, e_{\beta}^{b}$ run through $\{e_{i}^{b+}, e_{j}^{b-}\}$. Set $\theta = \mathbb{G}^{b}(x)$ for $x \in \mathbb{T}M$, then

$$\sum_{\alpha,\beta} \left\langle \mathbb{G}^{b}\left(e_{\alpha}^{b}\right) \wedge \iota_{e_{\beta}^{b}}\left(\mathcal{R}_{e_{\alpha}^{b},e_{\beta}^{b}}^{\phi,b}\mathbb{G}^{b}\left(x\right)\right), y \right\rangle = 2 \sum_{\alpha,\beta} \left\langle \mathbb{G}^{b}\left(\mathcal{R}_{e_{\alpha}^{b},e_{\beta}^{b}}^{\phi,b}x\right), e_{\beta}^{b} \right\rangle \mathbb{G}^{b}\left(e_{\alpha}^{b},y\right)$$
$$= 2 \sum_{\beta} \mathcal{R}^{\phi,b}\left(y, e_{\beta}^{b}, x, e_{\beta}^{b}\right) = -2\mathcal{R}c^{\phi,b}\left(y, x\right) = -2 \left\langle \mathbb{G}^{b}\mathcal{R}ic^{\phi,b}\left(x\right), y \right\rangle$$

from which (4.10) follows.

The symmetry of $\mathcal{R}c^{\phi,b}$ implies that $\mathbb{d}^{\phi,b*}$ is a differential at $\Omega^1_{\mathbb{T}}(M)$. For $x_i \in C^{\infty}(\mathbb{T}M)$, i = 1, ..., k, set $y_i = \mathbb{G}^b(x_i)$, then straightforward computation gives

$$\begin{split} & \left(\mathbb{d}^{\phi,b*} \circ \mathbb{d}^{\phi,b*}\right) \left[y_1 \wedge \ldots \wedge y_k\right] \\ &= \sum_{i < j} \left(-1\right)^{i+j} \left[\mathcal{R}c^{\phi,b}\left(x_j, x_i\right) - \mathcal{R}c^{\phi,b}\left(x_i, x_j\right)\right] y_1 \wedge \ldots \wedge \hat{y_i} \wedge \ldots \wedge \hat{y_j} \wedge \ldots \wedge y_k \\ &- \sum_{i < j < \ell} \left(-1\right)^{i+j+\ell} \mathbb{G}^b \left(\mathcal{R}^{\phi,b}_{x_i, x_j} x_\ell + c.p.\right) y_1 \wedge \ldots \wedge \hat{y_i} \wedge \ldots \wedge \hat{y_j} \wedge \ldots \wedge \hat{y_\ell} \wedge \ldots \wedge y_k \end{split}$$

Due to the symmetry of $\mathcal{R}c^{\phi,b}$, the terms in the second line vanish. Furthermore, when k = 2, the terms in the last line vanish as well.

Proposition 4.8. The operator $\mathbb{d}^{\phi,b*}$ is a differential at $\Omega^1_{\mathbb{T}}(M)$, i.e.

(4.12)
$$\theta \in \Omega^2_{\mathbb{T}}(M) \Longrightarrow \left(\mathbb{d}^{\phi, b*} \circ \mathbb{d}^{\phi, b*} \right) \theta = 0$$

If the connections $\nabla^{\pm\phi}$ are flat on TM, then $\left(\Omega^*_{\mathbb{T}}(M), \mathbb{d}^{\phi,b}\right)$ and $\left(\Omega^*_{\mathbb{T}}(M), \mathbb{d}^{\phi,b*}\right)$ are both chain complices.

Proof. For the last statement, that $(\Omega^*_{\mathbb{T}}(M), \mathbb{d}^{\phi,b})$ is a chain complex follows from (2.5), (2.8) and the algebraic Bianchi identity, which is item (2) in Theorem 4.4. The statement for $(\Omega^*_{\mathbb{T}}(M), \mathbb{d}^{\phi,b*})$ follows from (4.11) and the algebraic Bianchi identity.

4.3. **Bismut connection.** The generalized Bismut connection $\mathcal{D}^{\phi,B}$ introduced by Gualtieri [15] is a \mathbb{G}^{b} -metric connection and is the lift of a classical connection $\nabla^{\phi,B}$ on $\mathbb{T}M$, which depends on both ϕ and b:

(4.13)
$$\nabla_X^{\phi,B} y^{b+} := \nabla_X^{+\phi} Y + (b+g) \left(\nabla_X^{+\phi} Y \right)$$
 and $\nabla_X^{\phi,B} y^{b-} := \nabla_X^{-\phi} Y + (b-g) \left(\nabla_X^{-\phi} Y \right)$

Note that $\mathcal{D}^{\phi,\mathcal{B}}$ is compatible with the (almost) Dorfman bracket $*_{\phi-db}$ (Definition 2.7).

Since $\mathcal{D}^{\phi,B}$ is a lift of a classical connection, by (4.3), the (ϕ, b) -Bismut curvature $\mathcal{R}^{\phi,B}$ of $\mathcal{D}^{\phi,B}$ satisfies the differential Bianchi identity:

$$\mathbb{d}_{D}^{\phi,b}\mathcal{R}^{\phi,B}=0$$

More explicitly, $\mathcal{R}^{\phi,B}$ is determined by the classical curvature of $\nabla^{\phi,B}$, which in turn is given by $R^{\pm\phi}$:

(4.14)
$$\mathcal{R}^{\phi,\mathcal{B}}\left(x,y,z^{b\pm},w^{b\pm}\right) = R^{\pm\phi}\left(X,Y,Z,W\right)$$

The \mathbb{G}^{b} -eigendecomposition of the corresponding (ϕ, b) -Bismut Ricci tensor $\mathcal{R}c^{\phi, B}$ is thus

(4.15)
$$\mathcal{R}c^{\phi,B}\left(x,y^{b+}\right) = \mathcal{R}c^{+\phi}\left(X,Y\right) \text{ and } \mathcal{R}c^{\phi,B}\left(x,y^{b-}\right) = \mathcal{R}c^{-\phi}\left(X,Y\right)$$

4.4. **Scalar curvatures.** The traces of the (ϕ, b) -Ricci curvatures give the corresponding (ϕ, b) -scalar curvatures, which depend on the \mathbb{G}^b -metric connection $\mathcal{D}^{\mathbb{T}}$. For instance, the (ϕ, b) -Riemann scalar curvature $S^{\phi, b}$ is the trace of $\mathcal{R}ic^{\phi, b}$:

(4.16)
$$S^{\phi,b} = \sum_{j} \left[\mathcal{R}c^{\phi,b} \left(e_{i}^{b+}, e_{i}^{b+} \right) + \mathcal{R}c^{\phi,b} \left(e_{i}^{b-}, e_{i}^{b-} \right) \right] = 2S$$

where *S* is the classical scalar curvature of *g*. On the other hand, the (ϕ, b) -*Bismut scalar curvature S*^{ϕ,B} is the trace of $\mathcal{R}ic^{\phi,B}$:

$$S^{\phi,B} = \sum_{j} \left[\mathcal{R}c^{\phi,B} \left(e_{i}^{b+}, e_{i}^{b+} \right) + \mathcal{R}c^{\phi,B} \left(e_{i}^{b-}, e_{i}^{b-} \right) \right] = 2S - 3 \|\phi\|_{g}^{2}$$

where $\|\phi\|_g$ is the norm of ϕ with respect to g:

$$\|\phi\|_{g}^{2} = \sum_{i < j < k} \phi(X_{i}, X_{j}, X_{k})^{2}$$

5. GENERALIZED COMPLEX MANIFOLDS

Let \mathbb{J} be a generalized almost complex structure on M. It induces a polarization of $\mathbb{T}_{\mathbb{C}}M := \mathbb{T}M \otimes_{\mathbb{R}} \mathbb{C}$ as the direct sum of its $\pm \sqrt{-1}$ -eigenbundles:

(5.1)
$$\mathbb{T}_{\mathbb{C}}M = \mathbb{T}_{\mathbb{J}}^{1,0}M \oplus \mathbb{T}_{\mathbb{J}}^{0,1}M$$

Here $\mathbb{T}^{1,0}_{\mathbb{J}}M$ denotes the $\sqrt{-1}$ -eigenbundle of \mathbb{J} , and $\mathbb{T}^{0,1}_{\mathbb{J}}M$ its complex conjugate. They are maximally isotropic and are dual to each other under the pairing $2\langle,\rangle$ on $\mathbb{T}_{\mathbb{C}}M$. For instance, the space of (0, 1)-forms with respect to \mathbb{J} is identified with the sections of $\mathbb{T}^{1,0}_{\mathbb{J}}M$:

$$\Omega^{0,1}_{\mathbb{J}}(M) := C^{\infty} \left(\mathbb{T}^{1,0}_{\mathbb{J}} M \right)$$

Similar to the classical case, the (0, k)-forms with respect to \mathbb{J} are sections of $\wedge^k \mathbb{T}^{1,0}_{\mathbb{J}} M$:

$$\Omega^{0,k}_{\mathbb{J}}(M) := C^{\infty} \left(\wedge^{k} \mathbb{T}^{1,0}_{\mathbb{J}} M \right)$$

In general, the type decomposition of $\Omega^*_{\mathbb{T}}(M)$ with respect to \mathbb{J} is given by

(5.2)
$$\Omega^*_{\mathbb{T}}(M) = \bigoplus_{p,q} \Omega^{p,q}_{\mathbb{J}}(M) := \bigoplus_{p,q} C^{\infty} \left(\wedge^p \mathbb{T}^{0,1}_{\mathbb{J}} M \otimes \wedge^q \mathbb{T}^{1,0}_{\mathbb{J}} M \right)$$

For notational convenience, sometimes $\mathbb{T}^{1,0}_{\mathbb{J}}M$ is denoted *L*, while $\mathbb{T}^{0,1}_{\mathbb{J}}M$ is denoted *L*.

Definition 5.1. Let (M, \mathbb{J}) be a generalized almost complex manifold and let $\gamma \in \Omega^3(M)$ be a closed 3-form. Then a generalized connection $\mathcal{D}^{\mathbb{T}}$ on $\mathbb{T}M$ is \mathbb{J} -compatible with $*_{\gamma}$ if $\diamond_{\mathbb{T}}$ coincides with $*_{\gamma}$ on the sections from the same eigenbundle of \mathbb{J} , i.e.

(5.3)
$$x \diamond_{\mathbb{T}} y = x *_{\gamma} y \text{ and } \overline{x} \diamond_{\mathbb{T}} \overline{y} = \overline{x} *_{\gamma} \overline{y}$$

where $x, y \in C^{\infty}(\mathbb{T}^{1,0}_{\mathbb{J}}M)$. Such generalized connection $\mathcal{D}^{\mathbb{T}}$ is a γ - \mathbb{J} -connection if it furthermore is TM-torsion free.

It is straightforward to see that (5.3) is equivalent to the following, where $x, y \in C^{\infty}(\mathbb{T}M)$:

$$(\mathbb{J}x) \diamond_{\mathbb{T}} y + x \diamond_{\mathbb{T}} (\mathbb{J}y) = (\mathbb{J}x) *_{\gamma} y + x *_{\gamma} (\mathbb{J}y)$$

Thus, if non-empty, the space $\mathcal{D}_{\mathbb{J},\gamma}(\mathbb{T}M)$ of generalized connections that are \mathbb{J} -compatible with $*_{\gamma}$ is modeled on

$$\mathscr{D}_{\mathbb{J},\gamma}(\mathbb{T}M) \cong \left\{ A : \mathbb{T}M \otimes \mathbb{T}M \to \mathbb{T}M : A_{\mathbb{J}x}y - A_{y}(\mathbb{J}x) = A_{\mathbb{J}y}x - A_{x}(\mathbb{J}y) \right\}$$

It's then evident by (2.10) that the subspace $\mathscr{D}_{\mathbb{J},\gamma,\tau}(\mathbb{T}M)$ of the γ - \mathbb{J} -connections, if nonempty, is modeled on

(5.4)
$$\mathscr{D}_{\mathbb{J},\gamma,\tau}(\mathbb{T}M) \cong \left\{ A : \mathbb{T}M \odot \mathbb{T}M \to T^*M : A_{\mathbb{J}x}y - A_y(\mathbb{J}x) = A_{\mathbb{J}y}x - A_x(\mathbb{J}y) \right\}$$

Let $\gamma \in \Omega^3(M)$ be a closed 3-form, then \mathbb{J} is *integrable (with respect to* γ) if $\mathbb{T}^{1,0}_{\mathbb{J}}M$ is involutive under the Dorfman bracket $*_{\gamma}$, i.e.

$$x, y \in C^{\infty} \left(\mathbb{T}^{1,0}_{\mathbb{J}} M \right) \Longrightarrow x *_{\gamma} y \in C^{\infty} \left(\mathbb{T}^{1,0}_{\mathbb{J}} M \right)$$

In this case, $(M, \gamma; \mathbb{J})$ is a generalized complex manifold.

Lemma 5.2. Let $(M, \gamma; \mathbb{J})$ be a generalized complex manifold with a generalized connection $\mathcal{D}^{\mathbb{T}}$ on $\mathbb{T}M$ that is \mathbb{J} -compatible with $*_{\gamma}$. Then the operator $\mathbb{d}^{\mathbb{T}}$ decomposes into components as follows:

(5.5)
$$\mathbb{d}^{\mathbb{T}} = \partial_{\mathbb{J}}^{\mathbb{T}} + \overline{\partial}_{\mathbb{J}}^{\mathbb{T}} : \Omega_{\mathbb{J}}^{p,q}(M) \to \Omega_{\mathbb{J}}^{p+1,q}(M) \oplus \Omega_{\mathbb{J}}^{p,q+1}(M)$$

Proof. The proof is similar to that of Lemma 2.5, employing (5.3) and the integrability of \mathbb{J} . The details are left for the reader.

The integrability of \mathbb{J} implies that both of its eigenbundles are complex Lie algebroids, with their Lie brackets given by the restriction of $*_{\gamma}$. The corresponding Lie algebroid de Rham differential for $\overline{L} = \mathbb{T}_{\mathbb{J}}^{0,1} M$ will be denoted by $d_{\overline{L}}$:

(5.6)
$$d_{\overline{L}}: \Omega^{0,k}_{\mathbb{J}}(M) \to \Omega^{0,k+1}_{\mathbb{J}}(M)$$

If $\mathcal{D}^{\mathbb{T}}$ is \mathbb{J} -compatible with $*_{\gamma}$, then $d_{\overline{L}}$ coincides with the restriction of $\mathbb{d}^{\mathbb{T}}$ on $\Omega^{0,*}_{\mathbb{J}}(M)$. More precisely, for $\theta \in \Omega^{0,q}_{\mathbb{J}}(M)$, direct computation shows that

(5.7)
$$\left(d_{\overline{L}} \theta \right) \left(\overline{x}_0, \overline{x}_1, \dots, \overline{x}_q \right) = \sum_j (-1)^j \left(\mathcal{D}_{\overline{x}_j}^{\mathbb{T}} \theta \right) \left(\overline{x}_0, \dots, \widehat{\overline{x}_j}, \dots, \overline{x}_q \right)$$

where $x_j \in C^{\infty} \left(\mathbb{T}^{1,0}_{\mathbb{J}} M \right)$ for all *j*, which gives

(5.8)
$$\overline{\partial}_{\mathbb{J}}^{\mathbb{T}}\theta = d_{\overline{L}}\theta \in \Omega_{\mathbb{J}}^{0,q+1}(M)$$

The other component $\partial_{\mathbb{J}}^{\mathbb{T}} \theta \in \Omega_{\mathbb{J}}^{1,q}(M)$ is given by

(5.9)
$$\left(\partial_{\mathbb{J}}^{\mathbb{T}}\theta\right)\left(x_{0},\overline{x}_{1},\ldots,\overline{x}_{q}\right) = \left(\mathcal{D}_{x_{0}}^{\mathbb{T}}\theta\right)\left(\overline{x}_{1},\ldots,\overline{x}_{q}\right) + \sum_{j}\theta\left(\overline{x}_{1},\ldots,\mathcal{D}_{\overline{x}_{j}}^{\mathbb{T}}x_{0},\ldots,\overline{x}_{q}\right)$$

where $x_j \in C^{\infty}(\mathbb{T}^{1,0}_{\mathbb{J}}M)$. Via complex conjucation, the analogous versions of the identities (5.7), (5.8) and (5.9) are valid for $\overline{\theta} \in \Omega^{q,0}_{\mathbb{J}}(M)$ with $L = \mathbb{T}^{1,0}_{\mathbb{J}}M$ in place of \overline{L} .

Example 5.3. Suppose that $\mathcal{D}^{\mathbb{T}}$ is \mathbb{J} -compatible with $*_{\gamma}$ and consider $f \in C^{\infty}(M)$. It follows from Proposition 2.2 and Lemma 5.2 that

(5.10)
$$\partial_{\mathbb{J}}^{\mathbb{T}}\partial_{\mathbb{J}}^{\mathbb{T}}f = 0, \overline{\partial}_{\mathbb{J}}^{\mathbb{T}}\overline{\partial}_{\mathbb{J}}^{\mathbb{T}}f = 0, \text{ and } \partial_{\mathbb{J}}^{\mathbb{T}}\overline{\partial}_{\mathbb{J}}^{\mathbb{T}}f + \overline{\partial}_{\mathbb{J}}^{\mathbb{T}}\partial_{\mathbb{J}}^{\mathbb{T}}f = 0$$

The first two identities in the above can also be seen as the consequences of (5.8) and the corresponding version for d_L .

5.1. Generalized (almost) Hermitian manifolds. Let $(M; \mathbb{G}^b, \mathbb{J})$ be a *generalized almost Hermitian manifold*, i.e. \mathbb{J} and $\mathbb{J}_{-} := \mathbb{G}^b \mathbb{J}$ are commuting generalized almost complex structures. The eigenbundles of \mathbb{J} (and \mathbb{J}_{-}) decompose into the common eigenbundles of \mathbb{G}^b and \mathbb{J} . Let

(5.11)
$$\ell_{\pm} := \mathbb{T}^{1,0}_{\mathbb{J}} M \cap \left(C^b_{\pm} \otimes \mathbb{C} \right)$$

then for instance

$$\mathbb{T}^{1,0}_{\mathbb{J}}M = \mathscr{\ell}_{+} \oplus \mathscr{\ell}_{-} \text{ and } \mathbb{T}^{1,0}_{\mathbb{J}_{-}}M = \mathscr{\ell}_{+} \oplus \overline{\mathscr{\ell}}_{-}$$

28

The restriction of \mathbb{J} to C^b_{\pm} induces two almost complex structures I_{\pm} on TM:

(5.12)
$$I_{\pm}X + (b \pm g) \left(I_{\pm}X \right) := \mathbb{J}_{+} \left[X + (b \pm g)X \right]$$

It follows that $(M; \mathbb{G}^b, \mathbb{J})$ is equivalent to a pair of almost Hermitian structures (g, I_{\pm}) together with $b \in \Omega^2(M)$.

Lemma 5.4. For a generalized almost Hermitian manifold $(M; \mathbb{G}^b, \mathbb{J})$, the space of \mathbb{G}^b -metric γ - \mathbb{J} -connections, if non-empty, is modeled on the following subspace of $\Omega^3(M)$:

$$\left\{\phi \in \Omega^{3}(M) : \phi\left(I_{X}, Y, Z\right) + \phi\left(X, I_{+}Y, Z\right) = 0, \text{ for all } X, Y, Z \in C^{\infty}(TM)\right\}$$

Proof. It follows from (5.4) and Theorem 2.6.

Let $\phi = \gamma + db$, then by Theorem 2.8, $\mathcal{D}^{\phi,b}$ is metric compatible with $*_{\gamma}$. Hence, the \mathbb{J} -compatibility of $\mathcal{D}^{\phi,b}$ with $*_{\gamma}$ is equivalent to $\diamond_{\phi,b}$ and $*_{\gamma}$ coincide on sections of the same common eigenbundle of \mathbb{J} and \mathbb{G}^{b} , e.g.

(5.13)
$$x^{b\pm} *_{\gamma} y^{b\pm} = x^{b\pm} \diamond_{\phi,b} y^{b\pm}$$

where $x^{b\pm}, y^{b\pm} \in C^{\infty}(\ell_{\pm})$ with $\pi(x^{b\pm}) = X$ and so on.

Lemma 5.5. On a generalized almost Hermitian manifold $(M; \mathbb{G}^b, \mathbb{J})$, let $\gamma \in \Omega^3(M)$ be a closed 3-form and $\phi = \gamma + db$. Then $\mathcal{D}^{\phi,b}$ is \mathbb{J} -compatible with $*_{\gamma}$ iff $\nabla^{\pm \phi} I_{\pm} = 0$.

Proof. For any $X, Y, Z \in C^{\infty}(TM)$, let $x^{b\pm} \in C^{\infty}(C^{b}_{\pm})$ such that $\pi(x^{b\pm}) = X$ and so on, then

$$\left\langle x^{b\pm} *_{\gamma} y^{b\pm} - x^{b\pm} \diamond_{\phi,b} y^{b\pm}, z \right\rangle = \pm g \left(\nabla_{Z}^{\pm \phi} X, Y \right)$$

Thus (5.13) is equivalent to

$$g\left(\nabla_{Z}^{\pm\phi}X_{\pm},Y_{\pm}\right)=0$$

for all $Z \in C^{\infty}(TM)$ and $X_{\pm}, Y_{\pm} \in C^{\infty}(T_{\pm;1,0}M)$ respectively. Hence $\nabla^{\pm \phi}$ preserves $T_{\pm;1,0}M$ respectively, from which the statement follows.

Remark 5.6. The condition $\nabla^{\pm \phi} I_{\pm} = 0$ can be rewritten as

(5.14)
$$\left(\nabla_X I_{\pm} \right) Y = \pm \frac{1}{2} \left(I_{\pm} g^{-1} \iota_Y \iota_X \phi - g^{-1} \iota_{I_{\pm} Y} \iota_X \phi \right)$$

for $X, Y \in C^{\infty}(TM)$, which implies that I_{\pm} are of class $\mathcal{W}_1 \oplus \mathcal{W}_3 \oplus \mathcal{W}_4$ in the Gray-Hervella classification of almost Hermitian structures [10]. The Nijenhuis tensors of I_{\pm}

$$N_{I_{\pm}}(X,Y) := [X,Y] + I_{\pm}[I_{\pm}X,Y] + I_{\pm}[X,I_{\pm}Y] - [I_{\pm}X,I_{\pm}Y]$$

can be expressed in terms of ϕ :

$$g\left(N_{I_{\pm}}(X,Y),Z\right) = \phi\left(I_{\pm}X,I_{\pm}Y,Z\right) + \phi\left(I_{\pm}X,Y,I_{\pm}Z\right) + \phi\left(X,I_{\pm}Y,I_{\pm}Z\right) - \phi\left(X,Y,Z\right)$$

which implies that the integrability of I_{\pm} is equivalent to ϕ being of type (1, 2) + (2, 1) with respect to I_{\pm} respectively. Furthermore, by Friedrich-Ivanov [8], if almost Hermitian connections for (g, I_{\pm}) that admit completely skew torsions exist, they must be unique.

In $(M; \mathbb{G}^b, \mathbb{J})$, when \mathbb{J} is integrable with respect to $\gamma \in \Omega^3(M)$, the structure $(M, \gamma; \mathbb{G}^b, \mathbb{J})$ defines a *generalized Hermitian manifold*. In this case, the \mathbb{J} -compatibility of $\mathcal{D}^{\phi,b}$ with $*_{\gamma}$ is equivalent to the integrability of \mathbb{J}_- , i.e. it provides a *generalized Kähler condition*.

Theorem 5.7. On a generalized Hermitian manifold $(M, \gamma; \mathbb{G}^b, \mathbb{J})$, let $\phi = \gamma + db$. Then $(M, \gamma; \mathbb{G}^b, \mathbb{J})$ is a generalized Kähler manifold iff $\mathcal{D}^{\phi, b}$ is \mathbb{J} -compatible with $*_{\gamma}$.

Proof. Starting with the J-compatibility, then (5.13) and the integrability of J imply that ℓ_{\pm} and $\overline{\ell}_{\pm}$ are involutive with respect to $*_{\gamma}$, which further implies that I_{\pm} are integrable. Let $x^{b+} \in C^{\infty}(\ell_{+}), y^{b-} \in C^{\infty}(\ell_{-}), X_{+} = \pi(x^{b+})$ and $Y_{-} = \pi(y^{b-})$, then by (2.28) and Lemma 5.5

$$x^{b+} *_{\gamma} \overline{y^{b-}} = \left[\nabla_{X_{+}}^{-\phi} \overline{Y}_{-} + (b-g) \nabla_{X_{+}}^{-\phi} \overline{Y}_{-} \right] - \left[\nabla_{\overline{Y}_{-}}^{+\phi} X_{+} + (b+g) \nabla_{\overline{Y}_{-}}^{+\phi} X_{+} \right] \in \mathcal{E}_{+} \oplus \overline{\mathcal{E}}_{-}$$

Thus \mathbb{J}_{-} is integrable. The opposite direction is left to the reader.

Remark 5.8. The generalized Kähler condition in Theorem 5.7 relates to the condition given in [15] as follows. The condition $\nabla^{\pm \phi} I_{\pm} = 0$ on a generalized almost Hermitian manifold is equivalent to $\mathcal{D}^{\phi,B} \mathbb{J} = 0$, which implies the equivalence of the integrability of I_{\pm} to the type condition for the *generalized torsion* as defined in [15] – the integrability of \mathbb{J} then follows. In Theorem 5.7, the integrability of I_{\pm} follows from the integrability of \mathbb{J} and (5.13), obtaining the type of ϕ with respect to I_{\pm} as a consequence.

Corollary 5.9. On a generalized Hermitian manifold $(M, \gamma; \mathbb{G}^b, \mathbb{J})$, let $\phi = \gamma + db$. Then $(M, \gamma; \mathbb{G}^b, \mathbb{J})$ is a generalized Kähler manifold iff $\mathcal{D}^{\phi, B}\mathbb{J} = 0$.

Proof. As stated in Remark 5.8, $\mathcal{D}^{\phi,B} \mathbb{J} = 0$ is equivalent to $\nabla^{\pm \phi} I_{\pm} = 0$, which by Lemma 5.5 is equivalent to $\mathcal{D}^{\phi,b}$ being \mathbb{J} -compatible.

Let $\mathcal{D}^{\mathbb{T}}$ be any \mathbb{G}^{b} -metric connection on a generalized Hermitian manifold $(M, \gamma; \mathbb{G}^{b}, \mathbb{J})$. Its \mathbb{J} -*Ricci form* $\rho_{\mathbb{T}}^{\mathbb{T}} := \rho_{\mathbb{J}} (\mathcal{D}^{\mathbb{T}}) \in \Omega_{\mathbb{T}}^{2}(M)$ is defined as

$$\rho_{\mathbb{J}}^{\mathbb{T}}(x,y) := \sum_{i} \left[\mathcal{R}^{\mathbb{T}}\left(x,y,\mathbb{J}e_{i}^{b+},e_{i}^{b+}\right) + \mathcal{R}^{\mathbb{T}}\left(x,y,\mathbb{J}e_{i}^{b-},e_{i}^{b-}\right) \right]$$

The \mathbb{J} -scalar curvature for $\mathcal{D}^{\mathbb{T}}$ is

$$\mathcal{S}_{\mathbb{J}}^{\mathbb{T}} \mathrel{\mathop:}= \sum_{i} \left[\rho_{\mathbb{J}}^{\mathbb{T}} \left(\mathbb{J} e_{i}^{b+}, e_{i}^{b+} \right) + \rho_{\mathbb{J}}^{\mathbb{T}} \left(\mathbb{J} e_{i}^{b-}, e_{i}^{b-} \right) \right]$$

5.2. Generalized Kähler manifolds. Recall that the structure $(M, \gamma; \mathbb{G}^b, \mathbb{J})$ defines a *generalized Kähler manifold* if both \mathbb{J} and $\mathbb{J}_{-} = \mathbb{G}^b \mathbb{J}$ are integrable generalized almost complex structures with respect to γ ([17]). Let $\phi = \gamma + db$, Theorem 5.7 indicates that $\mathcal{D}^{\phi,b}$ is \mathbb{J} -compatible with $*_{\gamma}$. In particular, for $\theta \in \Omega^{p,0}_{\mathbb{T}}(M)$, (5.8) gives

(5.15)
$$\partial_{\mathbb{J}}^{\phi,b}\theta = d_L\theta \text{ and } \overline{\partial}_{\mathbb{J}}^{\phi,b}\overline{\theta} = d_{\overline{L}}\overline{\theta}$$

Example 5.10 (c.f. Example 5.3). Set $\mathcal{D}^{\mathbb{T}} = \mathcal{D}^{\phi,b}$ in (5.10), then the components in the third identity can further be rewritten in terms of the \mathbb{G}^{b} -eigendecomposition. For instance

(5.16)
$$\left(\partial_{\mathbb{J}}^{\phi,b}\overline{\partial}_{\mathbb{J}}^{\phi,b}f\right)\left(x_{\pm},\overline{y}_{\pm}\right) = \left(\partial_{\pm}\overline{\partial}_{\pm}f\right)\left(X_{\pm},\overline{Y}_{\pm}\right)$$

where ∂_{\pm} and $\overline{\partial}_{\pm}$ are the operators associated to the classical complex structures I_{\pm} , while $X_{\pm}, Y_{\pm} \in C^{\infty}(T_{\pm;1,0}M)$ are sections of the I_{\pm} -holomorphic tangent bundles respectively, and $x_{\pm} = X_{\pm} + (b \pm g) X_{\pm}$ and so on.

Since $d_{\overline{L}}^2 = 0$, as a consequence of (5.15), the algebraic Bianchi identity (4.2) for $\mathcal{R}^{\phi,b}$ implies that $\nabla^{+\phi}$ (resp. $\nabla^{-\phi}$) induces a natural I_{-} -holomorphic (resp. I_{+} -holomorphic) structure on the eigenbundles of I_{+} (resp. of I_{-}), providing an alternative proof of this well-known result in [15].

Proposition 5.11. Let $(M, \gamma; \mathbb{G}^b, \mathbb{J})$ be a generalized Kähler manifold. Let $\phi = \gamma + db$, $x, y, z, w \in C^{\infty}(\mathbb{T}^{1,0}_{\mathbb{J}}M)$, and \overline{x} , etc. are their complex conjugates. Then

(5.17)
$$\left\langle \mathcal{R}_{\overline{x},\overline{y}}^{\phi,b}\overline{z},w\right\rangle + c.p.\ in\ x,\ y,\ z = 0\ and\ \left\langle \mathcal{R}_{x,y}^{\phi,b}z,\overline{w}\right\rangle + c.p.\ in\ x,\ y,\ z = 0$$

which give rise to

$$R_{\overline{X}_+,\overline{Y}_+}^{-\phi}\overline{Z}_- = 0, R_{\overline{X}_+,\overline{Y}_+}^{-\phi}W_- = 0, R_{\overline{X}_-,\overline{Y}_-}^{+\phi}\overline{Z}_+ = 0 \text{ and } R_{\overline{X}_-,\overline{Y}_-}^{+\phi}W_+ = 0$$

as well as their complex conjugates, where $X_{\pm}, Y_{\pm}, Z_{\pm}, W_{\pm} \in C^{\infty} \left(T_{\pm;1,0} M \right)$.

Proof. To see the first identity in (5.17), note that $w \in \Omega^{0,1}_{\mathbb{J}}(M)$, then

$$\overline{\partial}_{\mathbb{J}}^{\phi,b} \circ \overline{\partial}_{\mathbb{J}}^{\phi,b} w = d_{\overline{L}}^2 w = 0$$

By (2.8) and (2.13), for $x, y, z \in C^{\infty} (\mathbb{T}^{1,0}_{\mathbb{J}} M)$

$$0 = \left(\overline{\partial}_{\mathbb{J}}^{\phi,b} \circ \overline{\partial}_{\mathbb{J}}^{\phi,b} w\right) \left(\overline{x}, \overline{y}, \overline{z}\right) = \left(\mathbb{d}^{\phi,b} \circ \mathbb{d}^{\phi,b} w\right) \left(\overline{x}, \overline{y}, \overline{z}\right)$$
$$= w\left(\left[\overline{x} \diamond_{\phi,b} \overline{y} \diamond_{\phi,b} \overline{z}\right]\right) = -2\left\langle \mathcal{R}_{\overline{x},\overline{y}}^{\phi,b} \overline{z}, w\right\rangle - c.p. \text{ in } x, y, z$$

The second identity then follows by taking complex conjugation.

To see the classical curvature identities, restrict x, y, z, w in (5.17) further to the \mathbb{G}^{b} eigenbundles (5.11). For instance, consider $x, y \in C^{\infty}(\ell_{+})$ and $z, w \in C^{\infty}(\ell_{-})$. Set $X_{+} = \pi(x), Z_{-} = \pi(z)$ and so on, then apply (4.2) to the first identity in (5.17) gives

$$-\left\langle 2g\left(R_{\overline{X}_{+},\overline{Y}_{+}}^{-\phi}\overline{Z}_{-}\right),w_{-}\right\rangle =g\left(R_{\overline{X}_{+},\overline{Y}_{+}}^{-\phi}W_{-},\overline{Z}_{-}\right)=0$$

Since (g, I_{-}) is Hermitian and $\nabla^{-\phi}$ preserves I_{-} , it gives the identities involving $R^{-\phi}$. The rest of the identities are obtained similarly.

For the generalized Bismut connection $\mathcal{D}^{\phi,B}$, with $\phi = \gamma + db$, (4.14) implies that $\rho_{\mathbb{J}}^{\phi,B}$ is computed by the sum of the respective classical Bismut-Ricci forms ρ_+ for (g, I_+) :

$$\rho_{\mathbb{J}}^{\phi,\mathcal{B}}(x,y) = \rho_{+}(X,Y) + \rho_{-}(X,Y)$$

The corresponding J-scalar curvature also decomposes:

$$\mathcal{S}_{\mathbb{J}}^{\phi,B} = \sum_{i} \left[\rho_{\mathbb{J}}^{\phi,B} \left(\mathbb{J}e_{i}^{b+}, e_{i}^{b+} \right) + \rho_{\mathbb{J}}^{\phi,B} \left(\mathbb{J}e_{i}^{b-}, e_{i}^{b-} \right) \right] = S_{+} + 2S_{+-} + S_{-}$$

where S_{\pm} are the respective classical Bismut scalar curvatures for $\left(g, I_{\pm}\right)$ and S_{+-} is the mixed Bismut scalar curvature:

$$S_{+-} := \sum_{i,j} R^{+\phi} \left(I_{-}X_{i}, X_{i}, I_{+}X_{j}, X_{j} \right) = \sum_{i,j} R^{-\phi} \left(I_{+}X_{i}, X_{i}, I_{-}X_{j}, X_{j} \right)$$

6. J-HOLOMORPHIC VECTOR BUNDLES

Let (V, h) be a Hermitian vector bundle on a generalized complex manifold $(M, \gamma; \mathbb{J})$. Recall that a \mathbb{J} -holomorphic structure on V is given by a flat $\mathbb{T}^{0,1}_{\mathbb{J}}M$ -connection:

$$\overline{\partial}_{\mathbb{J}}: C^{\infty}(V) \to \Omega^{0,1}_{\mathbb{J}}(V) := C^{\infty}\left(\mathbb{T}^{1,0}_{\mathbb{J}}M \otimes V\right): \overline{\partial}_{\mathbb{J}}(fv) = d_{\overline{L}}f \otimes v + f\overline{\partial}_{\mathbb{J}}v$$

for $f \in C^{\infty}(M)$ and $v \in C^{\infty}(V)$, such that

(6.1)
$$\overline{\partial}_{\mathbb{J}} \circ \overline{\partial}_{\mathbb{J}} = 0$$

where the extension to $\Omega_{\perp}^{0,k}(V)$ is given by

(6.2)
$$\overline{\partial}_{\mathbb{J}} : \Omega^{0,k}_{\mathbb{J}}(V) \to \Omega^{0,k+1}_{\mathbb{J}}(V) : \overline{\partial}_{\mathbb{J}}(\theta \otimes v) = d_{\overline{L}}\theta \otimes v + (-1)^k \theta \wedge \overline{\partial}_{\mathbb{J}}v$$

for $\theta \in \Omega^{0,k}_{\mathbb{J}}(M)$ and $v \in C^{\infty}(V)$. If $(M, \gamma; \mathbb{G}^{b}, \mathbb{J})$ is generalized Kähler, via the restriction to C^{b}_{\pm} , the \mathbb{J} -holomorphic structure $\overline{\partial}_{\mathbb{J}}$ induces on V an I_{\pm} -holomorphic structure, which will be denoted $\overline{\partial}_{\pm}$ respectively:

(6.3)
$$\overline{\partial}_{\pm,\overline{X}_{\pm}}v := \overline{\partial}_{\mathbb{J},\overline{X}_{\pm}}v$$

where $X_{\pm} \in T_{\pm 1.0}M$, $x_{\pm} = X_{\pm} + (b \pm g) X_{\pm}$ and $v \in C^{\infty}(V)$.

6.1. Connection on $\mathbb{T}^{1,0}_{\mathbb{J}}M$. Consider a generalized Kähler manifold $(M, \gamma; \mathbb{G}^b, \mathbb{J})$, a natural \overline{L} -connections can be defined on $\mathbb{T}^{1,0}_{\mathbb{J}}M$ using the diamond bracket $\diamond_{\mathbb{T}}$ associated to any \mathbb{G}^{b} -metric connection $\mathcal{D}^{\mathbb{T}}$:

(6.4)
$$\overline{\partial}_{\diamond}^{\mathbb{T}}: C^{\infty}\left(\mathbb{T}^{0,1}_{\mathbb{J}}M\right) \otimes C^{\infty}\left(\mathbb{T}^{1,0}_{\mathbb{J}}M\right) \to C^{\infty}\left(\mathbb{T}^{1,0}_{\mathbb{J}}M\right): \overline{\partial}_{\diamond,\overline{x}}^{\mathbb{T}}y := \left[\overline{x} \diamond_{\mathbb{T}}y\right]_{1,0}$$

where $x, y \in C^{\infty}(\mathbb{T}^{1,0}_{\mathbb{J}}M)$ and $[\bullet]_{1,0}$ denotes taking the (1, 0)-component with respect to \mathbb{J} .

It follows from Proposition 5.11 that $\overline{\partial}_{*}^{\phi,b}$ defined by $\mathcal{D}^{\phi,b}$ via (6.4) induces an I_{\pm} -holomorphic structure on $\mathbb{T}^{1,0}_{\mathbb{T}}M$ via its restriction to $\overline{\ell}_{\pm} \cong T_{\pm;0,1}M$. Moreover, by (2.8) and (5.13)

$$\left(\overline{\partial}^{\phi,b}_{\diamond} \circ \overline{\partial}^{\phi,b}_{\diamond}\right)_{\overline{x},\overline{y}} z = \left[\overline{x} \diamond_{\phi,b} \left(\overline{y} \diamond_{\phi,b} z\right) - \overline{y} \diamond_{\phi,b} \left(\overline{x} \diamond_{\phi,b} z\right) - \left(\overline{x} \diamond_{\phi,b} \overline{y}\right) \diamond_{\phi,b} z\right]_{1,0}$$
$$= \left[\mathcal{R}^{\phi,b}_{\overline{x},\overline{y}} z + \mathcal{R}^{\phi,b}_{\overline{y},z} \overline{y} + \mathcal{R}^{\phi,b}_{\overline{z},\overline{y}} \overline{x}\right]_{1,0}$$

Thus, from (4.2), $\overline{\partial}_{\diamond}^{\phi,b}$ defines a \mathbb{J} -holomorphic structure on $\mathbb{T}_{\mathbb{J}}^{1,0}M$ iff $\left[g\left(R_{\overline{X}_{+},Z_{+}}^{-\phi}\overline{Y}_{-}\right)\right]_{1,0} = \left[g\left(R_{\overline{Y}_{-},Z_{-}}^{+\phi}\overline{X}_{+}\right)\right]_{1,0} = 0$

for all $X_{\pm}, Y_{\pm}, Z_{\pm} \in C^{\infty}(T_{\pm;1,0}M)$. Since $\nabla^{\pm \phi}$ preserves I_{\pm} , together with (4.4), the above is equivalent to

$$R^{+\phi}\left(\overline{Y}_{-},\overline{W}_{+},\overline{X}_{+},Z_{+}\right) = R^{-\phi}\left(\overline{X}_{+},\overline{W}_{-},\overline{Y}_{-},Z_{-}\right) = 0$$

for all $X_{\pm}, Y_{\pm}, Z_{\pm}, W_{\pm} \in C^{\infty}(T_{\pm;1,0}M)$. The computations can be summarized as the following result.

Theorem 6.1. Let $(M, \gamma; \mathbb{G}^b, \mathbb{J})$ be a generalized Kähler manifold, and $\phi = \gamma + db$. Let $\overline{\partial}^{\phi, b}_{\diamond}$ be the \overline{L} -connection on $\mathbb{T}^{1,0}_{\mathbb{J}}M$ defined by

$$\overline{\partial}_{\diamond,\overline{x}}^{\phi,b} y := \left[\overline{x} \diamond_{\phi,b} y\right]_{1,0}$$

for $x, y \in C^{\infty} \left(\mathbb{T}^{1,0}_{\mathbb{J}} M \right)$. It is a \mathbb{J} -holomorphic structure on $\mathbb{T}^{1,0}_{\mathbb{J}} M$ iff

$$R^{\pm\phi}_{\overline{X}_{\mp},\overline{Y}_{\pm}}\overline{Z}_{\pm} = 0$$

for all $X_{\pm}, Y_{\pm}, Z_{\pm} \in C^{\infty}(T_{\pm;1,0}M)$. In particular, if $\nabla^{\pm\phi}$ are flat on TM, $\overline{\partial}_{\diamond}^{\phi,b}$ is a \mathbb{J} -holomorphic structure on $\mathbb{T}_{\mathbb{J}}^{1,0}M$.

6.2. Chern curvature. Analogous to the classical situation, on a \mathbb{J} -holomorphic Hermitian bundle (V, h), there exists a unique *generalized Chern connection*:

 $(6.5) D^{h,C} := \overline{\partial}_{\mathbb{J}} + \partial_{\mathbb{J}}$

where $\partial_{\mathbb{J}} : C^{\infty}(V) \to \Omega^{1,0}_{\mathbb{J}}(V) := C^{\infty} \left(\mathbb{T}^{0,1}_{\mathbb{J}} M \otimes V \right)$ is defined by

$$d_{\overline{L}}h(v_1, v_2) = h\left(\overline{\partial}_{\mathbb{J}}v_1, v_2\right) + h\left(v_1, \partial_{\mathbb{J}}v_2\right)$$

for all $v_i \in C^{\infty}(V)$.

Definition 6.2. On a generalized complex manifold $(M, \gamma; \mathbb{J})$, let $\mathcal{D}^{\mathbb{T}}$ be a *TM*-torsion free generalized connection on $\mathbb{T}M$. For a \mathbb{J} -holomorphic Hermitian bundle $(V, h, \overline{\partial}_{\mathbb{J}})$, its $\mathcal{D}^{\mathbb{T}}$ -*Chern curvature* $\mathcal{F}^{\mathbb{T},C}$ is the $\mathcal{D}^{\mathbb{T}}$ -curvature of its generalized Chern connection (6.5).

When $\mathcal{D}^{\mathbb{T}}$ is \mathbb{J} -compatible with $*_{\gamma}$, (5.8) implies that the extension (3.1) of $\mathcal{D}^{\mathbb{T}}$ to $\Omega^{*}_{\mathbb{T}}(V)$ by $\mathcal{D}^{h,C}$ is compatible with the extension (6.2) of $\overline{\partial}_{\mathbb{J}}$ to $\Omega^{0,*}_{\mathbb{T}}(V)$.

Proposition 6.3. Let $\mathcal{D}^{\mathbb{T}}$ be a γ - \mathbb{J} -connection on a generalized complex manifold $(M, \gamma; \mathbb{J})$, the corresponding $\mathcal{D}^{\mathbb{T}}$ -Chern curvature $\mathcal{P}^{\mathbb{T},C}$ is of type (1, 1) with respect to \mathbb{J} , i.e.:

(6.6)
$$\mathcal{F}^{\mathbb{T},C} \in \Omega^{1,1}_{\mathbb{J}} \left(\operatorname{End} \left(V \right) \right) := C^{\infty} \left(\mathbb{T}^{1,0}_{\mathbb{J}} M \wedge \mathbb{T}^{0,1}_{\mathbb{J}} M \otimes \operatorname{End} \left(V \right) \right)$$

In particular, over a generalized Kähler manifold $(M, \gamma; \mathbb{G}^b, \mathbb{J})$, the (ϕ, b) -Chern curvature $\mathcal{F}^{\phi,b,C}$ defined with $\mathcal{D}^{\mathbb{T}} = \mathcal{D}^{\phi,b}$, is of type (1, 1) with respect to \mathbb{J} .

Example 6.4 (c.f. Example 3.1). Consider a J-holomorphic Hermitian line bundle $(V, h, \overline{\partial}_{J})$ and set $\mathcal{D} = \mathcal{D}^{h,C}$, the generalized Chern connection. Choose a unitary local section $s \in C^{\infty}(V)$, there is a local section $u_{J}^{0,1} \in \Omega_{J}^{0,1}(M)$ such that

$$\overline{\partial}_{\mathbb{J}}s = u_{\mathbb{J}}^{0,1} \otimes s$$

Let $\mathcal{D}^{\mathbb{T}}$ be a γ - \mathbb{J} -connection, then $\mathcal{F}^{\mathbb{T},C}(\mathcal{D}) = \sqrt{-1} \mathbb{d}^{\mathbb{T}} u$ where $u \in \Omega^1_{\mathbb{T}}(M)$ and

$$\sqrt{-1}u = u_{\mathbb{J}}^{0,1} - \overline{u_{\mathbb{J}}^{0,1}}$$

By (5.8), the flatness of $\overline{\partial}_{\mathbb{J}}$ implies that

$$\overline{\partial}_{\mathbb{J}}^{\mathbb{T}}u^{0,1} = d_{\overline{L}}u_{\mathbb{J}}^{0,1} = 0$$

It then gives

$$\mathcal{F}^{\mathbb{T},C}\left(\mathcal{D}\right) = \partial_{\mathbb{J}}^{\mathbb{T}} u^{0,1} - \overline{\partial}_{\mathbb{J}}^{\mathbb{T}} \overline{u^{0,1}} \in \Omega_{\mathbb{J}}^{1,1}\left(M\right)$$

Over a generalized Kähler manifold, the (ϕ, b) -Chern connection and curvature are related to the classical Chern connections and curvatures via the \mathbb{G}^{b} -eigendecomposition.

Lemma 6.5. Let $(M, \gamma; \mathbb{G}^b, \mathbb{J})$ be a generalized Kähler manifold and $V \to M$ be a \mathbb{J} holomorphic vector bundle. Let $\nabla^{h,C}_{\pm}$ be the \mathbb{G}^b -eigendecomposition (3.11) of $\mathcal{D}^{h,C}$ (6.5). Then $\nabla^{h,C}_{\pm}$ are the Chern connections for the induced I_{\pm} -holomorphic structures (6.3) on V respectively. Furthermore, the classical Chern curvatures are components of the \mathbb{G}^b eigendecomposition of the (ϕ, b) -Chern curvature.

Proof. The statement about the connections follows from straightforward verification, while the statement about the curvature follows from Theorem 3.7.

Example 6.6. Continue from Example 6.4 and let $(M, \gamma; \mathbb{G}^b, \mathbb{J})$ be generalized Kähler. The I_{\pm} -holomorphic structures induced by $\overline{\partial}_{\mathbb{J}}$ are given locally by $\alpha_{\pm} \in \Omega_{\pm}^{0,1}(M)$:

$$u_{\mathbb{J}}^{0,1} = \left[g^{-1}(\alpha_{+}) + bg^{-1}(\alpha_{+}) + \alpha_{+}\right] - \left[g^{-1}(\alpha_{-}) + bg^{-1}(\alpha_{-}) - \alpha_{-}\right]$$

which satisfies

$$\overline{\partial}_{\pm}\alpha_{\pm} = 0 \text{ and } \left(\nabla_{\overline{X}_{+}}^{-\phi}\alpha_{-}\right)\left(\overline{Y}_{-}\right) - \left(\nabla_{\overline{Y}_{-}}^{+\phi}\alpha_{+}\right)\left(\overline{X}_{+}\right) = 0$$

where $X_{\pm} \in C^{\infty}(T_{\pm 1.0}M)$ and so on. The generalized Chern connection is then

$$\mathcal{D}^{h,C}s = \left(u_{\mathbb{J}}^{0,1} - \overline{u_{\mathbb{J}}^{0,1}}\right) \otimes s$$

while the classical Chern connections $\nabla_{\pm}^{b,C}$ are defined locally by $v_{\pm} = \alpha_{\pm} - \overline{\alpha}_{\pm}$ respectively. Then Example 3.8 gives the \mathbb{G}^{b} -eigendecomposition of the (ϕ, b) -Chern curvature. 6.3. J-Hermitian-Einstein equation. Let $(M, \gamma; \mathbb{G}^b, \mathbb{J})$ be a generalized Hermitian manifold. The analogue to the contraction by the Kähler form is the \mathbb{J}_- -contraction.

Definition 6.7. The $\mathbb{J}_{-contraction} \Lambda_{\mathbb{J}} : \wedge^2 \mathbb{T}_{\mathbb{C}} M \to \mathbb{R}$ is given by:

(6.7)
$$\Lambda_{\mathbb{J}_{-}}(x \wedge y) := \langle \mathbb{J}_{-}x, y \rangle = \mathbb{G}^{b}(\mathbb{J}x, y)$$

where $x, y \in C^{\infty}(\mathbb{T}_{\mathbb{C}}M)$.

In terms of the \mathbb{G}^{b} -eigendecomposition, it corresponds to

(6.8)
$$\Lambda_{\mathbb{J}_{-}}\left(x^{b\pm} \wedge y^{b\pm}\right) = \omega_{\pm}\left(X,Y\right) \text{ and } \Lambda_{\mathbb{J}_{-}}\left(x^{b\pm} \wedge y^{b\mp}\right) = 0$$

where $X = \pi (x^{b\pm})$, $Y = \pi (y^{b\pm})$ and $\omega_{\pm} = gI_{\pm}$. A version of the Hermitian-Einstein equation can thus be formulated in this context.

Definition 6.8. Let $(M, \gamma; \mathbb{G}^b, \mathbb{J})$ be a generalized Hermitian manifold and $\mathcal{D}^{\mathbb{T}}$ be a γ - \mathbb{J} connection on $\mathbb{T}M$. A Hermitian metric h on a J-holomorphic vector bundle $V \to M$ is $\mathcal{D}^{\mathbb{T}}$ - \mathbb{J} -Hermitian-Einstein if the corresponding $\mathcal{D}^{\mathbb{T}}$ -Chern curvature satisfies the following $\mathcal{D}^{\mathbb{T}}$ - \mathbb{J} -Hermitian-Einstein equation:

(6.9)
$$\sqrt{-1}\Lambda_{\mathbb{J}_{-}}\left(\mathcal{F}^{\mathbb{T},C}(V)\right) = 2c\mathrm{Id}_{V}$$

for some $c \in \mathbb{R}$. If $(M, \gamma; \mathbb{G}^b, \mathbb{J})$ is generalized Kähler and $\mathcal{D}^{\mathbb{T}} = \mathcal{D}^{\phi, b}$, the equation (6.9) will be simply called the J-Hermitian-Einstein equation, a solution of which is called a J-Hermitian-Einstein metric.

When $(M, \gamma; \mathbb{G}^b, \mathbb{J})$ is generalized Kähler and $\mathcal{D}^{\mathbb{T}} = \mathcal{D}^{\phi, b}$, by (3.18) and (6.8) the \mathbb{G}^b eigendecomposition of the left hand side of (6.9) is given by

(6.10)
$$\Lambda_{\mathbb{J}_{-}}\left(\mathcal{F}^{\phi,b,C}(V)\right) = \Lambda_{+}\left(F^{C}_{+}(V)\right) + \Lambda_{-}\left(F^{C}_{-}(V)\right)$$

where F_{\pm}^{C} are the curvatures of the classical Chern connections ∇_{\pm}^{h} on V. It follows that (6.9) is equivalent to an equation first proposed by Hitchin in [20].

Proposition 6.9. Over a generalized Kähler manifold $(M, \gamma; \mathbb{G}^b, \mathbb{J})$, the equation (6.9) is equivalent to

(6.11)
$$\frac{\sqrt{-1}}{2} \left(F_{+}^{C}(V) \wedge \omega_{+}^{m-1} + (-1)^{\varepsilon} F_{-}^{C}(V) \wedge \omega_{-}^{m-1} \right) = c (m-1)! \mathrm{Id}_{V} d \mathrm{vol}_{g}$$

where $\varepsilon = 0$ if I_{\pm} induce the same orientation on TM and $\varepsilon = 1$ otherwise.

Proof. Suppose that $d \operatorname{vol}_g = \frac{1}{m!} \omega_+^m$, then

$$mF_{+}^{C}(V) \wedge \omega_{+}^{m-1} = \Lambda_{+}\left(F_{+}^{C}(V)\right)\omega_{+}^{m} = m!\Lambda_{+}\left(F_{+}^{C}(V)\right)d\operatorname{vol}_{g}$$

 \square

Note that $\omega_{+}^{m} = (-1)^{\varepsilon} \omega_{-}^{m}$, it follows from (6.10) that (6.11) is equivalent to (6.9).

The J_-contraction naturally provides the definition of a degree.

Definition 6.10. On a generalized Hermitian manifold $(M, \gamma; \mathbb{G}^b, \mathbb{J})$ with a γ - \mathbb{J} -connection $\mathcal{D}^{\mathbb{T}}$, let $(V, h, \overline{\partial}_{\mathbb{J}})$ be a \mathbb{J} -holomorphic Hermitian vector bundle. The $\mathcal{D}^{\mathbb{T}}$ - \mathbb{G}^b -degree of V is given by:

(6.12)
$$\deg_{\mathbb{G}^{b}}^{\mathbb{T}}(V,h) := \frac{\sqrt{-1}}{4\pi} \int_{M} \operatorname{tr}_{V}\left[\Lambda_{\mathbb{J}_{-}}\left(\mathcal{F}^{\mathbb{T},C}(V)\right)\right] d\operatorname{vol}_{g}$$

When $(M, \gamma; \mathbb{G}^b, \mathbb{J})$ is generalized Kähler and $\mathcal{D}^{\mathbb{T}} = \mathcal{D}^{\phi, b}$, the $\mathcal{D}^{\phi, b}$ - \mathbb{G}^b -degree is simply called the \mathbb{G}^b -degree and denoted deg_{\mathbb{G}^b} (V, h).

Recall that the classical degrees of (V, h) with the induced I_+ -holomorphic structure are

(6.13)
$$\deg_{\pm}(V,h) := \frac{\sqrt{-1}}{2\pi} \int_{M} \operatorname{tr}_{h} \left[\Lambda_{\pm} \left(F_{\pm}^{C}(V) \right) \right] d\operatorname{vol}_{g}$$

Theorem 6.11. Let $(M, \gamma; \mathbb{G}^b, \mathbb{J})$ be a generalized Kähler manifold and $(V, h, \overline{\partial}_{\mathbb{J}})$ be a \mathbb{J} -holomorphic vector bundle on M. Then

(6.14)
$$\deg_{\mathbb{G}^{b}}(V,h) = \frac{1}{2} \left[\deg_{+}(V,h) + \deg_{-}(V,h) \right]$$

where $\deg_{\mathbb{G}^b}$ and \deg_{\pm} are as given in (6.12) and (6.13) respectively.

Proof. It follows from (6.10).

Example 6.12. Continue from Example 6.4 and work now on a generalized Hermitian manifold $(M, \gamma; \mathbb{G}^b, \mathbb{J})$, with a γ - \mathbb{J} -connection $\mathcal{D}^{\mathbb{T}}$. For $f \in C^{\infty}(M)$, let $h_1 = e^{2f}h$ be another Hermitian metric on the line bundle V. An h_1 -unitary local section is then given by $s_1 = e^{-f}s \in C^{\infty}(V)$, which leads to

$$\overline{\partial}_{\mathbb{J}}s_1 = \left(u_{\mathbb{J}}^{0,1} - d_{\overline{L}}f\right) \otimes s_1$$

Let \mathcal{D}_1 be the corresponding generalized Chern connection, whose $\mathcal{D}^{\mathbb{T}}$ -Chern curvature is given by

$$\mathcal{F}^{\mathbb{T},C}\left(\mathcal{D}_{1}\right) = \mathbb{d}^{\mathbb{T}}\left(\sqrt{-1}u + d_{L}f - d_{\overline{L}}f\right) = \mathcal{F}^{\mathbb{T},C}\left(\mathcal{D}\right) - 2\partial_{\mathbb{J}}^{\mathbb{T}}d_{\overline{L}}f$$

where the last step is due to (5.10). Hence

(6.15)
$$\deg_{\mathbb{G}^{b}}^{\mathbb{T}}\left(V,h_{1}\right) - \deg_{\mathbb{G}^{b}}^{\mathbb{T}}\left(V,h\right) = -\frac{\sqrt{-1}}{2\pi}\int_{M}\Lambda_{\mathbb{J}_{-}}\left(\partial_{\mathbb{J}}^{\mathbb{T}}d_{\overline{L}}f\right)d\operatorname{vol}_{\mathcal{S}^{b}}$$

It follows that $\deg_{\mathbb{G}^b}^{\mathbb{T}}$ is independent of the Hermitian metric on *V* iff the right hand side of (6.15) vanishes for all $f \in C^{\infty}(M)$.

The integrand in (6.15) gives rise to the second order operator for $f \in C^{\infty}(M)$:

$$P_{\mathbb{J}}(f) := -\sqrt{-1}\Lambda_{\mathbb{J}_{-}}\left(\partial_{\mathbb{J}}^{\mathbb{T}}d_{\overline{L}}f\right)$$

Similar to the classical case (e.g. [25]), P_{\downarrow} is elliptic since its symbol is given by

$$\sigma\left(P_{\mathbb{J}}\right) = 4\sqrt{-1}\Lambda_{\mathbb{J}_{-}}\left(\xi_{\mathbb{J}}^{0,1} \wedge \xi_{\mathbb{J}}^{1,0}\right) = 4\mathbb{G}\left(\xi_{\mathbb{J}}^{0,1}, \xi_{\mathbb{J}}^{1,0}\right) = \|\xi\|_{g}^{2}$$

where $\xi_{\parallel}^{0,1}$ is the projection of $\xi \in T^*M$ to $\mathbb{T}_{\parallel}^{1,0}M$ and $\xi_{\parallel}^{1,0}$ is the complex conjugate.

Definition 6.13. For a generalized Hermitian manifold $(M, \gamma; \mathbb{G}^b, \mathbb{J})$, the metric \mathbb{G}^b is $\mathcal{D}^{\mathbb{T}}$ - \mathbb{J} -*Gauduchon* if the right hand side of (6.15) vanishes for all $f \in C^{\infty}(M)$. When the structure is generalized Kähler, a $\mathcal{D}^{\phi,b}$ - \mathbb{J} -Gauduchon the metric \mathbb{G}^b is simply said to be \mathbb{J} -*Gauduchon*.

As in the classical situation, if $(M, \gamma; \mathbb{G}^b, \mathbb{J})$ is $\mathcal{D}^{\mathbb{T}}$ - \mathbb{J} -Gauduchon and (V, h) solves (6.9), then the constant *c* is given by

$$c = \frac{2\pi \deg_{\mathbb{G}^b}^{\downarrow}(V)}{\operatorname{rank}(V)\operatorname{Vol}_g(M)}$$

It is then natural to extend the notions of slope and stability to \mathbb{J} -holomorphic vector bundles over a $\mathcal{D}^{\mathbb{T}}$ - \mathbb{J} -Gauduchon generalized Hermitian manifold. The notion of coherent subsheaf in this context can be adopted from Definition 3.4 of [22].

Definition 6.14. Let \mathbb{G}^b be a $\mathcal{D}^{\mathbb{T}}$ - \mathbb{J} -Gauduchon metric. The $\mathcal{D}^{\mathbb{T}}$ - \mathbb{G}^b -slope of a \mathbb{J} -holomorphic vector bundle $\left(V, \overline{\partial}_{\mathbb{J}}\right)$ over M is

(6.16)
$$\mu_{\mathbb{G}^b}^{\mathbb{T}}(V) := \frac{\deg_{\mathbb{G}^b}^{\mathbb{T}}(V)}{\operatorname{rank}(V)}$$

The bundle V is $\mathcal{D}^{\mathbb{T}}$ - \mathbb{G}^{b} -semistable if for any coherent \mathbb{J} -holomorphic subsheaf W of V:

(6.17)
$$\mu_{\mathbb{G}^b}^{\mathbb{T}}(W) \leqslant \mu_{\mathbb{G}^b}^{\mathbb{T}}(V)$$

V is said to be $\mathcal{D}^{\mathbb{T}}$ - \mathbb{G}^{b} -stable if strict inequality holds in (6.17). Over a J-Gauduchon generalized Kähler manifold, the corresponding notions are simply referred to as the \mathbb{G}^{b} -slope, \mathbb{G}^{b} -semistable and \mathbb{G}^{b} -semistable respectively.

Recall that over a Hermitian manifold (M, g, I), the degree of any holomorphic vector bundle V is independent of a Hermitian metric on V iff g is Gauduchon, i.e.

(6.18)
$$\partial \overline{\partial} \left(\omega^{m-1} \right) = 0$$

where m is the complex dimension of M. On a generalized Kähler manifold, the J-Gauduchon condition can be expressed in a similar fashion.

Proposition 6.15. A generalized Kähler manifold $(M, \gamma; \mathbb{G}^b, \mathbb{J})$ is \mathbb{J} -Gauduchon iff

(6.19)
$$\partial_{+}\overline{\partial}_{+}\left(\omega_{+}^{m-1}\right) + (-1)^{\varepsilon} \partial_{-}\overline{\partial}_{-}\left(\omega_{-}^{m-1}\right) = 0$$

where $\varepsilon = 0$ if I_+ induce the same orientation on T M and $\varepsilon = 1$ otherwise.

Proof. Note that the degree of a Hermitian vector bundle coincides with that of its determinant line bundle, thus it's sufficient to consider line bundles. By (5.16) and (6.8)

$$\int_{M} \Lambda_{\mathbb{J}_{-}} \left(\partial_{\mathbb{J}}^{\phi, b} d_{\overline{L}} f \right) d\operatorname{vol}_{g} = \int_{M} \left[\Lambda_{+} \left(\partial_{+} \overline{\partial}_{+} f \right) + \Lambda_{-} \left(\partial_{-} \overline{\partial}_{-} f \right) \right] d\operatorname{vol}_{g}$$

The statement then follows from integration by parts.

Remark 6.16. Evidently, (6.19) holds for m < 3, since the 3-form below is closed:

$$\phi = \gamma + db = \mp d_{\pm}^c \omega_{\pm}$$

For $m \ge 3$, (6.19) is equivalent to

$$\phi \wedge d \left(\omega_+^{m-2} - (-1)^{\varepsilon} \, \omega_-^{m-2} \right) = 0$$

In particular, $(M, \gamma; \mathbb{G}^b, \mathbb{J})$ is \mathbb{J} -Gauduchon if the difference $\omega_+^{m-2} - (-1)^{\varepsilon} \omega_-^{m-2}$ is closed. When m = 3, (6.19) can also be rewritten as

$$\phi_+^{(2,1)} \wedge \phi_+^{(1,2)} + (-1)^{\epsilon} \phi_-^{(2,1)} \wedge \phi_-^{(1,2)} = 0$$

where, for instance, $\phi_{\pm}^{(2,1)}$ denote the (2, 1)-component of ϕ with respect to I_{\pm} respectively. It is clear that the generalized Kähler manifold is J-Gauduchon if g is Gauduchon with respect to both I_{\pm} .

7. GEOMETRIC LAX FLOWS

A Lax pair [24] consists of two families of operators $\{(L_t, P_t) : t \in I \subseteq \mathbb{R}\}$ such that

(7.1)
$$\frac{d}{dt}P_t = \left[L_t, P_t\right]$$

where $\{L_t\}$ is the *Lax operator* and it is assumed that $0 \in I$. The equation (7.1) is also said to be in *the Lax form*. Suppose that $\{\Psi_t\}$ is generated by $\{L_t\}$, i.e. it solves the equation

(7.2)
$$\frac{d}{dt}\Psi_t = L_t\Psi_t, \text{ with } \Psi_0 = \text{Id}$$

then $\{P_t\}$ can be obtained from pushing-forward of an initial operator P_0 by $\{\Psi_t\}$:

$$P_t = \Psi_t P_0 \Psi_t^{-1}$$

In particular, $\{P_t\}$ is an isospectral family.

Let $\{A_t\}$ be a smooth family of operators on the same space, then the *t*-differential of $\{\Psi_t^{-1}A_t\Psi_t\}$ describes the extend to which $\{(L_t, A_t)\}$ fails to be a Lax pair as well.

Definition 7.1. Suppose that $\{(L_t, P_t)\}$ is a Lax pair and $\{A_t\}$ be a smooth family of operators on the same space. The L_t -differential of A_t (along the Lax flow) is:

(7.3)
$$\delta_L A_t := \frac{d}{dt} A_t - \left[L_t, A_t \right]$$

It is straightforward to verify that commutativity with P_t is preserved by δ_L .

Lemma 7.2. If A_t and P_t commute for all t, then $\delta_L A_t$ also commute with P_t for all t. *Proof.* Notice that

$$\frac{d}{dt}\left(\Psi_t^{-1}A_t\Psi_t\right) = \Psi_t^{-1}\left(\delta_L A_t\right)\Psi_t$$

 \Box

from which the statement follows.

When a pair of geometric quantities forms a Lax pair, the corresponding equation (7.1) is said to generate a *geometric Lax flow*. Two main classes of examples will be described, where the operators $\{P_t\}$ are either generalized metrics or generalized almost complex structures. Such Lax pairs impose certain necessary conditions on the Lax operator.

Definition 7.3. Let \mathbb{G}^b be a generalized metric and \mathbb{J} a generalized (almost) complex structure. An operator \mathbb{L} on $C^{\infty}(\mathbb{T}M)$ is *Lax compatible with* \mathbb{G}^b if

(7.4)
$$\left\langle \mathbb{L}x^{b+}, y^{b-} \right\rangle + \left\langle x^{b+}, \mathbb{L}y^{b-} \right\rangle = 0$$

for all $x^{b+} \in C^{\infty}(C^b_+)$ and $y^{b-} \in C^{\infty}(C^b_-)$. The operator \mathbb{L} is *Lax compatible with* \mathbb{J} if

(7.5)
$$\langle \mathbb{L}x, y \rangle + \langle x, \mathbb{L}y \rangle = \langle \mathbb{L}\overline{x}, \overline{y} \rangle + \langle \overline{x}, \mathbb{L}\overline{y} \rangle = 0$$

for all $x, y \in C^{\infty} \left(\mathbb{T}^{1,0}_{\mathbb{J}} M \right)$.

Lemma 7.4. Suppose that $\{(\mathbb{L}_t, \mathbb{P}_t)\}$ is a Lax pair of operators on $C^{\infty}(\mathbb{T}M)$. If \mathbb{P}_t is a smooth family of generalized metrics or almost complex structures, then \mathbb{L}_t is Lax compatible with \mathbb{P}_t for all t.

Proof. It follow from the orthogonality of \mathbb{P}_t with respect to the pairing \langle, \rangle and that \mathbb{P}_t^2 are constant operators.

The following simplified criteria are useful in practice.

Corollary 7.5. Let $\mathbb{L} \in C^{\infty}$ (End $(\mathbb{T}M)$) and $x, y \in \mathbb{T}M$. Then

(1) (7.4) holds for \mathbb{L} if the bilinear forms $\mathbb{G}^{b}(\mathbb{L}x, y)$ are symmetric, i.e.

$$\mathbb{G}^{b}(\mathbb{L}x, y) = \mathbb{G}^{b}(\mathbb{L}y, x)$$

(2) (7.4) and (7.5) hold for \mathbb{L} if the bilinear forms $\langle \mathbb{L}x, y \rangle$ are skew-symmetric, i.e.

$$\langle \mathbb{L}x, y \rangle + \langle \mathbb{L}y, x \rangle = 0$$

 \Box

Proof. Left for the reader.

7.1. $\theta \in \Omega^2_{\mathbb{T}}(M)$ as the Lax operator. Let $\{P_t \in C^{\infty}(T^*M^{\otimes 2})\}$ be a smooth family of 2-tensors on M, whose symmetric and skew-symmetric parts are respectively P_t^s and P_t^a , i.e.

$$P_t = P_t^s + P_t^a$$
 with $P_t^s(X, Y) = P_t^s(Y, X), P_t^a(X, Y) = -P_t^a(Y, X)$

for $X, Y \in C^{\infty}(TM)$. Then $\{P_t\}$ defines an initial value problem for a family of generalized metrics $\mathbb{G}_t^{b_t}$ as follows:

(7.6)
$$\begin{cases} \frac{d}{dt}g_t &= -P_t^s \\ \frac{d}{dt}b_t &= P_t^a \end{cases}$$

The system (7.6) can be reformulated into a Lax flow for $\mathbb{G}_t^{b_t}$.

Lemma 7.6. Let $\{P_t \in C^{\infty}(T^*M^{\otimes 2})\}$ and $\{\theta_t \in \Omega^2_{\mathbb{T}}(M)\}$ be a smooth family of $\mathbb{T}M$ -forms such that

(7.7) $\theta_t\left(x^{b_t-}, y^{b_t+}\right) = P_t(X, Y)$

where $x^{b_t \pm} \in C^{\infty}\left(C^{b_t}_{\pm}\right)$ with $X = \pi\left(x^{b_t \pm}\right)$ and so on. Let $\theta_t : \mathbb{T}M \to \mathbb{T}M$ be given by $2\langle \theta_t(x), y \rangle := \theta_t(x, y)$

Then (7.6) is equivalent to the 2-tensor Lax flow

(7.8)
$$\frac{d}{dt} \mathbb{G}_t^{b_t} = \left[\theta_t, \mathbb{G}_t^{b_t}\right]$$

Proof. Note that θ_t satisfies Corollary 7.5 (2). Since $(\mathbb{G}^b)^2 = 1$, the differential of a smooth family of generalized metrics is skew with respect to the generalized metric at each time. Obviously the left hand side of (7.8) is skew with respect to $\mathbb{G}_t^{b_t}$. Thus only the mixed $\mathbb{G}_t^{b_t}$ -eigencomponents of the left hand side are non-trivial, one of which goes as follows:

$$0 = \frac{d}{dt} \left\langle \mathbb{G}_{t}^{b_{t}} x^{b_{t}-}, y^{b_{t}+} \right\rangle$$
$$= \left\langle \frac{d}{dt} \mathbb{G}_{t}^{b_{t}} x^{b_{t}-}, y^{b_{t}+} \right\rangle + \left\langle \mathbb{G}_{t}^{b_{t}} \frac{d}{dt} \left(b_{t} - g_{t} \right) X, y^{b_{t}+} \right\rangle + \left\langle \mathbb{G}_{t}^{b_{t}} x^{b_{t}-}, \frac{d}{dt} \left(b_{t} + g_{t} \right) Y \right\rangle$$
$$= \left\langle \frac{d}{dt} \mathbb{G}_{t}^{b_{t}} x^{b_{t}-}, y^{b_{t}+} \right\rangle + \frac{d}{dt} \left(b_{t} - g_{t} \right) (X, Y)$$

The right hand side is given by

$$\left\langle \left[\theta_{t}, \mathbb{G}_{t}^{b_{t}}\right] x^{b_{t}-}, y^{b_{t}+} \right\rangle = \left\langle \left(\theta_{t} \mathbb{G}_{t}^{b_{t}} - \mathbb{G}_{t}^{b_{t}} \theta_{t}\right) x^{b_{t}-}, y^{b_{t}+} \right\rangle = -2 \left\langle \theta_{t} \left(x^{b_{t}-}\right), y^{b_{t}+} \right\rangle = -P_{t} \left(X, Y\right)$$
Thus (7.8) gives rise to

Thus (7.8) gives rise to

$$\frac{d}{dt}\left(b_{t}-g_{t}\right)\left(X,Y\right)=P_{t}\left(X,Y\right)$$

from which (7.6) follows. The other direction is left for the reader.

A smooth conformal family of metrics $\{g_t\}$ can be seen as a solution to (7.6), by setting $P_t = P_t^s = f_t g_0$, where $\{f_t \in C^{\infty}(M)\}$ is a smooth family of functions. In this case, $b_t \equiv b_0$ is constant throughout the flow. The corresponding $\mathbb{T}M$ -forms as given in Lemma 7.6 can be chosen to be dependent only on the generalized metric \mathbb{G}^b .

Definition 7.7. Let \mathbb{G}^b be a generalized metric defined by (g, b) where g is a Riemannian metric. A $\mathbb{T}M$ -form $\theta \in \Omega^2_{\mathbb{T}}(M)$ is \mathbb{G}^b -conformal if there are $r, s \in C^{\infty}(M)$ such that

$$\theta\left(x^{b-}, y^{b+}\right) = rg\left(X, Y\right) + sb\left(X, Y\right) \text{ and } \theta\left(x^{b\pm}, y^{b\pm}\right) = 0$$

where $x^{b\pm} \in C^{\infty}(C^{b}_{\pm})$ with $X = \pi(x^{b\pm})$ and so on. *r*, *s* are called the *conformal weights*.

Remark 7.8. When the conformal weights coincide, i.e. r = s, a family of \mathbb{G}^{b} -conformal forms generates conformal deformations of the generalized metric \mathbb{G}^{b} . Otherwise, the metric *g* and *b* are deformed by different factors. In particular, when the conformal weight s = 0, it corresponds to classical conformal deformation of the metric *g*.

40

7.2. (ϕ, b) -curvature Lax flow. A special case of (7.8) is when θ_t are $\mathbb{d}^{\phi,b}$ -exact, i.e. $\theta_t = \mathbb{d}^{\phi_t,b_t}u_t$ for a smooth family of sections $\{u_t \in \Omega^1_{\mathbb{T}}(M)\}$ and 3-forms $\{\phi_t \in \Omega^3(M)\}$. Suppose that $u_t = Z_t + \zeta_t$ then

$$\begin{pmatrix} \mathbb{d}^{\phi_t, b_t} u_t \end{pmatrix} \begin{pmatrix} x^{b_t-}, y^{b_t+} \end{pmatrix} = X u_t \begin{pmatrix} y^{b_t+} \end{pmatrix} - Y u_t \begin{pmatrix} x^{b_t-} \end{pmatrix} - u_t \begin{pmatrix} x^{b_t-} \diamond_{\phi_t, b_t} y^{b_t+} \end{pmatrix}$$
$$= \left(\mathcal{L}_{Z_t} g_t \right) (X, Y) + \left[d \left(\zeta_t - \iota_{Z_t} b_t \right) - \iota_{Z_t} \phi_t \right] (X, Y)$$

It follows that in this case, the Lax flow (7.8) is equivalent to

(7.9)
$$\begin{cases} \frac{d}{dt}g_t = -\mathcal{L}_{Z_t}g_t \\ \frac{d}{dt}b_t = -\mathcal{L}_{Z_t}b_t + d\zeta_t - \iota_{Z_t}\left(\phi_t - db_t\right) \end{cases}$$

When $\gamma = \phi_t - db_t$ is a fixed closed 3-form, (7.9) describes the push-forward of an initial generalized metric by the family of generalized diffeomorphism of $\mathbb{T}M$ generated by u_t .

Proposition 7.9. Let $\gamma \in \Omega^3(M)$ be a closed 3-form and $u_t = Z_t + \zeta_t \in C^{\infty}(\mathbb{T}M)$ be a smooth family of sections. Let (λ_t, β_t) be the family of generalized diffeomorphisms generated by $\{u_t\}$ under $*_{\gamma}$ (recalled below). Set $\phi_t = \gamma + db_t$, then (7.9) coincides with the infinitesimal action by (λ_t, β_t) on a generalized metric \mathbb{G}^b via push-forward.

Proof. It's more straightforward to work with pull-back action on generalized metrics. Recall [23] that in (λ_t, β_t) , λ_t is the 1-parameter family of diffeomorphisms generated by Z_t , and

$$\beta_t := \int_0^t \lambda_s^* \left(d\zeta_s - \iota_{Z_s} \gamma \right) ds$$

The push-forward of $x = X + \xi \in \mathbb{T}M$ by (λ_t, β_t) is given by

$$\left(\lambda_{t},\beta_{t}\right)_{*}x=\lambda_{t*}\left(x+\iota_{X}\beta_{t}\right)=\lambda_{t*}X+\left(\lambda_{t}^{-1}\right)^{*}\left(\xi+\iota_{X}\beta_{t}\right)$$

with the corresponding infinitesimal action $x \mapsto -u_t *_{\gamma} x$.

The pull-back of \mathbb{G}^{b} by (λ_{t}, β_{t}) gives the family of generalized metrics

$$\mathbb{G}_{t}^{b_{t}}(x, y) := \mathbb{G}^{b}\left(\left(\lambda_{t}, \beta_{t}\right)_{*} x, \left(\lambda_{t}, \beta_{t}\right)_{*} y\right)$$

where $x, y \in C^{\infty}(\mathbb{T}M)$ are independent of *t*. Analogous to the computations in the classical case, differentiate the above with respect to *t* gives by the left hand side

$$\begin{pmatrix} \frac{d}{dt} \mathbb{G}_t^{b_t} \end{pmatrix} (x, y) = \frac{d}{dt} \left[g_t (X, Y) + g_t^{-1} \left(\xi - \iota_X b_t, \eta - \iota_Y b_t \right) \right]$$

=
$$\frac{1}{2} \left[\left(\frac{d}{dt} g_t \right) (X, Y) - \left(\frac{d}{dt} g_t \right) \left(g_t^{-1} \xi_t', g_t^{-1} \eta_t' \right) - \left(\frac{d}{dt} b_t \right) \left(Y, g_t^{-1} \xi_t' \right) - \left(\frac{d}{dt} b_t \right) \left(X, g_t^{-1} \eta_t' \right) \right]$$

where $\xi'_t = \xi - \iota_X b_t$ and $\eta'_t = \eta - \iota_Y b_t$; while by the right hand side

$$\begin{pmatrix} \frac{d}{dt} \mathbb{G}_{t}^{b_{t}} \end{pmatrix} (x, y) = Z_{t} \mathbb{G}_{t}^{b_{t}} (x, y) + \mathbb{G}_{t}^{b_{t}} \left(-u_{t} *_{\gamma} x, y \right) + \mathbb{G}_{t}^{b_{t}} \left(x, -u_{t} *_{\gamma} y \right)$$

$$= \frac{1}{2} \left[\left(\mathcal{L}_{Z_{t}} g_{t} \right) (X, Y) - \left(\mathcal{L}_{Z_{t}} g_{t} \right) \left(g_{t}^{-1} \xi_{t}', g_{t}^{-1} \eta_{t}' \right) \right]$$

$$+ \frac{1}{2} \left[\left(-\mathcal{L}_{Z_{t}} b + d\zeta_{t} - \iota_{Z_{t}} \gamma \right) \left(Y, g_{t}^{-1} \xi_{t}' \right) + \left(-\mathcal{L}_{Z_{t}} b + d\zeta_{t} - \iota_{Z_{t}} \gamma \right) \left(X, g_{t}^{-1} \eta_{t}' \right) \right]$$

Compare these two ways of computing $\left(\frac{d}{dt}\mathbb{G}_{t}^{b_{t}}\right)(x, y)$, the pull-back action on \mathbb{G}^{b} gives

$$\begin{cases} \frac{d}{dt}g_t = \mathcal{L}_{Z_t}g_t \\ \frac{d}{dt}b_t = \mathcal{L}_{Z_t}b_t - (d\xi_t - \iota_{Z_t}\gamma) \end{cases}$$

The push-forward action reverses the signs on the right hand side, which is (7.9).

A $d^{\phi,b}$ -exact $\mathbb{T}M$ -form in $\Omega^2_{\mathbb{T}}(M)$ can be seen as the (ϕ, b) -curvature of a unitary generalized connection on the trivial Hermitian line bundle. In general, let (V, h) be a Hermitian line bundle with a family of unitary generalized connections $\{D_t\}$. By Example 3.8, the Lax flow (7.8) defined by $\{\theta_t = \sqrt{-1}\mathcal{F}^{\phi_t,b_t}(D_t)\}$ where $\phi_t = \gamma + db_t$ is equivalent to the system

(7.10)
$$\begin{cases} \frac{d}{dt}g_t = -\mathcal{L}_{g_t^{-1}\psi_t}g_t \\ \frac{d}{dt}b_t = \sqrt{-1}F_0^{b_t} - \iota_{g_t^{-1}\psi_t}\phi_t \end{cases}$$

which reduces to (7.9) when V is trivial, and admits similar interpretation in terms of (not necessarily exact) generalized diffeomorphisms.

Theorem 7.10. Given a family of unitary generalized connections $\{D_t\}$ on a line bundle V with Hermitian metrics $\{h_t\}$, the Lax flow (7.8) defined by $\{\theta_t = \mathcal{F}^{\phi_t, b_t}(D_t)\}$, where $\phi_t = \gamma + db_t$, corresponds to the push-forward of an initial generalized metric \mathbb{G}^b by a family of generalized diffeomorphisms, which may not be exact, i.e. is not generated by global sections of $\mathbb{T}M$.

Proof. In the notations of Example 3.8, the local section defining D_t is given by $\sqrt{-1u_t} = \sqrt{-1}(Z_t + \zeta_t)$, where Z_t is a global vector field and ζ_t is a locally defined 1-form, with

$$Z_{t} = g_{t}^{-1}\psi_{t} = \frac{1}{2}g_{t}^{-1}\left(v_{t,+} - v_{t,-}\right) \text{ and } \zeta_{t} = \iota_{Z_{t}}b_{t} + \frac{1}{2}\left(v_{t,+} + v_{t,-}\right)$$

Then $\nabla_0^{b_t}$ is defined by the local section $\frac{1}{2}(v_{t,+} + v_{t,-})$, which implies that

$$\sqrt{-1}F_0^{b_t} = d\left(\zeta_t - \iota_{Z_t}b_t\right) = -\mathcal{L}_{Z_t}b_t + d\zeta_t + \iota_{Z_t}db_t$$

Locally, the second equation in (7.10) is thus

$$\frac{d}{dt}b_t = -\mathcal{L}_{Z_t}b_t + d\zeta_t - \iota_{Z_t}\gamma$$

noting that $\phi_t = \gamma + db_t$. By Proposition 7.9, (7.10) corresponds to pushing-forward by the local exact generalized diffeomorphisms generated by $\{(Z_t, \zeta_t)\}$. Globally, (7.10) corresponds to pushing-forward by possibly non-exact generalized diffeomorphisms.

Example 7.11. Let $(M, \gamma; \mathbb{J})$ be a generalized complex manifold and $\{\theta_t \in \Omega^2_{\mathbb{T}}(M)\}$ be a smooth family of $\mathbb{T}M$ -forms. The Lax flow with initial value $\mathbb{J}_0 = \mathbb{J}$

$$\frac{d}{dt}\mathbb{J}_t = \left[\theta_t, \mathbb{J}_t\right]$$

consists of generalized almost complex structures. The flow above preserves \mathbb{J} iff $[\theta_t, \mathbb{J}] = 0$ for all *t*, which is equivalent to $\theta_t \in \Omega^{1,1}_{\mathbb{J}}(M)$. Thus, starting with a generalized Hermitian manifold $(M, \gamma; \mathbb{G}^b, \mathbb{J})$, the Lax flow (7.8) defined by $\{\theta_t \in \Omega^{1,1}_{\mathbb{J}}(M)\}$ produces a family of generalized Hermitian structures with the same \mathbb{J} .

Suppose furthermore that $(M, \gamma; \mathbb{G}^b, \mathbb{J})$ admits a γ - \mathbb{J} -connection \mathcal{D}^T . Fix a \mathbb{J} -holomorphic line bundle V and let $\{\theta_t\}$ be the \mathcal{D}^T -Chern curvatures for a smooth family of Hermitian metrics $\{h_t\}$ on V. By Proposition 6.3, $\{\theta_t\}$ consists of (1, 1)-forms with respect to \mathbb{J} . By Theorem 7.10, the corresponding Lax flow (7.8) corresponds to the push-forward of \mathbb{G}^b by a family of generalized diffeomorphisms $\{(\lambda_t, \beta_t)\}$. Alternatively, it can be seen as a family of generalized Hermitian structures $(M, \gamma; \mathbb{G}^b, (\lambda_t, \beta_t)^* \mathbb{J})$ with \mathbb{G}^b fixed.

7.3. Ricci Lax flow. When the Ricci curvature of a generalized connection $\mathcal{D}^{\mathbb{T}}$ satisfies (7.4), it can serve as the Lax operator in a Lax pair involving the generalized metric.

Definition 7.12. A smooth family of pairs $\left\{ \left(\mathbb{G}_{t}^{b_{t}}, \mathcal{D}^{\mathbb{T}_{t}} \right) \right\}$ of generalized metrics $\mathbb{G}_{t}^{b_{t}}$ and $\mathbb{G}_{t}^{b_{t}}$ -metric connections is a solution to the $\mathcal{D}^{\mathbb{T}}$ -*Ricci Lax flow* if $\left\{ \mathcal{R}ic^{\mathbb{T}_{t}} \right\}$ satisfies (7.4) and the pair $\left\{ \left(\mathcal{R}ic^{\mathbb{T}_{t}}, \mathbb{G}_{t}^{b_{t}} \right) \right\}$ form a Lax pair, i.e.

(7.11)
$$\frac{d}{dt}\mathbb{G}_{t}^{b_{t}} = \left[\mathcal{R}ic^{\mathbb{T}_{t}}, \mathbb{G}_{t}^{b_{t}}\right]$$

When $\mathcal{D}^{\mathbb{T}_t}$ is prescribed to depend on $\mathbb{G}_t^{b_t}$, (7.11) becomes an equation for $\mathbb{G}_t^{b_t}$ only, in which case the family of generalized metrics $\mathbb{G}_t^{b_t}$ is said to be *a solution* to (7.11). If $\mathcal{D}^{\mathbb{T}_t} = \mathcal{D}^{\phi_t, b_t}$, then the flow (7.11) is simply called the *Ricci Lax flow*.

Since $\mathcal{R}c^{\phi,b}$ is symmetric, it satisfies the condition in Corollary 7.5. Theorem 7.13 below shows that the Ricci Lax flow is equivalent to

(7.12)
$$\frac{d}{dt} \left(g_t \mp b_t \right) = -2Rc_t^{\pm \phi_t} = -2Rc_t + \frac{1}{2}\phi_t^2 \pm d^*\phi_t$$

Let $\gamma \in \Omega^3(M)$ such that $d\gamma = 0$. The following generalized Ricci flow [9, 30] is a system for Riemannian metrics g_t and $b_t \in \Omega^2(M)$:

(7.13)
$$\begin{cases} \frac{d}{dt}g_t = -2Rc_t + \frac{1}{2}\left(\gamma + db_t\right)^2\\ \frac{d}{dt}b_t = -d^*\left(\gamma + db_t\right) \end{cases}$$

then (7.12) and (7.13) coincide when $\phi_t = \gamma + db_t$.

Theorem 7.13. Under the further constraints that $\mathcal{D}^{\mathbb{T}} = \mathcal{D}^{\phi,b}$ and is metric compatible with the Dorfman bracket $*_{\gamma}$ (2.31), i.e. $\phi_t - db_t = \gamma \in \Omega^3(M)$ for all t, the Ricci Lax flow (7.11) is equivalent to the generalized Ricci flow (7.13).

Proof. It's obvious that (7.13) follows from (7.12), by matching the symmetric and skew-symmetric terms on both sides. To obtain (7.12) from (7.11), fix $X, Y \in TM$ and consider

$$x^{b_t-} = X + (b_t - g_t) X$$
 and $y^{b_t+} = Y + (b_t + g_t) Y$

The left hand side of (7.12) is computed in Lemma 7.6, while the right hand side becomes:

$$\left\langle \left[\mathcal{R}ic^{\phi_t, b_t}, \mathbb{G}_t^{b_t} \right] x^{b_t^-}, y^{b_t^+} \right\rangle = \left\langle \left(\mathcal{R}ic^{\phi_t, b_t} \mathbb{G}_t^{b_t} - \mathbb{G}_t^{b_t} \mathcal{R}ic^{\phi_t, b_t} \right) x^{b_t^-}, y^{b_t^+} \right\rangle$$
$$= -2\mathcal{R}c^{\phi_t, b_t} \left(x^{b_t^-}, y^{b_t^+} \right) = -2\mathcal{R}c^{+\phi_t} \left(X, Y \right)$$

This gives half of (7.12). The other half is equivalent.

Example 7.14. Consider two 3-dimensional Lie groups: the 3-torus $T = (U(1))^3$ and G = SU(2), with their invariant metrics g and invariant volume forms ϕ . For both, set b = 0 and consider the corresponding $(\phi, 0)$ -Ricci curvature $\mathcal{R}ic^{\phi,0}$.

For *T*, the invariant metric is flat, thus Ric = 0 while $Ric^{\pm \phi} \neq 0$ by (4.8). In the \mathbb{G}^{b} -eigendecomposition, $\mathcal{R}c^{\phi,0}$ is of the form

$$\mathcal{R}c^{\phi,0} = \begin{pmatrix} 0 & R \\ R^T & 0 \end{pmatrix}$$

where $R = 1_3$, the 3 × 3 identity matrix.

For G, the invariant metric is the standard round metric on S^3 which is not flat, while the connections with torsion $\pm \phi$ are flat, hence $Ric^{\pm \phi} = 0$. In this case, $\mathcal{R}c^{\phi,0}$ is of the form

$$\mathcal{R}c^{\phi,0} = \begin{pmatrix} R & 0\\ 0 & R \end{pmatrix}$$

where *R* here is the classical Ricci tensor for the round metric (c.f. $\S2.5$).

Definition 7.7 and Theorem 7.13 lead to the natural generalization of Ricci soliton.

Definition 7.15. Let $\gamma \in \Omega^3(M)$ be a closed 3-form. A smooth family of generalized metrics $\{\mathbb{G}_t^{b_t}\}$ is a γ -*Ricci Lax soliton* if there exists a smooth family of sections $\{u_t \in C^\infty(\mathbb{T}M)\}$ and $\mathbb{G}_t^{b_t}$ -conformal forms $\{\theta_t \in \Omega^2_{\mathbb{T}}(M)\}$ with constant conformal weights r_t and s_t , and

(7.14)
$$\left[\mathcal{R}ic^{\phi_t,b_t} - \mathbb{d}^{\phi_t,b_t}u_t - \theta_t, \mathbb{G}_t^{b_t}\right] = 0$$

where $\phi_t = \gamma + db_t$. The family is a *gradient* γ -*Ricci Lax soliton* if furthermore there is $f \in C^{\infty}(M)$ such that $\left\{ u_t = -\frac{1}{2} \mathbb{G}_t^{b_t}(df) \right\}$.

Notice that $\mathbb{G}_t^{b_t}(df) = \nabla^t f + \iota_{\nabla^t f} b_t$, where $\nabla^t f$ is the gradient of f with respect to g_t . When $s_t = 0$, the gradient γ -Ricci Lax soliton equation is then equivalent to the system

(7.15)
$$\begin{cases} Rc_t - \frac{1}{4}\phi_t^2 + (\nabla^t)^2 f = r_t g_t \\ d^*\phi_t + \iota_{\nabla^t f}\phi_t = 0 \end{cases}$$

where $(\nabla^t)^2 f$ is the Hessian of f with respect to the Levi-Civita connection of g_t . When $r_t \equiv r$ is a constant function independent of t, the system (7.15) is exactly the *generalized Ricci soliton* equation (Apostolov-Streets-Ustinovskiy [3] and references therein).

7.4. **Bismut-Ricci Lax flow.** Even though $\mathcal{R}ic^{\phi,B}$ is neither symmetric nor skew-symmetric, it satisfies (7.4), and thus can be used to define a Lax flow for generalized metrics. In (7.11), taking $\mathcal{R}ic^{\phi,B}$ as the Lax operator leads to the *Bismut-Ricci Lax flow*:

(7.16)
$$\frac{d}{dt}\mathbb{G}_{t}^{b_{t}} = \left[\mathcal{R}ic_{t}^{\phi_{t},B_{t}},\mathbb{G}_{t}^{b_{t}}\right]$$

Since $\mathcal{R}c^{\phi,B}$ has the same mixed components as $\mathcal{R}c^{\phi,b}$, i.e.

$$\left[\mathcal{R}ic^{\phi,b},\mathbb{G}^{b}
ight]=\left[\mathcal{R}ic^{\phi,B},\mathbb{G}^{b}
ight]$$

by Theorem 7.13, (7.16) is equivalent to (7.12), so they generate the same flow for the generalized metric [29]. The $\mathcal{R}ic^{\phi,B}$ -differential of \mathcal{D}^{ϕ_t,B_t} takes a particularly simple form.

Proposition 7.16. Fix $X, Y, Z \in C^{\infty}(TM)$ and $x \in C^{\infty}(\mathbb{T}M)$, with $\pi(x) = X$. Let $y^{b_t\pm}, z^{b_t\pm} \in C^{b_t}_+$ so that $\pi(y^{b_t\pm}) = Y$ and $\pi(z^{b_t\pm}) = Z$. Then

(7.17)
$$\mathbb{G}_{t}^{b_{t}}\left(\left(\delta_{\mathcal{R}ic^{\phi,B}}\mathcal{D}_{x}^{\phi_{t},\mathcal{B}_{t}}\right)y^{b_{t}\pm},z^{b_{t}\pm}\right) = -g_{t}\left(Y,\frac{d}{dt}\nabla_{X}^{\pm\phi_{t}}Z\right)$$

Proof. Only the case for $C_{+}^{b_t}$ is shown here and the case for $C_{-}^{b_t}$ is similar.

$$\begin{split} & \mathbb{G}_{t}^{b_{t}}\left(\frac{d}{dt}\left(\mathcal{D}_{x}^{\phi_{t},B_{t}}\right)y^{b_{t}+},z^{b_{t}+}\right) \\ &= \frac{d}{dt}\left[\mathbb{G}_{t}^{b_{t}}\left(\mathcal{D}_{x}^{\phi_{t},B_{t}}y^{b_{t}+},z^{b_{t}+}\right)\right] - \mathbb{G}_{t}^{b_{t}}\left(\mathcal{D}_{x}^{\phi_{t},B_{t}}\left[\frac{d}{dt}\left(b_{t}+g_{t}\right)Y\right],z^{b_{t}+}\right) - \mathbb{G}_{t}^{b_{t}}\left(\mathcal{D}_{x}^{\phi_{t},B_{t}}y^{b_{t}+},\frac{d}{dt}\left(b_{t}+g_{t}\right)Z\right) \\ &= g_{t}\left(\frac{d}{dt}\nabla_{x}^{+\phi_{t}}Y,Z\right) + \left[XRc_{t}^{-\phi}\left(Y,Z\right) - Rc_{t}^{-\phi}\left(Y,\nabla_{x}^{+\phi_{t}}Z\right) - Rc_{t}^{-\phi}\left(\nabla_{x}^{+\phi_{t}}Y,Z\right)\right] \end{split}$$

where the last equality follows from (7.12) together with the fact that $\mathcal{D}^{\phi_t, B_t}$ preserves $\mathbb{G}_t^{b_t}$. Next, (4.15) implies that

$$\begin{split} & \mathbb{G}_{t}^{b_{t}}\left(\left[\mathcal{R}ic_{t}^{\phi_{t},B_{t}},\mathcal{D}_{x}^{\phi_{t},B_{t}}\right]y^{b_{t}+},z^{b_{t}+}\right) = \mathbb{G}_{t}^{b_{t}}\left(\mathcal{R}ic_{t}^{\phi_{t},B_{t}}\mathcal{D}_{x}^{\phi_{t},B_{t}}y^{b_{t}+} - \mathcal{D}_{x}^{\phi_{t},B_{t}}\mathcal{R}ic_{t}^{\phi_{t},B_{t}}y^{b_{t}+},z^{b_{t}+}\right) \\ = \mathcal{R}c_{t}^{\phi_{t},B_{t}}\left(\mathcal{D}_{x}^{\phi_{t},B_{t}}y^{b_{t}+},z^{b_{t}+}\right) - \mathcal{R}\mathcal{R}c_{t}^{\phi_{t},B_{t}}\left(y^{b_{t}+},z^{b_{t}+}\right) + \mathcal{R}c_{t}^{\phi_{t},B_{t}}\left(y^{b_{t}+},\mathcal{D}_{x}^{\phi_{t},B_{t}}z^{b_{t}+}\right) \\ = -\mathcal{R}\mathcal{R}c_{t}^{+\phi_{t}}\left(Y,Z\right) + \mathcal{R}c_{t}^{+\phi_{t}}\left(Y,\nabla_{x}^{+\phi_{t}}Z\right) + \mathcal{R}c_{t}^{+\phi_{t}}\left(\nabla_{x}^{+\phi_{t}}Y,Z\right) \end{split}$$

Combining the results above and applying (7.12) again lead to

$$\begin{split} & \mathbb{G}_{t}^{b_{t}}\left(\left(\delta_{\mathcal{R}ic^{\phi,B}}\mathcal{D}_{x}^{\phi_{t},B_{t}}\right)y^{b_{t}\pm},z^{b_{t}\pm}\right) \\ &= g_{t}\left(\frac{d}{dt}\nabla_{X}^{+\phi_{t}}Y,Z\right) - \left[X\left(\frac{d}{dt}g_{t}\right)(Y,Z) - \left(\frac{d}{dt}g_{t}\right)\left(Y,\nabla_{X}^{+\phi_{t}}Z\right) - \left(\frac{d}{dt}g_{t}\right)\left(\nabla_{X}^{+\phi_{t}}Y,Z\right)\right] \\ &= -g_{t}\left(Y,\frac{d}{dt}\nabla_{X}^{+\phi_{t}}Z\right) \end{split}$$

П

where the last equality follows from the fact that $\nabla^{+\phi_t}$ preserves g_t for all t.

Consider a smooth family of generalized almost Hermitian structures $(M, \gamma; \mathbb{G}_t^{b_t}, \mathbb{J}^t)$, where $\mathbb{G}_t^{b_t}$ is a solution to the Bismut-Ricci Lax flow (7.16). By Lemma 7.2, the $\mathcal{R}ic^{\phi,B}$ differential $\delta_{\mathcal{R}ic^{\phi,B}}\mathbb{J}^t$ preserves the $\mathbb{G}_t^{b_t}$ -eigenbundles. Following computations similar to those in Proposition 7.16 gives:

(7.18)
$$\mathbb{G}_{t}^{b_{t}}\left(\left(\delta_{\mathcal{R}ic^{\phi,\beta}}\mathbb{J}^{t}\right)y^{b\pm},z^{b\pm}\right) = -g_{t}\left(Y,\left(\frac{d}{dt}I_{\pm}^{t}\right)Z\right)$$

Set $\mathbb{J}_{-}^{t} := \mathbb{G}_{t}^{b_{t}} \mathbb{J}^{t}$ then

$$\delta_{\mathcal{R}ic^{\phi,B}}\mathbb{J}_{-}^{t} = \left(\delta_{\mathcal{R}ic^{\phi,B}}\mathbb{G}_{t}^{b_{t}}\right)\mathbb{J}^{t} + \mathbb{G}_{t}^{b_{t}}\left(\delta_{\mathcal{R}ic^{\phi,B}}\mathbb{J}^{t}\right) = \mathbb{G}_{t}^{b_{t}}\left(\delta_{\mathcal{R}ic^{\phi,B}}\mathbb{J}^{t}\right)$$

It implies that \mathbb{J}^t solves the following Lax flow (7.19), in which case as does \mathbb{J}^t_{\pm} , iff I^t_{\pm} are constant almost complex structures:

(7.19)
$$\frac{d}{dt}\mathbb{J}^{t} = \left[\mathcal{R}ic^{\phi_{t},B_{t}},\mathbb{J}^{t}\right]$$

In terms of the 2-forms ω_{\pm} , (7.19) is equivalent to the following simultaneous almost Hermitian Ricci flows:

(7.20)
$$\left(\frac{d}{dt}\omega_{\pm}^{t}\right)(X,Y) = 2Rc\left(X,I_{\pm}^{t}Y\right) - \frac{1}{2}\phi_{t}^{2}\left(X,I_{\pm}^{t}Y\right)$$

where $X, Y \in C^{\infty}(TM)$. It has the classical Kähler-Ricci flow as a special case.

Example 7.17. Take a family of classical Kähler structures (g_t, I, ω_t) , set

$$\mathbb{J}^{t} = \begin{bmatrix} I & 0 \\ 0 & -I^{*} \end{bmatrix}, \ \mathbb{J}^{t}_{-} = \begin{bmatrix} 0 & -\omega_{t}^{-1} \\ \omega_{t} & 0 \end{bmatrix}$$

The generalized Bismut connections, as well as the (ϕ, b) -connections, are simply the lift of the Levi-Civita connections for g_t . Let $X, Y \in TM$, since $\phi = 0$, (7.20) becomes

$$\left(\frac{d}{dt}\omega_t\right)(X,Y) = 2Rc_t(X,IY) = -2\rho_t(X,Y)$$

where $\rho_t(X, Y) = Rc_t(IX, Y)$ is the Ricci form. Thus (7.19) recovers exactly the equation for the classical Kähler-Ricci flow.

REFERENCES

- [1] Ilka Agricola and Thomas Friedrich, *A note on flat metric connections with antisymmetric torsion*, Diff. Geom. Appl. 28 (2010), 480 487.
- [2] Anton Alekseev, Henrique Bursztyn and Eckhard Meinrenken, *Pure spinors on Lie groups*, Astérisque No. 327 (2009), pp. 131 – 199.
- [3] Vestislav Apostolov, Jeffrey Streets and Yury Ustinovskiy, Variational structure and uniqueness of generalized Kähler-Ricci solitons, arXiv:2109.10295v1.
- [4] Jean-Michel Bismut, A local index theorem for non Kähler manifolds, Math. Ann. 284 (1989), pp. 681 699.
- [5] Kenneth A. Brakke, *The motion of a surface by its mean curvature*, Mathematical Notes, 20, Princeton University Press, Princeton, N.J., 1978.
- [6] Élie Cartan and Jan A. Schouten, On the geometry of the group manifold of simple and semisimple groups, Proc. Amsterdam 29 (1926), 803 815.
- [7] Élie Cartan and Jan A. Schouten, On Riemannian manifolds admitting an absolute parallelism, Proc. Amsterdam 29 (1926), 933 – 946.
- [8] Thomas Friedrich and Stefan Ivanov, *Parallel spinors and connections with skew-symmetric torsion in string theory*, Asian J. Math. Vol 6, No 2. (2002), pp. 303 336.
- [9] Mario Garcia-Fernandez and Jeffrey Streets, *Generalized Ricci Flow*, University Lecture Series, AMS, 2020.
- [10] Alfred Gray and Louis M. Hervella, *The sixteen classes of almost Hermitian manifolds and their linear invariants*, Ann. Mat. Pura Appl. (IV) CXXIII (1980), pp. 35 58.
- [11] Ryushi Goto, Scalar curvature as moment map in generalized Kahler geometry, Journal of symplectic geometry 18 (2020), pp. 147 – 190.
- [12] Ryushi Goto, Scalar curvature and the moment map in generalized Kahler geometry, arXiv:2105.13654
- [13] Phillip A. Griffiths, *Linearizing flows and a cohomological interpretation of Lax equations*, American Journal of Mathematics, Vol. 107, No. 6 (Dec., 1985), pp. 1445 1484.
- [14] Marco Gualtieri, Generalized complex geometry, Oxford Ph.D. thesis, 2003. arXiv:DG/0412211. https://doi.org/10.48550/arXiv.math/0401221
- [15] Marco Gualtieri, Branes on Poisson varieties, in The Many Facets of Geometry: A Tribute to Nigel Hitchin, Oxford University Press, 2010, pp. 368 – 394.
- [16] Marco Gualtieri, Generalized complex geometry, Annals of Mathematics. (2). 174 (1) (2011), pp. 75 123.
- [17] Marco Gualtieri, Generalized Kähler Geometry, Commun. Math. Phys. 331, 297 331 (2014).
- [18] Richard S. Hamilton, *Three-manifolds with positive Ricci curvature*. J. Differential Geom. 17 (1982), no. 2, pp. 255 – 306.
- [19] Nigel Hitchin, Generalized Calabi-Yau manifolds, Q. J. Math. 54 (2003), pp. 281 308.
- [20] Nigel Hitchin, *Generalized holomorphic bundles and the B-field action*, J. Geom. Phys. 61 (1) (2011), pp. 352 362.
- [21] Shengda Hu, On generalized Kähler geometry on compact Lie groups, preprint, arXiv:1501.00754v1. https://doi.org/10.48550/arXiv.1501.00754
- [22] Shengda Hu, Ruxandra Moraru and Reza Seyyidali, A Kobayashi-Hitchin correspondence for I_{\pm} holomorphic bundles, Advances in Mathematics 287 (2016), pp. 519 – 566.
- [23] Shengda Hu and Bernardo Uribe, *Extended manifolds and extended equivariant cohomology*, Journal of Geometry and Physics, 59 (1) (2009), pp. 104 131.
- [24] Peter Lax, *Integrals of nonlinear equations of evolution and solitary waves*, Communications on Pure and Applied Mathematics, 21 (5) (1968), pp. 467 490.
- [25] Martin Lübke and Andrei Teleman, *The Kobayashi-Hitchin Correspondence*, World Scientific Publishing Co., Inc., River Edge, NJ, 1995.
- [26] Pavol Severa and Fridrich Valach, Ricci flow, Courant algebroids, and renormalization of Poisson-Lie T-duality, Letters in Mathematical Physics, volume 107 (2017), pp. 1823 – 1835.

- [27] Pavol Ševera and Fridrich Valach, *Courant algebroids, Poisson-Lie T-duality, and type II supergravities,* Communications in Mathematical Physics, volume 375 (2020), pp. 307 – 344.
- [28] Jeffrey Streets, *Regularity and expanding entropy for connection Ricci flow*, Journal of Geometry and Physics 58 (2008), 900 912.
- [29] Jeffrey Streets, *Generalized geometry, T-duality, and renormalization group flow*, Journal of Geometry and Physics 114 (2017), pp. 506 522.
- [30] Jeffrey Streets and Gang Tian, *Generalized Kähler geometry and the pluriclosed flow*, Nuclear Physics B, 858 (2) (2012), pp. 366 376.
- [31] Yicao Wang, *Toric generalized Kähler structures III*, Journal of Geometry and Physics 151 (2020). *Email address*: shu@wlu.ca

DEPARTMENT OF MATHEMATICS, WILFRID LAURIER UNIVERSITY, 75 UNIVERSITY AVE. WEST, WATERLOO, CANADA

48