

PREPARED FOR SUBMISSION TO JHEP

An Approach to Quantum 2D Gravity

Vladimir V. Belokurov^{a,b} and Evgeniy T. Shavgulidze^a

^a*Lomonosov Moscow State University,*

^b*Institute for Nuclear Research of the Russian Academy of Sciences,*

E-mail: vvbelokurov@yandex.ru, shavgulidze@bk.ru

ABSTRACT: We consider a model of 2D gravity with the action quadratic in curvature and represent path integrals as integrals over the $SL(2, \mathbb{R})$ invariant Gaussian functional measure. We reduce these path integrals to the products of Wiener path integrals and calculate the correlation function of the metric in the first perturbative order.

KEYWORDS: 2D Gravity, Path integrals, Wiener measure

ARXIV EPRINT: [.](https://arxiv.org/abs/2206.05172)

Contents

1	Introduction	1
2	$SL(2, \mathbb{R})$ invariant Gaussian measure	2
3	An algorithm for path integrals calculation	3
4	Correlation function of the metric	5
5	Conclusion	7

1 Introduction

The enormous popularity of 2D gravity in the last several decades motivated by its role in string theory and studies of BH physics in the dimensional reduction approach has grown after realizing the Schwarzian nature of the JT dilaton gravity and the relation of this theory to SYK model (for some pedagogical reviews see, e.g., [1]-[3]).

The general form of the 2D gravity action up to the terms quadratic in curvature K is

$$\tilde{\mathcal{A}} = c_0 \int \sqrt{\mathcal{G}} d^2x + c_1 \int K \sqrt{\mathcal{G}} d^2x + c_2 \int K^2 \sqrt{\mathcal{G}} d^2x. \quad (1.1)$$

The first two terms do not determine the dynamics of 2D gravity. While the part of the action quadratic in the Gaussian curvature does. It is the model we consider in this paper.

Commonly it is transformed to the dilaton gravity action (see, e.g., the review [4]). An alternative way is to deal only with the geometric structures of the surface. (Some examples of this approach are given in [5]-[17]).

The action (1.1) is invariant under general coordinate transformations. Here, we reduce the set of coordinate transformations and consider the action restricted to the conformal gauge, where the metric of the 2D surface looks like

$$dl^2 = g(u, v) (du^2 + dv^2) = g(z, \bar{z}) dz d\bar{z} \quad \sqrt{\mathcal{G}} = g. \quad (1.2)$$

The Gaussian curvature of the surface is [18]

$$K = -\frac{1}{2g} \Delta \log g, \quad (1.3)$$

where Δ stands for the Laplacian.

Note that the action (1.1) is invariant under the complex analytic substitutions

$$z = \chi(\zeta), \quad \chi g(z, \bar{z}) = g(\chi(\zeta), \bar{\chi}(\bar{\zeta})) |\chi'_\zeta|^2, \quad dz d\bar{z} = |\chi'_\zeta|^2 d\zeta d\bar{\zeta}. \quad (1.4)$$

Therefore, we suppose the integration in the action (1.1) to be over the disc $d : (|z| \leq 1)$.

2 $SL(2, \mathbb{R})$ invariant Gaussian measure

We consider the specific form the action (1.1)

$$A = \frac{\lambda^2}{2} \int_d (K + 4)^2 g(z, \bar{z}) dz d\bar{z} = \frac{\lambda^2}{2} \int_d (\Delta\psi)^2 dz d\bar{z} \quad (2.1)$$

where

$$\Delta\psi = q \Delta \log q + \frac{4}{q}, \quad q = \frac{1}{\sqrt{g}}. \quad (2.2)$$

Now path integrals in the theory

$$\int \tilde{F}(g) \exp\{-\tilde{A}(g)\} dg \quad (2.3)$$

are path integrals

$$\int F(\psi) \exp\{-A(\psi)\} d\psi \quad (2.4)$$

over the Gaussian functional measure

$$\mu_\lambda(d\psi) = \frac{\exp\{-A(\psi)\} d\psi}{\int \exp\{-A(\psi)\} d\psi}. \quad (2.5)$$

The extremum of the action (2.1) is given by the equation $\Delta\psi = 0$, or

$$q \Delta \log q + \frac{4}{q} = 0. \quad (2.6)$$

We choose the boundary condition corresponding to the Poincare model of the Lobachevsky plane [18]

$$q_0|_{|z|=1} = 0.$$

The unique solution in the disk $d(|z| \leq 1)$ satisfying the boundary condition is

$$q_0 = 1 - z\bar{z}. \quad (2.7)$$

Let us rewrite the action (2.1) substituting $\psi \rightarrow f$ with

$$\Delta\psi = L[f] \quad (2.8)$$

where L is the differential operator

$$L \equiv q_0 \Delta - \frac{8}{q_0} \equiv \frac{1}{q_0} T^{-1}. \quad (2.9)$$

Note that $T^{-1} \equiv (q_0^2 \Delta - 8)$ is the Casimir operator of $SL(2, \mathbb{R})$ [19].

Now the action (2.1) is written as the integral over the measure $\frac{dz d\bar{z}}{(1-z\bar{z})^2}$ invariant under the action of the group $SL(2, \mathbb{R})$ in the disk

$$A = \frac{\lambda^2}{2} \int_d (T^{-1}[f])^2 \frac{dz d\bar{z}}{q_0^2}. \quad (2.10)$$

Therefore, we obtain the $SL(2, \mathbb{R})$ invariant Gaussian functional measure¹

$$\mu_\lambda(df) = \frac{\exp\{-A(f)\} df}{\int \exp\{-A(f)\} df} . \quad (2.11)$$

Note that the $SL(2, \mathbb{R})$ invariance of the measure guarantees the invariance for path integrals $\int F(f) \mu_\lambda(df)$.

3 An algorithm for path integrals calculation

In this section, we reduce path integrals over the measure (2.11) to the products of Wiener integrals. Here, it is convenient to use polar coordinates.

After the linear substitution (2.8) $\Delta\psi(\varrho, \varphi)$ is written as

$$\Delta\psi(\varrho, \varphi) = L[f(\varrho, \varphi)] = L_0[f(\varrho, \varphi)] + \frac{(1 - \varrho^2)}{\varrho^2} \frac{\partial^2}{\partial \varphi^2} f(\varrho, \varphi) . \quad (3.1)$$

where the differential operator L_0 is

$$L_0 \equiv (1 - \varrho^2) \frac{\partial^2}{\partial \varrho^2} + \frac{(1 - \varrho^2)}{\varrho} \frac{\partial}{\partial \varrho} - \frac{8}{(1 - \varrho^2)} . \quad (3.2)$$

Consider the Fourier series

$$\Delta\psi(\varrho, \varphi) = x_0(\varrho) + \sum_{n=1}^{\infty} (x_n \cos n\varphi + y_n \sin n\varphi) \quad (3.3)$$

and

$$f(\varrho, \varphi) = a_0(\varrho) + \sum_{n=1}^{\infty} (a_n \cos n\varphi + b_n \sin n\varphi) . \quad (3.4)$$

The coefficients of the first Fourier series can be expressed in terms of the coefficients of the second series

$$x_n(\varrho) = L_0[a_n(\varrho)] - n^2 \frac{(1 - \varrho^2)}{\varrho^2} a_n(\varrho) \quad (3.5)$$

and

$$y_n(\varrho) = L_0[b_n(\varrho)] - n^2 \frac{(1 - \varrho^2)}{\varrho^2} b_n(\varrho) . \quad (3.6)$$

Now the action (2.1) is written as

$$A = \frac{\lambda^2}{2} 2\pi \int_0^1 (x_0(\varrho))^2 \varrho d\varrho + \frac{\lambda^2}{2} \pi \sum_{n=1}^{\infty} \left[\int_0^1 (x_n(\varrho))^2 \varrho d\varrho + \int_0^1 (y_n(\varrho))^2 \varrho d\varrho \right] . \quad (3.7)$$

¹In what follows, we will not write down the normalization factor explicitly.

After the substitution

$$a_n(\varrho) = \varrho^n \frac{(1+n+(1-n)\varrho^2)}{(1-\varrho^2)} \int_{\varrho}^1 \frac{(1-\varrho_1^2)^2 V_n(\varrho_1)}{\varrho_1^{2n+1} (1+n+(1-n)\varrho_1^2)^2} d\varrho_1, \quad (3.8)$$

we obtain

$$\int_0^1 (x_n(\varrho))^2 \varrho d\varrho = \int_0^1 \frac{(1-\varrho^2)^4}{\varrho^{2n+1} (1+n+(1-n)\varrho^2)^2} (V'_n(\varrho))^2 d\varrho, \quad n=0,1,\dots, \quad (3.9)$$

where

$$V_n(\varrho) = \varrho^{2n+1} \frac{(1+n+(1-n)\varrho^2)^2}{(1-\varrho^2)^2} \left(\frac{(1-\varrho^2) a_n(\varrho)}{\varrho^n (1+n+(1-n)\varrho^2)} \right)'. \quad (3.10)$$

Therefore we have

$$\int_0^1 (x_n(\varrho))^2 \varrho d\varrho = \int_0^{+\infty} (U'_n(\tau_n))^2 d\tau_n \quad (3.11)$$

with

$$\tau_n = \int_0^{\varrho} \frac{\varrho_1^{2n+1}}{(1-\varrho_1^2)^2} \left(\frac{2}{(1-\varrho_1^2)} + n-1 \right)^2 d\varrho_1, \quad U_n(\tau_n) = V_n(\varrho). \quad (3.12)$$

The same equations are valid for the other terms in (3.7)

$$\int_0^1 (y_n(\varrho))^2 \varrho d\varrho = \int_0^{+\infty} (\tilde{U}'_n(\tau_n))^2 d\tau_n. \quad (3.13)$$

Now the measure $\mu_\lambda(df)$ is represented as the product of the Wiener measures²

$$\mu_\lambda(df) = w_{\frac{1}{\lambda\sqrt{2\pi}}}(dU_0) \prod_{n=1}^{\infty} w_{\frac{1}{\lambda\sqrt{\pi}}}(dU_n) w_{\frac{1}{\lambda\sqrt{\pi}}}(d\tilde{U}_n) \quad (3.14)$$

where

$$w_\sigma(dU) = \exp \left\{ -\frac{1}{2\sigma^2} \int_0^{+\infty} (U'(\tau))^2 d\tau \right\} dU. \quad (3.15)$$

For path integrals with integrands that depend on the modes with definite numbers, the product of the measures is reduced because of the cancelation of the same terms in the nominator and in the denominator.

The simplest basic path integrals over the measure $\mu_\lambda(df)$ are

$$\int_{C([0,1])} V_n(\varrho_1) V_n(\varrho_2) \mu_\lambda(df) \quad (3.16)$$

²See the footnote 1 in section (2).

with

$$0 \leq \varrho_1 \leq \varrho_2 < 1, \quad \tau_{n(1)} = \tau_n(\varrho_1) \leq \tau_n(\varrho_2) = \tau_{n(2)}.$$

For $n = 0$, (3.16) is equal to³)

$$\begin{aligned} \int_{C([0,1])} V_0(\varrho_1) V_0(\varrho_2) \mu_\lambda(df) &= \int_{C([0,+\infty))} U_0(\tau_{0(1)}) U_0(\tau_{0(2)}) w_{\frac{1}{\lambda\sqrt{2\pi}}} (dU_0) \\ &= \frac{1}{2\pi\lambda^2} \tau_{0(1)} \\ &= \frac{1}{2\pi\lambda^2} \int_0^{\varrho_1} \frac{\varrho}{(1-\varrho^2)^2} \left(\frac{2}{1-\varrho^2} - 1 \right)^2 d\varrho = \frac{1}{24\pi\lambda^2} \left(\frac{2}{1-\varrho_1^2} - 1 \right)^3. \end{aligned} \quad (3.17)$$

For $n \geq 1$,

$$\begin{aligned} \int_{C([0,1])} V_n(\varrho_1) V_n(\varrho_2) \mu_\lambda(df) &= \int_{C([0,+\infty))} U_n(\tau_{n(1)}) U_n(\tau_{n(2)}) w_{\frac{1}{\lambda\sqrt{\pi}}} (dU_n) \\ &= \frac{1}{\pi\lambda^2} \tau_{n(1)} = \frac{1}{\pi\lambda^2} \int_0^{\varrho} \frac{\varrho^{2n+1}}{(1-\varrho^2)^2} \left(\frac{2}{1-\varrho^2} + n - 1 \right)^2 d\varrho. \end{aligned} \quad (3.18)$$

More complicated path integrals with higher powers of integration variables are calculated using the Wick theorem.

4 Correlation function of the metric

In this section, we calculate the correlation function of the metric given by the integral

$$\langle g(\varrho_1, \varphi_1) g(\varrho_2, \varphi_2) \rangle_\mu = \int_{C(d)} g(\varrho_1, \varphi_1) g(\varrho_2, \varphi_2) \mu_\lambda(df) \quad (4.1)$$

in the first nontrivial perturbative order.

Due to the $SL(2, \mathbb{R})$ invariance, the correlation function of the metric can be rewritten as

$$\begin{aligned} \langle g(\varrho_1, \varphi_1) g(\varrho_2, \varphi_2) \rangle_\mu &= \frac{q_0^2(\varrho_*)}{q_0^2(\varrho_1) q_0^2(\varrho_2)} \int_{C(d)} g(0, 0) g(\varrho_*, \varphi_*) \mu_\lambda(df) \\ &= \frac{q_0^2(\varrho_*)}{q_0^2(\varrho_1) q_0^2(\varrho_2)} \langle g(0, 0) g(\varrho_*, \varphi_*) \rangle_\mu, \end{aligned} \quad (4.2)$$

where $(0, 0)$ and (ϱ_*, φ_*) are the results of the shift at the Lobachevsky plane of the coordinates (ϱ_1, φ_1) and (ϱ_2, φ_2) ($0 \leq \varrho_* \leq 1$, $0 \leq \varphi_* \leq 2\pi$). In particular,

$$\varrho_* = \frac{\sqrt{\varrho_1^2 + \varrho_2^2 - 2\varrho_1\varrho_2 \cos(\varphi_2 - \varphi_1)}}{\sqrt{1 + \varrho_1^2\varrho_2^2 - 4\varrho_1\varrho_2 \cos(\varphi_2 - \varphi_1)}}. \quad (4.3)$$

³See, e.g., [20]

The integration measure is invariant under rotations and averaged values depend only on the variable ϱ .

Let us represent the function $q(\varrho, \varphi)$ as the series in the powers of the variable f

$$q(\varrho, \varphi) = q_0(\varrho) (1 + p_1(f(\varrho, \varphi)) + p_2(f(\varrho, \varphi)) + \dots), \quad (4.4)$$

From (2.2) and (2.8), it follows that⁴

$$p_1(f) = f \quad (4.5)$$

and

$$L \left[p_2 - \frac{1}{2} f^2 \right] = f L[f] - \frac{8}{q_0} f^2. \quad (4.6)$$

For first nontrivial order calculations, we expand the metric up to the quadratic in f terms

$$g(\varrho_*, \varphi_*) = \frac{1}{q_0^2(\varrho_*)} \left(1 - 2f(\varrho_*, \varphi_*) + 3f^2(\varrho_*, \varphi_*) - 2 \frac{p_2(f(\varrho_*, \varphi_*))}{q_0(\varrho_*)} \right) \quad (4.7)$$

where p_2 is given by (4.6) and (ϱ_*, φ_*) is a point inside the disk d .

Note that, due to the $SL(2, \mathbb{R})$ invariance,

$$\langle g(\varrho, \varphi) \rangle_\mu = \int_{C(d)} g(\varrho, \varphi) \mu_\lambda(df) = \frac{1}{q_0^2(\varrho)} \langle g(0, 0) \rangle_\mu. \quad (4.8)$$

Taking into account (4.8) and (4.7), we obtain

$$\langle g(0, 0) g(\varrho_*, \varphi_*) \rangle_\mu = \frac{2}{q_0^2(\varrho_*)} \{ \langle g(0, 0) \rangle_\mu + 2 \langle f(0, 0) f(\varrho_*, \varphi_*) \rangle_\mu \}. \quad (4.9)$$

Now we calculate the first term in the sum in the r.h.s. of (4.9)

$$\langle g(0, 0) \rangle_\mu \equiv \langle g \rangle_\mu(0) = 1 + 3 \langle f^2 \rangle_\mu(0) - 2 \langle p_2 \rangle_\mu(0). \quad (4.10)$$

Using the $SL(2, \mathbb{R})$ invariance of the integration measure, we obtain from (4.6)

$$\langle p_2 \rangle_\mu(0) = \frac{1}{2} \langle f^2 \rangle_\mu(0) - 8 \langle f^2 \rangle_\mu(0) L_0^{-1} \left[\frac{1}{q_0} \right](0), \quad (4.11)$$

with

$$L_0^{-1} [h(\varrho_1)](\varrho) = -(1 + \varrho^2) \int_{\varrho}^1 \frac{(1 - \varrho_2^2)^2}{\varrho_2 (1 + \varrho_2^2)^2} \int_0^{\varrho_2} \frac{\varrho_1 (1 + \varrho_1^2)}{(1 - \varrho_1^2)^2} h(\varrho_1) d\varrho_1 d\varrho_2.$$

Obviously,

$$L_0^{-1} \left[\frac{1}{q_0} \right](0) = -\frac{1}{8},$$

and therefore

$$\langle g \rangle_\mu(0) = 1. \quad (4.12)$$

⁴Only the terms essential in the studied perturbative order ($\sim \frac{1}{\lambda^2}$) are explicitly written down.

The second term in (4.9) is reduced to

$$\langle f(0, 0) f(\varrho_*, \varphi_*) \rangle_\mu = \langle a_0(0) f(\varrho_*, \varphi_*) \rangle_\mu = \langle a_0(0) a_0(\varrho_*) \rangle_\mu \quad (4.13)$$

and can be calculated using the technique of the previous section.

Thus, the path integration in (4.13) results in

$$\begin{aligned} & \langle f(0, 0) f(\varrho_*, \varphi_*) \rangle_\mu \\ &= \frac{1}{6\pi\lambda^2} \int_{\varrho}^1 \frac{(1 - \varrho_2^2)^2}{\varrho_2 (1 + \varrho_2^2)^2} \int_{\varrho_*}^{\varrho_2^2} \frac{\varrho_1 (1 + \varrho_1^2)}{(1 - \varrho_1^2)^2} \frac{\varrho_1^2 (3 + \varrho_1^4)}{(1 - \varrho_1^2)^3} d\varrho_1 d\varrho_2 \\ &+ \frac{1}{12\pi\lambda^2} \int_{\varrho}^1 \frac{(1 - \varrho_2^2)^2}{\varrho_2 (1 + \varrho_2^2)^2} d\varrho_2 \int_0^{\varrho_*} \frac{\varrho_1 (1 + \varrho_1^2)}{(1 - \varrho_1^2)^2} \frac{\varrho_1^2 (3 + \varrho_1^4)}{(1 - \varrho_1^2)^3} d\varrho_1 \\ &= \frac{1}{12\pi\lambda^2} \left\{ -\frac{1}{2} \int_{\varrho_*}^1 \frac{\log t}{1 - t} dt + 2 \log 2 + \frac{1}{2} - \frac{1}{1 + \varrho_*^2} + \frac{1}{1 + \varrho_*^2} \log \left(\frac{1 - \varrho_*^2}{\varrho_*} \right) \right. \\ &\quad \left. + \log \varrho_* - 2 \log(1 + \varrho_*^2) - \log \sqrt{1 - \varrho_*^2} + \log \varrho_* \log \sqrt{1 - \varrho_*^2} \right\}. \quad (4.14) \end{aligned}$$

In the limit $\varrho_* \rightarrow 1$, we have an interesting (and expected) physical result. The quantum corrections to the correlation function of the metric tend to zero and⁵

$$\langle g(0, 0) g(\varrho_*, \varphi_*) \rangle_\mu \rightarrow \langle g(0, 0) \rangle_\mu \langle g(\varrho_*, \varphi_*) \rangle_\mu.$$

5 Conclusion

In this paper, we consider 2d gravity with the action quadratic in the Gaussian curvature of the surface. We rewrite the theory in the form invariant under the action of the group $SL(2, \mathbb{R})$ in the disk and then we treat path integrals in the theory as integrals over the $SL(2, \mathbb{R})$ invariant Gaussian functional measure. Then we demonstrate how these path integrals can be reduced to the products of Wiener path integrals and, as an example, calculate the correlation function of the metric in the first perturbation theory order.

In [21], we reduced path integrals of quadratic gravity in FLRW metric to Wiener path integrals. The elaborated in this paper method may be considered as a first step in generalizing the approach of [21] to higher dimensions.

In some other context, path integrals related with Casimir operator of $SL(2, \mathbb{R})$ where studied in [22]. However the measure used in [22] $\exp \{ (T^{-1}h, h) \} dh$ differs from the measure obtained in our paper $\exp \{ (T^{-1}h, T^{-1}h) \} dh$ (see (2.11), (2.10)).

⁵The factor 2 stands because of the symmetry $\varrho_* \Leftrightarrow 0$.

References

- [1] P. Ginsparg and G. W. Moore, Lecture on 2-d gravity and 2-d string theory, [arXiv:hep-th/9304011].
- [2] T. Banks, Lectures on black holes and information loss, Nucl. Phys. Proc. Suppl. **41** (1995) 21, [arXiv:hep-th/9412131].
- [3] G. Sarosi, AdS2 holography and the SYK model, arXiv:1711.08482.
- [4] D. Grumiller, W. Kummer, and D. V. Vassilevich, Dilaton gravity in two dimensions, Phys. Rept. **369** (2002) 327, [arXiv:hep-th/0204253].
- [5] M. O. Katanaev and I. V. Volovich, String model with dynamical geometry and torsion, Phys. Lett. **B175** (1986) 413, [arXiv:gr-qc/9511009].
- [6] M. O. Katanaev and I. V. Volovich, Two-dimensional gravity with dynamical torsion and strings, Ann. Phys. **197** (1990) 1, [arXiv:gr-qc/9511009].
- [7] J. G. Russo and A. A. Tseytlin, Scalar tensor quantum gravity in two dimensions, Nucl. Phys. **B382** (1992) 259, [arXiv:hep-th/9201021].
- [8] E. W. Mielke, F. Gronwald, Y. N. Obukhov, R. Tresguerres, and F. W. Hehl, Towards complete integrability of two-dimensional Poinkare gauge gravity, Phys. Rev. **D48** (1993) 3648, [arXiv:hep-th/9304043].
- [9] H. -J. Schmidt, The classical solutions of two-dimensional gravity, Gen. Rel. Grav. **31** (1999) 1187, [arXiv:gr-qc/9905051].
- [10] M. O. Katanaev, W. Kummer, and H. Liebl, Geometric interpretation and classification of global solutions in generalized dilaton gravity, Phys. Rev. **D53** (1996) 5609, [arXiv:gr-qc/9511009].
- [11] S. D. Odintsov and I. L. Shapiro, One loop renormalization of two-dimensional induced quantum gravity, Phys. Lett. **B263** (1991) 183.
- [12] S. D. Odintsov and I. L. Shapiro, Perturbative finiteness of 2-d induced quantum gravity, Mod. Phys. Lett. **A7** (1992) 437.
- [13] F. Haider and W. Kummer, Quantum functional integration of nonEinsteinian gravity in d=2, Int. J. Mod. Phys. **A9** (1994) 207.
- [14] W. Kummer, H. Liebl, and D. V. Vassilevich, Exact path integral quantization of generic 2-d dilaton gravity, Nucl. Phys. **B493** (1997) 491, [arXiv:gr-qc/9612012].
- [15] W. Kummer, H. Liebl, and D. V. Vassilevich, Non-perturbative path integral of 2d dilaton gravity and two-loop effects from scalar matter, Nucl. Phys. **B513** (1998) 723, [arXiv:hep-th/9707115].
- [16] W. Kummer, H. Liebl, and D. V. Vassilevich, Non-perturbative path integral of 2d dilaton gravity and two-loop effects from scalar matter, Nucl. Phys. **B544** (1999) 403, [arXiv:hep-th/9809168].
- [17] D. Grumiller, W. Kummer, and D. V. Vassilevich, The virtual black hole in 2d quantum gravity, Nucl. Phys. **B580** (2000) 438, [arXiv:gr-qc/0001038].
- [18] B. A. Dubrovin, S. P. Novikov and A. T. Fomenko, Modern Geometry - Methods and Applications Part I: The Geometry of Surfaces, Transformation Groups, and Fields (2 ed.), Springer-Verlag, 1992.

- [19] S. Lang, SL(2, R), Addison-Wesley Publishing, 1975.
- [20] Hui-Hsiung Kuo, Gaussian Measures in Banach Spaces, Springer, Berlin-Heidelberg-NY, 1975.
- [21] V. V. Belokurov and E. T. Shavgulidze, Path integrals in quadratic gravity, *JHEP* **02** (2022) 112 , [arXiv:2110.06041].
- [22] S. K. Ashok and J. Troost, Path integrals on $\mathfrak{sl}(2, \mathbb{R})$ orbits, arXiv:2204.00232.