ESSENTIAL DIMENSION OF CENTRAL SIMPLE ALGEBRAS OF DEGREE 8 AND EXPONENT 2 IN CHARACTERISTIC 2

ADAM CHAPMAN

ABSTRACT. The goal of this note is to reduce the existing upper bound for the essential dimension of central simple algebras of degree 8 and exponent 2 over fields of characteristic 2 from 10 to 9.

1. INTRODUCTION

Fixing a field k in the background, we consider the covariant functor Alg_{de} sending any field F containing k to the set of all (isomorphism classes of) central simple algebras of degree d and exponent dividing e over F. The essential dimension of an algebra A in $Alg_{d,e}(F)$ is the minimal possible transcendence degree of a field E over k with $k \subseteq E \subseteq F$ for which there exists a central simple algebra A_0 in $\operatorname{Alg}_{de}(E)$ with $A_0 \otimes F = A$. The essential dimension of $Alg_{d,e}$, denoted $ed(Alg_{d,e})$, is thus the supremum on the essential dimensions of A where A ranges over all algebras in $Alg_{de}(F)$ and F ranges over all fields containing k. The question of computing $ed(Alg_{d,e})$ is a very difficult one (see [Mer13, Section 10]) and the exact value is known only in specific cases: it is known that $ed(Alg_{2,2}) = ed(Alg_{3,3}) = 2$, but for any prime p > 3 it is still not known what $ed(Alg_{p,p})$ is. The value of $ed(Alg_{42})$ is 4 when $char(k) \neq 2$ and 3 when char(k) = 2. The value of $ed(Alg_{8,2})$ is 8 when $char(k) \neq 2$ ([BM12, Corollary 1.4]). The exact value of $ed(Alg_{8,2})$ when char(k) = 2 is not known, but was bounded from above by 10 and below by 3 in [Bae11]. The lower bound was improved to 4 in [McK17]. The goal of this short note is to prove the following theorem:

Theorem 1.1 (Main Theorem). When char(k) = 2, we have $ed(Alg_{8,2}) \le 9$.

This is proved in Section 3, and Section 2 describes the main ingredients.

2. Preliminaries

When F is a field of char(F) = 2, a quaternion algebra over F is of the form $[\alpha,\beta)_F = F\langle i,j : i^2 + i = \alpha, j^2 = \beta, jij^{-1} = i + 1 \rangle$ for some

²⁰¹⁰ Mathematics Subject Classification. 16K20 (primary); 16K50 (secondary).

Key words and phrases. Essential Dimension; Central Simple Algebras; Fields of Characteristic 2.

ADAM CHAPMAN

 $\alpha \in F$ and $\beta \in F^{\times}$. We write $\wp(t)$ for $t^2 + t$ and $F[\wp^{-1}(\alpha)]$ thus stands for $F[i:i^2+i=\alpha]$. By [GS17, Theorem 9.1.4] (credited to Teichmüller), every class in $_2Br(F)$ is represented by a tensor product of quaternion algebras. In particular, when A is a central simple algebra of degree 4 and exponent 2, it decomposes as a tensor product of two quaternion algebras ([Rac74]).

A tool that makes the study of the essential dimension of central simple algebras of degree 8 and exponent 2 easier is the chain lemma for quaternion algebras ([Dra83, Chapter 14, Theorem 7]): If $[\alpha, \beta)_F = [\alpha', \beta')_F$, then there exists $\delta \in F$ for which $[\alpha, \beta)_F = [\delta, \beta)_F = [\delta, \beta')_F = [\alpha', \beta')_F$. It is also good to have the necessary and sufficient condition for $[\alpha, \beta)_F$ to be split: $\alpha = x^2 + x + y^2\beta$ for some $x, y \in F$ ([Dra83, Page 104]). In [Bae11], Baek was using another necessary and sufficient condition for splitting: $\beta = x^2 + xy + y^2\alpha$ for some $x, y \in F$ ([GS17, Corollary 4.7.5]), but here we prefer to use the former condition.

Another important tool is the fact that the essential dimension of compositums of *n* cyclic quadratic field extensions of fields *F* containing *k* of char(*k*) = 2 is 1 (see [Led04, Lemma 2] and [BF03, Remark 3.8] for reference, and [CFM, Remark 4.6] for an explicit construction; despite the condition $|k| \ge 2^n$ appearing in [Led04, Lemma 2], the proof of the latter goes through under the weaker assumption that $|F| \ge 2^n$, and for our purpose this always applies, for in order for noncommutative division algebras to exist over *F*, the cardinality of *F* must be infinite). The meaning is that if $K = F[\wp^{-1}(\alpha), \wp^{-1}(\beta), \wp^{-1}(\gamma)]$ is a compositum of three cyclic quadratic extensions of a field *F* containing *k*, then there exists a field $L \supseteq k$ transcendence degree 1 over *k* containing some elements *a*, *b*, *c* such that the tensor product $L[\wp^{-1}(a), \wp^{-1}(b), \wp^{-1}(c)] \bigotimes_L F$ is *K*. In other words, the elements α, β, γ can be chosen to come from a single field *L* of transcendence degree at most 1 over *k*.

The last tool is the following lemma:

Lemma 2.1 ([Bae11, Lemma 3.3]). *Given a field F of characteristic 2 and* a field extension $E = F[i : i^2+i = \alpha]$ and $\beta \in F$, if $x^2+xy+y^2\alpha = u^2+uv+v^2\beta$ for some $x, y, u, v \in F$, then $[\beta, x + yi)_E = [\beta, y + v)_E$.

3. Proof of the main theorem

Consider a division algebra *A* of degree 8 and exponent 2 over a field *F* containing a subfield *k* of char(*k*) = 2. By [Row84], *A* contains a maximal subfield $K = F[\wp^{-1}(\alpha), \wp^{-1}(\beta), \wp^{-1}(\gamma)]$. By [Led04, Lemma 2], α and β and γ can be chosen to come from a single subfield $k \subseteq L \subseteq F$ of transcendence degree 1 over *k*. Let $E = F[\wp^{-1}(\alpha)] = F[\mu : \mu^2 + \mu = \alpha]$. Then A_E is Brauer equivalent to $[\beta, b]_E \otimes [\gamma, c)_E$ for some $b, c \in E$. The corestriction back to *F* is trivial, and so $[\beta, N_b)_F = [\gamma, N_c)_F$, where $N_b = b_0^2 + b_0b_1 + b_1^2\alpha$, and

 $N_c = c_0^2 + c_0c_1 + c_1^2\alpha$, given that $b = b_0 + b_1\mu$ and $c = c_0 + c_1\mu$ for some $b_0, b_1, c_0, c_1 \in F$ ([MM91]). By the chain lemma for quaternion algebras, there exists $\delta \in F$ for which $[\beta, N_b)_F = [\delta, N_b)_F = [\delta, N_c)_F = [\gamma, N_c)_F$. Therefore $\delta = \beta + u^2 + u + x^2N_b = \gamma + y^2 + y + z^2N_c = \lambda^2 + \lambda + t^2N_bN_c$ for some $u, x, y, z, \lambda, t \in F$. By replacing δ, y, λ with $\delta + u^2 + u, y + u, \lambda + u$ respectively, we can assume u = 0. Note that $y = \wp^{-1}(\beta + \gamma + x^2N_b + z^2N_c)$ and $\lambda = \wp^{-1}(\beta + x^2N_b + t^2N_bN_c)$. If x = 0 then $\delta = \beta$, which means $[\beta + \delta, b)_E$ is split. If $x \neq 0$, then from the equation $\delta = \beta + x^2N_b$ we obtain $(\beta + \delta)\frac{1}{x^2} = b_0^2 + b_0b_1 + b_1^2\alpha$. It follows then from Lemma 2.1 that $[\beta + \delta, b)_E = [\beta + \delta, b_1 + \frac{1}{x})_E$. Similarly, if z = 0 then $[\delta, bc)_E$ is split, and otherwise $[\gamma + \delta, c)_E = [\gamma + \delta, c_1 + \frac{1}{z})_E$, and if t = 0 then $[\delta, bc)_E$ is split and otherwise $[\delta, bc)_E = [\delta, \frac{1}{t} + b_0c_1 + b_1c_0 + b_1c_1)_E$.

Now, $A_E = [\beta, b)_E \otimes [\gamma, c) \sim_{Br} [\beta + \delta, b)_E \otimes [\gamma + \delta, c)_E \otimes [\delta, bc)_E$. In the expression $[\beta + \delta, b]_E \otimes [\gamma + \delta, c]_E \otimes [\delta, bc]_E$, each term is either split or isomorphic to $[\beta + \delta, b_1 + \frac{1}{r})_E$, $[\gamma + \delta, c_1 + \frac{1}{r})_E$ and $[\delta, \frac{1}{t} + b_0c_1 + b_1c_0 + b_1c_1)_E$ respectively. In either case, each term descends to F. If one of the terms is split, then A_E is the restriction of a biquaternion algebra $Q_1 \otimes Q_2$ from F to E, and thus $A = Q_1 \otimes Q_2 \otimes [\alpha, d]_F$ for some $d \in F^{\times}$. This means A is a decomposable algebra of degree 8 and exponent 2, and by [McK17, Corollary 5.9], A descends to a field of transcendence degree at most 4 over k. We may therefore continue with the case of $x, z, t \neq 0$. Hence, $A \sim_{Br} [\alpha, a)_F \otimes [\beta + \delta, b_1 + \frac{1}{x})_F \otimes [\gamma + \delta, c_1 + \frac{1}{z})_F \otimes [\delta, \frac{1}{t} + b_0 c_1 + b_1 c_0 + b_1 c_1)_F.$ Consider now the algebra $B = [\alpha, a]_T \otimes [\beta + \delta, b_1 + \frac{1}{r})_T \otimes [\gamma + \delta, c_1 + \frac{1}{r})_T$ $(\frac{1}{z})_T \otimes [\delta, \frac{1}{t} + b_0c_1 + b_1c_0 + b_1c_1)_T$ where $T = L(a, b_0, b_1, c_0, c_1, x, z, t, y, \lambda)$. Note that y and λ are algebraic over $L(a, b_0, b_1, c_0, c_1, x, z, t)$, and thus the transcendence degree of T over k is at most 9. In order to conclude the argument, we need to explain why B is of index 8 rather than 16, which means that it is $M_2(A_0)$ for some division algebra A_0 of degree 8 over T, and thus A descends to a degree 8 division algebra over a field of transcendence degree at most 9 over k. The restriction of B to $R = T[\varphi^{-1}(\alpha)]$ is Brauer equivalent to $[\beta + \delta, b_1 + \frac{1}{x})_R \otimes [\gamma + \delta, c_1 + \frac{1}{z})_R \otimes [\delta, \frac{1}{t} + b_0 c_1 + b_1 c_0 + b_1 c_1)_R$. Since $[\beta + \delta, b_1 + \frac{1}{x})_R = [\beta + \delta, b)_R$ (for the same equality $\delta = \beta + x^2 N_b$ holds true as before), $[\gamma + \delta, c_1 + \frac{1}{z})_R = [\gamma + \delta, c)_R$ and $[\delta, \frac{1}{t} + b_0c_1 + b_1c_0 + b_1c_1)_R = [\delta, bc)_R$, we get that $B \otimes R$ is Brauer equivalent to $[\beta, b)_R \otimes [\gamma, c)_R$. Thus, B is split by $T[\wp^{-1}(\alpha), \wp^{-1}(\beta), \wp^{-1}(\gamma)]$, and so its index is 8.

Remark 3.1. It seems that in the argument in [Bae11], the author is suggesting that one could assume x = z = 1 in our computation above. If we were talking only about the quadratic equations $\delta = \beta + x^2 N_b = \gamma + y^2 + y + z^2 N_c$ this would be correct, because replacing b_i and c_j with $\frac{b_i}{x}$ and $\frac{c_j}{z}$ would eliminate their occurrences in these equations. However, that would change

ADAM CHAPMAN

 $[\beta, b)_E$ to $[\beta, \frac{b}{x})_E$ and $[\gamma, c)_E$ to $[\gamma, \frac{c}{z})_E$. It is therefore not clear why one can assume x = z = 1. If this were true, that would reduce the upper bound of the essential dimension from 9 to 7.

Acknowledgements

The author wishes to thank the anonymous referee for the detailed reading of the manuscript and for the helpful comments and suggestions.

References

- [Bae11] Sanghoon Baek. Essential dimension of simple algebras in positive characteristic. C. R. Math. Acad. Sci. Paris, 349(7-8):375–378, 2011.
- [BF03] Grégory Berhuy and Giordano Favi. Essential dimension: a functorial point of view (after A. Merkurjev). Doc. Math., 8:279–330, 2003.
- [BM12] Sanghoon Baek and Alexander S. Merkurjev. Essential dimension of central simple algebras. *Acta Math.*, 209(1):1–27, 2012.
- [CFM] Adam Chapman, Mathieu Florence, and Kelly McKinnie. Common splitting fields of symbol algebras. *Manuscripta Math.* to appear.
- [Dra83] P.K. Draxl. Skew Fields, volume 81 of London Math. Soc. Lect. Notes. Cambridge University Press, 1983.
- [GS17] Philippe Gille and Tamás Szamuely. Central simple algebras and Galois cohomology, volume 165 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2017. Second edition of [MR2266528].
- [Led04] Arne Ledet. On the essential dimension of p-groups. In Galois theory and modular forms, volume 11 of Dev. Math., pages 159–172. Kluwer Acad. Publ., Boston, MA, 2004.
- [McK17] Kelly McKinnie. Essential dimension of generic symbols in characteristic *p*. *Forum Math. Sigma*, 5:e14, 30, 2017.
- [Mer13] Alexander S. Merkurjev. Essential dimension: a survey. *Transform. Groups*, 18(2):415–481, 2013.
- [MM91] P. Mammone and A. Merkurjev. On the corestriction of pⁿ-symbol. Israel J. Math., 76(1-2):73–79, 1991.
- [Rac74] M. L. Racine. A simple proof of a theorem of Albert. Proc. Amer. Math. Soc., 43:487–488, 1974.
- [Row84] Louis H. Rowen. Division algebras of exponent 2 and characteristic 2. J. Algebra, 90(1):71–83, 1984.

School of Computer Science, Academic College of Tel-Aviv-Yaffo, Rabenu Yeruham St., P.O.B 8401 Yaffo, 6818211, Israel

Email address: adam1chapman@yahoo.com