Coherent state excitations and string-added coherent states in gauge-gravity correspondence

Hai Lin

Shing-Tung Yau Center of Southeast University, Southeast University, Nanjing 210096, China

School of Mathematical Sciences, Southeast University, Nanjing 211189, China Yau Mathematical Sciences Center, Tsinghua University, Beijing 100084, China

Abstract

We analyze detailed properties of BPS coherent states and their connection to gravity. We interpret the group integral coherent state as a path integral over auxiliary variables coupled to the elementary letters of the theory. The eigenvalues of coherent state amplitudes can be viewed as collective coordinates of giant gravitons. Inspired by the above coherent states and by the integrability, we construct a new type of coherent states in the SL(2) sectors and their cousin PSU(1,1|2) sectors, analogous to the aforementioned coherent states. The large spin and small spin limits can be obtained by varying coherent state amplitudes. We add string words onto the BPS coherent states, and this gives rise to stringadded coherent states. The insertions of string multi-words can be viewed as operator-insertions in this path integral. We describe BPS states and near BPS states building upon these coherent states in gauge-gravity correspondence. For example, string-added coherent states and their near BPS spectra are analyzed. This approach is particularly convenient for heavy excited states.

1 Introduction

The gauge-gravity correspondence [1, 2, 3] is a nontrivial correspondence between a quantum theory with gravity in the bulk and a different quantum system on the boundary. The correspondence allows us to perform calculations related to superstring theory and quantum gravity from working on the quantum field theory side. On the other hand, the superstring theory provides the UV completion of supergravity, and is a UV-complete quantum gravity theory. Integrability [4] has greatly increased our understanding of the gauge-gravity correspondence. The correspondence also reveals the nature of the emergent spacetime e.g. [5]-[9]. As we see, the bulk emerges dynamically from the quantum mechanical description that lives in fewer dimensions.

On the gravity side, giant graviton branes [10]-[14] are excited states. In the context of gauge-gravity correspondence, emergent backreacted geometries correspond to highly excited states in the quantum field theory side, such as the bubbling geometries [15, 13, 14]. The states in the Hilbert space of the quantum field theory are explicitly mapped to the gravity side. Analysis in the field theory side shows that these different states live in the same Hilbert space. The dual large operators and their representation bases have been illuminated [13, 14]. These heavy operators involve emergent backreacted geometries in the dual quantum gravity system.

Here we focus on states which have interesting gravitational properties. One interesting type of states are coherent states [16, 17, 18]. Gravity dual of coherent states has been analyzed. The coherent states in this paper are related to the excited states of the gravitational spacetime. These set-ups help to address the question how do curved spacetimes emerge from dual quantum theory on the boundary. Coherent states also appear very widely in many other contexts of physics, and here we concentrate on special types of coherent states.

There are various representation bases for large operators. The large operators includes those describe giant gravitons and backreacted emergent geometries. The large operators also describe further excitations on these heavy excited states. The correlation functions between light operators and large operators can also be computed. The large operators can be expanded systematically in terms of representation bases. In some sense, the larger the operators are, there are more information that can be stored with the operators. For more details along some of these ideas, see e.g. [19]-[28],[8],[18],[16]. Various bases can be transformed into each other. Operators of giant gravitons are also analyzed by various important and insightful approaches [29]-[35] and [16]. These approaches are closely related to the scenarios needed for this paper. Various ideas have been put forward, in order to make the computations with large operators more efficient and convenient. Coherent state operators in gauge-gravity duality have also been considered in e.g. [36, 18, 37, 16, 17] and their related references. These different bases have different labelings due to their different symmetry properties of the multi-parameters of the coherent states. In particular, [16] has manifest permutation symmetry of the parameters. Mixed states and entangled states of coherent state operators in the gauge-gravity correspondence have also been considered [38]. Moreover, mixed states and entangled states of Young tableau states were considered [39]. Many interesting aspects of coherent states in relation to giant graviton states, were explored in [16, 17] and their related references. The multi-parameter in the coherent states are collective coordinates of giant gravitons.

The quantum gravitational system dual to these heavy excited state operators, involves backreacted emergent geometries. Analysis on the field theory side shows that these excited states live in the same Hilbert space of the gravity side. Since they live in the same Hilbert space, we can dynamically relate them using the Hamiltonian in the same Hilbert space. For example one can superpose states and compute transition probabilities between different states living in the gravity side, e.g. [40]-[44],[18],[37] and related discussions.

When the spacetime is dynamical and quantum, the local quantum field theory seems not adequate to describe these situations. On the other hand, string theory is able to describe these situations very well. The set-ups discussed in this paper can be put inside superstring theory, and the set-ups here are UV-complete. Somehow we have gone beyond local quantum field theory.

In Sec. 2, we analyze detailed properties of coherent states and their connection to gravity, among other things. We focus on above-mentioned heavy states. Inspired by the above coherent states and by the integrability, in Sec. 3, we construct a new type of coherent states in the SL(2) sectors and their cousins such as PSU(1,1|2) sectors, among other things. The large spin and small spin limits can be obtained by varying coherent state amplitudes.

In Sec. 4. we analyze string-added coherent states and their near BPS spectra, among other things. We can add some string words onto the BPS coherent states, and this gives the string-added coherent states. We identify and elaborate some classes of near BPS string-added coherent states, whose anomalous dimension energy can be extracted conveniently. The anormalous energy of the near BPS state is usually smaller than its bare dimension energy. Hence these energy are controllable on both gauge theory side and gravity side. One can expand around supersymmetric backgrounds to compute their anormalous energy from both gauge theory and gravity side. These states can be considered as excitations on the BPS background. This method is very useful for near BPS states.

Finally in Sec. 5, we make conclusions and discuss some closely related aspects. Note

added: Upon the completion of this work, we received the appearance of [17] which have worked out SO(N) and Sp(N) cases; Many analysis in the current paper here also works for these and other very interesting cases.

2 BPS coherent states and collective variables

We consider a new class of BPS coherent states with manifest permutation symmetries. This class of interesting coherent states in nonabelian gauge theories have been constructed in [16]. Here, we first analyze more detailed properties of the states. We use $4D \ \mathcal{N} = 4$ gauge theory as an important concrete example. This theory is an example of nonabelian gauge theories arising from multiple *D*-branes.

We begin by introducing some general set-ups. Operators O acting on the vacuum correspond to states $|\Psi\rangle$ in the Hilbert space, i.e. $|\Psi\rangle = O|0\rangle$. The operators can be built by the fundamental fields in the theory. For example, in nonabelian gauge theories, we can have a complex matrix field Z, where we have some U(N) gauge group. This U(N) symmetry originates from the symmetry on multiple D-branes. The field quanta are created and annihilated by ladder operators. When acting on the operators, we have the following correspondence

$$Z \leftrightarrow a_Z^{\dagger}, \quad \partial_Z \leftrightarrow a_Z$$
$$Z_j^i = (Z)_j^i \leftrightarrow (a_Z^{\dagger})_j^i$$
$$(\partial_Z)_j^i = \frac{\partial}{\partial (Z)_j^j} \leftrightarrow (a_Z)_j^i \qquad (2.1)$$

The action $(a_Z)_j^i$, $(\partial_Z)_j^i$ is equivalent to Wick contraction with $(Z)_i^j$. More detailedly, the right hand side of the correspondence contains $\frac{1}{\sqrt{N}}$ norm factor for each elementary field, and hence $\operatorname{Tr}(a_Z^{\dagger n}) |0\rangle$ corresponds to $\frac{1}{\sqrt{nN^n}} \operatorname{Tr}(Z^n)$. This is exact for any finite and fixed N. The convention here is that we have the canonical commutation relations and normalizations for a_Z^{\dagger} , a_Z .

Similarly for other complex matrix fields Y, X in the U(N) gauge theory, one has the correspondence

$$\begin{array}{rcl}
Y & \leftrightarrow & a_Y^{\dagger}, & \partial_Y \leftrightarrow a_Y \\
X & \leftrightarrow & a_X^{\dagger}, & \partial_X \leftrightarrow a_X
\end{array}$$
(2.2)

In the similar way, $\operatorname{Tr}(a_Y^{\dagger n})$ corresponds to $\operatorname{Tr}(Y^n)$ and $\operatorname{Tr}(a_X^{\dagger n})$ corresponds to $\operatorname{Tr}(X^n)$.

The new class of the BPS coherent states is

$$F[\Lambda] = \frac{1}{Vol_U} \int dU \exp\left(\operatorname{Tr}(U\Lambda U^{-1}a_Z^{\dagger})\right) |0\rangle.$$
(2.3)

Here $U \in U(N)$ is an unitary action on the *N D*-branes. The action of the unitary *U*, originates from the nonabelian gauge symmetry of the *N* D-branes. The integral is over the group manifold, with the condition $\frac{1}{Vol_U} \int dU \cdot 1 = 1$. This defines a coherent state $|F[\Lambda]\rangle$. The conjugate bra state is $\langle F[\Lambda]|$ and is defined by conjugating a_Z^{\dagger} (respectively Λ) to a_Z (respectively $\overline{\Lambda}$).

We can also write the integral as $\int dg \exp(\operatorname{Tr}(\Lambda(g^{-1}a_Z^{\dagger}g))) |0\rangle$ and view g as auxiliary variables coupled to the a_Z^{\dagger} fields. We then integrate out these auxiliary variables. Here we interpret the integration as a *path integral* of the auxiliary variables. As a path integral, it can be performed by a saddle point method.

The inner products of the states are

$$\langle F[\Lambda']|F[\Lambda]\rangle = \bar{F}[\bar{\Lambda}'] * F[\Lambda],$$
 (2.4)

$$\bar{F}[\bar{\Lambda}'] * F[\Lambda] = \frac{1}{Vol} \int d\tilde{U} \exp(\operatorname{Tr}(\tilde{U}^{-1}\Lambda \tilde{U}\bar{\Lambda}')).$$
(2.5)

The normalizations of the states are $\mathcal{N}_{\Lambda} = \langle F[\Lambda] | F[\Lambda] \rangle$, where

$$\mathcal{N}_{\Lambda} = \frac{1}{Vol} \int dU \exp(\mathrm{Tr}(U^{-1}\Lambda U\bar{\Lambda})).$$
 (2.6)

For more details, see [16]. Eqn. (2.5),(2.6) are Harish-Chandra-Itzykson-Zuber (HCIZ) integrals [45, 46, 47], whose computation can be very conveniently performed by localizations [47] and by saddle point methods.

We can define an unitary displacement operator

$$D(\Lambda) = \frac{1}{Vol_U} \int dU \exp\left(\operatorname{Tr}\left(U\Lambda U^{-1}a_Z^{\dagger} - U\bar{\Lambda}U^{-1}a_Z\right)\right).$$
(2.7)

Using Baker-Campbell-Hausdorff (BCH) formulas and commutation relations of the ladder operators, we write

$$F[\Lambda] = \mathcal{N}_{\Lambda}^{\frac{1}{2}} D(\Lambda) |0\rangle \tag{2.8}$$

and we show (2.3) and (2.8) are equivalent. $D(\Lambda)$ is an unitary operation in $U(\mathcal{H})$ acting on the Hilbert space \mathcal{H} of states. In this writing, the term in the exponent is manifestly anti-Hermitian, and hence the operation D is an unitary operation. The advantage is that it is manifestly unitary on the Hilbert space. Using BCH formulas and commutation relations of the ladder operators, we write D also in the following way

$$D(\Lambda) = \frac{\mathcal{N}_{\Lambda}^{-\frac{1}{2}}}{Vol_U} \int dU \exp\left(\operatorname{Tr}\left(U\Lambda U^{-1}a_Z^{\dagger}\right)\right) \exp\left(-\operatorname{Tr}(U\bar{\Lambda}U^{-1}a_Z)\right).$$
(2.9)

Eqn. (2.7) and (2.9) are equivalent. The state $F(\Lambda)$ is an eigenstate of the annihilation operators and we have $\text{Tr}(a_Z^n)F(\Lambda) = \text{Tr}(\Lambda^n)F(\Lambda)$ for gauge invariant observables.

We can also define phase shift operator

$$R(\Theta_Z) = \frac{1}{Vol} \int dU \exp(\operatorname{Tr}(iU\Theta_Z U^{-1} a_Z^{\dagger} a_Z)).$$
(2.10)

Here Θ_Z is a phase matrix, whose eigenvalues $\theta_z^i, i = 1, ..., N$, are phases rotating the eigenvalues of Λ_Z . Here we let Λ_Z, Θ_Z commute, i.e. $[\Lambda_Z, \Theta_Z] = 0$. Hence

$$\frac{1}{Vol} \int dU \exp(\operatorname{Tr}(iU\Theta_Z U^{-1}a_Z^{\dagger}a_Z)) |F[\Lambda_Z]\rangle = \frac{1}{Vol} \int dU \exp(\operatorname{Tr}(U\Lambda_Z e^{i\Theta_Z} U^{-1}a_Z^{\dagger})) |0\rangle$$
(2.11)

where we used BCH formulas. The eigenvalues are rotated by $\lambda_z^i \to \lambda_z^i e^{i\theta_z^i}$, and correspondingly, $\bar{\lambda}_z^i \to \bar{\lambda}_z^i e^{-i\theta_z^i}$.

The number operator is $\hat{N}_Z = a_Z^{\dagger} a_Z$. We have $\hat{J}_Z = \text{Tr}(a_Z^{\dagger} a_Z)$ and $J_Z = \langle \hat{J}_Z \rangle$ measures the expectation value of the number of Z fields in the state. The Hamiltonian operator is $\hat{H}_0 = \text{Tr}(a_Z^{\dagger} a_Z)$ and $H_0 = \langle \hat{H}_0 \rangle$ counts the excitation energy. Our Hamiltonian has subtracted out the Casimir energy of the ground state. The Hamiltonian on the state space of the coherent states is

$$E_{F[\Lambda]} = \langle \hat{H} \rangle_{F[\Lambda]} = \mathcal{N}_F^{-1} \langle F(\Lambda) | \hat{H} | F(\Lambda) \rangle = \mathcal{N}_F^{-1} \bar{F}[\bar{\Lambda}] * \hat{H} * F[\Lambda].$$
(2.12)

One can use a more sophisticated Lagrangian formalism [16] to obtain an effective action on coherent states derived by [16]. These constructions and expressions also works for the other matrix fields Y, X in the U(N) gauge theory by using their ladder operators $a_Y^{\dagger}, a_X^{\dagger}$ for (2.3).

The coherent states for coulomb branches also work similarly. Consider coulomb branch gauge group $G_1 \times G_2$ inside full gauge group G. In that case, we need to embed $U(N_1) \times U(N_2)$ into U(N), where $N_1 + N_2 = N$, and the matrix Λ_Z splits into blocks e.g. $\Lambda_Z^{(1)}, \Lambda_Z^{(2)}$ corresponding to the two gauge groups, and $\operatorname{rk}(\Lambda_Z^{(1)}) + \operatorname{rk}(\Lambda_Z^{(2)}) = \operatorname{rk}(\Lambda_Z)$. Coulomb branch operators have been considered in [41] which is related to Gelfand-Tsetlin patterns [48]. The integration $\int_G dU$ turns into $\int_{G_1 \times G_2} \prod dU_i$, where

 $U_i \in U(N_i), i = 1, 2$. In this case, the permutation symmetry S_N also reduces to $S_{N_1} \times S_{N_2}$.

Ref. [16] has also constructed a new class of eighth BPS coherent states, with manifest permutation symmetries. It is constructed by enlarging $U\Lambda_Z U^{-1}a_Z^{\dagger}$ in (2.3) to more terms, adding additional parameters Λ_Y and Λ_X . We can take a limit case of the eighth BPS states constructed by [16], by making $\Lambda_X \equiv 0$. For $\Lambda_{(Z,Y)} \neq 0$, these are quarter BPS states. They are in the kernel of the anormalous dimension dilatation operator, and this means $[\Lambda_Z, \Lambda_Y] = 0$. This is a good advantage of using BPS coherent states.

These states are

$$F[\Lambda_Y, \Lambda_Z] = \frac{1}{Vol} \int dU \exp(\operatorname{Tr}(U\Lambda_Y U^{-1} a_Y^{\dagger} + U\Lambda_Z U^{-1} a_Z^{\dagger})) |0\rangle.$$
 (2.13)

Now first, in the following, we check that the states are quarter BPS at one loop and two loop orders. The one-loop dilatation operator and effective Hamiltonian is given by

$$\Delta_2 = g_{YM}^2 \operatorname{Tr}\left([a_Z^{\dagger}, a_Y^{\dagger}][a_Y, a_Z]\right).$$
(2.14)

As pointed out by [16], when we have the dilatation operator act on F, we get a result that is equal to zero when the parameters Λ_Y, Λ_Z are commuting matrices. The a_Y, a_Z appears on the rightmost. When acting on the above coherent states, we have the identification $a_Y \leftrightarrow \Lambda_Y, a_Z \leftrightarrow \Lambda_Z$, and $[a_Y, a_Z] \leftrightarrow [\Lambda_Y, \Lambda_Z] = 0$. Hence we see explicitly that the action of the one-loop dilatation operator on (2.13) is zero.

The two-loop dilatation operator is

$$\Delta_{4} = -\frac{g^{2}}{2} : \operatorname{Tr}\left(\left[\left[a_{Y}^{\dagger}, a_{Z}^{\dagger}\right], a_{Z}\right]\left[\left[a_{Y}, a_{Z}\right], a_{Z}^{\dagger}\right]\right) : \\ -\frac{g^{2}}{2} : \operatorname{Tr}\left(\left[\left[a_{Y}^{\dagger}, a_{Z}^{\dagger}\right], a_{Y}\right]\left[\left[a_{Y}, a_{Z}\right], a_{Y}^{\dagger}\right]\right) : \\ -\frac{g^{2}}{2} : \operatorname{Tr}\left(\left[\left[a_{Y}^{\dagger}, a_{Z}^{\dagger}\right], T^{a}\right]\left[\left[a_{Y}, a_{Z}\right], T^{a}\right]\right) :$$
(2.15)

Here $g = \frac{g_{YM}^2}{8\pi^2}$ with our convention. The terms in the dilatation operators are normal ordered. The normal ordering symbols here indicate that the annihilation operators within the normal ordering symbols do not act on fields inside the normal ordering. We computed the action of the two-loop dilatation operator on the above coherent states, its action on (2.13) is again zero, due to $[\Lambda_Z, \Lambda_Y] = 0$. Then, by using nonrenormalization theorems, e.g. [49, 23], we are convinced that they are also higher-loop BPS.

The state is a generating function of single-trace and multi-trace states of the form $sTr(Y^{m_1}Z^{n_1}Y^{m_2}Z^{n_2}...Y^{m_l}Z^{n_l}...)$. The generated states are quarter BPS single-trace and multi-trace operators built by Z, Y. They have been investigated in e.g. [49]. The two matrices Z, Y correspond to C^2 of the transverse dimensions of N D-branes.

The unitary displacement operator is

$$D(\Lambda_Y, \Lambda_Z) = \frac{1}{Vol_U} \int dU \exp(\operatorname{Tr}(U\Lambda_Z U^{-1}a_Z^{\dagger} + U\Lambda_Y U^{-1}a_Y^{\dagger}) - \operatorname{Tr}(U\bar{\Lambda}_Z U^{-1}a_Z + U\bar{\Lambda}_Y U^{-1}a_Y))$$
(2.16)

and we write $F = \mathcal{N}_F^{\frac{1}{2}} D(\Lambda_Y, \Lambda_Z) |0\rangle$. It is equivalent to

$$D(\Lambda_Y, \Lambda_Z) = \frac{\mathcal{N}_F^{-\frac{1}{2}}}{Vol_U} \int dU \exp(\operatorname{Tr}(U\Lambda U^{-1}a_Z^{\dagger} + U\Lambda_Y U^{-1}a_Y^{\dagger})) \exp(-\operatorname{Tr}(U\bar{\Lambda}U^{-1}a_Z + U\bar{\Lambda}_Y U^{-1}a_Y)).$$
(2.17)

The phase shift operator is

$$R(\Theta_Y) = \frac{1}{Vol} \int dU \exp(\operatorname{Tr}(iU\Theta_Y U^{-1}a_Y^{\dagger}a_Y)).$$
(2.18)

Here we let Λ_Y, Θ_Y commute. The action of it rotates $\lambda_y^i \to \lambda_y^i e^{i\theta_y^i}$.

The Hamiltonian in the state space of coherent states is $\hat{H}_0 = \text{Tr}(a_Y^{\dagger}a_Y + a_Z^{\dagger}a_Z)$. This counts the BPS energy $H_0 = \langle \hat{H}_0 \rangle$. On the other hand, $\hat{J}_Y = \text{Tr}(a_Y^{\dagger}a_Y)$ and $J_Y = \langle \hat{J}_Y \rangle$ measures the expectation value of the number of Y fields in the state.

Now we perform contracting the ladder operators and the inner product is

$$\mathcal{N}_F = \bar{F}[\bar{\Lambda}_Z, \bar{\Lambda}_Y] * F[\Lambda_Z, \Lambda_Y] = I(U, \Lambda_\alpha, \bar{\Lambda}_\alpha)$$
(2.19)

where $\alpha = Z, Y$. The integral is

$$I(U, \Lambda_{\alpha}, \bar{\Lambda}_{\alpha}) = \frac{1}{Vol} \int dU \exp\left(\Gamma_{\text{HCIZ}}\left(U, \Lambda_{\alpha}, \bar{\Lambda}_{\alpha}\right)\right).$$
(2.20)

We denote the exponent in (2.20) Γ_{HCIZ} . The exponent is

$$\Gamma_{\text{HCIZ}}\left(U,\Lambda_{\alpha},\bar{\Lambda}_{\alpha}\right) = \text{Tr}(U\Lambda_{Y}U^{-1}\bar{\Lambda}_{Y} + U\Lambda_{Z}U^{-1}\bar{\Lambda}_{Z}).$$
(2.21)

The path integral (2.20) can be computed by a saddle point method. The saddle point equation is $d\Gamma_{\text{HCIZ}} = 0$. Now we write $dK = UdU^{-1}$, and we have

$$d\Gamma_{\text{HCIZ}} = \operatorname{Tr}\left(dK[\bar{\Lambda}_Y, U\Lambda_Y U^{-1}]\right) + \operatorname{Tr}\left(dK[\bar{\Lambda}_Z, U\Lambda_Z U^{-1}]\right) = 0.$$
(2.22)

The conditions for saddle points are

$$[\bar{\Lambda}_Y, U\Lambda_Y U^{-1}] + [\bar{\Lambda}_Z, U\Lambda_Z U^{-1}] = 0.$$
(2.23)

When U is a permutation matrix $P \in S_N$, these are saddle points. The group integral can be viewed as a path integral of the auxiliary variables U. Then the integral (2.20) can be computed by localization and saddle point method as described in [16].

We now turn to the eighth BPS case in more details. The states are

$$F[\Lambda_Z, \Lambda_X, \Lambda_Y] = \frac{1}{Vol} \int dU \exp\left(\operatorname{Tr}(U\Lambda_X U^{-1}a_X^{\dagger} + U\Lambda_Y U^{-1}a_Y^{\dagger} + U\Lambda_Z U^{-1}a_Z^{\dagger})\right) |0\rangle.$$
(2.24)

It can also be written as

$$F[\vec{\Lambda}] = \frac{1}{Vol} \int dU \exp\left(\operatorname{Tr}(U\vec{\Lambda}U^{-1}\vec{a}^{\dagger})\right) |0\rangle, \qquad (2.25)$$

where $\vec{\Lambda} = (\Lambda_X, \Lambda_Y, \Lambda_Z)$, $\vec{a}^{\dagger} = (a_X^{\dagger}, a_Y^{\dagger}, a_Z^{\dagger})$. This form of writing is also convenient for theories with global symmetries or flavor symmetries. As pointed out in [16], if the parameters $\Lambda_X, \Lambda_Y, \Lambda_Z$ mutually commute, the states are annihilated by the one-loop dilatation operator. The effective Hamiltonian is given by

$$\Delta = g_{YM}^2 \operatorname{Tr}([a_X^{\dagger}, a_Z^{\dagger}][a_Z, a_X]) + g_{YM}^2 \operatorname{Tr}[a_Y^{\dagger}, a_Z^{\dagger}][a_Z, a_Y]) + g_{YM}^2 \operatorname{Tr}([a_X^{\dagger}, a_Y^{\dagger}][a_Y, a_X]).$$
(2.26)

When acting on the coherent states, we have the correspondence $a_X \leftrightarrow \Lambda_X$, $[a_X, a_Z] \leftrightarrow [\Lambda_X, \Lambda_Z]$, $[a_X, a_Y] \leftrightarrow [\Lambda_X, \Lambda_Y]$, etc. Hence all the three terms acting on the states are zero, and the action of the dilatation is zero. Hence $\Lambda_X, \Lambda_Y, \Lambda_Z$ are required to commute pairwise. By using nonrenormalization theorems [49, 23], we can infer that they are in the kernel of the anormalous dilatation operator.

The unitary displacement operator is similar to the above (2.16) with more terms in the exponent added. And we assume $\Lambda_{(X,Y,Z)}$ are mutually commuting. By using BCH formulas, it is equivalent to the following, with the normalization factor,

$$D(\Lambda_X, \Lambda_Y, \Lambda_Z) = \frac{\mathcal{N}_F^{-\frac{1}{2}}}{Vol_U} \int dU \exp\left(\operatorname{Tr}(U\Lambda_Z U^{-1}a_Z^{\dagger} + U\Lambda_X U^{-1}a_X^{\dagger} + U\Lambda_Y U^{-1}a_Y^{\dagger})\right) \\ \exp\left(-\operatorname{Tr}(U\bar{\Lambda}_Z U^{-1}a_Z + U\bar{\Lambda}_X U^{-1}a_X + U\bar{\Lambda}_Y U^{-1}a_Y)\right).$$
(2.27)

We write $F = \mathcal{N}_F^{\frac{1}{2}} D(\Lambda_X, \Lambda_Y, \Lambda_Z) |0\rangle$. Similarly we define the phase shift operator $R(\Theta_X)$ similar to (2.10) and (2.18).

By performing usual manipulations contracting ladder operators, the overlap \mathcal{N}_F was computed in [16]. The overlap is a HCIZ integral, where the exponent is similar to (2.21) and have three terms, with $\Gamma_{\text{HCIZ}}(U, \Lambda_{\alpha}, \bar{\Lambda}_{\alpha}) = \text{Tr}(U\vec{\Lambda}U^{-1}\vec{\Lambda})$. The conditions for saddle points are in [16]. It is in a similar form as (2.23) with three terms. The integrals can be efficiently calculated by localization and saddle point method.

The Hamiltonian in the space of coherent states is $\hat{H}_0 = \text{Tr}(a_Z^{\dagger}a_Z + a_Y^{\dagger}a_Y + a_X^{\dagger}a_X)$, with Casimir energy subtracted. The energy in the space of coherent states is

$$E_F = \langle \hat{H} \rangle_F := \mathcal{N}_F^{-1} \langle F[\Lambda_{(X,Y,Z)}] | \hat{H} | F[\Lambda_{(X,Y,Z)}] \rangle = \mathcal{N}_F^{-1} \ \bar{F} * H * F.$$
(2.28)

The angular momentum operator is $\hat{J}_X = \text{Tr}(a_X^{\dagger}a_X)$ and $J_X = \langle \hat{J}_X \rangle$ measures the expectation value of the number of X fields in the state. We have that $J_Z = \text{Tr}(\bar{\Lambda}_Z \Lambda_Z)$, $J_Y = \text{Tr}(\bar{\Lambda}_Y \Lambda_Y)$, $J_X = \text{Tr}(\bar{\Lambda}_X \Lambda_X)$, and

$$H_0 = J_X + J_Y + J_Z. (2.29)$$

The vevs are

$$\langle \operatorname{Tr}(a_Z^{\dagger n} a_Z^n) \rangle_F = \langle \operatorname{Tr}(\bar{\Lambda}_Z \Lambda_Z)^n \rangle_F, \quad \langle \operatorname{Tr}(a_Y^{\dagger n} a_Y^n) \rangle_F = \langle \operatorname{Tr}(\bar{\Lambda}_Y \Lambda_Y)^n \rangle_F, \langle \operatorname{Tr}(a_X^{\dagger n} a_X^n) \rangle_F = \langle \operatorname{Tr}(\bar{\Lambda}_X \Lambda_X)^n \rangle_F.$$

$$(2.30)$$

Note that e.g. $(\bar{\Lambda}_Z \Lambda_Z)^n$ is conveniently a Hermitian matrix with real eigenvalues. We then look at the eigenvalue of the parameters $\lambda_i^x, \lambda_i^y, \lambda_i^z$. In the large N limit, $\rho(\lambda) = \rho(\lambda^{(x,y,z)})$ is an eigenvalue density. The coherent state $|F[\Lambda_{(X,Y,Z)}]\rangle$ may also be labeled as $|\rho(\lambda^{(x,y,z)})\rangle$ in the large N limit. We can also calculate the right hand sides of (2.30) for the case when n is not an integer, using eigenvalue formalism, e.g.

$$\langle \operatorname{Tr}(\bar{\Lambda}_Z \Lambda_Z)^n \rangle_F = \int d^6 \lambda \rho(\lambda) (\bar{\lambda}^z \lambda^z)^n.$$
 (2.31)

The methods developed in the approach of matrix eigenvalue effective models are very useful in this context [8],[50]-[54]. These vevs are dual to the multi-pole moments of the gravitational geometries in the gravity side, and can be measured in the gravity side via the methods of [55]-[61]. The n = 1 case of the similar quantities were analyzed in [60].

These states are heavy excited states in the gravity side. The particularly interesting regimes are J_X, J_Y, J_Z of order N^2 where there is backreaction of the excitation onto the spacetime geometry. In these regimes, single-trace and multi-traces are more difficult and less convenient. The coherent states are especially convenient in their large amplitude regimes. The gravity waves are collective excitations of these large N eigenvalue distributions. The order N regimes describe multi giant gravitons. The individual eigenvalue of the coherent state multi-parameter is the collective coordinate of giant gravitons on the gravity side [16], see also related observations [18, 37].

This new type of coherent states have been constructed by [16, 17]. The BPS coherent states are generating functions of single-trace and multi-trace states. The multi-trace states are multi-closed string states. Hence the BPS coherent states are also generating functions of multi-closed string states. It is possible to generalize to more matrices and more flavors, in other circumstances or in other types of gauge theories. A special class of quarter BPS coherent states were also constructed in [62], which has not made use of the manifest permutation symmetries.

3 The SL(2) sectors, their cousins, and meromorphic versions

We construct a new type of coherent states in the SL(2) sectors and their cousin PSU(1,1|2) sectors. We also construct a meromorphic version of coherent states, which can be transformed back to the full coherent states when det $\Lambda_Z \neq 0$. We give transformation relations between different types of coherent states, among other things.

Inspired by the coherent states in [16], whose aspects are analyzed in more details in Sec. 2, we construct a coherent state as generating function of

$$Tr(D_{+}^{n_{1}}ZD_{+}^{n_{2}}Z...D_{+}^{n_{\alpha}}Z...),$$
(3.1)

 $n_{\alpha} = 0, 1, 2, ...$ and so on. Here D_{+} is a light-cone covariant derivative operator. We write $D_{+} = n^{\mu}D_{\mu}$ where n^{μ} is a light-like vector. D_{+} is in the (1,0) representation of the local Lorentz group. When D_{+} are dilute, they can be viewed as impurities on top of Z. Similar conceptions are also raised in [26, 63]; see also [64]. For more details on SL(2) sectors and their relations to integrability and QCD, see for example [4, 65, 66].

We can construct the ladder operators corresponding to adding derivatives n times on top of Z,

$$c_D^{\dagger n} a_Z^{\dagger} \leftrightarrow D_+^n Z, \quad c_D^{\dagger} a_Z^{\dagger} \leftrightarrow D_+ Z, \quad a_Z^{\dagger} \leftrightarrow Z.$$
 (3.2)

We construct a type of coherent states as the generating function of (3.1),

$$K[R_Z, \Lambda_Z] = \frac{1}{Vol} \int dU \exp(\operatorname{Tr}(UR_Z U^{-1} c_D^{\dagger} a_Z^{\dagger} a_Z)) \exp(\operatorname{Tr}(U\Lambda_Z U^{-1} a_Z^{\dagger})) |0\rangle.$$
(3.3)

Here R_Z, Λ_Z commute. We make sure that before acting c_D^{\dagger} , there is already a background of Z-fields due to the right-most operator in (3.3). The angular momentum operator and spin operator are $\hat{J}_Z = \text{Tr}(a_Z^{\dagger}a_Z)$ and $\hat{S} = \text{Tr}(c_D^{\dagger}c_D)$. Here $S = \langle \hat{S} \rangle$ measures the expectation value of the number of D_+ . On this state (3.3), the expectation values are $J_Z = \langle \hat{J}_Z \rangle = \text{Tr}(\bar{\Lambda}_Z \Lambda_Z)$ and $S = \langle \hat{S} \rangle = \text{Tr}(R_Z \bar{R}_Z \bar{\Lambda}_Z^2 \Lambda_Z^2)$.

An alternative formulation is the state

$$G[T_{DZ}, \Lambda_Z] = \frac{1}{Vol} \int dU \exp(\operatorname{Tr}(UT_{DZ}U^{-1}c_D^{\dagger}a_Z) \exp(\operatorname{Tr}(U\Lambda_Z U^{-1}a_Z^{\dagger})) |0\rangle.$$
(3.4)

Here T_{DZ} , Λ_Z commute. We have that $J_Z = \text{Tr}(\bar{\Lambda}_Z \Lambda_Z)$ and $S = \text{Tr}(T_{DZ} \bar{T}_{DZ} \bar{\Lambda}_Z \Lambda_Z)$.

These are heavy states in the gravity side. If the D_+ is very dilute, it is the small spin limit. This is when $\alpha = N^{-1} \text{Tr}(T_{DZ} \overline{T}_{DZ})$ is much smaller than one. On the other hand, if the Z is dilute and D_+ is dense, it is the large spin limit. This is when α is much larger than one. The large spin limit has been discussed also in [26].

States of this type (3.3) can also be constructed for the operators analyzed in sections 2. We construct the following state

$$G[T_Y, \Lambda_Z] = \frac{1}{Vol} \int dU \exp(\operatorname{Tr}(UT_Y U^{-1} a_Y^{\dagger} a_Z)) \exp(\operatorname{Tr}(U\Lambda_Z U^{-1} a_Z^{\dagger})) |0\rangle.$$
(3.5)

Here T_Y, Λ_Z commute. The right-most operator in (3.5) is an eigenstate of a_Z . When acting on the coherent state background, we replace the annihilation operator by the vev. Here we replace the a_Z by vev Λ_Z , when the left operator in (3.5) acts on the right-most operator. The action $a_Y^{\dagger}a_Z$ annihilates a Z field and substitute it with a Y field, e.g. $Z^2 \to YZ$. Hence we have $J_Z = \text{Tr}(\bar{\Lambda}_Z \Lambda_Z)$ and $J_Y = \text{Tr}(T_Y \bar{T}_Y \bar{\Lambda}_Z \Lambda_Z)$.

The state (3.5) is a meromorphic version of (2.13). $G[T_Y, \Lambda_Z]$ is the excitation on top of $F[\Lambda_Z]$. It is a good approximation to $F[\Lambda_Y, \Lambda_Z]$, in the regime det $\Lambda_Z \neq 0$. We see that

$$G[T_Y, \Lambda_Z]|_{T_Y = \Lambda_Z^{-1} \Lambda_Y} \sim F[\Lambda_Y, \Lambda_Z],$$
(3.6)

when det $\Lambda_Z \neq 0$ and identifying $T_Y = \Lambda_Z^{-1} \Lambda_Y$. When det $\Lambda_Z = 0$, $G[T_Y, \Lambda_Z]$ can not be transformed back to $F[\Lambda_Y, \Lambda_Z]$, since there is pole in T_Y . Here T_Y is a meromorphic function of Λ_Z and not a holomorphic function, hence there is pole in T_Y . This is when det $\Lambda_Z = 0$. $F[\Lambda_Y, \Lambda_Z]$ is a smooth resolution of $G[T_Y, \Lambda_Z]$, extending it to cover the pole locus det $\Lambda_Z = 0$.

The meromorphic version provides new understandings of the coherent states and new understanding to the important question why we have the background fields Z. These background fields, e.g. Z fields, serve as new effective vacuum for new excitations. Similarly, for including X, we have the state

$$G[T_X, T_Y, \Lambda_Z] = \frac{1}{Vol} \int dU \exp(\operatorname{Tr}(UT_X U^{-1} a_X^{\dagger} a_Z + UT_Y U^{-1} a_Y^{\dagger} a_Z) \exp(\operatorname{Tr}(U\Lambda_Z U^{-1} a_Z^{\dagger})) |0\rangle.$$
(3.7)

Here T_X, T_Y, Λ_Z mutually commute. We have that $J_Z = \text{Tr}(\bar{\Lambda}_Z \Lambda_Z), J_Y = \text{Tr}(T_Y \bar{T}_Y \bar{\Lambda}_Z \Lambda_Z)$, and $J_X = \text{Tr}(T_X \bar{T}_X \bar{\Lambda}_Z \Lambda_Z)$.

In the regime det $\Lambda_Z \neq 0$,

$$G[T_X, T_Y, \Lambda_Z] \sim F[\Lambda_X, \Lambda_Y, \Lambda_Z], \tag{3.8}$$

with $T_X = \Lambda_Z^{-1} \Lambda_X, T_Y = \Lambda_Z^{-1} \Lambda_Y$. Hence state (3.7) is a meromorphic version of (2.24). $G[T_Y, \Lambda_Z], G[T_X, T_Y, \Lambda_Z]$ can be viewed as meromorphic versions of quarter and eighth BPS coherent states.

Here, it is very similar to having U(2) and U(3) flavor symmetry, or more generally $U(N_F)$ symmetry if there are N_F sets of fields with global symmetry. Similar expansions also works for quiver gauge theory with global symmetry, as well as their Coulomb branches. On the other hands, there are theories with these flavor symmetries.

Turning to the SL(2) sector, this sector can be enlarged to PSU(1, 1|2) sector. The PSU(1, 1|2) sector has been discussed from the point of view of integrability, e.g. [67]. Now we consider bosonic part of PSU(1, 1|2) sector, which is composed of Z, Y, D_+ and two fermions in the $(\frac{1}{2}, -\frac{1}{2})$ and $(\frac{1}{2}, \frac{1}{2})$ of the local Lorentz group. In the PSU(1, 1|2) sector, the bosonic part of the state is

$$G[T_{DZ}, T_{DY}, \Lambda_Z, \Lambda_Y] = \frac{1}{Vol} \int dU \exp(\operatorname{Tr}(UT_{DZ}U^{-1}c_D^{\dagger}a_Z) + \operatorname{Tr}(UT_{DY}U^{-1}c_D^{\dagger}a_Y)) \exp(\operatorname{Tr}(U\Lambda_Z U^{-1}a_Z^{\dagger} + U\Lambda_Y U^{-1}a_Y^{\dagger}) |0\rangle.$$
(3.9)

Here $T_{DZ}, T_{DY}, \Lambda_Z, \Lambda_Y$ mutually commute. The angular momentum and spin of the state is $J_Z = \text{Tr}(\bar{\Lambda}_Z \Lambda_Z), J_Y = \text{Tr}(\bar{\Lambda}_Y \Lambda_Y)$ and $S = \text{Tr}(|T_{DZ} \Lambda_Z + T_{DY} \Lambda_Y|^2)$.

An alternative formulation in the bosonic part of the PSU(1,1|2) sector, is

$$K[R_Z, R_Y, \Lambda_Z, \Lambda_Y] = \frac{1}{Vol} \int dU \exp(\operatorname{Tr}(UR_Z U^{-1} c_D^{\dagger} a_Z^{\dagger} a_Z) + \operatorname{Tr}(UR_Y U^{-1} c_D^{\dagger} a_Y^{\dagger} a_Y)) \exp(\operatorname{Tr}(U\Lambda_Z U^{-1} a_Z^{\dagger} + U\Lambda_Y U^{-1} a_Y^{\dagger}) |0\rangle.$$
(3.10)

Here $R_Z, \Lambda_Z, R_Y, \Lambda_Y$ mutually commute. We see that $J_Z = \text{Tr}(\bar{\Lambda}_Z \Lambda_Z), J_Y = \text{Tr}(\bar{\Lambda}_Y \Lambda_Y)$ and $S = \text{Tr}(|R_Z \bar{\Lambda}_Z \Lambda_Z + R_Y \bar{\Lambda}_Y \Lambda_Y|^2)$. Now we turn to the quiver case. For quiver gauge theories and orbifold daughters of $\mathcal{N} = 4$ theory, the coherent state construction is similar. Many orbifold daughters have integrability. Many aspects of them have been discussed and worked out in e.g. [68]-[74] and their related references. For example, we can make a projection of U(MN) theory to get the $U(N)^M$ theory. The construction of coherent states for quiver gauge theories have been worked out in [16]. For example, we have a $U(N_1) \times U(N_2)$ gauge group. We consider a pair of bifundamental fields $a_{12}^{\dagger}, a_{21}^{\dagger}$ in the (\bar{N}_1, N_2) and the (N_2, \bar{N}_1) representations. We make the same type of coherent states with their parameters $\Lambda_{21}, \Lambda_{12}$, whose roles are similar to Λ_Z, Λ_Y . The state is

$$F[\Lambda_{21}, \Lambda_{12}] = \frac{1}{Vol} \int \prod_{i=1}^{2} dU_i \exp\left(\operatorname{Tr}(\Lambda_{21}U_1 a_{12}^{\dagger} U_2^{-1} + \Lambda_{12}U_2 a_{21}^{\dagger} U_1^{-1}) |0\rangle.$$
(3.11)

Note that the conjugate of a_{21}^{\dagger} is a_{12} , and the conjugate of a_{12}^{\dagger} is a_{21} , with our convention. Now we construct a new state, which is the meromorphic version of (3.11),

$$G[T_{21}, \Lambda_{12}] = \frac{1}{Vol} \int \prod_{i=1}^{2} dU_i \exp(\operatorname{Tr}(U_2^{-1}T_{21}U_1a_{12}^{\dagger}a_{12})) \exp(\operatorname{Tr}(U_1^{-1}\Lambda_{12}U_2a_{21}^{\dagger})) |0\rangle.$$
(3.12)

The right-most operator in (3.12) is an eigenstate of a_{12} . We see that

$$G[T_{21}, \Lambda_{12}] \sim F[\Lambda_{21}, \Lambda_{12}],$$
 (3.13)

with $\Lambda_{12}T_{21} = \Lambda_{21}$, $T_{21} = \Lambda_{12}^{-1}\Lambda_{21}$. $G[T_{21}, \Lambda_{12}]$ is a meromorphic version of $F[\Lambda_{21}, \Lambda_{12}]$ and can be transformed to it when $N_1 = N_2$ and det $\Lambda_{12} \neq 0$. The inner products of (3.11) have been computed in [16], where localization techniques have been implemented.

4 String-added coherent states

The BPS coherent state can serve as the supersymmetric background for other excitations on top of it. We can consider further excitations on the BPS coherent state background, by adding strings on top of them. One important idea is splitting and identification of fields for the background and fields for the impurities. We begin with some general settings.

4.1 General settings with multi-words

The background BPS coherent state is (2.24). On this background, other fields are excited. This is to add words or multi-words onto the coherent state operator. These

other excited fields may be viewed as impurities, if their numbers are much fewer than the number of background fields. We generalize the ideas of [16] from single word to multi-words, and from single pair to multi-pairs.

The BPS coherent states are parametrized by $\vec{\Lambda} = (\Lambda_X, \Lambda_Y, \Lambda_Z)$. Consider $|v_x\rangle_i$, $|v_y\rangle_i$, $|v_z\rangle_i$ are eigenvectors, i.e.

$$\Lambda_X |v_x\rangle_i = \lambda_x^i |v_x\rangle_i, \quad \Lambda_Y |v_y\rangle_i = \lambda_y^i |v_y\rangle_i, \quad \Lambda_Z |v_z\rangle_i = \lambda_z^i |v_z\rangle_i.$$
(4.1)

When we multiply as $U\Lambda_X U^{-1}$, $U\Lambda_Y U^{-1}$, $U\Lambda_Z U^{-1}$, the vectors are rotated to $U |v_x\rangle_i$, $U |v_y\rangle_i$, $U |v_z\rangle_i$. The bra states are rotated to $\langle v_x|_i U^{-1}$, etc. The vectors are simultaneously rotated. There is also a global symmetry U(3) acting on the set of 3 vectors.

The analysis of [16] as well as the above analysis in Sec. 2 imply that $\Lambda_{(X,Y,Z)}$ are mutually commuting for BPS. Now we assume they mutually commute. Hence they can share the same eigenvectors. We consider there are two large eigenvalues $\lambda_{(X,Y,Z)}^i$ and $\lambda_{(X,Y,Z)}^j$. They share the same eigenvectors,

$$\Lambda_X \left| \vec{v} \right\rangle_{\alpha} = \lambda_x^{\alpha} \left| \vec{v} \right\rangle_{\alpha}, \quad \Lambda_Y \left| \vec{v} \right\rangle_{\alpha} = \lambda_y^{\alpha} \left| \vec{v} \right\rangle_{\alpha}, \quad \Lambda_Z \left| \vec{v} \right\rangle_{\alpha} = \lambda_z^{\alpha} \left| \vec{v} \right\rangle_{\alpha},$$

$$\text{for } \alpha = i, j.$$

$$(4.2)$$

Here only for two eigenvalues i, j having this property are needed. Hence $|v_x\rangle_{\alpha} = |v_y\rangle_{\alpha} = |v_z\rangle_{\alpha} = |\vec{v}\rangle_{\alpha}$, for $\alpha = i, j$. This means that in the two dimensional subspace involving $i, j, \Lambda_{(X,Y,Z)}$ share the same eigenvector $|\vec{v}\rangle_{i,j}$.

The added words on top is

$$\left\langle \vec{v} \right|_{i} U^{-1} W U \left| \vec{v} \right\rangle_{i} \quad (4.3)$$

and the state is

$$\frac{1}{Vol} \int dU \exp\left(\operatorname{Tr}(U\Lambda_X U^{-1}a_X^{\dagger} + U\Lambda_Y U^{-1}a_Y^{\dagger} + U\Lambda_Z U^{-1}a_Z^{\dagger})\right) \langle \vec{v}|_i U^{-1}WU | \vec{v} \rangle_j | 0 \rangle.$$
(4.4)

More generally, we add multi-words of this kind,

$$\frac{1}{Vol} \int dU \exp\left(\operatorname{Tr}(U\Lambda_X U^{-1}a_X^{\dagger} + U\Lambda_Y U^{-1}a_Y^{\dagger} + U\Lambda_Z U^{-1}a_Z^{\dagger})\right) \cdot \langle \vec{v}|_{i_1} U^{-1} W_1 U |\vec{v}\rangle_{j_1} \langle \vec{v}|_{i_2} U^{-1} W_2 U |\vec{v}\rangle_{j_2} \dots \langle \vec{v}|_{i_l} U^{-1} W_l U |\vec{v}\rangle_{j_l} |0\rangle.$$
(4.5)

The indices $(i_l, j_l) \in I_s$ form a set I_s . W is a word, in other words, a string. For example, W is schematically $D_+^{\dagger n_1} a_Z^{\dagger n_2} a_Y^{\dagger n_3} a_X^{\dagger n_4}$, with $n_{\alpha} = 0, 1, 2, ...$, and so on. W can be labelled by BMN operators or spin chain like operators, and we can also include $a_{\overline{Z}}^{\dagger}$ etc. The integral (4.5) can be viewed as a path integral of the auxiliary variable U which is coupled to $a_Z^{\dagger}, a_Y^{\dagger}, a_X^{\dagger}$ and W_{α} , in accord with the alternative interpretation in Sec 2. The insertions of W_{α} can be viewed as operator-insertion in this path integral (4.5). We have generalized the word insertions in (4.4) to multi-word insertions, and interpreted them as operator insertions in the path integral of auxiliary variables.

We extract the anormalous energy from the combined state, e.g. $F \cdot W$. The background energy are energy of BPS states. We extract the energy of non-BPS part of the excitation, corresponding to the anormalous dimension. The anomalous dimensions give rise to non-BPS energy on top of BPS energy. We works on the anormalous energy due to the quartic interaction vertices. The quartic interaction vertex connects one impurity field on the word part and one background field on the coherent state part, and they then Wick contract to their conjugates respectively. We can also use the dilatation operator.

The simplest cases are when the coherent state part F and the word part W are both BPS. The coherent state background F is BPS, hence there is no non-BPS energy from the interaction inside the coherent state part. Then the anomalous dimension energy is solely due to the interaction between the coherent state part and the word part. One of the simple cases is that W is an eighth BPS string. The entire state can be maintained as a near BPS state. Sec. 4.2 are special cases of these ideas.

4.2 Near BPS string-added coherent states

We first consider near BPS states of the form

$$\frac{1}{Vol} \int dU \exp\left(\operatorname{Tr}(U\Lambda_Z U^{-1}a_Z^{\dagger})\right) \cdot \langle \vec{v}|_{i_1} U^{-1} W_1 U | \vec{v} \rangle_{j_1} \langle \vec{v}|_{i_2} U^{-1} W_2 U | \vec{v} \rangle_{j_2} \dots \langle \vec{v}|_{i_l} U^{-1} W_l U | \vec{v} \rangle_{j_l} | 0 \rangle.$$
(4.6)

The entire state can be maintained as a near BPS state. For example we add impurities corresponding to X, and the word W is $a_X^{\dagger} a_Z^{\dagger m}$, m = 0, 1, ... and so on. In this case, W is BPS itself, and it can be viewed as a symmetrized state in the sector of Z, X. The anormalous energy is the energy corresponding to the anormalous dimension. Because W and F both has no anormalous dimension under the dilatation operator, the anormalous dimension energy is due to the interaction between the coherent state F part and the word W part.

We then consider near BPS states of the form

$$\frac{1}{Vol} \int dU \exp\left(\operatorname{Tr}(U\Lambda_Z U^{-1}a_Z^{\dagger} + U\Lambda_Y U^{-1}a_Y^{\dagger})\right) \cdot \langle \vec{v}|_{i_1} U^{-1} W_1 U |\vec{v}\rangle_{j_1} \langle \vec{v}|_{i_2} U^{-1} W_2 U |\vec{v}\rangle_{j_2} \dots \langle \vec{v}|_{i_l} U^{-1} W_l U |\vec{v}\rangle_{j_l} |0\rangle.$$
(4.7)

The word W is $a_X^{\dagger} \{a_Z^{\dagger} a_Y^{\dagger} a_Y^{\dagger} \}$, $m_{\alpha} = 0, 1, ...$ and so on. Here the curly bracket denotes symmetrized states in the sector of Z, Y, X with U(3) global symmetry, where the bracket denotes symmetrization over the global symmetry indices. See e.g. [49, 23]. The symmetrized states itself is BPS. We make both the coherent state F part and the word W part BPS, hence the non-BPS energy is coming from the interaction between the coherent state part and the word part. The above first example is a special case of the second example for $m_2 = 0$.

In the following, we compute for the case (4.7). The quartic vertices that are relevant here are $\text{Tr}|[Y,X]|^2$, $\text{Tr}|[Z,X]|^2$, $\text{Tr}|[Y,Z]|^2$. The anormalous dimension is due to the interaction of impurity a_X^{\dagger} with the a_Z^{\dagger} and a_Y^{\dagger} respectively, on the exponent. The interaction between a_Z^{\dagger} and a_Y^{\dagger} gives no anormalous dimension, due to that they are in the exponent of the BPS coherent state, as calculated in Sec 2. The word W is by itself BPS because it is symmetrized for a_Z^{\dagger} , a_Y^{\dagger} , a_X^{\dagger} . Hence the interaction between Z, Y in this case does not lead to anormalous energy. The Z, Y composite here are BPS.

The Hamiltonian involved for the excitation energy is

$$\Delta = g^2 \operatorname{Tr}([a_Y^{\dagger}, a_X^{\dagger}][a_X, a_Y]) + g^2 \operatorname{Tr}([a_Z^{\dagger}, a_X^{\dagger}][a_X, a_Z]).$$
(4.8)

Here, the interaction leading to the anormalous energy here are quartic vertices between the X in the word part and the Z and Y, respectively in the coherent state part. For example, X in the word part is attached to a quartic interaction vertex with a Z in the coherent state F part. And after interaction, they Wick contract to the conjugate of the word part and the conjugate of the coherent state part respectively. The coherent state part itself is BPS as explained.

The method of calculation follows from [16]. One can directly work with the quartic vertices as in BMN [75], or use the dilatation operator [76]. Here, $|v_y\rangle_{\alpha} = |v_z\rangle_{\alpha}$ for $\alpha = i, j$, as explained in Sec. 4.1, and we denote the pair of large eigenvalues $\vec{\lambda}_i, \vec{\lambda}_j$. When acting on the state, the extra annihilation operator a_Y brings down $U\Lambda_Y U^{-1}$, one to the left and the other to the right. These two pieces go as

$$g^{2} \langle v_{y}|_{i} U^{-1} U \Lambda_{Y} U^{-1}[a_{X}^{\dagger}, a_{Y}^{\dagger}] U | v_{y} \rangle_{j} - g^{2} \langle v_{y}|_{i} U^{-1}([a_{X}^{\dagger}, a_{Y}^{\dagger}] U \Lambda_{Y} U^{-1} U | v_{y} \rangle_{j}.$$
(4.9)

Since $|v_y\rangle_i$, $|v_y\rangle_i$ are eigenstates of Λ_Y , we get

$$g^{2}(\lambda_{i}^{y} - \lambda_{j}^{y})\langle v_{y}|_{i}U^{-1}[a_{X}^{\dagger}, a_{Y}^{\dagger}]U | v_{y} \rangle_{j}.$$
(4.10)

We now proceed the process for computing with the conjugate vector, and a_Y^{\dagger} brings down terms of $\bar{\Lambda}_Y$. We obtain an integral that involves \tilde{U}^{-1} , so we get a factor $(\bar{\lambda}_i^y - \bar{\lambda}_i^y)$ from (4.10). For a_Z , the calculation is similar, and we get

$$g^{2}(\lambda_{i}^{z}-\lambda_{j}^{z})\langle v_{z}|_{i}U^{-1}[a_{X}^{\dagger},a_{Z}^{\dagger}]U|v_{z}\rangle_{j}.$$
(4.11)

For the conjugate vector, a_Z^{\dagger} brings down terms of $\bar{\Lambda}_Z$. Again, we have the integral and get a factor $(\bar{\lambda}_i^z - \bar{\lambda}_j^z)$ from (4.11). Adding these two pieces from (4.10) and (4.11), the spectrum is

$$H = \frac{g^2}{8\pi^2} |\lambda_z^i - \lambda_z^j|^2 + \frac{g^2}{8\pi^2} |\lambda_y^i - \lambda_y^j|^2 = \frac{g^2}{8\pi^2} |\vec{\lambda}_i - \vec{\lambda}_j|^2.$$
(4.12)

The result subsumes the simplified case (4.6).

The extra energy is perturbation energy around the eigenvalue background, for a single pair of eigenvalues. We insert multiple words of this kind in (4.7). Hence for inserting multiple words of this kind,

$$H = \sum_{(i,j)} \frac{g^2}{8\pi^2} |\vec{\lambda}_i - \vec{\lambda}_j|^2,$$
(4.13)

where $(i, j) \in I_s$. This is in agreement with alternative observations [77, 78, 79]. The spectrum of this type of Hamiltonian was also studied by [79, 78, 77, 80, 81]; see also related methods [82].

In this case, by the method of centrally extended algebra [83],[84], the excitation energy are written in a square-root form, whose expansion gives (4.13). Hence the full square-root form is

$$\sum_{(i,j)} (\sqrt{1 + \frac{g^2}{4\pi^2}} |\vec{\lambda}_i - \vec{\lambda}_j|^2 - 1).$$
(4.14)

Here we subtract the bare dimension out in (4.14) to obtain the anormalous energy. The subtractions are the anormalous piece of the energy. The centrally extended algebra gives important insights in [84, 85, 72] and these are in agreement with the observations here.

These phenomena have very interesting dual interpretations on the gravity side. This is string spectra in backreacted excited state spacetime geometry. This analysis not only works for AdS background but also for backreacted backgrounds, in the context of string excitations in backreacted geometries. Eqn. (4.14) is in complete agreement with the analysis and observations [63, 86, 87, 88] on both the gauge theory side and the gravity side. This spectrum is in agreement with observations in the gravity side, in terms of magnon energy in backreacted geometries, e.g. magnon energy in annularly shaped droplet geometries.

The Eqn. (4.13) can be viewed as in the nonrelativistic regimes of (4.14) and it was considered as the near BPS excitation of heavy giant gravitons [78], in their near BPS limit. Eqn. (4.14) is the relativistic version of (4.13). This near BPS limit would be the same as the one in spin-matrix theory [89, 90] and they are hence closely related.

5 Discussion

The coherent state representation facilitates the calculation of ladder operators and hence the dilatation operator. For instance, it converts dilatation operator manipulations to algebraic manipulations. The coherent state representation of the operators have the advantage of simplifying the action of the above involved dilatation operators. We also checked higher loop dilatation operators on quarter BPS coherent states and the action is again vanishing as expected. By using nonrenormalization theorems, e.g. [49, 23], we infer that the BPS coherent states are in the kernel of the anomalous dilatation operator. This construction also works for coulomb branches.

In its original invention, [16] constructed coherent states averaged over group orbit, for the purpose of gauge invariance. Here we alternatively interpret this group average as a *path integral* of auxiliary variables coupled to the elementary letters of the theory. The insertions of string words on the coherent state can be viewed as operator-insertion in this path integral. We have generalized the word insertion to multi-word insertions, by conveniently interpreting them as operator insertions in this path integral. Hence, the group integral can be viewed as a path integral of the auxiliary variables U which are coupled to the elementary letters and added string words.

We also constructed a meromorphic version of coherent states, which can be transformed back to the full coherent states when det $\Lambda_Z \neq 0$. Here $G[T_Y, \Lambda_Z]$ is a meromorphic version of $F[\Lambda_Y, \Lambda_Z]$. The meromorphic versions are particularly useful when there are background Z fields. We also constructed meromorphic version of coherent states for product gauge groups and quivers. We give transformation relations between different types of coherent states, among other things.

We constructed new type of coherent states in the SL(2) sectors and bosonic part of PSU(1,1|2) sectors. These are operators with derivative insertions. The large spin limit and small spin limit can be reached by varying the coherent state amplitudes. We have also discussed other new states related to them.

The different coherent states differ by the symmetry properties of the multi-parameters of the coherent states. The coherent states constructed in [16] have manifest permutation symmetries among the eigenvalues. Other coherent states have gauge-fixed these symmetries [37], and can be viewed as gauge-fixed versions. Although different construction of coherent states may have different norms, there are many norm-independent features and observables of coherent state construction, for example, when calculating the vev of $a_Z^{\dagger n} a_Z^l$, the norm factor of the numerator and of the denominator cancels each other and is hence norm-independent.

We have used these coherent states to describe BPS and near BPS states, in gaugestring correspondence. We analyzed string-added coherent states, in general a product of a coherent state and a word, and calculated anormalous dimension energy. The string-added coherent state captures a class of near BPS spectra. One can use more general words, by using BMN like operators or spin chain like operators. The method of [16] gives great insights for computing excitations on supersymmetric backgrounds. Magnon excitations have also been computed for both gauge theory side and gravity side in related works. The agreement to these calculations is astounding, because the underlying methods and physical perspectives of these different approaches are different and alternative to each other. This approach is also intimately related to the effective eigenvalue model. In these involved circumstances, the eigenvalues provide some string theory background, and the off-diagonal fluctuations provide string excitations [8]. Non-BPS excitations are added as a perturbation of the BPS excitations. These observations pointed out to a way to solve the strong coupling dynamics of these near-BPS systems, and we have encountered some new strong coupling phenomena in those regimes. On a very different perspective, the approach of adding excitations may be relevant for |91, 92|.

On the gravity side, these coherent states are related to giant gravitons and backreacted geometries. The individual eigenvalue of coherent state is the collective coordinate of giant gravitons [16], see also related discussions [18, 37]. The coherent state construction is an important step in this regime. These are regimes where the single-trace and multi-trace operators are very difficult to handle efficiently. The effective action of the coherent state collective coordinate in the Lagrangian formalism has been constructed [16], where properties of giant gravitons and geometric quotient and projection of the transverse space in the dual gravity geometry, in the case of quiver theory, have been observed by this approach [16]. These are important for giant gravitons and backreacted geometries and related to various important discussions of e.g. [93]-[97].

Our approach shield new lights to near BPS states and near BPS sectors. We conveniently look at eighth BPS states and near BPS states, in this set-up. Another way to understand near BPS states is by spin-matrix theory e.g. [89, 90], which is closely related to the approaches in this paper. The spin-matrix theory is very insightful for both near BPS string states and near BPS giant gravitons. This is intimately related to the consistent reductions. The related consistent reductions are also very interestingly discussed in [98] to fewer dimensions.

The involved path integral can be computed by saddle point method and it is an exact computation in the context of localization. The localization technique is also very useful for Wilson operators. Localization methods have occurred remarkably in Wilson loops in e.g. [99] and related references. Moreover, emergent geometries are also dual to large Wilson loop operators, e.g. [100]-[104] as some examples. It would be good to have an unified understanding together with these circumstances.

The coherent state operators, Young tableau operators, and large Wilson operators are heavy operators. They have gravitational dual descriptions. The gravitational dual system to these very heavy operators, involves backreacted emergent geometries, e.g. [105]-[114] as some examples, and see their related references. These heavy states induce backreactions in the dual quantum gravity system. The coherent states are also heavy excited states. The coherent state approach leads to higher multi-pole moments in the gravity side, which can be measured on the boundary of the gravity side, as discussed in Sec. 2. Our scenarios are closely related to fuzzball geometries and their related discussions, e.g. [115]-[121] and [122].

Near BPS states can also describe near BPS black holes, which are related to various giant configurations and intersecting giants. The string configurations [123] on giants are relevant. A deeper understanding of the quarter and eighth BPS sector as well as near BPS sector will lead to implications for physics of extremal and near-extremal black holes e.g. [124, 125].

The SO(N) and Sp(N) cases of this approach have been worked out in [17]. The related Young tableaux basis have been addressed and developed in details in e.g. [27, 126, 127] and their related references. The case of other gauge groups and product gauge groups have also been considered [17, 16], as well as in this paper. This approach also works for SO/Sp theories, quiver theories, and coulomb branches. Although the analysis of the current paper is mainly for U(N) and SU(N) case, many features can be similar in the case of these other gauge groups.

Acknowledgments

We would like to thank B. Czech, R. de Mello Koch, L.-Y. Hung, Y. Jiang, S. Lin, Y. Liu, S. Ramgoolam, J. Simon, M. Sperling, R. Suzuki for useful discussions or communications. The work was supported in part by National Key R&D Program of China grant No. 2020YFA0713000, by Overseas high-level talents program, and by Fundamental Research Funds for the Central Universities of China.

References

 J. M. Maldacena, "The Large N limit of superconformal field theories and supergravity," Adv. Theor. Math. Phys. 2 (1998), 231 [hep-th/9711200].

- S. S. Gubser, I. R. Klebanov and A. M. Polyakov, "Gauge theory correlators from noncritical string theory," Phys. Lett. B 428 (1998), 105 [arXiv:hep-th/9802109 [hep-th]].
- [3] E. Witten, "Anti-de Sitter space and holography," Adv. Theor. Math. Phys. 2 (1998), 253 [arXiv:hep-th/9802150 [hep-th]].
- [4] N. Beisert, C. Ahn, L. F. Alday, Z. Bajnok, J. M. Drummond, L. Freyhult, N. Gromov, R. A. Janik, V. Kazakov and T. Klose, *et al.* "Review of AdS/CFT Integrability: An Overview," Lett. Math. Phys. **99** (2012) 3 [arXiv:1012.3982 [hep-th]].
- [5] M. Rangamani and T. Takayanagi, "Holographic Entanglement Entropy," Lect. Notes Phys. 931 (2017), pp.1-246 [arXiv:1609.01287 [hep-th]].
- [6] G. T. Horowitz and J. Polchinski, "Gauge/gravity duality," In: D. Oriti, Approaches to quantum gravity, Cambridge University Press [gr-qc/0602037].
- [7] R. de Mello Koch and J. Murugan, "Emergent Spacetime," In: G. Ellis et. al., Foundations of Space and Time: Reflections on Quantum Gravity, Cambridge University Press [arXiv:0911.4817 [hep-th]].
- [8] D. Berenstein, "Large N BPS states and emergent quantum gravity," JHEP 01 (2006), 125 [arXiv:hep-th/0507203 [hep-th]].
- [9] V. Balasubramanian, J. de Boer, V. Jejjala and J. Simon, "The Library of Babel: On the origin of gravitational thermodynamics," JHEP 12 (2005), 006 [arXiv:hep-th/0508023 [hep-th]].
- [10] J. McGreevy, L. Susskind and N. Toumbas, "Invasion of the giant gravitons from Anti-de Sitter space," JHEP 06 (2000), 008 [arXiv:hep-th/0003075 [hep-th]].
- [11] M. T. Grisaru, R. C. Myers and O. Tafjord, "SUSY and goliath," JHEP 08 (2000), 040 [arXiv:hep-th/0008015 [hep-th]].
- [12] A. Hashimoto, S. Hirano and N. Itzhaki, "Large branes in AdS and their field theory dual," JHEP 08 (2000), 051 [arXiv:hep-th/0008016 [hep-th]].
- [13] S. Corley, A. Jevicki and S. Ramgoolam, "Exact correlators of giant gravitons from dual N=4 SYM theory," Adv. Theor. Math. Phys. 5 (2002), 809 [arXiv:hepth/0111222 [hep-th]].

- [14] D. Berenstein, "A Toy model for the AdS/CFT correspondence," JHEP 07 (2004), 018 [arXiv:hep-th/0403110 [hep-th]].
- [15] H. Lin, O. Lunin and J. M. Maldacena, "Bubbling AdS space and 1/2 BPS geometries," JHEP 10 (2004), 025 [arXiv:hep-th/0409174 [hep-th]].
- [16] D. Berenstein and S. Wang, "BPS coherent states and localization," [arXiv:2203.15820 [hep-th]].
- [17] A. Holguin and S. Wang, "Giant Gravitons, Harish-Chandra integrals, and BPS states in symplectic and orthogonal $\mathcal{N}=4$ SYM," [arXiv:2206.00020 [hep-th]].
- [18] D. Berenstein and A. Miller, "Superposition induced topology changes in quantum gravity," JHEP 11 (2017), 121 [arXiv:1702.03011 [hep-th]].
- [19] S. Ramgoolam, "Schur-Weyl duality as an instrument of Gauge-String duality," AIP Conf. Proc. 1031 (2008) 255 [arXiv:0804.2764 [hep-th]].
- [20] T. W. Brown, P. J. Heslop and S. Ramgoolam, "Diagonal free field matrix correlators, global symmetries and giant gravitons," JHEP 04 (2009), 089 [arXiv:0806.1911 [hep-th]].
- [21] R. Bhattacharyya, S. Collins and R. de Mello Koch, "Exact Multi-Matrix Correlators," JHEP 03 (2008), 044 [arXiv:0801.2061 [hep-th]].
- [22] R. de Mello Koch, P. Diaz and N. Nokwara, "Restricted Schur Polynomials for Fermions and integrability in the su(2|3) sector," JHEP 03 (2013), 173 [arXiv:1212.5935 [hep-th]].
- [23] J. Pasukonis and S. Ramgoolam, "From counting to construction of BPS states in N=4 SYM," JHEP 02 (2011), 078 [arXiv:1010.1683 [hep-th]].
- [24] R. de Mello Koch, N. Ives and M. Stephanou, "Correlators in Nontrivial Backgrounds," Phys. Rev. D 79 (2009), 026004 [arXiv:0810.4041 [hep-th]].
- [25] R. Suzuki, "Three-point functions in $\mathcal{N} = 4$ SYM at finite N_c and background independence," JHEP **05** (2020), 118 [arXiv:2002.07216 [hep-th]].
- [26] R. de Mello Koch, P. Diaz and H. Soltanpanahi, "Non-planar Anomalous Dimensions in the sl(2) Sector," Phys. Lett. B 713 (2012), 509-513 [arXiv:1111.6385 [hep-th]].

- [27] P. Caputa, R. de Mello Koch and P. Diaz, "A basis for large operators in N=4 SYM with orthogonal gauge group," JHEP **03** (2013), 041 [arXiv:1301.1560 [hep-th]].
- [28] Y. Kimura and S. Ramgoolam, "Branes, anti-branes and brauer algebras in gaugegravity duality," JHEP 11 (2007), 078 [arXiv:0709.2158 [hep-th]].
- [29] D. Gaiotto and J. H. Lee, "The Giant Graviton Expansion," [arXiv:2109.02545 [hep-th]].
- [30] K. Budzik and D. Gaiotto, "Giant gravitons in twisted holography," [arXiv:2106.14859 [hep-th]].
- [31] Y. Jiang, S. Komatsu and E. Vescovi, "Structure constants in $\mathcal{N} = 4$ SYM at finite coupling as worldsheet g-function," JHEP **07** (2020) no.07, 037 [arXiv:1906.07733 [hep-th]].
- [32] G. Chen, R. de Mello Koch, M. Kim and H. J. R. Van Zyl, "Absorption of closed strings by giant gravitons," JHEP 10 (2019), 133 [arXiv:1908.03553 [hep-th]].
- [33] S. Murthy, "Unitary matrix models, free fermion ensembles, and the giant graviton expansion," [arXiv:2202.06897 [hep-th]].
- [34] R. de Mello Koch, J. Smolic and M. Smolic, "Giant Gravitons with Strings Attached (I)," JHEP 06 (2007), 074 [arXiv:hep-th/0701066 [hep-th]].
- [35] H. Lin, "Giant gravitons and correlators," JHEP 12 (2012), 011 [arXiv:1209.6624 [hep-th]].
- [36] J. Simon, "Correlations vs connectivity in R-charge," JHEP 10 (2018), 048 [arXiv:1805.11279 [hep-th]].
- [37] H. Lin and K. Zeng, "Detecting topology change via correlations and entanglement from gauge/gravity correspondence," J. Math. Phys. 59 (2018) no.3, 032301 [arXiv:1705.10776 [hep-th]].
- [38] H. Lin and Y. Zhu, "Coherent state superpositions, entanglement and gauge/gravity correspondence," J. Math. Phys. 62 (2021) no.5, 052301 [arXiv:2011.13919 [hep-th]].
- [39] H. Lin and Y. Zhu, "Entanglement and mixed states of Young tableau states in gauge/gravity correspondence," Nucl. Phys. B 972 (2021), 115572 [arXiv:2107.14219 [hep-th]].

- [40] T. W. Brown, R. de Mello Koch, S. Ramgoolam and N. Toumbas, "Correlators, Probabilities and Topologies in N=4 SYM," JHEP 03 (2007), 072 [arXiv:hep-th/0611290 [hep-th]].
- [41] P. Diaz, H. Lin and A. Veliz-Osorio, "Graph duality as an instrument of Gauge-String correspondence," J. Math. Phys. 57 (2016) no.5, 052302 [arXiv:1505.04837 [hep-th]].
- [42] V. Balasubramanian, B. Czech, K. Larjo, D. Marolf and J. Simon, "Quantum geometry and gravitational entropy," JHEP 12 (2007), 067 [arXiv:0705.4431 [hepth]].
- [43] D. Berenstein and A. Miller, "Can Topology and Geometry be Measured by an Operator Measurement in Quantum Gravity?," Phys. Rev. Lett. **118** (2017) no.26, 261601 [arXiv:1605.06166 [hep-th]].
- [44] S. D. Mathur, "Tunneling into fuzzball states," Gen. Rel. Grav. 42 (2010), 113 [arXiv:0805.3716 [hep-th]].
- [45] Harish-Chandra, "Differential operators on a semisimple Lie algebra," Amer. J. Math. 79 (1957), no. 1, 87.
- [46] C. Itzykson and J. B. Zuber, "The Planar Approximation. 2.," J. Math. Phys. 21 (1980), 411.
- [47] J. J. Duistermaat and G. J. Heckman, "On the Variation in the cohomology of the symplectic form of the reduced phase space," Invent. Math. 69 (1982) 259.
- [48] A. Borodin, G. Olshanski, "The Young bouquet and its boundary," Mosc. Math. J. 13 (2013) 193-232 [arXiv:1110.4458].
- [49] C. Lewis-Brown and S. Ramgoolam, "Quarter-BPS states, multi-symmetric functions and set partitions," JHEP 03 (2021), 153 [arXiv:2007.01734 [hep-th]].
- [50] M. Masuku and J. P. Rodrigues, "Laplacians in polar matrix coordinates and radial fermionization in higher dimensions," J. Math. Phys. 52 (2011), 032302 [arXiv:0911.2846 [hep-th]].
- [51] D. Berenstein, D. H. Correa and S. E. Vazquez, "All loop BMN state energies from matrices," JHEP 02 (2006), 048 [arXiv:hep-th/0509015 [hep-th]].

- [52] D. H. Correa and M. Wolf, "Shaping up BPS States with Matrix Model Saddle Points," J. Phys. A 43 (2010), 465402 [arXiv:1007.5284 [hep-th]].
- [53] R. de Mello Koch, D. Gossman, L. Nkumane and L. Tribelhorn, "Eigenvalue Dynamics for Multimatrix Models," Phys. Rev. D 96 (2017) no.2, 026011 [arXiv:1608.00399 [hep-th]].
- [54] D. E. Berenstein, M. Hanada and S. A. Hartnoll, "Multi-matrix models and emergent geometry," JHEP 02 (2009), 010 [arXiv:0805.4658 [hep-th]].
- [55] K. Skenderis and M. Taylor, "Anatomy of bubbling solutions," JHEP 09 (2007), 019 [arXiv:0706.0216 [hep-th]].
- [56] K. Skenderis and M. Taylor, "Holographic Coulomb branch vevs," JHEP 08 (2006), 001 [arXiv:hep-th/0604169 [hep-th]].
- [57] V. Balasubramanian, B. Czech, K. Larjo and J. Simon, "Integrability versus information loss: A Simple example," JHEP **11** (2006), 001 [arXiv:hep-th/0602263 [hep-th]].
- [58] L. Grant, L. Maoz, J. Marsano, K. Papadodimas and V. S. Rychkov, "Minisuperspace quantization of Bubbling AdS and free fermion droplets," JHEP 08 (2005), 025 [arXiv:hep-th/0505079 [hep-th]].
- [59] V. Balasubramanian, B. Czech, V. E. Hubeny, K. Larjo, M. Rangamani and J. Simon, "Typicality versus thermality: An Analytic distinction," Gen. Rel. Grav. 40 (2008), 1863-1890 [arXiv:hep-th/0701122 [hep-th]].
- [60] B. Chen, S. Cremonini, A. Donos, F. L. Lin, H. Lin, J. T. Liu, D. Vaman and W. Y. Wen, "Bubbling AdS and droplet descriptions of BPS geometries in IIB supergravity," JHEP 10 (2007), 003 [arXiv:0704.2233 [hep-th]].
- [61] J. T. Liu, H. Lu, C. N. Pope and J. F. Vazquez-Poritz, "Bubbling AdS black holes," JHEP 10 (2007), 030 [arXiv:hep-th/0703184 [hep-th]].
- [62] H. Lin and K. Zeng, "A construction of quarter BPS coherent states and Brauer algebras," Adv. Theor. Math. Phys. 24 (2020) no.5, 1111-1169 [arXiv:1709.10093 [hep-th]].
- [63] D. Berenstein and A. Holguin, "Open giant magnons on LLM geometries," JHEP 01 (2021), 080 [arXiv:2010.02236 [hep-th]]

- [64] M. Kim and H. J. R. Van Zyl, "Semiclassical SL(2) strings on LLM backgrounds," Phys. Lett. B **784** (2018), 62-67 [arXiv:1805.12460 [hep-th]].
- [65] A. A. Tseytlin, "Quantum strings in AdS(5) x S*5 and AdS/CFT duality," Int. J. Mod. Phys. A 25 (2010) 319.
- [66] A. V. Belitsky, A. S. Gorsky and G. P. Korchemsky, "Gauge/string duality for QCD conformal operators," Nucl. Phys. B 667 (2003), 3-54 [arXiv:hep-th/0304028 [hep-th]].
- [67] N. Beisert and B. I. Zwiebel, "On Symmetry Enhancement in the psu(1,1|2) Sector of N=4 SYM," JHEP 10 (2007), 031 [arXiv:0707.1031 [hep-th]].
- [68] D. Berenstein and R. Cotta, "Aspects of emergent geometry in the AdS/CFT context," Phys. Rev. D 74 (2006), 026006 [arXiv:hep-th/0605220 [hep-th]].
- [69] D. E. Berenstein and S. A. Hartnoll, "Strings on conifolds from strong coupling dynamics: Quantitative results," JHEP 03 (2008), 072 [arXiv:0711.3026 [hep-th]].
- [70] K. J. Larsen and N. A. Obers, "Phases of Thermal N=2 Quiver Gauge Theories," JHEP 01 (2008), 057 [arXiv:0708.3199 [hep-th]].
- [71] V. Mitev and E. Pomoni, "Exact effective couplings of four dimensional gauge theories with $\mathcal{N} = 2$ supersymmetry," Phys. Rev. D **92** (2015) no.12, 125034 [arXiv:1406.3629 [hep-th]].
- [72] A. Gadde and L. Rastelli, "Twisted Magnons," JHEP 04 (2012), 053 [arXiv:1012.2097 [hep-th]]
- [73] E. Conde, H. Lin, J. M. Penin, A. V. Ramallo and D. Zoakos, "D3–D5 theories with unquenched flavors," Nucl. Phys. B 914 (2017), 599-622 [arXiv:1607.04998 [hep-th]].
- [74] J. Pasukonis and S. Ramgoolam, "Quivers as Calculators: Counting, Correlators and Riemann Surfaces," JHEP 04 (2013), 094 [arXiv:1301.1980 [hep-th]].
- [75] D. E. Berenstein, J. M. Maldacena and H. S. Nastase, "Strings in flat space and pp waves from N=4 superYang-Mills," JHEP 04 (2002), 013 [arXiv:hep-th/0202021 [hep-th]].

- [76] N. Beisert, C. Kristjansen and M. Staudacher, "The Dilatation operator of conformal N=4 superYang-Mills theory," Nucl. Phys. B 664 (2003), 131-184 [arXiv:hepth/0303060 [hep-th]].
- [77] R. de Mello Koch, G. Kemp and S. Smith, "From Large N Nonplanar Anomalous Dimensions to Open Spring Theory," Phys. Lett. B 711 (2012), 398-403 [arXiv:1111.1058 [hep-th]].
- [78] W. Carlson, R. de Mello Koch and H. Lin, "Nonplanar Integrability," JHEP 03 (2011), 105 [arXiv:1101.5404 [hep-th]].
- [79] D. Berenstein, "Giant gravitons: a collective coordinate approach," Phys. Rev. D 87 (2013) no.12, 126009 [arXiv:1301.3519 [hep-th]].
- [80] R. de Mello Koch and S. Ramgoolam, "A double coset ansatz for integrability in AdS/CFT," JHEP **06** (2012), 083 [arXiv:1204.2153 [hep-th]].
- [81] R. de Mello Koch, M. Dessein, D. Giataganas and C. Mathwin, "Giant Graviton Oscillators," JHEP 10 (2011), 009 [arXiv:1108.2761 [hep-th]].
- [82] M. Cook and J. P. Rodrigues, "Strongly coupled large N spectrum of two matrices coupled via a Yang-Mills interaction," Phys. Rev. D 78 (2008), 065024 [arXiv:0710.0073 [hep-th]].
- [83] N. Beisert, "The SU(2|2) dynamic S-matrix," Adv. Theor. Math. Phys. 12 (2008), 945-979 [arXiv:hep-th/0511082 [hep-th]].
- [84] D. Berenstein, "On the central charge extension of the $\mathcal{N} = 4$ SYM spin chain," JHEP **05** (2015), 129 [arXiv:1411.5921 [hep-th]].
- [85] S. de Carvalho, R. de Mello Koch and M. Kim, "Central Charges for the Double Coset," JHEP 05 (2020), 007 [arXiv:2001.10181[hep-th]].
- [86] D. Berenstein and A. Holguin, "Open giant magnons suspended between dual giant gravitons in $\mathcal{N} = 4$ SYM," JHEP **09** (2020), 019 [arXiv:2006.08649 [hep-th]].
- [87] R. de Mello Koch, C. Mathwin and H. J. R. Van Zyl, "LLM Magnons," JHEP 03 (2016), 110 [arXiv:1601.06914 [hep-th]].
- [88] R. de Mello Koch, J. H. Huang and L. Tribelhorn, "Exciting LLM Geometries," JHEP 07 (2018), 146 [arXiv:1806.06586 [hep-th]].

- [89] S. Baiguera, T. Harmark and Y. Lei, "Spin Matrix Theory in near $\frac{1}{8}$ -BPS corners of $\mathcal{N} = 4$ super-Yang-Mills," JHEP **02** (2022), 191 [arXiv:2111.10149 [hep-th]].
- [90] T. Harmark, "Interacting Giant Gravitons from Spin Matrix Theory," Phys. Rev. D 94 (2016) no.6, 066001 [arXiv:1606.06296 [hep-th]].
- [91] A. Morozov, "Pair correlator in the Itzykson-Zuber integral," Mod. Phys. Lett. A 7 (1992), 3503-3508 [arXiv:hep-th/9209074 [hep-th]].
- [92] V. A. Kazakov and A. A. Migdal, "Induced QCD at large N," Nucl. Phys. B 397 (1993) 214 [arXiv:hep-th/9206015 [hep-th]].
- [93] I. Biswas, D. Gaiotto, S. Lahiri and S. Minwalla, "Supersymmetric states of N=4 Yang-Mills from giant gravitons," JHEP 12 (2007), 006 [arXiv:hep-th/0606087 [hep-th]].
- [94] G. Mandal and N. V. Suryanarayana, "Counting 1/8-BPS dual-giants," JHEP 03 (2007), 031 [arXiv:hep-th/0606088 [hep-th]].
- [95] A. Basu and G. Mandal, "Dual giant gravitons in AdS(m) x Y(n) (Sasaki-Einstein)," JHEP 07 (2007), 014 [arXiv:hep-th/0608093 [hep-th]].
- [96] D. Martelli and J. Sparks, "Dual Giant Gravitons in Sasaki-Einstein Backgrounds," Nucl. Phys. B 759 (2006), 292 [arXiv:hep-th/0608060 [hep-th]].
- [97] J. Pasukonis and S. Ramgoolam, "Quantum states to brane geometries via fuzzy moduli spaces of giant gravitons," JHEP **04** (2012), 077 [arXiv:1201.5588 [hep-th]].
- [98] G. Ishiki, Y. Takayama and A. Tsuchiya, "N=4 SYM on R x S*3 and theories with 16 supercharges," JHEP 10 (2006), 007 [arXiv:hep-th/0605163 [hep-th]].
- [99] V. Pestun, "Localization of the four-dimensional N=4 SYM to a two-sphere and 1/8 BPS Wilson loops," JHEP 12 (2012), 067 [arXiv:0906.0638 [hep-th]].
- [100] S. Yamaguchi, "Bubbling geometries for half BPS Wilson lines," Int. J. Mod. Phys. A 22 (2007), 1353-1374 [arXiv:hep-th/0601089 [hep-th]].
- [101] O. Lunin, "On gravitational description of Wilson lines," JHEP 0606 (2006) 026 [hep-th/0604133].
- [102] J. Gomis and F. Passerini, "Holographic Wilson Loops," JHEP 08 (2006), 074 [arXiv:hep-th/0604007 [hep-th]].

- [103] E. D'Hoker, J. Estes and M. Gutperle, "Gravity duals of half-BPS Wilson loops," JHEP 0706 (2007) 063 [arXiv:0705.1004 [hep-th]].
- [104] J. Gomis, S. Matsuura, T. Okuda and D. Trancanelli, "Wilson loop correlators at strong coupling: From matrices to bubbling geometries," JHEP 08 (2008), 068 [arXiv:0807.3330 [hep-th]].
- [105] E. Gava, G. Milanesi, K. S. Narain and M. O'Loughlin, "1/8 BPS states in Ads/CFT," JHEP 05 (2007), 030 [arXiv:hep-th/0611065 [hep-th]].
- [106] Y. Kimura and H. Lin, "Young diagrams, Brauer algebras, and bubbling geometries," JHEP 01 (2012), 121 [arXiv:1109.2585 [hep-th]].
- [107] J. P. Gauntlett, N. Kim and D. Waldram, "Supersymmetric AdS(3), AdS(2) and Bubble Solutions," JHEP 04 (2007), 005 [arXiv:hep-th/0612253 [hep-th]].
- [108] N. Kim, "AdS(3) solutions of IIB supergravity from D3-branes," JHEP 01 (2006), 094 [arXiv:hep-th/0511029 [hep-th]].
- [109] J. T. Liu, H. Lu, C. N. Pope and J. F. Vazquez-Poritz, "New supersymmetric solutions of N=2, D=5 gauged supergravity with hyperscalars," JHEP 10 (2007), 093 [arXiv:0705.2234 [hep-th]].
- [110] Z.-W. Chong, H. Lu and C. N. Pope, "BPS geometries and AdS bubbles," Phys. Lett. B 614 (2005) 96 [hep-th/0412221].
- [111] S. Gukov and E. Witten, "Rigid Surface Operators," Adv. Theor. Math. Phys. 14 (2010) no.1, 87-178 [arXiv:0804.1561 [hep-th]].
- [112] N. Drukker, J. Gomis and S. Matsuura, "Probing N=4 SYM With Surface Operators," JHEP 10 (2008), 048 [arXiv:0805.4199 [hep-th]].
- [113] J. Gomis and S. Matsuura, "Bubbling surface operators and S-duality," JHEP 0706 (2007) 025 [arXiv:0704.1657 [hep-th]].
- [114] S. Mukhi and M. Smedback, "Bubbling orientifolds," JHEP 08 (2005), 005 [arXiv:hep-th/0506059 [hep-th]].
- [115] S. D. Mathur, "The Quantum structure of black holes," Class. Quant. Grav. 23 (2006), R115 [arXiv:hep-th/0510180 [hep-th]].

- [116] I. Bena and N. P. Warner, "Black holes, black rings and their microstates," Lect. Notes Phys. 755 (2008), 1-92 [arXiv:hep-th/0701216 [hep-th]].
- [117] K. Skenderis and M. Taylor, "The fuzzball proposal for black holes," Phys. Rept. 467 (2008), 117-171 [arXiv:0804.0552 [hep-th]].
- [118] V. Balasubramanian, J. de Boer, S. El-Showk and I. Messamah, "Black Holes as Effective Geometries," Class. Quant. Grav. 25 (2008), 214004 [arXiv:0811.0263 [hep-th]].
- [119] I. Bena, E. J. Martinec, R. Walker and N. P. Warner, "Early Scrambling and Capped BTZ Geometries," JHEP 04 (2019), 126 [arXiv:1812.05110 [hep-th]].
- [120] D. R. Mayerson, "Fuzzballs and Observations," Gen. Rel. Grav. 52 (2020) no.12, 115 [arXiv:2010.09736 [hep-th]].
- [121] B. Chakrabarty, S. Rawash and D. Turton, "Shockwaves in black hole microstate geometries," JHEP 02 (2022), 202 [arXiv:2112.08378 [hep-th]].
- [122] A. Belin and B. Withers, "From sources to initial data and back again: on bulk singularities in Euclidean AdS/CFT," JHEP 12 (2020), 185 [arXiv:2007.10344 [hep-th]].
- [123] D. Berenstein and A. Holguin, "String junctions suspended between giants," [arXiv:2202.11729 [hep-th]].
- [124] V. Balasubramanian, J. de Boer, V. Jejjala and J. Simon, "Entropy of nearextremal black holes in AdS(5)," JHEP 05 (2008), 067 [arXiv:0707.3601 [hep-th]].
- [125] R. Fareghbal, C. N. Gowdigere, A. E. Mosaffa and M. M. Sheikh-Jabbari, "Nearing Extremal Intersecting Giants and New Decoupled Sectors in N = 4 SYM," JHEP 08 (2008), 070 [arXiv:0801.4457 [hep-th]].
- [126] P. Caputa, R. de Mello Koch and P. Diaz, "Operators, Correlators and Free Fermions for SO(N) and Sp(N)," JHEP 06 (2013), 018 [arXiv:1303.7252 [hep-th]].
- [127] C. Lewis-Brown and S. Ramgoolam, "BPS operators in $\mathcal{N} = 4$ SO(N) super Yang-Mills theory: plethysms, dominoes and words," JHEP **11** (2018), 035 [arXiv:1804.11090 [hep-th]].