

Period Integrals (Givental's I -function) of Calabi-Yau Hypersurface in CP^{N-1} and Intersection Numbers of Moduli Space of Quasimaps from CP^1 with Two Marked Points to CP^{N-1}

Masao Jinzenji⁽¹⁾, Kohki Matsuzaka⁽²⁾

⁽¹⁾ Department of Mathematics,
Okayama University
Okayama, 700-8530, Japan

⁽²⁾ Division of Mathematics, Graduate School of Science
Hokkaido University
Kita-ku, Sapporo, 060-0810, Japan

e-mail address: ⁽¹⁾ pcj70e4e@okayama-u.ac.jp
⁽²⁾ kohki@math.sci.hokudai.ac.jp

June 22, 2022

Abstract

In this paper, we derive the generalized hypergeometric functions (period integrals) used in mirror computation of Calabi-Yau hypersurface in CP^{N-1} as generating functions of intersection numbers of the moduli space of quasimaps from CP^1 with two marked points to CP^{N-1} .

1 Introduction

The aim of this paper is to clarify geometrical meaning of period integrals or generalized hypergeometric functions that are given as solutions of the following ordinary differential equation:

$$\left(\left(\frac{d}{dx} \right)^{N-1} - N \cdot e^x \cdot \left(N \frac{d}{dx} + (N-1) \right) \left(N \frac{d}{dx} + (N-2) \right) \cdots \left(N \frac{d}{dx} + 1 \right) \right) W(x) = 0. \quad (1.1)$$

These were used in the mirror computation of genus 0 Gromov-Witten invariants of Calabi-Yau hypersurface in CP^{N-1} [3, 8]. By using Frobenius's method, the solutions are explicitly given as follows.

$$\begin{aligned} W_j(x) &:= \frac{\partial^j}{\partial \epsilon^j} \left(\sum_{d=0}^{\infty} \left(\frac{\prod_{r=1}^{Nd} (r+N\epsilon)}{\prod_{r=1}^d (r+\epsilon)^N} \right) e^{(d+\epsilon)x} \right) \Big|_{\epsilon=0} \\ &= \sum_{i=0}^j \binom{j}{i} x^{j-i} \left(\sum_{d=0}^{\infty} \frac{\partial^i}{\partial \epsilon^i} \left(\frac{\prod_{r=1}^{Nd} (r+N\epsilon)}{\prod_{r=1}^d (r+\epsilon)^N} \right) \Big|_{\epsilon=0} e^{dx} \right), \end{aligned}$$

$$=: \sum_{i=0}^j \binom{j}{i} x^{j-i} w_i(x) \quad (j = 0, 1, \dots, N-2). \quad (1.2)$$

On the other hand, geometrical study of mirror symmetry from the point of view of moduli space of quasimaps has been pursued by several groups of researchers [1, 5]. Let us focus here the line of study in this direction pursued by our group. Here, we restrict our attention to the Calabi-Yau hypersurface in CP^{N-1} . In [5], we construct the moduli space $\widetilde{Mp}_{0,2}(N, d)$ of quasimaps (polynomial maps) of degree d from CP^1 with two marked points to CP^{N-1} and defined the intersection number $w(\mathcal{O}_{h^a} \mathcal{O}_{h^b})_{0,d}$ ($a+b=N-3$ and h is the hyperplane class in $H^*(CP^{N-1}, \mathbf{C})$). This corresponds to B-model analogue of the genus 0 and degree d Gromov-Witten invariant $\langle \mathcal{O}_{h^a} \mathcal{O}_{h^b} \rangle_{0,d}$ of the Calabi-Yau hypersurface in CP^{N-1} . Let us introduce the following generating functions of these intersection numbers.

$$\begin{aligned} w(\mathcal{O}_{h^a} \mathcal{O}_{h^b})_0(x) &:= Nx + \sum_{d=1}^{\infty} w(\mathcal{O}_{h^a} \mathcal{O}_{h^b})_{0,d} e^{dx}, \\ \langle \mathcal{O}_{h^a} \mathcal{O}_{h^b} \rangle_0(t) &:= Nt + \sum_{d=1}^{\infty} \langle \mathcal{O}_{h^a} \mathcal{O}_{h^b} \rangle_{0,d} e^{dt}. \end{aligned} \quad (1.3)$$

We have proved the following equalities:

$$\begin{aligned} w(\mathcal{O}_{h^{N-3}} \mathcal{O}_{h^0})_0(x) &= Nt(x) = N \frac{W_1(x)}{W_0(x)} = N(x + \frac{w_1(x)}{w_0(x)}), \\ \langle \mathcal{O}_{h^a} \mathcal{O}_{h^b} \rangle_0(t(x)) &= w(\mathcal{O}_{h^a} \mathcal{O}_{h^b})_0(x). \end{aligned} \quad (1.4)$$

The first equality tells us that $w(\mathcal{O}_{h^{N-3}} \mathcal{O}_{h^0})_0(x)$ gives us the mirror map used in mirror computation of genus 0 Gromov-Witten invariants of the hypersurface. It was proved in [5]. The second equality tells that $w(\mathcal{O}_{h^a} \mathcal{O}_{h^b})_0(x)$ is translated into generating function of the corresponding Gromov-Witten invariants via the mirror map. Geometrical proof of it was given in [6]. In [13], Saito gave explicit toric construction of $\widetilde{Mp}_{0,2}(N, d)$ and showed that it is a compact orbifold. Moreover, he determined Chow ring of $\widetilde{Mp}_{0,2}(N, d)$. It is generated by $(d+1)$ generators H_0, H_1, \dots, H_d and relations of the generators are given by,

$$(H_0)^N = 0, (H_i)^N (2H_i - H_{i-1} - H_{i+1}) = 0 \quad (i = 1, 2, \dots, d-1), (H_d)^N = 0. \quad (1.5)$$

With these notations, $w(\mathcal{O}_{h^a} \mathcal{O}_{h^b})_{0,d}$ can be explicitly written as follows:

$$w(\mathcal{O}_{h^a} \mathcal{O}_{h^b})_{0,d} = \int_{\widetilde{Mp}_{0,2}(N, d)} (H_0)^a \left(\frac{\prod_{j=1}^d e(H_{j-1}, H_j)}{\prod_{j=1}^{d-1} (NH_j)} \right) (H_d)^b, \quad (1.6)$$

where $e(x, y) = \prod_{j=0}^N (jx + (N-j)y)$. This expression was already derived in [5] and was effectively used to prove the first equality of (1.4). Generalization of his explicit toric construction to the case of moduli space of quasimaps from CP^1 with two marked points to other toric manifolds have been given in [10, 12].

Then we are naturally led to the question:

“ Can we express the generalized hypergeometric function $W_j(x)$ in (1.2) as generating function of intersection numbers of $\widetilde{Mp}_{0,2}(N, d)$? ”

Of course, we have already obtained some results on this question in the $N=5$ case, which were presented in Chapter 5 of [7]. Moreover, we can obtain many hints from Givental’s theory of I -function and J -function [2]. In [2], Givental suggested that the hypergeometric functions are closely related to the two point Gromov-Witten invariants:

$$\langle \sigma_j(\mathcal{O}_{h^{N-3-j}}) \mathcal{O}_{h^0} \rangle_{0,d} := \int_{\overline{M}_{0,2}(CP^{N-1}, d)} (\psi_1)^j \wedge ev_1^*(h^{N-3-j}) \wedge c_T(\mathcal{E}_d). \quad (1.7)$$

where $\overline{M}_{0,2}(CP^{N-1}, d)$ is the moduli space of stable maps from genus 0 stable curve with two marked points to CP^{N-1} , ψ_1 is the Mumford-Morita-Miller class associated with the first marked point, $ev_1 : \overline{M}_{0,2}(CP^{N-1}, d) \rightarrow CP^{N-1}$ is the evaluation map at the first marked point and \mathcal{E}_d is the rank $Nd + 1$ vector bundle that imposes condition that image of the stable curve lies inside the Calabi-Yau hypersurface. Moreover, we have known that the element in Chow ring of $\widetilde{Mp}_{0,2}(N, d)$ that corresponds to ψ_1 in the Chow ring of $\overline{M}_{0,2}(CP^{N-1}, d)$ is given by $(H_1 - H_0)$ [11, 13].

In this paper, we define the following intersection number on $\widetilde{Mp}_{0,2}(N, d)$:

$$w(\sigma_j(\mathcal{O}_{h^a})\mathcal{O}_{h^b})_{0,d} = \int_{\widetilde{Mp}_{0,2}(N,d)} (H_0)^a (H_1 - H_0)^j \left(\frac{\prod_{j=1}^d e(H_{j-1}, H_j)}{\prod_{j=1}^{d-1} (NH_j)} \right) (H_d)^b, \quad (1.8)$$

and prove the following theorem.

Theorem 1

$$\begin{aligned} \frac{1}{N} w(\sigma_j(\mathcal{O}_{h^{N-2-j}})\mathcal{O}_{h^{-1}}|\mathcal{O}_h)_{0,2|1} &:= \frac{d}{N} w(\sigma_j(\mathcal{O}_{h^{N-2-j}})\mathcal{O}_{h^{-1}})_{0,2} + \frac{1}{N} w(\sigma_{j-1}(\mathcal{O}_{h^{N-1-j}})\mathcal{O}_{h^{-1}})_{0,2} \\ &=: \frac{1}{j!} \frac{\partial^j}{\partial \epsilon^j} \left(\frac{\prod_{r=1}^{Nd} (r + N\epsilon)}{\prod_{r=1}^d (r + \epsilon)^N} \right) \Big|_{\epsilon=0}. \end{aligned} \quad (1.9)$$

We have several remarks on the above theorem. Since we have insertion of $\mathcal{O}_{h^{-1}}$, we have to insert $\frac{1}{H_d}$ at the corresponding position in the formula (1.8). But this negative power cancels with H_d in the polynomial $e^N(H_{d-1}, H_d) = NH_{d-1}((N-1)H_{d-1} + H_d) \cdots (H_{d-1} + (N-1)H_d)NH_d$. Hence the integrands used in evaluating $w(\sigma_j(\mathcal{O}_{h^{N-2-j}})\mathcal{O}_{h^{-1}})_{0,2}$ and $w(\sigma_{j-1}(\mathcal{O}_{h^{N-1-j}})\mathcal{O}_{h^{-1}})_{0,2}$ are polynomials in H_0, \dots, H_d ¹. In (1.9), we formally introduce the notation:

$$w(\sigma_j(\mathcal{O}_{h^{N-2-j}})\mathcal{O}_{h^{-1}}|\mathcal{O}_h)_{0,d} := d \cdot w(\sigma_j(\mathcal{O}_{h^{N-2-j}})\mathcal{O}_{h^{-1}})_{0,d} + w(\sigma_{j-1}(\mathcal{O}_{h^{N-1-j}})\mathcal{O}_{h^{-1}})_{0,d}. \quad (1.10)$$

But we think that the intersection number $w(\sigma_j(\mathcal{O}_{h^{N-2-j}})\mathcal{O}_{h^{-1}}|\mathcal{O}_h)_{0,d}$ should be defined as three-pointed intersection number of the moduli space $\widetilde{Mp}_{0,2|1}(N, d)$ used in [9]. Since we haven't explicitly determined Chow ring of $\widetilde{Mp}_{0,2|1}(N, d)$ and don't know well-defined Mumford-Morita-Miller class of the moduli space, the above notation is formal. If the notation $w(\sigma_j(\mathcal{O}_{h^{N-2-j}})\mathcal{O}_{h^{-1}}|\mathcal{O}_h)_{0,d}$ is justified as the corresponding intersection number of $\widetilde{Mp}_{0,2|1}(N, d)$, (1.10) corresponds to "Hori's equation" [4] for a three-pointed gravitational virtual structure constant:

$$w(\sigma_j(\mathcal{O}_{h^a})\mathcal{O}_{h^b}|\mathcal{O}_h)_{0,d} = d \cdot w(\sigma_j(\mathcal{O}_{h^a})\mathcal{O}_{h^b})_{0,d} + w(\sigma_{j-1}(\mathcal{O}_{h^{a+1}})\mathcal{O}_{h^b})_{0,d}. \quad (1.11)$$

Using this notation and Theorem 1, we can derive the following corollary by straightforward computation.

Corollary 1 $W_j(x)$ is written as the following generating function of intersection numbers of $\widetilde{Mp}_{0,2}(N, d)$.

$$\begin{aligned} \frac{1}{j!} W_j(x) &= \sum_{d=0}^{\infty} \sum_{m=0}^{\infty} \sum_{k=0}^{\infty} \frac{x^m}{m!} \frac{1}{N} w(\sigma_k(\mathcal{O}_{h^{N-2-j+m}})\mathcal{O}_{h^{-1}}|\mathcal{O}_h)_{0,d} e^{dx} \\ &=: \sum_{d=0}^{\infty} \frac{1}{N} w\left(\frac{e^{hx} h^{N-2-j}}{1 - \tilde{\psi}_1}, h^{-1} | h\right)_{0,d} e^{dx}, \end{aligned}$$

¹The operator insertion of $\mathcal{O}_{h^{-1}}$ was also considered by Zinger [14] in defining the vector bundle \mathcal{V}'_0 on $\overline{M}_{0,2}(CP^{N-1}, d)$ in his notation.

where $\tilde{\psi}_1$ denotes “expected” Mumford-Morita-Miller class of $\widetilde{Mp}_{0,2|1}(N, d)$. Then, we can also express Givental’s I-function [2] as a generating function of intersection numbers of $\widetilde{Mp}_{0,2}(N, d)$.

$$\begin{aligned} I(P, x) := \sum_{j=0}^{N-2} \frac{1}{j!} W_j(x) P^j &= \sum_{j=0}^{N-2} \sum_{d=0}^{\infty} \frac{1}{N} w\left(\frac{e^{hx} h^{-j} P^j h^{N-2}}{1 - \tilde{\psi}_1}, h^{-1}|h\right)_{0,d} e^{dx} \\ &= \frac{P^{N-2}}{N} \sum_{d=0}^{\infty} w\left(\frac{e^{hx}}{(1 - \tilde{\psi}_1)(1 - \frac{h}{P})}, h^{-1}|h\right)_{0,d} e^{dx}. \end{aligned}$$

Acknowledgment We would like to thank Prof. B. Kim and Prof. T. Ohmoto for valuable discussions. Our research is partially supported by JSPS grant No. 22K03289.

2 Proof of the Main Theorem

Our proof of Theorem 1 is based on the following formula derived in [5, 13]:

$$\begin{aligned} w(\sigma_j(\mathcal{O}_{h^a})\mathcal{O}_{h^b})_{0,2} &= \frac{1}{(2\pi\sqrt{-1})^{d+1}} \oint_{C_0} \frac{dz_0}{(z_0)^N} \oint_{C_1} \frac{dz_1}{(z_1)^N} \cdots \oint_{C_d} \frac{dz_d}{(z_d)^N} \\ &\quad \times (z_0)^a (z_1 - z_0)^j \prod_{l=1}^d e(z_{l-1}, z_l) \prod_{l=1}^{d-1} \frac{1}{N z_l (2z_l - z_{l-1} - z_{l+1})} \\ &\quad \times (z_d)^b, \end{aligned} \tag{2.12}$$

where the operation $\frac{1}{2\pi\sqrt{-1}} \oint_{C_i} dz_i$ means taking residues at $z_i = 0$ for $i = 0, d$ and at $z_i = 0, \frac{z_{i-1} + z_{i+1}}{2}$ for $i = 1, \dots, d-1$. We introduce here a notation:

$$w_{j,d} := w(\sigma_j(\mathcal{O}_{h^{N-2-j}})\mathcal{O}_{h^{-1}})_{0,2},$$

for brevity. $e(x, y)$ is given as,

$$e(x, y) = \prod_{r=0}^N (rx + (N-r)y).$$

Note that $e(x, y) = e(y, x)$. For later use, we introduce,

$$\begin{aligned} a_i^{(l)} &:= \frac{1}{i!} \frac{\partial^i}{\partial \epsilon^i} \left(\prod_{r=1}^N (N(l-1) + r + N\epsilon) \right) \Big|_{\epsilon=0} \\ &= \frac{1}{i!} \frac{\partial^i}{\partial \epsilon^i} \left(\prod_{r=N(l-1)+1}^{Nl} (r + N\epsilon) \right) \Big|_{\epsilon=0} \quad (i = 0, 1, 2, \dots; l = 1, \dots, d). \end{aligned}$$

We can easily see,

$$a_i^{(l)} := 0 \quad (i = N+1, N+2, \dots; l = 1, \dots, d).$$

Before turning into the proof of Theorem 1, we prepare several lemmas.

Lemma 1 For $l = 1, \dots, d$, we can express $e(x, y)$ by using $a_i^{(l)}$ ($i = 1, 2, \dots$) as follows.

$$e(x, y) = Nx \sum_{i=1}^{\infty} a_i^{(l)} l^i \left(x - \frac{l-1}{l} y \right)^i (y-x)^{N-i}. \tag{2.13}$$

Proof. Note that $\prod_{r=1}^N (N(l-1) + r + N\epsilon)$ is expanded as,

$$\prod_{r=1}^N (N(l-1) + r + N\epsilon) = \sum_{i=0}^{\infty} a_i^{(l)} \epsilon^i.$$

By using this expression, we can derive (2.13) in the following way.

$$\begin{aligned} e(x, y) &= e(y, x) \\ &= \prod_{r=0}^N (ry + (N-r)x) \\ &= Nx \prod_{r=1}^N (r(y-x) + Nx) \\ &= Nx(y-x)^N \prod_{r=1}^N \left(r + N \frac{x}{y-x} \right) \\ &= Nx(y-x)^N \prod_{r=1}^N \left(N(l-1) + r + N \left(\frac{x}{y-x} - (l-1) \right) \right) \\ &= Nx(y-x)^N \prod_{r=1}^N \left(N(l-1) + r + N \left(\frac{lx - (l-1)y}{y-x} \right) \right) \\ &= Nx(y-x)^N \sum_{i=0}^{\infty} a_i^{(l)} \left(\frac{lx - (l-1)y}{y-x} \right)^i \\ &= Nx \sum_{i=0}^{\infty} a_i^{(l)} (lx - (l-1)y)^i (y-x)^{N-i}. \\ &= Nx \sum_{i=0}^{\infty} a_i^{(l)} l^i \left(x - \frac{l-1}{l} y \right)^i (y-x)^{N-i}. \quad \square \end{aligned}$$

Next, we rewrite the r.h.s. of (1.9).

Lemma 2

$$\frac{1}{j!} \frac{\partial^j}{\partial \epsilon^j} \left(\frac{\prod_{r=1}^{Nd} (r+N\epsilon)}{\prod_{r=1}^d (r+\epsilon)^N} \right) \Big|_{\epsilon=0} = \sum_{\substack{k_1+\dots+k_d=j \\ k_1, \dots, k_d \geq 0}} A_{k_1}^{(1)} \cdots A_{k_d}^{(d)}, \quad (2.14)$$

where

$$A_k^{(l)} := \sum_{i=0}^k a_i^{(l)} \binom{N+k-i-1}{k-i} \cdot \frac{(-1)^{k-i}}{l^{N+k-i}}.$$

Proof. For $l = 1, \dots, d$, we set,

$$F^{(l)}(\epsilon) := \prod_{r=N(l-1)+1}^{Nl} (r+N\epsilon) = \prod_{r=1}^N (N(l-1) + r + N\epsilon) = \sum_{i=0}^{\infty} a_i^{(l)} \epsilon^i,$$

and

$$G^{(l)}(\epsilon) := (l+\epsilon)^{-N}.$$

Then $(\partial^j/\partial\epsilon^j)\{F^{(1)}(\epsilon)G^{(1)}(\epsilon)\cdots F^{(d)}(\epsilon)G^{(d)}(\epsilon)\}|_{\epsilon=0}$ is nothing but the l.h.s. of (2.14). By using the following identity:

$$\begin{aligned} G^{(l)}(\epsilon) &= (l+\epsilon)^{-N} \\ &= \frac{1}{l^N} \left(1 + \frac{\epsilon}{l}\right)^{-N} \\ &= \frac{1}{l^N} \sum_{i=0}^{\infty} \binom{-N}{i} \left(\frac{\epsilon}{l}\right)^i \\ &= \frac{1}{l^N} \sum_{i=0}^{\infty} (-1)^i \binom{N+i-1}{N-1} \left(\frac{\epsilon}{l}\right)^i \\ &= \sum_{i=0}^{\infty} \frac{1}{l^N} \binom{N+i-1}{i} \left(-\frac{1}{l}\right)^i \epsilon^i, \end{aligned}$$

we can rewrite $F^{(l)}(\epsilon)G^{(l)}(\epsilon)$ as,

$$\begin{aligned} F^{(l)}(\epsilon)G^{(l)}(\epsilon) &= \left(\sum_{k=0}^{\infty} a_k^{(l)} \epsilon^k \right) \left(\sum_{k=0}^{\infty} \frac{1}{l^N} \binom{N+k-1}{i} \left(-\frac{1}{l}\right)^k \epsilon^k \right) \\ &= \sum_{k=0}^{\infty} \left(\sum_{i=0}^k a_i^{(l)} \cdot \frac{1}{l^N} \binom{N+k-i-1}{k-i} \left(-\frac{1}{l}\right)^{k-i} \right) \epsilon^k \\ &= \sum_{k=0}^{\infty} \left(\sum_{i=0}^k a_i^{(l)} \binom{N+k-i-1}{k-i} \cdot \frac{(-1)^{k-i}}{l^{N+k-i}} \right) \epsilon^k \\ &= \sum_{k=0}^{\infty} A_k^{(l)} \epsilon^k. \end{aligned}$$

Hence we obtain,

$$\begin{aligned} F^{(1)}(\epsilon)G^{(1)}(\epsilon)\cdots F^{(d)}(\epsilon)G^{(d)}(\epsilon) &= \left(\sum_{k_1=0}^{\infty} A_{k_1}^{(1)} \epsilon^{k_1} \right) \cdots \left(\sum_{k_d=0}^{\infty} A_{k_d}^{(d)} \epsilon^{k_d} \right) \\ &= \sum_{j=0}^{\infty} \left(\sum_{\substack{k_1+\cdots+k_d=j \\ k_1,\dots,k_d \geq 0}} A_{k_1}^{(1)} \cdots A_{k_d}^{(d)} \right) \epsilon^j. \end{aligned}$$

This directly leads us to,

$$\frac{1}{j!} \frac{\partial^j}{\partial\epsilon^j} \left(\frac{\prod_{r=1}^{Nd} (r+N\epsilon)}{\prod_{r=1}^d (r+\epsilon)^N} \right) \Big|_{\epsilon=0} = \sum_{\substack{k_1+\cdots+k_d=j \\ k_1,\dots,k_d \geq 0}} A_{k_1}^{(1)} \cdots A_{k_d}^{(d)}. \quad \square$$

The next two lemmas are useful in evaluating the residue integral in the r.h.s of (2.12),

Lemma 3 *Let α be a non-negative integer. For $l = 0, \dots, d-2$, we have,*

$$\begin{aligned} &\frac{1}{2\pi\sqrt{-1}} \oint_{z_l=\frac{l}{l+1}z_{l+1}} dz_l \frac{(z_{l+1}-z_l)^\alpha}{\left(z_l - \frac{l}{l+1}z_{l+1}\right)^{\alpha+1}} \cdot \frac{e(z_l, z_{l+1})}{Nz_l(2z_{l+1}-z_l-z_{l+2})} \\ &= (z_{l+1})^N \left(\frac{l+1}{l+2}\right)^{\alpha+1} \sum_{s=0}^{\alpha} (l+2)^s A_s^{(l+1)} \frac{(z_{l+2}-z_{l+1})^{\alpha-s}}{\left(z_{l+1} - \frac{l+1}{l+2}z_{l+2}\right)^{\alpha-s+1}}, \end{aligned} \quad (2.15)$$

where the symbol $\frac{1}{2\pi\sqrt{-1}} \oint_{z_l=\frac{l}{l+1}z_{l+1}} dz_l$ means taking a residue at $z_l = \frac{l}{l+1}z_{l+1}$.

Proof. In this proof, we denote by I the integral in the l.h.s of (2.15). By using Lemma 1, the integrand of I is rewritten as follows,

$$\begin{aligned} & \frac{(z_{l+1} - z_l)^\alpha}{\left(z_l - \frac{l}{l+1}z_{l+1}\right)^{\alpha+1}} \cdot \frac{e(z_l, z_{l+1})}{Nz_l(2z_{l+1} - z_l - z_{l+2})} \\ &= \frac{1}{\left(z_l - \frac{l}{l+1}z_{l+1}\right)^{\alpha+1}} \cdot \frac{Nz_l \sum_{i=0}^{\infty} a_i^{(l+1)} (l+1)^i \left(z_l - \frac{l}{l+1}z_{l+1}\right)^i (z_{l+1} - z_l)^{N+\alpha-i}}{Nz_l(2z_{l+1} - z_l - z_{l+2})} \\ &= \frac{1}{\left(z_l - \frac{l}{l+1}z_{l+1}\right)^{\alpha+1}} \cdot \frac{\sum_{i=0}^{\infty} a_i^{(l+1)} (l+1)^i \left(z_l - \frac{l}{l+1}z_{l+1}\right)^i (z_{l+1} - z_l)^{N+\alpha-i}}{2z_{l+1} - z_l - z_{l+2}}. \end{aligned}$$

Since we have,

$$\begin{aligned} & \left. \frac{\sum_{i=0}^{\infty} a_i^{(l+1)} (l+1)^i \left(z_l - \frac{l}{l+1}z_{l+1}\right)^i (z_{l+1} - z_l)^{N+\alpha-i}}{2z_{l+1} - z_l - z_{l+2}} \right|_{z_l=\frac{l}{l+1}z_{l+1}} \\ &= \frac{a_0^{(l+1)} \left(\frac{z_{l+1}}{l+1}\right)^{N+\alpha-i}}{\frac{l+2}{l+1}z_{l+1} - z_{l+2}} \\ &= \frac{\left(\frac{z_{l+1}}{l+1}\right)^{N+\alpha-i}}{\frac{l+2}{l+1}z_{l+1} - z_{l+2}} \cdot \prod_{r=1}^N (Nl + r) \neq 0, \end{aligned}$$

I is explicitly evaluated as follows.

$$\begin{aligned} I &= \frac{1}{\alpha!} \frac{\partial^\alpha}{\partial z_l^\alpha} \left. \frac{\sum_{i=0}^{\infty} a_i^{(l+1)} (l+1)^i \left(z_l - \frac{l}{l+1}z_{l+1}\right)^i (z_{l+1} - z_l)^{N+\alpha-i}}{2z_{l+1} - z_l - z_{l+2}} \right|_{z_l=\frac{l}{l+1}z_{l+1}} \\ &= \frac{1}{\alpha!} \sum_{p=0}^{\alpha} \binom{\alpha}{p} \frac{\partial^p}{\partial z_l^p} \left. \left(\sum_{i=0}^{\infty} a_i^{(l+1)} (l+1)^i \left(z_l - \frac{l}{l+1}z_{l+1}\right)^i (z_{l+1} - z_l)^{N+\alpha-i} \right) \right|_{z_l=\frac{l}{l+1}z_{l+1}} \\ &\quad \cdot \left(\frac{\partial^{\alpha-p}}{\partial z_l^{\alpha-p}} \left. (2z_{l+1} - z_l - z_{l+2})^{-1} \right|_{z_l=\frac{l}{l+1}z_{l+1}} \right) \\ &= \frac{1}{\alpha!} \sum_{p=0}^{\alpha} \binom{\alpha}{p} \\ &\quad \cdot \sum_{i=0}^{\infty} a_i^{(l+1)} (l+1)^i \sum_{q=0}^p \binom{p}{q} \left(\frac{\partial^q}{\partial z_l^q} \left. \left(z_l - \frac{l}{l+1}z_{l+1} \right)^i \right|_{z_l=\frac{l}{l+1}z_{l+1}} \right) \left(\frac{\partial^{p-q}}{\partial z_l^{p-q}} \left. (z_{l+1} - z_l)^{N+\alpha-i} \right|_{z_l=\frac{l}{l+1}z_{l+1}} \right) \\ &\quad \cdot \left(\frac{\partial^{\alpha-p}}{\partial z_l^{\alpha-p}} \left. (2z_{l+1} - z_l - z_{l+2})^{-1} \right|_{z_l=\frac{l}{l+1}z_{l+1}} \right). \end{aligned}$$

By basic calculus, we can easily see,

$$\begin{aligned} \left. \frac{\partial^q}{\partial z_l^q} \right|_{z_l = \frac{l}{l+1} z_{l+1}} \left(z_l - \frac{l}{l+1} z_{l+1} \right)^i &= \frac{i!}{(i-q)!} \delta_0^{i-q}, \\ \left. \frac{\partial^{p-q}}{\partial z_l^{p-q}} \right|_{z_l = \frac{l}{l+1} z_{l+1}} (z_{l+1} - z_l)^{N+\alpha-i} &= (-1)^{p-q} \frac{(N+\alpha-i)!}{(N+\alpha-i-p+q)!} \left(\frac{z_{l+1}}{l+1} \right)^{N+\alpha-i-p+q}, \\ \left. \frac{\partial^{\alpha-p}}{\partial z_l^{\alpha-p}} \right|_{z_l = \frac{l}{l+1} z_{l+1}} (2z_{l+1} - z_l - z_{l+2})^{-1} &= (\alpha-p)! \left(\frac{l+2}{l+1} \right)^{-\alpha+p-1} \left(z_{l+1} - \frac{l+1}{l+2} z_{l+2} \right)^{-\alpha+p-1}, \end{aligned}$$

where δ_0^{i-q} is Kronecker's delta. Therefore, we obtain,

$$I = (z_{l+1})^N \sum_{p=0}^{\alpha} \sum_{i=0}^p a_i^{(l+1)} (l+1)^{-N+i+1} (l+2)^{-\alpha+p-1} (-1)^{p-i} \binom{N+\alpha-i}{p-i} \frac{(z_{l+1})^{\alpha-p}}{\left(z_{l+1} - \frac{l+1}{l+2} z_{l+2} \right)^{\alpha-p+1}}.$$

Now we expand $(z_{l+1})^{\alpha-p}$ in the following form:

$$\begin{aligned} (z_{l+1})^{\alpha-p} &= (l+2)^{\alpha-p} \left(\left(z_{l+1} - \frac{l+1}{l+2} z_{l+2} \right) + \frac{l+1}{l+2} (z_{l+2} - z_{l+1}) \right)^{\alpha-p} \\ &= (l+2)^{\alpha-p} \sum_{q=0}^{\alpha-p} \binom{\alpha-p}{\alpha-p-q} \left(z_{l+1} - \frac{l+1}{l+2} z_{l+2} \right)^{\alpha-p-q} \left(\frac{l+1}{l+2} \right)^q (z_{l+2} - z_{l+1})^q. \end{aligned}$$

Then I is rewritten by,

$$\begin{aligned} I &= (z_{l+1})^N \sum_{p=0}^{\alpha} \sum_{q=0}^{\alpha-p} \sum_{i=0}^p a_i^{(l+1)} (l+1)^{-N+i+q+1} (l+2)^{-q-1} (-1)^{p-i} \binom{N+\alpha-i}{p-i} \binom{\alpha-p}{\alpha-p-q} \\ &\quad \cdot \frac{(z_{l+2} - z_{l+1})^q}{\left(z_{l+1} - \frac{l+1}{l+2} z_{l+2} \right)^{q+1}}. \end{aligned}$$

Note that the indices (p, q, i) runs over the set $\{(p, q, i) \in \mathbb{Z}^3 \mid 0 \leq p \leq \alpha, 0 \leq q \leq \alpha-p, 0 \leq i \leq p\}$. Now we rename the indices as follows:

$$\begin{cases} s = q \\ t = i \\ u = p - i \end{cases}$$

Then (s, t, u) runs over $\{(s, t, u) \in \mathbb{Z}^3 \mid 0 \leq s \leq \alpha, 0 \leq t \leq \alpha-s, 0 \leq u \leq \alpha-s-t\}$. By using the identity,

$$(-1)^{\alpha-s-t-u} \binom{\alpha-t-u}{\alpha-s-t-u} = \binom{-s-1}{\alpha-s-t-u},$$

we can further rewrite I into,

$$\begin{aligned} I &= (z_{l+1})^N \sum_{s=0}^{\alpha} \sum_{t=0}^{\alpha-s} \sum_{u=0}^{\alpha-s-t} a_t^{(l+1)} (l+1)^{-N+s+t+1} (l+2)^{-s-1} (-1)^{\alpha-s-t} \binom{N+\alpha-t}{u} (-1)^{\alpha-s-t-u} \binom{\alpha-t-u}{\alpha-s-t-u} \\ &\quad \cdot \frac{(z_{l+2} - z_{l+1})^s}{\left(z_{l+1} - \frac{l+1}{l+2} z_{l+2} \right)^{s+1}} \end{aligned}$$

$$\begin{aligned}
&= (z_{l+1})^N \sum_{s=0}^{\alpha} \sum_{t=0}^{\alpha-s} a_t^{(l+1)} (l+1)^{-N+s+t+1} (l+2)^{-s-1} (-1)^{\alpha-s-t} \\
&\quad \cdot \left(\sum_{u=0}^{\alpha-s-t} \binom{N+\alpha-t}{u} \binom{-s-1}{\alpha-s-t-u} \right) \\
&\quad \cdot \frac{(z_{l+2} - z_{l+1})^s}{\left(z_{l+1} - \frac{l+1}{l+2} z_{l+2} \right)^{s+1}}.
\end{aligned}$$

Then by using Chu–Vandermonde identity,

$$\sum_{u=0}^{\alpha-s-t} \binom{N+\alpha-t}{u} \binom{-s-1}{\alpha-s-t-u} = \binom{N+\alpha-s-t-1}{\alpha-s-t},$$

we finally reach the r.h.s. of (2.15):

$$\begin{aligned}
I &= (z_{l+1})^N \sum_{s=0}^{\alpha} \sum_{t=0}^{\alpha-s} a_t^{(l+1)} (l+1)^{-N+s+t+1} (l+2)^{-s-1} (-1)^{\alpha-s-t} \binom{N+\alpha-s-t-1}{\alpha-s-t} \\
&\quad \cdot \frac{(z_{l+2} - z_{l+1})^s}{\left(z_{l+1} - \frac{l+1}{l+2} z_{l+2} \right)^{s+1}} \\
&= (z_{l+1})^N \left(\frac{l+1}{l+2} \right)^{\alpha+1} \sum_{s=0}^{\alpha} (l+2)^{\alpha-s} A_{\alpha-s}^{(l+1)} \frac{(z_{l+2} - z_{l+1})^s}{\left(z_{l+1} - \frac{l+1}{l+2} z_{l+2} \right)^{s+1}} \\
&= (z_{l+1})^N \left(\frac{l+1}{l+2} \right)^{\alpha+1} \sum_{s=0}^{\alpha} (l+2)^s A_s^{(l+1)} \frac{(z_{l+2} - z_{l+1})^{\alpha-s}}{\left(z_{l+1} - \frac{l+1}{l+2} z_{l+2} \right)^{\alpha-s+1}}. \square
\end{aligned}$$

Lemma 4

$$\begin{aligned}
&\frac{1}{2\pi\sqrt{-1}} \oint_{z_{d-1}=\frac{d-1}{d}z_d} dz_{d-1} \frac{(z_d - z_{d-1})^\alpha}{\left(z_{d-1} - \frac{d-1}{d} z_d \right)^{\alpha+1}} \cdot \frac{e(z_{d-1}, z_d)}{N z_{d-1}} \\
&= (z_d)^N \cdot d^\alpha \cdot \sum_{l=0}^{\alpha} A_l^{(d)} \cdot (-d)^{-\alpha+l}.
\end{aligned}$$

Proof. The proof goes in a similar manner as the one of Lemma 3 except for use of the relation:

$$\binom{N+a}{N} = \sum_{l=0}^a \binom{N+l-1}{N-1}.$$

□

Proof of Theorem 1.

The integrand in the r.h.s. of (2.12) is given by,

$$\begin{aligned}
&\frac{1}{(z_0)^N \cdots (z_d)^N} \cdot (z_0)^{N-2-j} (z_1 - z_0)^j \cdot \frac{e(z_0, z_1)}{N z_1 (2z_1 - z_0 - z_2)} \cdots \frac{e(z_{d-2}, z_{d-1})}{N z_{d-1} (2z_{d-1} - z_{d-2} - z_d)} \cdot \frac{e(z_{d-1}, z_d)}{z_d} \\
&= \frac{1}{(z_1)^N \cdots (z_d)^N} \cdot \frac{(z_1 - z_0)^j}{(z_0)^{j+1}} \cdot \frac{e(z_0, z_1)}{N z_0 (2z_1 - z_0 - z_2)} \cdots \frac{e(z_{d-2}, z_{d-1})}{N z_{d-2} (2z_{d-1} - z_{d-2} - z_d)} \cdot \frac{e(z_{d-1}, z_d)}{N z_{d-1}} \cdot \frac{N}{z_d}.
\end{aligned}$$

By using Lemma 3, integration over z_0 results in the following integrand.

$$\begin{aligned}
& \frac{1}{(z_1)^N \cdots (z_d)^N} \cdot (z_1)^N \left(\frac{1}{2}\right)^{j+1} \sum_{s_1=0}^j 2^{s_1} A_{s_1}^{(1)} \frac{(z_2 - z_1)^{j-s_1}}{(z_1 - \frac{1}{2}z_2)^{j-s_1+1}} \\
& \cdot \frac{e(z_1, z_2)}{Nz_1(2z_2 - z_1 - z_3)} \cdots \frac{e(z_{d-2}, z_{d-1})}{Nz_{d-2}(2z_{d-1} - z_{d-2} - z_d)} \cdot \frac{e(z_{d-1}, z_d)}{Nz_{d-1}} \cdot \frac{N}{z_d} \\
& = \frac{1}{(z_2)^N \cdots (z_d)^N} \sum_{s_1=0}^j 2^{-j+s_1-1} A_{s_1}^{(1)} \frac{(z_2 - z_1)^{j-s_1}}{(z_1 - \frac{1}{2}z_2)^{j-s_1+1}} \cdot \frac{e(z_1, z_2)}{Nz_1(2z_2 - z_1 - z_3)} \\
& \cdot \frac{e(z_2, z_3)}{Nz_2(2z_3 - z_2 - z_4)} \cdots \frac{e(z_{d-2}, z_{d-1})}{Nz_{d-2}(2z_{d-1} - z_{d-2} - z_d)} \cdot \frac{e(z_{d-1}, z_d)}{Nz_{d-1}} \cdot \frac{N}{z_d}
\end{aligned}$$

Since this integrand is holomorphic at $z_1 = 0$, we only have to take residue at $z_1 = \frac{0+z_2}{2} = \frac{z_2}{2}$. Thus integration over z_1 gives us,

$$\begin{aligned}
& \frac{1}{(z_2)^N \cdots (z_d)^N} \sum_{s_1=0}^j 2^{-j+s_1-1} A_{s_1}^{(1)}(z_2)^N \left(\frac{2}{3}\right)^{j-s_1+1} \sum_{s_2=0}^{j-s_1} 3^{s_2} A_{s_2}^{(2)} \frac{(z_3 - z_2)^{j-s_1-s_2}}{(z_2 - \frac{2}{3}z_3)^{j-s_1-s_2+1}} \\
& \cdot \frac{e(z_2, z_3)}{Nz_2(2z_3 - z_2 - z_4)} \cdots \frac{e(z_{d-2}, z_{d-1})}{Nz_{d-2}(2z_{d-1} - z_{d-2} - z_d)} \cdot \frac{e(z_{d-1}, z_d)}{Nz_{d-1}} \cdot \frac{N}{z_d} \\
& = \frac{1}{(z_3)^N \cdots (z_d)^N} \sum_{s_1=0}^j A_{s_1}^{(1)} \sum_{s_2=0}^{j-s_1} 3^{-j+s_1+s_2-1} A_{s_2}^{(2)} \frac{(z_3 - z_2)^{j-s_1-s_2}}{(z_2 - \frac{2}{3}z_3)^{j-s_1-s_2+1}} \cdot \frac{e(z_2, z_3)}{Nz_2(2z_3 - z_2 - z_4)} \\
& \cdot \frac{e(z_3, z_4)}{Nz_3(2z_4 - z_3 - z_5)} \cdots \frac{e(z_{d-2}, z_{d-1})}{Nz_{d-2}(2z_{d-1} - z_{d-2} - z_d)} \cdot \frac{e(z_{d-1}, z_d)}{Nz_{d-1}} \cdot \frac{N}{z_d}.
\end{aligned}$$

In the same manner, we integrate out z_2, \dots, z_{d-2} . Then we obtain,

$$\begin{aligned}
w_{j,d} &= \frac{1}{(2\pi\sqrt{-1})^2} \oint_{C_{d-1}} dz_{d-1} \oint_{C_d} dz_d \frac{1}{(z_d)^N} \sum_{s_1=0}^j A_{s_1}^{(1)} \sum_{s_2=0}^{j-s_1} A_{s_2}^{(2)} \\
&\quad \cdots \sum_{s_{d-1}=0}^{j-s_1-\cdots-s_{d-2}} d^{-j+s_1+\cdots+s_{d-1}-1} A_{s_{d-1}}^{(d-1)} \frac{(z_d - z_{d-1})^{j-s_1-\cdots-s_{d-1}}}{(z_{d-1} - \frac{d-1}{d}z_d)^{j-s_1-\cdots-s_{d-1}+1}} \cdot \frac{e(z_{d-1}, z_d)}{Nz_{d-1}} \cdot \frac{N}{z_d}.
\end{aligned}$$

Since this integrand is holomorphic at $z_{d-1} = 0$, we only have to take residue at $z_{d-1} = \frac{d-1}{d}z_d$. With the aid of Lemma 4, remaining integration goes as follows.

$$\begin{aligned}
w_{j,d} &= \frac{1}{2\pi\sqrt{-1}} \oint_{C_d} dz_d \sum_{s_1=0}^j A_{s_1}^{(1)} \sum_{s_2=0}^{j-s_1} A_{s_2}^{(2)} \\
&\quad \cdots \sum_{s_{d-1}=0}^{j-s_1-\cdots-s_{d-2}} d^{-j+s_1+\cdots+s_{d-1}-1} A_{s_{d-1}}^{(d-1)} \cdot d^{j-s_1-\cdots-s_{d-1}} \cdot \sum_{s_d=0}^{j-s_1-\cdots-s_{d-1}} A_{s_d}^{(d)} \cdot (-d)^{-j+s_1+\cdots+s_d} \cdot \frac{N}{z_d} \\
&= \frac{N}{d} \sum_{s_1=0}^j A_{s_1}^{(1)} \sum_{s_2=0}^{j-s_1} A_{s_2}^{(2)} \\
&\quad \cdots \sum_{s_{d-1}=0}^{j-s_1-\cdots-s_{d-2}} A_{s_{d-1}}^{(d-1)} \sum_{s_d=0}^{j-s_1-\cdots-s_{d-1}} A_{s_d}^{(d)} \cdot (-d)^{-j+s_1+\cdots+s_d}
\end{aligned}$$

$$\begin{aligned}
&= \frac{N}{d} \left(-\frac{1}{d} \right)^j \sum_{r=0}^j \sum_{\substack{s_1+\dots+s_d=r \\ s_1, \dots, s_d \geq 0}} A_{s_1}^{(1)} \cdots A_{s_d}^{(d)} (-d)^r \\
&= \frac{N}{d} \left(-\frac{1}{d} \right)^j \sum_{r=0}^j B_r (-d)^r,
\end{aligned}$$

where we used Lemma 2 and introduced the notation:

$$B_r := \frac{1}{r!} \frac{\partial^r}{\partial \epsilon^r} \left. \left(\frac{\prod_{l=1}^{Nd} (l + N\epsilon)}{\prod_{l=1}^d (l + \epsilon)^N} \right) \right|_{\epsilon=0}.$$

Therefore, the l.h.s. of (1.9) reduces to,

$$\begin{aligned}
\frac{d}{N} w_{j,d} + \frac{1}{N} w_{j-1,d} &= \left(-\frac{1}{d} \right)^j \sum_{r=0}^j B_r (-d)^r + \frac{1}{d} \left(-\frac{1}{d} \right)^{j-1} \sum_{r=0}^{j-1} B_r (-d)^r \\
&= \left(-\frac{1}{d} \right)^j \sum_{r=0}^j B_r (-d)^r - \left(-\frac{1}{d} \right)^{j-1} \sum_{r=0}^{j-1} B_r (-d)^r \\
&= \left(-\frac{1}{d} \right)^j B_j (-d)^j \\
&= \frac{1}{j!} \frac{\partial^j}{\partial \epsilon^j} \left. \left(\frac{\prod_{r=1}^{Nd} (r + N\epsilon)}{\prod_{r=1}^d (r + \epsilon)^N} \right) \right|_{\epsilon=0},
\end{aligned}$$

which completes the proof of Theorem 1. \square

References

- [1] I.Ciocan-Fontanine, B. Kim. *Wall-crossing in genus zero quasimap theory and mirror maps*. Algebr. Geom. 1 (2014), no. 4, 400–448.
- [2] A. B. Givental. *Equivariant Gromov-Witten invariants*. Internat. Math. Res. Notices 1996, no. 13, 613–663.
- [3] B. R. Greene, D. R. Morrison, M. R. Plesser. *Mirror manifolds in higher dimension*. Comm. Math. Phys. 173 (1995), no. 3, 559–597.
- [4] K. Hori. *Constraints For Topological Strings In $D \geq 1$* . Nucl. Phys. B439 (1995) 395–420.
- [5] M. Jinzenji. *Mirror Map as Generating Function of Intersection Numbers: Toric Manifolds with Two Kähler Forms*, arXiv:1006.0607, Comm. Math. Phys. 323(2013), no. 2, 747–811.
- [6] M. Jinzenji. *Geometrical Proof of Generalized Mirror Transformation of Projective Hypersurfaces*. arXiv:1712.09819, Preprint.
- [7] M. Jinzenji. *Classical Mirror Symmetry*. SpringerBriefs in Mathematical Physics, 29. Springer, Singapore, 2018. viii+140 pp. ISBN: 978-981-13-0055-4; 978-981-13-0056-1 81-01.
- [8] M. Jinzenji, M. Nagura. *Mirror symmetry and an exact calculation of an $(N-2)$ -point correlation function on a Calabi-Yau manifold embedded in CP^{N-1}* . Internat. J. Modern Phys. A 11 (1996), no. 7, 1217–1252.
- [9] M. Jinzenji, M. Shimizu. *Multi-Point Virtual Structure Constants and Mirror Computation of CP^2 -model*. Commun. Number Theory Phys. 7 (2013), no. 3, 411–468.

- [10] M. Jinzenji, H.Saito. *Moduli Space of Quasi-Maps from \mathbb{P}^1 with Two Marked Points to $\mathbb{P}(1, 1, 1, 3)$ and j -invariant.* J. Math. Soc. Japan 73 (2021), no. 4, 995–1018.
- [11] B. Kim. *Private Communication.* Sept. in 2016.
- [12] K. Matsuzaka. *Toric construction and Chow ring of moduli space of quasi maps from \mathbb{P}^1 with two marked points to $\mathbb{P}^1 \times \mathbb{P}^1$.* Internat. J. Math. 31 (2020), no. 12, 2050094, 29 pp.
- [13] H. Saito. *Chow Rings of $\widetilde{Mp}_{0,2}(N, d)$ and $\overline{M}_{0,2}(\mathbb{P}^{N-1}, d)$ and Gromov-Witten Invariants of Projective Hypersurfaces of Degree 1 and 2,* Internat. J. Math. 28(2017), no. 12, 1750090.
- [14] A. Zinger. *The reduced genus-one Gromov-Witten invariants of Calabi-Yau hypersurfaces.* J. Amer. Math. Soc. 22 (2009), 691–737.