Interacting dark energy in curved FLRW spacetime from Weyl Integrable Spacetime

S. Chatzidakis,¹ A. Giacomini,^{2,*} P.G.L. Leach,³ G.

Leon,^{4,3,†} A. Paliathanasis,^{2,3,‡} and Supriya Pan^{5,§}

¹Applied Technology High School, PO Box 66866, Al Ain, United Arab Emirates ²Instituto de Ciencias Físicas y Matemáticas, Universidad Austral de Chile, Valdivia 5090000, Chile

³Institute of Systems Science, Durban University of Technology, PO Box 1334, Durban 4000, South Africa

⁴Departamento de Matemáticas, Universidad Católica del Norte, Avda. Angamos 0610, Casilla 1280 Antofagasta, Chile ⁵Department of Mathematics. Presidency University.

86/1 College Street, Kolkata 700073, India

(Dated: June 15, 2022)

In the present article, we show that a simple modification to the Einstein-Hilbert action can explain the possibility of mutual interaction between the cosmic fluids. That is achieved considering the Weyl Integrable Spacetime in the background of a nonflat Friedmann-Lemnaître-Robertson-Walker geometry for the universe. We show that widely-known phenomenological interacting cosmological scenarios can naturally appear in this context. We then performed the dynamical system analysis of the underlying cosmological scenario and explored many possibilities extracted from this gravitational theory.

PACS numbers: 98.80.-k, 95.35.+d, 95.36.+x

Keywords: Cosmology; Weyl integrable space; scalar field; dynamical analysis

^{*}Electronic address: alexgiacomini@uach.cl

[†]Electronic address: genly.leon@ucn.cl

[‡]Electronic address: anpaliat@phys.uoa.gr

[§]Electronic address: supriya.maths@presiuniv.ac.in

1. INTRODUCTION

The detection of the late-time accelerating phase of the universe |48, 49| is one of the significant discoveries in cosmology which abruptly changed our ideas on the evolution of our universe. To describe this accelerating expansion, usually two well-known approaches are considered. One is the introduction of some hypothetical dark energy (DE) fluid in Einstein's General Relativity (GR), and the other is to either modify the Einstein's GR or introduce new gravitational theories. In the latter approach, the additional (new) geometrical terms appearing from the modified (new) gravitational theory mimic the role of a DE. Following both the approaches, several cosmological models have been proposed in the last several years to explain the dynamics of the universe [12, 14, 16, 18, 40, 41, 54]. Among these models, the Λ -Cold Dark Matter (Λ CDM) cosmological model, designed in the context of GR, has been quite successful in describing a large span of astronomical and cosmological data. However, the physics of the primary two ingredients in the ΛCDM paradigm, namely, the cold dark matter and cosmological constant is not yet clearly understood despite many astronomical missions. Apart from that, the cosmological constant problem [58], and the coincidence problem [59] require explanation in this cosmological scenario. These are not the only limitations in this successful cosmological model. It has been consistently observed that some key cosmological parameters estimated from the early time measurements by Planck within this minimal ACDM cosmology [3] significantly differ from other astronomical missions [1, 32, 33, 35, 50]. The most statistically significant discrepancy (at 5σ) is the Hubble constant tension between Planck [3] and Supernovae H_0 for the Equation of State (SH0ES) collaboration [50] which demands a revision of the ACDM cosmology. On the other hand, the tension in the growth of structures, quantified through the parameter S_8 , defined as a combination of the amplitude of the matter power spectrum σ_8 with the matter density at present Ω_m $(S_8 = \sigma_8 \sqrt{\Omega_m/0.3})$ between Planck and cosmic shear measurements [1, 32, 33, 35] has been another issue which has further strengthened the revision of the ACDM cosmology. The cosmological discrepancies have been one of the hot topics at present moment which fueled the scientific community to find an alternative to the ACDM cosmology, see Refs. [2, 21, 22, 25, 47] which include a variety of cosmological models attempting to explain these discrepancies. However, despite many attempts performed by a large number of investigators (see again [2, 21, 22, 25, 47]), the tension free cosmological model is yet to be discovered. In order to be more transparent in this direction, it should be mentioned that the simultaneous solutions to both the tensions is hard to achieve. Due to the existing correlation between H_0 and S_8 , the solution of the H_0 tension in a specific cosmological model may worsen the tension in the S_8 parameter in the same cosmological model. Moreover, the solution to the Hubble H_0 tension in a cosmological model can increase the tension in the sound horizon r_d derived from the cosmic microwave background, and other cosmological probes [6]. As the full cosmic picture is yet to be discovered, there is certainly no reason to favor any cosmological model over the others. Motivated by this, in the present work we investigate a cosmological dynamics which arises from a generalized Einstein-Hilbert action, and interestingly, this offers a novel route to reproduce the interacting DM-DE models without any phenomenological basis¹.

In this work we consider the cosmological dynamics in the Weyl Integrable geometry, a generalization of the Einstein-Hilbert action. We note that the Weyl Integrable spacetime did not get much attention in the literature without any specific reason (only a few works discussing the astrophysical and cosmological consequences are available [5, 28, 42-44, 51-53). While it is interesting to notice that within this cosmological domain, one can mimic the late-time cosmic acceleration [4, 39] and also the early inflationary phase [26]. Here we have explicitly found here, the Weyl Integrable geometry has a very novel feature which enlightens the cosmology of interacting DM-DE models by offering a field theoretical prescription. It is widely known that the theory of interacting dark matter - dark energy (known as interacting dark energy or coupled dark energy) has gained significant attention for solving several cosmological problems, such as the cosmic coincidence problem (see Refs. [8, 19, 20, 34]), cosmological tensions (see Ref. [25]). Here we explicitly show that the Weyl Integrable geometry can offer a solid foundation to the theory of interacting dark energy. The interacting dark energy models are mostly governed by some phenomenological choices of the interaction function, which modifies the expansion history at the background and perturbation levels. Although attempts to construct the interaction function from the action formalisms have been considered in the literature [10, 11, 17, 29, 45, 56], nevertheless, we are not done yet. In the present work, we show that the dark sectors' interaction naturally arises

¹ Let us note that the interacting DM-DE models are very promising to alleviate the Hubble tension, see Tables B1 and B2 of [25].

in the context of Weyl Integrable spacetime quite very naturally without any mathematical complexities. This result is one of the exciting outcomes of this work. Having this novel feature, we performed the dynamical system analysis of the underlying cosmological scenario in the background of a nonflat Friedmann-Lemaître-Robertson-Walker (FLRW) line element. The assumption of the nonflat FLRW universe model is not an artist's imagination but rather the analyses of the cosmic microwave background anisotropies from Planck (considering the Pilk likelihood) favour a closed universe at several standard deviations [3, 23, 24, 31].

Interactive matter-scalar field schemes have been explored before where the conservation equations have the structure

$$\dot{\rho}_m + 3H\left(\rho_m + p_m\right) = Q, \quad \dot{\phi}\left[\ddot{\phi} + 3H\dot{\phi} + V'(\phi)\right] = -Q, \tag{1}$$

where a dot means derivative with respect to cosmic time t, comma derivative with respect to ϕ , ρ_m is the energy density of matter, ϕ is the scalar field, $V(\phi)$ its potential, Q is the interaction term, and $H = \dot{a}/a = \theta/3$ stands for the *Hubble parameter* (which is a general measure of the isotropic rate of spatial expansion), where a denotes the scale factor of the universe.

In Ref. [37] such a scalar field cosmology with a generalized harmonic potential was investigated in flat and negatively curved Friedmann-Lemaître-Robertson-Walker and Bianchi I metrics. An interaction between the scalar field and matter is considered in the form $Q = -\mu/2\rho_m \dot{\phi}$. Asymptotic methods and averaging theory were used to obtain relevant information about the solution space. In this approach, the Hubble parameter played the role of a time-dependent perturbation parameter which controls the magnitude of the error between full-system and time-averaged solutions as it decreases. This approach shows that full and time-averaged systems have the same asymptotic behaviour. Numerical simulations were presented as evidence of such behaviour. Moreover, the asymptotic behaviour of the solutions is independent of the coupling function.

Interactive schemes like (1) naturally appears in Scalar Tensor theory (STT) of gravity. Say, a general class of STTs, written in the so-called Einstein frame (EF), which is given by [36]

$$S_{EF} = \int_{M_4} d^4x \sqrt{|g|} \left\{ \frac{1}{2} R - \frac{1}{2} g^{\mu\nu} \nabla_\mu \phi \nabla_\nu \phi - V(\phi) + \chi(\phi)^{-2} \mathcal{L}_{\text{matter}}(\mu, \nabla \mu, \chi(\phi)^{-1} g_{\alpha\beta}) \right\},\tag{2}$$

where R is the curvature scalar, ϕ is the scalar field, ∇_{α} is the covariant derivative, $V(\phi)$ is the quintessence self-interacting potential, $\chi(\phi)^{-2}$ is the coupling function, $\mathcal{L}_{\text{matter}}$ is the matter Lagrangian, and μ is a collective name for the matter degrees of freedom, repeated indexes mean sum over them. The energy-momentum tensor of matter is defined by

$$T_{\alpha\beta} = -\frac{2}{\sqrt{|g|}} \frac{\delta}{\delta g^{\alpha\beta}} \left\{ \sqrt{|g|} \chi^{-2}(\phi) \mathcal{L}(\mu, \nabla \mu, \chi^{-1}(\phi) g_{\alpha\beta}) \right\}.$$
 (3)

By considering the conformal transformation $\overline{g}_{\alpha\beta} = \chi(\phi)^{-1}g_{\alpha\beta}$, defining the Brans-Dicke (BD) coupling "constant" $\omega(\chi)$ in such way that $d\phi = \pm \sqrt{\omega(\chi) + 3/2}\chi^{-1}d\chi$ and recalling $\overline{V}(\chi) = \chi^2 V(\phi(\chi))$, the action (2) can be written in the Jordan frame (JF) as [15]

$$S_{JF} = \int_{M_4} d^4 x \sqrt{|\overline{g}|} \left\{ \frac{1}{2} \chi \overline{R} - \frac{1}{2} \frac{\omega(\chi)}{\chi} (\overline{\nabla}\chi)^2 - \overline{V}(\chi) + \mathcal{L}_{\text{matter}}(\mu, \nabla\mu, \overline{g}_{\alpha\beta}) \right\}.$$
(4)

Here the bar is used to denote geometrical objects defined with respect to the metric $\overline{g}_{\alpha\beta}$. In the STT given by (4), the energy-momentum of the matter fields,

$$\overline{T}_{\alpha\beta} = -\frac{2}{\sqrt{|\overline{g}|}} \frac{\delta}{\delta \overline{g}^{\alpha\beta}} \left\{ \sqrt{|\overline{g}|} \mathcal{L}(\mu, \nabla \mu, \overline{g}_{\alpha\beta}) \right\},\tag{5}$$

is separately conserved. That is $\overline{\nabla}^{\alpha} \overline{T}_{\alpha\beta} = 0$. However, when is written in the EF (2), with a matter energy-momentum tensor given by (3), this is no longer the case (although the overall energy density is conserved). In fact in the EF we find that

$$Q_{\beta} \equiv \nabla^{\alpha} T_{\alpha\beta} = -\frac{1}{2} T \chi(\phi)^{-1} \frac{\mathrm{d}\chi(\phi)}{\mathrm{d}\phi} \nabla_{\beta}\phi, \quad T = T_{\alpha}^{\alpha}.$$
 (6)

For action (2), the strength of the coupling between the perfect fluid and the scalar field is $Q = \frac{1}{2}(4 - 3\gamma)\rho_m \dot{\phi} \frac{d\ln\chi(\phi)}{d\phi}$, where $\chi(\phi)$ is an input function. In reference [9] the interaction terms (in the flat FLRW geometry) $Q = \alpha \dot{\phi} \rho_m$ and $Q = \alpha \rho_m H$ were investigated, here α is a constant, ϕ is the scalar field, ρ is the energy density of background matter and H is the Hubble parameter. The first choice corresponds to an exponential coupling function $\chi(\phi) = \chi_0 \exp(2\alpha\phi/(4 - 3\gamma))$. The second case corresponds to the choice $\chi = \chi_0 a^{-2\alpha/(4-3\gamma)}$ (and then, $\rho \propto a^{\alpha-3\gamma}$). Other models within the general setup (1), incorporates effective interaction terms $Q = 3\alpha H \rho_m$, $Q = 3\beta H \rho_\phi$ and $Q = 3H(\alpha\rho_m + \beta\rho_\phi)$ [13, 38].

The article has been organized as follows. In section 2, we provide the gravitational equations of the Weyl Integrable spacetime in the FLRW universe model. In section 3 we perform the dynamical system analysis of the cosmological scenario considering the spatial curvature of the universe. Finally, in section 4 we describe the main findings of this article in brief.

2. WEYL INTEGRABLE SPACETIME

In Weyl Integrable Spacetime (WIST) or Weyl Integrable Geometry (WIG), the Einstein-Hilbert Action is modified as

$$S_W = \int dx^4 \sqrt{-g} \left(\widetilde{R} + \xi \left(\widetilde{\nabla}_{\nu} \left(\widetilde{\nabla}_{\mu} \phi \right) \right) g^{\mu\nu} - V \left(\phi \right) + \mathcal{L}_m \right), \tag{7}$$

in which $g_{\mu\nu}$ is the metric tensor for the physical space, $\widetilde{\nabla}_{\mu}$ denotes covariant derivative defind by the symbols $\widetilde{\Gamma}^{\kappa}_{\mu\nu}$, where $\widetilde{\Gamma}^{\kappa}_{\mu\nu}$ are the Christoffel symbols for the conformally related metric $\widetilde{g}_{\mu\nu} = \phi g_{\mu\nu}$. Parameter ξ is an arbitrary coupling constant, and \mathcal{L}_m is the Lagrangian function for the matter source.

When \mathcal{L}_m describes a perfect fluid with energy density ρ and pressure component p, the field equations in the Einstein-Weyl theory are derived to be

$$\widetilde{G}_{\mu\nu} + \widetilde{\nabla}_{\nu} \left(\widetilde{\nabla}_{\mu} \phi \right) - (2\xi - 1) \left(\widetilde{\nabla}_{\mu} \phi \right) \left(\widetilde{\nabla}_{\nu} \phi \right) + \xi g_{\mu\nu} g^{\kappa\lambda} \left(\widetilde{\nabla}_{\kappa} \phi \right) \left(\widetilde{\nabla}_{\lambda} \phi \right) - V \left(\phi \right) g_{\mu\nu}$$

$$= - \left(\widetilde{\rho} + \widetilde{p} \right) u_{\mu} u_{\nu} - \widetilde{p} g_{\mu\nu},$$
(8)

where $\widetilde{G}_{\mu\nu}$ is the Einstein tensor with respect the metric $\widetilde{g}_{\mu\nu}$ and u^{μ} is the comovig observer. The new parameters $\widetilde{\rho}$, \widetilde{p} are the energy density and pressure components for the matter source multiplied by the factor $e^{-\frac{\phi}{2}}$, that means $(\widetilde{\rho}, \widetilde{p}) = (e^{-\frac{\phi}{2}}\rho_m, e^{-\frac{\phi}{2}}p_m)$.

The field equations (8) can be written in the equivalent form

$$G_{\mu\nu} - \lambda \left(\phi_{,\mu} \phi_{,\nu} - \frac{1}{2} g_{\mu\nu} \phi^{,\kappa} \phi_{,\kappa} \right) - V \left(\phi \right) g_{\mu\nu} = -\left(\widetilde{\rho} + \widetilde{p} \right) u_{\mu} u_{\nu} - \widetilde{p} g_{\mu\nu} \tag{9}$$

where $G_{\mu\nu}$ is the Einstein tensor for the background space $g_{\mu\nu}$ and $\lambda = 2\xi - \frac{3}{2}$.

Now, to proceed with the cosmological dynamics, we consider a homogeneous and isotropic universe characterized by the Friedmann-Lemaître-Robertson-Walker (FLRW) line element

$$ds^{2} = -dt^{2} + a^{2}(t) \left[\frac{dr^{2}}{1 - Kr^{2}} + r^{2} \left(d\theta^{2} + \sin^{2}\theta d\varphi^{2} \right] \right),$$
(10)

where a(t) is the expansion scale factor of the universe and K is the curvature scalar of the universe which can take any of the values of $\{0, +1, -1\}$. For K = 0, +1, -1, spatially flat, closed and open geometries of the universe are represented. For the above FLRW metric, with lapse function N = 1, one can write down the Friedmann equations for dust matter $(p_m = 0)$ as follows

$$\left(\frac{\theta^2}{3} + \frac{3K}{a^2}\right) - \frac{\lambda}{2}\dot{\phi}^2 - V(\phi) - e^{-\frac{\phi}{2}}\rho_m = 0,$$
(11)

$$\dot{\theta} + \frac{\theta^2}{2} + \frac{3K}{2a^2} - \frac{3}{2}V(\phi) + \frac{3}{4}\lambda\dot{\phi}^2 = 0,$$
(12)

$$\ddot{\phi} + \theta \dot{\phi} + \frac{1}{\lambda} V_{,\phi} + \frac{1}{2\lambda} e^{-\frac{\phi}{2}} \rho_m = 0, \qquad (13)$$

$$\dot{\rho_m} + \theta \rho_m - \rho_m \dot{\phi} = 0. \tag{14}$$

where we have considered the comoving observer $u^{\mu} = \delta_t^{\mu}$, an overhead dot represents the cosmic time differentiation and $\theta = 3\dot{a}/a$ is the expansion rate of the FLRW universe. The conservation equations for the scalar field and the matter sector are of special importance in this context. The eqns. (13) and (14) can be alternatively expressed as

$$\dot{\rho}_{\phi} + \theta(\rho_{\phi} + p_{\phi}) = -\frac{1}{2} \,\widetilde{\rho} \,\dot{\phi},\tag{15}$$

$$\dot{\widetilde{\rho}} + \theta(\widetilde{\rho} + \widetilde{p}) = \frac{1}{2} \,\widetilde{\rho} \,\dot{\phi},\tag{16}$$

where $\rho_{\phi} = \frac{\lambda}{2} (\dot{\phi}^2 + V(\phi)), p_{\phi} = \frac{\lambda}{2} (\dot{\phi}^2 - V(\phi))$ and $\tilde{p} = 0$. One can clearly see that the above two equations represent an interacting two-fluid system: $\nabla_{\nu} T_{\phi}^{\mu\nu} = -Q(t) = -\nabla_{\nu} T_{\text{matter}}^{\mu\nu}$ where the interaction function Q(t) is given by

$$Q(t) = \frac{1}{2}\tilde{\rho}(t)\,\dot{\phi}(t).\tag{17}$$

Eqs. (15)-(16) realizes such an interactive scheme like (1), with $p_m = 0$ (dust matter). It is a very intriguing result in this context because the above formalism naturally induces an interaction between the scalar field and the matter field without mathematical complexities. We note that in General Relativity, usually one needs to introduce the interaction between the dark matter and dark energy by hand, but here in the WIST, one can see that an interaction term Q is already present. It is one of the essences of WIST where we naturally have an interaction term between the scalar field and the matter sector. We further note that the above interaction may allow a changeable sign property depending on the sign of $\dot{\phi}$. We refer to Refs. [7, 27, 30, 46, 55, 57] where the sign changeable interaction functions have been investigated from the phenomenological ground.

In the line of cosmological parameters, the equation of state parameter for the effective fluid is defined to be $w_{tot}(t) := \frac{p_{tot}(t)}{\rho_{tot}(t)}$, and it is related to the deceleration parameter q(t), in the presence of curvature, by

$$q(t) := -1 - \frac{3\dot{\theta}(t)}{\theta(t)^2} = \frac{(1+3w_{tot}(t))}{2} \left(1 + \frac{9K}{a(t)\theta(t)^2}\right).$$
(18)

Since in this study we consider that the matter source is that of the dust fluid, i.e $p_m = 0$, from equation (14), it follows $\rho_m = \rho_{m0}a^{-3}e^{\phi}$, thus we end with the set of differential equations (11), (12) and (13).

In the following section we present a detailed analysis of the dynamics of the field equations (11), (12), (13) and (14) by investigating the stationary points and their stability in new dimensionless variables. Such an analysis is essential to understand further the effects of the curvature term in the cosmological dynamics of the WIST.

3. DYNAMICAL ANALYSIS

We consider the dimensionless dependent variables $\{x, y, \omega_m, \eta, \Omega_K\}$ defined as

$$\dot{\phi} = \sqrt{2\left(1 + \frac{\theta^2}{3}\right)}x,\tag{19}$$

$$V(\phi) = \left(1 + \frac{\theta^2}{3}\right) y^2 , \qquad (20)$$

$$\rho_m = \left(1 + \frac{\theta^2}{3}\right) e^{\frac{\phi}{2}} \Omega_m , \qquad (21)$$

$$K = \left(1 + \frac{\theta^2}{3}\right) a^2 \Omega_K,\tag{22}$$

$$\theta = \frac{\sqrt{3\eta}}{\sqrt{1-\eta^2}},\tag{23}$$

with inverse

$$x = \frac{\dot{\phi}}{\sqrt{2}D},\tag{24}$$

$$y = \frac{\sqrt{V(\phi)}}{D},\tag{25}$$

$$\Omega_m = \frac{e^{-\frac{\phi}{2}}\rho_m}{D^2},\tag{26}$$

$$\Omega_K = \frac{K}{a^2 D^2}, \quad D = \sqrt{1 + \theta^2/3}$$
(27)

$$\eta = \frac{\theta/\sqrt{3}}{\sqrt{1+\theta^2/3}},\tag{28}$$

and the new independent variable $dt = \sqrt{1 + \frac{\theta^2}{3}} d\tau$. As $\eta \to \pm 1$ we have $\theta \to \pm \infty$. In the new variables the field equations are described by the following algebraic-differential system

$$\Omega_K - \frac{1}{3} \left(\lambda x^2 + y^2 - \eta^2 + \Omega_m \right) = 0$$
(29)

with differential equations

$$\frac{dx}{d\tau} = \frac{1}{12} \left(8\sqrt{3}\eta \lambda x^3 + 2\sqrt{3}\eta x \left(2\eta^2 - 2y^2 + \Omega_m - 6 \right) - \frac{3\sqrt{2} \left(2\mu y^2 + \Omega_m \right)}{\lambda} \right), \quad (30)$$

$$\frac{dy}{d\tau} = \frac{y}{6} \left(\sqrt{3}\eta \left(2\eta^2 + 4\lambda x^2 - 2y^2 + \Omega_m \right) + 3\sqrt{2}\mu x \right), \tag{31}$$

$$\frac{d\Omega_m}{d\tau} = \frac{1}{6} \Omega_m \left(8\sqrt{3}\eta \lambda x^2 + 3\sqrt{2}x + 2\sqrt{3}\eta \left(2\eta^2 - 2y^2 + \Omega_m - 3 \right) \right), \tag{32}$$

$$\frac{d\eta}{d\tau} = \frac{\sqrt{3}}{6} \left(\eta^2 - 1\right) \left(2\eta^2 + 4\lambda x^2 - 2y^2 + \Omega_m\right).$$
(33)

The new variable $\mu = V_{,\phi}/V$. In the following we shall consider that $V(\phi)$ is the exponential potential such that $\mu = \text{const.}$

We remark that under the discrete transformation $y \to -y$ the dynamical system (29)-(33) is invariant. Thus, without any loss of generality we focus our analysis in the branch $y \ge 0$, while by definition $\Omega_m \ge 0$ and $\eta^2 \le 1$. Moreover, in the new variables the effective equation of state parameter w_{tot} and the deceleration parameters read

$$w_{tot}\left(x, y, \Omega_m, \eta\right) = \frac{\lambda x^2 - y^2}{\lambda x^2 + y^2 + \Omega_m},\tag{34}$$

and

$$q(x, y, \Omega_m, \eta) = \frac{4\lambda x^2 - 2y^2 + \Omega_m}{2\eta^2}.$$
(35)

The stationary points of the dynamical system (29)-(33) describe exact asymptotic solutions for the field equations. Indeed from (34) we can determine explicitly the scale factor at every point $P = (x(P), y(P), \Omega_m(P), \eta(P))$. Indeed for $w_{tot} \neq -1$, the exact solution for the scale factor at a given point is $a(t) = a_0 t^{\frac{2}{3(1+w_{tot})}}$, while for $w_{tot} = -1$ we calculate $a(t) = a_0 e^{\theta_0 t}$.

We proceed with our analysis to investigate the stationary points for the field equations and study their stability properties. Such analysis is essential to understand the evolution of the physical variables and construct the cosmological history.

3.1. Stationary points

The stationary points for the dynamical system (29)-(33) can be categorized in the following families of points $P = (x(P), y(P), \Omega_m(P), \eta(P))$.

3.1.1. Points
$$P_1^{\pm}$$

The stationary points P_1^{\pm} with coordinates

$$P_1^{\pm} = \left(\frac{1}{\sqrt{\lambda}}, 0, 0, \pm 1\right). \tag{36}$$

describe asymptotic solutions where only the kinetic component of the scalar field potential contributes to the total cosmological solution. The stationary points are real and physically accepted when $\lambda > 0$. Moreover we derive $\Omega_K(P_1^{\pm}) = 0$, $q(P_1^{\pm}) = 2$ and $w_{tot}(P_1^{\pm}) = 1$. Hence the background space is that of the spatially flat FLRW space with scale factor $a(t) = a_0 t^{\frac{1}{3}}$.

In order to investigate the stability of the stationary points we derive the eigenvalues of the following matrix around the stationary points

$$\mathbf{A} = \begin{pmatrix} \frac{\partial}{\partial x} \begin{pmatrix} dx \\ d\tau \end{pmatrix} & \frac{\partial}{\partial y} \begin{pmatrix} dx \\ d\tau \end{pmatrix} & \frac{\partial}{\partial \Omega_m} \begin{pmatrix} dx \\ d\tau \end{pmatrix} & \frac{\partial}{\partial \eta} \begin{pmatrix} dx \\ d\tau \end{pmatrix} \\ \frac{\partial}{\partial x} \begin{pmatrix} dy \\ d\tau \end{pmatrix} & \frac{\partial}{\partial y} \begin{pmatrix} dy \\ d\tau \end{pmatrix} & \frac{\partial}{\partial \Omega_m} \begin{pmatrix} dy \\ d\tau \end{pmatrix} & \frac{\partial}{\partial \eta} \begin{pmatrix} dy \\ d\tau \end{pmatrix} \\ \frac{\partial}{\partial x} \begin{pmatrix} d\Omega_m \\ d\tau \end{pmatrix} & \frac{\partial}{\partial y} \begin{pmatrix} d\Omega_m \\ d\tau \end{pmatrix} & \frac{\partial}{\partial \Omega_m} \begin{pmatrix} d\Omega_m \\ d\tau \end{pmatrix} & \frac{\partial}{\partial \eta} \begin{pmatrix} d\Omega_m \\ d\tau \end{pmatrix} \\ \frac{\partial}{\partial x} \begin{pmatrix} d\eta \\ d\tau \end{pmatrix} & \frac{\partial}{\partial y} \begin{pmatrix} d\eta \\ d\tau \end{pmatrix} & \frac{\partial}{\partial \Omega_m} \begin{pmatrix} d\eta \\ d\tau \end{pmatrix} & \frac{\partial}{\partial \eta} \begin{pmatrix} d\eta \\ d\tau \end{pmatrix} \end{pmatrix}.$$
(37)

Indeed for the point P_1^+ the eigenvalues of the matrix **A** read

$$e_1(P_1^+) = \frac{4}{\sqrt{3}}, e_1(P_1^+) = 6\sqrt{3}, e_3(P_1^+) = \frac{\sqrt{2} + 2\sqrt{3\lambda}}{2\sqrt{\lambda}}, e_4(P_1^+) = \sqrt{3} + \frac{\mu}{\sqrt{2\lambda}}.$$
 (38)

Because $e_1(P_1^+)$, $e_2(P_1^+)$ and $e_3(P_1^+)$ have always poisitive real parts, the stationary point can not be an attractor which means that the asymptotic solution at the point is unstable. Specifically for $\mu > -\sqrt{6\lambda}$, P_1^+ is a source while for $\mu < \sqrt{6\lambda}$, P_1^+ is a saddle point.

In a similar way the eigenvalues for the stationary point P_1^- are

$$e_1(P_1^-) = -\frac{4}{\sqrt{3}}, e_1(P_1^-) = -6\sqrt{3}, e_3(P_1^-) = \frac{\sqrt{2} - 2\sqrt{3\lambda}}{2\sqrt{\lambda}}, e_4(P_1^-) = -\sqrt{3} + \frac{\mu}{\sqrt{2\lambda}}.$$
 (39)

Thus for $\{\lambda > \frac{1}{6}, \mu \leq 1\}$ and $\{\mu > 0, \lambda > \frac{\mu^2}{6}\}$ all the eigenvalues have negative real parts from where we can infer that P_1^- is an attractor while the asymptotic solution at the specific point is a stable solution.

3.1.2. Points P_2^{\pm}

The family of points P_2^{\pm} is consisted by the stationary points

$$P_2^{\pm} = \left(-\frac{1}{\sqrt{\lambda}}, 0, 0, \pm 1\right). \tag{40}$$

The physical properties of the asymptotic solution at the points P_2^{\pm} are similar to that of the points P_1^{\pm} ; thus, we omit the discussion. We proceed with the derivation of the eigenvalues for the linearized matrix and the investigation of the stability properties.

Around the stationary point P_2^+ the eigenvalues of the linearized system are

$$e_1(P_2^+) = \frac{4}{\sqrt{3}}, e_2(P_2^+) = 6\sqrt{3}, e_3(P_2^+) = \frac{-\sqrt{2} + 2\sqrt{3\lambda}}{2\sqrt{\lambda}}, e_4(P_2^+) = \sqrt{3} - \frac{\mu}{\sqrt{2\lambda}}, \quad (41)$$

from where we infer that when $0 < \lambda < \frac{1}{6}$ or $\mu > 0$, $0 < \lambda < \frac{\mu^2}{6}$ point P_1^+ is a saddle point, otherwise is a source.

For point P_2^- we derive the eigenvalues

$$e_1(P_2^-) = -\frac{4}{\sqrt{3}}, e_2(P_2^-) = -6\sqrt{3}, e_3(P_2^-) = -\frac{\sqrt{2} + 2\sqrt{3\lambda}}{2\sqrt{\lambda}}, e_4(P_2^-) = -\sqrt{3} - \frac{\mu}{\sqrt{2\lambda}}, \quad (42)$$

from where we conclude that when $\{\lambda > 0, \mu < 0\}$, or $\{\mu > 0, \lambda > \frac{\mu^2}{6}\}$ all the eigenvalues have negative real parts from where we infer that point P_2^- is an attractor.

3.1.3. Points
$$P_3^{\pm}$$

Points P_3^{\pm} with coordinates

$$P_3^{\pm} = \left(-\frac{\sqrt{6}}{\mu}, \frac{\sqrt{2(6\lambda - \mu^2)}}{\mu}, 0, \pm 1\right)$$
(43)

describe asymptotic exact solutions where both the kinetic part and the potential terms contributes to the cosmological fluid. The points are physically accepted when $6\lambda \ge \mu^2$, which means that $\lambda > 0$. The effective parameter for the equation of state is $w_{tot} \left(P_3^{\pm}\right) = 1$, while the deceleration parameter is derived $q \left(P_3^{\pm}\right) = 4 \left(-1 + \frac{6\lambda}{\mu^2}\right)$. Furthermore, we derive $\Omega_K \left(P_3^{\pm}\right) = -1 + \frac{6\lambda}{\mu^2}$. Consequently, the stationary points when exist describe exact solutions of FLRW universe with positive curvature, $\Omega_K \left(P_3^{\pm}\right) > 0$, while always $q \left(P_3^{\pm}\right) > 0$. The corresponding eigenvalues are

$$e_1(P_3^+) = 6\sqrt{3}, e_2(P_3^+) = \sqrt{3}\frac{(\mu - 1)}{\mu},$$
(44)

$$e_{3}\left(P_{3}^{+}\right) = \frac{2}{\sqrt{3}}\left(1 - \frac{\sqrt{2\left(9\lambda - \mu^{2}\right)}}{\mu}\right), \ e_{4}\left(P_{3}^{+}\right) = \frac{2}{\sqrt{3}}\left(1 + \frac{\sqrt{2\left(9\lambda - \mu^{2}\right)}}{\mu}\right), \tag{45}$$

and

$$e_1(P_3^-) = -6\sqrt{3}, e_2(P_3^-) = -\sqrt{3}\frac{(\mu - 1)}{\mu},$$
(46)

$$e_{3}\left(P_{3}^{-}\right) = -\frac{2}{\sqrt{3}}\left(1 - \frac{\sqrt{2\left(9\lambda - \mu^{2}\right)}}{\mu}\right), \ e_{4}\left(P_{3}^{-}\right) = -\frac{2}{\sqrt{3}}\left(1 + \frac{\sqrt{2\left(9\lambda - \mu^{2}\right)}}{\mu}\right), \quad (47)$$

We conclude that points P_3^\pm are always saddle points when they exist.

3.1.4. Points P_4^{\pm}

The stationary points which form the family of points P_4^{\pm} have coordinates

$$P_4^{\pm} = \left(\pm\sqrt{\frac{2}{3}}, 0, -\frac{8}{3}\lambda, \pm 1\right).$$
(48)

The points are physically accepted for $\lambda < 0$. The asymptotic solutions at the points describe FLRW spacetimes with curvature $\Omega_K \left(P_4^{\pm} \right) = -\frac{(1+2\lambda)}{3}$, that is $-\frac{1}{2} < \lambda < 0$, $\Omega_K \left(P_4^{\pm} \right) < 0$ otherwise $\Omega_K \left(P_4^{\pm} \right) > 0$. The deceleration parameter and the effective equation of state parameter are calculated $q \left(P_4^{\pm} \right) = 0$ and $w_{tot} \left(P_4^{\pm} \right) = -\frac{1}{3}$.

As far as the stability properties of the stationary points are concerned, we derive the eigenvalues for the linearized system near to the points, they are

$$e_1\left(P_4^+\right) = 2\sqrt{3}, e_2\left(P_4^+\right) = \frac{1+\mu}{\sqrt{3}}, e_3\left(P_4^+\right) = -\frac{1+\sqrt{3+4\lambda}}{\sqrt{3}}, e_4\left(P_4^+\right) = -\frac{1-\sqrt{3+4\lambda}}{\sqrt{3}}, (49)$$

and

$$e_1(P_4^-) = -2\sqrt{3}, e_2(P_4^-) = -\frac{1+\mu}{\sqrt{3}}, e_3(P_4^-) = \frac{1+\sqrt{3+4\lambda}}{\sqrt{3}}, e_4(P_4^-) = \frac{1-\sqrt{3+4\lambda}}{\sqrt{3}}.$$
(50)

It is straightforward to conclude that P_4^{\pm} are saddle points.

3.1.5. Points P_5^{\pm}

Points P_5^{\pm} have coordinates

$$P_5^{\pm} = \left(\mp \frac{1}{\sqrt{6\lambda}}, 0, 1 - \frac{1}{6\lambda}, \pm 1\right).$$

From the constraint $\Omega_m \ge 0$, it follows that parameter λ is constraint as, $\lambda < 0$ or $\lambda > \frac{1}{6}$. The asymptotic solution at the points describe spatially flat FLRW solutions, i.e. $\Omega_K \left(P_5^{\pm}\right) = 0$, with $q\left(P_5^{\pm}\right) = \frac{1}{4}\left(2 + \frac{1}{\lambda}\right)$ and $w_{tot}\left(P_5^{\pm}\right) = \frac{1}{6\lambda}$. Hence, acceleration exists when $-\frac{1}{2} < \lambda < 0$.

The eigenvalues of the linearized system around the stationary points are determined

$$e_1(P_5^+) = \frac{\sqrt{3}}{2\lambda} (1+6\lambda), \ e_2(P_5^+) = \frac{1+2\lambda}{2\sqrt{3}\lambda}, e_3(P_5^+) = \frac{1-6\lambda}{4\sqrt{3}\lambda}, e_4(P_5^+) = \frac{1+6\lambda-2\mu}{4\sqrt{3}\lambda}, (51)$$

and

$$e_1(P_5^-) = -\frac{\sqrt{3}}{2\lambda}(1+6\lambda), \ e_2(P_5^-) = -\frac{1+2\lambda}{2\sqrt{3}\lambda}, \ e_3(P_5^-) = -\frac{1-6\lambda}{4\sqrt{3}\lambda}, \ e_4(P_5^-) = -\frac{1+6\lambda-2\mu}{4\sqrt{3}\lambda}$$
(52)

Hence, point P_5^+ is an attractor when $\left\{\mu \leq 0, -\frac{1}{6} < \lambda < 0\right\}$, $\left\{0 < \mu < \frac{1}{2}, \frac{2\mu-1}{6} < \lambda < 0\right\}$, while point P_5^- is always a saddle point. We observe that point P_5^+ can describe an accelerated universe as a future attractor for the dynamical system

3.1.6. Points
$$P_6^{\pm}$$

The family of points P_6^{\pm} has coordinates

$$P_{6}^{\pm} = \left(\pm \frac{\sqrt{6}}{1-2\mu}, \frac{\sqrt{(1+6\lambda-2\mu)(1+\mu)}}{1-2\mu}, \frac{12\lambda(\mu-1)}{(1-2\mu)^{2}}, \pm 1\right).$$

The points are real and physically accepted when the free parameters λ and μ are constraint as $\left\{\frac{1}{2} + 3\lambda < \mu < -1\right\}$ or $\left\{\lambda < 0, -1 < \mu < \frac{1}{2} + 3\lambda\right\}$. The physical parameters at the stationary points are derived $\Omega_K\left(P_6^{\pm}\right) = \frac{(1+6\lambda-2\mu)\mu}{(1-2\mu)^2}$, $q\left(P_6^{\pm}\right) = \frac{(1+\mu)\left(\mu-1-18\lambda\mu+2\mu^2\right)}{(1-2\mu)^3}$ and $w_{tot}\left(P_6^{\pm}\right) = \frac{1}{2\mu-1}$. In Fig. 1 we present the region plots in the two-dimensional space for the variables $\{\lambda, \mu\}$, where the stationary points P_6^{\pm} are real, the spatially curvature is negative, i.e. $\Omega_K\left(P_6^{\pm}\right) < 0$, and the exact solution describes an accelerated universe $q\left(P_6^{\pm}\right) < 0$. We observe that when $\omega_R\left(P_6^{\pm}\right) < 0$, then the exact solution describes acceleration.




FIG. 1: Region plots in the two-dimensional space for the variables $\{\lambda, \mu\}$, where the stationary points P_6^{\pm} are real (left plot), the spatially curvature is negative, $\Omega_K \left(P_6^{\pm}\right) < 0$ (center plot), and the exact solution describes an accelerated universe $q \left(P_6^{\pm}\right) < 0$ (right plot)




FIG. 2: Region plots in the two-dimensional space for the variables $\{\lambda, \mu\}$, where the real parts of the eivenvalues for the linearized system at the points P_6^{\pm} are negatives. We observe that points P_6^{\pm} are always saddle points when they exist.

Furthermore, we determine the eigenvalues of the dynamical system numerically around the stationary points P_6^{\pm} and we found that the stationary points P_6^{\pm} are always saddle points when they exist. In Fig. 2 we present the regions in the space $\{\lambda, \mu\}$ in which the real part of the eigenvalues is negative.




FIG. 3: Region plots in the two-dimensional space for the variables $\{\lambda, \mu\}$, where the real parts of the eivenvalues for the linearized system at the points P_7^{\pm} are negatives. We observe that points P_7^{\pm} are always saddle points.

3.1.7. Points
$$P_7^{\pm}$$

Points P_7^{\pm} are similar with P_6^{\pm} where now

$$P_7^{\pm} = \left(\pm \frac{\sqrt{6}}{1-2\mu}, -\frac{\sqrt{(1+6\lambda-2\mu)(1+\mu)}}{1-2\mu}, \frac{12\lambda(\mu-1)}{(1-2\mu)^2}, \pm 1\right).$$

points are physically accepted when $\{1 < \mu < \frac{1}{2} + 3\lambda, \lambda > \frac{1}{6}\}$. The physical parameters Ω_K , q and w_{tot} have similar functional form with that of points P_6^{\pm} . However in contrary with before, when the points exist, the asymptotic solutions describe FLRW universes with positive spatial curvature and no acceleration.

In Fig. 3 we present the regions in the space $\{\lambda, \mu\}$ where the eigenvalues of the linearized system around the points P_7^{\pm} have negative real components. We conclude that the stationary points P_7^{\pm} when they exist, are always saddle points. 3.1.8. Points P_8^{\pm}

The set of stationary points P_8^{\pm} with coordinates

$$P_8^{\pm} = (0, 0, 0, \pm 1), \qquad (53)$$

describes the Milne solution, $\Omega_K(P_8^{\pm}) = -\frac{1}{3}$, $q(P_8^{\pm}) = 0$ and $w_{tot}(P_8^{\pm}) = -\frac{1}{3}$.

The eigenvalues of the linearized system around P_8^{\pm} are

$$e_1(P_8^+) = 2\sqrt{3}, e_2(P_8^+) = -\frac{2}{\sqrt{3}}, e_3(P_8^+) = -\frac{1}{\sqrt{3}}, \ e_4(P_8^+) = \frac{1}{\sqrt{3}}$$
 (54)

and

$$e_1(P_8^-) = -2\sqrt{3}, e_2(P_8^-) = \frac{2}{\sqrt{3}}, e_3(P_8^-) = \frac{1}{\sqrt{3}}, \ e_4(P_8^-) = -\frac{1}{\sqrt{3}}$$
(55)

that is, the Milne solutions are unstable solutions, since points P_8^{\pm} are always saddle points.

3.1.9. Point P_9

Point $P_9 = (0, 0, 0, 0)$, describes the Minkowski spacetime, $\Omega_K(P_9) = 0$. From numerical simulations it is easy to observe that the Minkowski vacuum solution is always unstable.

3.1.10. Line of points P_{10}

Finally, the line of points P_{10} ,

$$P_{10} = (0, \eta, 0, \eta) \tag{56}$$

describe de Sitter universes, $\Omega_K(P_{10}) = 0$, $q(P_{10}) = -1$ and $w_{tot}(P_{10}) = -1$. At the asymptotic solution only the scalar field potential contributes to the cosmological evolution.

The eigenvalues of the linearized system are calculated

$$e_1(P_{10}) = 0, e_2(P_{10}) = -\sqrt{3}\eta, e_3(P_{10}) = -\sqrt{3}\eta, e_4(P_{10}) = \frac{2\eta(2\eta^2 - 3)}{\sqrt{3}}$$

Thus for $\eta < 0$, the points always describe unstable de Sitter universes (saddle or local source), while for $0 < \eta < \sqrt{3/2}$, there exists a submanifold in which the stationary point describes the de Sitter universe as a future attractor. The derivation of the submanifold is a mathematical calculation which does not contribute to the physical discussion. Hence, it is omitted. The cases $1 < \eta < \sqrt{3/2}$ and $\eta \ge \sqrt{3/2}$ are not allowed due to the existence condition $-1 \le \eta \le 1$.

4. CONCLUSIONS

The discrepancies in the cosmological data and the universe's curvature are two major issues in cosmology at present. The first one indicates the need of revision of the standard cosmological model Λ CDM, and the second issue directly questions the geometry of the universe by pointing toward a preference for a closed universe at several standard deviations [3, 23, 24, 31].

The revision of the Λ CDM model has been performed in various ways, and modification of Einstein's gravitational theory is one of them. Even though a cluster of cosmological models has been proposed in the literature, none of them can explain all the observational discrepancies. Thus, from this ground, all cosmological models are equivalent. In the present article, we have considered a modification of Einstein's gravity, namely, the Weyl Integrable geometry and explored the cosmological scenarios in the presence of the nonflat FLRW line element as this describes the most generalized cosmic structure.

We have found some exciting results from this modified gravitational theory. In particular, we found that the dark sector's interaction naturally arises in this context without any mathematical complexities. That fills a gap in the literature on interacting cosmology, which is mainly driven by the phenomenological choices of the interaction function. In order to understand the nature of the cosmological scenario derived from this gravitational framework, we performed the dynamical system analysis and found a variety of possibilities.

The admitted stationary points of the dynamical system corresponding to ten families of points, which can describe all the possible FLRW universes with zero and non-zero spatial curvature. We conclude that the future attractors of the dynamical system describe spatially flat FLRW geometries. Moreover, exact solutions with acceleration are recovered. We conclude that Weyl Integrable geometry can solve the flatness problem. Additionally, this is one of the first studies in literature where interacting models in the dark sector are discussed in the presence of curvature.

In future, we plan to investigate further this cosmological scenario by studying the evolution of the perturbations near the stationary points for the background geometry.

Acknowledgments

The research of AG was funded by Agencia Nacional de Investigación y Desarrollo -ANID through the program FONDECYT Regular grant no. 1200293. GL was funded by Vicerrectoría de Investigación y Desarrollo Tecnológico at UCN. AP was partially supported by the National Research Foundation of South Africa (Grant Numbers 131604). AP thanks Prof. G. Michalakopoulos and the Ionian University for the hospitality provided while part of this work carried out. SP acknowledges the financial support from the Department of Science and Technology (DST), Govt. of India, under the Scheme "Fund for Improvement of S&T Infrastructure (FIST)" [File No. SR/FST/MS-I/2019/41].

- T. M. C. Abbott et al. Dark Energy Survey year 1 results: Cosmological constraints from galaxy clustering and weak lensing. *Phys. Rev. D*, 98(4):043526, 2018.
- [2] Elcio Abdalla et al. Cosmology intertwined: A review of the particle physics, astrophysics, and cosmology associated with the cosmological tensions and anomalies. JHEAp, 34:49–211, 2022.
- [3] N. Aghanim et al. Planck 2018 results. VI. Cosmological parameters. Astron. Astrophys., 641:A6, 2020. [Erratum: Astron.Astrophys. 652, C4 (2021)].
- [4] Ricardo Aguila, José Edgar Madriz Aguilar, Claudia Moreno, and Mauricio Bellini. Present accelerated expansion of the universe from new Weyl-Integrable gravity approach. *Eur. Phys.* J. C, 74(11):3158, 2014.
- [5] J. E. Madriz Aguilar, C. Romero, J. B. Fonseca Neto, T. S. Almeida, and J. B. Formiga. (2+1)-Dimensional Gravity in Weyl Integrable Spacetime. *Class. Quant. Grav.*, 32(21):215003, 2015.
- [6] Nikki Arendse et al. Cosmic dissonance: are new physics or systematics behind a short sound horizon? Astron. Astrophys., 639:A57, 2020.
- [7] Fabiola Arevalo, Antonella Cid, Luis P. Chimento, and Patricio Mella. On sign-changeable interaction in FLRW cosmology. *Eur. Phys. J. C*, 79(4):355, 2019.
- [8] Micheal S. Berger and Hamed Shojaei. Interacting dark energy and the cosmic coincidence problem. Phys. Rev. D, 73:083528, 2006.
- [9] Andrew P. Billyard and Alan A. Coley. Interactions in scalar field cosmology. Phys. Rev. D,

61:083503, 2000.

- [10] Christian G. Boehmer, Nicola Tamanini, and Matthew Wright. Interacting quintessence from a variational approach Part I: algebraic couplings. *Phys. Rev. D*, 91(12):123002, 2015.
- [11] Christian G. Boehmer, Nicola Tamanini, and Matthew Wright. Interacting quintessence from a variational approach Part II: derivative couplings. *Phys. Rev. D*, 91(12):123003, 2015.
- [12] Yi-Fu Cai, Salvatore Capozziello, Mariafelicia De Laurentis, and Emmanuel N. Saridakis. f(T) teleparallel gravity and cosmology. *Rept. Prog. Phys.*, 79(10):106901, 2016.
- [13] Víctor H. Cárdenas, Daniela Grandón, and Samuel Lepe. Dark energy and Dark matter interaction in light of the second law of thermodynamics. *Eur. Phys. J. C*, 79(4):357, 2019.
- [14] Timothy Clifton, Pedro G. Ferreira, Antonio Padilla, and Constantinos Skordis. Modified Gravity and Cosmology. *Phys. Rept.*, 513:1–189, 2012.
- [15] A. A. Coley. *Dynamical systems and cosmology*. Kluwer, Dordrecht, Netherlands, 2003.
- [16] Edmund J. Copeland, M. Sami, and Shinji Tsujikawa. Dynamics of dark energy. Int. J. Mod. Phys. D, 15:1753–1936, 2006.
- [17] Guido D'Amico, Teresa Hamill, and Nemanja Kaloper. Quantum field theory of interacting dark matter and dark energy: Dark monodromies. *Phys. Rev. D*, 94(10):103526, 2016.
- [18] Antonio De Felice and Shinji Tsujikawa. f(R) theories. Living Rev. Rel., 13:3, 2010.
- [19] Sergio del Campo, Ramon Herrera, German Olivares, and Diego Pavon. Interacting models of soft coincidence. *Phys. Rev. D*, 74:023501, 2006.
- [20] Sergio del Campo, Ramon Herrera, and Diego Pavon. Toward a solution of the coincidence problem. Phys. Rev. D, 78:021302, 2008.
- [21] Eleonora Di Valentino et al. Cosmology intertwined III: $f\sigma_8$ and S_8 . Astropart. Phys., 131:102604, 2021.
- [22] Eleonora Di Valentino et al. Snowmass2021 Letter of interest cosmology intertwined II: The hubble constant tension. Astropart. Phys., 131:102605, 2021.
- [23] Eleonora Di Valentino, Alessandro Melchiorri, and Joseph Silk. Planck evidence for a closed Universe and a possible crisis for cosmology. *Nature Astron.*, 4(2):196–203, 2019.
- [24] Eleonora Di Valentino, Alessandro Melchiorri, and Joseph Silk. Investigating Cosmic Discordance. Astrophys. J. Lett., 908(1):L9, 2021.
- [25] Eleonora Di Valentino, Olga Mena, Supriya Pan, Luca Visinelli, Weiqiang Yang, Alessandro Melchiorri, David F. Mota, Adam G. Riess, and Joseph Silk. In the realm of the Hubble

tension—a review of solutions. Class. Quant. Grav., 38(15):153001, 2021.

- [26] J. C. Fabris, J. M. Salim, and S. L. Sautu. Inflationary cosmological solutions in Weyl integrable geometry. *Mod. Phys. Lett. A*, 13:953–959, 1998.
- [27] Mónica Forte. On extended sign-changeable interactions in the dark sector. Gen. Rel. Grav., 46(10):1811, 2014.
- [28] Radouane Gannouji, Hemwati Nandan, and Naresh Dadhich. FLRW cosmology in Weyl-Integrable Space-Time. JCAP, 11:051, 2011.
- [29] Jérôme Gleyzes, David Langlois, Michele Mancarella, and Filippo Vernizzi. Effective Theory of Interacting Dark Energy. JCAP, 08:054, 2015.
- [30] Juan-Juan Guo, Jing-Fei Zhang, Yun-He Li, Dong-Ze He, and Xin Zhang. Probing the signchangeable interaction between dark energy and dark matter with current observations. Sci. China Phys. Mech. Astron., 61(3):030011, 2018.
- [31] Will Handley. Curvature tension: evidence for a closed universe. *Phys. Rev. D*, 103(4):L041301, 2021.
- [32] H. Hildebrandt et al. KiDS-450: Cosmological parameter constraints from tomographic weak gravitational lensing. Mon. Not. Roy. Astron. Soc., 465:1454, 2017.
- [33] H. Hildebrandt et al. KiDS+VIKING-450: Cosmic shear tomography with optical and infrared data. Astron. Astrophys., 633:A69, 2020.
- [34] Greg Huey and Benjamin D. Wandelt. Interacting quintessence. The Coincidence problem and cosmic acceleration. *Phys. Rev. D*, 74:023519, 2006.
- [35] Shahab Joudaki et al. KiDS-450 + 2dFLenS: Cosmological parameter constraints from weak gravitational lensing tomography and overlapping redshift-space galaxy clustering. Mon. Not. Roy. Astron. Soc., 474(4):4894–4924, 2018.
- [36] Nemanja Kaloper and Keith A. Olive. Singularities in scalar tensor cosmologies. Phys. Rev. D, 57:811–822, 1998.
- [37] Genly Leon, Esteban González, Alfredo D. Millano, and Felipe Orlando Franz Silva. A Perturbative Analysis of Interacting Scalar Field Cosmologies, 3 2020.
- [38] Samuel Lepe and Francisco Peña. Interacting cosmic fluids and phase transitions under a holographic modeling for dark energy. *Eur. Phys. J. C*, 76(9):507, 2016.
- [39] John Miritzis. Acceleration in Weyl integrable spacetime. Int. J. Mod. Phys. D, 22:1350019, 2013.

- [40] S. Nojiri, S.D. Odintsov, and V.K. Oikonomou. Modified Gravity Theories on a Nutshell: Inflation, Bounce and Late-time Evolution. *Phys. Rept.*, 692:1–104, 2017.
- [41] Shin'ichi Nojiri and Sergei D. Odintsov. Introduction to modified gravity and gravitational alternative for dark energy. eConf, C0602061:06, 2006.
- [42] Andronikos Paliathanasis. Dynamical Analysis and Cosmological Evolution in Weyl Integrable Gravity. Universe, 7(12):468, 2021.
- [43] Andronikos Paliathanasis, Genly Leon, and John D. Barrow. Einstein-aether theory in Weyl integrable geometry. Eur. Phys. J. C, 80(12):1099, 2020.
- [44] Andronikos Paliathanasis, Genly Leon, and John D. Barrow. Inhomogeneous spacetimes in Weyl integrable geometry with matter source. *Eur. Phys. J. C*, 80(8):731, 2020.
- [45] Supriya Pan, German S. Sharov, and Weiqiang Yang. Field theoretic interpretations of interacting dark energy scenarios and recent observations. *Phys. Rev. D*, 101(10):103533, 2020.
- [46] Supriya Pan, Weiqiang Yang, Chiranjeeb Singha, and Emmanuel N. Saridakis. Observational constraints on sign-changeable interaction models and alleviation of the H₀ tension. *Phys. Rev.*, D100(8):083539, 2019.
- [47] Leandros Perivolaropoulos and Foteini Skara. Challenges for ACDM: An update. 5 2021.
- [48] S. Perlmutter et al. Measurements of the cosmological parameters Omega and Lambda from the first 7 supernovae at z>=0.35. Astrophys. J., 483:565, 1997.
- [49] Adam G. Riess et al. Observational evidence from supernovae for an accelerating universe and a cosmological constant. Astron. J., 116:1009–1038, 1998.
- [50] Adam G. Riess et al. A Comprehensive Measurement of the Local Value of the Hubble Constant with 1 km/s/Mpc Uncertainty from the Hubble Space Telescope and the SH0ES Team. 12 2021.
- [51] Jesús Martín Romero, Mauricio Bellini, and José Edgar Madriz Aguilar. Gravitational waves from a Weyl-Integrable manifold: a new formalism. *Phys. Dark Univ.*, 13:1–6, 2016.
- [52] J. M. Salim and S. L. Sautu. Gravitational collapse in Weyl integrable space-times. Class. Quant. Grav., 16:3281–3295, 1999.
- [53] Erhard Scholz. MOND-like acceleration in integrable Weyl geometric gravity. Found. Phys., 46(2):176–208, 2016.
- [54] Thomas P. Sotiriou and Valerio Faraoni. f(R) Theories Of Gravity. Rev. Mod. Phys., 82:451–497, 2010.

- [55] Cheng-Yi Sun and Rui-Hong Yue. New Interaction between Dark Energy and Dark Matter Changes Sign during Cosmological Evolution. *Phys. Rev. D*, 85:043010, 2012.
- [56] C. van de Bruck and J. Morrice. Disformal couplings and the dark sector of the universe. JCAP, 04:036, 2015.
- [57] Hao Wei. Cosmological Constraints on the Sign-Changeable Interactions. Commun. Theor. Phys., 56:972–980, 2011.
- [58] Steven Weinberg. The Cosmological Constant Problem. Rev. Mod. Phys., 61:1–23, 1989.
- [59] Ivaylo Zlatev, Li-Min Wang, and Paul J. Steinhardt. Quintessence, cosmic coincidence, and the cosmological constant. *Phys. Rev. Lett.*, 82:896–899, 1999.