

# $\mathcal{R}(p, q)$ -trinomial probability distribution: properties and particular cases

Fridolin Melong

*Institut für Mathematik, Universität Zürich,  
Winterthurerstrasse 190, CH-8057 Zürich, Switzerland  
fridomelong@gmail.com*

June 16, 2022

## Abstract

In this paper, we investigate the trinomial probability distribution of the first and second kind from the  $\mathcal{R}(p, q)$ -quantum algebras. Moreover, we compute their  $\mathcal{R}(p, q)$ -factorial moments and derive the corresponding covariance. Particular cases of trinomial probability distribution are deduced from the formalism developed.

**Keywords.**  $\mathcal{R}(p, q)$ -calculus, quantum algebras, trinomial distribution, factorial moments,  $\mathcal{R}(p, q)$ -covariance.  
MSC (2020)17B37, 60E05, 05A30

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Generalized trinomial probability distribution</b>	<b>4</b>
2.1	$\mathcal{R}(p, q)$ -trinomial distribution of the first kind . . . . .	4
2.2	Negative $\mathcal{R}(p, q)$ -trinomial distribution of the first kind . . . . .	7
2.3	$\mathcal{R}(p, q)$ -trinomial distribution of the second kind . . . . .	10
2.4	Negative $\mathcal{R}(p, q)$ -trinomial distribution of the second kind . . . . .	13
<b>3</b>	<b>Trinomial distributions and particular quantum algebras</b>	<b>16</b>
3.1	Trinomial distribution and Biedenharn-Macfarlane algebra [1, 13] . . . . .	16
3.2	Trinomial distribution and Jagannathan-Srinivasa algebra [12] . . . . .	19
3.3	Trinomial distribution and Chakrabarty and Jagannathan algebra [2] . . . . .	21
3.4	Trinomial distribution and Hounkonnou-Ngompe generalization of $q$ -Quesne algebra [11] . . . . .	24

## 1 Introduction

Charalambos presented the  $q$ - deformed Vandermonde and Cauchy formulae. Moreover, the  $q$ - deformed univariate discrete probability distributions were investigated. Their properties and limiting distributions were derived [3].

Furthermore, the  $q$ - deformed multinomial coefficients was defined and their recurrence relations were deduced. Also, the  $q$ - deformed multinomial and negative  $q$ - deformed multinomial probability distributions of the first and second kind were presented [4].

The same author extended the multivariate  $q$ - deformed Vandermonde and Cauchy formulae, and the multivariate  $q$ - Polya and inverse  $q$ - Polya were constructed [5]. Moreover, the  $q$ - factorial moments of the bivariate discrete distributions were investigated [6].

Let  $p$  and  $q$  be two positive real numbers such that  $0 < q < p < 1$ . We consider a meromorphic function  $\mathcal{R}$  defined on  $\mathbb{C} \times \mathbb{C}$  by [9]:

$$\mathcal{R}(u, v) = \sum_{s,t=-l}^{\infty} r_{st} u^s v^t, \quad (1.1)$$

with an eventual isolated singularity at the zero, where  $r_{st}$  are complex numbers,  $l \in \mathbb{N} \cup \{0\}$ ,  $\mathcal{R}(p^n, q^n) > 0, \forall n \in \mathbb{N}$ , and  $\mathcal{R}(1, 1) = 0$  by definition. We denote by  $\mathbb{D}_R$  the bidisk

$$\mathbb{D}_R = \{w = (w_1, w_2) \in \mathbb{C}^2 : |w_j| < R_j\},$$

where  $R$  is the convergence radius of the series (1.1) defined by Hadamard formula as follows [14]:

$$\limsup_{s+t \rightarrow \infty} \sqrt[s+t]{|r_{st}| R_1^s R_2^t} = 1.$$

We denote by  $\mathcal{O}(\mathbb{D}_R)$  the set of holomorphic functions defined on  $\mathbb{D}_R$ .

The  $\mathcal{R}(p, q)$ - deformed numbers is defined by [9]:

$$[n]_{\mathcal{R}(p,q)} := \mathcal{R}(p^n, q^n), \quad n \in \mathbb{N}, \quad (1.2)$$

the  $\mathcal{R}(p, q)$ - deformed factorials and binomial coefficients are given as:

$$[n]!_{\mathcal{R}(p,q)} := \begin{cases} 1 & \text{for } n = 0 \\ \mathcal{R}(p, q) \cdots \mathcal{R}(p^n, q^n) & \text{for } n \geq 1, \end{cases} \quad (1.3)$$

and

$$\begin{bmatrix} n \\ m \end{bmatrix}_{\mathcal{R}(p,q)} := \frac{[n]!_{\mathcal{R}(p,q)}}{[m]!_{\mathcal{R}(p,q)} [n-m]!_{\mathcal{R}(p,q)}}, \quad (n, m) \in \mathbb{N} \cup \{0\}, \quad n \geq m. \quad (1.4)$$

We consider the following linear operators on  $\mathcal{O}(\mathbb{D}_R)$  given by:

$$\begin{aligned} Q &: \varphi \longmapsto Q\varphi(z) := \varphi(qz) \\ P &: \varphi \longmapsto P\varphi(z) := \varphi(pz), \end{aligned} \quad (1.5)$$

leading to define the  $\mathcal{R}(p, q)$ - deformed derivative:

$$\partial_{\mathcal{R},p,q} := \partial_{p,q} \frac{p-q}{P-Q} \mathcal{R}(P, Q) = \frac{p-q}{pP-qQ} \mathcal{R}(pP, qQ) \partial_{p,q}, \quad (1.6)$$

where  $\partial_{p,q}$  is the  $(p, q)$ - derivative:

$$\partial_{p,q} : \varphi \longmapsto \partial_{p,q}\varphi(z) := \frac{\varphi(pz) - \varphi(qz)}{z(p - q)}. \quad (1.7)$$

The quantum algebra associated with the  $\mathcal{R}(p, q)$ - deformation, denoted by  $\mathcal{A}_{\mathcal{R}(p,q)}$  is generated by the set of operators  $\{1, A, A^\dagger, N\}$  satisfying the following commutation relations [10]:

$$\begin{aligned} AA^\dagger &= [N + 1]_{\mathcal{R}(p,q)}, & A^\dagger A &= [N]_{\mathcal{R}(p,q)}. \\ [N, A] &= -A, & [N, A^\dagger] &= A^\dagger \end{aligned} \quad (1.8)$$

with its realization on  $\mathcal{O}(\mathbb{D}_R)$  given by:

$$A^\dagger := z, \quad A := \partial_{\mathcal{R}(p,q)}, \quad N := z\partial_z,$$

where  $\partial_z := \frac{\partial}{\partial z}$  is the usual derivative on  $\mathbb{C}$ .

The  $\mathcal{R}(p, q)$ - deformed numbers (1.2) can be rewritten as follows [7]:

$$[n]_{\mathcal{R}(p,q)} = \frac{\phi_1^n - \phi_2^n}{\phi_1 - \phi_2}, \quad \phi_1 \neq \phi_2, \quad (1.9)$$

where  $(\phi_i)_{i \in \{1,2\}}$  are functions of the parameters deformations  $p$  and  $q$ .

The following relations hold [8]:

$$[x]_{\mathcal{R}(p^{-1}, q^{-1})} = (\phi_1 \phi_2)^{1-x} [x]_{\mathcal{R}(p,q)}, \quad (1.10)$$

$$[r]_{\mathcal{R}(p^{-1}, q^{-1})}! = (\phi_1 \phi_2)^{-\binom{r}{2}} [r]_{\mathcal{R}(p,q)}!, \quad (1.11)$$

$$[x]_{r, \mathcal{R}(p^{-1}, q^{-1})} = (\phi_1 \phi_2)^{-xr + \binom{r+1}{2}} [x]_{r, \mathcal{R}(p,q)}. \quad (1.12)$$

For  $a, b \in \mathbb{N}$ , the  $\mathcal{R}(p, q)$ - deformed shifted factorial is defined by [7]:

$$(u \oplus v)_{\mathcal{R}(p,q)}^n := \prod_{i=1}^n (u \phi_1^{i-1} + v \phi_2^{i-1}), \quad \text{with} \quad (u \oplus v)_{\mathcal{R}(p,q)}^0 := 1.$$

Analogously,

$$(u \ominus v)_{\mathcal{R}(p,q)}^n := \prod_{i=1}^n (u \phi_1^{i-1} - b \phi_2^{i-1}), \quad \text{with} \quad (u \ominus v)_{\mathcal{R}(p,q)}^0 := 1.$$

The  $\mathcal{R}(p, q)$ - deformed factorial of  $u$  of order  $r$  is defined by [8]:

$$[u]_{r, \mathcal{R}(p,q)} = \prod_{i=1}^r [u - i + 1]_{\mathcal{R}(p,q)}, \quad r \in \mathbb{N},$$

Furthermore, the generalized Vandermonde, Cauchy formulae and univariate probability distributions induced from the  $\mathcal{R}(p, q)$ - deformed quantum algebras are investigated in [7].

Our aim is to determine the trinomial probability distribution of the first and second kind with properties associated to the  $\mathcal{R}(p, q)$ - deformed quantum algebras [10].

This paper is organized as follows: In section 2, we present the  $\mathcal{R}(p, q)$ - trinomial probability distribution of the first and second kind. Also their negative counterparts. Besides, the properties namely ( $\mathcal{R}(p, q)$ -factorial moments and covariance) are investigated. Section 3 is reserved to deduce the relevant particular cases corresponding to some quantum algebras known in the literature.

## 2 Generalized trinomial probability distribution

The trinomial probability distribution of the first and second kind from the  $\mathcal{R}(p, q)$ -deformed quantum algebras are investigated. Their factorial moments are also computed. Besides, the  $\mathcal{R}(p, q)$ - covariance is deduced.

### 2.1 $\mathcal{R}(p, q)$ -trinomial distribution of the first kind

The probability function of the  $\mathcal{R}(p, q)$ -random vector  $\underline{Y} = (Y_1, Y_2)$  of the  $\mathcal{R}(p, q)$ -trinomial probability distribution of the first kind, with parameters  $n, \underline{\alpha} = (\alpha_1, \alpha_2), p$ , and,  $q$  is given by:

$$P(Y_1 = y_1, Y_2 = y_2) = \begin{bmatrix} n \\ y_1, y_2 \end{bmatrix}_{\mathcal{R}(p,q)} \frac{\alpha_1^{y_1} \alpha_2^{y_2} \phi_1^{\binom{n-y_1}{2} + \binom{n-y_2}{2}} \phi_2^{\binom{y_1}{2} + \binom{y_2}{2}}}{(1 \oplus \alpha_1)_{\mathcal{R}(p,q)}^n (1 \oplus \alpha_2)_{\mathcal{R}(p,q)}^{n-y_1}},$$

where  $y_j \in \{0, 1, \dots, n\}, y_1 + y_2 \leq n, s_j = \sum_{i=1}^j y_i, 0 < \alpha_j < 1$ , and  $j \in \{1, 2\}$ .

**Theorem 2.1** *The  $\mathcal{R}(p, q)$ -factorial moments of the  $\mathcal{R}(p, q)$ -trinomial probability distribution of the first kind, with parameters  $n, \underline{\alpha} = (\alpha_1, \alpha_2), p$ , and,  $q$  are determined by the following relations:*

$$E([Y_1]_{m_1, \mathcal{R}(p,q)}) = \frac{\alpha_1^{m_1} \phi_2^{\binom{m_1}{2}} [n]_{m_1, \mathcal{R}(p,q)}}{(1 \oplus \alpha_1)_{\mathcal{R}(p,q)}^{m_1}}, \quad m_1 \in \{0, 1, \dots, n\}, \quad (2.1)$$

$$E([Y_2]_{m_2, \mathcal{R}(p,q)} | Y_1 = y_1) = \frac{\alpha_2^{m_2} \phi_2^{\binom{m_2}{2}} [n - y_1]_{m_2, \mathcal{R}(p,q)}}{(1 \oplus \alpha_2)_{\mathcal{R}(p,q)}^{m_2}}, \quad (2.2)$$

with  $m_2 \in \{0, 1, \dots, n - y_1\}$ .

$$E([Y_2]_{m_2, \mathcal{R}(p,q)}) = \frac{\alpha_2^{m_2} \phi_1^{\binom{m_2}{2} + m_2(n-m_2)} \phi_2^{\binom{m_2}{2}} [n]_{m_2, \mathcal{R}(p,q)}}{(1 \oplus \alpha_2)_{\mathcal{R}(p,q)}^{m_2} (\phi_1^{n-m_2} \oplus \alpha_1 \phi_2^{n-m_2})_{\mathcal{R}(p,q)}^{m_2}}, \quad (2.3)$$

with  $m_2 \in \{0, 1, \dots, n\}$  and

$$E([Y_1]_{m_1, \mathcal{R}(p,q)} [Y_2]_{m_2, \mathcal{R}(p,q)}) = \frac{\phi_1^{\binom{m_2}{2} + m_2(n-m_2)} \phi_2^{\binom{m_1}{2} + \binom{m_2}{2}} \alpha_1^{m_1} \alpha_2^{m_2} [n]_{m_1+m_2, \mathcal{R}(p,q)}}{(1 \oplus \alpha_1)_{\mathcal{R}(p,q)}^{m_1} (1 \oplus \alpha_2)_{\mathcal{R}(p,q)}^{m_2} (\phi_1^{n-m_2} \oplus \alpha_1 \phi_2^{n-m_2})_{\mathcal{R}(p,q)}^{m_2}}, \quad (2.4)$$

where  $m_1 \in \{0, 1, \dots, n - m_2\}$  and  $m_2 \in \{0, 1, \dots, n\}$ .

**Proof 2.2** The  $\mathcal{R}(p, q)$ - random variable  $Y_1$  follows the  $\mathcal{R}(p, q)$ - binomial probability distribution of the first kind as:

$$P(Y_1 = y_1) = \begin{bmatrix} n \\ y_1 \end{bmatrix}_{\mathcal{R}(p,q)} \frac{\alpha_1^{y_1} \phi_1^{\binom{n-y_1}{2}} \phi_2^{\binom{y_1}{2}}}{(1 \oplus \alpha_1)_{\mathcal{R}(p,q)}^n}, \quad y_1 \in \{0, 1, \dots, n\}. \quad (2.5)$$

From [7], the  $\mathcal{R}(p, q)$ - factorial moments of  $Y_1$  are given by (2.1). Moreover, the conditional probability distribution of  $Y_2$ , given that  $Y_1 = y_1$ , is the  $\mathcal{R}(p, q)$ - binomial probability distribution of the first kind, with mass function :

$$P(Y_2 = y_2 | Y_1 = y_1) = \begin{bmatrix} n - y_1 \\ y_2 \end{bmatrix}_{\mathcal{R}(p,q)} \frac{\alpha_2^{y_2} \phi_1^{\binom{n-y_1-y_2}{2}} \phi_2^{\binom{y_2}{2}}}{(1 \oplus \alpha_2)_{\mathcal{R}(p,q)}^{n-y_1}}, \quad y_2 \in \{0, 1, \dots, n - y_1\}.$$

Using also [7], the conditional  $\mathcal{R}(p, q)$ -factorial moments of  $Y_2$ , given that  $Y_1 = y_1$ , are furnished by (2.2). Besides, we determine the  $\mathcal{R}(p, q)$ - factorial moments of  $Y_2$  according to the formula:

$$\begin{aligned} E([Y_2]_{m_2, \mathcal{R}(p,q)}) &= E\left(E([Y_2]_{m_2, \mathcal{R}(p,q)} | Y_1)\right) \\ &= \frac{\alpha_2^{m_2} \phi_2^{\binom{m_2}{2}}}{(1 \oplus \alpha_2)_{\mathcal{R}(p,q)}^{m_2}} E([n - Y_1]_{m_2, \mathcal{R}(p,q)}). \end{aligned}$$

Obviously,

$$E([n - Y_1]_{m_2, \mathcal{R}(p,q)}) = \sum_{y_1=0}^{n-m_2} [n - y_1]_{m_2, \mathcal{R}(p,q)} \begin{bmatrix} n \\ y_1 \end{bmatrix}_{\mathcal{R}(p,q)} \frac{\alpha_1^{y_1} \phi_1^{\binom{n-y_1}{2}} \phi_2^{\binom{y_1}{2}}}{(1 \oplus \alpha_1)_{\mathcal{R}(p,q)}^n}.$$

From the relation

$$[n - y_1]_{m_2, \mathcal{R}(p,q)} \begin{bmatrix} n \\ y_1 \end{bmatrix}_{\mathcal{R}(p,q)} = [n]_{m_2, \mathcal{R}(p,q)} \begin{bmatrix} n - m_2 \\ y_1 \end{bmatrix}_{\mathcal{R}(p,q)}$$

and the  $\mathcal{R}(p, q)$ -binomial formula [8]:

$$(1 \oplus t)_{\mathcal{R}(p,q)}^n = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_{\mathcal{R}(p,q)} \phi_1^{\binom{n-k}{2}} \phi_2^{\binom{k}{2}} t^k, \quad t \in \mathbb{R},$$

we have:

$$\begin{aligned}
E([n - Y_1]_{m_2, \mathcal{R}(p, q)}) &= \sum_{y_1=0}^{n-m_2} [n]_{m_2, \mathcal{R}(p, q)} \begin{bmatrix} n - m_2 \\ y_1 \end{bmatrix}_{\mathcal{R}(p, q)} \frac{\alpha_1^{y_1} \phi_1^{\binom{n-y_1}{2}} \phi_2^{\binom{y_1}{2}}}{(1 \oplus \alpha_1)_{\mathcal{R}(p, q)}^n} \\
&= \frac{\phi_1^{\binom{m_2}{2} + m_2(n-m_2)} [n]_{m_2, \mathcal{R}(p, q)} (1 \oplus \alpha_1)_{\mathcal{R}(p, q)}^{n-m_2}}{(1 \oplus \alpha_1)_{\mathcal{R}(p, q)}^n} \\
&= \frac{\phi_1^{\binom{m_2}{2} + m_2(n-m_2)} [n]_{m_2, \mathcal{R}(p, q)}}{(\phi_1^{n-m_2} \oplus \alpha_1 \phi_2^{n-m_2})_{\mathcal{R}(p, q)}^{m_2}}.
\end{aligned}$$

Thus,

$$E([Y_2]_{m_2, \mathcal{R}(p, q)}) = \frac{\alpha_2^{m_2} \phi_2^{\binom{m_2}{2}}}{(1 \oplus \alpha_2)_{\mathcal{R}(p, q)}^{m_2}} \frac{[n]_{m_2, \mathcal{R}(p, q)}}{(\phi_1^{n-m_2} \oplus \alpha_1 \phi_2^{n-m_2})_{\mathcal{R}(p, q)}^{m_2}}.$$

So, the joint  $\mathcal{R}(p, q)$ -factorial moments  $E([Y_1]_{m_1, \mathcal{R}(p, q)} [Y_2]_{m_2, \mathcal{R}(p, q)})$ ,  $m_2 \in \{0, 1, \dots, n - m_1\}$ ,  $m_1 \in \{0, 1, \dots, n\}$  can be calculated by applying the relation:

$$\begin{aligned}
E([Y_1]_{m_1, \mathcal{R}(p, q)} [Y_2]_{m_2, \mathcal{R}(p, q)}) &= E\left(E([Y_1]_{m_1, \mathcal{R}(p, q)} [Y_2]_{m_2, \mathcal{R}(p, q)} | Y_1)\right) \\
&= \frac{\alpha_2^{m_2} \phi_2^{\binom{m_2}{2}}}{(1 \oplus \alpha_2)_{\mathcal{R}(p, q)}^{m_2}} E([Y_1]_{m_1, \mathcal{R}(p, q)} [n - Y_1]_{m_2, \mathcal{R}(p, q)}).
\end{aligned}$$

Since

$$\begin{aligned}
E([Y_1]_{m_1, \mathcal{R}(p, q)} [n - Y_1]_{m_2, \mathcal{R}(p, q)}) &= \sum_{y_1=m_1}^{n-m_2} [y_1]_{m_1, \mathcal{R}(p, q)} [n - y_1]_{m_2, \mathcal{R}(p, q)} \\
&\quad \times \begin{bmatrix} n \\ y_1 \end{bmatrix}_{\mathcal{R}(p, q)} \frac{\alpha_1^{y_1} \phi_1^{\binom{n-y_1}{2}} \phi_2^{\binom{y_1}{2}}}{(1 \oplus \alpha_1)_{\mathcal{R}(p, q)}^n}.
\end{aligned}$$

Using the relation

$$[y_1]_{m_1, \mathcal{R}(p, q)} [n - y_1]_{m_2, \mathcal{R}(p, q)} \begin{bmatrix} n \\ y_1 \end{bmatrix}_{\mathcal{R}(p, q)} = [n]_{m_1+m_2, \mathcal{R}(p, q)} \begin{bmatrix} n - m_1 - m_2 \\ y_1 - m_1 \end{bmatrix}_{\mathcal{R}(p, q)}$$

and the  $\mathcal{R}(p, q)$ -binomial formula [8], we obtain:

$$\begin{aligned}
E([Y_1]_{m_1, \mathcal{R}(p, q)} [n - Y_1]_{m_2, \mathcal{R}(p, q)}) &= [n]_{m_1+m_2, \mathcal{R}(p, q)} \alpha_1^{m_1} \phi_2^{\binom{m_1}{2}} \\
&\quad \times \sum_{y_1=m_1}^{n-m_2-m_1} \begin{bmatrix} n - m_1 - m_2 \\ y_1 - m_1 \end{bmatrix}_{\mathcal{R}(p, q)} \frac{(\alpha_1 \phi_2^{m_1})^{y_1-m_1} \phi_2^{\binom{y_1-m_1}{2}}}{(1 \oplus \alpha_1)_{\mathcal{R}(p, q)}^n} \\
&= \frac{[n]_{m_1+m_2, \mathcal{R}(p, q)} \alpha_1^{m_1} \phi_2^{\binom{m_1}{2}} (\phi_1^{m_1} \oplus \alpha_1 \phi_2^{m_1})_{\mathcal{R}(p, q)}^{n-m_1-m_2}}{(1 \oplus \alpha_1)_{\mathcal{R}(p, q)}^n}.
\end{aligned}$$

Thus,

$$\begin{aligned} E([Y_1]_{m_1, \mathcal{R}(p,q)} [Y_2]_{m_2, \mathcal{R}(p,q)}) &= \frac{[n]_{m_1+m_2, \mathcal{R}(p,q)} \alpha_1^{m_1} \alpha_2^{m_2} \phi_2^{\binom{m_1}{2} + \binom{m_2}{2}} (\phi_1^{m_1} \oplus \alpha_1 \phi_2^{m_1})_{\mathcal{R}(p,q)}^{n-m_1-m_2}}{(1 \oplus \alpha_2)_{\mathcal{R}(p,q)}^{m_2} (1 \oplus \alpha_1)_{\mathcal{R}(p,q)}^n} \\ &= \frac{[n]_{m_1+m_2, \mathcal{R}(p,q)} \alpha_1^{m_1} \alpha_2^{m_2} \phi_2^{\binom{m_1}{2} + \binom{m_2}{2}}}{(1 \oplus \alpha_1)_{\mathcal{R}(p,q)}^{m_1} (1 \oplus \alpha_2)_{\mathcal{R}(p,q)}^{m_2} (\phi_1^{n-m_2} \oplus \alpha_1 \phi_2^{n-m_2})_{\mathcal{R}(p,q)}^{m_2}}. \end{aligned}$$

**Corollary 2.3** The  $\mathcal{R}(p, q)$ - covariance of the  $\mathcal{R}(p, q)$ - trinomial probability distribution of the first kind is presented by:

$$\text{Cov}([Y_1]_{\mathcal{R}(p,q)}, [Y_2]_{\mathcal{R}(p,q)}) = \frac{\phi_1^{n-1} \alpha_1 \alpha_2 [n]_{\mathcal{R}(p,q)} ([n-1]_{\mathcal{R}(p,q)} - [n]_{\mathcal{R}(p,q)})}{(1 + \alpha_1)(1 + \alpha_2)(\phi_1^{n-1} + \alpha_1 \phi_2^{n-1})}.$$

**Proof 2.4** By definition, the covariance of  $[Y_1]_{\mathcal{R}(p,q)}$  and  $[Y_2]_{\mathcal{R}(p,q)}$  is given by

$$\text{Cov}([Y_1]_{\mathcal{R}(p,q)}, [Y_2]_{\mathcal{R}(p,q)}) = E([Y_1]_{\mathcal{R}(p,q)} [Y_2]_{\mathcal{R}(p,q)}) - E([Y_1]_{\mathcal{R}(p,q)}) E([Y_2]_{\mathcal{R}(p,q)}).$$

Taking  $m_1 = m_2 = 1$ , in the relations (2.1), (2.2), and (2.3), we get

$$E([Y_1]_{\mathcal{R}(p,q)}) E([Y_2]_{\mathcal{R}(p,q)}) = \frac{\alpha_1 [n]_{\mathcal{R}(p,q)}}{(1 + \alpha_1)} \frac{\alpha_2 \phi_1^{n-1} [n]_{\mathcal{R}(p,q)}}{(1 + \alpha_2)(\phi_1^{n-1} + \alpha_1 \phi_2^{n-1})}$$

and

$$E([Y_1]_{\mathcal{R}(p,q)} [Y_2]_{\mathcal{R}(p,q)}) = \frac{\phi_1^{n-1} \alpha_1 \alpha_2 [n]_{2, \mathcal{R}(p,q)}}{(1 + \alpha_1)(1 + \alpha_2)(\phi_1^{n-1} + \alpha_1 \phi_2^{n-1})}.$$

After computation, the result follows.

## 2.2 Negative $\mathcal{R}(p, q)$ -trinomial distribution of the first kind

Let  $U_n$  be the number of successes until the occurrence of the  $n^{\text{th}}$  failure, in a sequence of independent Bernoulli trials, with probability of success at the  $i^{\text{th}}$  trial given as:

$$p_i = \frac{\alpha \phi_2^{i-1}}{\phi_1^{i-1} + \alpha \phi_2^{i-1}}, \quad i \in \mathbb{N} \cup \{0\}.$$

**Lemma 2.5** The  $\mathcal{R}(p, q)$ - random variable  $U_n$  obeys the negative  $\mathcal{R}(p, q)$ - binomial probability distribution of the first kind :

$$P(U = u) = \left[ \begin{matrix} n + u - 1 \\ u \end{matrix} \right]_{\mathcal{R}(p,q)} \frac{\alpha^u \phi_1^{\binom{n-u}{2}} \phi_2^{\binom{u}{2}}}{(1 \oplus \alpha)_{\mathcal{R}(p,q)}^{n+u}}, \quad u \in \mathbb{N} \cup \{0\}. \quad (2.6)$$

Moreover, the  $\mathcal{R}(p^{-1}, q^{-1})$ - factorial moments are presented by the relation

$$E([U_n]_{m, \mathcal{R}(p^{-1}, q^{-1})}) = [n + m - 1]_{m, \mathcal{R}(p^{-1}, q^{-1})} \alpha^m, \quad m \in \mathbb{N} \cup \{0\}. \quad (2.7)$$

We denote by  $W_j$  the number of successes of the  $j^{\text{th}}$  kind until the occurrence of the  $n$ th failure of the second kind, in a sequence of Bernoulli trials with chain-composite failures. Then, the  $\mathcal{R}(p, q)$ -random vector  $\underline{W} = (W_1, W_2)$  follows the negative  $\mathcal{R}(p, q)$ -trinomial probability distribution of the first kind, with parameters  $n$ ,  $\underline{\alpha} = (\alpha_1, \alpha_2)$ ,  $p$ , and,  $q$ . Its mass function is presented by:

$$P(\underline{W} = \underline{w}) = \binom{n + w_1 + w_2 - 1}{w_1, w_2}_{\mathcal{R}(p, q)} \frac{\alpha_1^{w_1} \alpha_2^{w_2} \phi_1^{\binom{n-w_1}{2} + \binom{n-w_2}{2}} \phi_2^{\binom{w_1}{2} + \binom{w_2}{2}}}{(1 \oplus \alpha_1)_{\mathcal{R}(p, q)}^{n+w_1+w_2} (1 \oplus \alpha_2)_{\mathcal{R}(p, q)}^{n+w_2}}, \quad (2.8)$$

where  $w_j \in \mathbb{N} \cup \{0\}$ ,  $0 < \alpha_j < 1$ , and  $j \in \{1, 2\}$ .

**Theorem 2.6** For  $m_1 \in \mathbb{N} \cup \{0\}$  and  $m_2 \in \mathbb{N} \cup \{0\}$ , the  $\mathcal{R}(p^{-1}, q^{-1})$ -factorial moments of the negative  $\mathcal{R}(p, q)$ -trinomial probability distribution of the first kind, with parameters  $n$ ,  $\underline{\alpha} = (\alpha_1, \alpha_2)$ ,  $p$ , and,  $q$  are given as follows:

$$E([W_2]_{m_2, \mathcal{R}(p^{-1}, q^{-1})}) = \alpha_2^{m_2} [n + m_2 - 1]_{m_2, \mathcal{R}(p, q)}, \quad (2.9)$$

$$E([W_1]_{m_1, \mathcal{R}(p^{-1}, q^{-1})} | W_2 = w_2) = \alpha_1^{m_1} [n + w_2 + m_1 - 1]_{m_1, \mathcal{R}(p, q)}, \quad (2.10)$$

$$E\left(\frac{[W_1]_{m_1, \mathcal{R}(p^{-1}, q^{-1})}}{(\phi_1^{n+W_2} \oplus \alpha_2 \phi_2^{n+W_2})_{\mathcal{R}(p, q)}^{m_1}}\right) = \alpha_1^{m_1} [n + m_1 - 1]_{m_1, \mathcal{R}(p, q)}, \quad (2.11)$$

and

$$E\left(\frac{[W_1]_{m_1, \mathcal{R}(p^{-1}, q^{-1})} [W_2]_{m_2, \mathcal{R}(p^{-1}, q^{-1})}}{(\phi_1^{n+W_2} \oplus \alpha_2 \phi_2^{n+W_2})_{\mathcal{R}(p, q)}^{m_1}}\right) = \frac{[n + m_1 + m_2 - 1]_{m_1+m_2, \mathcal{R}(p, q)}}{\alpha_1^{-m_1} \alpha_2^{-m_2}}, \quad (2.12)$$

**Proof 2.7** According to the relations (2.6) and (2.7), we derive the  $\mathcal{R}(p^{-1}, q^{-1})$ -factorial moments of  $W_2$  given by (2.9). Moreover, the conditional distribution of the  $\mathcal{R}(p, q)$ -random variable  $W_1$ , given that  $W_2 = w_2$ , is a negative  $\mathcal{R}(p, q)$ -trinomial probability distribution of the first kind, with mass function:

$$P(W_1 = w_1 | W_2 = w_2) = \binom{n + w_1 + w_2 - 1}{w_1}_{\mathcal{R}(p, q)} \frac{\alpha_1^{w_1} \phi_1^{\binom{n-w_1}{2}} \phi_2^{\binom{w_1}{2}}}{(1 \oplus \alpha_1)_{\mathcal{R}(p, q)}^{n+w_1+w_2}}, \quad m_1 \in \mathbb{N} \cup \{0\}.$$

Using, once the relations (2.6) and (2.7), the conditional  $\mathcal{R}(p^{-1}, q^{-1})$ -factorial moments of  $W_1$ , given that  $W_2 = w_2$ , are given by (2.10).

Furthermore, the expected value of the  $\mathcal{R}(p^{-1}, q^{-1})$ -function of  $\underline{W} = (W_1, W_2)$

$$\Gamma := \frac{[W_1]_{m_1, \mathcal{R}(p^{-1}, q^{-1})}}{(\phi_1^{n+W_2} \oplus \alpha_2 \phi_2^{n+W_2})_{\mathcal{R}(p, q)}^{m_1}}, \quad m_1 \in \mathbb{N} \cup \{0\}$$

can be calculated according to the relation

$$\begin{aligned} E(\Gamma) &= E(E(\Gamma | W_2)) \\ &= \alpha_1^{m_1} E\left(\frac{[n + W_2 + m_1 - 1]_{m_1, \mathcal{R}(p, q)}}{(\phi_1^{n+W_2} \oplus \alpha_2 \phi_2^{n+W_2})_{\mathcal{R}(p, q)}^{m_1}}\right). \end{aligned}$$



Since

$$E\left(\frac{[n + W_2 + m_1 - 1]_{m_1, \mathcal{R}(p, q)}}{(\phi_1^{n+W_2} \oplus \alpha_2 \phi_2^{n+W_2})_{\mathcal{R}(p, q)}^{m_1}}\right) = \sum_{w_2=0}^{\infty} [n + w_2 + m_1 - 1]_{m_1, \mathcal{R}(p, q)} \\ \times \begin{bmatrix} n + w_2 - 1 \\ w_2 \end{bmatrix}_{\mathcal{R}(p, q)} \frac{\alpha_2^{w_2} \phi_1^{\binom{n-w_2}{2}} \phi_2^{\binom{w_2}{2}}}{(1 \oplus \alpha_2)_{\mathcal{R}(p, q)}^{n+m_1+w_2}}.$$

From the relation

$$[n + w_2 + m_1 - 1]_{m_1, \mathcal{R}(p, q)} \begin{bmatrix} n + w_2 - 1 \\ w_2 \end{bmatrix}_{\mathcal{R}(p, q)} = [n + m_1 - 1]_{m_1, \mathcal{R}(p, q)} \begin{bmatrix} n + m_1 + w_2 - 1 \\ w_2 \end{bmatrix}_{\mathcal{R}(p, q)}$$

and the negative  $\mathcal{R}(p, q)$ -binomial formula [8], we have:

$$E\left(\frac{[n + W_2 + m_1 - 1]_{m_1, \mathcal{R}(p, q)}}{(\phi_1^{n+W_2} \oplus \alpha_2 \phi_2^{n+W_2})_{\mathcal{R}(p, q)}^{m_1}}\right) = [n + m_1 - 1]_{m_1, \mathcal{R}(p, q)} \sum_{w_2=0}^{\infty} \begin{bmatrix} n + m_1 + w_2 - 1 \\ w_2 \end{bmatrix}_{\mathcal{R}(p, q)} \\ \times \frac{\alpha_2^{w_2} \phi_1^{\binom{n-w_2}{2}} \phi_2^{\binom{w_2}{2}}}{(1 \oplus \alpha_2)_{\mathcal{R}(p, q)}^{n+m_1+w_2}} \\ = [n + m_1 - 1]_{m_1, \mathcal{R}(p, q)}.$$

Thus, the relation (2.11) holds. Analogously, the expected value of the  $\mathcal{R}(p^{-1}, q^{-1})$ -function of  $\underline{W} = (W_1, W_2)$

$$\Lambda := \frac{[W_1]_{m_1, \mathcal{R}(p^{-1}, q^{-1})} [W_2]_{m_2, \mathcal{R}(p^{-1}, q^{-1})}}{(\phi_1^{n+W_2} \oplus \alpha_2 \phi_2^{n+W_2})_{\mathcal{R}(p, q)}^{m_1}}, \quad m_1 \in \mathbb{N} \cup \{0\}, \quad m_2 \in \mathbb{N} \cup \{0\}$$

may be computed using the relation

$$E(\Lambda) = E(E(\Lambda|W_2)) \\ = \alpha_1^{m_1} E\left(\frac{[W_2]_{m_2, \mathcal{R}(p^{-1}, q^{-1})} [n + W_2 + m_1 - 1]_{m_1, \mathcal{R}(p, q)}}{(\phi_1^{n+W_2} \oplus \alpha_2 \phi_2^{n+W_2})_{\mathcal{R}(p, q)}^{m_1}}\right).$$

Since

$$E\left(\frac{[W_2]_{m_2, \mathcal{R}(p^{-1}, q^{-1})} [n + W_2 + m_1 - 1]_{m_1, \mathcal{R}(p, q)}}{(\phi_1^{n+W_2} \oplus \alpha_2 \phi_2^{n+W_2})_{\mathcal{R}(p, q)}^{m_1}}\right) = \sum_{w_2=m_2}^{\infty} \frac{[w_2]_{m_2, \mathcal{R}(p^{-1}, q^{-1})} [n + w_2 + m_1 - 1]_{m_1, \mathcal{R}(p, q)}}{(1 \oplus \alpha_2)_{\mathcal{R}(p, q)}^{m_1}} \\ \times \begin{bmatrix} n + w_2 - 1 \\ w_2 \end{bmatrix}_{\mathcal{R}(p, q)} \frac{\alpha_2^{w_2} \phi_1^{\binom{n-w_2}{2}} \phi_2^{\binom{w_2}{2}}}{(1 \oplus \alpha_2)_{\mathcal{R}(p, q)}^{n+w_2}}.$$

Using the relations

$$[w_2]_{m_2, \mathcal{R}(p^{-1}, q^{-1})} = (\phi_1 \phi_2)^{-w_2 m_2 + \binom{m_2+1}{2}} [w_2]_{m_2, \mathcal{R}(p, q)},$$

$$\binom{w_2}{2} + \binom{m_2 + 1}{2} - w_2 m_2 = \binom{w_2 - m_2}{2}$$

and

$$\begin{aligned} [w_2]_{m_2, \mathcal{R}(p, q)} [n + w_2 + m_1 - 1]_{m_1, \mathcal{R}(p, q)} \begin{bmatrix} n + w_2 - 1 \\ w_2 \end{bmatrix}_{\mathcal{R}(p, q)} &= [n + m_1 + m_2 - 1]_{m_1 + m_2, \mathcal{R}(p, q)} \\ &\times \begin{bmatrix} n + m_1 + w_2 - 1 \\ w_2 - m_2 \end{bmatrix}_{\mathcal{R}(p, q)}, \end{aligned}$$

together with the negative  $\mathcal{R}(p, q)$ -binomial formula [8], we get

$$E\left(\frac{[W_2]_{m_2, \mathcal{R}(p^{-1}, q^{-1})} [n + W_2 + m_1 - 1]_{m_1, \mathcal{R}(p, q)}}{(\phi_1^{n+W_2} \oplus \alpha_2 \phi_2^{n+W_2})_{\mathcal{R}(p, q)}^{m_1}}\right) = [n + m_2 + m_1 - 1]_{m_1 + m_2, \mathcal{R}(p, q)} \alpha_2^{m_2}$$

and the proof is achieved.

**Corollary 2.8** The  $\mathcal{R}(p, q)$ -covariance of the  $\mathcal{R}(p, q)$ -random variables  $\widehat{W} := (\phi_1^{n+W_2} \oplus \alpha_2 \phi_2^{n+W_2})^{-1} [W_1]_{m_1, \mathcal{R}(p^{-1}, q^{-1})}$  and  $\overline{W} := [W_2]_{m_2, \mathcal{R}(p^{-1}, q^{-1})}$  is determined by:

$$\text{Cov}(\widehat{W}, \overline{W}) = [n]_{\mathcal{R}(p, q)} \alpha_1 \alpha_2 ([n + 1]_{\mathcal{R}(p, q)} - [n]_{\mathcal{R}(p, q)}),$$

where the  $\mathcal{R}(p, q)$ -random vector  $\underline{W} = (W_1, W_2)$  follows the negative  $\mathcal{R}(p, q)$ -trinomial probability distribution of the first kind, with parameters  $n$ ,  $\underline{\alpha} = (\alpha_1, \alpha_2)$ ,  $p$ , and,  $q$ .

**Proof 2.9** By definition, the  $\mathcal{R}(p, q)$ -covariance of  $\widehat{W}$  and  $\overline{W}$  is given by

$$\text{Cov}(\widehat{W}, \overline{W}) = E\left(\widehat{W} \overline{W}\right) - E(\widehat{W}) E(\overline{W}).$$

Taking  $m_1 = m_2 = 1$ , in the relations (2.9), (2.11), and (2.12), the result follows.

### 2.3 $\mathcal{R}(p, q)$ -trinomial distribution of the second kind

The probability function of the  $\mathcal{R}(p, q)$ -random vector  $\underline{X} = (X_1, X_2)$  of the  $\mathcal{R}(p, q)$ -trinomial probability distribution of the second kind, with parameters  $n$ ,  $\underline{\beta} = (\beta_1, \beta_2)$ ,  $p$ , and,  $q$  is given by:

$$P(\underline{X} = \underline{x}) = \begin{bmatrix} n \\ x_1, x_2 \end{bmatrix}_{\mathcal{R}(p, q)} \beta_1^{x_1} \beta_2^{x_2} (1 \ominus \beta_1)_{\mathcal{R}(p, q)}^{n-x_1} (1 \ominus \beta_2)_{\mathcal{R}(p, q)}^{n-x_1-x_2}, \quad (2.13)$$

where  $x_j \in \{0, 1, \dots, n\}$ ,  $x_1 + x_2 \leq n$ ,  $s_j = \sum_{i=1}^j x_i$ ,  $0 < \beta_j < 1$ , and  $j \in \{1, 2\}$ .

**Theorem 2.10** The  $\mathcal{R}(p, q)$ -factorial moments of the  $\mathcal{R}(p, q)$ -trinomial probability distribution of the second kind, with parameters  $n$ ,  $\underline{\beta} = (\beta_1, \beta_2)$ ,  $p$ , and,  $q$ , are given by:

$$E([X_1]_{m_1, \mathcal{R}(p, q)}) = \beta_1^{m_1} [n]_{m_1, \mathcal{R}(p, q)}, m_1 \in \{0, 1, \dots, n\}, \quad (2.14)$$

$$E([X_2]_{m_2, \mathcal{R}(p, q)} | X_1 = x_1) = \beta_2^{m_2} [n - x_1]_{m_2, \mathcal{R}(p, q)}, \quad (2.15)$$

where  $m_2 \in \{0, 1, \dots, n - x_1\}$ ,

$$E\left(\frac{[X_2]_{m_2, \mathcal{R}(p, q)}}{\left(\phi_1^{n-m_2-X_1} \ominus \beta_1 \phi_2^{n-m_2-X_1}\right)_{\mathcal{R}(p, q)}^{m_2}}\right) = \beta_2^{m_2} [n]_{m_2, \mathcal{R}(p, q)}, \quad (2.16)$$

with  $m_2 \in \{0, 1, \dots, n\}$ , and

$$E\left(\frac{[X_1]_{m_1, \mathcal{R}(p, q)} [X_2]_{m_2, \mathcal{R}(p, q)}}{\left(\phi_1^{n-m_2-X_1} \ominus \beta_1 \phi_2^{n-m_2-X_1}\right)_{\mathcal{R}(p, q)}^{m_2}}\right) = \beta_1^{m_1} \beta_2^{m_2} [n]_{m_1+m_2, \mathcal{R}(p, q)}, \quad (2.17)$$

where  $m_1 \in \{0, 1, \dots, n - m_2\}$  and  $m_2 \in \{0, 1, \dots, n\}$ .

**Proof 2.11** The  $\mathcal{R}(p, q)$ - random variable  $X_1$  obey the  $\mathcal{R}(p, q)$ - binomial probability distribution of the second kind, with mass function:

$$P(X_1 = x_1) = \begin{bmatrix} n \\ x_1 \end{bmatrix}_{\mathcal{R}(p, q)} \beta_1^{x_1} (1 \ominus \beta_1)_{\mathcal{R}(p, q)}^{n-x_1}, x_1 \in \{0, 1, \dots, n\}.$$

Thus, using [7], the relation (2.14) follows. Furthermore, the conditional distribution of the  $\mathcal{R}(p, q)$ - random variable  $X_2$ , given that  $X_1 = x_1$ , is a  $\mathcal{R}(p, q)$ -binomial probability distribution of the second kind, with mass density

$$P(X_2 = x_2 | X_1 = x_1) = \begin{bmatrix} n - x_1 \\ x_2 \end{bmatrix}_{\mathcal{R}(p, q)} \beta_2^{x_2} (1 \ominus \beta_2)_{\mathcal{R}(p, q)}^{n-x_1-x_2}.$$

According again to [7], the conditional  $\mathcal{R}(p, q)$ - factorial moments of  $X_2$ , given that  $X_1 = x_1$ , are furnished by (2.15). The expected value of the  $\mathcal{R}(p, q)$ -function of  $\underline{X} = (X_1, X_2)$

$$\Delta := \frac{[X_2]_{m_2, \mathcal{R}(p, q)}}{\left(\phi_1^{n-m_2-X_1} \ominus \beta_1 \phi_2^{n-m_2-X_1}\right)_{\mathcal{R}(p, q)}^{m_2}}, m_2 \in \{0, 1, \dots, n\}$$

can be computed according to the relation

$$\begin{aligned} E(\Delta) &= E[E(\Delta | X_1)] \\ &= \beta_2^{m_2} E\left[\frac{[n - X_1]_{m_2, \mathcal{R}(p, q)}}{\left(\phi_1^{n-m_2-X_1} \ominus \beta_1 \phi_2^{n-m_2-X_1}\right)_{\mathcal{R}(p, q)}^{m_2}}\right]. \end{aligned}$$

Since

$$E \left[ \frac{[n - X_1]_{m_2, \mathcal{R}(p, q)}}{(\phi_1^{n-m_2-X_1} \ominus \beta_1 \phi_2^{n-m_2-X_1})_{\mathcal{R}(p, q)}^{m_2}} \right] = \sum_{x_1=0}^{n-m_2} [n - x_1]_{m_2, \mathcal{R}(p, q)} \begin{bmatrix} n \\ x_1 \end{bmatrix}_{\mathcal{R}(p, q)} \times \frac{\beta_1^{x_1} (1 \ominus \beta_1)_{\mathcal{R}(p, q)}^{n-x_1}}{(\phi_1^{n-m_2-x_1} \ominus \beta_1 \phi_2^{n-m_2-x_1})_{\mathcal{R}(p, q)}^{m_2}}.$$

Using the relations:

$$[n - x_1]_{m_2, \mathcal{R}(p, q)} \begin{bmatrix} n \\ x_1 \end{bmatrix}_{\mathcal{R}(p, q)} = [n]_{m_2, \mathcal{R}(p, q)} \begin{bmatrix} n - m_2 \\ x_1 \end{bmatrix}_{\mathcal{R}(p, q)},$$

$$(1 \ominus \beta_1)_{\mathcal{R}(p, q)}^{n-x_1} = (1 \ominus \beta_1)_{\mathcal{R}(p, q)}^{n-m_2-x_1} (\phi_1^{n-m_2-x_1} \ominus \beta_1 \phi_2^{n-m_2-x_1})_{\mathcal{R}(p, q)}^{m_2}, \quad (2.18)$$

and the  $\mathcal{R}(p, q)$ -binomial formula [8], we have:

$$E \left[ \frac{[n - X_1]_{m_2, \mathcal{R}(p, q)}}{(\phi_1^{n-m_2-X_1} \ominus \beta_1 \phi_2^{n-m_2-X_1})_{\mathcal{R}(p, q)}^{m_2}} \right] = [n]_{m_2, \mathcal{R}(p, q)} \sum_{x_1=0}^{n-m_2} \begin{bmatrix} n - m_2 \\ x_1 \end{bmatrix}_{\mathcal{R}(p, q)} \times \beta_1^{m_1} (1 \ominus \beta_1)_{\mathcal{R}(p, q)}^{n-m_2-x_1}.$$

Thus,

$$E \left( \frac{[X_2]_{m_2, \mathcal{R}(p, q)}}{(\phi_1^{n-m_2-X_1} \ominus \beta_1 \phi_2^{n-m_2-X_1})_{\mathcal{R}(p, q)}^{m_2}} \right) = \beta_2^{m_2} [n]_{m_2, \mathcal{R}(p, q)}.$$

Similarly, the expected value of the  $\mathcal{R}(p, q)$ -function of  $\underline{X} = (X_1, X_2)$

$$\nabla := \frac{[X_1]_{m_1, \mathcal{R}(p, q)} [X_2]_{m_2, \mathcal{R}(p, q)}}{(\phi_1^{n-m_2-X_1} \ominus \beta_1 \phi_2^{n-m_2-X_1})_{\mathcal{R}(p, q)}^{m_2}}, \quad m_1 \in \{0, 1, \dots, n - m_2\}, \quad m_2 \in \{0, 1, \dots, n\}$$

may be calculated using the relation:

$$\begin{aligned} E(\nabla) &= E[E(\nabla | X_1)] \\ &= \beta_2^{m_2} E \left[ \frac{[X_1]_{m_1, \mathcal{R}(p, q)} [n - X_1]_{m_2, \mathcal{R}(p, q)}}{(\phi_1^{n-m_2-X_1} \ominus \beta_1 \phi_2^{n-m_2-X_1})_{\mathcal{R}(p, q)}^{m_2}} \right]. \end{aligned}$$

Since

$$E \left[ \frac{[X_1]_{m_1, \mathcal{R}(p, q)} [n - X_1]_{m_2, \mathcal{R}(p, q)}}{(\phi_1^{n-m_2-X_1} \ominus \beta_1 \phi_2^{n-m_2-X_1})_{\mathcal{R}(p, q)}^{m_2}} \right] = \sum_{x_1=0}^{n-m_2} [x_1]_{m_1, \mathcal{R}(p, q)} [n - x_1]_{m_2, \mathcal{R}(p, q)} \begin{bmatrix} n \\ x_1 \end{bmatrix}_{\mathcal{R}(p, q)} \times \frac{\beta_1^{x_1} (1 \ominus \beta_1)_{\mathcal{R}(p, q)}^{n-x_1}}{(\phi_1^{n-m_2-x_1} \ominus \beta_1 \phi_2^{n-m_2-x_1})_{\mathcal{R}(p, q)}^{m_2}}.$$

From the  $\mathcal{R}(p, q)$ -binomial formula [8], the relations (2.18), and

$$[x_1]_{m_1, \mathcal{R}(p, q)} [n - x_1]_{m_2, \mathcal{R}(p, q)} \begin{bmatrix} n \\ x_1 \end{bmatrix}_{\mathcal{R}(p, q)} = [n]_{m_1 + m_2, \mathcal{R}(p, q)} \begin{bmatrix} n - m_1 - m_2 \\ x_1 - m_1 \end{bmatrix}_{\mathcal{R}(p, q)},$$

we obtain:

$$E \left[ \frac{[X_1]_{m_1, \mathcal{R}(p, q)} [n - X_1]_{m_2, \mathcal{R}(p, q)}}{(\phi_1^{n - m_2 - X_1} \ominus \beta_1 \phi_2^{n - m_2 - X_1})_{\mathcal{R}(p, q)}^{m_2}} \right] = [n]_{m_1 + m_2, \mathcal{R}(p, q)} \beta_1^{m_1}$$

and the relation (2.17) follows.

**Corollary 2.12** The  $\mathcal{R}(p, q)$ - covariance of the functions  $\widehat{X} := [X_1]_{\mathcal{R}(p, q)}$  and  $\overline{X} := (\phi_1^{n-1-X_1} - \beta_1 \phi_2^{n-1-X_1})^{-1} [X_2]_{\mathcal{R}(p, q)}$  is given by:

$$\text{Cov}(\widehat{X}, \overline{X}) = [n]_{\mathcal{R}(p, q)} \beta_1 \beta_2 ([n - 1]_{\mathcal{R}(p, q)} - [n]_{\mathcal{R}(p, q)}),$$

with  $\underline{X} = (X_1, X_2)$  a  $\mathcal{R}(p, q)$ -random vector satisfying the  $\mathcal{R}(p, q)$ -trinomial probability distribution of the second kind, with parameters  $n, \underline{\beta} = (\beta_1, \beta_2), p$ , and,  $q$ .

**Proof 2.13** Taking  $m_1 = m_2 = 1$  in the relations (2.14), (2.16), and (2.17), we compute  $E(\widehat{X} \overline{X})$  and  $E(\overline{X}) E(\widehat{X})$ . Therefore, the proof is achieved.

## 2.4 Negative $\mathcal{R}(p, q)$ -trinomial distribution of the second kind

Let  $T_n$  be the number of failures until the occurrence of the  $n^{\text{th}}$  success, in a sequence of independent geometric sequences of trials. Then, the distribution of the  $\mathcal{R}(p, q)$ - random variable  $T_n$  is called negative  $\mathcal{R}(p, q)$ -binomial distribution of the second kind, with parameters  $n, \beta, p$ , and,  $q$ .

**Lemma 2.14** The probability function of the negative  $\mathcal{R}(p, q)$ - binomial probability distribution of the second kind, with parameters  $n, \beta, p$ , and,  $q$ , is given by :

$$P(T = t) = \begin{bmatrix} n + t - 1 \\ t \end{bmatrix}_{\mathcal{R}(p, q)} \beta^t (1 \ominus \beta)_{\mathcal{R}(p, q)}^n, t \in \mathbb{N} \cup \{0\}. \quad (2.19)$$

Furthermore, its  $\mathcal{R}(p, q)$ -factorial moments are presented as follows:

$$E([T]_{m, \mathcal{R}(p, q)}) = \frac{[n + m - 1]_{m, \mathcal{R}(p, q)} \beta^m}{(\phi_1^n \ominus \beta_2 \phi_2^n)_{\mathcal{R}(p, q)}^m}, m \in \mathbb{N}. \quad (2.20)$$

Let  $\underline{V} = (V_1, V_2)$  be a  $\mathcal{R}(p, q)$ -random vector obeying the negative  $\mathcal{R}(p, q)$ - trinomial probability distribution of the second kind, with parameters  $n, \underline{\beta} = (\beta_1, \beta_2), p$ , and,  $q$ . Then, its mass function is given by:

$$P(\underline{V} = \underline{v}) = \begin{bmatrix} n + v_1 + v_2 - 1 \\ v_1, v_2 \end{bmatrix}_{\mathcal{R}(p, q)} \beta_1^{v_1} \beta_2^{v_2} (1 \ominus \beta_1)_{\mathcal{R}(p, q)}^{n+v_2} (1 \ominus \beta_2)_{\mathcal{R}(p, q)}^n, v_j \in \mathbb{N} \cup \{0\},$$

where  $0 < \beta_j < 1$ , and  $j \in \{1, 2\}$ .

**Theorem 2.15** For  $m_1 \in \mathbb{N} \cup \{0\}$  and  $m_2 \in \mathbb{N} \cup \{0\}$ , the  $\mathcal{R}(p, q)$ -factorial moments of the negative  $\mathcal{R}(p, q)$ -trinomial probability distribution of the second kind, with parameters  $n, \underline{\beta} = (\beta_1, \beta_2), p$ , and,  $q$  are presented as follows:

$$E([V_2]_{m_2, \mathcal{R}(p, q)}) = \frac{\beta_2^{m_2} [n + m_2 - 1]_{m_2, \mathcal{R}(p, q)}}{(\phi_1^n \ominus \beta_2 \phi_2^n)_{\mathcal{R}(p, q)}^{m_2}}, \quad (2.21)$$

$$E([V_1]_{m_1, \mathcal{R}(p, q)} | V_2 = v_2) = \frac{\beta_1^{m_1} [n + v_2 + m_1 - 1]_{m_1, \mathcal{R}(p, q)}}{(\phi_1^{n+v_2} \ominus \beta_1 \phi_2^{n+v_2})_{\mathcal{R}(p, q)}^{m_1}}, \quad (2.22)$$

$$E\left(\frac{[V_1]_{m_1, \mathcal{R}(p, q)}}{(\phi_1^{n+V_2} \ominus \beta_1 \phi_2^{n+V_2})_{\mathcal{R}(p, q)}^{-m_1}}\right) = \frac{\beta_1^{m_1} [n + m_1 - 1]_{m_1, \mathcal{R}(p, q)}}{(\phi_1^n \ominus \beta_2 \phi_2^n)_{\mathcal{R}(p, q)}^{m_1}}, \quad (2.23)$$

and

$$E\left(\frac{[V_1]_{m_1, \mathcal{R}(p, q)} [V_2]_{m_2, \mathcal{R}(p, q)}}{(\phi_1^{n+V_2} \ominus \beta_1 \phi_2^{n+V_2})_{\mathcal{R}(p, q)}^{-m_2}}\right) = \frac{\beta_1^{m_1} \beta_2^{m_2} [n + m_1 + m_2 - 1]_{m_1+m_2, \mathcal{R}(p, q)}}{(\phi_1^n \ominus \beta_2 \phi_2^n)_{\mathcal{R}(p, q)}^{m_1+m_2}}. \quad (2.24)$$

**Proof 2.16** The relation (2.21) comes from (2.19) and (2.20). Furthermore, the conditional distribution of the  $\mathcal{R}(p, q)$ - random variable  $V_1$ , given that  $V_2 = v_2$ , is a negative  $\mathcal{R}(p, q)$ -binomial distribution of the second kind, with probability function:

$$P(V_1 = v_1 | V_2 = v_2) = \left[ \begin{matrix} n + v_1 + v_2 - 1 \\ v_1 \end{matrix} \right]_{\mathcal{R}(p, q)} \beta_1^{v_1} (1 \ominus \beta_1)_{\mathcal{R}(p, q)}^{n+v_2}, \quad v \in \mathbb{N} \cup \{0\}.$$

Using once the relations (2.19) and (2.20), the conditional  $\mathcal{R}(p, q)$ - factorial moments of  $V_1$ , given that  $V_2 = v_2$ , are given by (2.22). Besides, the expected value of the  $\mathcal{R}(p, q)$ -function of  $\underline{V} = (V_1, V_2)$

$$\widehat{\Delta} := \frac{[V_1]_{m_1, \mathcal{R}(p, q)}}{(\phi_1^{n+V_2} \ominus \beta_2 \phi_2^{n+V_2})_{\mathcal{R}(p, q)}^{m_1}}, \quad m_1 \in \mathbb{N} \cup \{0\}$$

may be evaluated according to the relation:

$$\begin{aligned} E(\widehat{\Delta}) &= E[E(\widehat{\Delta} | V_2)] \\ &= \beta_1^{m_1} E\left([n + V_2 + m_1 - 1]_{m_1, \mathcal{R}(p, q)}\right). \end{aligned}$$

Since

$$\begin{aligned} E\left([n + V_2 + m_1 - 1]_{m_1, \mathcal{R}(p, q)}\right) &= \sum_{v_2=0}^{\infty} [n + v_2 + m_1 - 1]_{m_1, \mathcal{R}(p, q)} \\ &\quad \times \left[ \begin{matrix} n + v_2 - 1 \\ v_2 \end{matrix} \right]_{\mathcal{R}(p, q)} \beta_2^{m_2} (1 \ominus \beta_2)_{\mathcal{R}(p, q)}^n. \end{aligned}$$

From the relation

$$[n + v_2 + m_1 - 1]_{m_1, \mathcal{R}(p, q)} \begin{bmatrix} n + v_2 - 1 \\ v_2 \end{bmatrix}_{\mathcal{R}(p, q)} = [n + m_1 - 1]_{m_1, \mathcal{R}(p, q)} \begin{bmatrix} n + m_1 + v_2 - 1 \\ v_2 \end{bmatrix}_{\mathcal{R}(p, q)},$$

and the negative  $\mathcal{R}(p, q)$ -binomial formula [8], we have:

$$\begin{aligned} E\left([n + V_2 + m_1 - 1]_{m_1, \mathcal{R}(p, q)}\right) &= [n + m_1 - 1]_{m_1, \mathcal{R}(p, q)} (1 \ominus \beta_2)_{\mathcal{R}(p, q)}^n \sum_{v_2=0}^{\infty} \\ &\times \begin{bmatrix} n + v_2 - 1 \\ v_2 \end{bmatrix}_{\mathcal{R}(p, q)} \beta_2^{m_2} \\ &= \frac{[n + m_1 - 1]_{m_1, \mathcal{R}(p, q)}}{(\phi_1^n \ominus \beta_2 \phi_2^n)_{\mathcal{R}(p, q)}^{m_1}}. \end{aligned}$$

Thus, the relation (2.23) holds.

So, the expected value of the  $\mathcal{R}(p, q)$ -function of  $\underline{V} = (V_1, V_2)$

$$\widehat{\nabla} := \frac{[V_1]_{m_1, \mathcal{R}(p, q)} [V_2]_{m_2, \mathcal{R}(p, q)}}{(\phi_1^{n+V_2} \ominus \beta_1 \phi_2^{n+V_2})_{\mathcal{R}(p, q)}^{-m_1}}, \quad m_1 \in \mathbb{N} \cup \{0\}, \quad m_2 \in \mathbb{N} \cup \{0\}$$

can be calculated using the relation:

$$\begin{aligned} E(\widehat{\nabla}) &= E[E(\widehat{\nabla} | V_2)] \\ &= \beta_1^{m_1} E\left[[V_2]_{m_2, \mathcal{R}(p, q)} [n + V_2 + m_1 - 1]_{m_1, \mathcal{R}(p, q)}\right]. \end{aligned}$$

Since

$$\begin{aligned} E\left[[V_2]_{m_2, \mathcal{R}(p, q)} [n + V_2 + m_1 - 1]_{m_1, \mathcal{R}(p, q)}\right] &= \sum_{v_2=m_2=0}^{\infty} [v_2]_{m_2, \mathcal{R}(p, q)} [n + v_2 + m_1 - 1]_{m_1, \mathcal{R}(p, q)} \\ &\times \begin{bmatrix} n + v_2 - 1 \\ v_2 \end{bmatrix}_{\mathcal{R}(p, q)} \beta_2^{m_2} (1 \ominus \beta_2)_{\mathcal{R}(p, q)}^n. \end{aligned}$$

From the  $\mathcal{R}(p, q)$ -binomial formula [8], the relations (2.18), and

$$\begin{aligned} [v_2]_{m_2, \mathcal{R}(p, q)} [n + v_2 + m_1 - 1]_{m_1, \mathcal{R}(p, q)} \begin{bmatrix} n + v_2 - 1 \\ v_2 \end{bmatrix}_{\mathcal{R}(p, q)} &= [n + m_1 + m_2 - 1]_{m_1 + m_2, \mathcal{R}(p, q)} \\ &\times \begin{bmatrix} n + m_1 + v_2 - 1 \\ v_2 - m_2 \end{bmatrix}_{\mathcal{R}(p, q)}, \end{aligned}$$

we obtain:

$$E\left[[V_2]_{m_2, \mathcal{R}(p, q)} [n + V_2 + m_1 - 1]_{m_1, \mathcal{R}(p, q)}\right] = \frac{[n + m_1 + m_2 - 1]_{m_1 + m_2, \mathcal{R}(p, q)} \beta_2^{m_2}}{(\phi_1^n \ominus \beta_2 \phi_2^n)_{\mathcal{R}(p, q)}^{m_1 + m_2}}$$

and the relation (2.24) follows.

**Corollary 2.17** *The  $\mathcal{R}(p, q)$ -covariance of the functions and  $\widehat{V} := (\phi_1^{n+V_2} \ominus \beta_2 \phi_2^{n+V_2})[V_1]_{m_2, \mathcal{R}(p, q)}$  and  $\overline{V} := [V_2]_{m_1, \mathcal{R}(p, q)}$  is given by:*

$$Cov(\widehat{V}, \overline{V}) = \frac{[n]_{\mathcal{R}(p, q)} \beta_1 \beta_2}{(\phi_1^n - \beta_2 \phi_2^n)} \left( \frac{[n+1]_{\mathcal{R}(p, q)}}{(\phi_1^{n+1} - \beta_2 \phi_2^{n+1})} - \frac{[n]_{\mathcal{R}(p, q)}}{(\phi_1^n - \beta_2 \phi_2^n)} \right), \quad (2.25)$$

where  $\underline{V} = (V_1, V_2)$  is a  $\mathcal{R}(p, q)$ -random vector satisfying the negative  $\mathcal{R}(p, q)$ -trinomial probability distribution of the second kind, with parameters  $n$ ,  $\underline{\beta} = (\beta_1, \beta_2)$ ,  $p$ , and,  $q$ .

**Proof 2.18** *Putting  $m_1 = m_2 = 1$ , in the relations (2.21), (2.23), and (2.24), we obtain:*

$$E(\widehat{V}\overline{V}) = \frac{[n+1]_{\mathcal{R}(p, q)} [n]_{\mathcal{R}(p, q)} \beta_1 \beta_2}{(\phi_1^n - \beta_2 \phi_2^n)(\phi_1^{n+1} - \beta_2 \phi_2^{n+1})} \text{ and } E(\widehat{V})E(\overline{V}) = \frac{[n]_{\mathcal{R}(p, q)} [n]_{\mathcal{R}(p, q)} \beta_2 \beta_1}{(\phi_1^n - \beta_2 \phi_2^n)^2}.$$

After computation, the result follows.

### 3 Trinomial distributions and particular quantum algebras

In this section, we deduce the  $\mathcal{R}(p, q)$ -trinomial probability distribution of the first and second kind, their negative and properties from some quantum algebra existing in the literature.

#### 3.1 Trinomial distribution and Biedenharn-Macfarlane algebra [1, 13]

Taking  $\mathcal{R}(x) = \frac{x-x^{-1}}{q-q^{-1}}$ , we obtain:

(i) The  $q$ -trinomial distribution of the first kind is given by

$$P(Y_1 = y_1, Y_2 = y_2) = \begin{bmatrix} n \\ y_1, y_2 \end{bmatrix}_q \frac{\alpha_1^{y_1} \alpha_2^{y_2} q^{\binom{n-y_1}{2} + \binom{n-y_2}{2}} q^{-\binom{y_1}{2} - \binom{y_2}{2}}}{(1 \oplus \alpha_1)_q^n (1 \oplus \alpha_2)_q^{n-y_1}} \quad (3.1)$$

and the  $q$ -factorial moments are:

$$\begin{aligned} E([Y_1]_{m_1, q}) &= \frac{[n]_{m_1, q} \alpha_1^{m_1} q^{-\binom{m_1}{2}}}{(1 \oplus \alpha_1)_q^{m_1}}, \quad m_1 \in \{0, 1, \dots, n\}, \\ E([Y_2]_{m_2, q} | Y_1 = y_1) &= \frac{[n - y_1]_{m_2, q} \alpha_2^{m_2} q^{-\binom{m_2}{2}}}{(1 \oplus \alpha_2)_q^{m_2}}, \quad m_2 \in \{0, 1, \dots, n - y_1\}, \\ E([Y_2]_{m_2, p, q}) &= \frac{[n]_{m_2, p, q} \alpha_2^{m_2} q^{\binom{m_2}{2}} p^{\binom{m_2}{2} + m_2(n-m_2)}}{(1 \oplus \alpha_2)_{p, q}^{m_2} (p^{n-m_2} \oplus \alpha_1 q^{n-m_2})_{p, q}^{m_2}}, \quad m_2 \in \{0, 1, \dots, n\}, \end{aligned}$$

and

$$E([Y_1]_{m_1, q} [Y_2]_{m_2, q}) = \frac{[n]_{m_1+m_2, q} \alpha_1^{m_1} \alpha_2^{m_2} q^{\binom{m_2}{2} + m_2(n-m_2)} q^{-\binom{m_1}{2} - \binom{m_2}{2}}}{(1 \oplus \alpha_1)_q^{m_1} (1 \oplus \alpha_2)_q^{m_2} (p^{n-m_2} \oplus \alpha_1 q^{n-m_2})_q^{m_2}},$$



where  $m_1 \in \{0, 1, \dots, n - m_2\}$  and  $m_2 \in \{0, 1, \dots, n\}$ . Moreover, the  $q$ -covariance is derived as follows:

$$Cov([Y_1]_q, [Y_2]_q) = \frac{q^{n-1} \alpha_1 \alpha_2 [n]_q ([n-1]_q - [n]_q)}{(1 + \alpha_1)(1 + \alpha_2)(q^{n-1} + \alpha_1 q^{-n+1})}.$$

- (ii) The negative  $q$ -trinomial distribution of the first kind, with parameters  $n$ ,  $\underline{\alpha} = (\alpha_1, \alpha_2)$ , and,  $q$  is presented by:

$$P(\underline{W} = \underline{w}) = \binom{n + w_1 + w_2 - 1}{w_1, w_2}_q \frac{\alpha_1^{w_1} \alpha_2^{w_2} q^{\binom{n-w_1}{2} + \binom{n-w_2}{2}} q^{-\binom{w_1}{2} - \binom{w_2}{2}}}{(1 \oplus \alpha_1)_q^{n+w_1+w_2} (1 \oplus \alpha_2)_q^{n+w_2}},$$

where  $w_j \in \mathbb{N} \cup \{0\}$ ,  $0 < \alpha_j < 1$ , and  $j \in \{1, 2\}$ . Besides, for  $m_1 \in \mathbb{N} \cup \{0\}$  and  $m_2 \in \mathbb{N} \cup \{0\}$ ,

its  $q^{-1}$ -factorial moments are given as follows:

$$E([W_2]_{m_2, q^{-1}}) = [n + m_2 - 1]_{m_2, q} \alpha_2^{m_2}, \quad m_2 \in \mathbb{N} \cup \{0\},$$

$$E([W_1]_{m_1, q^{-1}} | W_2 = w_2) = [n + w_2 + m_1 - 1]_{m_1, q} \alpha_1^{m_1}, \quad m_1 \in \mathbb{N} \cup \{0\},$$

$$E\left(\frac{[W_1]_{m_1, q^{-1}}}{(p^{n+W_2} \oplus \alpha_2 q^{-n-W_2})_q^{m_1}}\right) = [n + m_1 - 1]_{m_1, q} \alpha_1^{m_1}, \quad m_1 \in \mathbb{N} \cup \{0\},$$

and

$$E\left(\frac{[W_1]_{m_1, q^{-1}} [W_2]_{m_2, q^{-1}}}{(p^{n+W_2} \oplus \alpha_2 q^{-n-W_2})_q^{m_1}}\right) = [n + m_1 + m_2 - 1]_{m_1+m_2, q} \alpha_1^{m_1} \alpha_2^{m_2},$$

Furthermore, the covariance of  $\widehat{W} := (q^{n+W_2} \oplus \alpha_2 q^{-n-W_2})^{-1} [W_1]_{m_1, q^{-1}}$  and  $\overline{W} := [W_2]_{m_2, q^{-1}}$  is determined by:

$$Cov(\widehat{W}, \overline{W}) = [n]_q \alpha_1 \alpha_2 ([n+1]_q - [n]_q).$$

- (iii) The  $q$ -trinomial distribution of the second kind, with parameters  $n$ ,  $\underline{\beta} = (\beta_1, \beta_2)$ , and,  $q$  is given by:

$$P(X_1 = x_1, X_2 = x_2) = \binom{n}{x_1, x_2}_q \beta_1^{x_1} \beta_2^{x_2} (1 \ominus \beta_1)_q^{n-x_1} (1 \ominus \beta_2)_q^{n-x_1-x_2},$$

where  $x_j \in \{0, 1, \dots, n\}$ ,  $x_1 + x_2 \leq n$ ,  $s_j = \sum_{i=1}^j x_i$ ,  $0 < \beta_j < 1$ , and  $j \in \{1, 2\}$ .

Besides, its  $q$ -factorial moments are given by:

$$E([X_1]_{m_1, q}) = [n]_{m_1, q} \beta_1^{m_1}, \quad m_1 \in \{0, 1, \dots, n\},$$

$$E([X_2]_{m_2, q} | X_1 = x_1) = [n - x_1]_{m_2, q} \beta_2^{m_2}, \quad m_2 \in \{0, 1, \dots, n - x_1\},$$

$$E\left(\frac{[X_2]_{m_2,q}}{(p^{n-m_2-X_1} \ominus \beta_1 q^{n-m_2-X_1})_q^{m_2}}\right) = [n]_{m_2,q} \beta_2^{m_2}, \quad m_2 \in \{0, 1, \dots, n\},$$

and

$$E\left(\frac{[X_1]_{m_1,q}[X_2]_{m_2,q}}{(p^{n-m_2-X_1} \ominus \beta_1 q^{n-m_2-X_1})_q^{m_2}}\right) = [n]_{m_1+m_2,q} \beta_1^{m_1} \beta_2^{m_2},$$

where  $m_1 \in \{0, 1, \dots, n - m_2\}$  and  $m_2 \in \{0, 1, \dots, n\}$ . Moreover, the  $q$ -covariance of the functions  $(q^{n-1-X_1} - \beta_1 q^{-n+1+X_1})^{-1}[X_2]_q$  and  $[X_1]_q$  is given by:

$$Cov\left([X_1]_q, (q^{n-1-X_1} - \beta_1 q^{-n+1+X_1})^{-1}[X_2]_q\right) = [n]_q \beta_1 \beta_2 ([n-1]_q - [n]_q).$$

(iii) The negative  $q$ -trinomial probability distribution of the second kind, with parameters  $n, \underline{\beta} = (\beta_1, \beta_2)$ , and,  $q$  is given by:

$$P(\underline{V} = \underline{v}) = \binom{n+v_1+v_2-1}{v_1, v_2}_q \beta_1^{v_1} \beta_2^{v_2} (1 \ominus \beta_1)_q^{n+v_2} (1 \ominus \beta_2)_q^n, \quad v_j \in \mathbb{N} \cup \{0\},$$

where  $0 < \beta_j < 1$ , and  $j \in \{1, 2\}$ . Moreover, its  $q$ -factorial moments of are presented as follows:

$$E([V_2]_{m_2,q}) = \frac{[n+m_2-1]_{m_2,q} \beta_2^{m_2}}{(p^n \ominus \beta_2 q^n)_q^{m_2}},$$

$$E([V_1]_{m_1,q} | V_2 = v_2) = \frac{[n+v_2+m_1-1]_{m_1,q} \beta_1^{m_1}}{(p^{n+v_2} \ominus \beta_1 q^{n+v_2})_q^{m_1}},$$

$$E\left(\frac{[V_1]_{m_1,q}}{(p^{n+V_2} \ominus \beta_1 q^{n+V_2})_q^{-m_1}}\right) = \frac{[n+m_1-1]_{m_1,q} \beta_1^{m_1}}{(p^n \ominus \beta_2 q^n)_q^{m_1}},$$

and

$$E\left(\frac{[V_1]_{m_1,q}[V_2]_{m_2,q}}{(p^{n+V_2} \ominus \beta_1 q^{n+V_2})_q^{-m_2}}\right) = \frac{[n+m_1+m_2-1]_{m_1+m_2,q} \beta_1^{m_1} \beta_2^{m_2}}{(p^n \ominus \beta_2 q^n)_q^{m_1+m_2}},$$

where  $m_1 \in \mathbb{N} \cup \{0\}$  and  $m_2 \in \mathbb{N} \cup \{0\}$ . Furthermore, the  $q$ -covariance of the functions and  $\widehat{V} := (q^{n+V_2} - \beta_2 q^{-n-V_2})[V_1]_{m_2,q}$  and  $\overline{V} := [V_2]_{m_1,q}$  is given by:

$$Cov(\widehat{V}, \overline{V}) = \frac{[n]_q \beta_1 \beta_2}{(q^n - \beta_2 q^{-n})} \left( \frac{[n+1]_q}{(q^{n+1} - \beta_2 q^{-n-1})} - \frac{[n]_q}{(q^n - \beta_2 q^{-n})} \right).$$

### 3.2 Trinomial distribution and Jagannathan-Srinivasa algebra [12]

For illustration, we investigate by taking  $\mathcal{R}(x, y) = (p - q)^{-1}(x - y)$  :

(i) The  $(p, q)$ -trinomial distribution of the first kind is given by

$$P(Y_1 = y_1, Y_2 = y_2) = \left[ \begin{matrix} n \\ y_1, y_2 \end{matrix} \right]_{p,q} \frac{\alpha_1^{y_1} \alpha_2^{y_2} p^{\binom{n-y_1}{2} + \binom{n-y_2}{2}} q^{\binom{y_1}{2} + \binom{y_2}{2}}}{(1 \oplus \alpha_1)_{p,q}^n (1 \oplus \alpha_2)_{p,q}^{n-y_1}} \quad (3.2)$$

and the  $(p, q)$ -factorial moments are:

$$\begin{aligned} E([Y_1]_{m_1, p, q}) &= \frac{[n]_{m_1, p, q} \alpha_1^{m_1} q^{\binom{m_1}{2}}}{(1 \oplus \alpha_1)_{p,q}^{m_1}}, \quad m_1 \in \{0, 1, \dots, n\}, \\ E([Y_2]_{m_2, p, q} | Y_1 = y_1) &= \frac{[n - y_1]_{m_2, p, q} \alpha_2^{m_2} q^{\binom{m_2}{2}}}{(1 \oplus \alpha_2)_{p,q}^{m_2}}, \quad m_2 \in \{0, 1, \dots, n - y_1\}, \\ E([Y_2]_{m_2, p, q}) &= \frac{[n]_{m_2, p, q} \alpha_2^{m_2} q^{\binom{m_2}{2}} p^{\binom{m_2}{2} + m_2(n - m_2)}}{(1 \oplus \alpha_2)_{p,q}^{m_2} (p^{n - m_2} \oplus \alpha_1 q^{n - m_2})_{p,q}^{m_2}}, \quad m_2 \in \{0, 1, \dots, n\}, \end{aligned}$$

and

$$E([Y_1]_{m_1, p, q} [Y_2]_{m_2, p, q}) = \frac{[n]_{m_1 + m_2, p, q} \alpha_1^{m_1} \alpha_2^{m_2} p^{\binom{m_2}{2} + m_2(n - m_2)} q^{\binom{m_1}{2} + \binom{m_2}{2}}}{(1 \oplus \alpha_1)_{p,q}^{m_1} (1 \oplus \alpha_2)_{p,q}^{m_2} (p^{n - m_2} \oplus \alpha_1 q^{n - m_2})_{p,q}^{m_2}},$$

where  $m_1 \in \{0, 1, \dots, n - m_2\}$  and  $m_2 \in \{0, 1, \dots, n\}$ . Moreover, the  $(p, q)$ -covariance is derived as follows:

$$Cov([Y_1]_{p,q}, [Y_2]_{\mathcal{R}(p,q)}) = \frac{p^{n-1} \alpha_1 \alpha_2 [n]_{p,q} ([n-1]_{p,q} - [n]_{p,q})}{(1 + \alpha_1)(1 + \alpha_2)(p^{n-1} + \alpha_1 q^{n-1})}.$$

(ii) The negative  $(p, q)$ -trinomial distribution of the first kind, with parameters  $n, \underline{\alpha} = (\alpha_1, \alpha_2), p$ , and  $q$  is presented by:

$$P(\underline{W} = \underline{w}) = \left[ \begin{matrix} n + w_1 + w_2 - 1 \\ w_1, w_2 \end{matrix} \right]_{p,q} \frac{\alpha_1^{w_1} \alpha_2^{w_2} p^{\binom{n-w_1}{2} + \binom{n-w_2}{2}} q^{\binom{w_1}{2} + \binom{w_2}{2}}}{(1 \oplus \alpha_1)_{p,q}^{n+w_1+w_2} (1 \oplus \alpha_2)_{p,q}^{n+w_2}},$$

where  $w_j \in \mathbb{N} \cup \{0\}, 0 < \alpha_j < 1$ , and  $j \in \{1, 2\}$ . Besides, for  $m_1 \in \mathbb{N} \cup \{0\}$  and  $m_2 \in \mathbb{N} \cup \{0\}$ ,

its  $(p^{-1}, q^{-1})$ -factorial moments are given as follows:

$$E([W_2]_{m_2, p^{-1}, q^{-1}}) = [n + m_2 - 1]_{m_2, p, q} \alpha_2^{m_2}, \quad m_2 \in \mathbb{N} \cup \{0\},$$

$$E([W_1]_{m_1, p^{-1}, q^{-1}} | W_2 = w_2) = [n + w_2 + m_1 - 1]_{m_1, p, q} \alpha_1^{m_1}, \quad m_1 \in \mathbb{N} \cup \{0\},$$

$$E\left(\frac{[W_1]_{m_1, p^{-1}, q^{-1}}}{(p^{n+W_2} \oplus \alpha_2 q^{n+W_2})_{p,q}^{m_1}}\right) = [n + m_1 - 1]_{m_1, p, q} \alpha_1^{m_1}, \quad m_1 \in \mathbb{N} \cup \{0\},$$

and

$$E\left(\frac{[W_1]_{m_1, p^{-1}, q^{-1}} [W_2]_{m_2, p^{-1}, q^{-1}}}{(p^{n+W_2} \oplus \alpha_2 q^{n+W_2})_{p, q}^{m_1}}\right) = [n + m_1 + m_2 - 1]_{m_1+m_2, p, q} \alpha_1^{m_1} \alpha_2^{m_2},$$

Furthermore, the covariance of  $\widehat{W} := (p^{n+W_2} \oplus \alpha_2 q^{n+W_2})^{-1} [W_1]_{m_1, p^{-1}, q^{-1}}$  and  $\overline{W} := [W_2]_{m_2, p^{-1}, q^{-1}}$  is determined by:

$$Cov(\widehat{W}, \overline{W}) = [n]_{p, q} \alpha_1 \alpha_2 ([n + 1]_{p, q} - [n]_{p, q}).$$

(iii) The  $(p, q)$ -trinomial distribution of the second kind, with parameters  $n, \underline{\beta} = (\beta_1, \beta_2)$ ,  $p$ , and  $q$  is given by:

$$P(X_1 = x_1, X_2 = x_2) = \begin{bmatrix} n \\ x_1, x_2 \end{bmatrix}_{p, q} \beta_1^{x_1} \beta_2^{x_2} (1 \ominus \beta_1)_{p, q}^{n-x_1} (1 \ominus \beta_2)_{p, q}^{n-x_1-x_2},$$

where  $x_j \in \{0, 1, \dots, n\}$ ,  $x_1 + x_2 \leq n$ ,  $s_j = \sum_{i=1}^j x_i$ ,  $0 < \beta_j < 1$ , and  $j \in \{1, 2\}$ .

Besides, its  $(p, q)$ -factorial moments are given by:

$$E([X_1]_{m_1, p, q}) = [n]_{m_1, p, q} \beta_1^{m_1}, \quad m_1 \in \{0, 1, \dots, n\},$$

$$E([X_2]_{m_2, p, q} | X_1 = x_1) = [n - x_1]_{m_2, p, q} \beta_2^{m_2}, \quad m_2 \in \{0, 1, \dots, n - x_1\},$$

$$E\left(\frac{[X_2]_{m_2, p, q}}{(p^{n-m_2-X_1} \ominus \beta_1 q^{n-m_2-X_1})_{p, q}^{m_2}}\right) = [n]_{m_2, p, q} \beta_2^{m_2}, \quad m_2 \in \{0, 1, \dots, n\},$$

and

$$E\left(\frac{[X_1]_{m_1, p, q} [X_2]_{m_2, p, q}}{(p^{n-m_2-X_1} \ominus \beta_1 q^{n-m_2-X_1})_{p, q}^{m_2}}\right) = [n]_{m_1+m_2, p, q} \beta_1^{m_1} \beta_2^{m_2},$$

where  $m_1 \in \{0, 1, \dots, n - m_2\}$  and  $m_2 \in \{0, 1, \dots, n\}$ . Moreover, the  $(p, q)$ -covariance of the functions  $(p^{n-1-X_1} - \beta_1 q^{n-1-X_1})^{-1} [X_2]_{p, q}$  and  $[X_1]_{p, q}$  is given by:

$$Cov\left([X_1]_{p, q}, (p^{n-1-X_1} - \beta_1 q^{n-1-X_1})^{-1} [X_2]_{p, q}\right) = [n]_{p, q} \beta_1 \beta_2 ([n - 1]_{p, q} - [n]_{p, q}).$$

(iii) The negative  $(p, q)$ -trinomial probability distribution of the second kind, with parameters  $n, \underline{\beta} = (\beta_1, \beta_2)$ ,  $p$ , and  $q$  is given by:

$$P(\underline{V} = \underline{v}) = \begin{bmatrix} n + v_1 + v_2 - 1 \\ v_1, v_2 \end{bmatrix}_{p, q} \beta_1^{v_1} \beta_2^{v_2} (1 \ominus \beta_1)_{p, q}^{n+v_2} (1 \ominus \beta_2)_{p, q}^n, \quad v_j \in \mathbb{N} \cup \{0\},$$

where  $0 < \beta_j < 1$ , and  $j \in \{1, 2\}$ . Moreover, its  $(p, q)$ -factorial moments are presented as follows:

$$E([V_2]_{m_2, p, q}) = \frac{[n + m_2 - 1]_{m_2, p, q} \beta_2^{m_2}}{(p^n \ominus \beta_2 q^n)_{p, q}^{m_2}},$$

$$E([V_1]_{m_1, p, q} | V_2 = v_2) = \frac{[n + v_2 + m_1 - 1]_{m_1, p, q} \beta_1^{m_1}}{(p^{n+v_2} \ominus \beta_1 q^{n+v_2})_{p, q}^{m_1}},$$

$$E\left(\frac{[V_1]_{m_1, p, q}}{(p^{n+V_2} \ominus \beta_1 q^{n+V_2})_{p, q}^{-m_1}}\right) = \frac{[n + m_1 - 1]_{m_1, p, q} \beta_1^{m_1}}{(p^n \ominus \beta_2 q^n)_{p, q}^{m_1}},$$

and

$$E\left(\frac{[V_1]_{m_1, p, q} [V_2]_{m_2, p, q}}{(p^{n+V_2} \ominus \beta_1 q^{n+V_2})_{p, q}^{-m_2}}\right) = \frac{[n + m_1 + m_2 - 1]_{m_1+m_2, p, q} \beta_1^{m_1} \beta_2^{m_2}}{(p^n \ominus \beta_2 q^n)_{p, q}^{m_1+m_2}},$$

where  $m_1 \in \mathbb{N} \cup \{0\}$  and  $m_2 \in \mathbb{N} \cup \{0\}$ . Furthermore, the  $(p, q)$ -covariance of the functions and  $\widehat{V} := (p^{n+V_2} \ominus \beta_2 q^{n+V_2}) [V_1]_{m_2, p, q}$  and  $\overline{V} := [V_2]_{m_1, p, q}$  is given by:

$$Cov(\widehat{V}, \overline{V}) = \frac{[n]_{p, q} \beta_1 \beta_2}{(p^n - \beta_2 q^n)} \left( \frac{[n+1]_{p, q}}{(p^{n+1} - \beta_2 q^{n+1})} - \frac{[n]_{p, q}}{(p^n - \beta_2 q^n)} \right).$$

### 3.3 Trinomial distribution and Chakrabarty and Jagannathan algebra [2]

Putting  $\mathcal{R}(x, y) = \frac{1-xy}{(p^{-1}-q)x}$ , we obtain:

(i) The  $(p^{-1}, q)$ -trinomial distribution of the first kind is given by

$$P(Y_1 = y_1, Y_2 = y_2) = \left[ \begin{matrix} n \\ y_1, y_2 \end{matrix} \right]_{p^{-1}, q} \frac{\alpha_1^{y_1} \alpha_2^{y_2} p^{-\binom{n-y_1}{2} - \binom{n-y_2}{2}} q^{\binom{y_1}{2} + \binom{y_2}{2}}}{(1 \oplus \alpha_1)_{p^{-1}, q}^n (1 \oplus \alpha_2)_{p^{-1}, q}^{n-y_1}}$$

and the  $(p^{-1}, q)$ -factorial moments are:

$$E([Y_1]_{m_1, p^{-1}, q}) = \frac{[n]_{m_1, p^{-1}, q} \alpha_1^{m_1} q^{\binom{m_1}{2}}}{(1 \oplus \alpha_1)_{p^{-1}, q}^{m_1}}, \quad m_1 \in \{0, 1, \dots, n\},$$

$$E([Y_2]_{m_2, p^{-1}, q} | Y_1 = y_1) = \frac{[n - y_1]_{m_2, p^{-1}, q} \alpha_2^{m_2} q^{\binom{m_2}{2}}}{(1 \oplus \alpha_2)_{p^{-1}, q}^{m_2}}, \quad m_2 \in \{0, 1, \dots, n - y_1\},$$

$$E([Y_2]_{m_2, p^{-1}, q}) = \frac{[n]_{m_2, p^{-1}, q} \alpha_2^{m_2} q^{\binom{m_2}{2}} p^{-\binom{m_2}{2} - m_2(n-m_2)}}{(1 \oplus \alpha_2)_{p^{-1}, q}^{m_2} (p^{n-m_2} \oplus \alpha_1 q^{n-m_2})_{p^{-1}, q}^{m_2}}, \quad m_2 \in \{0, 1, \dots, n\},$$

and

$$E([Y_1]_{m_1, p^{-1}, q} [Y_2]_{m_2, p^{-1}, q}) = \frac{[n]_{m_1+m_2, p^{-1}, q} \alpha_1^{m_1} \alpha_2^{m_2} p^{-\binom{m_2}{2} - m_2(n-m_2)} q^{\binom{m_1}{2} + \binom{m_2}{2}}}{(1 \oplus \alpha_1)_{p^{-1}, q}^{m_1} (1 \oplus \alpha_2)_{p^{-1}, q}^{m_2} (p^{n-m_2} \oplus \alpha_1 q^{n-m_2})_{p^{-1}, q}^{m_2}},$$

where  $m_1 \in \{0, 1, \dots, n - m_2\}$  and  $m_2 \in \{0, 1, \dots, n\}$ . Moreover, the  $(p^{-1}, q)$ -covariance is derived as follows:

$$Cov([Y_1]_{p^{-1}, q}, [Y_2]_{p^{-1}, q}) = \frac{p^{-n+1} \alpha_1 \alpha_2 [n]_{p^{-1}, q} ([n-1]_{p^{-1}, q} - [n]_{p^{-1}, q})}{(1 + \alpha_1)(1 + \alpha_2)(p^{-n+1} + \alpha_1 q^{n-1})}.$$

- (ii) The negative  $(p^{-1}, q)$ -trinomial distribution of the first kind, with parameters  $n$ ,  $\underline{\alpha} = (\alpha_1, \alpha_2)$ ,  $p$ , and,  $q$  is presented by:

$$P(\underline{W} = \underline{w}) = \binom{n + w_1 + w_2 - 1}{w_1, w_2}_{p^{-1}, q} \frac{\alpha_1^{w_1} \alpha_2^{w_2} p^{-\binom{n-w_1}{2} - \binom{n-w_2}{2}} q^{\binom{w_1}{2} + \binom{w_2}{2}}}{(1 \oplus \alpha_1)_{p^{-1}, q}^{n+w_1+w_2} (1 \oplus \alpha_2)_{p^{-1}, q}^{n+w_2}},$$

where  $w_j \in \mathbb{N} \cup \{0\}$ ,  $0 < \alpha_j < 1$ , and  $j \in \{1, 2\}$ . Besides, for  $m_1 \in \mathbb{N} \cup \{0\}$  and  $m_2 \in \mathbb{N} \cup \{0\}$ ,

its  $(p, q^{-1})$ -factorial moments are given as follows:

$$E([W_2]_{m_2, p, q^{-1}}) = [n + m_2 - 1]_{m_2, p^{-1}, q} \alpha_2^{m_2}, \quad m_2 \in \mathbb{N} \cup \{0\},$$

$$E([W_1]_{m_1, p, q^{-1}} | W_2 = w_2) = [n + w_2 + m_1 - 1]_{m_1, p^{-1}, q} \alpha_1^{m_1}, \quad m_1 \in \mathbb{N} \cup \{0\},$$

$$E\left(\frac{[W_1]_{m_1, p, q^{-1}}}{(p^{n+W_2} \oplus \alpha_2 q^{n+W_2})_{p^{-1}, q}^{m_1}}\right) = [n + m_1 - 1]_{m_1, p^{-1}, q} \alpha_1^{m_1}, \quad m_1 \in \mathbb{N} \cup \{0\},$$

and

$$E\left(\frac{[W_1]_{m_1, p, q^{-1}} [W_2]_{m_2, p, q^{-1}}}{(p^{n+W_2} \oplus \alpha_2 q^{n+W_2})_{p^{-1}, q}^{m_1}}\right) = [n + m_1 + m_2 - 1]_{m_1+m_2, p^{-1}, q} \alpha_1^{m_1} \alpha_2^{m_2},$$

Furthermore, the covariance of  $\widehat{W} := (p^{-n-W_2} - \alpha_2 q^{n+W_2})^{-1} [W_1]_{m_1, p, q^{-1}}$  and  $\overline{W} := [W_2]_{m_2, p, q^{-1}}$  is determined by:

$$Cov(\widehat{W}, \overline{W}) = [n]_{p^{-1}, q} \alpha_1 \alpha_2 ([n+1]_{p^{-1}, q} - [n]_{p^{-1}, q}).$$

- (iii) The  $(p^{-1}, q)$ -trinomial distribution of the second kind, with parameters  $n$ ,  $\underline{\beta} = (\beta_1, \beta_2)$ ,  $p$ , and,  $q$  is given by:

$$P(X_1 = x_1, X_2 = x_2) = \binom{n}{x_1, x_2}_{p^{-1}, q} \beta_1^{x_1} \beta_2^{x_2} (1 \ominus \beta_1)_{p^{-1}, q}^{n-x_1} (1 \ominus \beta_2)_{p^{-1}, q}^{n-x_1-x_2},$$

where  $x_j \in \{0, 1, \dots, n\}$ ,  $x_1 + x_2 \leq n$ ,  $s_j = \sum_{i=1}^j x_i$ ,  $0 < \beta_j < 1$ , and  $j \in \{1, 2\}$ .

Besides, its  $(p^{-1}, q)$ -factorial moments are given by:

$$E([X_1]_{m_1, p^{-1}, q}) = [n]_{m_1, p^{-1}, q} \beta_1^{m_1}, \quad m_1 \in \{0, 1, \dots, n\},$$

$$E([X_2]_{m_2, p^{-1}, q} | X_1 = x_1) = [n - x_1]_{m_2, p^{-1}, q} \beta_2^{m_2}, \quad m_2 \in \{0, 1, \dots, n - x_1\},$$

$$E\left(\frac{[X_2]_{m_2, p^{-1}, q}}{(p^{n-m_2-X_1} \ominus \beta_1 q^{n-m_2-X_1})_{p^{-1}, q}^{m_2}}\right) = [n]_{m_2, p^{-1}, q} \beta_2^{m_2}, \quad m_2 \in \{0, 1, \dots, n\},$$

and

$$E\left(\frac{[X_1]_{m_1, p^{-1}, q} [X_2]_{m_2, p^{-1}, q}}{(p^{n-m_2-X_1} \ominus \beta_1 q^{n-m_2-X_1})_{p^{-1}, q}^{m_2}}\right) = [n]_{m_1+m_2, p^{-1}, q} \beta_1^{m_1} \beta_2^{m_2},$$

where  $m_1 \in \{0, 1, \dots, n - m_2\}$  and  $m_2 \in \{0, 1, \dots, n\}$ . Moreover, the  $(p^{-1}, q)$ -covariance of the functions  $(p^{-n+1+X_1} - \beta_1 q^{n-1-X_1})^{-1} [X_2]_{p^{-1}, q}$  and  $[X_1]_{p^{-1}, q}$  is given by:

$$Cov\left[[X_1]_{p^{-1}, q}, (p^{-n+1+X_1} - \beta_1 q^{n-1-X_1})^{-1} [X_2]_{p^{-1}, q}\right] = [n]_{p^{-1}, q} \beta_1 \beta_2 ([n-1]_{p^{-1}, q} - [n]_{p^{-1}, q}).$$

(iii) The negative  $(p^{-1}, q)$ -trinomial probability distribution of the second kind, with parameters  $n, \underline{\beta} = (\beta_1, \beta_2), p$ , and,  $q$  is given by:

$$P(\underline{V} = \underline{v}) = \left[ \begin{matrix} n + v_1 + v_2 - 1 \\ v_1, v_2 \end{matrix} \right]_{p^{-1}, q} \beta_1^{v_1} \beta_2^{v_2} (1 \ominus \beta_1)_{p^{-1}, q}^{n+v_2} (1 \ominus \beta_2)_{p^{-1}, q}^n, \quad v_j \in \mathbb{N} \cup \{0\},$$

where  $0 < \beta_j < 1$ , and  $j \in \{1, 2\}$ . Moreover, its  $(p^{-1}, q)$ -factorial moments are presented as follows:

$$E([V_2]_{m_2, p^{-1}, q}) = \frac{[n + m_2 - 1]_{m_2, p^{-1}, q} \beta_2^{m_2}}{(p^n \ominus \beta_2 q^n)_{p^{-1}, q}^{m_2}},$$

$$E([V_1]_{m_1, p^{-1}, q} | V_2 = v_2) = \frac{[n + v_2 + m_1 - 1]_{m_1, p^{-1}, q} \beta_1^{m_1}}{(p^{n+v_2} \ominus \beta_1 q^{n+v_2})_{p^{-1}, q}^{m_1}},$$

$$E\left(\frac{[V_1]_{m_1, p^{-1}, q}}{(p^{n+V_2} \ominus \beta_1 q^{n+V_2})_{p^{-1}, q}^{-m_1}}\right) = \frac{[n + m_1 - 1]_{m_1, p^{-1}, q} \beta_1^{m_1}}{(p^n \ominus \beta_2 q^n)_{p^{-1}, q}^{m_1}},$$

and

$$E\left(\frac{[V_1]_{m_1, p^{-1}, q} [V_2]_{m_2, p^{-1}, q}}{(p^{n+V_2} \ominus \beta_1 q^{n+V_2})_{p^{-1}, q}^{-m_2}}\right) = \frac{[n + m_1 + m_2 - 1]_{m_1+m_2, p^{-1}, q} \beta_1^{m_1} \beta_2^{m_2}}{(p^n \ominus \beta_2 q^n)_{p^{-1}, q}^{m_1+m_2}},$$

where  $m_1 \in \mathbb{N} \cup \{0\}$  and  $m_2 \in \mathbb{N} \cup \{0\}$ . Furthermore, the  $(p^{-1}, q)$ -covariance of the functions and  $\widehat{V} := (p^{-n-V_2} - \beta_2 q^{n+V_2}) [V_1]_{m_2, p^{-1}, q}$  and  $\overline{V} := [V_2]_{m_1, p^{-1}, q}$  is given by:

$$Cov(\widehat{V}, \overline{V}) = \frac{[n]_{p^{-1}, q} \beta_1 \beta_2}{(p^{-n} - \beta_2 q^n)} \left( \frac{[n+1]_{p^{-1}, q}}{(p^{-n-1} - \beta_2 q^{n+1})} - \frac{[n]_{p^{-1}, q}}{(p^{-n} - \beta_2 q^n)} \right).$$

### 3.4 Trinomial distribution and Hounkonnou-Ngompe generalization of $q$ - Quesne algebra [11]

Setting  $\mathcal{R}(x, y) = \frac{xy-1}{(q-p^{-1})y}$ , we obtain the following results:

- (i) The Hounkonnou-Ngompe generalization of  $q$ - Quesne-trinomial distribution of the first kind is given by:

$$P(Y_1 = y_1, Y_2 = y_2) = \left[ \begin{matrix} n \\ y_1, y_2 \end{matrix} \right]_{p,q}^Q \frac{\alpha_1^{y_1} \alpha_2^{y_2} p^{\binom{n-y_1}{2} + \binom{n-y_2}{2}} q^{-\binom{y_1}{2} - \binom{y_2}{2}}}{(1 \oplus \alpha_1)_{p,q}^n (1 \oplus \alpha_2)_{p,q}^{n-y_1}}$$

and their factorial moments are:

$$\begin{aligned} E([Y_1]_{m_1, p, q}^Q) &= \frac{[n]_{m_1, p, q}^Q \alpha_1^{m_1} q^{-\binom{m_1}{2}}}{(1 \oplus \alpha_1)_{p,q}^{m_1}}, \quad m_1 \in \{0, 1, \dots, n\}, \\ E([Y_2]_{m_2, p, q}^Q | Y_1 = y_1) &= \frac{[n - y_1]_{m_2, p, q}^Q \alpha_2^{m_2} q^{-\binom{m_2}{2}}}{(1 \oplus \alpha_2)_{p,q}^{m_2}}, \quad m_2 \in \{0, 1, \dots, n - y_1\}, \\ E([Y_2]_{m_2, p, q}^Q) &= \frac{[n]_{m_2, p, q}^Q \alpha_2^{m_2} p^{\binom{m_2}{2} + m_2(n-m_2)} q^{-\binom{m_2}{2}}}{(1 \oplus \alpha_2)_{p,q}^{m_2} (p^{n-m_2} \oplus \alpha_1 q^{n-m_2})_{p,q}^{m_2}}, \quad m_2 \in \{0, 1, \dots, n\}, \end{aligned}$$

and

$$E([Y_1]_{m_1, p, q}^Q [Y_2]_{m_2, p, q}^Q) = \frac{[n]_{m_1+m_2, p, q}^Q \alpha_1^{m_1} \alpha_2^{m_2} p^{\binom{m_2}{2} + m_2(n-m_2)} q^{-\binom{m_1}{2} - \binom{m_2}{2}}}{(1 \oplus \alpha_1)_{p,q}^{m_1} (1 \oplus \alpha_2)_{p,q}^{m_2} (p^{n-m_2} \oplus \alpha_1 q^{n-m_2})_{p,q}^{m_2}},$$

where  $m_1 \in \{0, 1, \dots, n - m_2\}$  and  $m_2 \in \{0, 1, \dots, n\}$ . Moreover, the covariance is derived as follows:

$$Cov([Y_1]_{p,q}^Q, [Y_2]_{\mathcal{R}(p,q)}^Q) = \frac{p^{n-1} [n]_{p,q}^Q \alpha_1 \alpha_2 ([n-1]_{p,q}^Q - [n]_{p,q}^Q)}{(1 + \alpha_1)(1 + \alpha_2)(p^{n-1} + \alpha_1 q^{1-n})}.$$

- (ii) The negative Hounkonnou-Ngompe generalization of  $q$ - Quesne-trinomial distribution of the first kind, with parameters  $n$ ,  $\underline{\alpha} = (\alpha_1, \alpha_2)$ ,  $p$ , and,  $q$  is presented by:

$$P(\underline{W} = \underline{w}) = \left[ \begin{matrix} n + w_1 + w_2 - 1 \\ w_1, w_2 \end{matrix} \right]_{p,q}^Q \frac{\alpha_1^{w_1} \alpha_2^{w_2} p^{\binom{n-w_1}{2} + \binom{n-w_2}{2}} q^{-\binom{w_1}{2} - \binom{w_2}{2}}}{(1 \oplus \alpha_1)_{p,q}^{n+w_1+w_2} (1 \oplus \alpha_2)_{p,q}^{n+w_2}},$$

where  $w_j \in \mathbb{N} \cup \{0\}$ ,  $0 < \alpha_j < 1$ , and  $j \in \{1, 2\}$ . Besides, its factorial moments are given as follows:

$$\begin{aligned} E([W_2]_{m_2, p^{-1}, q^{-1}}^Q) &= \alpha_2^{m_2} [n + m_2 - 1]_{m_2, p, q}^Q, \\ E([W_1]_{m_1, p^{-1}, q^{-1}}^Q | W_2 = w_2) &= \alpha_1^{m_1} [n + w_2 + m_1 - 1]_{m_1, p, q}^Q, \\ E\left(\frac{[W_1]_{m_1, p^{-1}, q^{-1}}^Q}{(p^{n+W_2} \oplus \alpha_2 q^{n+W_2})_{p,q}^{m_1}}\right) &= \alpha_1^{m_1} [n + m_1 - 1]_{m_1, p, q}^Q, \end{aligned}$$



and

$$E\left(\frac{[W_1]_{m_1, p^{-1}, q^{-1}}^Q [W_2]_{m_2, p^{-1}, q^{-1}}^Q}{(p^{n+W_2} \oplus \alpha_2 q^{n+W_2})_{p, q}^{m_1}}\right) = \alpha_1^{m_1} \alpha_2^{m_2} [n + m_1 + m_2 - 1]_{m_1 + m_2, p, q}^Q,$$

where  $m_1 \in \mathbb{N} \cup \{0\}$  and  $m_2 \in \mathbb{N} \cup \{0\}$ . Furthermore, the covariance of  $\widehat{W} := (p^{n+W_2} \oplus \alpha_2 q^{n+W_2})^{-1} [W_1]_{m_1, p^{-1}, q^{-1}}^Q$  and  $\overline{W} := [W_2]_{m_2, p^{-1}, q^{-1}}^Q$  is determined by:

$$Cov(\widehat{W}, \overline{W}) = [n]_{p, q}^Q \alpha_1 \alpha_2 ([n + 1]_{p, q}^Q - [n]_{p, q}^Q).$$

(iii) The Hounkonnou-Ngompe generalization of  $q$ - Quesne-trinomial distribution of the second kind, with parameters  $n, \underline{\beta} = (\beta_1, \beta_2), p$ , and,  $q$  is given by:

$$P(X_1 = x_1, X_2 = x_2) = \left[ \begin{matrix} n \\ x_1, x_2 \end{matrix} \right]_{p, q}^Q \beta_1^{x_1} \beta_2^{x_2} (1 \ominus \beta_1)_{p, q}^{n-x_1} (1 \ominus \beta_2)_{p, q}^{n-x_1-x_2},$$

where  $x_j \in \{0, 1, \dots, n\}, x_1 + x_2 \leq n, s_j = \sum_{i=1}^j x_i, 0 < \beta_j < 1$ , and  $j \in \{1, 2\}$ .

Besides, its factorial moments are given by:

$$E([X_1]_{m_1, p, q}^Q) = [n]_{m_1, p, q}^Q \beta_1^{m_1}, \quad m_1 \in \{0, 1, \dots, n\},$$

$$E([X_2]_{m_2, p, q}^Q | X_1 = x_1) = [n - x_1]_{m_2, p, q}^Q \beta_2^{m_2}, \quad m_2 \in \{0, 1, \dots, n - x_1\},$$

$$E\left(\frac{[X_2]_{m_2, p, q}^Q}{(p^{n-m_2-X_1} \ominus \beta_1 q^{n-m_2-X_1})_{p, q}^{m_2}}\right) = [n]_{m_2, p, q}^Q \beta_2^{m_2}, \quad m_2 \in \{0, 1, \dots, n\},$$

and

$$E\left(\frac{[X_1]_{m_1, p, q}^Q [X_2]_{m_2, p, q}^Q}{(p^{n-m_2-X_1} \ominus \beta_1 q^{n-m_2-X_1})_{p, q}^{m_2}}\right) = [n]_{m_1 + m_2, p, q}^Q \beta_1^{m_1} \beta_2^{m_2},$$

where  $m_1 \in \{0, 1, \dots, n - m_2\}$  and  $m_2 \in \{0, 1, \dots, n\}$ . Moreover, the covariance of the functions  $(p^{n-1-X_1} - \beta_1 q^{1+X_1-n})^{-1} [X_2]_{p, q}^Q$  and  $[X_1]_{p, q}^Q$  is given by:

$$Cov\left[[X_1]_{p, q}^Q, (p^{n-1-X_1} - \beta_1 q^{1+X_1-n})^{-1} [X_2]_{p, q}^Q\right] = [n]_{p, q}^Q \beta_1 \beta_2 ([n - 1]_{p, q}^Q - [n]_{p, q}^Q).$$

(iii) The negative Hounkonnou-Ngompe generalization of  $q$ - Quesne- trinomial probability distribution of the second kind, with parameters  $n, \underline{\beta} = (\beta_1, \beta_2), p$ , and,  $q$  is given by:

$$P(\underline{Y} = \underline{v}) = \left[ \begin{matrix} n + v_1 + v_2 - 1 \\ v_1, v_2 \end{matrix} \right]_{p, q}^Q \beta_1^{v_1} \beta_2^{v_2} (1 \ominus \beta_1)_{p, q}^{n+v_2} (1 \ominus \beta_2)_{p, q}^n, \quad v_j \in \mathbb{N} \cup \{0\},$$

where  $0 < \beta_j < 1$ , and  $j \in \{1, 2\}$ . Moreover, for  $m_1 \in \mathbb{N} \cup \{0\}$  and  $m_2 \in \mathbb{N} \cup \{0\}$ , its factorial moments of are presented as follows:

$$E([V_2]_{m_2,p,q}^Q) = \frac{[n + m_2 - 1]_{m_2,p,q}^Q \beta_2^{m_2}}{(p^n \ominus \beta_2 q^n)_{p,q}^{m_2}},$$

$$E([V_1]_{m_1,p,q}^Q | V_2 = v_2) = \frac{[n + v_2 + m_1 - 1]_{m_1,p,q}^Q \beta_1^{m_1}}{(p^{n+v_2} \ominus \beta_1 q^{n+v_2})_{p,q}^{m_1}},$$

$$E\left(\frac{[V_1]_{m_1,p,q}^Q}{(p^{n+V_2} \ominus \beta_1 q^{n+V_2})_{p,q}^{-m_1}}\right) = \frac{[n + m_1 - 1]_{m_1,p,q}^Q \beta_1^{m_1}}{(p^n \ominus \beta_2 q^n)_{p,q}^{m_1}},$$

and

$$E\left(\frac{[V_1]_{m_1,p,q}^Q [V_2]_{m_2,p,q}^Q}{(p^{n+V_2} \ominus \beta_1 q^{n+V_2})_{p,q}^{-m_2}}\right) = \frac{[n + m_1 + m_2 - 1]_{m_1+m_2,p,q}^Q \beta_1^{m_1} \beta_2^{m_2}}{(p^n \ominus \beta_2 q^n)_{p,q}^{m_1+m_2}}.$$

Furthermore, the covariance of the functions  $\widehat{V} := (p^{n+V_2} \ominus \beta_2 q^{n+V_2}) [V_1]_{m_2,p,q}^Q$  and  $\overline{V} := [V_2]_{m_1,p,q}^Q$  is given by:

$$Cov(\widehat{V}, \overline{V}) = \frac{[n]_{p,q}^Q \beta_1 \beta_2}{(p^n - \beta_2 q^n)} \left( \frac{[n+1]_{p,q}^Q}{(p^{n+1} - \beta_2 q^{n+1})} - \frac{[n]_{p,q}^Q}{(p^n - \beta_2 q^n)} \right).$$

## Acknowledgments

This research was partly supported by the SNF Grant No. IZSEZO\_206010.

## References

- [1] L. C. Biedenharn, The quantum group  $SU_q(2)$  and a  $q$ -analogue of the boson operators, *J. Phys. A* **22** (1989), L873-L878.
- [2] R. Chakrabarti and R. Jagannathan, A  $(p, q)$ -oscillator realisation of two-parameter quantum algebras, *J. Phys. A: Math. Gen.* **24**, L711-L718, IMSC-91-15 (1991).
- [3] Ch. A. Charalambides, *Discrete  $q$ -distributions*. John Wiley and Sons, Inc., Hoboken, New Jersey, 2016.
- [4] Charalambides, Ch. A.  $q$ -Multinomial and negative  $q$ -multinomial distributions, *Comm. Statist. Theory Meth.* (2020).
- [5] Charalambides, Ch. A. Multivariate  $q$ -Pólya and inverse  $q$ -Pólya distributions, *Comm. Statist. Theory Meth.* (2020).

- [6] Charalambides, Ch. A.  $q$ - Factorial moments of bivariate discrete distributions, *Comm. Statist. Theory Meth.* (2022).
- [7] M. N. Hounkonnou and F. Melong,  $\mathcal{R}(p, q)$  Analogs of Discrete Distributions: General Formalism and Applications, *Journal of Stochastic Analysis*: Vol. 1 : No. 4 , Article 11. DOI: 10.31390/josa.1.4.11, (2020).
- [8] M. N. Hounkonnou and F. Melong,  $\mathcal{R}(p, q)$ -deformed combinatorics: full characterization and illustration, arXiv:1906.03059.
- [9] M. N. Hounkonnou and J. D. Kyemba Bukweli,  $\mathcal{R}(p, q)$ -calculus: differentiation and integration, *SUT. J. Math.* **49**, 145-167 (2013).
- [10] M. N. Hounkonnou and J. D. Kyemba Bukweli,  $(R, p, q)$ -deformed quantum algebras: Coherent states and special functions, *J. Math. Phys.* **51** , 063518-063518.20 (2010).
- [11] M. N. Hounkonnou and E. B. Ngompe Nkouankam, New  $(p, q, \mu, \nu, f)$ -deformed states, *J. Phys. A: Math. Theor.* **40**, 12113-12130 (2007).
- [12] R. Jagannathan and K. Srinivasa Rao, Two-parameter quantum algebras, twin-basic numbers, and associated generalized hypergeometric series, *Proceedings of the International Conference on Number Theory and Mathematical Physics*, 20-21 December 2005.
- [13] A. J. Macfarlane, On  $q$ - analogues of the quantum harmonic oscillator and quantum group  $SU_q(2)$ , *J. Phys. A* **22** (1989), 4581-4588.
- [14] T. Nishino, *Function theory in several complex variables*, Translations of mathematical monographs, Volume 193, American Mathematical Society, Providence, Rhode Island, 2001.