A two-component Bose-Einstein condensate can 'bypass' the nocloning theorem

Shouvik Datta*

 $Department\ of\ Physics,\ Indian\ Institute\ of\ Science\ Education\ and\ Research,$

Pune 411008, Maharashtra, India

*Email: shouvik@iiserpune.ac.in

Keywords: Exciton, Bose-Einstein Condensation, Quantum Optics, Quantum Information

Abstract

No-cloning theorem in quantum cryptography prevents an eavesdropper from perfectly duplicating any arbitrary quantum state. Here we argue that an experimental scheme to produce a quantum superposition of interacting Bose-Einstein condensates can generate N bosonic clones of any arbitrary single quantum state at large N limit of thermodynamic equilibrium with high fidelity and thus operationally 'bypass' the restrictions imposed by the above mentioned theorem. This is possible because quantum statistical nature of this 'cloning operation' does not require strict unitary evolution of standard quantum mechanics within a single Hilbert space.

1. INTRODUCTION

The original no-cloning theorem [1,2] states that linearity and unitarity of quantum mechanics prevents perfect copying of an arbitrary quantum superposition state. As such various replicating schemes were proposed and reviewed [3] for cloning 'albeit imperfectly' [4,5] up to an optimal level. However, most reports of such 'optimal' cloning of photons in quantum cryptography relied on the process of duplication within the constraints of a single, 1st quantized Hilbert space of a quantum system. This is, to the best of our understanding, need not be followed exactly from an 'operational' perspective to execute quantum cloning involving a series of intermediate quantum statistical mechanical processes. We will also argue that the fidelity and coherence of the final quantum state with respect to the initial quantum state can, however, be preserved by ensuring 'identical' amplifications of both quantum channels (i.e. two orthogonal polarizations) and subsequently generating quantum superposition of interacting Bose-Einstein condensates (BEC) having identical bosonic clones in the limit of large numbers.

To elaborate further, here we take the example of forming quantum superposition state of a two interacting BECs [6] in the mean field approximation and extend that to excitons or bound electron-hole pairs to execute such quantum cloning. In this case, we will not be violating any linearity, unitarity of 1st quantized formalism of norm preserving Hilbert space while executing the quantum cloning routes using the creation of a quantum statistical object like two interacting excitonic BECs. It is mainly because the 'quantum copier' which processes this cloning mechanism by generating two-component interacting BECs will be 'operationally' mapped from the 1st quantized state of the incoming photon to a 2nd quantized Fock space or many-body Hilbert space of excitonic BEC through light matter interactions. There the creations and destructions of particles (e.g. excitons) are allowed through a sequence of transitions, first mediated by separate

amplifications of individual polarization states and subsequently by generation of two sets of excitons using an excitonic material/structure. These two groups of excitons can then be driven to the quantum ground state of a two-component, interacting BECs via 'weak' Josephson like interactions [6] as prescribed in the past for atomic BECs. We will argue that once such two-component, interacting BECs are formed under the mean field approximation [6], then it can be used to germinate multiple clones of the incoming photon to 'bypass' the no-cloning theorem and thereby successfully clone any arbitrary quantum superposition state. Therefore, the whole 'operational' processes of quantum cloning can happen by stepping outside the stringent theoretical framework of 'unitary' evolutions of quantum mechanics within a single Hilbert space in the first place, which is the all-important basis for the well-known no-cloning theorem [1,2].

2. EXPERIMENTAL PROPOSAL

The schematic of this quantum mechanical cloning machine (QCM) is described in Fig. 1. We assume that the incoming photon is linearly polarized such that the resultant polarization vector is in an unknown quantum superposition state

$$|\varphi\rangle = \alpha |\uparrow\rangle + \beta |\rightarrow\rangle \tag{1}$$

where α and β are 'unknown' complex numbers such that $(\alpha)^2 + (\beta)^2 = 1$. The states $|\uparrow\rangle$ and $|\rightarrow\rangle$ are vertical and horizontal polarizations respectively and form an orthonormal, complete basis in Hilbert space of those two states. The net polarization of state $|\phi\rangle$ is also at an unknown angle θ with respect to the vertical axis. In actual experimental realization, one may also use a superposition of left and right circularly polarized light instead of $|\uparrow\rangle$ and $|\rightarrow\rangle$. Please note that




Fig. 1. Schematic of the quantum cloning process using a two-component, interacting Bose-Einstein condensates of excitons. We purposely desist from knowing/tracking any time resolved individual outcome(s) of the beam splitting, reflections from the mirrors as well as that of the amplification processes. We treat the whole system as a black box till the quantum ground state of the two-component, interacting BECs emit those cloned photons in a phase coherent way at the large N limit of thermodynamic equilibrium. This is necessary so that we do not to disturb the phase coherence between the vertical | ↑ ⟩ and horizontal | → ⟩ polarizations by the process of measurements. The only requirement is large and identical amplification of both polarization channels using periodically pooled non-linear crystals. Green squares are 50:50 polarizing beam splitters. Polarizing beam splitters # 2,3 are used to filter out any unwanted polarization states due to the presence of spontaneous emissions during amplification processes. These last two 'classical' beam splitters can ignore negligible quantum fluctuations entering through their unused ports while filtering these highly amplified beams in respective polarization channels. Thicker arrows indicate the amplified beams. Details of proposed amplification of single photons and associated issues are discussed in the text.

the choice of $|\uparrow\rangle$ and $|\rightarrow\rangle$ as basis states here is merely to bring notational similarity with original no-cloning literatures in quantum cryptography.

Operationally speaking, one can design the QCM in the following manner as shown in Fig. 1. First - (a) a polarizing beam splitter can separate the respective vertical $|\uparrow\rangle$ and horizontal $|\rightarrow\rangle$ polarizations of the incoming photon in two different channels, likely with some additional quantum noise entering through the unused ports of the beam splitter numbered '1'. Although, it was argued in the no cloning literature that there exists no QCM which can perfectly amplify an arbitrary superposition of polarization states. However, it is always recognized [1,7] that those same arguments do not actually rule out the possibilities of having some devices which can separately amplify two different orthogonally polarized states $|\uparrow\rangle$ and $|\rightarrow\rangle$. Therefore, one can first split these orthogonal polarizations and - (b) thereafter amplify the weak signals of $|\uparrow\rangle$ and $|\rightarrow\rangle$ separately. For example, one may use polarization sensitive amplification processes using degenerate parametric amplifier with a strong pump beam such that $\omega_{signal} = \omega_{idler} = \frac{\omega_{Pump}}{2} =$ ω . Technical details of this amplification for single photons will be discussed below. Finally, one can - (c) allow these two amplified beams to be incident on a cloning device 'D' (blue rectangle in Fig. 1) without any additional phase lag introduced between the two polarization channels. The final stage of this QCM is the device 'D' which can be made of any light sensitive excitonic material/structure having high quantum yields of optical absorption in the first place. Specifically, if we are using 'D' which can generate excitons using linear optical absorption of light and if all of those excitons generated by amplified light having orthogonal polarizations $|\uparrow\rangle$ and $|\rightarrow\rangle$ can be driven into two separate but interacting BECs, then one will be able [6] to create a quantum superposition of this two-component excitonic BEC along with some additional light-matter interactions.

It is to be strictly noted here that one cannot detect and/or track these photon(s) during the beam splitting, amplification and final mixing process in the excitonic device 'D'. As a result, we will not be destroying the quantum information embedded within vertical $|\uparrow\rangle$ and horizontal $|\rightarrow\rangle$ polarization states of the incoming photon by excluding these measurement induced direct disturbances to the quantum state. Such amplifications of (say) 1 to N photons will only be required to generate sufficiently large numbers of photons to drive the excitonic BEC in this cloning device 'D'. In fact, because both photons and excitons are bosons themselves. The polarization state of input signal at the two amplification stages (red rectangles in Fig. 1) are known. The processes of identical amplifications of vertical $|\uparrow\rangle$ or horizontal $|\rightarrow\rangle$ polarization states in both channels to large photon numbers will be discussed in the next paragraph. Subsequent mixing of these amplified beams having $|\uparrow\rangle$ or $|\rightarrow\rangle$ polarization states by creating exciton BECs will be described in the next section.

As such the number-phase uncertainty principle certainly complicates the precise evaluation of quantum optical phase of a single photon before the amplification. These challenges [8-10] were well documented and subsequently a lot of in-depth studies were reported on this particular topic [11-15] in the last couple of decades. However, in our case we would like to operate outside these domains of strong quantum optical fluctuations of 'relative' phase of both polarization channels, which will further be used to create two separate but interacting excitonic BECs. As long as the physical amplification process of vertical $|\uparrow\rangle$ and horizontal $|\rightarrow\rangle$ polarization states are large and 'identical', the density of photoexcitations generated in D, presumably with a 'known' electronic structure, can remain proportional to incident amplified intensities from both polarization channels. Such 'identical' amplification of both channels can be ensured by having $\alpha' = \eta \alpha$ and $\beta' = \eta \beta$ respectively, where $|\eta|^2$ is the common multiplication

factor during the 'intensity' amplification process. Any quantum phase fluctuations at the limit of large N number of amplified photons or equivalently with large amplifications as $|\eta|^2 >> 1$ can then be ignored [16, 17] for each polarization channels and subsequently the relative phase of vertical $|\uparrow\rangle$ and horizontal $|\rightarrow\rangle$ polarization channels can remain well defined with such identical η . In fact, we will see that ignoring [6,18] such quantum phase fluctuations of each polarization channels can be useful in developing a mean-field description of the excitonic matter wave following Ref. 6 which will be important for this particular cloning process as well. For example, one may try of noiseless linear amplification (NLA) [11] using quantum scissors along with single photons of either $|\uparrow\rangle$ or $|\rightarrow\rangle$ to drive these NLA stages for respective vertical $|\uparrow\rangle$ and horizontal $|\rightarrow\rangle$ polarization channels (see Fig. 1a of Ref. 11). However, the gains of NLAs will never be large enough to generate many photons and subsequently many excitons to produce a BEC in device 'D'.

A single photon is an ideal amplitude squeezed number state. One can even try to use squeezed states [10,19] as the output of such amplification stages. However, in such phase sensitive parametric amplification for any single photon Fock state, one does not have a well-defined phase in quadrature space. Nevertheless, phase lock can be maintained between the signal and pump beam using the well-known quasi phase matching (QPM) [20] technique of non-linear optics. Here the momentum conservation necessary for phase-matching is adjusted using a periodically pooled structures. In our case, the relevant phase (Φ) between the pump beam and weak input beam can further be controlled during the amplification process by periodical phase modulation of the weak input signal with a mirror on a piezoelectric transducer stage (i.e. Fig 1 of Ref. 10, 21). Alternatively, one can use electro-optic modulator based phase modulation [22] of single-photon wave packets along with the above mentioned use of QPM to execute this. Such

external, periodic, phase modulation [10, 21, 22] of the weak input signal can be the key for relative phase locking between the strong pump beam and weak signal beam along quasi phase matching [20]. It will also be helpful in nullifying the effect of any unwelcome coupling to random vacuum modes entering through the first beam splitter marked as '1' in Fig. 1. Moreover, such phase sensitive, degenerate parametric amplification can be pre-calibrated to have large, identical gains $|\eta|^2$ for both $|\uparrow\rangle$ and $|\rightarrow\rangle$ polarization channels. Unlike the phase insensitive amplification, in phase sensitive amplification procedure, the gain is always independent of the number of input photons even for few or single photons. Prior experimental calibrations can, therefore, be used to identically amplify one particular quadrature component of both polarization states $|\uparrow\rangle$ and $|\rightarrow\rangle$ at the cost of extra noise in the other quadrature. Choice of respective polarization configurations (type I, type II etc.) of pump, signal and idler with respect to the non-linear crystal and the angle between the optic axis and propagation direction can be chosen during the actual experiment. Spontaneous parametric emission is a problem in case of phase insensitive amplifier for small photon numbers 'n'. However, in a phase sensitive amplification, the noisy quadrature components also decay exponentially [10] with enhanced propagation distance. This can be further enhanced using an optical cavity as well. For significantly high gains a phase sensitive amplifier can also compensate for inevitable losses to approach a noise figure close to one [10] for the amplifying quadrature. In this way, these phase sensitive, degenerate parametric amplifiers can generate 'noiseless' [10] amplification separately and independently for both vertical |↑⟩ and horizontal $|\rightarrow\rangle$ polarization states in respective channels of Fig. 1. Additionally, any noise from spontaneous emission in such a parametric down-conversion process can further be filtered [23] out using two additional 50:50 polarizing beam splitters (#2 & 3 in Fig. 1) before these are incident on excitonic material/structure D. Moreover, with strongly amplified beams in both the polarization channels,

one can finally ignore the effects of vacuum field fluctuations entering through the open ports of these beam splitters before generating spin selective excitons in D. As a result, down the line, these 2^{nd} and 3^{rd} beam splitters can further prevent any significant quantum fluctuations of relative phase of both polarization channels $|\uparrow\rangle$ and $|\rightarrow\rangle$. In the next section, we will discuss how these amplified $|\uparrow\rangle$ and $|\rightarrow\rangle$ states can be used to generate two separate sets of excitons (electron hole pairs) which can subsequently be tuned in a phase coherent way to form the two-component BEC of excitons as prescribed in the past [6].

Moreover, we can assume that each photon incident on 'D' can generate one exciton in the device 'D' through linear optical absorption processes. We also assume that all or same fraction of these two different ensembles of excitons photo generated individually with vertical $|\uparrow\rangle$ and horizontal $|\rightarrow\rangle$ polarizations can be driven into separate BECs having the quantum ground states as $|\psi_{\uparrow}\rangle$ and $|\psi_{\rightarrow}\rangle$ respectively. This is necessary to ensure that density of excitons in quantum ground states of $|\psi_{\uparrow}\rangle$ and $|\psi_{\rightarrow}\rangle$ can also remain proportional to $|\alpha|^2$ and $|\beta|^2$ respectively by using identical amplifications to preserve the relative phase in both polarization channels as mentioned in the previous section. However, by saying this, we are also assuming that $|\psi_{\uparrow}\rangle$ and $|\psi_{\rightarrow}\rangle$ are two orthogonal states and not any coherent states! Then another light source can be tuned in to generate weak 'Josephson' like coupling between $|\psi_{\uparrow}\rangle$ and $|\psi_{\rightarrow}\rangle$ to form a two-component BEC under mean field approximation [6]. All these above assumptions will prevent any phase randomization during the formative stages of these two BECs. The material/structure of the cloning device 'D' must also be sensitive for such spin/polarization selective optical transitions as well as allow for coherent interactions between these two excitonic BEC states of $|\psi_{\uparrow}\rangle$ and $|\psi_{\rightarrow}\rangle$.

3. EXPECTED OUTCOMES

Consequently, the resultant one particle ground state of this two-component interacting BEC will look like [6],

$$|\psi_1\rangle = \alpha'|\psi_1\rangle + \beta'|\psi_2\rangle \tag{2}$$

where coefficients are $\alpha' = \eta \alpha$ and $\beta' = \eta \beta$ respectively, the amplification factor η can be complex number and $(\alpha')^2 + (\beta')^2 = |\eta|^2$ as the single particle normalization factor. We are only assuming that $|\eta|^2$ can be kept identical for both polarization channels. Such calibrations are operationally possible with prior information of the nature of polarization states of the incoming photon as $|\uparrow\rangle$ and $|\rightarrow\rangle$ as the basis states of the input photon. Correspondingly, if 'N' such excitons can be produced in the cloning device 'D' with N amplified photons having either vertical $|\uparrow\rangle$ and horizontal $|\rightarrow\rangle$ polarization and then tuned with additional Josephson like 'weak' interactions to form a two-component BEC ground state of N excitons under mean-field approximation as prescribed earlier [6]. Finally, one gets the quantum ground state of two-component excitonic BEC under weak interaction as shown below in Eq. 3. This similar to that described by Eq. 7 of Ref. 6 in the context of cold atoms.

$$\psi_{2-component-BEC}\rangle = |\psi_1\rangle \otimes |\psi_1\rangle \otimes ... |\psi_1\rangle = \frac{1}{\sqrt{N!}} [\alpha'\alpha^{\dagger} + \beta'b^{\dagger}]^N |0\rangle$$
 (3)

where $|0\rangle$ is the vacuum state of the two-component quantum superposition of excitonic BEC states and α^{\dagger} , b^{\dagger} are creation operators of excitons in $|\psi_{\uparrow}\rangle \otimes |\psi_{\rightarrow}\rangle$ BEC states respectively and $(\alpha_N)^2 + (\beta_N)^2 = N|\eta|^2$ where $\alpha_N = \sqrt{N}\alpha'$, $\beta_N = \sqrt{N}\beta'$ for respective polarizations. As mentioned above, here we assume that each incoming photon having either vertical $|\uparrow\rangle$ and horizontal $|\rightarrow\rangle$ polarizations produce only and only one exciton in the device D. Most importantly,

the fidelity of such mapping can also be maintained by ensuring that $\alpha' = \eta \alpha$ and $\beta' = \eta \beta$ respectively and it will be discussed later in detail. As such, it was known that the large N thermodynamic limit, in general, will always produce [24,25] this particular quantum state $|\psi_{2-component-BEC}\rangle$. Detailed theoretical calculations to yield this final state of $|\psi_{2-component-BEC}\rangle$ were already reported [18,24,25] in a different context for condensation of Helium-3 as well. Therefore, here we refrain from writing down the same equations already elaborated in these past reports [6,18,24,25] as well as for sake of keeping a much wider generality in our quantum cloning procedure using such a two-component BEC at this stage. As described in the past, this final quantum state $|\psi_{2-component-BEC}\rangle$ can be the harmonic oscillator like ground state of this two-component BEC of N excitons as direct product of identical $|\psi_1
angle=$ $\alpha'|\psi_{\uparrow}\rangle + \beta'|\psi_{\rightarrow}\rangle$ as single particle excitonic excitations which are tuned via Josephson like coupling. The relative phase fluctuations of $|\psi_{\uparrow}\rangle$, $|\psi_{\rightarrow}\rangle$ in the $|\psi_{2-component-BEC}\rangle$ will decrease as $N^{-\frac{1}{2}}$ [18,24,25] and will be practically negligible at the large N limit of thermodynamic equilibrium. As a result, it will be possible [6,18,24,25] to generate this quantum state $|\psi_{2-component-BEC}\rangle = \frac{1}{\sqrt{N!}} [\alpha' \alpha^{\dagger} + \beta' b^{\dagger}]^{N} |0\rangle$ as multiple cloned state of the incoming photon with $|\varphi\rangle$.

There can be two degenerate energy levels [6] of this interacting BECs in case of strong coupling under mean field approximations. It was shown [6] that for beyond mean field approximations, there can be further splitting of those degenerate energy levels [Eq. 22 & 39 in Ref. 6] leading to a two-level quantum system. It was also described there how the ground-state wave function can take the form of a Schrodinger-cat state under such circumstances. In a way, going beyond the mean field approximation, one can even create [6] a macroscopic "Schrodinger-catinger-cating approximation".

cat" state formed by two interacting Bose condensates of excitons made separately from amplified vertical $|\uparrow\rangle$ and horizontal $|\rightarrow\rangle$ polarizations of single photons. However, exploring such experimental situations are currently beyond the scope of this study.

At this stage, we must also note that the linearity of optical absorption process in the excitonic cloning device 'D' (blue rectangle in Fig. 1) is, however, important. This will be required to prevent the generation of any higher order, multi particle superposition such as $[|\psi_{\uparrow\uparrow}\rangle + |\psi_{\rightrightarrows}\rangle]$ etc., which can also form during the mixing of amplified vertical $|\uparrow\rangle$ and horizontal $|\rightarrow\rangle$ polarizations while generating two distinct sets of exciton BECs with quantum states $|\psi_{\uparrow}\rangle$, $|\psi_{\rightarrow}\rangle$ respectively. This is because superposition states like $[|\psi_{\uparrow}\rangle + |\psi_{\rightrightarrows}\rangle]$ and $[|\psi_{\uparrow\uparrow}\rangle + |\psi_{\rightrightarrows}\rangle]$ etc., are qualitatively different from states $\frac{1}{\sqrt{N!}}[\alpha'\alpha^{\dagger} + \beta'b^{\dagger}]^{N}|0\rangle$ which is required for the universal cloning operations using such two-component BEC [6,18,24,25].

Finally, we expect that M of these excitons in identical quantum state of $|\psi_1\rangle = \alpha'|\psi_1\rangle + \beta'|\psi_{\rightarrow}\rangle$ within that two-component BEC ground state having long range spatial coherence to radiatively decay in to photons having vertical $|\uparrow\rangle$ and horizontal $|\rightarrow\rangle$ polarizations respectively with quantum efficiency (ξ). This is because, photons emitted through the recombination of all these excitons in the 'identical' quantum ground state of this two-component, interacting BEC will have a narrow spectral line widths and will be spontaneously phase coherent by themselves as consequence of BEC. This can also happen via polarization conserving stimulated emissions. One can also use an additional optical cavity to trigger these stimulated emissions as well. Thereafter, we can finally recover the initial quantum superposition state $|\phi\rangle = \alpha |\uparrow\rangle + \beta |\rightarrow\rangle$ of the incoming photon using the generic spatio-temporally coherent emissions from the quantum ground state of the two-component, interacting BECs of excitons as

$$|\psi_{Cloned}\rangle = |\varphi\rangle \otimes |\varphi\rangle \otimes \dots |\varphi\rangle = \frac{1}{\sqrt{M!}} [\alpha''c^{\dagger} + \beta''d^{\dagger}]^{M} |0\rangle$$
 (4)

where c^{\dagger} , d^{\dagger} are creation operators of photons in $|\uparrow\rangle$ & $|\rightarrow\rangle$ states respectively and generated from the two-component BEC as defined in equation (3) and $\alpha_{M'} = \sqrt{M} \alpha'' = \sqrt{M} \xi \alpha' = \sqrt{M} \xi \eta \alpha = C\alpha$ and $\beta_{M'} = \sqrt{M} \beta'' = \sqrt{M} \xi \beta' = \sqrt{M} \xi \eta \beta = C\beta$ such that $(\alpha_{M'})^2 + (\beta_{M'})^2 = |C|^2$ and $C = \sqrt{M} \xi \eta$ where C can be the experimentally determined constant factor. As a result, apart from a different normalization factor due to non-ideal quantum yields of the optical absorption and emission at the cloning device 'D', one can, in principle, duplicate any incoming pure state in arbitrary quantum superposition like $|\phi\rangle = \alpha |\uparrow\rangle + \beta |\rightarrow\rangle$ as

$$|\psi_{Cloned}\rangle \equiv |\varphi\rangle \otimes |\varphi\rangle \otimes ... |\varphi\rangle \cong \frac{(\xi\eta)^M}{\sqrt{M!}} [\alpha c^{\dagger} + \beta d^{\dagger}]^M |0\rangle$$
 (5)

Obviously, in practice, there will also be some Stokes shifts and the energy of the emitted photons can be red shifted from those of the absorbed one in the excitonic device 'D'. However, a prior knowledge of the photon energy and polarization basis of the incoming signal being used for communication and choosing a known/tailored electronic structure of D can be helpful to match the energies of amplified photons suitably with respect to the incoming one. This energy matching process is not going to affect the overall fidelity of the quantum cloning process of the incoming arbitrary superposition state. Here we are also assuming that the quantum ground state of this two-component exciton BEC in 'D' is not 'dark' or forbidden for light emissions due to spin selection rules. However, incident angles of amplified photon beams of two different polarizations can always be tuned [26] for a desired outcome.

4. DISCUSSIONS

Above mentioned quantum duplication scheme is an 'interdisciplinary' experimental proposal using quantum optics, light matter interactions and quantum statistical effects in condensed matter physics of BEC of excitons or electron-hole pairs. This is possible because these excitons as bosons within a BEC are quantum clones of each other in the first place. The overall processes of amplification followed by photo generation of excitons (bosons) driven towards a macroscopically large two-component, interacting BEC state and subsequently their radiative recombinations are certainly not any norm conserving unitary process within a single Hilbert space. Therefore, in this proposed QCM, we are not directly violating the no-cloning theorem but just 'bypassing' it to make quantum clones of any arbitrary superposition of quantum states. We are only using a series of quantum statistical light-matter interactions to generate the two-component, interacting excitonic BECs at the large N limit of thermodynamic equilibrium. This is possible once the individual polarization modes can be first separated, identically amplified and then eventually mixed to produce the quantum superposition state of two interacting BECs [6,18,24,25] of excitons which are tuned by Josephson like external light-matter interactions. We also argued how this scheme of mapping an arbitrary quantum superposition on to the quantum ground state of a twocomponent BEC can prevent unwanted multi-particle superposition(s) and produce quantum cloning of photons through generation and re-emission processes mediated by this quantum ground state of two-component BEC having large number of phase coherent excitons. This is because, formation of a superfluid BEC of excitons will resist usual dissipation and decoherence. Moreover, one can also extend this scheme beyond such photo generated excitonic ensembles to any other two-component BEC systems including exciton-polaritons, photons and even to atomic systems as well. Moreover, the relative errors of generating the $|\psi_{2-component-BEC}\rangle$ state as shown in Eq.

(3) will practically vanish at the large N thermodynamic limit. Moreover, the density matrix ρ_{input} for the input quantum state of Eq. 1 and the ρ_{output} for output quantum state of each single particle components of the above Eq. 4 are same $\left(\rho_{input} = \rho_{output} = \rho\right)$ along with the normalization prescribed in the text. As a result, the fidelity of such cloning in terms of $F\left(\rho_{input}, \rho_{output}\right) = \left[Tr\left(\sqrt{\sqrt{\rho_{input}}\rho_{output}\sqrt{\rho_{input}}}\right)\right]^2 = F(\rho,\rho) = 1$. Therefore, having such abundantly many bosonic clones of BEC can increase the cloning fidelity and reduce errors for a perfect cloning to an insignificant level by producing a large number of clones.

The security of quantum cryptography can still be recovered by 'operationally' moving from a simple 'qubit' like 2-level quantum system to a quantum system having large and unknown 'd' dimensional 'qudits' or even to an infinite dimensional continuum basis states to transfer information. This can make the process of identifying a suitable material system to execute a well-defined d-component BEC mediated QCM enormously complicated, if not impossible for large, unknown and possibly a randomly variable 'd' for successive usages. Such qudit based quantum processors [27,28] and communication [29] devices are certainly being developed in the recent past.

5. Conclusions

Finally, formation of such a two-component, interacting BEC is not a norm conserving, linear process restricted within the domain of a single Hilbert space, but belongs to the domain of quantum statistical mechanics of interactive, quantum system of light-matter interactions and excitons. Thus generating bosonic clones using the quantum ground state of an excitonic BEC is

a suitable medium for operational execution of 'perfect' cloning of any arbitrary quantum state without having any direct conflict with basic premises of the no-cloning theorem [1,2]. This whole experimental proposal is centered around (i) identical and (ii) large amplifications of both vertical $|\uparrow\rangle$ and horizontal $|\rightarrow\rangle$ polarization channels separately, then (iii) ignoring the phase fluctuations at large N thermodynamic limit, (iv) photo generation of a two-component excitonic BEC tuned with external but 'weak' [6] light-matter interactions, finally (v) the radiative recombination of phase coherent excitons from the BEC ground state leading to 'cloned' photons. We also note that instead of pulsed [10], degenerate parametric amplifier as proposed above, use of a continuous wave [30] pumped degenerate parametric amplifier may also be beneficial to work along with our experimental proposal. The large N thermodynamic limit helps us in two ways – (a) firstly, it helps us to ignore phase fluctuations during the amplification of single photon, (b) secondly, it also facilitates in generating [6,18,24,25] the superposition state [Eq. 3-5] of two interacting BECs in the mean field approximation as the 'suitable' mediator for quantum cloned state. Having a phase coherent BEC also helps in preventing the quantum phase information from dephasing [6,18,24,25] within an experimentally realizable time scale.

Furthermore, perfect cloning is also a sufficient condition for allowing communication of information between space-like separated points using entanglement. Therefore, by that same token, such possibilities of 'perfect' quantum cloning of any maximally entangled state of photon, which are 'not' limited [31,32] by the constraints of the linearity, unitarity and completeness of the 1st quantized wave functions within a single Hilbert space, can, in principle, no longer prohibit superluminal [33,34], EPR [35] like signal communications using the above mentioned scheme involving 2nd quantized state of field operators intermediated via the formation of a two-component BEC.

Funding. The author acknowledges the Department of Science and Technology (DST), India (Grants # DIA/2018/000029, CRG/2019/000412) for supports.

Disclosures. The author has no Financial or Non-financial interests to disclose. The author has no conflicts to disclose.

Data availability. No data was used in writing this manuscript.

REFERENCES

1. W. K. Wooters and W. H. Zurek, "A single quantum cannot be cloned," Nature **299**, 802 (1982).

- 2. W. K. Wooters and W. H. Zurek, "The no-cloning theorem," Physics Today **62**, 76 (2009).
- 3. N. J. Cerf and J. Fiurasek, "Optical Quantum Cloning," J. Progress in Optics **49**, 455 (2006).
- 4. V. Buzek and M. Hillery, "Quantum copying: Beyond the no-cloning theorem," *Phys. Rev.* A **54** 1844 (1996).
- 5. V. Scarani, S. Iblisdir, N. Gisin and A. Acín, "Quantum cloning," *Rev. Mod. Phys.* **77** 1225 (2005).
- 6. J. I. Cirac, M. Lewenstein, K. Mølmer and P. Zoller, "Quantum superposition states of Bose-Einstein condensates," Phys. Rev. A **57**, 1208 (1998).
- 7. L. Mandel, "Is a photon amplifier always polarization dependent?," Nature **30**4, 188 (1983).
- 8. C. M. Caves, "Quantum-mechanical noise in an interferometer," Phys. Rev. D 23, 1693 (1981).
- 9. C. M. Caves, "Quantum limits on noise in linear amplifiers," Phys. Rev. D **26**, 1817 1982).
- J. A. Levenson, K. Bencheikh, D. J. Lovering, P. Vidakovic and C. Simonneau, "Quantum noise in optical parametric amplification: a means to achieve noiseless optical functions," Quantum. Semiclass. Opt. 9, 221 (1997).
- 11. S. Kocsis, G. Y. Xiang, T. C. Ralph and G. J. Pryde, "Heralded noiseless amplification of a photon," Nat. Phys. **9**, 23 (2013).
- 12. A. Zavatta, J. Fiurasek and M. Bellini, "A high-fidelity noiseless amplifier for quantum light states," Nat. Photon. **5**, 52 (2011).
- 13. G. Y. Xiang, T. C. Ralph, A. P. Lund, N. Walk and G. J. Pryde, "Heralded noiseless linear amplification and distillation of entanglement," Nat. Photon. **4**, 316 (2010).
- 14. M. A. Usuga, C. R. Müller, C. Wittmann, P. Marek, R. Filip, C. Marquardt, Gerd Leuchs, and U. L. Andersen, "Noise-powered probabilistic concentration of phase information," Nat. Phys. **6**, 767 (2010).

- 15. F. Ferreyrol, M. Barbieri, R. Blandino, S. Fossier, R. Tualle-Brouri and P. Grangier, "Implementation of a Nondeterministic Optical Noiseless Amplifier," Phys. Rev. Lett. **104**, 123603 (2010).
- 16. Z. Y. Ou, "Fundamental quantum limit in precision phase measurement," Phys. Rev. A **55**, 2598 (1997).
- 17. C. I. Osorio, N. Bruno, N. Sangouard, H. Zbinden, N. Gisin, and R. T. Thew "Heralded photon amplification for quantum communication," Phys. Rev. A 86, 023815 (2012).
- 18. A. J. Leggett, "Broken Gauge Symmetry in a Bose Condensate, in Bose-Einstein Condensation," edited by A. Griffin, D. W. Snoke and S. Stringari, Cambridge University Press, (1995).
- 19. R. Schnabel, "Squeezed states of light and their applications in laser Interferometers," Physics Reports **684**,1 (2017).
- 20. P. Xu, S. N. Zhu, "Review Article: Quasi-phase-matching engineering of entangled photons," AIP Advances 2, 041401 (2012).
- 21. W. Tittel, J. Brendel, H. Zbinden, and N. Gisin, "Quantum Cryptography Using Entangled Photons in Energy-Time Bell States," Phys. Rev. Lett. **84**, 4737 (2000).
- 22. H. P. Specht, J. Bochmann, M. Mucke, B. Weber, E. Figueroa, D. L. Moehring and G. Rempe, "Phase shaping of single-photon wave packets," Nat. Photonics 3, 469 (2009).
- 23. Sylvain Fasel, Nicolas Gisin, Grégoire Ribordy, Valerio Scarani, and Hugo Zbinden, "Quantum Cloning with an Optical Fiber Amplifier," Phys. Rev. Lett. **89**, 107901 (2002).
- 24. A. J. Leggett and F. Sols, "On the concept of spontaneously broken gauge symmetry in condensed matter physics," Found. Phys. **21**, 353 (1991).
- 25. A. J. Leggett, "Number-Phase Fluctuations in Two-Band Superconductors" Prog. Theor. Phys. **36**, 901 (1966).
- 26. G. Wang, C. Robert, M. M. Glazov, F. Cadiz, E. Courtade, T. Amand, D. Lagarde, T. Taniguchi, K. Watanabe, B. Urbaszek and X. Marie, "In-Plane Propagation of Light in Transition Metal Dichalcogenide Monolayers: Optical Selection Rules," Phys. Rev. Lett. **119**, 047401 (2017).
- 27. Y. Chi et. al, "A programmable qudit-based quantum processor," Nat. Comm. **13**, 1166 (2022).
- 28. Y. Wang, Z. Hu, B. C. Sanders and S. Kais, "Qudits and High-Dimensional Quantum Computing," Front. Phys. **8**, 589504 (2020).
- 29. D. Cozzolino, B. D. Lio, D. Bacco and L. K. Oxenløwe, "High-Dimensional Quantum Communication: Benefits, Progress, and Future Challenges," Adv. Quantum Technol. **2**, 1900038 (2019).
- 30. Takeshi Umeki, Osamu Tadanaga, Atsushi Takada, and Masaki Asobe, "Phase sensitive degenerate parametric amplification using directly-bonded PPLN ridge waveguides" Opt. Express 19, 6326 (2011).
- 31. D. Bruss, G. M. D'Ariano, C. Macchiavello and M. F. Sacchi, "Approximate quantum cloning and the impossibility of superluminal information transfer," Phys. Rev. A **62**, 062302 (2000).
- 32. C. Simon, V. Bužek and N. Gisin, "The no-signaling condition and quantum dynamics," Phys. Rev. Lett. **87**, 170405 (2001).

- 33. N. Herbert, "FLASH—A superluminal communicator based upon a new kind of quantum measurement," Found. Phys. **12**, 1171 (1982).
- 34. D. Dieks, "Communication by EPR devices," Phys. Lett. A 92, 271 (1982).
- 35. A. Einstein, B. Podolsky and N. Rosen, "Can Quantum-Mechanical Description of Physical Reality Be Considered Complete?," Phys. Rev. A **47**, 777 (1935).