

Global Gevrey regularity and analyticity of a weakly dissipative Camassa-Holm equation

Zhiying Meng¹ * and Zhaoyang Yin^{1,2} †

¹Department of Mathematics, Sun Yat-sen University,
Guangzhou, 510275, China

²Faculty of Information Technology,
Macau University of Science and Technology, Macau, China

Abstract

This work is concerned with the Gevrey regularity and analyticity of the solution to a weakly dissipative Camassa-Holm system. We first demonstrate the local Gevrey regularity and analyticity of this equation. Then, we discuss the continuity of the data-to-solution map. Finally, we obtain the global Gevrey regularity of this system in Gevrey class G_σ with $\sigma \geq 1$ in time.

Keywords: A weakly dissipative Camassa-Holm equation; analyticity; Gevrey class; global Gevrey regularity

Mathematics Subject Classification: 35Q53, 35B30, 35C07, 35G25

Contents

1	Introduction	1
2	Preliminaries	2
3	Local Gevrey regularity and analyticity	5
4	Continuity of the data-to-solution map	7
5	Global Gevrey regularity and analyticity	8
	References	11

1 Introduction

*email: mengzhy3@mail2.sysu.edu.cn

†email: mcsyzy@mail.sysu.edu.cn

In this paper, we study the following weakly dissipative Camassa-Holm equation [18]

$$\begin{cases} u_t - u_{txx} + 3uu_x + \lambda(u - u_{xx}) = 2uu_x + uu_{xx} + \alpha u + \beta u^2 u_x + \gamma u^3 u_x + \Gamma u_{xxx}, & t > 0, \\ u(0, x) = u_0, & x \in \mathbb{R}, \end{cases} \quad (1.1)$$

where $\alpha, \beta, \gamma, \Gamma$ are real number and λ is a positive number. Let $\Lambda^{-2} = (1 - \partial_{xx})^{-1}$, $h(u) = (\alpha + \Gamma)u + \frac{\beta}{3}u^3 + \frac{\gamma}{4}u^4$. The above equation can be reformulated in the following form

$$\begin{cases} u_t + (u + \Gamma)u_x + \lambda u = Q, & t > 0, \quad x \in \mathbb{R}, \\ u(0, x) = u_0, \end{cases} \quad (1.2)$$

where $Q = -\Lambda^{-2}\partial_x\left(-h(u) + u^2 + \frac{1}{2}u_x^2\right)$. For $\lambda = \alpha = \beta = \gamma = \Gamma = 0$, the equation (1.2) reads as the Camassa-Holm (CH) equation

$$(1 - \partial_x^2)u_t = 3uu_x - 2u_x u_{xx} - uu_{xxx}, \quad (1.3)$$

which has a bi-Hamiltonian structure and is completely integrable [6, 10]. The local well-posedness of CH equation for initial data $u_0 \in H^s$ with $s > \frac{3}{2}$ has been proved in [7, 8, 31]. In addition, many researchers established the local well-posedness of CH equation in Besov spaces $B_{p,r}^s$ with (i) $s > 1 + \frac{1}{p}$; (ii) $s = 1 + \frac{1}{p}$, $p \in [1, \infty)$, $r = 1$ [11, 12, 20, 23, 32]. The CH equation has not only global strong solutions [7–9] but also the solutions will blow up in finite time [3, 7, 8]. The authors established the existence of the unique global solutions [5]. In [4], Bressan and Constantin proved the global dissipative solutions of the CH equation. Its local ill-posedness problem was presented in [14, 15, 21]. The analyticity for the solutions of CH equation were studied in [2, 16]. In [24], the authors discussed the Gevrey regularity for the Camassa-Holm type systems by a generalized Ovsyannikov theorem.

Recently, the author established the local well-posedness of system (1.2) in Sobolev spaces H^s with $s > \frac{3}{2}$ [18]. In [25], the authors proved that this equation is locally well-posed in Besov spaces and the existence of global strong solutions under the condition that small initial datum. And the existence and uniqueness of weak solutions were presented in [26]. The analyticity and Gevrey regularity of the system (1.2) has not been investigated. Hence, in this paper, using the generalized Ovsyannikov theorem [24], we get the local analyticity and Gevrey regularity of the solutions to system (1.2). And we see that the continuity of the data-to-solution map. On the other hand, following the idea [19], we employ the global analyticity and Gevrey regularity of this system.

Our paper is organized as follows. In Section 2, we give some preliminary results. In Sections 3, 4, we discuss the the local analyticity and Gevrey regularity, and the continuity of the data-to-solution map of the equation (1.2). In Section 5, we study the global analyticity and Gevrey regularity.

2 Preliminaries

In this section, we give some theorems and some lemmas that will be of use to prove our main results. Now, we study the Cauchy problem for the above system which can be rewritten in the following abstract form

$$\begin{cases} \frac{du}{dt} = F(t, u(t)), \\ u(0, x) = u_0. \end{cases} \quad (2.1)$$

Theorem 2.1. [1, 27, 29] Let $\{X_\delta\}_{0 < \delta < 1}$ be a scale of decreasing Banach spaces, namely, for any $0 < \delta' < \delta$, we have $X_\delta \subset X_{\delta'}$ and $\|\cdot\|_{\delta'} \leq \|\cdot\|_\delta$. Given $T, R > 0$. Let $u_0 \in X_1$, assume that
(1) For any $0 < \delta' < \delta < 1$, the function $t \mapsto u(t)$ is holomorphic in $|t| < T$ and continuous on $|t| < T$ with values in X_δ and

$$\sup_{|t| < T} \|u(t)\|_\delta < R.$$

Then $t \mapsto F(t, u(t))$ is a holomorphic function on $|t| < T$ with values in $X_{\delta'}$.

(2) For any $0 < \delta' < \delta < 1$, and any $u, v \in \overline{B(u_0, R)} \subset X_\delta$, there exists a positive constant L depending on u_0 and R such that

$$\sup_{|t| < T} \|F(t, u(t)) - F(t, v(t))\|_{\delta'} \leq \frac{L}{\delta' - \delta} \|u - v\|_\delta.$$

(3) For any $0 < \delta < 1$, there exists a positive constant M depending on u_0 such that

$$\sup_{|t| < T} \|F(t, u_0(t))\|_{\delta'} \leq \frac{M}{1 - \delta}.$$

Then there exist a $T_0 \in (0, T)$ and a unique solution to the Cauchy problem (2.1), which for any $\delta \in (0, 1)$ is holomorphic in $|t| < T_0(1 - \delta)$ with values in X_δ .

The above theorem was first put forwarded by Ovsyannikov in [28–30]. Note that the original Ovsyannikov theorem is not valid for the Gerver class. Owing to this kind of space cannot such that the condition (2) hold in Theorem 2.1. Hence, for the Gerver class, we prove that

$$\sup_{|t| < T} \|F(t, u(t)) - F(t, v(t))\|_{\delta'} \leq \frac{L}{(\delta' - \delta)^\sigma} \|u - v\|_\delta,$$

where $\sigma \geq 1$. As $\sigma > 1$, the above inequality is weaker than the condition (2). Recently, Luo and Yin build a new auxiliary function to get a generalized Ovsyannikov theorem by modifying the proof of [24].

Theorem 2.2. [24, 33] Let $\{X_\delta\}_{0 < \delta < 1}$ be a scale of decreasing Banach spaces, namely, for any $0 < \delta' < \delta$, we have $X_\delta \subset X_{\delta'}$ and $\|\cdot\|_{\delta'} \leq \|\cdot\|_\delta$. Given $T, R > 0$. Let $u_0 \in X_1$, assume that
(1) For any $0 < \delta' < \delta < 1$, the function $t \mapsto u(t)$ is holomorphic in $|t| < T$ and continuous on $|t| < T$ with values in X_δ and

$$\sup_{|t| < T} \|u(t)\|_\delta < R.$$

Then $t \mapsto F(t, u(t))$ is a holomorphic function on $|t| < T$ with values in $X_{\delta'}$.

(2) For any $0 < \delta' < \delta < 1$, and any $u, v \in \overline{B(u_0, R)} \subset X_\delta$, there exists a positive constant L depending on u_0 and R such that

$$\sup_{|t| < T} \|F(t, u(t)) - F(t, v(t))\|_{\delta'} \leq \frac{L}{(\delta' - \delta)^\sigma} \|u - v\|_\delta.$$

(3) For any $0 < \delta < 1$, there exists a positive constant M depending on u_0 such that

$$\sup_{|t| < T} \|F(t, u_0(t))\|_{\delta'} \leq \frac{M}{1 - \delta}.$$

Then there exists a $T_0 \in (0, T)$ and a unique solution to the Cauchy problem (2.1), which for any $\delta \in (0, 1)$ is holomorphic in $|t| < \frac{T_0(1-\delta)^\sigma D_\sigma}{2^\sigma - 1}$ with values in X_δ and $D_\sigma = \frac{1}{2^\sigma - 2 + \frac{1}{2^{\sigma+1}}}$.

Now, we give a Banach spaces, which can be used to prove Theorem 2.2 by using the fixed point argument.

Definition 2.3. [24, 33] Let $\sigma \geq 1$. For any $a > 0$, we represent by E_a the space of functions $u(t)$ which for each $0 < \delta < 1$ and $|t| < \frac{a(1-\delta)^\sigma D_\sigma}{2^\sigma - 1}$ with $D_\sigma = \frac{1}{2^\sigma - 2 + \frac{1}{2^\sigma + 1}}$ are holomorphic and continuous functions of t with values in X_δ such that

$$\|u\|_{E_T} := \sup_{|t| < \frac{a(1-\delta)^\sigma}{2^\sigma - 1}} \left(\|u(t)\|_{G_{\sigma,s}^\delta} (1-\delta)^\sigma \sqrt{1 - \frac{|t|}{a(1-\delta)^\sigma}} \right) < +\infty.$$

Remark 2.4. [24, 33] Indeed, $T_0 = \min\left\{\frac{1}{2^{2\sigma+4}L}, \frac{(2^\sigma-1)R}{(2^\sigma-1)2^{2\sigma+3}LR+MD_\sigma}\right\}$ with $D_\sigma = \frac{1}{2^\sigma - 2 + \frac{1}{2^\sigma + 1}}$, which provides a lower bound of the lifespan.

Remark 2.5. If $\sigma = 1$, Theorem 2.1 reduces to the so-called abstract Cauchy-Kovalevsky theorem.

Now, we give the definition of Sobolev-Gevrey spaces and recall some properties are often used.

Definition 2.6. [13] Let $\sigma, \delta > 0$ and s be a real number. A function $f \in G_{\sigma,s}^\delta(\mathbb{R}^d)$ if and only of $f \in \mathcal{C}^\infty(\mathbb{R}^d)$ and satisfies

$$\|f\|_{G_{\sigma,s}^\delta} = \left(\int_{\mathbb{R}} (1 + |\xi|^2)^s e^{2\delta(1+|\xi|^2)^{\frac{1}{2\sigma}}} |\hat{f}(\xi)|^2 d\xi \right)^{\frac{1}{2}} < \infty.$$

A function $f \in G_{\sigma,s}^\delta(\mathbb{R}^d)$ if and only of $f \in \mathcal{C}^\infty(\mathbb{R}^d)$ and satisfies

$$\|f\|_{\tilde{G}_{\sigma,s}^\delta} = \left(\int_{\mathbb{R}} (1 + |\xi|^2)^s e^{2\delta|\xi|^{\frac{1}{\sigma}}} |\hat{f}(\xi)|^2 d\xi \right)^{\frac{1}{2}} < \infty.$$

Definition 2.7. [22] A function is of Gevrey class $\sigma \geq 1$, if there exist $\delta > 0$, $s \geq 0$ such that $f \in G_{\sigma,s}^\delta(\mathbb{R}^d)$. Denote the the functions of Gevrey class σ

$$f \in G_\sigma(\mathbb{R}^d) := \cup_{\delta > 0, s \in \mathbb{R}} G_{\sigma,s}^\delta(\mathbb{R}^d).$$

Remark 2.8. [17] Denote the Fourier multiplier $e^{\delta\Lambda^{\frac{1}{\sigma}}}$ and $e^{\delta(-\Delta)^{\frac{1}{2\sigma}}}$ by

$$e^{\delta\Lambda^{\frac{1}{\sigma}}} f = \mathcal{F}^{-1} \left(e^{\delta(1+|\xi|^2)^{\frac{1}{2\sigma}}} \hat{f} \right) \quad \text{and} \quad e^{\delta(-\Delta)^{\frac{1}{2\sigma}}} f = \mathcal{F}^{-1} \left(e^{\delta|\xi|^{\frac{1}{\sigma}}} \hat{f} \right).$$

Then we have $\|f\|_{G_{\sigma,s}^\delta(\mathbb{R}^d)} = \|e^{\delta\Lambda^{\frac{1}{\sigma}}} f\|_{H^s(\mathbb{R}^d)}$ and $\|f\|_{\tilde{G}_{\sigma,s}^\delta} = \|e^{\delta(-\Delta)^{\frac{1}{2\sigma}}} f\|_{H^s(\mathbb{R}^d)}$. Moreover, for any $\sigma \geq 1$, we get

$$\|f\|_{\tilde{G}_{\sigma,s}^\delta} \leq \|f\|_{G_{\sigma,s}^\delta} \leq e^\delta \|f\|_{\tilde{G}_{\sigma,s}^\delta}.$$

If $0 < \sigma < 1$, it is called ultra-analytic function. If $\sigma = 1$, it is usual analytic function and δ is called the radius of analyticity. If $\delta > 1$, it is the Gevrey class function.

Proposition 2.9. If $0 < \delta' < \delta$, $0 < \sigma' < \sigma$ and $s' < s$. Then we have $G_{\sigma,s}^\delta(\mathbb{R}^d) \hookrightarrow G_{\sigma,s}^{\delta'}(\mathbb{R}^d)$, $G_{\sigma',s}^\delta(\mathbb{R}^d) \hookrightarrow G_{\sigma,s}^\delta(\mathbb{R}^d)$ and $G_{\sigma,s}^\delta(\mathbb{R}^d) \hookrightarrow G_{\sigma,s'}^\delta(\mathbb{R}^d)$.

Proposition 2.10. [24] Let $f \in G_{\sigma,s}^\delta(\mathbb{R})$. Assume that $0 < \delta' < \delta$, $\sigma > 0$ and $s \in \mathbb{R}$. Then we have

$$\begin{aligned} \|\partial_x f\|_{G_{\sigma,s}^{\delta'}(\mathbb{R})} &\leq \frac{e^{-\sigma} \sigma^\sigma}{2(\delta' - \delta)^\sigma} \|f\|_{G_{\sigma,s}^\delta(\mathbb{R})}, \\ \|(1 - \partial_{xx})^{-1} f\|_{G_{\sigma,s}^{\delta'}(\mathbb{R})} &\leq \|f\|_{G_{\sigma,s-2}^\delta(\mathbb{R})} \leq \|f\|_{G_{\sigma,s}^\delta(\mathbb{R})}, \\ \|(1 - \partial_{xx})^{-1} \partial_x f\|_{G_{\sigma,s}^{\delta'}(\mathbb{R})} &\leq \|f\|_{G_{\sigma,s-1}^\delta(\mathbb{R})} \leq \|f\|_{G_{\sigma,s}^\delta(\mathbb{R})}. \end{aligned}$$

Proposition 2.11. *Let $s > \frac{1}{2}$, $\sigma \geq 1$ and $\delta > 0$. Then $G_{\sigma,s}^\delta(\mathbb{R})$ is an algebra. Moreover, there exists two constants C_s and \bar{C}_s such that*

$$\begin{aligned} \|fg\|_{G_{\sigma,s}^\delta(\mathbb{R})} &\leq C_s \|f\|_{G_{\sigma,s}^\delta(\mathbb{R})} \|g\|_{G_{\sigma,s}^\delta(\mathbb{R})}, \\ \|fg\|_{G_{\sigma,s-1}^\delta(\mathbb{R})} &\leq \bar{C}_s \|f\|_{G_{\sigma,s}^\delta(\mathbb{R})} \|g\|_{G_{\sigma,s-1}^\delta(\mathbb{R})}. \end{aligned}$$

Lemma 2.12. *[24, 33] Let $\sigma \geq 1$. For each $a > 0$, $u \in E_a$, $0 < \delta < 1$ and $0 \leq t < \frac{a(1-\delta)^\sigma D_\sigma}{2^\sigma - 1}$ with $D_\sigma = \frac{1}{2^\sigma - 2 + \frac{1}{2^{\sigma+1}}}$. Then we have*

$$\int_0^t \frac{\|u(\tau)\|_{\delta(\tau)}}{(\delta(\tau) - \delta)^\sigma} \leq \frac{a2^{2\delta+3}\|u\|_{E_a}}{(1-\delta)^\sigma} \sqrt{\frac{a(1-\delta)^\sigma}{a(1-\delta)^\sigma - t}},$$

where

$$\delta(\tau) = \frac{1}{2}(1+\delta) + \left(\frac{1}{2}\right)^{2+\frac{1}{\sigma}} \left([(1-\delta)^\sigma - \frac{t}{a}]^{\frac{1}{\sigma}} - [(1-\delta)^\sigma + (2^{\sigma+1} - 1)\frac{t}{a}]^{\frac{1}{\sigma}} \right) \in (0, 1).$$

Lemma 2.13. *[25] Let $u_0 \in B_{p,r}^s$ with $s > \frac{3}{2}$, $p = r = 2$. A constant ϵ such that if*

$$H_0 \triangleq |\alpha| + |\Gamma| + \|u_0\|_{B_{p,r}^s} + \frac{|\beta|}{3} \|u_0\|_{B_{p,r}^s}^2 + \frac{|\gamma|}{4} \|u_0\|_{B_{p,r}^s}^3 \leq \lambda\epsilon, \quad (2.2)$$

for all real numbers $\alpha, \beta, \gamma, \Gamma$ and $\lambda > 0$. Then the solution to the system (1.2) exists globally in time. Moreover, for any $t \in [0, \infty)$ we have

$$H(t) \triangleq |\alpha| + |\Gamma| + \|u(t)\|_{B_{p,r}^s} + \frac{|\beta|}{3} \|u(t)\|_{B_{p,r}^s}^2 + \frac{|\gamma|}{4} \|u(t)\|_{B_{p,r}^s}^3 \leq H_0. \quad (2.3)$$

3 Local Gevery regularity and analyticity

This section is devoted to establishing the local Gevery regularity and analyticity. Our main results can be stated as follows.

Theorem 3.1. *Let $\sigma \geq 1$, $s > \frac{3}{2}$. Assume that $u_0 \in G_{\sigma,s}^1$. Then for any $0 < \delta < 1$, there exists a $T_0 > 0$ such that the system (1.2) has a unique solution u which is holomorphic in $|t| < \frac{T_0(1-\delta)^\sigma}{2^\sigma - 1}$ with values in $G_{\sigma,s}^\delta(\mathbb{R})$. Moreover $T_0 = \frac{1}{2^{2\sigma+8}C'(e^{-\sigma\sigma}+2)(1+\|u_0\|_{G_{\sigma,s}^1})^4}$, where the positive constant C' depending on $s, \alpha, \beta, \gamma, \lambda, \Gamma$.*

Proof. The equation (1.2) can be written as follows

$$\begin{cases} \frac{d}{dt}u = F(t, u(t)), \\ u(0, x) = u_0, \end{cases} \quad (3.1)$$

where $F(t, u(t)) = -(u + \Gamma)u_x - \lambda u - P_x$. Fixed a $\sigma \geq 1$ and $s > \frac{3}{2}$. Theorem 2.2 entails that $\{G_{\sigma,s}^\delta\}_{0 < \delta < 1}$ is a scale of decreasing Banach spaces. For any $0 < \delta' < \delta$, it follows from Propositions 2.9-2.11 that

$$\|F(t, u(t))\|_{G_{\sigma,s}^{\delta'}} \leq \frac{1}{2} \|(u^2)_x\|_{G_{\sigma,s}^{\delta'}} + |\Gamma| \|u_x\|_{G_{\sigma,s}^{\delta'}} + C_s \left(\|h(u)\|_{G_{\sigma,s-1}^{\delta'}} + \|u^2\|_{G_{\sigma,s-1}^{\delta'}} + \|u_x^2\|_{G_{\sigma,s-1}^{\delta'}} \right)$$

$$\leq \frac{C_s e^{-\sigma} \sigma^\sigma}{2(\delta' - \delta)^\sigma} \|u^2\|_{G_{\sigma,s}^\delta} + \frac{C_{s,\Gamma} e^{-\sigma} \sigma^\sigma}{(\delta' - \delta)^\sigma} \|u\|_{G_{\sigma,s}^\delta} + \|P_x\|_{G_{\sigma,s}^{\delta'}}. \quad (3.2)$$

Using Propositions 2.9-2.11 again, we have

$$\begin{aligned} \|P_x\|_{G_{\sigma,s}^{\delta'}} &\leq C_s \left(\|h(u)\|_{G_{\sigma,s-1}^{\delta'}} + \|u^2\|_{G_{\sigma,s-1}^{\delta'}} + \|u_x^2\|_{G_{\sigma,s-1}^{\delta'}} \right) \\ &\leq C' \left(\|u\|_{G_{\sigma,s}^\delta} + \|u\|_{G_{\sigma,s}^\delta}^2 + \|u\|_{G_{\sigma,s}^\delta}^3 + \|u\|_{G_{\sigma,s}^\delta}^4 + \|u\|_{G_{\sigma,s}^\delta}^5 \right) \\ &\leq C' \|u\|_{G_{\sigma,s}^\delta} (1 + \|u\|_{G_{\sigma,s}^\delta})^4, \end{aligned}$$

where C' depending on $s, \alpha, \beta, \gamma, \lambda, \Gamma$.

Adding up the above estimate to (3.2), one can deduce that

$$\|F(t, u(t))\|_{G_{\sigma,s}^{\delta'}} \leq \frac{C'(e^{-\sigma} \sigma^\sigma + 2)}{2(\delta' - \delta)^\sigma} \|u\|_{G_{\sigma,s}^\delta} (1 + \|u\|_{G_{\sigma,s}^\delta})^4, \quad (3.3)$$

which means that $F(t, u(t))$ satisfies the condition (1) of Theorem 2.1. Adopting the similar procedure for $F(u_0)$ as above, we arrive at

$$\|F(u_0)\|_{G_{\sigma,s}^\delta} \leq \frac{C'(e^{-\sigma} \sigma^\sigma + 2)}{2(1 - \delta)^\sigma} \|u_0\|_{G_{\sigma,s}^1} (1 + \|u_0\|_{G_{\sigma,s}^1})^4.$$

Therefore, we obtain that $F(t, u(t))$ satisfies the condition (3) in Theorem 2.1 with

$$M = \frac{C'(e^{-\sigma} \sigma^\sigma + 2)}{2} \|u_0\|_{G_{\sigma,s}^1} (1 + \|u_0\|_{G_{\sigma,s}^1})^4.$$

On the other hand, we have to check that $F(t, u(t))$ satisfies the condition (2) of Theorem 2.1. Let $\|u - u_0\|_{G_{\sigma,s}^\delta} \leq R$ and $\|v - v_0\|_{G_{\sigma,s}^\delta} \leq R$. One can get from Theorem 2.1 and Propositions 2.9-2.11 that

$$\begin{aligned} \|F(u) - F(v)\|_{G_{\sigma,s}^{\delta'}} &\leq \frac{C_s e^{-\sigma} \sigma^\sigma}{2(\delta' - \delta)^\sigma} \|u^2 - v^2\|_{G_{\sigma,s}^\delta} + \frac{C_{s,\Gamma} e^{-\sigma} \sigma^\sigma}{(\delta' - \delta)^\sigma} \|u - v\|_{G_{\sigma,s}^\delta} + \|P_x(u) - P_x(v)\|_{G_{\sigma,s}^{\delta'}} \\ &\leq \frac{C_s e^{-\sigma} \sigma^\sigma}{2(\delta' - \delta)^\sigma} \|u - v\|_{G_{\sigma,s}^\delta} \|u + v\|_{G_{\sigma,s}^\delta} + \frac{C_{s,\Gamma} e^{-\sigma} \sigma^\sigma}{(\delta' - \delta)^\sigma} \|u - v\|_{G_{\sigma,s}^\delta} \\ &\quad + C' \|u - v\|_{G_{\sigma,s}^\delta} (1 + \|u + v\|_{G_{\sigma,s}^\delta})^4 \\ &\leq \frac{C_s e^{-\sigma} \sigma^\sigma}{2(\delta' - \delta)^\sigma} \|u - v\|_{G_{\sigma,s}^\delta} (\|u_0\|_{G_{\sigma,s}^\delta} + R + 1) + C' \|u - v\|_{G_{\sigma,s}^\delta} (1 + \|u_0\|_{G_{\sigma,s}^1} + R)^4 \\ &\leq \frac{C'(e^{-\sigma} \sigma^\sigma + 2)}{2(\delta' - \delta)^\sigma} \|u - v\|_{G_{\sigma,s}^\delta} (1 + \|u_0\|_{G_{\sigma,s}^1} + R)^4. \end{aligned} \quad (3.4)$$

Hence, we get F satisfies the condition (2) of Theorem 2.1 with

$$L = C'(e^{-\sigma} \sigma^\sigma + 2)(\|u_0\|_{G_{\sigma,s}^1} + R + 1)^4.$$

Then, we conclude that the local existence result of (3.1) with the Gevrey regularity or analyticity, and

$$T_0 = \min\left\{\frac{1}{2^{2\sigma+4}L}, \frac{(2^\sigma - 1)R}{(2^\sigma - 1)^{2^{2\sigma+3}}LR + MD_\sigma}\right\},$$

with $D_\sigma = \frac{1}{2^\sigma - 2 + \frac{1}{2^{\sigma+1}}}$.

On the other hand, choosing $R = 1 + \|u_0\|_{G_{\sigma,s}^1}$, we infer that $L = 2^4 C'(e^{-\sigma} \sigma^\sigma + 2)(1 + \|u_0\|_{G_{\sigma,s}^1})^4$ and $M \leq 2^{2\sigma+3}LR$. Moreover, we get

$$T_0 = \frac{1}{2^{2\sigma+8}C'(e^{-\sigma} \sigma^\sigma + 2)(1 + \|u_0\|_{G_{\sigma,s}^1})^4}.$$

Thus, we finish the proof of Theorem 3.1. \square

4 Continuity of the data-to-solution map

In this section, we consider the continuity of the data-to-solution map for initial data and solutions in Theorem 3.1. We first introduce a definition to illustrate what means the data-to-solution map is continuous from $G_{\sigma,s}^1(\mathbb{R})$ into the solutions space.

Definition 4.1. *Let $\sigma \geq 1$ and $s > \frac{3}{2}$. Then, the data-to-solution map $u_0 \mapsto u$ of the system (1.2) is continuous if for a given initial data $u_0^\infty \in G_{\sigma,s}^1$ there exists a $T = T(\|u_0\|_{G_{\sigma,s}^1}, \|u_0^\infty\|_{G_{\sigma,s}^1}) > 0$, such that for any sequence $u_0^n \in G_{\sigma,s}^1$ and $\|u_0^n - u_0^\infty\|_{G_{\sigma,s}^1} \rightarrow 0$ as $n \rightarrow \infty$, the corresponding solutions u^n of system (1.2) satisfy $\|u^n - u^\infty\|_{E_T} \rightarrow 0$ as $n \rightarrow \infty$, where*

$$\|u\|_{E_T} := \sup_{|t| < \frac{T(1-\delta)^\sigma}{2^\sigma - 1}} \left(\|u(t)\|_{G_{\sigma,s}^\delta} (1-\delta)^\sigma \sqrt{1 - \frac{|t|}{T(1-\delta)^\sigma}} \right).$$

Theorem 4.2. *Given $\sigma \geq 1$ and $s > \frac{3}{2}$. Let $u_0 \in G_{\sigma,s}^1(\mathbb{R})$. Then the data-to-solution map $u_0 \mapsto u$ of the system (1.2) is continuous from $G_{\sigma,s}^1(\mathbb{R})$ into the solutions space.*

Proof. In general, assume that $t \geq 0$. Defining

$$T^\infty = \frac{1}{2^{2\sigma+8}C'(e^{-\sigma}\sigma^\sigma + 2)(1 + \|u_0^\infty\|_{G_{\sigma,s}^1})^4}, \quad T^n = \frac{1}{2^{2\sigma+8}C'(e^{-\sigma}\sigma^\sigma + 2)(1 + \|u_0^n\|_{G_{\sigma,s}^1})^4}.$$

Owing to $\|u_0^n - u_0^\infty\|_{G_{\sigma,s}^1} \rightarrow 0$ as $n \rightarrow \infty$, there exists a constant N such that if $n \geq N$, we obtain

$$\|u_0^n\|_{G_{\sigma,s}^1} \leq \|u_0^\infty\|_{G_{\sigma,s}^1} + 1. \quad (4.1)$$

Set

$$T = \frac{1}{2^{2\sigma+8}C'(e^{-\sigma}\sigma^\sigma + 2)(2 + \|u_0^n\|_{G_{\sigma,s}^1})^4}. \quad (4.2)$$

This means that $T \leq \min\{T^n, T^\infty\}$ for any $n \geq N$. A similar argument based on the proof of Theorem 3.1, we claim that T^n and T^∞ are the existence time of the solutions u^n and u^∞ corresponding to u_0^n and u_0^∞ respectively. Therefore, we see that for any $n \geq N$,

$$u^\infty(t, x) = u_0^\infty + \int_0^t F(\tau, u^\infty(t, \tau)) d\tau, \quad 0 \leq t < \frac{T(1-\delta)^\sigma}{2^\sigma - 1}, \quad (4.3)$$

$$u^n(t, x) = u_0^n + \int_0^t F(\tau, u^n(t, \tau)) d\tau, \quad 0 \leq t < \frac{T(1-\delta)^\sigma}{2^\sigma - 1}, \quad (4.4)$$

with F is given in Theorem 3.1. Thus, for any $0 \leq t \leq \frac{T(1-\delta)^\sigma}{2^\sigma - 1}$ and $0 \leq \delta \leq 1$, we can obtain

$$\|u^n(t) - u^\infty(t)\|_\delta \leq \|u_0^n(t) - u_0^\infty(t)\|_\delta + \int_0^t \|F(u^n(\tau)) - F(u^\infty(\tau))\|_\delta d\tau. \quad (4.5)$$

Define that

$$\delta(t) = \frac{1}{2}(1 + \delta) + \left(\frac{1}{2}\right)^{2+\frac{1}{\sigma}} \left([(1-\delta)^\sigma - \frac{t}{T}]^{\frac{1}{\sigma}} - [(1-\delta)^\sigma + (2^{\sigma+1} - 1)\frac{t}{T}]^{\frac{1}{\sigma}} \right).$$

Making use of Lemma 2.12, we know that $\delta < \delta(t) < 1$. One can get from (4.5) that

$$\|F(u^n(\tau)) - F(u^\infty(\tau))\|_\delta \leq \frac{L\|u^n(\tau) - u^\infty(\tau)\|_{\delta(\tau)}}{(\delta(\tau) - \delta)^\sigma}, \quad (4.6)$$

where $L = 2^4 C'(e^{-\sigma} \sigma^\sigma + 2)(1 + \|u_0\|_1)^4$. Hence, we can deduce that

$$\|u^n(t) - u^\infty(t)\|_\delta \leq \|u_0^n(t) - u_0^\infty(t)\|_\delta + L \int_0^t \frac{\|u^n(\tau) - u^\infty(\tau)\|_{\delta(\tau)}}{(\delta(\tau) - \delta)^\sigma} d\tau. \quad (4.7)$$

It follows from Lemma 2.12 that

$$\|u^n(t) - u^\infty(t)\|_\delta \leq \|u_0^n(t) - u_0^\infty(t)\|_\delta + \frac{T2^{2\delta+3}\|u^n(t) - u^\infty(t)\|_{E_T}}{(1 - \delta)^\sigma} \sqrt{\frac{T(1 - \delta)^\sigma}{T(1 - \delta)^\sigma - t}}. \quad (4.8)$$

Combining (4.2) and $LT2^{2\sigma+3} < \frac{1}{2}$. Therefore, we obtain

$$\|u^n(t) - u^\infty(t)\|_\delta \leq \|u_0^n(t) - u_0^\infty(t)\|_\delta + \frac{\|u^n(t) - u^\infty(t)\|_{E_T}}{2(1 - \delta)^\sigma} \sqrt{\frac{T(1 - \delta)^\sigma}{T(1 - \delta)^\sigma - t}},$$

which implies

$$\begin{aligned} & \|u^n(t) - u^\infty(t)\|_{E_T} (1 - \delta)^\sigma \sqrt{1 - \frac{t}{T(1 - \delta)^\sigma}} \\ & \leq \|u_0^n - u_0^\infty\|_\delta (1 - \delta)^\sigma \sqrt{1 - \frac{t}{T(1 - \delta)^\sigma}} + \frac{1}{2} \|u^n(t) - u^\infty(t)\|_{E_T} \\ & \leq \|u_0^n - u_0^\infty\|_1 + \frac{1}{2} \|u^n(t) - u^\infty(t)\|_{E_T}. \end{aligned}$$

In view of the right hand side of the above estimate is independent of t and δ . By taking the supremum over $0 < \delta < 1$, $0 < t < \frac{T(1-\delta)^\sigma}{2^\sigma - 1}$, we see that

$$\|u^n(t) - u^\infty(t)\|_{E_T} \leq \|u_0^n - u_0^\infty\|_1 + \frac{1}{2} \|u^n(t) - u^\infty(t)\|_{E_T},$$

which means that

$$\|u^n(t) - u^\infty(t)\|_{E_T} \leq 2\|u^n(t) - u^\infty(t)\|_{E_T}.$$

This verifies the above inequality is valid for any $n \geq N$. \square

5 Global Gevery regularity and analyticity

In this section, we focus on the global Gevery regularity of solution to the system (1.2). Before stating the main result of this section, we introduce some lemmas.

Lemma 5.1. [17] *Let $\delta \geq 0$, $\sigma \geq 1$ and $s > 1$. Then, for any $\xi, \eta \in \mathbb{R}$, we have*

$$\begin{aligned} & |(1 + \xi^2)^{\frac{s}{2}} e^{\delta(1+\xi^2)^{\frac{1}{2\sigma}}} - (1 + \eta^2)^{\frac{s}{2}} e^{\delta(1+\eta^2)^{\frac{1}{2\sigma}}}| \\ & \leq C_s |\xi - \eta| \left((1 + |\xi - \eta|^2)^{\frac{s-1}{2}} + (1 + |\eta|^2)^{\frac{s-1}{2}} \right. \\ & \quad \left. + \delta[(1 + |\xi - \eta|^2)^{\frac{s-1}{2} + \frac{1}{2\sigma}} + (1 + |\eta|^2)^{\frac{s-1}{2} + \frac{1}{2\sigma}}] e^{\delta(1+|\xi-\eta|^2)^{\frac{1}{2\sigma}}} \right). \end{aligned}$$

Lemma 5.2. [17] Let $\delta \geq 0$, $\sigma \geq 1$ and $s > 1 + \frac{d}{2}$. Suppose that $u \in G_{\sigma, \frac{1}{2\sigma}}^\delta(\mathbb{R})$, $v \in G_{\sigma, \frac{1}{2\sigma}}^\delta(\mathbb{R})$. Then, we have

$$\begin{aligned} |\langle \Lambda^s e^{\delta \Lambda^{\frac{1}{\sigma}}} (u \cdot v), \Lambda^s e^{\delta \Lambda^{\frac{1}{\sigma}}} v \rangle| &\leq C \|\Lambda^s u\| \|\Lambda^s v\|^2 + C \delta \left(\|\Lambda^s e^{\delta \Lambda^{\frac{1}{\sigma}}} u\| \|\Lambda^{s+\frac{1}{\sigma}} e^{\delta \Lambda^{\frac{1}{\sigma}}} v\|^2 \right. \\ &\quad \left. + \|\Lambda^{s+\frac{1}{\sigma}} e^{\delta \Lambda^{\frac{1}{\sigma}}} u\| \|\Lambda^{s+\frac{1}{\sigma}} e^{\delta \Lambda^{\frac{1}{\sigma}}} v\| \|\Lambda^s e^{\delta \Lambda^{\frac{1}{\sigma}}} v\| \right), \end{aligned}$$

where $\langle \cdot, \cdot \rangle$ represents the scalar product of $L^2(\mathbb{R}^d)$, and $\|\cdot\| := \|\cdot\|_{L^2}$, Λ^s denotes for the inverse Fourier transform of $(1 + |\xi|^2)^{\frac{s}{2}} \hat{f}$.

Lemma 5.3. [17] For all $\delta \geq 0$, $\sigma \geq 1$, $s \in \mathbb{R}$ and $l > 0$. Let $u \in G_{\sigma, \frac{1}{2\sigma}}^\delta(\mathbb{R})$. Then we have the following estimate

$$\|u\|_{G_{\sigma, s}^\delta} \leq \sqrt{e} \|u\|_{H^s} + (2\delta)^{\frac{1}{2}} \|u\|_{G_{\sigma, s+\frac{1}{2\sigma}}^\delta}.$$

We can now state the main result of this section.

Theorem 5.4. Given $s > \frac{3}{2}$, $\sigma \geq 1$. Let $u_0 \in G_\sigma(\mathbb{R})$. Assume that $|\alpha| + |\Gamma| + \|u_0\|_{H^s} + \frac{|\beta|}{3} \|u_0\|_{H^s}^2 + \frac{|\gamma|}{4} \|u_0\|_{H^s}^3 \leq \lambda\epsilon$. Then there exists a unique global solution of system (1.2) in Gevery class σ , namely, for any $t \geq 0$, $u(t, \cdot)$ is of Gevery class σ .

Proof. We aim to get the global priori estimate of u in the time-dependent space $G_{\sigma, s}^{\delta(t)}$. One can apply the Fourier Galerkin approximating method to construct local solutions in $G_{\sigma, s}^{\delta(t)}$, globalize the result by the later estimate (5.9).

According to the definition of the Gevery class, we see that $G_{\sigma, s}^\delta \hookrightarrow G_{\sigma, \infty}^{\delta-\epsilon}$ for any $\epsilon > 0$. Notice that $u_0 \in G_\sigma(\mathbb{R})$. In general, we can suppose that $u_0 \in G_{\sigma, s}^1$. In the following, we will claim that there exists a $\delta(t)$ such that the solutions remain belong to the Gevery class G_σ . For any $t \in [0, T]$, it follows from the system (1.2) that

$$\begin{aligned} \frac{d}{dt} \|u\|_{G_{\sigma, s}^{\delta(t)}}^2 &= \frac{d}{dt} \int (1 + |\xi|^2)^s e^{2\delta(t)|\xi|^{\frac{1}{\sigma}}} \hat{u}(\xi) \bar{\hat{u}}(\xi) d\xi \\ &= 2\dot{\delta}(t) \int (1 + |\xi|^2)^s |\xi|^{\frac{1}{\sigma}} e^{2\delta(t)|\xi|^{\frac{1}{\sigma}}} \hat{u}(\xi) \bar{\hat{u}}(\xi) d\xi \\ &\quad + \mathbf{Re} \int (1 + |\xi|^2)^s e^{2\delta(t)|\xi|^{\frac{1}{\sigma}}} \left(-\widehat{uu_x} - \Gamma \hat{u} - \lambda \hat{u} - \widehat{P_x} \right) (\xi) \bar{\hat{u}}(\xi) d\xi, \end{aligned} \quad (5.1)$$

with \mathbf{Re} stands the real part of a complex number.

Taking advantage of Lemma 5.2, we get

$$\begin{aligned} \left| \int (1 + |\xi|^2)^s e^{2\delta(t)|\xi|^{\frac{1}{\sigma}}} \widehat{uu_x}(\xi) \bar{\hat{u}}(\xi) d\xi \right| &= |\langle \Lambda^s e^{\delta(t)\Lambda^{\frac{1}{\sigma}}} (uu_x), \Lambda^s e^{\delta(t)\Lambda^{\frac{1}{\sigma}}} u \rangle| \\ &\leq C \left(\|u\|_{H^s}^3 + \delta(t) \|u\|_{G_{\sigma, s}^{\delta(t)}} \|u\|_{G_{\sigma, s+\frac{1}{2\sigma}}^{\delta(t)}}^2 \right). \end{aligned} \quad (5.2)$$

Likewise, one can get the following estimates

$$\begin{aligned} &\left| \int (1 + |\xi|^2)^s e^{2\delta(t)(1+|\xi|^2)^{\frac{1}{2\sigma}}} \widehat{-P_x}(\xi) \bar{\hat{u}}(\xi) d\xi \right| \\ &= |\langle \Lambda^s e^{\delta(t)\Lambda^{\frac{1}{\sigma}}} \Lambda^{-2} \left(\partial_x (-h(u) - u^2 - \frac{1}{2} u_x^2) \right), \Lambda^s e^{\delta(t)\Lambda^{\frac{1}{\sigma}}} u \rangle| \\ &\leq \|\Lambda^{s-1} e^{\delta(t)\Lambda^{\frac{1}{\sigma}}} \left(\partial_x (-h(u) - u^2 - \frac{1}{2} u_x^2) \right)\| \|\Lambda^s e^{\delta(t)\Lambda^{\frac{1}{\sigma}}} u\| \end{aligned}$$

$$\leq C \left(\|u\|_{G_{\sigma,s}^{\delta(t)}}^2 + \|u\|_{G_{\sigma,s}^{\delta(t)}}^3 + \|u\|_{G_{\sigma,s}^{\delta(t)}}^4 + \|u\|_{G_{\sigma,s}^{\delta(t)}}^5 \right). \quad (5.3)$$

Combining Lemma 5.3 with $l = 1$, $l = \frac{2}{3}$, $l = \frac{1}{2}$ and $l = \frac{2}{5}$, we obtain

$$\begin{aligned} \|u\|_{G_{\sigma,s}^{\delta(t)}}^2 &\leq C \left(\|u\|_{H^s}^2 + \delta(t) \|u\|_{G_{\sigma,s+\frac{1}{2\sigma}}^{\delta(t)}}^2 \right), \\ \|u\|_{G_{\sigma,s}^{\delta(t)}}^3 &\leq C \left(\|u\|_{H^s}^3 + \delta(t) \|u\|_{G_{\sigma,s+\frac{1}{3\sigma}}^{\delta(t)}}^3 \right) \leq C \left(\|u\|_{H^s}^3 + \delta(t) \|u\|_{G_{\sigma,s}^{\delta(t)}} \|u\|_{G_{\sigma,s+\frac{1}{2\sigma}}^{\delta(t)}}^2 \right), \\ \|u\|_{G_{\sigma,s}^{\delta(t)}}^4 &\leq C \left(\|u\|_{H^s}^4 + \delta(t) \|u\|_{G_{\sigma,s+\frac{1}{4\sigma}}^{\delta(t)}}^4 \right) \leq C \left(\|u\|_{H^s}^4 + \delta(t) \|u\|_{G_{\sigma,s}^{\delta(t)}}^2 \|u\|_{G_{\sigma,s+\frac{1}{2\sigma}}^{\delta(t)}}^2 \right), \\ \|u\|_{G_{\sigma,s}^{\delta(t)}}^5 &\leq C \left(\|u\|_{H^s}^5 + \delta(t) \|u\|_{G_{\sigma,s+\frac{1}{5\sigma}}^{\delta(t)}}^5 \right) \leq C \left(\|u\|_{H^s}^5 + \delta(t) \|u\|_{G_{\sigma,s}^{\delta(t)}}^3 \|u\|_{G_{\sigma,s+\frac{1}{2\sigma}}^{\delta(t)}}^2 \right). \end{aligned} \quad (5.4)$$

By the same token, it yields

$$\left| \int (1 + |\xi|^2)^s e^{2\delta(t)(1+|\xi|^2)\frac{1}{2\sigma}} (-\Gamma\hat{u} - \lambda\hat{u})(\xi) \bar{\hat{u}}(\xi) d\xi \right| \leq \left(\|u\|_{H^s}^2 + C\delta(t) \|u\|_{G_{\sigma,s+\frac{1}{2\sigma}}^{\delta(t)}}^2 \right). \quad (5.5)$$

Plugging (5.2)-(5.5) into (5.1), we obtain

$$\frac{1}{2} \frac{d}{dt} \|u\|_{G_{\sigma,s}^{\delta(t)}}^2 \leq \left(\dot{\delta}(t) + C\delta(t)(1 + \|u\|_{G_{\sigma,s}^{\delta(t)}}^3) \right) \|u\|_{G_{\sigma,s+\frac{1}{2\sigma}}^{\delta(t)}}^2 + C \underbrace{(1 + \|u\|_{H^s})^5}_{b(t)},$$

which implies

$$\frac{1}{2} \frac{d}{dt} (1 + \|u\|_{G_{\sigma,s}^{\delta(t)}})^2 \leq \left(\dot{\delta}(t) + C\delta(t)(1 + \|u\|_{G_{\sigma,s}^{\delta(t)}}^3) \right) (1 + \|u\|_{G_{\sigma,s+\frac{1}{2\sigma}}^{\delta(t)}})^2 + Cb^5(t).$$

For any $t \in [0, T_0]$, we assume that

$$(1 + \|u\|_{G_{\sigma,s}^{\delta(t)}})^2 \leq 4f^2(t), \quad (5.6)$$

where $f^2(t) := 2(1 + \|u_0\|_{G_{\sigma,s}^{\delta_0}})^2 + 2C \int_0^t b^5(t') dt'$ and take

$$\dot{\delta}(t) = -8C\delta(t)f^3(t). \quad (5.7)$$

Noting that $B_{2,2}^s = H^s$, Lemma 2.13 guarantees that the existence of global classical solution $u \in \mathcal{C}(\mathbb{R}^+; H^s)$. Hence, it follows from (5.7) that

$$\delta(t) = \delta_0 \exp \left(-8C \int_0^t f^3(t') dt' \right) > 0, \quad \forall t \in [0, \infty), \quad (5.8)$$

where $0 < \delta_0 < 1$.

Combining (5.6) and (5.7), we get for any $[0, T_0]$

$$(1 + \|u(t)\|_{G_{\sigma,s}^{\delta(t)}})^2 \leq f^2(t). \quad (5.9)$$

Applying Theorem 3.1, we have the unique global solution on time interval $[0, T_0]$. Moreover, we get

$$(1 + \|u(T_0)\|_{G_{\sigma,s}^{\delta(T_0)}})^2 \leq f^2(T_0). \quad (5.10)$$

Using (5.6)-(5.7) and (5.10) again, then there exists a $T_1 > T_0$ such that $(1 + \|u\|_{G_{\sigma,s}^{\delta(t)}})^2 \leq 2f^2(t)$ on $[T_0, T_1]$. Therefore, we have $\sup_{t \in [T_0, T_1]} (1 + \|u\|_{G_{\sigma,s}^{\delta(t)}})^2 \leq f^2(t)$. Moreover, we conclude that

$$\sup_{t \in [0, T_1]} (1 + \|u(t)\|_{G_{\sigma,s}^{\delta(t)}})^2 \leq f^2(t).$$

Repeating the bootstrap argument, we see that the global existence of strong solution of the system (1.2). \square

Acknowledgements. This work was partially supported by NNSFC (Grant No. 12171493), FDCT (Grant No. 0091/2018/A3), the Guangdong Special Support Program (Grant No.8-2015).

References

- [1] M. S. Baouendi, C. Goulaouic. Remarks on the abstract form of nonlinear Cauchy-Kovalevsky theorems. *Comm. Partial Differential Equations*, 2(11):1151–1162, 1977.
- [2] F. Rafael, A. Barostichi, Alexandrou Himonas, and Gerson Petronilho. Autonomous Ovsyannikov theorem and applications to nonlocal evolution equations and systems. *J. Funct. Anal.*, 270(1):330–358, 2016.
- [3] A. Bressan, A. Constantin. Global conservative solutions of the Camassa-Holm equation. *Arch. Ration. Mech. Anal.*, 183(2):215–239, 2007.
- [4] A. Bressan, A. Constantin. Global dissipative solutions of the Camassa-Holm equation. *Anal. Appl. (Singap.)*, 5(1):1–27, 2007.
- [5] A. Bressan, G. Chen, Q. Zhang. Uniqueness of conservative solutions to the Camassa-Holm equation via characteristics. *Discrete Contin. Dyn. Syst.*, 35(1):25–42, 2015.
- [6] R. Camassa, D. D. Holm. An integrable shallow water equation with peaked solitons. *Phys. Rev. Lett.*, 71(11):1661–1664, 1993.
- [7] A. Constantin, J. Escher. Global existence and blow-up for a shallow water equation. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 26(2):303–328, 1998.
- [8] A. Constantin, J. Escher. Well-posedness, global existence, and blowup phenomena for a periodic quasi-linear hyperbolic equation. *Comm. Pure Appl. Math.*, 51(5):475–504, 1998.
- [9] A. Constantin. Existence of permanent and breaking waves for a shallow water equation: a geometric approach. *Ann. Inst. Fourier (Grenoble)*, 50(2):321–362, 2000.
- [10] A. Constantin. On the scattering problem for the Camassa-Holm equation. *R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci.*, 457(2008):953–970, 2001.
- [11] R. Danchin. A few remarks on the Camassa-Holm equation. *Differential Integral Equations*, 14(8):953–988, 2001.
- [12] R. Danchin. A note on well-posedness for Camassa-Holm equation. *J. Differential Equations*, 192(2):429–444, 2003.
- [13] C. Foias, R. Temam. Gevrey class regularity for the solutions of the Navier-Stokes equations. *J. Funct. Anal.*, 87(2):359–369, 1989.
- [14] Y. Guo, W. Ye, Z. Yin. Ill-posedness for the cauchy problem of the Camassa-Holm equation in $B_{\infty,1}^1(\mathbb{R})$. *327: 127-144*, 2022.
- [15] Z. Guo, X. Liu, L. Molinet, Z. Yin. Ill-posedness of the Camassa-Holm and related equations in the critical space. *J. Differential Equations*, 266(2-3):1698–1707, 2019.
- [16] A. A. Himonas, G. Misioek. Analyticity of the Cauchy problem for an integrable evolution equation. *Math. Ann.*, 327(3):575–584, 2003.
- [17] H. He, Z. Yin. The global Gevrey regularity and analyticity of a two-component shallow water system with higher-order inertia operators. *J. Differential Equations*, 267(4):2531–2559, 2019.

- [18] I. L. Freire. Wave breaking for shallow water models with time decaying solutions. *J. Differential Equations*, 269(4):3769–3793, 2020.
- [19] C. D. Levermore, M. Oliver. Analyticity of solutions for a generalized Euler equation. *J. Differential Equations*, 133(2):321–339, 1997.
- [20] J. Li, Z. Yin. Remarks on the well-posedness of Camassa-Holm type equations in Besov spaces. *J. Differential Equations*, 261(11):6125–6143, 2016.
- [21] J. Li, Y. Yu, W. Zhu. Ill-posedness for the Camassa-Holm and related equations in Besov spaces. *J. Differential Equations*, 306:403–417, 2022.
- [22] T. Lyons. Particle trajectories in extreme Stokes waves over infinite depth. *Discrete Contin. Dyn. Syst.*, 34(8):3095–3107, 2014.
- [23] Y. Li, P. Olver. Well-posedness and blow-up solutions for an integrable nonlinearly dispersive model wave equation. *J. Differential Equations*, 162(1):27–63, 2000.
- [24] W. Luo, Z. Yin. Gevrey regularity and analyticity for Camassa-Holm type systems. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 18(3):1061–1079, 2018.
- [25] Z. Meng, Z. Yin, On the Cauchy problem for a weakly dissipative Camassa-Holm equation in critical Besov spaces. *arXiv: 2206.05013*, 2022.
- [26] Z. Meng, Z. Yin. Existence and uniqueness of the globally conservative solutions for a weakly dissipative Camassa-Holm equation in time weighted $H^1(\mathbb{R})$ space. *arXiv:2206.06523*, 2022.
- [27] L. Nirenberg. An abstract form of the nonlinear Cauchy-Kowalewski theorem. *J. Differential Geometry*, 6:561–576, 1972.
- [28] L. V. Ovsjannikov. Nonlocal Cauchy problems in fluid dynamics. 137–142, 1971.
- [29] L. V. Ovsjannikov. Singular operator in the scale of Banach spaces. *Dokl. Akad. Nauk SSSR*, 163:819–822, 1965.
- [30] L. V. Ovsjannikov. A nonlinear Cauchy problem in a scale of Banach spaces. *Dokl. Akad. Nauk SSSR*, 200:789–792, 1971.
- [31] G. Rodríguez-Blanco. On the Cauchy problem for the Camassa-Holm equation. *Nonlinear Anal.*, 46(3):309–327, 2001.
- [32] W. Ye, Z. Yin, Y. Guo. A new result for the local well-posedness of the Camassa-Holm type equations in critical Besov spaces $B_{p,1}^{1+\frac{1}{p}}$, $1 \leq p < +\infty$. *arXiv: 2101.00803*, 2021.
- [33] L. Zhang, B. Liu. On the Luo-Yin results concerning Gevrey regularity and analyticity for Camassa-Holm-type systems. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 21:1741–1744, 2020.