

# Decays $h \rightarrow e_a e_b$ , $e_b \rightarrow e_a \gamma$ , and $(g - 2)_{e,\mu}$ in a 3-3-1 model with inverse seesaw neutrinos

T.T. Hong,<sup>1,\*</sup> N.H.T. Nha,<sup>2,†</sup> T. Phong Nguyen,<sup>2,‡</sup> L. T. T. Phuong,<sup>1,§</sup> and L.T. Hue<sup>¶3,4,\*\*</sup>

<sup>1</sup>*An Giang University, VNU - HCM, Ung Van Khiem Street,  
Long Xuyen, An Giang 88000, Vietnam*

<sup>2</sup>*Department of Physics, Can Tho University, 3/2 Street, Can Tho, Vietnam*

<sup>3</sup>*Subatomic Physics Research Group,  
Science and Technology Advanced Institute,  
Van Lang University, Ho Chi Minh City 70000, Vietnam*

<sup>4</sup>*Faculty of Technology, Van Lang University,  
Ho Chi Minh City 70000, Vietnam*

## Abstract

We will show that the 3-3-1 model with new heavy right handed neutrinos as  $SU(3)_L$  singlets can explain simultaneously the lepton flavor violating decays of the SM-like Higgs boson, cLFV decays  $e_b \rightarrow e_a \gamma$ , and the electron  $(g - 2)_e$  anomalies under recent experimental data. The discrepancy of  $(g - 2)_\mu$  predicted by the model under consideration and that of the standard model can reach  $10^{-9}$ . The decay rates of the standard model-like Higgs boson  $h \rightarrow \tau e, \tau \mu$  can reach the values of  $\mathcal{O}(10^{-4})$ .

PACS numbers:

---

¶ Corresponding author

\*Electronic address: tthong@agu.edu.vn

†Electronic address: nhtnha95@gmail.com

‡Electronic address: thanhphong@ctu.edu.vn

§Electronic address: ltthuong@agu.edu.vn

\*\*Electronic address: lethohue@vlu.edu.vn

## I. INTRODUCTION

The experimental evidence of neutrino oscillation [1–5] confirms that the lepton flavor number is violated in the neutral lepton sector. This is a great motivation to search for many lepton flavor violating (LFV) processes, namely the promoting ones we will focus on this work are the LFV decays of the charged leptons  $e_b \rightarrow e_a \gamma$  and the standard model-like (SM-like) Higgs boson (LFVH)  $h \rightarrow e_a^\pm e_b^\mp$ . The charged lepton flavor violating (cLFV) decays  $e_b \rightarrow e_a \gamma$  are constrained by experiments as follows [6, 7]:

$$\text{Br}(\tau \rightarrow \mu \gamma) < 4.4 \times 10^{-8}, \quad \text{Br}(\tau \rightarrow e \gamma) < 3.3 \times 10^{-8}, \quad \text{Br}(\mu \rightarrow e \gamma) < 4.2 \times 10^{-13}. \quad (1)$$

Upcoming sensitivities will be orders of  $10^{-9}$  and  $10^{-14}$  for decays  $\tau \rightarrow \mu \gamma, e \gamma$  [8, 9] and  $\mu \rightarrow e \gamma$  [10], respectively. LFVH decays have been investigated in many models beyond the standard model (BSM). On the other hand, the latest experimental constraints are:  $\text{Br}(h \rightarrow \tau^\pm \mu^\mp) < 2.5 \times 10^{-3}$  [11],  $\text{Br}(h \rightarrow \tau^\pm e^\mp) < 4.7 \times 10^{-3}$  [12], and  $\text{Br}(h \rightarrow \mu^\pm e^\mp) < 6.1 \times 10^{-3}$  [13]. The future experimental sensitivities may be  $1.4 \times 10^{-4}$ ,  $1.6 \times 10^{-4}$ , and  $1.2 \times 10^{-5}$ , respectively [14]. The small upper bounds of the cLFV branching rates prefer the explanation that they come from loop corrections relevant to LFV sources, including ones available in the neutral lepton sector. For models consisting of these necessary tree level couplings to accommodate neutrino oscillation data such as the Zee model [15], constraints on the LFV sources such as Yukawa couplings are very strict [16]. Therefore, new scalar masses must not be heavier than 300 GeV in order to explain successfully the recent  $(g - 2)$  data [16].

To explain the neutrino oscillation data, the BSMs with the general seesaw (GSS) mechanism also result in LFV decays. But the versions adding only heavy seesaw neutrinos type-I predict suppressed LFV rates that are much smaller than the upcoming experimental sensitivities [17, 18]. In contrast, the models with only new inverse seesaw (ISS) neutrinos can predict large LFV rates. In addition, LFVH rates may be large in the regions satisfying constraints of  $\text{Br}(e_b \rightarrow e_a \gamma)$  [19–22]. On the other hand, LFVH rates may be smaller when other constraints are considered [23, 24]. In the supersymmetric (SUSY) versions of these models with new LFV sources from superparticles, LFVH rates may reach large order of  $\mathcal{O}(10^{-5})$  [17, 25–33]. LFVH decays were also mentioned in many other non-SUSY extensions of the SM [19, 34–64].

Unless some specific condition of the appearance of very light new bosons, the above cLFV constraints will result in small new one-loop contributions to anomalous magnetic moments (AMM) of charged leptons  $(g - 2)_{e_a}/2 \equiv a_{e_a}$ , in contrast with recent experimental data. Namely, the  $4.2\sigma$  deviation between standard model (SM) prediction based on dispersion calculation [65] and experiments [66, 67] for muon is

$$\Delta a_\mu^{\text{NP}} \equiv a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = (2.51 \pm 0.59) \times 10^{-9}. \quad (2)$$

This result is a bit inconsistent with the lattice QCD approach used to calculate the SM prediction, giving a value closer to the experimental value [68]. Regarding the electron anomaly, the  $1.6\sigma$  discrepancy between SM and experiment was reported [6]

$$\Delta a_e^{\text{NP}} \equiv a_e^{\text{exp}} - a_e^{\text{SM}} = (4.8 \pm 3.0) \times 10^{-13}. \quad (3)$$

The recent studies of cLFV decays in the regions satisfying the AMM data were done in some specific models such as SUSY with largest  $\text{Br}(h \rightarrow \tau\mu) \sim \mathcal{O}(10^{-4})$  [69]. Other BSM with the presence of leptoquarks can explain the large  $\Delta a_\mu^{\text{NP}} \sim \mathcal{O}(10^{-9})$  [70].

A recent work discussing an extension of the 3-3-1 with right-handed neutrinos [71–73], named the 3-3-1 model with inverse seesaw neutrinos (331ISS) [74], with the aim of giving an explanation of both the  $(g - 2)_\mu$  data and the neutrino oscillation data through the ISS mechanism. The model needs new  $SU(3)_L$  gauge singlets including three neutral leptons  $X_{aR}$  and a new singly charged Higgs boson  $h^\pm$  to accommodate all experimental data of neutrino oscillation, cLFV bounds (1) and  $\Delta a_\mu$  in  $1\sigma$  deviation given in Eq. (2). Although cLFV and/or LFBV decays were investigated previously with promoting predictions for the 331ISS [75–78], the AMM data was not included. Our aim in this work is filling this gap. We note that other 3-3-1 models [79–82] constructed previously can accommodate the  $(g - 2)_\mu$  data only when they are extended such as adding new vector-like fermions, or/and scalars [83–87]. But none of them paid attention to the correlations between LFBV decays and  $(g - 2)_{e_a}$  anomalies.

Our paper is organized as follows. In Sec. II we discuss the necessary ingredients of a 331ISS model for studying LFBVDs and how the ISS mechanism works to generate active neutrino masses and mixing consistent with current experimental data. In Sec. III we present all couplings needed to determine the one-loop contributions to the LFBV decay amplitudes of the SM-like Higgs boson, cLFV decays, and  $(g - 2)_{e_a}$ . In section IV, we provide detailed

numerical illustrations and discussions. Section V contains our conclusions. Finally, the Appendix lists all of the analytic formulas expressing one-loop contributions to LFBVH decay amplitudes calculated in the unitary gauge.

## II. THE 331ISS MODEL FOR TREE-LEVEL NEUTRINO MASSES

### A. Particle content and lepton masses

The particle content of the 331ISS model will be summarized in this section. We will ignore the quark sector, which was discussed previously, but irrelevant in our work [74, 76, 88]. We also ignore many detailed calculations presented in Ref. [74]. The electric charge operator defined by the gauge group  $SU(3)_L \times U(1)_X$  is  $Q = T_3 - \frac{1}{\sqrt{3}}T_8 + X$ , where  $T_{3,8}$  are diagonal  $SU(3)_L$  generators. Each lepton family consists of a  $SU(3)_L$  triplet  $L_{aL} = (\nu_a, e_a, N_a)_L^T \sim (3, -\frac{1}{3})$  and a right-handed charged lepton  $e_{aR} \sim (1, -1)$  with  $a = 1, 2, 3$ . The 331ISS model contains three neutral lepton  $X_{aR} \sim (1, 0)$ ,  $a = 1, 2, 3$ , and a singly charged Higgs boson  $\sigma^\pm \sim (1, 1)$ . The three Higgs triplets are  $\rho = (\rho_1^+, \rho^0, \rho_2^+)^T \sim (3, \frac{2}{3})$ ,  $\eta = (\eta_1^0, \eta^-, \eta_2^0)^T \sim (3, -\frac{1}{3})$ , and  $\chi = (\chi_1^0, \chi^-, \chi_2^0)^T \sim (3, -\frac{1}{3})$ . The vacuum expectation values (vev) for generating all tree-level quark masses and leptons are  $\langle \rho \rangle = (0, \frac{v_1}{\sqrt{2}}, 0)^T$ ,  $\langle \eta \rangle = (\frac{v_2}{\sqrt{2}}, 0, 0)^T$  and  $\langle \chi \rangle = (0, 0, \frac{w}{\sqrt{2}})^T$ . Two neutral Higgs components having zero vevs because of their non-zero generalized lepton numbers [74] corresponding to a new  $U(1)_\mathcal{L}$  [89].

In the 331ISS, nine gauge bosons get masses through the covariant kinetic Lagrangian of the Higgs boson triplets,  $\mathcal{L}^H = \sum_{H=\chi, \eta, \rho} (D_\mu H)^\dagger (D^\mu H)$ , where  $D_\mu = \partial_\mu - igW_\mu^a T^a - ig_X T^9 X X_\mu$ ,  $a = 1, 2, \dots, 8$ , and  $T^9 \equiv \frac{I_3}{\sqrt{6}}$  and  $\frac{1}{\sqrt{6}}$  for (anti)triplets and singlets [90]. There are two pairs of singly charged gauge bosons, denoted as  $W^\pm$  and  $Y^\pm$ , defined as

$$W_\mu^\pm = \frac{W_\mu^1 \mp iW_\mu^2}{\sqrt{2}}, \quad Y_\mu^\pm = \frac{W_\mu^6 \pm iW_\mu^7}{\sqrt{2}}, \quad (4)$$

with the respective masses  $m_W^2 = \frac{g^2}{4}(v_1^2 + v_2^2)$  and  $m_Y^2 = \frac{g^2}{4}(w^2 + v_1^2)$ . The breaking pattern of the model is  $SU(3)_L \times U(1)_X \rightarrow SU(2)_L \times U(1)_Y \rightarrow U(1)_Q$ , leading to the matching condition that  $W^\pm$  are the SM gauge boson, implying that

$$v_1^2 + v_2^2 \equiv v^2 = (246\text{GeV})^2, \quad \frac{g_X}{g} = \frac{3\sqrt{2}s_W}{\sqrt{3 - 4s_W^2}}, \quad g_{SW} = e, \quad (5)$$

where  $e$  and  $s_W$  are, respectively, the electric charge and sine of the Weinberg angle. Similarly to the 2HDM, the following parameters will be used,

$$t_\beta \equiv \tan \beta = \frac{v_2}{v_1}, \quad (6)$$

which leads to  $v_1 = v c_\beta$  and  $v_2 = v s_\beta$ .

The Yukawa Lagrangian generating lepton masses is:

$$\begin{aligned} \mathcal{L}_l^Y = & -h_{ab}^e \overline{L_a} \rho e_{bR} + h_{ab}^\nu \epsilon^{ijk} \overline{(L_a)_i} (L_b)_j \rho_k^* - y_{ba}^\chi \overline{X_{bR}} \chi^\dagger L_a - \frac{1}{2} (\mu_X)_{ab} \overline{X_{aR}} (X_{bR})^c \\ & - Y_{ab}^\sigma \overline{(X_{aR})^c} e_{bR} \sigma^+ + \text{H.c.}, \end{aligned} \quad (7)$$

where  $a, b = 1, 2, 3$ . The first term generates charged lepton masses as  $m_{e_a} \equiv \frac{h_{ab}^e v_1}{\sqrt{2}} \delta_{ab}$ , with the assumption that the flavor states are also physical.

In the basis  $n'_L = (\nu_L, N_L, (X_R)^c)^T$ , Lagrangian (7) generates a neutrino mass term written in terms of the total  $9 \times 9$  mass matrix consisting of nine  $3 \times 3$  sub-matrices [76], namely

$$-\mathcal{L}_{\text{mass}}^\nu = \frac{1}{2} \overline{(n'_L)^c} \mathcal{M}^\nu n'_L + \text{H.c.}, \quad \text{where} \quad \mathcal{M}^\nu = \begin{pmatrix} \mathcal{O}_3 & m_D^T & \mathcal{O}_3 \\ m_D & \mathcal{O}_3 & M_R^T \\ \mathcal{O}_3 & M_R & \mu_X \end{pmatrix}, \quad (8)$$

where  $(n'_L)^c = ((\nu_L)^c, (N_L)^c, X_R)^T$ ,  $(M_R)_{ab} \equiv y_{ab}^\chi \frac{w}{\sqrt{2}}$ , and  $(m_D^T)_{ab} = -(m_D)_{ab} \equiv \sqrt{2} h_{ab}^\nu v_1$  with  $a, b = 1, 2, 3$ . The matrix  $\mu_X$  in Eq. (7) is symmetric, which will be considered as a diagonal matrix without loss of generality.

The mass matrix  $\mathcal{M}^\nu$  is diagonalized by a  $9 \times 9$  unitary matrix  $U^\nu$ ,

$$U^{\nu T} \mathcal{M}^\nu U^\nu = \hat{M}^\nu = \text{diag}(m_{n_1}, m_{n_2}, \dots, m_{n_9}) = \text{diag}(\hat{m}_\nu, \hat{M}_N), \quad (9)$$

where  $m_{n_i}$  ( $i = 1, 2, \dots, 9$ ) are masses corresponding to the physical states  $n_{iL}$ . The two mass matrices  $\hat{m}_\nu = \text{diag}(m_{n_1}, m_{n_2}, m_{n_3})$  and  $\hat{M}_N = \text{diag}(m_{n_4}, m_{n_5}, \dots, m_{n_9})$  consist of the masses of the active  $n_{aL}$  ( $a = 1, 2, 3$ ) and extra neutrinos  $n_{iL}$  ( $i = 1, 2, \dots, 6$ ), respectively. The following approximation solution of  $U^\nu$  is valid for any specific seesaw mechanisms,

$$U^\nu = \Omega \begin{pmatrix} U_{\text{PMNS}} & \mathcal{O}_{3 \times 6} \\ \mathcal{O}_{6 \times 3} & V \end{pmatrix}, \quad \Omega \simeq \begin{pmatrix} I_3 - \frac{1}{2} R R^\dagger & R \\ -R^\dagger & I_6 - \frac{1}{2} R^\dagger R \end{pmatrix}, \quad (10)$$

where  $R, V$  are  $3 \times 6, 3 \times 6$  matrices, respectively. All entries of  $R$  must satisfy  $|R_{aI}| \ll 1$ , so that all ISS relations can be derived perturbatively.

The relations between the flavor and mass eigenstates are

$$n'_L = U^\nu n_L, \quad \text{and } (n'_L)^c = U^{\nu*} (n_L)^c \equiv U^{\nu*} n_R, \quad (11)$$

where  $n_L \equiv (n_{1L}, n_{2L}, \dots, n_{9L})^T$ , and the Majorana states are  $n_i = (n_{iL}, n_{iR})^T$ .

The ISS relations are

$$R_2^* = m_D^T M_R^{-1}, \quad R_1^* = -R_2^* \mu_X (M_R^T)^{-1} \simeq \mathcal{O}_3, \quad (12)$$

$$m_\nu = R_2^* \mu_X R_2^\dagger = U_{\text{PMNS}}^* \hat{m}_\nu U_{\text{PMNS}}^\dagger = m_D^T M_R^{-1} \mu_X (M_R^{-1})^T m_D, \quad (13)$$

$$V^* \hat{M}_N V^\dagger = M_N + \frac{1}{2} M_N R^\dagger R + \frac{1}{2} R^T R^* M_N. \quad (14)$$

From experimental data of  $m_\nu$ , we can determine all independent parameters in  $m_D$  and three entries of  $M^{-1} \equiv M_R^{-1} \mu_X (M_R^{-1})^T$  [76, 88]. Namely the Dirac mass matrix has antisymmetric form as follows

$$m_D = z e^{i\alpha_{23}} \times \tilde{m}_D, \quad (15)$$

where  $\alpha_{23} \equiv \arg[h_{32}^\nu]$ ,  $\tilde{m}_D$  is an anti-symmetric matrix with  $(\tilde{m}_D)_{23} = 1$ , and

$$z = \sqrt{2} v_1 |h_{32}^\nu| = \sqrt{2} v_1 |h_{23}^\nu| \equiv z_0 c_\beta \quad (16)$$

is a positive and real parameter. Eq. (13) gives  $(m_\nu)_{ij} = [m_D^T M^{-1} m_D]_{ij}$  for all  $i, j = 1, 2, 3$ , leading to six independent equations. Solving three of them with  $i \neq j$ , the non-diagonal entries of  $M^{-1}$  are functions of  $M_{ii}^{-1}$  and  $x_{12,13}$ . Inserting these functions into the three remaining relations with  $i = j$ , we obtain

$$(\tilde{m}_D)_{32} = \frac{m_\nu)_{13}^2 - (m_\nu)_{11} (m_\nu)_{33}}{(m_\nu)_{13} (m_\nu)_{23} - (m_\nu)_{12} (m_\nu)_{33}}, \quad (\tilde{m}_D)_{21} = \frac{(m_\nu)_{12} (m_\nu)_{13} - (m_\nu)_{11} (m_\nu)_{23}}{(m_\nu)_{13} (m_\nu)_{23} - (m_\nu)_{12} (m_\nu)_{33}}, \quad (17)$$

and  $\text{Det}[m_\nu] = 0$ . From  $M^{-1} = M_R^{-1} \mu_X (M_R^{-1})^T$  we derive that three parameters of the matrix  $\mu_X$  as certain but lengthy functions of  $(z e^{i\alpha_{23}})$ , all entries of  $M_R$  and  $m_\nu$ . While  $m_\nu$  are fixed by experiments, all entries of  $M_R$  are free parameters. We will fix  $\alpha_{23} = 0$ , because it is absorbed into the  $\mu_X$ .

In the limit that  $|R_2| \ll 1$ , the heavy neutrino masses can be determined approximately based on Eq. (14), namely

$$V^* \hat{M}_N V^\dagger \simeq M_N. \quad (18)$$

We define the reduced matrix  $M_R \equiv z\tilde{M}_R$ ,  $(\tilde{M}_R)_{ij} \equiv k_{ij}$ , provided that  $R_2^* = -\tilde{m}_D/\tilde{M}_R$ . The matrix  $M_R$  is always diagonalized by two unitary transformations  $V_{L,R}$  [91]:

$$V_L^T M_R V_R = z \times \hat{k} = z \times \text{diag}(\hat{k}_1, \hat{k}_2, \hat{k}_3), \quad (19)$$

where all  $\hat{k}_{1,2,3}$  are always positive and  $\hat{k}_a \gg 1$  so that all ISS relations are valid. Therefore,  $M_R$  is expressed in terms of  $\hat{k}$  and  $V_{L,R}$ . Then the matrix  $V$  in Eq. (14) can be found approximately as follows

$$V = \frac{1}{\sqrt{2}} \begin{pmatrix} V_R & iV_R \\ V_L & -iV_L \end{pmatrix} \rightarrow V^T M_N V = z \times \begin{pmatrix} \hat{k} & \mathcal{O}_{3 \times 3} \\ \mathcal{O}_{3 \times 3} & \hat{k} \end{pmatrix}. \quad (20)$$

As a consequence, for any qualitative estimations, we will use the approximation that heavy neutrinos masses are  $m_{n_{a+3}} = m_{n_{a+6}} \simeq z\hat{k}_a$  with  $a=1,2,3$ ;  $R_1 \simeq \mathcal{O}_3$ ; and

$$U^\nu \simeq \begin{pmatrix} \left( I_3 - \frac{1}{2} R_2 R_2^\dagger \right) U_{\text{PMNS}} & \frac{1}{\sqrt{2}} R_2 V_L & \frac{-i}{\sqrt{2}} R_2 V_L \\ \mathcal{O}_3 & \frac{V_R}{\sqrt{2}} & \frac{iV_R}{\sqrt{2}} \\ -R_2^\dagger U_{\text{PMNS}} & \left( I_3 - \frac{R_2^\dagger R_2}{2} \right) \frac{V_R}{\sqrt{2}} & \left( I_3 - \frac{R_2^\dagger R_2}{2} \right) \frac{-iV_R}{\sqrt{2}} \end{pmatrix}. \quad (21)$$

We have checked and confirm that the above approximations give numerical results consistent with those discussed in Ref. [74]. Therefore, these approximate formulas will be used in this work.  $m_\nu$  is chosen as the input with  $3\sigma$  neutrino oscillation data to fix  $\tilde{m}_D$ . The free parameters  $z_0$  and  $\hat{k}_{1,2,3}$ ,  $V_R$  will be scanned in the valid ranges to construct the total neutrino mixing matrix  $U^\nu$  defined in Eq. (21). Because  $V_L$  satisfies that

$$R_2 V_L = \tilde{m}_D^\dagger V_R^* \hat{k}^{-1}, \quad R_2 R_2^\dagger = \tilde{m}_D^\dagger V_R \hat{k}^{-2} \tilde{m}_D, \quad (22)$$

which do not depend explicitly on  $V_L$ . Hence it affects weakly on all relevant processes. We will fix  $V_L = I_3$  from now on.

Lagrangian for quark masses were discussed previously [89]. Here, we just remind that the Yukawa couplings of the top quark must satisfy the perturbative limit  $h_{33}^u < \sqrt{4\pi}$ , leading to a lower bound of  $v_2$ :  $v_2 > \frac{\sqrt{2}m_t}{\sqrt{4\pi}}$ . Combined with the relations in Eqs. (5) and (6), the lower bound of  $t_\beta$  is  $t_\beta \geq 0.3$ . The upper bound of  $t_\beta$  can be derived from the tau mass,  $m_\tau = h_{33}^3 \times v c_\beta \sqrt{2} \rightarrow h_{33}^3 = m_\tau \sqrt{2} / (v c_\beta) < \sqrt{4\pi}$ , leading to a rather weak upper bound  $t_\beta = \sqrt{1/c_\beta^2 - 1} \leq 346$ .

## B. Higgs bosons

The Higgs potential used here respect the new lepton number defined in ref. [89], namely

$$\begin{aligned}
V_h = & \sum_S \left[ \mu_S^2 S^\dagger S + \lambda_S (S^\dagger S)^2 \right] + \lambda_{12}(\eta^\dagger \eta)(\rho^\dagger \rho) + \lambda_{13}(\eta^\dagger \eta)(\chi^\dagger \chi) + \lambda_{23}(\rho^\dagger \rho)(\chi^\dagger \chi) \\
& + \tilde{\lambda}_{12}(\eta^\dagger \rho)(\rho^\dagger \eta) + \tilde{\lambda}_{13}(\eta^\dagger \chi)(\chi^\dagger \eta) + \tilde{\lambda}_{23}(\rho^\dagger \chi)(\chi^\dagger \rho) + \sqrt{2}\omega f (\epsilon_{ijk} \eta^i \rho^j \chi^k + \text{h.c.}) \\
& + \sigma^+ \sigma^- \left[ \mu_\sigma^2 + \sum_S \lambda_S^\sigma S^\dagger S \right] + [f_\eta(\rho^\dagger \eta)\sigma^+ + f_\chi(\rho^\dagger \chi)\sigma^+ + \text{h.c.}], \tag{23}
\end{aligned}$$

where  $f$  is a dimensionless parameter,  $f_{\eta,\chi}$  are mass dimensional,  $S = \eta, \rho, \chi$ . These three trilinear couplings break softly the general lepton number  $\mathcal{L}$ . For simplicity, we fix  $f_\chi = 0$  by applying a suitable discrete symmetry. The last line in Eq. (23) contains all additional terms couplings with new charged Higgs singlets compared with Higgs potential considered in previous works [74]. They do not affect the squared mass matrices of both neutral CP-odd and -even Higgs bosons. The minimum conditions of the Higgs potential as well as the identification of the SM-like Higgs were discussed in detailed previously [30, 92], hence we just list the necessary results here. The model contains three pairs of singly charged Higgs bosons  $h_{1,2,3}^\pm$  and two Goldstone bosons  $G_{W,Y}^\pm$  of the singly charged gauge bosons  $W^\pm$  and  $Y^\pm$ , respectively.  $m_{H_1^\pm}^2 = \left( \frac{\tilde{\lambda}_{12}v^2}{2} + \frac{fw^2}{s_\beta c_\beta} \right)$ ,  $m_{h_3^\pm}^2 = (v^2 c_\beta^2 + w^2) \left( \frac{\tilde{\lambda}_{23}}{2} + ft_\beta \right)$ , and  $m_{G_W^\pm}^2 = m_{G_Y^\pm}^2 = 0$ . The relations between the original and mass eigenstates of the charged Higgs bosons are [92]

$$\begin{pmatrix} \eta^\pm \\ \rho_1^\pm \\ \sigma^\pm \end{pmatrix} = \begin{pmatrix} -s_\beta & c_\alpha c_\beta & s_\alpha c_\beta \\ c_\beta & c_\alpha s_\beta & s_\alpha s_\beta \\ 0 & -s_\alpha & c_\alpha \end{pmatrix} \begin{pmatrix} G_W^\pm \\ h_1^\pm \\ h_2^\pm \end{pmatrix}, \quad \begin{pmatrix} \rho_2^\pm \\ \chi^\pm \end{pmatrix} = \begin{pmatrix} -s_\theta & c_\theta \\ c_\theta & s_\theta \end{pmatrix} \begin{pmatrix} G_Y^\pm \\ h_3^\pm \end{pmatrix}, \tag{24}$$

where  $t_\theta = v_1/w$ , and

$$\begin{aligned}
f &= \frac{c_\beta s_\beta \left( 2c_\alpha^2 m_{h_1^\pm}^2 + 2s_\alpha^2 m_{h_2^\pm}^2 - \tilde{\lambda}_{12}v^2 \right)}{2\omega^2}, \quad f_\eta = \frac{\sqrt{2}c_\alpha s_\alpha (m_{h_2^\pm}^2 - m_{h_1^\pm}^2)}{v}, \\
\mu_\sigma^2 &= \frac{1}{2} \left( 2c_\alpha^2 m_{h_2^\pm}^2 - v^2 (c_\beta^2 \lambda_2^\sigma + s_\beta^2 \lambda_1^\sigma) + 2s_\alpha^2 m_{h_1^\pm}^2 - \lambda_3^\sigma \omega^2 \right). \tag{25}
\end{aligned}$$

These result are consistent with ref. [90, 92, 93] in the limits of  $s_\alpha = 0, \pm 1$ . The results given in Eqs. (24) and (25) obtained by solving the following  $3 \times 3$  squared mass matrix in



the basis  $(\eta^\pm, \rho_1^\pm, \sigma^\pm)$ :

$$\mathcal{M}_c^2 = \begin{pmatrix} \frac{f\omega^2}{t_\beta} + \frac{1}{2}c_\beta^2\tilde{\lambda}_{12}v^2 & f\omega^2 + \frac{1}{2}c_\beta\tilde{\lambda}_{12}s_\beta v^2 & \frac{c_\beta f_\eta v}{\sqrt{2}} \\ f\omega^2 + \frac{1}{2}c_\beta\tilde{\lambda}_{12}s_\beta v^2 & ft_\beta\omega^2 + \frac{1}{2}\tilde{\lambda}_{12}s_\beta^2 v^2 & \frac{f_\eta s_\beta v}{\sqrt{2}} \\ \frac{c_\beta f_\eta v}{\sqrt{2}} & \frac{f_\eta s_\beta v}{\sqrt{2}} & \frac{v^2}{2}(c_\beta^2\lambda_2^\sigma + s_\beta^2\lambda_1^\sigma) + \frac{\lambda_3^\sigma\omega^2}{2} + \mu_\sigma^2 \end{pmatrix}. \quad (26)$$

We will find out that the Higgs masses  $m_{h_{1,2}^\pm}$  and the mixing angle  $\alpha$  are functions of the Higgs parameters in the Higgs potential.

The model contains five CP-odd neutral scalar components included in the five neutral Higgs bosons  $\eta_1^0 = (v_2 + R_1 + iI_1)/\sqrt{2}$ ,  $\rho^0 = (v_1 + R_2 + iI_2)/\sqrt{2}$ ,  $\chi_2^0 = (\omega + R_3 + iI_3)/\sqrt{2}$ ,  $\eta_2^0 = (R_4 + iI_4)/\sqrt{2}$ , and  $\chi_1^0 = (R_5 + iI_5)/\sqrt{2}$ . Three of them are Goldstones bosons of the neutral gauge bosons  $Z, Z'$  and  $X^0$ . The two remaining are physical states with masses

$$m_{a_1}^2 = (s_\beta^2 v^2 + \omega^2) \left( ft_\beta^{-1} + \frac{1}{2}\tilde{\lambda}_{13} \right), \quad m_{a_2}^2 = f \left( \frac{\omega^2}{c_\beta s_\beta} + c_\beta s_\beta v^2 \right). \quad (27)$$

As a consequence, the parameter  $f$  must satisfies  $f > 0$ .

Considering the CP-even scalars, there are two sub-matrices  $2 \times 2$  and  $3 \times 3$  for masses of these Higgs bosons in two bases  $(\eta_2^0, \chi_1^0)$  and  $(\eta_1^0, \rho_1^0, \chi_1^0)$ , namely

$$M_{0,3}^2 = \begin{pmatrix} \frac{c_\beta f \omega^2}{s_\beta} + 2s_\beta^2 \lambda_1 v^2 & c_\beta s_\beta \lambda_{12} v^2 - \omega^2 f & \omega(s_\beta \lambda_{13} - c_\beta f)v \\ c_\beta s_\beta \lambda_{12} v^2 - \omega^2 f & \frac{s_\beta f \omega^2}{c_\beta} + 2c_\beta^2 \lambda_2 v^2 & \omega(c_\beta \lambda_{23} - s_\beta f)v \\ \omega(s_\beta \lambda_{13} - c_\beta f)v & \omega(c_\beta \lambda_{23} - s_\beta f)v & 2\lambda_3 \omega^2 + c_\beta s_\beta f v^2 \end{pmatrix},$$

$$M_{0,2}^2 = \begin{pmatrix} \frac{1}{2}\omega^2 \left( \tilde{\lambda}_{13} + \frac{2c_\beta f}{s_\beta} \right) & \frac{1}{2}\omega(\tilde{\lambda}_{13}s_\beta + 2c_\beta f)v \\ \frac{1}{2}\omega(\tilde{\lambda}_{13}s_\beta + 2c_\beta f)v & \frac{1}{2}s_\beta(\tilde{\lambda}_{13}s_\beta + 2c_\beta f)v^2 \end{pmatrix}. \quad (28)$$

The matrix  $M_{0,2}^2$  has one zero value and  $m_{h_4}^2 = \left( \frac{f}{t_\beta} + \frac{\tilde{\lambda}_{13}}{2} \right) (s_\beta^2 v^2 + \omega^2)$  corresponding to one goldstone boson of  $X^0$  and a heavy neutral Higgs boson  $h_4^0$  with mass at the  $SU(3)_L$  breaking scale. On the other hand, we see that  $\text{Det}[M_{0,3}^2] \neq 0$  but  $\text{Det}[M_{0,3}^2]|_{v=0} = 0$ , which implies that there is at least one Higgs boson mass at the electroweak scale that can be identified with the SM-like Higgs boson. In particular, it can be proved that

$$C_1^h M_{0,3}^2 C_1^{hT}|_{v=0} = \text{diag} \left( 0, 2\lambda_3 \omega^2, f\omega^2/(s_\beta c_\beta) \right), \quad C_1^h = \begin{pmatrix} s_\beta & c_\beta & 0 \\ -c_\beta & s_\beta & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (29)$$

and  $C_1^h M_{0,3}^2 C_1^{hT} \equiv M_{0,3}'^2$  satisfying:

$$(M_{0,3}'^2)_{11} = 2v^2 (c_\beta^4 \lambda_2 + c_\beta^2 \lambda_{12} s_\beta^2 + \lambda_1 s_\beta^4),$$

$$\begin{aligned}
(M'_{0,3})_{22} &= 2c_\beta^2 s_\beta^2 v^2 (\lambda_1 - \lambda_{12} + \lambda_2) + \frac{f\omega^2}{c_\beta s_\beta}, \\
(M'_{0,3})_{33} &= fc_\beta s_\beta v^2 + 2\lambda_3 \omega^2, \\
(M'_{0,3})_{12} &= (M'_{0,3})_{21} = c_\beta s_\beta v^2 (s_\beta^2 (\lambda_{12} - 2\lambda_1) - c_\beta^2 (\lambda_{12} - 2\lambda_2)), \\
(M'_{0,3})_{13} &= (M'_{0,3})_{31} = v\omega (-2fc_\beta s_\beta + c_\beta^2 \lambda_{23} + \lambda_{13} s_\beta^2), \\
(M'_{0,3})_{32} &= (M'_{0,3})_{23} = v\omega (fc_\beta^2 - fs_\beta^2 + c_\beta s_\beta (\lambda_{23} - \lambda_{13})). \tag{30}
\end{aligned}$$

Therefore, there is a unitary transformation  $C_2^h$  with  $(C_2^h)_{ij} \sim \mathcal{O}(v/w)$  ( $i \neq j$ ) so that  $C_2^h M'_{0,3} C_2^{hT} = \text{diag} (m_{h_1^0}^2, m_{h_2^0}^2, m_{h_3^0}^2)$  and  $m_{h_1^0}^2 \sim \mathcal{O}(v^2)$  [76, 94, 95]. Hence  $h_1^0$  is identified with the SM-like Higgs boson found at the LHC, namely  $h_1^0 \equiv h$ . For simplicity, we will fix  $C_2^h = I_3$  in this work, and use the relations  $(\eta_1^0, \rho_1^0, \chi_1^0) = C_1^{hT} (h_1^0, h_2^0, h_3^0)$  in our numerical investigation, where only  $\eta_1^0 \sim R_1$  and  $\rho_1^0 \sim R_2$  give contributions to  $h_1^0$ , namely

$$R_1 = s_\beta h_1^0 - c_\beta h_2^0, \quad R_2 = c_\beta h_1^0 + s_\beta h_2^0. \tag{31}$$

This assumption leads to a consequence that the  $m_{h_1^0}$  is independent with the Higgs self-couplings relating with one-loop decays  $h_1^0 \rightarrow e_a e_b$ , as it will be seen as follows:

$$-\mathcal{L}_h = V_h = \sum_{i,j=1}^3 -g_{hij} h_1^0 h_i^+ h_j^- + \dots, \tag{32}$$

where non-zero  $g_{h_1^0 ij} = g_{hji}$  are

$$\begin{aligned}
g_{h11} &= -vc_\alpha^2 \left[ \left( 2c_\beta^2 s_\beta^2 (\lambda_1 - \lambda_{12} + \lambda_2) + \lambda_{12} + \tilde{\lambda}_{12} \right) + t_\alpha^2 (c_\beta^2 \lambda_2^\sigma + s_\beta^2 \lambda_1^\sigma) + \frac{2s_\alpha^2 (m_{h_1^\pm}^2 - m_{h_2^\pm}^2)}{v^2} \right], \\
g_{h22} &= -vc_\alpha^2 \left[ t_\alpha^2 \left( 2c_\beta^2 s_\beta^2 (\lambda_1 - \lambda_{12} + \lambda_2) + \lambda_{12} + \tilde{\lambda}_{12} \right) + (c_\beta^2 \lambda_2^\sigma + s_\beta^2 \lambda_1^\sigma) - \frac{2s_\alpha^2 (m_{h_1^\pm}^2 - m_{h_2^\pm}^2)}{v^2} \right], \\
g_{h12} &= -c_\alpha s_\alpha v \left[ 2s_\beta^2 c_\beta^2 (\lambda_1 - \lambda_{12} + \lambda_2) - s_\beta^2 \lambda_1^\sigma - c_\beta^2 \lambda_2^\sigma + \lambda_{12} + \tilde{\lambda}_{12} - \frac{(c_\alpha^2 - s_\alpha^2)(m_{h_1^\pm}^2 - m_{h_2^\pm}^2)}{v^2} \right], \\
g_{h33} &= -v \left[ c_\beta^2 \left( 2c_\theta^2 \lambda_2 + s_\theta^2 (\lambda_{23} + \tilde{\lambda}_{23}) \right) + s_\beta^2 (c_\theta^2 \lambda_{12} + \lambda_{13} s_\theta^2) + c_\beta c_\theta^2 (2fs_\beta + c_\beta \tilde{\lambda}_{23}) \right]. \tag{33}
\end{aligned}$$

In the next section, we will derive all of the remaining couplings giving one-loop contributions of decays mentioned in this work.

### III. COUPLINGS AND ANALYTIC FORMULAS

#### A. Decays $e_b \rightarrow e_a \gamma$ and $(g-2)_{e_a}$

Couplings of charged gauge bosons giving one-loop contributions to LFV amplitudes are:

$$L_{V\pm ff} = \frac{g}{\sqrt{2}} \sum_{a=1}^3 \sum_{i=1}^9 \bar{n}_i \gamma^\mu P_L e_a [U_{ai}^{\nu*} W_\mu^+ + U_{(a+3)i}^{\nu*} Y_\mu^+] + \text{h.c.}, \quad (34)$$

All steps for calculation to derive these couplings were presented in ref. [76]. From now on, we always choose that  $m_{e_b} > m_{e_a}$ , equivalently  $b > a = 1, 2, 3$ , to define the decays  $e_b \rightarrow e_a \gamma$ . One-loop form factors from charged gauge bosons are [96]:

$$\begin{aligned} c_{(ab)R}(W) &= \frac{eg^2}{32\pi^2 m_W^2} \sum_{i=1}^9 U_{ai}^{\nu*} U_{bi}^{\nu*} \tilde{F}_V(x_{W,i}), \\ c_{(ab)R}(Y) &= \frac{eg^2}{32\pi^2 m_Y^2} \sum_{i=1}^9 U_{(a+3)i}^{\nu} U_{(b+3)i}^{\nu*} \tilde{F}_V(x_{Y,i}), \end{aligned} \quad (35)$$

where  $x_{v,i} = m_{n_i}^2 / m_v^2$ ;  $v = W, Y$ ;

$$\tilde{F}_V(x) = -\frac{10 - 43x + 78x^2 - 49x^3 + 4x^4 + 18x^3 \ln(x)}{24(x-1)^4}; \quad (36)$$

$e = \sqrt{4\pi\alpha_{\text{em}}}$  being the electromagnetic coupling constant; and  $g = e/s_W$ .

The Yukawa couplings of charged Higgs bosons with leptons are defined by

$$\mathcal{L}^{\text{tnh}^\pm} = -\frac{g}{\sqrt{2}m_W} \sum_{k=1}^3 \sum_{a=1}^3 \sum_{i=1}^9 h_k^+ \bar{n}_i \left( \lambda_{ai}^{L,k} P_L + \lambda_{ai}^{R,k} P_R \right) e_a + \text{h.c.}, \quad (37)$$

where

$$\begin{aligned} \lambda_{ai}^{R,1} &= m_{e_a} c_\alpha t_\beta U_{ai}^{\nu*} - \sum_{c=1}^3 \frac{v Y_{ca}^\sigma s_\alpha}{\sqrt{2}} U_{(c+6)i}^{\nu*}, & \lambda_{ai}^{L,1} &= c_\alpha s_\beta z_0 e^{i\alpha_{23}} \sum_{c=1}^3 (\tilde{m}_D)_{ac} U_{(c+3)i}^\nu, \\ \lambda_{ai}^{R,2} &= m_{e_a} s_\alpha t_\beta U_{ai}^{\nu*} + \sum_{c=1}^3 \frac{v Y_{ca}^\sigma c_\alpha}{\sqrt{2}} U_{(c+6)i}^{\nu*}, & \lambda_{ai}^{L,2} &= s_\alpha s_\beta z_0 e^{i\alpha_{23}} \sum_{c=1}^3 (\tilde{m}_D)_{ac} U_{(c+3)i}^\nu, \\ \lambda_{ai}^{R,3} &= \frac{m_{e_a} c_\theta U_{(a+3)i}^{\nu*}}{c_\beta}, & \lambda_{ai}^{L,3} &= c_\theta z_0 \sum_{c=1}^3 \left[ -e^{i\alpha_{23}} (\tilde{m}_D)_{ac} U_{ci}^\nu + t_\theta^2 (\tilde{M}_R^T)_{ac} U_{(c+6)i}^\nu \right]. \end{aligned} \quad (38)$$

The one-loop form factors are [96]:

$$c_{(ab)R}(h_k^\pm) = \frac{eg^2}{32\pi^2 m_W^2 m_{e_b} m_{h_k^\pm}^2} \sum_{i=1}^9 \left[ \lambda_{ai}^{L,k*} \lambda_{bi}^{R,k} m_{n_i} F_H(x_{k,i}) \right]$$

$$+ \left( m_{e_b} \lambda_{ai}^{L,k*} \lambda_{bi}^{L,k} + m_{e_a} \lambda_{ai}^{R,k*} \lambda_{bi}^{R,k} \right) \tilde{F}_H(x_{k,i}) \Big], \quad (39)$$

where  $b \geq a$ ,  $x_{k,i} = m_{n_i}^2/m_{h_k^\pm}^2$ , and the one-loop functions  $F_H(x)$  and  $\tilde{F}_H(x)$  are

$$F_H(x) = -\frac{1-x^2+2x \ln(x)}{4(x-1)^3}, \quad \tilde{F}_H(x) = -\frac{-1+6x-3x^2-2x^3+6x^2 \ln(x)}{24(x-1)^4}. \quad (40)$$

The total one-loop contributions to the cLFV amplitude  $e_b \rightarrow e_a \gamma$  and  $\Delta a_{e_a}^{331\text{ISS}}$  are

$$\begin{aligned} c_{(ab)R} &= \sum_{x=W,Y} c_{(ab)R}(x) + \sum_{k=1}^3 c_{(ab)R}(h_k^\pm), \\ c_{(ba)R} &= (c_{(ab)R}[a \leftrightarrow b]) \times \frac{m_{e_a}}{m_{e_b}}. \end{aligned} \quad (41)$$

The second line of Eq. (41) is derived from the equality that  $c_{(ba)R}(x) = (c_{(ab)R}(x)[b \leftrightarrow a]) \times (m_{e_a}/m_{e_b})$  for all  $x = W, Y, h_{1,2,3}^\pm$ . Formulas of contributions to  $a_{e_a}$  are:

$$a_{e_a} = -\frac{4m_{e_a}^2}{e} \text{Re}[c_{(aa)R}] = -\frac{4m_{e_a}^2}{2\pi^2 v^2} \text{Re}[c'_{(aa)R}], \quad c'_{(ab)R} = c_{(ab)R} \times \left( \frac{eg^2}{32\pi^2 m_W^2} \right)^{-1}. \quad (42)$$

One-loop contributions from heavy neutral Higgs bosons are very suppressed hence they are ignored here. The deviation of  $a_{e_a}$  between predictions by the two models 331ISS and SM are

$$\Delta a_{e_a} = \Delta a_{e_a}^{331\text{ISS}} \equiv a_{e_a} - a_{e_a}^{\text{SM}}(W), \quad (43)$$

where  $a_{e_a}^{\text{SM}}(W) = 5g^2 m_\mu^2 / (96\pi^2 m_W^2)$  is the SM's prediction [97]. In this work,  $\Delta a_{e_a}$  will be considered as new physics (NP) predicted by the 331ISS, used to compare with experimental data in numerical investigations.

The branching ratios of the cLFV processes are [96]

$$\text{Br}(e_b \rightarrow e_a \gamma) \simeq \frac{6\alpha_{em}}{\pi} \left( |c'_{(ab)R}|^2 + |c'_{(ba)R}|^2 \right) \text{Br}(e_b \rightarrow e_a \bar{\nu}_a \nu_b), \quad (44)$$

where  $G_F = 1/(\sqrt{2}v^2)$ , consistent with previous results [76, 93] for 3-3-1 models.

The formulas of  $U^\nu$  given in Eq. (21) results in approximate expressions of  $c_{(ab)R}$  and  $c_{(ba)R}$  with  $b \geq a$  as follows:

$$\begin{aligned} c'_{(ab)R}(W) &= -\frac{5}{12} \left[ \delta_{ab} - (\tilde{m}_D^\dagger V_R \hat{k}^{-2} \tilde{m}_D)_{ab} \right] + \sum_{e=1}^3 (\tilde{m}_D^\dagger V_R^* \hat{k}^{-1})_{ae} (\tilde{m}_D^T V_R \hat{k}^{-1})_{be} F_V(x'_{W,e}), \\ c'_{(ab)R}(Y) &= \frac{m_W^2}{m_Y^2} \sum_{e=1}^3 (V_R)_{ae}^* (V_R)_{be} F_V(x'_{Y,e}), \end{aligned}$$

$$\begin{aligned}
c'_{(ab)R}(h_1^\pm) &= \frac{z_0^2}{m_{h_1^\pm}^2} \sum_{e=1}^3 (\tilde{m}_D^* V_R^*)_{ae} \left\{ c_\alpha^2 s_\beta^2 (\tilde{m}_D^T V_R \hat{k}^{-1})_{be} - \frac{v s_{2\alpha} s_{2\beta}}{4m_b} [Y^{\sigma T} V_R]_{be} \right\} \hat{k}_e F_H(x'_{e,1}) \\
&+ \frac{c_\alpha^2 s_\beta^2 z_0^2}{m_{h_1^\pm}^2} \sum_{e=1}^3 (\tilde{m}_D V_R)_{ae}^* (\tilde{m}_D V_R)_{be} \tilde{F}_H(x'_{e,1}) \\
&+ \frac{1}{24} \left\{ \frac{m_{e_a}^2 c_\alpha^2 t_\beta^2}{m_{h_1^\pm}^2} \delta_{ab} + \frac{m_{e_a} v s_\alpha c_\alpha t_\beta}{\sqrt{2} m_{h_1^\pm}^2} \left[ \frac{m_{e_a}}{m_{e_b}} (R_2 Y^\sigma)_{ab} + (Y^{\sigma\dagger} R_2^\dagger)_{ab} \right] \right\} \\
&+ \sum_{e=1}^3 \tilde{F}_H(x'_{e,1}) c_\alpha^2 \left\{ \frac{m_{e_a}^2 t_\beta^2}{m_{h_1^\pm}^2} [(R_2 V_L)_{ae} (R_2 V_L)_{be}^*] + \frac{v^2 t_\alpha^2 m_{e_a}}{m_{e_b} m_{h_1^\pm}^2} [(Y^{\sigma T} V_R)_{ae} (Y^{\sigma T} V_R)_{be}^*] \right. \\
&\left. - \frac{m_{e_a} v s_\alpha c_\alpha t_\beta}{m_{h_1^\pm}^2} \left[ \frac{m_{e_a}}{m_{e_b}} (R_2 V_L)_{ae} (Y^{\sigma T} V_R)_{be}^* + (Y^{\sigma T} V_R)_{ae} (R_2 V_L)_{be}^* \right] \right\}, \quad (45)
\end{aligned}$$

$$c'_{(ab)R}(h_2^\pm) = c'_{(ab)R}(h_1^\pm) \left[ m_{h_1^\pm} \rightarrow m_{h_2^\pm}, c_\alpha \rightarrow s_\alpha, s_\alpha \rightarrow -c_\alpha \right],$$

$$\begin{aligned}
c'_{(ab)R}(h_3^\pm) &= \frac{z_0^2}{m_{h_3^\pm}^2} \left\{ \sum_{e=1}^3 \left[ (\tilde{m}_D^* \tilde{m}_D V_R \hat{k}^{-1})_{ae} (V_R^*)_{be} k_e F_H(x'_{e,3}) \right] - \frac{1}{24} (\tilde{m}_D^* \tilde{m}_D)_{ab} \right. \\
&\left. + \sum_{e=1}^3 \left[ (\tilde{m}_D \tilde{m}_D^* V_R \hat{k}^{-1})_{ae} (\tilde{m}_D^* \tilde{m}_D V_R^* \hat{k}^{-1})_{be} + \frac{m_{e_a}^2}{z_0^2 c_\beta^2} (V_R^*)_{be} (V_R)_{ae} \right] \tilde{F}_H(x'_{e,3}) \right\},
\end{aligned}$$

where the equalities in Eq. (22) were used. In addition, we ignore the minor contributions proportional to  $R_2^\dagger R_2$ , and  $R_2 R_2^\dagger$ . Because only two terms relating to  $R_2 Y^\sigma$  and  $R_2^\dagger R_2$  depend on  $V_L$ , but give small one-loop contributions to  $\Delta a_{e_a}$ , we fix  $V_L = I_3$  without loss of generality.

The above expressions of  $c_{(ab)R}$  and  $c_{(ba)R}$  given in Eq. (45) give some interesting properties. First, all terms are proportional to  $1/m_{h_k^\pm}^2$ , hence large  $|\Delta a_{e_a}|$  corresponding to large  $|c_{(aa)R}|$  will prefer small  $m_{h_k^\pm}^2$ . In contrast, experimental constraints on cLFV decay rates require small  $|c_{(ab)R}|$  and  $|c_{(ba)R}|$ , hence  $m_{h_k^\pm}^2$  should be large. It is easy to get small  $\text{Br}(e_b \rightarrow e_a \gamma)$  with enough large  $m_{h_k^\pm}$ , but difficult to get large  $|\Delta a_{e_a}|$ . Previously numerical investigation showed another situation [74], where small  $m_{h_k^\pm}^2$  are needed for large  $\Delta a_\mu$ , and the destructive correlations between particular terms in  $c_{(ab)R}$  and  $c_{(ba)R}$  must appear to result in small  $\text{Br}(e_b \rightarrow e_a \gamma)$ . The structure of the mass Dirac matrix  $\tilde{m}_D$  strongly affects these destructive correlations. As we will see, the antisymmetric property of  $\tilde{m}_D$  and the neutrino oscillation data fix a certain form of  $\tilde{m}_D$ , namely the fixed values considered in this work are  $(\tilde{m}_D)_{32} = -(\tilde{m}_D)_{23} = 1$ ,  $-(\tilde{m}_D)_{12} = (\tilde{m}_D)_{21} \simeq 0.613$ , and  $-(\tilde{m}_D)_{13} = (\tilde{m}_D)_{31} \simeq 0.357$ , and  $(\tilde{m}_D)_{11} = (\tilde{m}_D)_{22} = (\tilde{m}_D)_{33} = 0$ . They do not support large absolute values of the

diagonal entries relating to  $c_{(aa)R}$ . Therefore, the simple case of  $V_R = I_3$ , degenerate values of heavy neutrino masses  $\hat{k}_{11} = \hat{k}_{22} = \hat{k}_{33}$ , and  $Y^\sigma = \mathcal{O}_{3 \times 3}$  will give  $c_{(ab)R} \sim \tilde{m}_D \tilde{m}_D^*$ . As a result, constraints on cLFV decays always exclude the regions of parameter space predicting large  $(g-2)_{e,\mu}$ . This conclusion is completely consistent with the numerical results reported in Ref. [74]. In addition, the presence of  $\sigma^\pm$  and non-zero Yukawa coupling matrix  $Y^\sigma$  is necessary to explain  $1\sigma$  range of  $(g-2)_\mu$  obtained by experiment. Additionally, the formulas given in Eq. (45) explain explicitly that large  $\Delta a_\mu$  also needs large  $z_0$ . And, large  $t_\beta$  and non-zero  $Y^\sigma$  support more strong destructive correlations to guarantee that  $(e_b \rightarrow e_a)$  satisfies the current constraints.

Finally, We emphasize that the  $(g-2)_e$  data and LFVHD were not discussed previously for the 331ISS model. Our numerical investigation showed that large  $(g-2)_e$  requires nonzero values of  $s_\alpha$ , which was not considered in Ref. [74]. In addition, large values of  $Y_{22,33,23,32}^\sigma$  should be investigated carefully because they may result in too large  $\text{Br}(h \rightarrow \tau\mu)$  that may be excluded by the experimental constraints.

## B. Decays $h_1^0 \rightarrow e_a e_b$

The Yukawa couplings  $h_1^0 f f$  is , namely

$$\mathcal{L}_{h_1^0 f f}^Y = -\frac{g}{2m_W} h_1^0 \left[ \frac{1}{2} \sum_{i,j=1}^9 \bar{n}_i (\lambda_{ij}^0 P_L + \lambda_{ij}^{0*} P_R) n_j + m_{e_a} \bar{e}_a e_a \right], \quad (46)$$

where

$$\lambda_{ij}^0 = \sum_{c=1}^3 (U_{ci}^\nu U_{cj}^{\nu*} m_{n_i} + U_{ci}^{\nu*} U_{cj}^\nu m_{n_j}), \quad (47)$$

is a symmetric coefficient  $\lambda_{ij}^0 = \lambda_{ji}^0$  corresponding to the Feynman rules given in Ref. [91]. All of the Feynman rules for couplings involved in LFV processes at one-loop level are listed in Table I, where we used  $s_\theta = gv_1/(2m_Y)$ . We focus on the limit of tiny  $t_\theta \simeq s_\theta = 0$ , and the suppressed deviation of the SM-like Higgs mixing mentioned previously [76, 94]. Namely, they will be fixed to be zeros in the numerical calculations.

The effective Lagrangian and partial decay width of the decay  $h_1^0 \rightarrow e_a^\pm e_b^\mp$  are

$$\begin{aligned} \mathcal{L}^{\text{LFVH}} &= h_1^0 (\Delta_{(ab)L} \bar{e}_a P_L e_b + \Delta_{(ab)R} \bar{e}_a P_R e_b) + \text{H.c.}, \\ \Gamma(h_1^0 \rightarrow e_a e_b) &= \Gamma(h_1^0 \rightarrow e_a^- e_b^+) + \Gamma(h_1^0 \rightarrow e_a^+ e_b^-) = \frac{m_{h_1^0}}{8\pi} (|\Delta_{(ab)L}|^2 + |\Delta_{(ab)R}|^2), \end{aligned} \quad (48)$$

Vertex	Coupling
$h_1^0 \bar{e}_a e_a$	$-\frac{igm_{ea}}{2m_W}$
$h_1^0 \bar{n}_i n_j$	$-\frac{ig}{2m_W} \left( \lambda_{ij}^0 P_L + \lambda_{ij}^{0*} P_R \right)$
$h_k^+ \bar{n}_i e_b, h_k^- \bar{e}_a n_i$	$\frac{-ig}{\sqrt{2}m_W} \left( \lambda_{bi}^{L,k} P_L + \lambda_{bi}^{R,k} P_R \right), \frac{-ig}{\sqrt{2}m_W} \left( \lambda_{ai}^{L,k*} P_R + \lambda_{ai}^{R,k*} P_L \right)$
$W_\mu^+ \bar{n}_i e_b, W_\mu^- \bar{e}_a n_i$	$\frac{ig}{\sqrt{2}} U_{ai}^{\nu*} \gamma^\mu P_L, \frac{ig}{\sqrt{2}} U_{ai}^\nu \gamma^\mu P_L$
$Y_\mu^+ \bar{n}_i e_b, Y_\mu^- \bar{e}_a n_i$	$\frac{ig}{\sqrt{2}} U_{(a+3)i}^{\nu*} \gamma^\mu P_L, \frac{ig}{\sqrt{2}} U_{(a+3)i}^\nu \gamma^\mu P_L$
$h_3^+ h_1^0 Y_\mu^-, h_3^- Y_\mu^+ h_1^0$	$\frac{i}{2} g c_\beta c_\theta (p_+ - p_0)^\mu, -\frac{i}{2} g c_\beta c_\theta (p_- - p_0)^\mu$
$h_1^0 W_\mu^+ W_\nu^-$	$igm_W g^{\mu\nu}$
$h_1^0 Y_\mu^+ Y_\nu^-$	$ig c_\beta s_\theta m_Y g^{\mu\nu}$

TABLE I: Feynman rules for one-loop contributions to  $(g-2)$  anomalies,  $e_b \rightarrow e_a \gamma$ , and  $h_1^0 \rightarrow e_a e_b$  in the unitary gauge.  $p_0$  and  $p_\pm$  are the incoming momenta of  $h_1^0$  and  $h_3^\pm$ , respectively.

where the scalar factors  $\Delta_{(ab)L,R}$  are loop contributions in this work. In the unitary gauge, the one-loop Feynman diagrams contributing to  $\Delta_{(ab)L,R}$  are shown in Fig. 1. The valid




FIG. 1: One-loop Feynman diagrams contributing to the decay  $h_1^0 \rightarrow e_a e_b$  in the unitary gauge. Here  $V^\pm = W^\pm, Y^\pm$ ;  $k, l = 1, 2, 3$ .

condition  $m_{h_1^0} \gg m_{a,b}$  was used in Eq. (48), where  $m_{a,b}$  are the lepton masses satisfying  $p_{1,2}^2 = m_{a,b}^2$  and  $p_{h_1^0}^2 \equiv (p_1 + p_2)^2 = m_{h_1^0}^2$ . The branching ratio of a LfVHD is  $\text{Br}(h_1^0 \rightarrow e_a e_b) = \Gamma(h_1^0 \rightarrow e_a e_b) / \Gamma_{h_1^0}^{\text{total}}$ , where  $\Gamma_{h_1^0}^{\text{total}} \simeq 4.1 \times 10^{-3} \text{ GeV}$  [98, 99]. The  $\Delta_{(ab)L,R}$  can be written as

$$\Delta_{(ab)L,R} = \sum_{i=1,5,7,8} \Delta_{(ab)L,R}^{(i)W} + \sum_{i=1}^{10} \Delta_{(ab)L,R}^{(i)Y} \quad (49)$$

where the analytic forms of  $\Delta_{(ab)L,R}^{(i)W}$  and  $\Delta_{(ab)L,R}^{(i)Y}$  are shown in the Appendix. There are numbers of tiny one-loop contributions, which we will ignore in the numerical calculations. They are calculated using the unitary gauge with the same techniques given in Refs. [22, 76]. The contributions from diagrams (2), (3), and (5) with  $Y^\pm$  exchanges have suppressed factors  $c_\beta m_W^3/m_Y^3$ . The one-loop contributions from diagram (6) are suppressed with heavy singly charged Higgs bosons, which we checked consistently with the result mentioned in ref. [76].

#### IV. NUMERICAL DISCUSSION

In this work, we will use the neutrino oscillation data given in Ref. [99, 100]. The standard form of the lepton mixing matrix  $U_{\text{PMNS}}$  is the function of three angles  $\theta_{ij}$ , one Dirac phase  $\delta$  and two Majorana phases  $\alpha_1$  and  $\alpha_2$  [101], namely

$$U_{\text{PMNS}}^{\text{PDG}} = f(s_{12}, s_{13}, s_{23}, \delta) \times \text{diag}(1, e^{i\alpha_1}, e^{i\alpha_2}),$$

$$f(s_{12}, s_{13}, s_{23}, \delta) \equiv \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & s_{13}e^{-i\delta} \\ 0 & 1 & 0 \\ -s_{13}e^{i\delta} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (50)$$

where  $s_{ij} \equiv \sin \theta_{ij}$ ,  $c_{ij} \equiv \cos \theta_{ij} = \sqrt{1 - s_{ij}^2}$ ,  $i, j = 1, 2, 3$  ( $i < j$ ),  $0 \leq \theta_{ij} < 90$  [Deg.] and  $0 < \delta \leq 720$  [Deg.]. The Majorana phases are chosen in the range  $-180 \leq \alpha_i \leq 180$  [Deg.]. For numerical investigation, we will choose a benchmark corresponding to the normal order of the neutrino oscillation data as the input to fix  $\tilde{m}_D$  that  $s_{12}^2 = 0.32$ ,  $s_{23}^2 = 0.547$ ,  $s_{13}^2 = 0.0216$ ,  $\Delta m_{21}^2 = 7.55 \times 10^{-5}[\text{eV}^2]$ ,  $\Delta m_{32}^2 = 2.424 \times 10^{-3}[\text{eV}^2]$ ,  $\delta = 180$  [Deg], and  $\alpha_1 = \alpha_2 = 0$ . Consequently, the reduced Dirac mass matrix  $\tilde{m}_D$  is fixed as follows

$$\tilde{m}_D = \begin{pmatrix} 0 & 0.613 & 0.357 \\ -0.613 & 0 & 1 \\ -0.357 & -1 & 0 \end{pmatrix}. \quad (51)$$

The mixing matrix  $V_R$  is parameterized using the same formulas given in Eq. (50),  $V_R = f(s_{12}^r, s_{13}^r, s_{23}^r, 0)$  with  $|s_{ij}^r| \leq 1$ . The remaining free parameters are scanned in the following ranges:

$$\hat{k}_{1,2,3} \geq 5, \quad 600 \text{ [GeV]} \leq m_{h_{1,2,3}^\pm} \leq 1500 \text{ [GeV]}, \quad |s_\alpha| \leq 1, \quad \max[|Y_{ij}^\sigma|] \leq 1.5,$$



$$t_\beta \in [30, 70], 400 \text{ [GeV]} \leq z \leq 1200 \text{ [GeV]}. \quad (52)$$

The experimental parameters are  $G_F = 1.663787 \times 10^{-5} \text{ [GeV}^{-2}\text{]}$ ,  $g = 0.652$ ,  $\alpha_{em} = e^2/(4\pi) = 1/137$ ,  $s_W^2 = 0.231$ ,  $m_e = 5 \times 10^{-4} \text{ [GeV]}$ ,  $m_\mu = 0.105 \text{ [GeV]}$ ,  $m_\tau = 1.776 \text{ [GeV]}$ , and  $m_W = 80.385 \text{ [GeV]}$ ,  $\text{Br}(\mu \rightarrow e\bar{\nu}_e\nu_\mu) \simeq 1$ ,  $\text{Br}(\tau \rightarrow e\bar{\nu}_e\nu_\tau) \simeq 0.1782$ , and  $\text{Br}(\tau \rightarrow \mu\bar{\nu}_\mu\nu_\tau) \simeq 0.1739$ . We note that the upper bounds of  $m_{h_k^\pm}$  and  $t_\beta$  based on the previous work to accommodate large values of  $\Delta a_\mu$ . Chosen scanning range of  $t_\beta$  also satisfies the perturbative limit mentioned above.

We comment here the results obtained previously in ref. [74], where large  $t_\beta \geq 50$  and small values of singly charged Higgs bosons  $h_k^\pm$  ( $k = 1, 2, 3$ ) are required for large  $(g-2)_\mu$  satisfying  $1\sigma$  experimental data of  $(g-2)_\mu$  and all constraints from cLFV decays  $e_b \rightarrow e_a\gamma$ . But only the case of  $s_\alpha = 0$  and non-zero  $Y_{11,22,23,32}^\sigma$  was mentioned. Our numerical investigation shows that this case results in small  $\Delta a_e$  which cannot satisfy  $1\sigma$  range of the experimental data given in Eq. (3). Without  $\sigma^\pm$ , we obtain two maximal values of  $\Delta a_e$  that  $\Delta a_e \leq 2.5 \times 10^{-14}$  and  $1.5 \times 10^{-14}$  for the NO and IO schemes, respectively. Hence, determining the regions of parameter space giving large  $\Delta a_e$  will be very interesting.

Because of the above reasons, we focus on the regions of parameter space giving large  $\Delta a_e$  that satisfies the  $1\sigma$  experimental data of  $(g-2)_e$  as well as all current constraints of cLFV decay rates  $\text{Br}(e_b \rightarrow e_a\gamma)$ . The investigation shows that the  $1\sigma$  range of  $\Delta a_e \in [1.8 \times 10^{-13}, 7.8 \times 10^{-13}]$  can be obtained easily in a wide range of the parameter space, for example the following fixed values of  $z_0 = 500 \text{ GeV}$ ,  $m_{h_1^\pm} = 639.08 \text{ GeV}$ ,  $m_{h_2^\pm} = 902.1 \text{ GeV}$ ,  $m_{h_3^\pm} = 730.2 \text{ GeV}$ ,  $m_{n_4} = m_{n_7} = 2.0126 \text{ TeV}$ ,  $m_{n_5} = m_{n_8} = 2.2318 \text{ TeV}$ ,  $t_\beta = 50$ , and  $s_\alpha = 0.5$ , and scanning the remaining parameters, we have a benchmark point that  $m_{n_6} = m_{n_9} = 1.801 \text{ TeV}$ ,  $s_{12}^r = -0.3924$ ,  $s_{13}^r = -0.8304$ ,  $s_{23}^r = -0.1148$ , and  $0 \leq |Y_{ab}^\sigma| \leq 0.02$ , which result in the following allowed values of relevant physical processes:  $\Delta a_e = 2.529 \times 10^{-13}$ ,  $\Delta a_\mu = 0.666 \times 10^{-9}$ ,  $\text{Br}(\mu \rightarrow e\gamma) = 2.513 \times 10^{-13}$ ,  $\text{Br}(\tau \rightarrow e\gamma) = 8.015 \times 10^{-10}$ ,  $\text{Br}(\tau \rightarrow \mu\gamma) = 1.71 \times 10^{-8}$ ,  $\text{Br}(h_1^0 \rightarrow \mu e) = 4.511 \times 10^{-9}$ ,  $\text{Br}(h_1^0 \rightarrow \tau e) = 1.72 \times 10^{-6}$ , and  $\text{Br}(h_1^0 \rightarrow \tau\mu) = 1.64 \times 10^{-5}$ .

We list here other interesting benchmark points of the parameter space corresponding to large  $t_\beta = 60$ , that satisfy  $1\sigma$  range of  $(g-2)_e$ ,  $\Delta a_\mu \geq 0.6 \times 10^{-9}$ , and all current LFV upper bounds. For other large  $t_\beta$  values, the results are the same.

1. A benchmark point giving large  $\text{Br}(h_1^0 \rightarrow \tau e) \sim \mathcal{O}(10^{-5})$ :

$$\begin{aligned} \{z_0[\text{GeV}], t_\beta, s_\alpha\} &= \{539.6, 60, 0.336\}, \{s_{12,13,23}^r\} = \{-0.1757, -0.0051, 0.2142\}, \\ \{m_{h_{1,2,3}^\pm}[\text{TeV}]\} &= \{0.624, 0.709, 0.98\}, \{m_{4,5,6} = m_{7,8,9}[\text{TeV}]\} = \{2.084, 3.286, 2.245\}, \\ Y^\sigma &= \begin{pmatrix} 0.00693 & -0.0186 & 0.0336 \\ 0.0278 & -0.0402 & 0.0062 \\ 0.0039 & 0.0384 & 0.0056 \end{pmatrix}. \end{aligned}$$

The corresponding values of  $\Delta a_{e,\mu}$  and LFV decay rates are

$$\begin{aligned} \Delta a_e &= 5.931 \times 10^{-13}, \Delta a_\mu = 0.516 \times 10^{-9}, \\ \text{Br}\{(\mu \rightarrow e\gamma), (\tau \rightarrow e\gamma), (\tau \rightarrow \mu\gamma)\} &= \{2.6 \times 10^{-13}, 4.22 \times 10^{-9}, 1.03 \times 10^{-8}\} \\ \text{Br}(h_1^0 \rightarrow \{\mu e, \tau e, \tau\mu\}) &= \{1.23 \times 10^{-8}, 1.95 \times 10^{-5}, 1.38 \times 10^{-5}\}. \end{aligned}$$

2. There exists benchmark point that allows large  $\text{Br}(h_1^0 \rightarrow \tau e) \sim \mathcal{O}(10^{-5})$ , but small  $\text{Br}(h_1^0 \rightarrow \tau e) < (10^{-8})$ :

$$\begin{aligned} \{z_0[\text{GeV}], t_\beta, s_\alpha\} &= \{623.9, 60, 0.444\}, \{s_{12,13,23}^r\} = \{0.006, 0.0157, 0.3661\}, \\ \{m_{h_{1,2,3}^\pm}[\text{TeV}]\} &= \{0.624, 0.709, 0.980\}, \{m_{4,5,6} = m_{7,8,9}[\text{TeV}]\} = \{2.958, 3.432, 2.888\}, \\ Y^\sigma &= \begin{pmatrix} 0.004 & -0.912 & 0.068 \\ 0.0351 & -0.0342 & -0.130 \\ -0.0004 & 0.0454 & 0.0213 \end{pmatrix}, \end{aligned}$$

$$\begin{aligned} \Delta a_e &= 7.781 \times 10^{-13}, \Delta a_\mu = 0.977 \times 10^{-9}, \\ \text{Br}\{(\mu \rightarrow e\gamma), (\tau \rightarrow e\gamma), (\tau \rightarrow \mu\gamma)\} &= \{1.67 \times 10^{-13}, 2.29 \times 10^{-8}, 4.0 \times 10^{-8}\}, \\ \text{Br}(h_1^0 \rightarrow \{\mu e, \tau e, \tau\mu\}) &= \{3.96 \times 10^{-7}, 5.7 \times 10^{-5}, 1.96 \times 10^{-9}\}. \end{aligned}$$

3. A larger  $|Y^\sigma|$  giving a benchmark point with large  $\text{Br}(h \rightarrow \tau\mu) \sim \mathcal{O}(10^{-4})$ :

$$\begin{aligned} \{z_0[\text{GeV}], t_\beta, s_\alpha\} &= \{741.2, 60, 0.931\}, \{s_{12,13,23}^r\} = \{0.685, -0.67, 0.127\}, \\ \{m_{h_{1,2,3}^\pm}[\text{TeV}]\} &= \{0.861, 0.871, 0.98\}, \{m_{4,5,6} = m_{7,8,9}[\text{TeV}]\} = \{2.301, 3.150, 2.355\}, \\ Y^\sigma &= \begin{pmatrix} 0.0536 & 0.07 & -0.409 \\ 0.108 & -0.385 & 0.019 \\ 0.036 & 0.656 & 0.007 \end{pmatrix}, \end{aligned}$$

$$\begin{aligned}\Delta a_e &= 3.437 \times 10^{-13}, \quad \Delta a_\mu = 0.883 \times 10^{-9}, \\ \text{Br}\{(\mu \rightarrow e\gamma), (\tau \rightarrow e\gamma), (\tau \rightarrow \mu\gamma)\} &= \{2.1 \times 10^{-14}, 1.4 \times 10^{-9}, 2.73 \times 10^{-8}\}, \\ \text{Br}(h_1^0 \rightarrow \{\mu e, \tau e, \tau\mu\}) &= \{1.02 \times 10^{-8}, 1.08 \times 10^{-5}, 1.05 \times 10^{-4}\}.\end{aligned}$$

From our numerical investigation, we found that the regions allowing  $1\sigma$  range of  $\Delta a_e$  data and cLFV constraints are very wide. But the regions allowed large  $(g-2)_\mu$  are difficult to control. This is because of large free parameters in the model that our code is not still smart to collect these points. Because of the special form of  $\tilde{m}_D$  that require the non-degenerate matrix  $\hat{k}$  and the strong destructive correlations between the mixing angles  $s_{ab}^x$  and the entries of  $Y^\sigma$  in order to get small  $\text{Br}(e_b \rightarrow e_a\gamma)$ , in the regions allow large  $\Delta a_\mu$ . There may exist some certain relations between these parameters for collecting more interesting points allowing large  $(g-2)_\mu$  at  $1\sigma$  experimental range. We will determine them in a future work.

## V. CONCLUSION

In this work, we have constructed the analytic formulas for one-loop contributions to the LFV decays of the SM-like Higgs boson  $h_1^0 \rightarrow e_a e_b$  in the 3-3-1 model with inverse seesaw neutrinos. We also give an analytic formulas to explain qualitatively the results of large  $(g-2)_\mu$  reported previously. Numerical tests is used to confirm the consistency between the two calculations. We introduced new parameterisation of the heavy neutrino mass matrix to reduce the number of free parameters used to investigate  $(g-2)_{e,\mu}$  anomalies and LFV decays  $e_b \rightarrow e_a\gamma$  and  $h_1^0 \rightarrow e_a e_b$ . Our numerical investigation shows that the model can predict easily the  $1\sigma$  range of experimental data for  $(g-2)_e$  and satisfy simultaneously the cLFV constrains  $\text{Br}(e_b \rightarrow e_a\gamma)$ . But we only obtain the regions of parameter space that give largest values of  $\Delta a_\mu \simeq 10^{-9}$ , which rather smaller than the lower bound of  $1\sigma$  range reported recently. The reason is that recent numerical code used in our investigation only works in the limit of small  $|y^\sigma| < 0.1$ . In the regions of the parameter space, the largest values of  $\text{Br}(h_1^0 \rightarrow \tau e, \tau\mu)$  are order of  $\mathcal{O}(10^{-5})$ , while  $\text{Br}(h_1^0 \rightarrow \mu e) < 10^{-7}$  which is still smaller than the recent experimental bounds. The regions with large  $|Y^\sigma|_{ab}$  may be more interesting, which is our future work, where many other LFV processes such as  $Z \rightarrow e_b e_a$ ,  $e_b \rightarrow e_c e_d e_f$ , and the  $\mu - e$  conversion in nuclei will be discussed together.

## Acknowledgments

This research is funded by An Giang University under grant number 21.01.TB. L. T. Hue is thankful to Van Lang University.

## Appendix A: Form factors of LFVHD in the unitary gauge

The one-loop contributions here are calculated using the notations of Passarino functions [102, 103] given in Ref. [24], consistent with LoopTools [104], see a detailed discussion in ref. [30, 105]. The Passarino-Veltman (PV) functions used in this work are defined as follows:  $B_\mu^{(i)} \equiv B_1^{(i)} \times (-1)^i p_{i\mu}$  with  $i = 1, 2$ , and  $C_\mu \equiv \sum_{i=1}^2 (-1)^i p_{i\mu} \times C_i$ . As mentioned in Ref. [24], the two  $B_1^{(1)}$  and  $C_1$  have opposite signs with those introduced in Ref. [30]. They come from the signs of  $p_{1,2}$  in the internal momenta  $(k - p_1)$  and  $(k + p_2)$  shown in Fig. 1, which  $p_1$  has an opposite sign, which is different from the standard notation of  $k + p_1$  defined in LoopTools. The PV-functions used in our formulas are:  $B_{0,1}^{(i)} = B_{0,1}(p_i^2; M_0^2, M_i^2)$ ,  $C_{0,1,2} = C_{0,1,2}(p_1^2, (p_1 + p_2)^2, p_2^2; M_0^2, M_1^2, M_2^2)$ , and  $B_0^{(12)} = B_0((p_1 + p_2)^2; M_1^2, M_2^2)$ . In the below, when the external momenta are fixed as  $p_1^2 = m_{e_a}^2$ ,  $p_2^2 = m_{e_b}^2$ , and  $(p_1 + p_2)^2 = m_{h_1^0}^2$ , we use the simpler notations as follows  $C_{0,1,2}(p_1^2, m_{h_1^0}^2, p_2^2; M_0^2, M_1^2, M_2^2) \equiv C_{0,1,2}(M_0^2, M_1^2, M_2^2)$ ,  $B_{0,1}^{(i)}(M_0^2, M_i^2) = B_{0,1}(p_i^2; M_0^2, M_i^2)$ , and  $B_0(m_{h_1^0}^2; M_1^2, M_2^2) = B_0^{(12)}(M_1^2, M_2^2)$ .

The analytic expressions  $\Delta_{L,R}^{(i)W} \equiv \Delta_{(ab)L,R}^{(i)W}$  for one-loop contributions from the diagram (i) in Fig. 1 are

$$\begin{aligned} \Delta_L^{(1)W} &= \frac{g^3 m_a}{64\pi^2 m_W^3} \sum_{i=1}^9 U_{ai}^\nu U_{bi}^{*\nu} \left\{ m_{n_i}^2 \left( B_0^{(1)} + B_0^{(2)} + B_1^{(1)} \right) + m_b^2 B_1^{(2)} - \left( 2m_W^2 + m_{h_1^0}^2 \right) m_{n_i}^2 C_0 \right. \\ &\quad - \left[ m_{n_i}^2 \left( 2m_W^2 + m_{h_1^0}^2 \right) + 2m_W^2 \left( 2m_W^2 + m_a^2 - m_b^2 \right) \right] C_1 \\ &\quad \left. - \left[ 2m_W^2 \left( m_a^2 - m_{h_1^0}^2 \right) + m_b^2 m_{h_1^0}^2 \right] C_2 \right\}, \\ \Delta_R^{(1)W} &= \frac{g^3 m_b}{64\pi^2 m_W^3} \sum_{i=1}^9 U_{ai}^\nu U_{bi}^{*\nu} \left\{ m_{n_i}^2 \left( B_0^{(1)} + B_0^{(2)} + B_1^{(2)} \right) + m_a^2 B_1^{(1)} - \left( 2m_W^2 + m_{h_1^0}^2 \right) m_{n_i}^2 C_0 \right. \\ &\quad - \left[ m_{n_i}^2 \left( 2m_W^2 + m_{h_1^0}^2 \right) + 2m_W^2 \left( 2m_W^2 - m_a^2 + m_b^2 \right) \right] C_2 \\ &\quad \left. - \left[ 2m_W^2 \left( m_b^2 - m_{h_1^0}^2 \right) + m_a^2 m_{h_1^0}^2 \right] C_1 \right\}, \\ \Delta_L^{(7+8)W} &= \frac{g^3 m_a m_b^2}{64\pi^2 m_W^3 (m_b^2 - m_a^2)} \end{aligned}$$

$$\times \sum_{i=1}^9 U_{ai}^\nu U_{bi}^{\nu*} \left[ 2m_{n_i}^2 \left( B_0^{(2)} - B_0^{(1)} \right) + (2m_W^2 + m_{n_i}^2) \left( B_1^{(2)} - B_1^{(1)} \right) + m_b^2 B_1^{(2)} - m_a^2 B_1^{(1)} \right],$$

$$\Delta_R^{(7+8)W} = \frac{m_a}{m_b} \Delta_L^{(7+8)W},$$

where  $B_{0,1}^{(k)} = B_{0,1}^{(k)}(m_{n_i}^2, m_W^2)$  and  $C_{0,1,2} = C_{0,1,2}(m_{n_i}^2, m_W^2, m_W^2)$ ,

$$\Delta_L^{(5)W} = \frac{g^3 m_a}{64\pi^2 m_W^3} \sum_{i,j=1}^9 U_{ai}^{\nu*} U_{bj}^\nu \left\{ D_{ij} \left[ -m_{n_j}^2 B_0^{(12)} + m_{n_i}^2 B_1^{(1)} + m_j^2 m_W^2 C_0 \right. \right. \\ \left. \left. + \left( 2m_W^2 (m_{n_i}^2 + m_{n_j}^2) + 2m_{n_i}^2 m_{n_j}^2 - m_a^2 m_{n_j}^2 - m_b^2 m_{n_i}^2 \right) C_1 \right] \right. \\ \left. + D_{ij}^* m_{n_i} m_{n_j} \left[ -B_0^{(12)} + B_1^{(1)} + m_W^2 C_0 + \left( 4m_W^2 + m_{n_i}^2 + m_{n_j}^2 - m_a^2 - m_b^2 \right) C_1 \right] \right\},$$

$$\Delta_R^{(5)W} = \frac{g^3 m_b}{64\pi^2 m_W^3} \sum_{i,j=1}^9 U_{ai}^{\nu*} U_{bj}^\nu \left\{ D_{ij} \left[ -m_{n_i}^2 B_0^{(12)} + m_{n_j}^2 B_1^{(2)} + m_{n_i}^2 m_W^2 C_0 \right. \right. \\ \left. \left. + \left( 2m_W^2 (m_{n_i}^2 + m_{n_j}^2) + 2m_{n_i}^2 m_{n_j}^2 - m_a^2 m_{n_j}^2 - m_b^2 m_{n_i}^2 \right) C_2 \right] \right. \\ \left. + D_{ij}^* m_{n_i} m_{n_j} \left[ -B_0^{(12)} + B_1^{(2)} + m_W^2 C_0 + \left( 4m_W^2 + m_{n_i}^2 + m_{n_j}^2 - m_a^2 - m_b^2 \right) C_2 \right] \right\},$$

where  $D_{ij} = \sum_{c=1}^3 U_{ci}^\nu U_{cj}^{\nu*}$ ,  $B_0^{(12)} = B_0^{(12)}(m_{n_i}^2, m_{n_j}^2)$ ,  $B_1^{(1)} = B_1^{(1)}(m_W^2, m_{n_i}^2)$ ,  $B_1^{(2)} = B_1^{(2)}(m_W^2, m_{n_j}^2)$ , and  $C_{0,1,2} = C_{0,1,2}(m_W^2, m_{n_i}^2, m_{n_j}^2)$ . The analytic expressions  $\Delta_{L,R}^{(i)Y} \equiv \Delta_{(ab)L,R}^{(i)Y h_3^\pm}$  with  $i = 4, 6, 9, 10$ , are

$$\Delta_L^{(1)Y} = \frac{g^3 m_a c_\beta s_\theta}{64\pi^2 m_Y^3} \sum_{i=1}^9 U_{(a+3)i}^\nu U_{(b+3)i}^{\nu*} \left\{ m_{n_i}^2 \left( B_0^{(1)} + B_0^{(2)} + B_1^{(1)} \right) + m_b^2 B_1^{(2)} \right. \\ \left. - \left( 2m_Y^2 + m_{h_1^0}^2 \right) m_{n_i}^2 C_0 - \left[ 2m_Y^2 (2m_Y^2 + m_a^2 - m_b^2) + m_{n_i}^2 (2m_Y^2 + m_{h_1^0}^2) \right] C_1 \right. \\ \left. + \left[ 2m_Y^2 (m_a^2 - m_{h_1^0}^2) + m_b^2 m_{h_1^0}^2 \right] C_2 \right\},$$

$$\Delta_R^{(1)Y} = \frac{g^3 m_b c_\beta s_\theta}{64\pi^2 m_Y^3} \sum_{i=1}^9 U_{(a+3)i}^{\nu*} U_{(b+3)i}^\nu \left\{ m_{n_i}^2 \left( B_0^{(1)} + B_0^{(2)} + B_1^{(2)} \right) + m_a^2 B_1^{(1)} \right. \\ \left. - \left( 2m_Y^2 + m_{h_1^0}^2 \right) m_{n_i}^2 C_0 - \left[ 2m_Y^2 (2m_Y^2 - m_a^2 + m_b^2) + m_{n_i}^2 (2m_Y^2 + m_{h_1^0}^2) \right] C_2 \right. \\ \left. + \left[ 2m_Y^2 (m_b^2 - m_{h_1^0}^2) + m_b^2 m_{h_1^0}^2 \right] C_1 \right\},$$

$$\Delta_L^{(7+8)Y} = \frac{g^3 m_a m_b^2}{64\pi^2 m_W m_Y^2 (m_b^2 - m_a^2)} \sum_{i=1}^9 U_{(a+3)i}^{\nu*} U_{(b+3)i}^\nu \\ \times \left[ 2m_{n_i}^2 \left( B_0^{(2)} - B_0^{(1)} \right) + (2m_Y^2 + m_{n_i}^2) \left( B_1^{(2)} - B_1^{(1)} \right) - m_a^2 B_1^{(1)} + m_b^2 B_1^{(2)} \right],$$

$$\Delta_R^{(7+8)Y} = \frac{m_a}{m_b} \Delta_L^{(7+8)Y},$$

where  $B_{0,1}^{(k)} = B_{0,1}^{(k)}(m_{n_i}^2, m_Y^2)$  and  $C_{0,1,2} = C_{0,1,2}(m_{n_i}^2, m_Y^2, m_Y^2)$ . One-loop contributions from diagram 5 are

$$\begin{aligned}\Delta_L^{(5)Y} &= \frac{g^3 m_a}{64\pi^2 m_W m_Y^2} \\ &\times \sum_{i,j=1}^9 U_{(a+3)i}^\nu U_{(b+3)j}^{\nu*} \left\{ D_{ij} \left[ -m_{n_j}^2 B_0^{(12)} + m_{n_i}^2 B_1^{(1)} + m_j^2 m_W^2 C_0 \right. \right. \\ &+ \left. \left. \left( 2m_W^2 (m_{n_i}^2 + m_{n_j}^2) + 2m_{n_i}^2 m_{n_j}^2 - m_a^2 m_{n_j}^2 - m_b^2 m_{n_i}^2 \right) C_1 \right] \right. \\ &+ \left. D_{ij}^* m_{n_i} m_{n_j} \left[ -B_0^{(12)} + B_1^{(1)} + m_W^2 C_0 + \left( 4m_W^2 + m_{n_i}^2 + m_{n_j}^2 - m_a^2 - m_b^2 \right) C_1 \right] \right\}, \\ \Delta_R^{(5)Y} &= \frac{g^3 m_b}{64\pi^2 m_W m_Y^2} \\ &\times \sum_{i,j=1}^9 U_{(a+3)i}^\nu U_{(b+3)j}^{\nu*} \left\{ D_{ij} \left[ -m_{n_i}^2 B_0^{(12)} + m_{n_j}^2 B_1^{(2)} + m_{n_i}^2 m_W^2 C_0 \right. \right. \\ &+ \left. \left. \left( 2m_W^2 (m_{n_i}^2 + m_{n_j}^2) + 2m_{n_i}^2 m_{n_j}^2 - m_a^2 m_{n_j}^2 - m_b^2 m_{n_i}^2 \right) C_2 \right] \right. \\ &+ \left. D_{ij}^* m_{n_i} m_{n_j} \left[ -B_0^{(12)} + B_1^{(2)} + m_W^2 C_0 + \left( 4m_W^2 + m_{n_i}^2 + m_{n_j}^2 - m_a^2 - m_b^2 \right) C_2 \right] \right\},\end{aligned}$$

where  $B_0^{(12)} = B_0^{(12)}(m_{n_i}^2, m_{n_j}^2)$ ,  $B_1^{(1)} = B_1^{(1)}(m_Y^2, m_{n_i}^2)$ ,  $B_1^{(2)} = B_1^{(2)}(m_Y^2, m_{n_j}^2)$ , and  $C_{0,1,2} = C_{0,1,2}(m_Y^2, m_{n_i}^2, m_{n_j}^2)$ .

$$\begin{aligned}\Delta_L^{(2)Y} &= -\frac{g^3 m_a c_\theta c_\beta}{64\pi^2 m_W m_Y^2} \sum_{i=1}^9 U_{(a+3)i}^\nu \\ &\times \left\{ \lambda_{bi}^{L,1} m_{n_i} \left[ B_0^{(1)} + B_1^{(1)} + \left( m_Y^2 + m_{h_3^\pm}^2 - m_{h_1^0}^2 \right) C_0 - \left( m_Y^2 - m_{h_3^\pm}^2 + m_{h_1^0}^2 \right) C_1 \right] \right. \\ &- \left. \lambda_{bi}^{R,1} m_b \left[ 2m_Y^2 C_1 + \left( m_Y^2 + m_{h_3^\pm}^2 - m_{h_1^0}^2 \right) C_2 \right] \right\}, \\ \Delta_R^{(2)Y} &= \frac{g^3 c_\theta c_\beta}{64\pi^2 m_W m_Y^2} \sum_{i=1}^9 U_{(a+3)i}^\nu \\ &\times \left\{ \lambda_{bi}^{L,1} m_b m_{n_i} \left[ 2m_Y^2 C_0 + \left( m_Y^2 - m_{h_3^\pm}^2 + m_{h_1^0}^2 \right) C_2 \right] \right. \\ &+ \lambda_{bi}^{R,1} \left[ m_{n_i}^2 B_0^{(1)} + m_a^2 B_1^{(1)} - m_{n_i}^2 \left( m_Y^2 - m_{h_3^\pm}^2 + m_{h_1^0}^2 \right) C_0 \right. \\ &+ \left. \left. \left[ 2m_Y^2 \left( m_{h_1^0}^2 - m_b^2 \right) - m_a^2 \left( m_Y^2 - m_{h_3^\pm}^2 + m_{h_1^0}^2 \right) \right] C_1 - 2m_b^2 m_Y^2 C_2 \right] \right\},\end{aligned}$$

where  $B_k^{(1)} = B_k^{(1)}(m_Y^2, m_{n_i}^2)$  ( $k = 0, 1$ ) and  $C_{0,1,2} = C_{0,1,2}(m_{n_i}^2, m_Y^2, m_{h_3^\pm}^2)$ ,

$$\begin{aligned}\Delta_L^{(3)Y} &= \frac{g^3 c_\theta c_\beta}{64\pi^2 m_W m_Y^2} \sum_{i=1}^9 U_{(b+3)i}^{\nu*} \\ &\times \left\{ \lambda_{ai}^{L,1*} m_a m_{n_i} \left[ 2m_Y^2 C_0 + \left( m_Y^2 - m_{h_3^\pm}^2 + m_{h_1^0}^2 \right) C_1 \right] \right\}\end{aligned}$$

$$\begin{aligned}
& + \lambda_{ai}^{R,1*} \left[ m_{n_i}^2 B_0^{(2)} + m_b^2 B_1^{(2)} - m_{n_i}^2 \left( m_Y^2 - m_{h_3^\pm}^2 + m_{h_1^0}^2 \right) C_0 \right. \\
& \left. - 2m_a^2 m_Y^2 C_1 + \left[ 2m_Y^2 \left( m_{h_1^0}^2 - m_a^2 \right) - m_b^2 \left( m_Y^2 - m_{h_3^\pm}^2 + m_{h_1^0}^2 \right) \right] C_2 \right] \Big\}, \\
\Delta_R^{(3)Y} &= \frac{g^3 m_b c_\theta c_\beta}{64\pi^2 m_W m_Y^2} \\
& \times \sum_{i=1}^9 U_{(b+3)i}^{\nu*} \left\{ \lambda_{ai}^{L,1*} m_{n_i} \left[ B_0^{(2)} + B_1^{(2)} + \left( m_Y^2 + m_{h_3^\pm}^2 - m_{h_1^0}^2 \right) C_0 - \left( m_Y^2 - m_{h_3^\pm}^2 + m_{h_1^0}^2 \right) C_2 \right] \right. \\
& \left. - \lambda_{ai}^{R,1*} m_a \left[ \left( m_Y^2 + m_{h_3^\pm}^2 - m_{h_1^0}^2 \right) C_1 + 2m_Y^2 C_2 \right] \right\},
\end{aligned}$$

where  $B_k^{(2)} = B_k^{(2)}(m_Y^2, m_{n_i}^2)$  and  $C_{0,1,2} = C_{0,1,2}(m_{n_i}^2, m_{h_3^\pm}^2, m_Y^2)$ ,

$$\begin{aligned}
\Delta_L^{(4)h_{k,l}^\pm} &= \frac{g^2 g_{hkl}}{32\pi^2 m_W^2} \sum_{i=1}^9 \left[ -\lambda_{ai}^{R,k*} \lambda_{bi}^{L,k} m_{n_i} C_0 + \lambda_{ai}^{L,k*} \lambda_{bi}^{L,k} m_a C_1 + \lambda_{ai}^{R,k*} \lambda_{bi}^{R,k} m_b C_2 \right], \\
\Delta_R^{(4)h_{k,l}^\pm} &= \frac{g^2 g_{hkl}}{32\pi^2 m_W^2} \sum_{i=1}^9 \left[ -\lambda_{ai}^{L,k*} \lambda_{bi}^{R,k} m_{n_i} C_0 + \lambda_{ai}^{R,k*} \lambda_{bi}^{R,k} m_a C_1 + \lambda_{ai}^{L,k*} \lambda_{bi}^{L,k} m_b C_2 \right],
\end{aligned}$$

where  $\{k, l\} = \{1, 2\}, \{2, 1\}, \{1, 1\}, \{2, 2\}, \{3, 3\}$ ,  $g_{h21} = g_{h12}$  and  $C_{0,1,2} = C_{0,1,2}(m_{n_i}^2, m_{h_k^\pm}^2, m_{h_l^\pm}^2)$ ,

$$\begin{aligned}
\Delta_L^{(6)h_k^\pm} &= \frac{g^3}{64\pi^2 m_W^3} \sum_{i,j=1}^9 \left\{ \lambda_{ij}^{0*} \left[ \lambda_{ai}^{R,k*} \lambda_{bj}^{L,k} \left( B_0^{(12)} + m_{H_k^\pm}^2 C_0 + m_a^2 C_1 + m_b^2 C_2 \right) \right. \right. \\
& \left. \left. + \lambda_{ai}^{R,k*} \lambda_{bj}^{R,k} m_b m_{n_j} C_2 + \lambda_{ai}^{L,k*} \lambda_{bj}^{L,k} m_a m_{n_i} C_1 \right] \right. \\
& \left. + \lambda_{ij}^0 \left[ \lambda_{ai}^{R,k*} \lambda_{bj}^{L,k} m_{n_i} m_{n_j} C_0 + \lambda_{ai}^{R,k*} \lambda_{bj}^{R,k} m_{n_i} m_b (C_0 + C_2) \right. \right. \\
& \left. \left. + \lambda_{ai}^{L,k*} \lambda_{bj}^{L,k} m_a m_{n_j} (C_0 + C_1) + \lambda_{ai}^{L,k*} \lambda_{bj}^{R,k} m_a m_b (C_0 + C_1 + C_2) \right] \right\}, \\
\Delta_R^{(6)h_k^\pm} &= \frac{g^3}{64\pi^2 m_W^3} \sum_{i,j=1}^9 \left\{ \lambda_{ij}^0 \left[ \lambda_{ai}^{L,k*} \lambda_{bj}^{R,k} \left( B_0^{(12)} + m_{H_k^\pm}^2 C_0 + m_a^2 C_1 + m_b^2 C_2 \right) \right. \right. \\
& \left. \left. + \lambda_{ai}^{L,k*} \lambda_{bj}^{L,k} m_b m_{n_j} C_2 + \lambda_{ai}^{R,k*} \lambda_{bj}^{R,k} m_a m_{n_i} C_1 \right] \right. \\
& \left. + \lambda_{ij}^{0*} \left[ \lambda_{ai}^{L,k*} \lambda_{bj}^{R,k} m_{n_i} m_{n_j} C_0 + \lambda_{ai}^{L,k*} \lambda_{bj}^{L,k} m_{n_i} m_b (C_0 + C_2) \right. \right. \\
& \left. \left. + \lambda_{ai}^{R,k*} \lambda_{bj}^{R,k} m_a m_{n_j} (C_0 + C_1) + \lambda_{ai}^{R,k*} \lambda_{bj}^{L,k} m_a m_b (C_0 + C_1 + C_2) \right] \right\},
\end{aligned}$$

where  $k = 1, 2, 3$ ,  $B_0^{(12)} = B_0^{(12)}(m_{n_i}^2, m_{n_j}^2)$ , and  $C_{0,1,2} = C_{0,1,2}(m_{n_i}^2, m_{n_j}^2, m_{h_k^\pm}^2)$ ,

$$\begin{aligned}
\Delta_L^{(9+10)h_k^\pm} &= \frac{g^3}{64\pi^2 m_W^3 (m_a^2 - m_b^2)} \\
& \times \sum_{i=1}^9 \left[ m_a m_b m_{n_i} \lambda_{ai}^{L,k*} \lambda_{bi}^{R,k} \left( B_0^{(1)} - B_0^{(2)} \right) + m_{n_i} \lambda_{ai}^{R,k*} \lambda_{bi}^{L,k} \left( m_b^2 B_0^{(1)} - m_a^2 B_0^{(2)} \right) \right]
\end{aligned}$$

$$\begin{aligned}
& +m_a m_b \left( \lambda_{ai}^{L,k*} \lambda_{bi}^{L,k} m_b + \lambda_{ai}^{R,k*} \lambda_{bi}^{R,k} m_a \right) \left( -B_1^{(1)} + B_1^{(2)} \right) \Big], \\
\Delta_R^{(9+10)h_k^\pm} &= \frac{g^3}{64\pi^2 m_W^3 (m_a^2 - m_b^2)} \\
& \times \sum_{i=1}^9 \left[ m_a m_b m_{n_i} \lambda_{ai}^{R,k*} \lambda_{bi}^{L,k} \left( B_0^{(1)} - B_0^{(2)} \right) + m_{n_i} \lambda_{ai}^{L,k*} \lambda_{bi}^{R,k} \left( m_b^2 B_0^{(1)} - m_a^2 B_0^{(2)} \right) \right. \\
& \left. + m_a m_b \left( \lambda_{ai}^{R,k*} \lambda_{bi}^{R,k} m_b + \lambda_{ai}^{L,k*} \lambda_{bi}^{L,k} m_a \right) \left( -B_1^{(1)} + B_1^{(2)} \right) \right],
\end{aligned}$$

where  $k = 1, 2, 3$ ,  $B_{0,1}^{(k)} = B_{0,1}^{(k)}(m_{n_i}^2, m_{h_k^\pm}^2)$ . The details to derive the above formulas of  $\Delta_{L,R}^{(i)}$  were shown in Refs. [22, 88], and hence we do not present them in this work. We note that the scalar functions  $\Delta_{L,R}^{(1)W}$  and  $\Delta_{L,R}^{(1,2,3)Y}$  include parts that do not depend on  $m_{n_i}$ , and therefore they vanish because of the Glashow-Iliopoulos-Maiani mechanism.

The divergent cancellation in the total  $\Delta_{L,R}$  is shown as follows.

$$\begin{aligned}
\text{div} \left[ \Delta_L^{(1)W} \right] &= m_a \Delta_\epsilon \times \frac{3}{2} \times \sum_{i=1}^9 U_{ai}^{\nu*} U_{bi}^\nu m_{n_i}^2, \\
\text{div} \left[ \Delta_L^{(5)W} \right] &= m_a \Delta_\epsilon \times \sum_{i,j=1}^9 U_{ai}^{\nu*} U_{bj}^\nu \left( -D_{ij}^* m_{n_j}^2 - \frac{1}{2} D_{ij} m_{n_i}^2 \right), \\
\text{div} \left[ \Delta_L^{(7+8)W} \right] &= \text{div} \left[ \Delta_L^{(4)Y} \right] = \text{div} \left[ \Delta_L^{(7+8)Y} \right] = 0, \\
\text{div} \left[ \Delta_L^{(1)Y} \right] &= m_a \Delta_\epsilon \times \left( \frac{3s_\theta^4}{2c_\beta^2} \right) \sum_{i=1}^9 U_{(a+3)i}^\nu U_{(b+3)i}^\nu m_{n_i}^2, \\
\text{div} \left[ \Delta_L^{(2)Y} \right] &= m_a \Delta_\epsilon \times \left( -\frac{c_\theta s_\theta^2}{2c_\beta} \right) \sum_{i=1}^9 U_{(a+3)i}^\nu \lambda_{bi}^{L,1} m_{n_i}, \\
\text{div} \left[ \Delta_L^{(3)Y} \right] &= \Delta_\epsilon \times \left( \frac{c_\theta s_\theta^2}{c_\beta} \right) \sum_{i=1}^9 U_{(a+3)i}^{\nu*} L_{ai}^{R,1} m_{n_i}^2, \\
\text{div} \left[ \Delta_L^{(5)Y} \right] &= m_a \Delta_\epsilon \times \frac{s_\theta^2}{c_\beta^2} \sum_{i,j=1}^9 U_{(a+3)i}^{\nu*} U_{(b+3)j}^\nu \left( -D_{ij}^* m_{n_j}^2 - \frac{1}{2} D_{ij} m_{n_i}^2 \right), \\
\text{div} \left[ \Delta_L^{(6)Y h_k^\pm} \right] &= m_a \Delta_\epsilon \times \sum_{i,j=1}^9 U_{(a+3)i}^{\nu*} \lambda_{ij}^{0*} \lambda_{bj}^{L,k}, \\
\text{div} \left[ \Delta_L^{(9+10)Y h_k^\pm} \right] &= -m_a \Delta_\epsilon \times \sum_{i=1}^9 U_{(a+3)i}^{\nu*} \lambda_{bi}^{L,k} m_{n_i}, \tag{A1}
\end{aligned}$$

where:  $\text{div} B_0^{(1)} = \text{div} B_0^{(2)} = \text{div} B_0^{(12)} = -2 \text{div} B_1^{(1)} = -2 \text{div} B_1^{(2)} = \Delta_\epsilon$  and  $1/m_Y = s_\theta / (c_\beta m_W)$ .

Easy to see that  $\text{div} \left[ \Delta_L^{(1)W} \right] + \text{div} \left[ \Delta_L^{(5)W} \right] = \text{div} \left[ \Delta_L^{(6)Y h_k^\pm} \right] + \text{div} \left[ \Delta_L^{(9+10)Y h_k^\pm} \right] = 0$  and



the sum of the remaining divergent parts is zero in case we are focusing on investigating  $c_\theta = 1$ .

- 
- [1] Y. Fukuda *et al.* [Super-Kamiokande], Phys. Rev. Lett. **81** (1998), 1562-1567 [arXiv:hep-ex/9807003 [hep-ex]].
  - [2] S. Fukuda *et al.* [Super-Kamiokande], Phys. Rev. Lett. **86** (2001), 5651-5655 [arXiv:hep-ex/0103032 [hep-ex]].
  - [3] S. Fukuda *et al.* [Super-Kamiokande], Phys. Rev. Lett. **86** (2001), 5656-5660 [arXiv:hep-ex/0103033 [hep-ex]].
  - [4] Q. R. Ahmad *et al.* [SNO], Phys. Rev. Lett. **89** (2002), 011301 [arXiv:nucl-ex/0204008 [nucl-ex]].
  - [5] Q. R. Ahmad *et al.* [SNO], Phys. Rev. Lett. **89** (2002), 011302 [arXiv:nucl-ex/0204009 [nucl-ex]].
  - [6] L. Morel, Z. Yao, P. Cladé and S. Guellati-Khélifa, Nature **588**, no.7836, 61-65 (2020) doi:10.1038/s41586-020-2964-7.
  - [7] B. Aubert *et al.* [BaBar], Phys. Rev. Lett. **104**, 021802 (2010) [arXiv:0908.2381 [hep-ex]].
  - [8] E. Kou *et al.* [Belle-II], PTEP **2019**, no.12, 123C01 (2019) [erratum: PTEP **2020**, no.2, 029201 (2020)] [arXiv:1808.10567 [hep-ex]].
  - [9] T. Aushev, W. Bartel, A. Bondar, J. Brodzicka, T. E. Browder, P. Chang, Y. Chao, K. F. Chen, J. Dalseno and A. Drutskoy, *et al.* [arXiv:1002.5012 [hep-ex]].
  - [10] A. M. Baldini *et al.* [MEG II], Eur. Phys. J. C **78**, no.5, 380 (2018) [arXiv:1801.04688 [physics.ins-det]].
  - [11] A. M. Sirunyan *et al.* [CMS], JHEP **06** (2018), 001.
  - [12] G. Aad *et al.* [ATLAS], Phys. Lett. B **800**, 135069 (2020) [arXiv:1907.06131 [hep-ex]].
  - [13] [ATLAS], ATLAS-CONF-2019-037.
  - [14] Q. Qin, Q. Li, C. D. Lü, F. S. Yu and S. H. Zhou, Eur. Phys. J. C **78** (2018) no.10, 835 [arXiv:1711.07243 [hep-ph]].
  - [15] A. Zee, Phys. Lett. B **93**, 389 (1980) [erratum: Phys. Lett. B **95**, 461 (1980)] doi:10.1016/0370-2693(80)90349-4
  - [16] R. K. Barman, R. Dcruz and A. Thapa, JHEP **03**, 183 (2022) [arXiv:2112.04523 [hep-ph]].

- [17] E. Arganda, A. M. Curiel, M. J. Herrero and D. Temes, *Phys. Rev. D* **71** (2005), 035011.
- [18] X. Marcano and R. A. Morales, *Front. in Phys.* **7**, 228 (2020) doi:10.3389/fphy.2019.00228 [arXiv:1909.05888 [hep-ph]].
- [19] A. Ilakovac, *Phys. Rev. D* **62** (2000), 036010.
- [20] E. Arganda, M. J. Herrero, X. Marcano and C. Weiland, *Phys. Rev. D* **91** (2015) no.1, 015001.
- [21] E. Arganda, M. J. Herrero, X. Marcano, R. Morales and A. Szyrkman, *Phys. Rev. D* **95** (2017) no.9, 095029.
- [22] N. H. Thao, L. T. Hue, H. T. Hung and N. T. Xuan, *Nucl. Phys. B* **921**, 159-180 (2017), arXiv:1703.00896 [hep-ph].
- [23] G. Hernández-Tomé, J. I. Illana and M. Masip, *Phys. Rev. D* **102**, no.11, 113006 (2020) [arXiv:2005.11234 [hep-ph]].
- [24] T. P. Nguyen, T. T. Thuc, D. T. Si, T. T. Hong and L. T. Hue, *PTEP* **2022**, 023 (2022), arXiv:2011.12181 [hep-ph].
- [25] A. Brignole and A. Rossi, *Phys. Lett. B* **566** (2003), 217-225.
- [26] A. Brignole and A. Rossi, *Nucl. Phys. B* **701** (2004), 3-53.
- [27] J. L. Diaz-Cruz, *JHEP* **05** (2003), 036.
- [28] P. T. Giang, L. T. Hue, D. T. Huong and H. N. Long, *Nucl. Phys. B* **864** (2012), 85-112.
- [29] M. Arana-Catania, E. Arganda and M. J. Herrero, *JHEP* **09** (2013), 160 [erratum: *JHEP* **10** (2015), 192].
- [30] L. T. Hue, H. N. Long, T. T. Thuc and T. Phong Nguyen, *Nucl. Phys. B* **907** (2016), 37, arXiv:1512.03266 [hep-ph].
- [31] E. Arganda, M. J. Herrero, R. Morales and A. Szyrkman, *JHEP* **03** (2016), 055.
- [32] E. Arganda, M. J. Herrero, X. Marcano and C. Weiland, *Phys. Rev. D* **93** (2016) no.5, 055010.
- [33] M. Zeleny-Mora, J. L. Díaz-Cruz and O. Félix-Beltrán, [arXiv:2112.08412 [hep-ph]].
- [34] B. Yang, J. Han and N. Liu, *Phys. Rev. D* **95** (2017) no.3, 035010.
- [35] J. Herrero-García, T. Ohlsson, S. Riad and J. Wirén, *JHEP* **04** (2017), 130.
- [36] H. K. Guo, Y. Y. Li, T. Liu, M. Ramsey-Musolf and J. Shu, *Phys. Rev. D* **96** (2017) no.11, 115034.
- [37] M. Aoki, S. Kanemura, K. Sakurai and H. Sugiyama, *Phys. Lett. B* **763** (2016), 352-357.

- [38] K. Cheung, W. Y. Keung and P. Y. Tseng, Phys. Rev. D **93** (2016) no.1, 015010.
- [39] K. Huitu, V. Keus, N. Koivunen and O. Lebedev, JHEP **05** (2016), 026.
- [40] C. H. Chen and T. Nomura, Eur. Phys. J. C **76** (2016) no.6, 353.
- [41] C. F. Chang, C. H. V. Chang, C. S. Nugroho and T. C. Yuan, Nucl. Phys. B **910** (2016), 293-308.
- [42] W. Altmannshofer, S. Gori, A. L. Kagan, L. Silvestrini and J. Zupan, Phys. Rev. D **93** (2016) no.3, 031301.
- [43] Y. Omura, E. Senaha and K. Tobe, Phys. Rev. D **94** (2016) no.5, 055019.
- [44] A. Lami and P. Roig, Phys. Rev. D **94** (2016) no.5, 056001.
- [45] D. Das and A. Kundu, Phys. Rev. D **92** (2015) no.1, 015009.
- [46] A. Crivellin, G. D'Ambrosio and J. Heeck, Phys. Rev. Lett. **114** (2015), 151801.
- [47] M. D. Campos, A. E. Cárcamo Hernández, H. Päs and E. Schumacher, Phys. Rev. D **91** (2015) no.11, 116011.
- [48] Y. Omura, E. Senaha and K. Tobe, JHEP **05** (2015), 028.
- [49] L. de Lima, C. S. Machado, R. D. Matheus and L. A. F. do Prado, JHEP **11** (2015), 074.
- [50] J. Heeck, M. Holthausen, W. Rodejohann and Y. Shimizu, Nucl. Phys. B **896** (2015), 281-310.
- [51] I. Doršner, S. Fajfer, A. Greljo, J. F. Kamenik, N. Košnik and I. Nišandžić, JHEP **06** (2015), 108.
- [52] X. G. He, J. Tandean and Y. J. Zheng, JHEP **09** (2015), 093.
- [53] A. Dery, A. Efrati, Y. Nir, Y. Soreq and V. Susič, Phys. Rev. D **90** (2014), 115022.
- [54] A. Celis, V. Cirigliano and E. Passemar, Phys. Rev. D **89** (2014), 013008.
- [55] A. Falkowski, D. M. Straub and A. Vicente, JHEP **05** (2014), 092.
- [56] R. Harnik, J. Kopp and J. Zupan, JHEP **03** (2013), 026.
- [57] P. S. Bhupal Dev, R. Franceschini and R. N. Mohapatra, Phys. Rev. D **86** (2012), 093010.
- [58] A. Goudelis, O. Lebedev and J. h. Park, Phys. Lett. B **707** (2012), 369-374.
- [59] J. L. Diaz-Cruz and J. J. Toscano, Phys. Rev. D **62** (2000), 116005.
- [60] J. G. Korner, A. Pilaftsis and K. Schilcher, Phys. Rev. D **47** (1993), 1080-1086.
- [61] A. Pilaftsis, Z. Phys. C **55** (1992), 275-282.
- [62] A. Pilaftsis, Phys. Lett. B **285** (1992), 68-74.
- [63] J. Herrero-Garcia, N. Rius and A. Santamaria, JHEP **11** (2016), 084.

- [64] G. Blankenburg, J. Ellis and G. Isidori, Phys. Lett. B **712** (2012), 386-390.
- [65] T. Aoyama, N. Asmussen, M. Benayoun, J. Bijnens, T. Blum, M. Bruno, I. Caprini, C. M. Carloni Calame, M. Cè and G. Colangelo, *et al.* Phys. Rept. **887**, 1-166 (2020) doi:10.1016/j.physrep.2020.07.006 [arXiv:2006.04822 [hep-ph]].
- [66] B. Abi *et al.* [Muon g-2], Phys. Rev. Lett. **126**, no.14, 141801 (2021) doi:10.1103/PhysRevLett.126.141801 [arXiv:2104.03281 [hep-ex]].
- [67] G. W. Bennett *et al.* [Muon g-2], Phys. Rev. D **73** (2006), 072003 [arXiv:hep-ex/0602035 [hep-ex]].
- [68] S. Borsanyi, Z. Fodor, J. N. Guenther, C. Hoelbling, S. D. Katz, L. Lellouch, T. Lippert, K. Miura, L. Parato and K. K. Szabo, *et al.* Nature **593** (2021) no.7857, 51-55 doi:10.1038/s41586-021-03418-1 [arXiv:2002.12347 [hep-lat]].
- [69] Z. N. Zhang, H. B. Zhang, J. L. Yang, S. M. Zhao and T. F. Feng, Phys. Rev. D **103** (2021) no.11, 115015 doi:10.1103/PhysRevD.103.115015 [arXiv:2105.09799 [hep-ph]].
- [70] S. Baek and K. Nishiwaki, Phys. Rev. D **93** (2016) no.1, 015002 doi:10.1103/PhysRevD.93.015002 [arXiv:1509.07410 [hep-ph]].
- [71] R. Foot, H. N. Long and T. A. Tran, Phys. Rev. D **50** (1994) no.1, R34-R38 doi:10.1103/PhysRevD.50.R34 [arXiv:hep-ph/9402243 [hep-ph]].
- [72] H. N. Long, Phys. Rev. D **54** (1996), 4691-4693 doi:10.1103/PhysRevD.54.4691 [arXiv:hep-ph/9607439 [hep-ph]].
- [73] H. N. Long, Phys. Rev. D **53** (1996), 437-445 doi:10.1103/PhysRevD.53.437 [arXiv:hep-ph/9504274 [hep-ph]].
- [74] L. T. Hue, H. T. Hung, N. T. Tham, H. N. Long and T. P. Nguyen, Phys. Rev. D **104** (2021) no.3, 033007 doi:10.1103/PhysRevD.104.033007 [arXiv:2104.01840 [hep-ph]].
- [75] H. B. Zhang, T. F. Feng, S. M. Zhao, Y. L. Yan and F. Sun, Chin. Phys. C **41** (2017) no.4, 043106 doi:10.1088/1674-1137/41/4/043106 [arXiv:1511.08979 [hep-ph]].
- [76] T. P. Nguyen, T. T. Le, T. T. Hong and L. T. Hue, Phys. Rev. D **97** (2018) no.7, 073003 doi:10.1103/PhysRevD.97.073003 [arXiv:1802.00429 [hep-ph]].
- [77] A. E. Cárcamo Hernández, L. T. Hue, S. Kovalenko and H. N. Long, Eur. Phys. J. Plus **136** (2021) no.11, 1158 doi:10.1140/epjp/s13360-021-02146-9 [arXiv:2001.01748 [hep-ph]].
- [78] H. T. Hung, N. T. Tham, T. T. Hieu and N. T. T. Hang, PTEP **2021** (2021) no.8, 083B01 doi:10.1093/ptep/ptab082 [arXiv:2103.16018 [hep-ph]].

- [79] J. C. Montero, F. Pisano and V. Pleitez, *Phys. Rev. D* **47** (1993), 2918-2929 doi:10.1103/PhysRevD.47.2918 [arXiv:hep-ph/9212271 [hep-ph]].
- [80] M. Singer, J. W. F. Valle and J. Schechter, *Phys. Rev. D* **22** (1980), 738 doi:10.1103/PhysRevD.22.738
- [81] P. H. Frampton, *Phys. Rev. Lett.* **69** (1992), 2889-2891 doi:10.1103/PhysRevLett.69.2889
- [82] F. Pisano and V. Pleitez, *Phys. Rev. D* **46** (1992), 410-417 doi:10.1103/PhysRevD.46.410 [arXiv:hep-ph/9206242 [hep-ph]].
- [83] M. Lindner, M. Platscher and F. S. Queiroz, *Phys. Rept.* **731** (2018), 1-82 doi:10.1016/j.physrep.2017.12.001 [arXiv:1610.06587 [hep-ph]].
- [84] A. S. De Jesus, S. Kovalenko, F. S. Queiroz, C. Siqueira and K. Sinha, *Phys. Rev. D* **102** (2020) no.3, 035004 doi:10.1103/PhysRevD.102.035004 [arXiv:2004.01200 [hep-ph]].
- [85] Á. S. de Jesus, S. Kovalenko, F. S. Queiroz, C. A. de S. Pires and Y. S. Villamizar, *Phys. Lett. B* **809** (2020), 135689 doi:10.1016/j.physletb.2020.135689 [arXiv:2003.06440 [hep-ph]].
- [86] A. E. C. Hernández, D. T. Huong and I. Schmidt, *Eur. Phys. J. C* **82** (2022) no.1, 63 doi:10.1140/epjc/s10052-022-10011-x [arXiv:2109.12118 [hep-ph]].
- [87] L. T. Hue, K. H. Phan, T. P. Nguyen, H. N. Long and H. T. Hung, [arXiv:2109.06089 [hep-ph]].
- [88] S. M. Boucenna, J. W. F. Valle and A. Vicente, *Phys. Rev. D* **92** (2015) no.5, 053001 doi:10.1103/PhysRevD.92.053001 [arXiv:1502.07546 [hep-ph]].
- [89] D. Chang and H. N. Long, *Phys. Rev. D* **73** (2006), 053006 doi:10.1103/PhysRevD.73.053006 [arXiv:hep-ph/0603098 [hep-ph]].
- [90] A. J. Buras, F. De Fazio, J. Girrbach and M. V. Carlucci, *JHEP* **02** (2013), 023 doi:10.1007/JHEP02(2013)023 [arXiv:1211.1237 [hep-ph]].
- [91] H. K. Dreiner, H. E. Haber and S. P. Martin, *Phys. Rept.* **494** (2010), 1-196 doi:10.1016/j.physrep.2010.05.002 [arXiv:0812.1594 [hep-ph]].
- [92] L. Ninh and H. N. Long, *Phys. Rev. D* **72** (2005), 075004 doi:10.1103/PhysRevD.72.075004 [arXiv:hep-ph/0507069 [hep-ph]].
- [93] L. T. Hue, L. D. Ninh, T. T. Thuc and N. T. T. Dat, *Eur. Phys. J. C* **78** (2018) no.2, 128 doi:10.1140/epjc/s10052-018-5589-3 [arXiv:1708.09723 [hep-ph]].
- [94] H. Okada, N. Okada, Y. Orikasa and K. Yagyu, *Phys. Rev. D* **94** (2016) no.1, 015002 doi:10.1103/PhysRevD.94.015002 [arXiv:1604.01948 [hep-ph]].

- [95] H. T. Hung, T. T. Hong, H. H. Phuong, H. L. T. Mai and L. T. Hue, Phys. Rev. D **100** (2019) no.7, 075014 doi:10.1103/PhysRevD.100.075014 [arXiv:1907.06735 [hep-ph]].
- [96] A. Crivellin, M. Hoferichter and P. Schmidt-Wellenburg, Phys. Rev. D **98** (2018) no.11, 113002 doi:10.1103/PhysRevD.98.113002 [arXiv:1807.11484 [hep-ph]].
- [97] F. Jegerlehner and A. Nyffeler, Phys. Rept. **477** (2009), 1-110 doi:10.1016/j.physrep.2009.04.003 [arXiv:0902.3360 [hep-ph]].
- [98] A. Denner, S. Heinemeyer, I. Puljak, D. Rebuszi and M. Spira, Eur. Phys. J. C **71** (2011), 1753 doi:10.1140/epjc/s10052-011-1753-8 [arXiv:1107.5909 [hep-ph]].
- [99] P. A. Zyla *et al.* [Particle Data Group], PTEP **2020** (2020) no.8, 083C01 doi:10.1093/ptep/ptaa104
- [100] K. Abe *et al.* [T2K], Nature **580** (2020) no.7803, 339-344 [erratum: Nature **583** (2020) no.7814, E16] doi:10.1038/s41586-020-2177-0 [arXiv:1910.03887 [hep-ex]].
- [101] M. Tanabashi *et al.* [Particle Data Group], Phys. Rev. D **98** (2018) no.3, 030001 doi:10.1103/PhysRevD.98.030001
- [102] G. 't Hooft and M. J. G. Veltman, Nucl. Phys. B **44** (1972), 189-213 doi:10.1016/0550-3213(72)90279-9
- [103] A. Denner and S. Dittmaier, Nucl. Phys. B **734** (2006), 62-115 doi:10.1016/j.nuclphysb.2005.11.007 [arXiv:hep-ph/0509141 [hep-ph]].
- [104] T. Hahn and M. Perez-Victoria, Comput. Phys. Commun. **118** (1999), 153-165 doi:10.1016/S0010-4655(98)00173-8 [arXiv:hep-ph/9807565 [hep-ph]].
- [105] K. H. Phan, H. T. Hung and L. T. Hue, PTEP **2016** (2016) no.11, 113B03 doi:10.1093/ptep/ptw158 [arXiv:1605.07164 [hep-ph]].