

High angular resolution near-IR view of the Orion Bar revealed by Keck/NIRC2

Emilie Habart¹, Romane Le Gal^{2,3}, Carlos Alvarez⁴, Els Peeters^{5,6,7}, Olivier Berné⁸, Mark G. Wolfire⁹, Javier R. Goicoechea¹⁰, Thiébaud Schirmer¹¹, Emeric Bron¹², and Markus Röllig¹³

¹ Université Paris-Saclay, CNRS, Institut d'Astrophysique Spatiale, 91405 Orsay, France

² Institut de Planétologie et d'Astrophysique de Grenoble (IPAG), Université Grenoble Alpes, CNRS, F-38000 Grenoble, France

³ Institut de Radioastronomie Millimétrique (IRAM), 300 rue de la piscine, F-38406 Saint-Martin d'Hères, France

⁴ W.M. Keck Observatory, 65-1120 Mamalahoa Hwy, Kamuela, HI, USA

⁵ Department of Physics & Astronomy, The University of Western Ontario, London ON N6A 3K7, Canada

⁶ Institute for Earth and Space Exploration, The University of Western Ontario, London ON N6A 3K7, Canada

⁷ Carl Sagan Center, SETI Institute, 339 Bernardo Avenue, Suite 200, Mountain View, CA 94043, USA

⁸ Institut de Recherche en Astrophysique et Planétologie, Université de Toulouse, CNRS, CNES, UPS, 9 Av. du colonel Roche, 31028 Toulouse Cedex 04, France

⁹ Astronomy Department, University of Maryland, College Park, MD 20742, USA

¹⁰ Instituto de Física Fundamental (IFF), CSIC, Calle Serrano 121-123, 28006, Madrid, Spain

¹¹ Department of Space, Earth and Environment, Chalmers University of Technology, Onsala Space Observatory, 439 92 Onsala, Sweden

¹² LERMA, Observatoire de Paris, PSL Research University, CNRS, Sorbonne Universités, F-92190 Meudon, France

¹³ I. Physikalisches Institut der Universität zu Köln, Zùlpicher Straße 77, 50937 Köln, Germany

2022

ABSTRACT

Context. Nearby Photo-Dissociation Regions (PDRs), where the gas and dust are heated by the far UV-irradiation emitted from stars, are ideal templates to study the main stellar feedback processes.

Aims. With this study we aim to probe the detailed structures at the interfaces between ionized, atomic, and molecular gas in the Orion Bar. This nearby prototypical strongly irradiated PDR will be among the first targets of the James Webb Space Telescope (JWST) within the framework of the PDRs4All Early Release Science program.

Methods. We employed the sub-arcsec resolution accessible with Keck-II NIRC2 and its adaptive optics system to obtain the most detailed and complete images, ever performed, of the vibrationally excited line H_2 1-0 S(1) at $2.12 \mu\text{m}$, tracing the dissociation front, and the [FeII] and Br γ lines, at 1.64 and $2.16 \mu\text{m}$ respectively, tracing the ionization front. The former is a powerful tracer of the FUV radiation field strength and gas density distribution at the PDR edge, while the last two trace the temperature and density distribution from the ionized gas to the PDR. We obtained narrow-band filter images in these key gas line diagnostic over $\sim 40''$ at spatial scales of $\sim 0.1''$ (~ 0.0002 pc or ~ 40 AU at 414 pc).

Results. The Keck/NIRC2 observations spatially resolve a plethora of irradiated sub-structures such as ridges, filaments, globules and propyls. This portends what JWST should accomplish and how it will complement the highest-resolution ALMA maps of the molecular cloud. A remarkable spatial coincidence between the H_2 1-0 S(1) vibrational and HCO^+ J=4-3 rotational emission previously obtained with ALMA is observed. This likely indicates the intimate link between these two molecular species and highlights that in high pressure PDR the H/H_2 and $C^+/C/CO$ transitions zones come closer as compared to a typical layered structure of a constant density PDR. The H/H_2 dissociation front appears as a highly structured dense region with over-dense sub-structures with a typical thickness of a few $\sim 10^{-3}$ pc. This is in agreement with several previous studies that claimed that the Orion Bar edge is composed of very small, dense, highly irradiated PDRs at high thermal pressure immersed in a more diffuse environment.

Key words. <photodissociation regions - interstellar medium - infrared emission - star forming regions>

1. Introduction

Stellar feedback influences the star formation activity through molecular cloud dissolution, compression processes and increase in ionization fraction, which slows down ambipolar diffusion. However, the precise nature and importance of all these mechanisms relative to other processes that influence star formation remains largely unknown. Stellar feedback can take the form of radiative heating and pressure, thermal expansion and evaporation, and stellar winds and shocks (e.g., Pabst et al. 2020; Schneider et al. 2020, and reference therein). The combined radiative and dynamical feedback of massive stars on their parental

cloud is not yet understood. Therefore, direct observation and quantification of the radiative heating efficiency (e.g., the coupling of gas and dust to the intense stellar-ultraviolet fields) and the mechanical energy injection is necessary.

Recent near-infrared images obtained with the Gemini telescope at high angular resolution ($\sim 0.06 - 0.11''$) revealed, at a spectacular level of detail, unexpected structures within the strongly irradiated molecular cloud at the Western Wall in Carina at a distance of 2.3 kpc (Hartigan et al. 2020). These authors found series of ridges, fragments, and waves with sizes in the ~ 200 - 2000 AU range that may result from the complex

dynamical and radiative processes that sculpt these systems. Indeed, while regularly-spaced ridges that run parallel to the photodissociation front may suggest large-scale magnetic fields are dynamically-important (Mackey & Lim 2011), several fragments and waves could resemble Kelvin-Helmholtz instabilities. Berné et al. (2010) reported the presence of a series of five surprisingly regular wavelets (separated by 0.1 pc or 20000 AU) at the surface of the Orion molecular cloud. These authors proposed that these waves formed by the mechanical interaction of high-velocity plasma/gas, produced by massive stars, with the dense molecular gas, which has provoked hydrodynamical instabilities. Sub-arcsecond resolution is required to resolve detailed structures even for the nearest regions of massive star formation. This capability now exists with adaptive optics (AO) imaging, and will be further developed/expanded in the near future with JWST.

The main goal of the present study is to observe and quantify the stellar feedback in the closest site of ongoing massive star-formation: the Orion molecular cloud. This region is located about 5 times closer than the Carina star forming region. The dominant stellar feedback processes can be probed by observations of Photo-Dissociation Regions (PDRs) where far-ultraviolet photons of massive stars create warm regions of gas and dust in the neutral atomic and molecular gas (e.g., for a recent review see Wolfire et al. 2022, and references therein). Nearby, edge-on PDRs such as the Orion Bar are ideal targets due to their wide extension on the sky, bright emission, and proximity that allow to probe these processes on a small physical scale. PDRs are particularly bright in the infrared (IR).

However, until now a very high angular (sub-arcsec resolution) IR molecular view of the Orion Bar PDR was missing. In this study, we present observations obtained with Keck/NIRC2 using AO, which provide the most complete and detailed maps of the complex UV-irradiated region where the conversion from ionized to atomic to molecular gas occurs. We mapped the vibrationally excited line of H_2 at $2.12 \mu\text{m}$, tracing the dissociation front, and the [FeII] line at $1.64 \mu\text{m}$ and the $\text{Br}\gamma$ line at $2.16 \mu\text{m}$, tracing the ionization front. The proximity of the region¹ (414 pc) combined with recent advances in instrumentation allows for the first time to conduct studies from the (sub-)parsec scales to the smallest structures of ~ 100 AU. Observations with AO provide nearly diffraction-limited images in the near-IR over a field of view of $1'$ in size.

The paper is organized as follows. The target the Orion Bar is described in Sect. 2. The observations and data reduction are described in Sect. 3. In Sect. 4, the spatial morphology of the line intensities is discussed and compared to previous observations. In Sects. 5 and 6, the observed line intensities are corrected for dust extinction and compared to the model predictions. Finally, a short summary and prospects for JWST are given in Section 7.

2. The Orion Bar

The Orion Bar, a strongly ultraviolet (UV) irradiated PDR, is an escarpment of the Orion molecular cloud (OMC), the closest site of ongoing massive star formation. The Bar is illuminated by the O7-type star θ^1 Ori C, the most massive and luminous member of the Trapezium cluster at the heart of the Orion Nebula (e.g., O'Dell 2001). The trapezium cluster creates a blister $H\text{ II}$ region that is eating its way into the parental cloud. A large cavity

has been carved out of the molecular gas and the inner concave structure tilts to form the Orion Bar (Wen & O'Dell 1995; O'Dell 2001). Weillbacher et al. (2015) presented MUSE/VLT integral spectroscopic data of the central part of the Orion Nebula with an angular resolution of ~ 0.7 to $1.2''$. They derived an extinction map and estimated physical properties (electron temperature and density) of the Orion Nebula HII region. The Far-UV (FUV) radiation field incident on the Orion Bar PDR is $G_0 = 1 - 4 \times 10^4$ in Habing units (e.g., Marconi et al. 1998). Beyond the ionization front (IF), where the gas converts from fully ionized to fully neutral, only FUV photons with energies below 13.6 eV penetrate the cloud. This corresponds to the edge of the PDR. Many previous works have studied the Orion Bar PDR's spatial stratification. The first PDR layers are predominantly neutral and atomic: $[H] > [H_2] \gg [H^+]$. They display NIR atomic emission lines from low ionization potential elements (Walmsley et al. 2000) as well as carbon and sulfur radio recombination lines (Wyrowski et al. 1997; Cuadrado et al. 2019; Goicoechea & Cuadrado 2021). This warm and moderately dense gas (n_H of a few 10^4 cm^{-3}) is mainly cooled by the very bright FIR $[C^+]$ $158 \mu\text{m}$ and $[O^0]$ 63 and $145 \mu\text{m}$ fine-structure lines (Tielens et al. 1993; Herrmann et al. 1997; Bernard-Salas et al. 2012; Ossenkopf et al. 2013). The atomic PDR zone also hosts the peak of the Mid-IR aromatic particles emission (e.g., Bregman et al. 1989; Sellgren et al. 1990; Tielens et al. 1993; Giard et al. 1994; Knight et al. 2021; Schirmer et al. in prep).

At about $10\text{--}15''$ (or $0.02\text{--}0.03$ pc) from the IF (at $A_V \approx 0.5\text{--}2$ mag of visual extinction into the neutral cloud), the dissociating FUV photons are sufficiently attenuated and most of the hydrogen becomes molecular² (e.g., van der Werf et al. 2013). This corresponds to the dissociation front (DF) where the H/H_2 transition takes place. The DF displays a forest of IR rotational and vibrationally excited H_2 lines (e.g., Parmar et al. 1991; Luhman et al. 1994; van der Werf et al. 1996; Allers et al. 2005; Shaw et al. 2009; Zhang et al. 2021), including FUV-pumped levels up to $v = 12$ (Kaplan et al. 2017, 2021) and HD rotational lines (Wright et al. 1999; Joblin et al. 2018). FUV-pumped H_2 drives the formation of molecular ions such as CH^+ and SH^+ through reactions of C^+ and S^+ with vibrationally excited (e.g., Goicoechea et al. 2017; Parikka et al. 2017; Lehmann et al. 2022, and references therein). Analysis of the IR H_2 and 21 cm H I lines suggests warm temperatures ($T_k \approx 400\text{--}700 \text{ K}$) at the DF. Using stationary PDR models (i.e. PDR models with stationary chemistry and without dynamics), the gas density in the atomic gas of $n_H = 4 - 5 \times 10^4 \text{ cm}^{-3}$ is consistent with the [OI] and [CII] line emission and the separation between the IF and DF (Tielens et al. 1993; Hogerheijde et al. 1995; Marconi et al. 1998).

The IF and DF separation depends also on the dust extinction. Larger grains than found in diffuse ISM that have lower FUV absorption cross-sections need to be invoked in isobaric PDR models in order to be consistent with the observed IF and DF separation of $\sim 10\text{--}15''$ (e.g., Allers et al. 2005). Reduction in the FUV attenuation cross-section is in agreement with the recent work of Schirmer et al. (in prep) modeling dust emission and extinction throughout the Orion Bar PDR. Analyzing the Spitzer and Herschel observations with the radiative transfer code SOC (Juvella 2019), together with the dust THEMIS model (Jones et al. 2013, 2017), Schirmer et al. (in prep) found that nano-grains are in fact strongly depleted compared to diffuse ISM (by a factor ≥ 10), leading to a FUV attenuation reduction,

¹ The most commonly adopted distance to the Bar is 414 pc (Menten et al. 2007) although more recent observations, including Gaia, point to slightly lower values (Kounkel et al. 2017; Großschedl et al. 2018).

² This range of extinction value is only value for high G_0/n_H PDRs like the Orion Bar, but not in lower G_0/n_H PDRs where the H/H_2 transitions occur at much lower A_V due to the H_2 self-shielding.

and a minimum grain size that is larger than in the diffuse ISM. In their model, the gas density is estimated to be about $4 \times 10^4 \text{ cm}^{-3}$ at the mid-IR aromatic nano-dust emission peak, and about $1.5 \times 10^5 \text{ cm}^{-3}$ on average between the mid-IR emission peak and the H_2 emission peak. This is consistent with previous estimates.

Beyond the DF, between $A_V = 2$ and 4 mag, the $\text{C}^+/\text{C}/\text{CO}$ transition takes place (Tauber et al. 1995) and the PDR becomes molecular. Some observations and models suggest the presence of relatively large (0.01-0.02 pc) high density clumps ($n_{\text{H}} = 10^6\text{--}10^7 \text{ cm}^{-3}$) in the molecular PDR (e.g., Burton et al. 1990; van der Werf et al. 1996; Young Owl et al. 2000; Lis & Schilke 2003; Andree-Labsch et al. 2017). These over-dense components must be embedded in a lower density medium mainly responsible for the extended PDR emission. Alternatively, the observed H_2 and high-J CO lines emission close to the DF may be explained by a roughly isobaric PDR, at high thermal-pressure $P_{\text{th}}/k \approx 10^8 \text{ cm}^{-3} \text{ K}$ (e.g., Allers et al. 2005; Joblin et al. 2018). We note that high density and pressure appear as a common feature in high FUV field PDRs, as demonstrated by a number of observations (e.g., Ossenkopf et al. 2010; Sheffer et al. 2011; Köhler et al. 2015; Wu et al. 2018). In the high pressure isobaric model, a constant pressure from the atomic to the warm molecular region results in an atomic region and a very narrow emission for the warm molecular tracers. In the Joblin et al. (2018) model with $P=2.8 \times 10^8 \text{ cm}^{-3} \text{ K}$ the atomic region had a size of $\sim 3.5''$ while the emission zone of the H_2 (1-0) S(1) line has a width of $\sim 0.25''$. The predicted atomic region width is nevertheless significantly smaller than the one observed ($\sim 10\text{--}15''$) and the very narrow molecular emission is unresolved in most tracers until now.

While most of the previous molecular studies relied on modest angular resolution observations, ALMA has provided $\sim 1''$ resolution images of the molecular emission (Goicoechea et al. 2016). Instead of an homogeneous PDR with well-defined and spatially separated H/H_2 and $\text{C}^+/\text{C}/\text{CO}$ transition zones, ALMA revealed rich small-scale ($\sim 0.004 \text{ pc}$) over-dense structures (akin to filaments and globules), sharp edges, and bright emission from an embedded proplyd (object 203-506; Champion et al. 2017). These observations have challenged the traditional view of PDRs (and their models, e.g., Kirsanova & Wiebe 2019) and revealed a steeply varying interface between the atomic and the molecular gas (Fig. 1c, Goicoechea et al. 2016). The observed over-dense substructures may have been induced by UV radiation-driven compression (Gorti & Hollenbach 2002; Tremblin et al. 2012). Advection of the molecular gas through the DF has been suggested by CO emission observed by ALMA in the form of globules or plumes apparently extending into the atomic gas (Goicoechea et al. 2016).

Recent FIR observations of the dust emission polarization from the Orion Bar PDR suggest a relatively modest plane-of-the-sky magnetic field strength of $B_0 \approx 300 \mu\text{G}$ (e.g., Chuss et al. 2019; Guerra et al. 2021). Hence, magnetic pressure of $3 \times 10^7 \text{ K cm}^{-3}$ (comparable to the ambient and inter-clump thermal pressure of $1\text{--}3 \times 10^7 \text{ K cm}^{-3}$) may also play a role in the PDR dynamics (e.g., Pellegrini et al. 2009; Pabst et al. 2020).

3. Observations and data reduction

3.1. Observations

On December 4, 2020 (Program ID N004), we used the NIRC2 wide field camera in combination with the Keck-II AO Laser Guide Star (LGS) system to observe a representative Field-of-View (FoV) of $40'' \times 40''$ of the Orion Bar centered on




Fig. 1: Hubble Space Telescope WFC2 mosaic of the Orion Bar (Credit: NASA/STScI/Rice Univ./C.O'Dell et al. - Program ID: PRC95-45a). The NIRC2 wide camera FoV is shown with the cyan square. The yellow square zooms into the two proplyds lying in the targeted FoV.




Fig. 2: The PanSTARRS (z band) image showing the NIRC2 wide camera FoV (cyan square, $40'' \times 40''$) centered on RA=05 : 35 : 20.662, DEC=-05 : 25 : 19.38 (J2000). The green circles enclose tip-tilt stars that can be used to perform the LGS observations.

RA=05:35:20.662, DEC=-05:25:19.38 (J2000) with a pixel size of $0.04''$, as illustrated in Figs. 1 and 2. This instrumental setup provides the optimal combination of FoV and diffraction-limited imaging capabilities to fulfill our science goals. The FoV was rotated by 52.5° (North to East). Observations were obtained in the narrow-band filters $\text{H}_2 v = 1 - 0$ ($\lambda_c = 2.128 \mu\text{m}$, $\Delta\lambda = 0.034 \mu\text{m}$), $[\text{FeII}]$ ($\lambda_c = 1.645 \mu\text{m}$, $\Delta\lambda = 0.026 \mu\text{m}$) and $\text{Br } \gamma$ ($\lambda_c = 2.168 \mu\text{m}$, $\Delta\lambda = 0.0326 \mu\text{m}$). We also obtained observations in the Kcont ($\lambda_c = 2.271 \mu\text{m}$, $\Delta\lambda = 0.033 \mu\text{m}$) and Hcont ($\lambda_c = 1.580 \mu\text{m}$, $\Delta\lambda = 0.023 \mu\text{m}$) filters to subtract the

continuum from the $H_2 v = 1 - 0$, Br γ , and [FeII] line maps respectively.

Fig. 2 shows the different point sources within $60''$ of the center of the region that can be used as tip-tilt stars for the LGS system. The executed observation sequence was a 5-point dither pattern using a $\sim 1''$ dither step with $H_2 v = 1 - 0$ and Kcont consecutive images on each dither position. The same sequence was executed for the [FeII] and Hcont filters, and the Br γ and Kcont filters. The exposure time per image was 150s ($15s \times 10$ coadds) in multiple-correlated double-sampling mode with 8 reads per coadd.

We integrated 750s on-source for each filter. The observing strategy was successful although it was a bit challenging due to the variable nebular background on the tip-tilt star. In fact, the nebular background can significantly increase the effective magnitude of the tip-tilt star, which in turn can degrade the AO correction. In an environment with such a complex background as the Orion Bar, the automatic AO acquisition sequence may need to be modified to manually take a background for the tip-tilt sensor away from the nebular region. Such manual background acquisition reduces the efficiency of the AO observations. We took images in the $H_2 v = 1 - 0$ and Kcont filters with V377 Ori as the tip-tilt star. However, due to the high nebular background, we used θ^2 Ori A as the tip-tilt star for the [FeII]/Hcont and Br- γ /Kcont sequences. The factor of ~ 2 larger separation between θ^2 Ori A and the science FoV with respect to V377 Ori was compensated by the fact that θ^2 Ori A is 7.5 magnitudes brighter on the wave front sensor. Our observations clearly benefited from good seeing conditions and AO correction (K-band Strehl ratio $\sim 30\%$). The full width at half maximum (FWHM) of the point-spread function (PSF) was $\sim 0.11''$. For comparison, JWST will achieve a resolution of $0.085''$ in the K-band. In summary, these unusual bright extended source observations for NIRC2 were successful.

3.2. Data reduction

The basic data reduction of the imaging frames (i.e., dark correction, flat field, sky subtraction) were performed using standard procedures for near-IR imaging with a modified version of the reduction pipeline from the UCLA Galactic Center group based on Pyraf (Lu et al. 2009). The set of frames slightly offset in telescope position were realigned and co-added to produce the final image. We calculated star centroids and offsets from one frame to the other using the IRAF task IMALIGN in order to line up the images. Thus, the images were co-added by weighted median which suppresses the ghosts and the diffraction spike due to the bright θ^2 Ori A star. The final images are of dimension of $38.64'' \times 38.88''$ or $0.08 \text{ pc} \times 0.08 \text{ pc}$ at 414 pc.

In order to overcome the telescope pointing error (which is of the order of $0.5''$) and determine the proper World Coordinate System (WCS), we matched the stars in our NIRC2 image to Gaia coordinates. We used the Gaia catalog data early release 3 (<https://www.cosmos.esa.int/web/gaia>).

For the flux calibration, we determined the conversion factor between the detector counts and physical units using previous observations of the $H_2 1-0 \text{ S}(1)$ line obtained with NTT/SOFI (Walmsley et al. 2000). We convolved the continuum-subtracted Keck/NIRC2 H_2 line emission to the beam of the NTT/SOFI observation (seeing limited $\sim 1''$). We estimated the flux conversion factor in regions of the maps with high signal to noise ratio. Thus, the reduced science maps (in Analog Digital Unit per second) were multiplied by this conversion factor to get the flux-calibrated science maps in $\text{erg cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$. We assigned

an accuracy of about 25% to the measurements of the line intensity. The same conversion factor was used for the $H_2 v = 1 - 0$, Br γ and [FeII] lines maps since they were obtained with almost the same weather conditions and the detector response is similar. We compared the fluxes of each of the lines we obtained with those measured by Walmsley et al. (2000) through the PDR (see their position C, Table 1) and found a good agreement considering the different spatial resolution. Maps that are not continuum-subtracted were divided by the filter width and are in units $\text{erg cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1} \mu\text{m}^{-1}$. We must underline that we could have fluxed-calibrated the observations using the photometric standard FS-7 observed in the H and Ks filters. However, we would obtained similar uncertainties due to the sky background variations between the standard star observations and the science observations, which was significant in the K band.

4. Spatial distribution

In this section, the spatial morphology of the hydrogen Brackett γ , iron and molecular hydrogen lines are presented and compared to previous observations.

4.1. Br γ and iron lines

The maps obtained in the Br γ and [FeII] filters and the continuum-subtracted emission are shown in Fig. 3, while the H_2 filter maps are shown in Fig. 4. Most of the structures seen in these lines and the continuum are very similar to those visible in the $H\alpha$ HST map (Fig. 1). The ionization front is clearly seen in the Br γ and [FeII] line emission maps. However, there are some differences between the Br γ and [FeII] line emission morphologies. In particular, in the area surrounding the two identified proplyds near the Bar, 203-506 and 203-504 (Bally et al. 2000 and located in the yellow squares in Fig. 1): both proplyds emit in [FeII] along with bright streamers, whereas only the proplyd 203-504 shows strong Br γ emission as further described in Section 4.5.

The surface brightness profiles measured in the [FeII], Br γ and H_2 lines along 4 cuts perpendicular to the Bar are shown in Fig. 5. These cuts use a 3-pixel average corresponding to the typical FWHM of the PSF. Cut 1 goes through the left edge of bar, cut 2 through the position of the JWST/NIRSpec-IFU and MIRI-IFU mosaics of the ERS program (see Fig. 3 in Berné et al. 2022), cuts 3 and 4 through the right edge of the bar (see Fig. 5). To investigate the stratification of the interfaces from ionized to molecular emission, we also constructed averaged emission cut over the entire FoV and perpendicular to the Orion Bar. As expected for a PDR seen edge-on, we found that the H_2 emission is displaced further into the cloud than the Br γ and [FeII] line emission. A precise determination of the distance between the IF delineated by the [FeII] and Br γ lines and the DF delineated by the H_2 line can be obtained. From the averaged emission cuts, we estimated that the offset between the IF and the main peak of the H_2 vibrational emission is between 12 and $12.5''$ (0.024 – 0.025 pc). Finally, the [FeII] and Br γ surface brightness profiles also show extended emission towards the PDR. This emission may originate in the flattened region beyond the bar that is still illuminated directly by the ionizing star (Wen & O'dell 1995; O'Dell 2001).




Fig. 3: Integrated intensity maps of the [FeII] and continuum-subtracted [FeII] line emission (left column) and of the Bry and continuum-subtracted Bry line emission (right column). Units are $\text{erg cm}^{-2} \text{s}^{-1} \text{sr}^{-1} \mu\text{m}^{-1}$ for the [FeII] and Bry maps and $\text{erg cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$ for the continuum-subtracted line emission map. The horizontal dashed lines indicate the average position of the ionization and dissociation fronts (as determined from the averaged emission cut over the entire FoV presented in Fig. 5).

4.2. Vibrationally excited H_2 line

The maps obtained in the H_2 2.12μm filter and the continuum-subtracted H_2 line emission are shown in Fig. 4 and also in Fig. 6 for comparison with the ALMA $\text{HCO}^+ J = 4 - 3$ line emission map (see Section 4.4). In the H_2 line emission map, several bright H_2 emission peaks are spatially resolved and show small scale structures. The H/H_2 dissociation front appears highly structured with several ridges and extremely sharp filaments with a width of 1 to 2'' (0.002-0.004 pc or 400-800 AU, see Figs. 4 and 5). Ridges and filaments run parallel to the dissociation front but a succession of bright sub-structures in H_2 is also observed from the edge of the DF towards the molecular region. The H_2 emission peaks appear in an area that starts at about 10'' from the IF and up to 20'' and even 25-30'' in some places from the IF (see Figs. 4 and 5). The bright sub-structures result in several peaks in the H_2 brightness profiles shown in Fig. 5 obtained along the cuts perpendicular to the Orion Bar. The peaks are roughly spaced by a few arcseconds (~ 0.005 -

0.01 pc or ~ 1000 -2000 AU). The surface area occupied by the bright sub-structures (ridges, filaments), i.e. with H_2 line intensity $> 2 \times 10^{-4} \text{ erg s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$, corresponds to about 40-45% of the H_2 emission zone in the Bar.

Since the H_2 emission is very sensitive to both the FUV radiation field and the gas density (as explained below), these very narrow and bright sub-structures must be due to dense material directly irradiated. Towards more tenuous material, the H_2 emission is spatially much more extended and weaker. The multiple H_2 emission peaks along and across the Bar may be associated with a multitude of small highly irradiated and dense PDRs. A single edge-on dense PDR produces one narrow H_2 emission peak. The slight tilt of the Bar along the line of sight combined with the presence of over-dense structures may explain the observed complex emission distribution, and especially the succession of H_2 peaks across the Bar.

The intensity variations in the different H_2 emission peaks ranging from ~ 2 to $\sim 10 \times 10^{-4} \text{ erg s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$ (Fig. 5) may result from a combination of effects due to the local gas densi-




Fig. 4: Same as for Fig. 3 for the H₂ 1-0 S(1) line emission.

ties, geometry and dust extinction effects. Density effects must be important. In fact, PDR models show that for the conditions prevailing in the Orion Bar, the intensity of the H₂ $v=1-0$ S(1) line is approximately proportional to the gas density (e.g., Burton et al. 1990). At equilibrium between the formation of H₂ and the

photodissociation of H₂ by FUV flux, the intensity of the 1-0 S(1) line is proportional to $R_f n_H N(H_0)$ where R_f is the H₂ formation rate, n_H is the total hydrogen gas density and $N(H_0)$ the column density of atomic H atoms from the PDR edge. For low values of n_H/G_0 ($< 40 \text{ cm}^{-3}$), which is the case in the atomic




Fig. 5: Top row: Superposition of the four cuts perpendicular to the Orion Bar described in Sect. 4.1 on the [FeII] $1.64\mu\text{m}$ and H_2 $2.12\mu\text{m}$ continuum subtracted emission line maps. Cuts 1, 2, 3, 4 are positioned from left to right. The squares area are used to compute the lines intensity in the key regions presented in Table 1. Second and third rows: H_2 (blue), Bry (red) and [FeII] (green) line intensity (in $\text{erg s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$) profiles shown as function of the distance from the IF for each cut along with the HCO^+ line integrated intensity profile (in K km s^{-1} , grey). The intensity profiles of the Bry and [FeII] lines have been scaled by the factors indicated in the legends. The line intensities have not been corrected for dust extinction. Bottom row: The averaged emission cut calculated by taking the mean over the entire FoV and perpendicular to the Orion Bar. The vertical dashed lines indicate the average position of the IF, DF and HCO^+ peak.

zone of the Orion Bar with $G_0 \sim 10^4$ and $n_{\text{H}} = 10^4 - 10^5 \text{ cm}^{-3}$, the H^0/H_2 is driven by dust opacity and $\text{N}(\text{H}^0)$ is a constant equal to a few 10^{21} cm^{-2} . The intensity is thus linearly proportional to the density. Moreover, the H_2 and $\text{HCO}^+ \text{ J}=4-3$ emission show a remarkable similar spatial distribution as shown in Fig. 6 and further described in Sect. 4.4. This rotational line is a good indicator of dense gas (Goicoechea et al. 2016).

The dense sub-structures seen in H_2 may be surrounded by a lower-density gas component producing an extended emission. In the H_2 data, a more widespread and extended emission (with an intensity of about $2 \times 10^{-4} \text{ erg s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$) seems to be in fact observed at the photodissociation front at about $10-15''$ ($0.02-0.03 \text{ pc}$) and up to $20-30''$ ($0.04-0.06 \text{ pc}$) from the IF. Moreover, the H_2 map and brightness profiles show some emission in front of the DF (see Fig. 5 and Table 1). Part of this emission likely originates from the surface of the Orion molecular cloud-1 (OMC1). This surface is perpendicular to the line of sight and is

illuminated by the Trapezium cluster, making it a face-on PDR. The emission from this background face-on PDR was also observed with Herschel in other PDR tracers, especially in high-J CO lines (Parikka et al. 2018) and [OI] 63 and $145 \mu\text{m}$ and [CII] $158 \mu\text{m}$ (Bernard-Salas et al. 2012). An increase in the H_2 emission at the ionisation front is also observed. This was also visible in the previous data of Walmsley et al. (2000) and could originate from the background molecular cloud.

4.3. Comparison with previous H_2 observations

Our Keck observations convolved at the angular resolution of $\sim 1''$ of the previous SOFI/NTT observations by Walmsley et al. (2000) show good agreement in terms of line distribution and intensity (see Table 1). It can be noted that in the SOFI/NTT observations, the brightest sub-structures were already observed but not spatially resolved. The TEXES observations of the pure ro-

tational H_2 0-0 S(1) and S(2) lines by Allers et al. (2005), with a resolution of $\sim 2''$ cover the south-west part of our map. The comparison of these observations with the H_2 1-0 S(1) line map obtained previously by van der Werf et al. (1996) show that the spatial distribution of the pure rotational lines and the vibrationally excited line agree in remarkable detail (see Fig. 2 in Allers et al. 2005). This is predicted by high pressure models (Allers et al. 2005; Joblin et al. 2018), where a separation between the H_2 lines is expected to be very small ($\leq 0.5''$). Thus, one would expect to observe an overall spatial coincidence on the whole Keck map. Nevertheless, detailed variations are expected because the pure rotational lines that result from collisional excitation have a different dependence on the local physical conditions. Future JWST-ERS observations (Berné et al. 2022) will allow such an investigation in the parts of our maps observed with the IFU-MIRI spectroscopy.

4.4. Comparison between H_2 and HCO^+ from ALMA

In this section, the distribution and correlation of the H_2 line emission with the ALMA observation of the HCO^+ J=4-3 line are analysed. Considering the position of the compact sources (e.g. proplyds 203-506), the ALMA observations showed an offset to our Keck observations in coordinates of R.A.=-0.15" and DEC.=0.75". By correcting this offset, a good overlap of the position of the compact sources and of the Bar is obtained.

In Fig. 6, we compare the maps obtained in the H_2 1-0 S(1) and HCO^+ J=4-3 lines across the same field of view. The middle and bottom figures zoom into two parts of the map to be able to compare in detail the spatial distribution of the emission from each of the detected sub-structures. Most of the sub-structures are common to both maps and show a very similar distribution. Due to a high-dipole moment and a high critical density, the HCO^+ 4-3 line (n_{critical} of a few 10^6 cm^{-3}) is a good proxy of the gas density (Goicoechea et al. 2016). Its integrated intensity is roughly proportional to the density in the $n_{\text{H}} = 10^4 - 10^6 \text{ cm}^{-3}$ range. The average density that reproduces the mean HCO^+ towards the dissociation front (at about $15''$ from the IF) is about $n_{\text{H}} = 0.5 - 1.5 \cdot 10^6 \text{ cm}^{-3}$. Thus, some of the densest portions of the Bar lie along the dissociation front. We note that the small H_2 and HCO^+ structures discussed here localized at the DF are shifted by a about $\sim 20''$ relative to the bigger ($5''$ - $10''$) condensations seen more inside the molecular cloud (Young Owl et al. 2000; Lis & Schilke 2003).

Although there is a very good spatial coincidence between the H_2 and HCO^+ sub-structures, the H_2 line emission decreases faster in the Bar than the HCO^+ line emission (see Fig. 5). This can be explained as follows. Firstly, the vibrationally excited H_2 line is more sensitive to the FUV field flux and the most intense H_2 emission must come from the transition region between the diffuse and very dense medium where the FUV radiation is not yet attenuated. The FUV shielding produced by the ridge of high-density sub-structures may significantly decrease the vibrational excited H_2 in the deeper regions. Secondly, extinction due to dust in the Orion Bar and to foreground dust may affect the apparent morphology of the near-IR H_2 images. Small-scale dust extinction differences could in fact result in morphological differences between the near-IR and millimeter-wave images. Dust extinction effects on the line intensities is discussed in Sect. 5.

The overall remarkable spatial coincidence between the H_2 and HCO^+ line emission shows that they both come from high-densities but also indicates their strong chemical link. Detection of both bright HCO^+ and CO emission by ALMA towards the H_2 vibrational emission layers (Fig. 2c in Goicoechea et al. 2016)

suggests that the C^+/CO transition nearly coincides with the H/H_2 transition. This fact was predicted in Joblin et al. (2018), where the best fitting Orion Bar model has a separation between these transitions of $\sim 0.25''$. We obtained a precise determination of the average offset between the peak of the H_2 vibrational emission (delineating the H/H_2 transition) and the edge of the observed HCO^+ emission (delineating the C^+/CO transition). From the averaged emission cuts, we estimated that this average offset is less than $1''$, about $\sim 0.6''$ (or 0.0012 pc). However considering that the angular resolution of the ALMA data is $\sim 1''$, higher angular resolution observations are required to measure the exact value of this offset. The presence of a large quantity of irradiated H_2 may have important consequences for the chemical structure. Due to the high densities and enhanced H_2 gas heating via formation and collisional de-excitation of FUV pumped levels, the chemistry is triggered by high temperature and FUV pumped levels. Endothermic reactions and reactions with energy barriers become faster (Agúndez et al. 2010; Nagy et al. 2013). Reactions of H_2 with abundant atoms and ions, e.g. C^+ , shift the carbon bearing molecular gas towards the cloud edge.

We note that although a spatial coincidence is observed between the H_2 and HCO^+ over-dense sub-structures, a spatial shift of the emission peak is measured for several sub-structures (e.g., sub-structure 3 in the middle panels and sub-structures 2, 3 in the bottom panels of Fig. 6). This may essentially result from the fact that the two emission lines vary differently with the local conditions (FUV field flux and gas density) and that the H_2 near-IR line is affected by dust extinction. The most important differences in the spatial distribution of the H_2 and HCO^+ emission are found in the atomic region. In fact, the brighter structures seen in the atomic regions in HCO^+ (like globulets) are faintly visible in H_2 ro-vibrational emission (e.g., the sub-structures denoted 1 and 6 in the middle panels and the sub-structure denoted 1 in the bottom panels of Fig. 6). These structures are probably located in the background or foreground. Indeed, the emission velocity of these HCO^+ structures ($v_{\text{LSR}} = 8-9 \text{ km s}^{-1}$) is more consistent with emission from the background OMC1 than from the Bar ($v_{\text{LSR}} = 10.5 \text{ km s}^{-1}$). Some extended H_2 line emission seen in the atomic zone to the south-west part of the Bar (see bottom panel of Fig. 6) has also no counterpart in the HCO^+ or CO line emission maps. Finally, plume-like CO features seen with ALMA in the atomic zone by Goicoechea et al. (2016) which may be advected from the surface of the molecular cloud have no evident correspondences with the H_2 emission structures observed by Keck/NIRC2.

4.5. Proplyds and collapsing protostars

We are able to spatially resolve the FUV irradiated Orion Bar PDR edge but also additional compact sources corresponding to over-dense irradiated sub-structures such as proplyds near the IF, and perhaps collapsing protostars near the DF. In the following, we briefly describe the observations towards these sources.

4.5.1. Proplyds

Fig. 7 shows proplyds 203-504 and 203-506 (Bally et al. 2000) which are present in the FoV of our Keck/NIRC2 observations. The star of 203-504 is detected in all filters, however the disk/envelope is not clearly identifiable in these images. The 203-506 proplyd is clearly visible in emission in both the H_2 and [FeII] lines, whereas it is not detected in the Bry line.




Fig. 6: Left panels: Integrated intensity maps (continuum-subtracted) of the H_2 1-0 S(1) line emission in $\text{erg cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$. Right panels: Integrated intensity maps of the HCO^+ J=4-3 lines in K km s^{-1} .

The H_2 emission appears to have a similar morphology to the [OI] 6300 Å line observed by Bally et al. (2000). This is expected since both lines trace warm neutral gas (~ 1000 K) near the dissociation front, inside the dense PDR that is created at the disk surface (Champion et al. 2017). The [FeII] line shows a very different morphology, with an elongated structure perpen-

dicular to the proplyd disk, and is reminiscent of a jet, such as those identified for proplyds by Bally et al. (2015). The [FeII] emission would then be associated with irradiated shocked gas. The absence of $\text{Br}\gamma$ emission could have different explanations. The coming PDR4alls JWST data should bring additional clues to better understand it.

4.5.2. Over-dense sub-structures

Numerous over-dense small structures (1-2'') appear in the H₂ and HCO⁺ emission that could be self-gravitating, or transient, turbulently compressed features or compressed by FUV photoevaporation (Gorti & Hollenbach 2002; Tremblin et al. 2012). Recently, Röllig & Ossenkopf-Okada (2022) showed that the intensity levels of the HCO⁺ emission map as well as its spatial distribution is consistent with predictions from nonstationary, clumpy PDR model ensembles. Whether they can become star-forming clumps (for example, by merging into massive clumps) is uncertain. Gravitational collapse appears not apparent from their density distribution (no high-density power-law tail, Goicoechea et al. 2016). In fact, the mass of a cylinder with a density $n_H \sim 10^6 \text{ cm}^{-3}$, a width of 1'' and a length of 2''-6'' is $< 0.005 M_\odot$ (Goicoechea et al. 2016), which is much lower than the virial and critical masses needed to make them gravitationally unstable (approximately $5 M_\odot$ from the inferred gas temperature, density, and velocity dispersion, Inutsuka & Miyama 1997). However, the very bright substructure located in the northeast of our map and denoted clump in Fig. 6 shows a very particular structure (with a size of $\sim 1000 \text{ AU}$), likely a YSO candidate (Goicoechea et al. in prep).

5. Line intensities and extinction correction

5.1. Line intensities in the key zones

In Table 1, we give the intensities of the [FeII], Br γ , and H₂ lines measured towards the ionized region, IF, PDR atomic region, DF, and from the molecular region. The line intensities of each of these characteristic regions have been estimated in the 2'' by 2'' square areas shown in Fig. 5 (top panels). The areas for the atomic, DF, and molecular zones which are located at about 5.5'', 12.5'', 21'' from the IF, are positioned within the FoV of the NIRSpec-IFU and MIRI-IFU mosaics (see Fig. 3 of Berné et al. 2022) which are along the direction of cut 2 in Fig. 5. On the other hand, the areas chosen for the ionized and IF regions are not aligned along the JWST mosaics (i.e. cut 2) because the Keck maps do not entirely cover these ionized gas and IF regions along the JWST mosaic (i.e. cut 2). We therefore chose areas for the ionized gas and IF regions further south-west (towards the direction of cuts 3 and 4, see Fig. 5).

Comparison with previous measurements of the intensities of these lines with SOFI/NTT by Walmsley et al. (2000) in the ionized, IF, and DF regions are given in Table 1. The measurements are in good agreement within a factor of about 2 for the [FeII] and Br γ lines at the IF and the H₂ line at the DF. However, we find some larger differences for Br γ in the HII and DF regions and H₂ in the HII and IF regions. This might be mostly due to the fact that the position regions of Walmsley et al. (2000) (shown as A, B, C in their Fig. 1) do not correspond to our positions. Their HII and IF regions are outside of the Keck/NIRC2 field-of-view.

5.2. Extinction correction

Corrections for extinction due to the foreground dust and internal dust in the Bar itself of the line intensities were estimated and reported in Table 1. The foreground visual extinction A_V produced by the Veil was estimated to be not greater than about 2 mag by O'Dell & Yusef-Zadeh (2000) using a radio-to-optical surface brightness comparison. Using flux measurements of the Balmer lines (H α to H β flux ratio), Weillbacher et al. (2015) es-

timated that the foreground extinction is about $A_V \sim 1.3 \text{ mag}$ (or $A_K \sim 0.15 \text{ mag}$) towards the ionized gas in the line of sight of the Orion Bar (see their Fig. 15). The foreground dust structure appears smooth and devoid of significant substructure. By this dust screen, the line emission will be attenuated by $e^{-\tau_\lambda}$ with $\tau_\lambda = 0.921 A_\lambda$ the optical depth at the wavelength of the line. To adopt a dust reddening appropriate to Orion, we follow Goicoechea et al. (2015) and used the extinction curve of Cardelli et al. (1989) as refined by Blagrave et al. (2007) with $R_V = A_V/E(B-V) = 5.5$. The reddening factors are 0.216, 0.143 and 0.138 at 1.64, 2.12 and 2.16 μm respectively. Given this degree of foreground extinction, we estimated that the [FeII], Br γ , and H₂ lines are approximately 23%, 15%, and 16% brighter respectively if foreground extinction correction is taken into account.

For the H₂ line in the PDR, internal extinction due to the dust in the Bar itself may significantly attenuate the emission. An additional magnitude of extinction due to dust in the Bar between the ionized gas and the region of excited H₂ may result in a stronger increase in line intensity. In order to estimate this extinction, we assumed that the column density of gas along the line of sight is equal to the column density between the IF and the region of excited H₂. Thus, we used the density profile towards the Bar as constrained from comparison between Spitzer and Herschel observations and a radiative transfer code with the THEMIS dust model (Schirmer et al. in prep). The corresponding calculated column density is $1.5 \cdot 10^{21} \text{ H cm}^{-2}$ from the IF to the beginning of the H₂ emission (at 10'' or 0.02 pc from the IF) and $10 \cdot 10^{21} \text{ H cm}^{-2}$ from the IF to the end of the H₂ emission (at 20'' or 0.04 pc from the IF). Adopting the same dust reddening as described before from Cardelli et al. (1989) and $A_V/N_H = 3.5 \times 10^{-22} \text{ mag cm}^{-2}$, we estimated an extinction of $A_V = 1$ to 4 or $A_K = 0.13$ to 0.54. If this internal dust extinction correction is taken into account and a correction factor of $e^{-\tau_\lambda}$ is applied, the H₂ line is approximately 12 to 40% brighter. Regarding the assumed extinction law, the Cardelli extinction curve with $R_V = 5.5$ is similar to that calculated by the THEMIS dust model in the Orion bar with nano-grains depletion (Schirmer et al. in prep). For this model, the reddening factors are 0.224, 0.139 and 0.134 at 1.64, 2.12 and 2.16 μm respectively. The correction factor on the line intensities assuming their extinction law will differ by less than 3%.

We must underline that if we assume that the gas and the dust are well mixed in the H₂ emission zone, one must apply a correction factor of $(1 - e^{-\tau_\lambda})/\tau_\lambda$ (e.g. Thronson et al. 1990). Assuming a visual extinction of $A_V = 3$ for the total H₂ emission zone (including the succession of H₂ peaks) as calculated using the dust density model, we computed the corresponding correction factor. This visual extinction is equivalent to consider a gas density of $1\text{-}1.5 \cdot 10^5 \text{ cm}^{-3}$ and a length of 0.02 pc (as the width of the H₂ emission zone in Fig. 5). This visual extinction corresponding to a large column density of UV pumped H₂ can be considered as an upper limit. With these assumptions, the line is approximately 18% brighter after extinction correction in the H₂ emission zone. Considering the increase of 12% due to the extinction correction of the atomic zone from the IF to the beginning of the H₂ emission zone, the line will thus be 30% brighter. This is relatively close to what we found previously assuming no mixing which is of 40% (due to the internal extinction correction from the IF to the end of the H₂ emission zone).

In Table 1 and for the comparison to the model predictions in the following section, we assumed that the H₂ line will be in total 56% brighter after correcting for foreground (16%) and internal extinction (40%) in the Bar. Our total (foreground and internal)

extinction corrections for H_2 are thus about $A_K = 0.17 + 0.54 = 0.71$ which is very close to the one derived by Kaplan et al. (2021) of $A_K = 0.72 \pm 0.1$. These authors used an effective way to measure extinction which consists of comparing the observed to theoretical H_2 ro-vibrational line flux ratios from pairs of lines arising from the same upper level that are widely separated in wavelength. Their errors on A_K were estimated from the assumptions made on the color correction. Their IGRINS observation at high spectral resolution were made along a slit positioned across the Bar and centered on at the southwest of the Bar (corresponding to the position of the sub-structure 5 in the bottom zoom of Fig. 6). However, we must underline that their extinction measurement is lower than the values derived by Luhman et al. (1998) also using H_2 ro-vibrational transitions with common upper levels but observed a long a slit passing roughly through the center of our Keck maps. Luhman et al. (1998) found averaged values of $A_K = 2.6 \pm 0.7$ and 2.3 ± 0.8 respectively for the H_2 peak and beyond. The H_2 line will be approximately 90% brighter if an extinction correction of $A_K = 2.6$ (or $A_V = 19$) is taken into account. The internal extinction might be variable depending on the sightline and the density of the region crossed. From the radio and NIR H_2 line maps, Walmsley et al. (2000) suggest in fact that extinction can vary rapidly as a function of position in the Bar. Specifically, precise spatial estimates of the internal PDR extinction are needed. This will be possible with the high angular resolution near-IR line maps that will be obtained with JWST's NIRCAM and NIRSpec IFUs. This will constrain in detail how dust extinction affects the apparent morphology of the H_2 line emission.

6. Comparison to model predictions

To guide our interpretation, in this section, we compare our extinction-corrected observations of the emission gas lines to the model predictions in the HII region, the IF, the atomic region, the DF, and the molecular zone. The model predictions are those we used to make the template spectra described in Berné et al. (2022) and to predict line intensities for the JWST Exposure Time Calculator. The gas line intensities have been computed using (i) the Cloudy code for the ionized gas in the H II region and the ionization front (Ferland et al. 2017); and (ii) the Meudon PDR code for the contribution from the atomic and molecular lines in the neutral PDR gas (Le Petit et al. 2006). The parameters and calculation requirements used for each model and region are described in detail in Berné et al. (2022). In the following subsections, only some computational results important for the comparison predictions/observations are given.

In this comparison with model predictions, our primary goal was to ensure that the models reproduce the observed intensity peak of the ionised and neutral gas lines emission by Keck. This allowed us to adjust the required JWST integration time in order

Table 1: Observed and predicted line intensities in $\text{erg cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$ in the ionized, IF, atomic, DF, and molecular regions. For Keck observations, the median and standard deviation of the intensities observed in the $2''$ by $2''$ square areas shown in Fig. 5 are given. The intensity peak of the lines is also given. The observed Keck intensities have been corrected for dust extinction and the correction factor applied is given in % (in the row "Corr. Extinction"). Measurements of the intensities of these lines with SOFI/NTT by Walmsley et al. (2000) in the ionized, IF, and DF regions (corresponding to their A, B, and C regions in their Table 1) are also reported.

Line	λ (μm)	A (s^{-1})	E_u (K)	Intensities				
				ionized	IF	atomic	DF	molecular
[FeII]	1.64440	0.006	11445.9					
			Obs. Keck	$3.6\pm 0.7 \cdot 10^{-4}$	$7.0\pm 1.6 \cdot 10^{-4}$	$1.7\pm 0.6 \cdot 10^{-4}$	$1.1\pm 0.6 \cdot 10^{-4}$	$0.7\pm 0.6 \cdot 10^{-4}$
			Max. Intensity	$6 \cdot 10^{-4}$	$1.2 \cdot 10^{-3}$			
			Corr. Extinction	+23%	+23%	+23%	+23%	+23%
			Obs. NTT	$4.7 \cdot 10^{-4}$	$6.0 \cdot 10^{-4}$		$1.2 \cdot 10^{-4}$	
			Cloudy model	$1.9 \cdot 10^{-7}$	$6 \cdot 10^{-4}$	-	-	-
H ₂	2.12183	$3.47 \cdot 10^{-7}$	6951.3					
			Obs. Keck	$1.8\pm 0.6 \cdot 10^{-4}$	$2.2\pm 0.7 \cdot 10^{-4}$	$0.46\pm 0.46 \cdot 10^{-4}$	$4.7\pm 2.3 \cdot 10^{-4}$	$1\pm 0.8 \cdot 10^{-4}$
			Max. Intensity				$13.6 \cdot 10^{-4}$	
			Corr. Extinction	+ 16%	+16%	+16%	+56%	+56%
			Obs. NTT	$0.3 \cdot 10^{-4}$	$0.7 \cdot 10^{-4}$		$3.0 \cdot 10^{-4}$	
			Meudon model	-	-	$0.1 \cdot 10^{-4}$	$22 \cdot 10^{-4}$	$3.7 \cdot 10^{-9}$
Br γ	2.16612	$3.045 \cdot 10^5$	154582.7					
			Obs. Keck	$15.4\pm 0.1 \cdot 10^{-4}$	$26.3\pm 0.3 \cdot 10^{-4}$	$9.2\pm 0.8 \cdot 10^{-4}$	$4.8\pm 0.8 \cdot 10^{-4}$	$2.6\pm 0.8 \cdot 10^{-4}$
			Max. Intensity	$20.9 \cdot 10^{-4}$	$32.4 \cdot 10^{-4}$			
			Corr. Extinction	+15%	+15%	+15%	+15%	+15%
			Obs. NTT	$36.6 \cdot 10^{-4}$	$39.4 \cdot 10^{-4}$		$11.7 \cdot 10^{-4}$	
			Cloudy model	$4.7 \cdot 10^{-4}$	$36 \cdot 10^{-4}$	-	-	-

to get a high signal-to-noise ratio and avoid saturation problems, a challenging aspect for JWST observation of bright targets. We emphasize that our model predictions were obtained by making simple assumptions and by separately estimating the ionized and neutral molecular gas lines. The radiative transfer and the thermal balance were not calculated continuously from the HII region to the molecular region. We have not attempted to reproduce the separation between the IF and the DF.

6.1. Predicted $\text{Br}\gamma$ and [FeII] lines emission

The input parameters of Pellegrini et al. (2009) and Shaw et al. (2009) were adopted for the Cloudy model of the Orion Bar. This is an isobaric model with initial electron density of $n_e \sim 3500 \text{ cm}^{-3}$ and temperature of $T_e \sim 9000 \text{ K}$. The electron density reaches a maximum of $n_e \sim 6600 \text{ cm}^{-3}$ before the IF and drops sharply at the IF. At the IF, the total proton density increases to $n = 28,000 \text{ cm}^{-3}$ while the electronic temperature decreases to $T_e \sim 4000 \text{ K}$. The $\text{Br}\gamma$ $2.16 \mu\text{m}$ line emissivity is spatially extended in the HII region, following the electronic density profile, and then decreases strongly when H^+ becomes H^0 at the IF. On the other hand, the [FeII] $1.64 \mu\text{m}$ line emissivity peaks strongly at the IF.

In Table 1, we report the predicted intensities of the lines. The [FeII] $1.64 \mu\text{m}$ and $\text{Br}\gamma$ $2.16 \mu\text{m}$ lines model predictions are in good agreement with the observations at the IF front (by a factor less than 1.4). On the other hand, the emission of these lines in the HII region are significantly lower in the model predictions. This difference applies for several other ionized gas lines

predicted by Cloudy. The observations may overlap the IF where the observed intensities are bright.

6.2. Predicted H_2 line emission

For the Meudon code, we considered an isobaric model with a thermal pressure $P = 4 \cdot 10^8 \text{ K cm}^{-3}$. This model exhibits gas physical conditions expected near the dissociation front of the Orion Bar (e.g., Goicoechea et al. 2016; Joblin et al. 2018). We fixed the radiation field impinging on the PDR so that, at the edge of the PDR, $G_0 = 2 \cdot 10^4$ in Mathis units. We adopted the extinction curve of HD 38087 of ? and $R_V = 5.62$ which is close to the value determined for Orion Bar of 5.5 (Marconi et al. 1998). The model includes an exact radiative transfer calculation for the UV pumping lines originating from the first 30 levels of H_2 , while the other lines are treated using the FGK approximation. This allows to account for mutual shielding effects between overlapping H_2 lines and can significantly affect the position of the H/H_2 transition. The H_2 1-0 S(1) line emissivity peaks near the H/H_2 transition layer where the gas density is of the order of few 10^5 cm^{-3} . As in Joblin et al. (2018), the model does not reproduce the observed width of the atomic region, i.e. the distance between the position of the IF and the DF. It is significantly smaller than in the observation. We note that as suggested by Allers et al. (2005), a PDR model with a lower gas pressure in the atomic zone (i.e., a lower gas density) and a reduction of the FUV dust cross section will better match the observed spatial position of the H_2 emission peak. In the Schirmer et al. (in prep) dust model with nano-grain depletion (i.e., FUV extinction reduction), the visual extinction integrated from the PDR edge to

the observed H_2 emission peak position at 12-12.5'' (or 0.024-0.025 pc) is of $A_V \sim 1$. The required model medium density in the atomic zone is of $5 \times 10^4 \text{ cm}^{-3}$, close to the inter-clump density.

We calculated the cumulative line intensities from the atomic and H^0/H_2 transition region ($0 < A_V < 0.4$ and $0.4 < A_V < 2.5$) and from the molecular region (which starts at the C/CO transition, $2.5 < A_V < 8.5$). The optically thin line surface brightness are enhanced by a geometrical factor of $1/\cos(\theta)=2$ relative to the face-on surface brightness. The viewing angle θ between the line-of-sight and the normal to the PDR is taken equal to $\sim 60^\circ$ which gives an approximation of a nearly edge-on PDR and is the maximum inclination that can be used to derive line intensities in the 1D PDR Meudon code. The uncertainty on this angle could lead to an additional scaling factor on the line intensity.

The H_2 2.12 μm line intensity as predicted by the model in the DF is in agreement with the measured peak intensity at the DF by a factor of 1.6 (too high for the model). This could partly be due to a geometrical factor but also to the fact that the isobaric model considered here with a high pressure of $P = 4 \times 10^8 \text{ K cm}^{-3}$ gives a too high density and over-estimates the H_2 1-0 S(1) line (see section 6.3). The H_2 1-0 S(1) line emissivity width predicted by the model is of $\sim 0.0006 \text{ pc}$ or $\sim 0.3''$. This is lower than the observed thickness of the sharpest H_2 filaments (see Fig. 5). The observed width of the bright H_2 ridge (about $1''$ to $2''$) may be increased by geometrical effects or a lower thermal pressure.

In the atomic zone, the model predicts a much lower intensity by a factor of ~ 10 with respect to the DF because the H_2 density is much lower. This is in agreement with the data but the large uncertainties on the measured intensity make a quantitative comparison difficult. In the molecular zone, the predicted intensities are extremely low by a factor $> 10^4$ with respect to the DF. The measured intensity decreases much less. It is comparable to the intensity measured in the HII region divided by a factor of 2. The observed H_2 emission in the molecular zone must come from the flattened region behind the main ionization front on the far side of the Orion Bar.

6.3. Thermal pressure and dynamic effects

Considering the TEXES H_2 0-0 S(1) line observation with an angular resolution of $2''$ (Allers et al. 2005) and the 1-0 S(1) line Keck observation convolved at the same resolution and corrected for extinction, the 1-0 S(1) / 0-0 S(1) line ratio is estimated to be about $\sim 1 \pm 0.4$ at the DF (position A in Allers et al. 2005). The ratio predicted by the model is about 2.5 times higher. A pressure of $4 \times 10^8 \text{ K cm}^{-3}$ is probably too high to reproduce this ratio and more detailed modeling would be needed. The previous H_2 study by Allers et al. (2005) show in fact that an isobaric model with $P = 8 \times 10^7 \text{ K cm}^{-3}$ (with a pressure 5 times lower than the one we assumed) matches the observed intensities of the ground-state rotational lines, the H_2 1-0 S(1) and 2-1 S(1) lines.

To determine the pressure and density variations at the PDR edge, future detailed spatial studies of both the H_2 pure rotational and rovibrational lines are required. Excitation diagram from the numerous H_2 lines observable by JWST MIRI and NIRSpec will permit to trace the warm and hot (UV-pumped) excitation temperatures, local gas density and pressure gradients at small spatial scales. It will be possible to probe the physical conditions in most of the dense sub-structures detected with Keck or ALMA as well as the very thin surface layers that are sufficiently heated to photo-evaporate from the PDR. On the other hand, one could marginally spatially resolve the temperature gradient inside the over-dense substructures themselves. For a high pressure PDR,

the H_2 emission thickness is predicted to be $< 1''$ and spatial shift between the H_2 rotational line emission peaks is comparable or smaller to the JWST diffraction limit which is about $0.2''$ to $1''$ in the 5-28 μm range where are the pure rotational lines.

These constraints on the physical conditions may allow to better understand the dynamical effects in PDRs, e.g. compression waves, photo-evaporative flows, and IF and DF instabilities. Recently, Maillard et al. (2021) modelled the dynamical effects of the radiative stellar feedback on the H-to- H_2 transition. These authors built a semi-analytical model of the H/ H_2 transition in a 1D plane-parallel PDR illuminated by FUV radiation with advancing IF resulting from the photoevaporation mechanism. For the high-excitation PDRs such as the Orion Bar, moderate effects are however predicted. The H^0/H_2 transition will be slightly closer to the IF and sharper. Much stronger effects of the advection dynamics with merged, or close to merged, IF and DF are predicted for low excitation PDRs illuminated by late O stars such as the Horsehead nebula.

7. Summary

In this section, a short summary and prospects for JWST are given.

1. We obtained with Keck/NIRC2 using AO the most complete and detailed maps of the complex UV-irradiated region of the Orion Bar where the conversion from ionized to atomic to molecular gas occurs. We mapped the vibrationally excited line of H_2 at 2.12 μm , tracing the dissociation front, and the [FeII] line at 1.64 μm and the Br γ line at 2.16 μm , tracing the ionization front. This allows for the first time to spatially resolve the dissociation front with a resolution of $0.1''$ (40 AU) and obtain a very precise determination of the offset between the peak of the ionization and dissociation front of 12 and 12.5'' (0.024-0.025 pc).
2. The H^0/H_2 dissociation front appears highly structured with several ridges and extremely sharp filaments with a width of $1 - 2''$ (0.002-0.004 pc or 400-800 AU). Ridges and filaments run parallel to the dissociation front but a succession of bright sub-structures in H_2 is also observed from the edge of the DF towards the molecular region. The multiple H_2 emission peaks along and across the Bar may be associated with a multitude of small highly irradiated and dense PDRs.
3. The comparison with ALMA data of the HCO^+ J=4-3 line emission, a good indicator of dense gas, show a remarkable similar spatial distribution between the H_2 and HCO^+ emission. This suggests that they both come from high densities but also indicates their strong chemical link and that the H/ H_2 and C $^+$ /C/CO transition zones are very close.
4. We spatially resolved additional compact sources corresponding to over-dense irradiated sub-structures such as proplyds near the ionization front, and perhaps collapsing protostars near the dissociation front.
5. We compared our extinction-corrected observations of the emission gas lines to the model predictions using the Cloudy code for the ionized gas and the Meudon PDR code for the neutral PDR gas. The models reproduce well (by a factor less than 2) the observed intensity peak of the gas lines emission. The predicted vibrationally excited H_2 come from the H^0/H_2 transition layer where the gas density is of the order of few 10^5 cm^{-3} .
6. This work confirms without ambiguity that one of the densest portions of the Bar lie along the dissociation front and

that the Orion Bar edge is composed of small, dense sub-structures at high thermal pressure with P about 10^8 K cm^{-3} immersed in a more diffuse environment. Further studies are required to accurately determine the gas physical conditions (pressure and density gradients) and dynamics of these over-dense irradiated sub-structures and better understand their physical origin and future.

The JWST ERS proposal on the Orion Bar PDR (Berné et al. 2022) will very soon give access to multiple spectroscopic images with a similar spatial resolution to the Keck/NIRC2 observations but a signal-to-noise gain that will increase contrast and allow a sharper detection of all less bright substructures. JWST with IFU spectroscopy will give insight into the local gas physical conditions (temperature, density and pressure) and chemical composition of the warm very structured irradiated medium. It will be possible to probe the physical conditions in most of the dense sub-structures detected with Keck or ALMA as well as the very thin surface layers that are sufficiently heated to photo-evaporate from the Orion Bar edge. Furthermore, JWST will spatially resolve at near and mid-IR wavelengths dust emission and scattering simultaneously with the gas lines. Variation of the dust properties (e.g., size distribution, extinction) as function of PDR depth will be constrained and will be taken into account in the UV shielding, thermal balance and H_2 formation process.

Acknowledgements. The data presented herein were obtained at the W. M. Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W. M. Keck Foundation.

The authors wish to recognize and acknowledge the very significant cultural role and reverence that the summit of Maunakea has always had within the indigenous Hawaiian community. We are most fortunate to have the opportunity to conduct observations from this mountain.

This research has made use of the Keck Observatory Archive (KOA), which is operated by the W. M. Keck Observatory and the NASA Exoplanet Science Institute (NExScI), under contract with the National Aeronautics and Space Administration. Javier R. Goicoechea thanks the Spanish MCIYU for funding support under grant PID2019-106110GB-I00.

References

- Agúndez, M., Goicoechea, J. R., Cernicharo, J., Faure, A., & Roueff, E. 2010, *ApJ*, 713, 662
- Allers, K. N., Jaffe, D. T., Lacy, J. H., Draine, B. T., & Richter, M. J. 2005, *ApJ*, 630, 368
- Andree-Labsch, S., Ossenkopf-Okada, V., & Röllig, M. 2017, *A&A*, 598, A2
- Bally, J., Ginsburg, A., Silvia, D., & Youngblood, A. 2015, *A&A*, 579, A130
- Bally, J., O'Dell, C., & McCaughrean, M. J. 2000, *The Astronomical Journal*, 119, 2919
- Bernard-Salas, J., Habart, E., Arab, H., et al. 2012, *A&A*, 538, A37
- Berné, O., Habart, É., Peeters, E., et al. 2022, arXiv e-prints, arXiv:2201.05112
- Berné, O., Marcelino, N., & Cernicharo, J. 2010, *Nature*, 466, 947
- Blagrove, K. P. M., Martin, P. G., Rubin, R. H., et al. 2007, *ApJ*, 655, 299
- Bregman, J. D., Allamandola, L. J., Tielens, A. G. G. M., Geballe, T. R., & Witteborn, F. C. 1989, *ApJ*, 344, 791
- Burton, M. G., Hollenbach, D. J., & Tielens, A. G. G. M. 1990, *ApJ*, 365, 620
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
- Champion, J., Berné, O., Vicente, S., et al. 2017, *A&A*, 604, A69
- Chuss, D. T., Andersson, B. G., Bally, J., et al. 2019, *ApJ*, 872, 187
- Cuadrado, S., Salas, P., Goicoechea, J. R., et al. 2019, *A&A*, 625, L3
- Ferland, G. J., Chatzikos, M., Guzmán, F., et al. 2017, *Rev. Mexicana Astron. Astrofis.*, 53, 385
- Giard, M., Bernard, J. P., Lacombe, F., Normand, P., & Rouan, D. 1994, *A&A*, 291, 239
- Goicoechea, J. R. & Cuadrado, S. 2021, *A&A*, 647, L7
- Goicoechea, J. R., Cuadrado, S., Pety, J., et al. 2017, *A&A*, 601, L9
- Goicoechea, J. R., Pety, J., Cuadrado, S., et al. 2016, *Nature*, 537, 207
- Goicoechea, J. R., Teyssier, D., Etzaluze, M., et al. 2015, *ApJ*, 812, 75
- Gorti, U. & Hollenbach, D. 2002, *ApJ*, 573, 215
- Großschedl, J. E., Alves, J., Meingast, S., et al. 2018, *A&A*, 619, A106
- Guerra, J. A., Chuss, D. T., Dowell, C. D., et al. 2021, *ApJ*, 908, 98
- Hartigan, P., Downes, T., & Isella, A. 2020, *ApJ*, 902, L1
- Herrmann, F., Madden, S. C., Nikola, T., et al. 1997, *ApJ*, 481, 343
- Hogerheijde, M. R., Jansen, D. J., & van Dishoeck, E. F. 1995, *A&A*, 294, 792
- Inutsuka, S.-i. & Miyama, S. M. 1997, *ApJ*, 480, 681
- Joblin, C., Bron, E., Pinto, C., et al. 2018, *A&A*, 615, A129
- Jones, A. P., Fanciullo, L., Köhler, M., et al. 2013, *A&A*, 558, A62
- Jones, A. P., Köhler, M., Ysard, N., Bocchio, M., & Verstraete, L. 2017, *A&A*, 602, A46
- Juvela, M. 2019, *A&A*, 622, A79
- Kaplan, K. F., Dinerstein, H. L., Kim, H., & Jaffe, D. T. 2021, *ApJ*, 919, 27
- Kaplan, K. F., Dinerstein, H. L., Oh, H., et al. 2017, *ApJ*, 838, 152
- Kirsanova, M. S. & Wiebe, D. S. 2019, *MNRAS*, 486, 2525
- Knight, C., Peeters, E., Stock, D. J., Vacca, W. D., & Tielens, A. G. G. M. 2021, arXiv e-prints, arXiv:2106.01519
- Köhler, M., Ysard, N., & Jones, A. P. 2015, *A&A*, 579, A15
- Kounkel, M., Hartmann, L., Loinard, L., et al. 2017, *ApJ*, 834, 142
- Le Petit, F., Nehmé, C., Le Bourlot, J., & Roueff, E. 2006, *ApJS*, 164, 506
- Lehmann, A., Godard, B., Pineau des Forêts, G., Vidal-García, A., & Falgarone, E. 2022, *A&A*, 658, A165
- Lis, D. C. & Schilke, P. 2003, *ApJ*, 597, L145
- Lu, J. R., Ghez, A. M., Hornstein, S. D., et al. 2009, *ApJ*, 690, 1463
- Luhman, K. L., Engelbracht, C. W., & Luhman, M. L. 1998, *ApJ*, 499, 799
- Luhman, M. L., Jaffe, D. T., Keller, L. D., & Pak, S. 1994, *ApJ*, 436, L185
- Mackey, J. & Lim, A. J. 2011, *MNRAS*, 412, 2079
- Maillard, V., Bron, E., & Le Petit, F. 2021, *A&A*, 656, A65
- Marconi, A., Testi, L., Natta, A., & Walmsley, C. M. 1998, *A&A*, 330, 696
- Menten, K. M., Reid, M. J., Forbrich, J., & Brunthaler, A. 2007, *A&A*, 474, 515
- Nagy, Z., Van der Tak, F. F. S., Ossenkopf, V., et al. 2013, *A&A*, 550, A96
- O'Dell, C. R. 2001, *ARA&A*, 39, 99
- O'Dell, C. R., Kollatschny, W., & Ferland, G. J. 2017, *ApJ*, 837, 151
- O'Dell, C. R. & Yusef-Zadeh, F. 2000, *AJ*, 120, 382
- Ossenkopf, V., Röllig, M., Neufeld, D. A., et al. 2013, *A&A*, 550, A57
- Ossenkopf, V., Röllig, M., Simon, R., et al. 2010, *A&A*, 518, L79
- Pabst, C. H. M., Goicoechea, J. R., Teyssier, D., et al. 2020, *A&A*, 639, A2
- Parikka, A., Habart, E., Bernard-Salas, J., et al. 2017, *A&A*, 599, A20
- Parikka, A., Habart, E., Bernard-Salas, J., Köhler, M., & Abergel, A. 2018, *A&A*, 617, A77
- Parmar, P. S., Lacy, J. H., & Achtermann, J. M. 1991, *ApJ*, 372, L25
- Pellegrini, E. W., Baldwin, J. A., Ferland, G. J., Shaw, G., & Heathcote, S. 2009, *ApJ*, 693, 285
- Röllig, M. & Ossenkopf-Okada, V. 2022, arXiv e-prints, arXiv:2205.04233
- Schirmer, T., Ysard, N., Habart, E., et al. in prep
- Schneider, N., Simon, R., Guevara, C., et al. 2020, *PASP*, 132, 104301
- Sellgren, K., Tokunaga, A. T., & Nakada, Y. 1990, *ApJ*, 349, 120
- Shaw, G., Ferland, G. J., Henney, W. J., et al. 2009, *ApJ*, 701, 677
- Sheffer, Y., Wolfire, M. G., Hollenbach, D. J., Kaufman, M. J., & Cordier, M. 2011, *ApJ*, 741, 45
- Tauber, J. A., Lis, D. C., Keene, J., Schilke, P., & Buettgenbach, T. H. 1995, *A&A*, 297, 567
- Thronson, Harley A., J., Majewski, S., Descartes, L., & Hereld, M. 1990, *ApJ*, 364, 456
- Tielens, A. G. G. M., Meixner, M. M., van der Werf, P. P., et al. 1993, *Science*, 262, 86
- Tremblin, P., Audit, E., Minier, V., Schmidt, W., & Schneider, N. 2012, *A&A*, 546, A33
- van der Werf, P. P., Goss, W. M., & O'Dell, C. R. 2013, *ApJ*, 762, 101
- van der Werf, P. P., Stutzki, J., Sternberg, A., & Krabbe, A. 1996, *A&A*, 313, 633
- Walmsley, C. M., Natta, A., Oliva, E., & Testi, L. 2000, *A&A*, 364, 301
- Weilbacher, P. M., Monreal-Ibero, A., Kollatschny, W., et al. 2015, *A&A*, 582, A114
- Wen, Z. & O'dell, C. R. 1995, *ApJ*, 438, 784
- Wolfire, M. G., Vallini, L., & Chevance, M. 2022, arXiv e-prints, arXiv:2202.05867
- Wright, C. M., van Dishoeck, E. F., Cox, P., Sidher, S. D., & Kessler, M. F. 1999, *ApJ*, 515, L29
- Wu, R., Bron, E., Onaka, T., et al. 2018, *A&A*, 618, A53
- Wyrowski, F., Schilke, P., Hofner, P., & Walmsley, C. M. 1997, *ApJ*, 487, L171
- Young Owl, R. C., Meixner, M. M., Wolfire, M., Tielens, A. G. G. M., & Tauber, J. 2000, *ApJ*, 540, 886
- Zhang, Z. E., Cummings, S. J., Wan, Y., Yang, B., & Stancil, P. C. 2021, *ApJ*, 912, 116