

# Conservation laws for potential vorticity in a salty ocean or cloudy atmosphere

Parvathi Kooloth<sup>1</sup>, Leslie M. Smith<sup>1,2</sup>, Samuel N. Stechmann<sup>1,3</sup>

<sup>1</sup>Department of Mathematics,

<sup>2</sup>Department of Engineering Physics,

<sup>3</sup>Department of Atmospheric and Oceanic Sciences,

University of Wisconsin–Madison, Madison, WI, 53706, USA

## Key Points:

- Important conservation laws of circulation and potential vorticity are known only under idealized assumptions.
- Here, conservation laws are described with additional realism, namely salinity in the ocean or moisture and clouds in the atmosphere.
- Potential vorticity is not conserved for each fluid parcel but for certain pancake-shaped volumes.

arXiv:2206.08254v1 [physics.ao-ph] 16 Jun 2022

**Abstract**

One of the most important conservation laws in atmospheric and oceanic science is conservation of potential vorticity. The original derivation is approximately a century old, in the work of Rossby and Ertel, and it is related to the celebrated circulation theorems of Kelvin and Bjerknes. However, the laws apply to idealized fluids, and extensions to more realistic scenarios have been problematic. Here, these laws are extended to hold with additional fundamental complexities, including salinity in the ocean, or moisture and clouds in the atmosphere. In the absence of these additional complexities, it is known that potential vorticity is conserved following each fluid parcel; here, for a salty ocean or cloudy atmosphere, the general conserved quantity is potential vorticity integrated over certain pancake-shaped volumes. Furthermore, the conservation laws are also related to a symmetry in the Lagrangian, which brings a connection to the symmetry-conservation relationships seen in other areas of physics.

**Plain Language Summary**

A vortex can be seen in the atmosphere or ocean in a variety of settings, including the intense vortices that are familiar from tornadoes and hurricanes. A quantity called vorticity is a measure of the strength of a vortex, or, more generally, of the amount of swirl in a fluid. Intimately connected to vorticity is an important conserved quantity referred to as potential vorticity. This quantity remains unchanged following the trajectory of a fluid parcel. Owing to its conserved nature, potential vorticity is widely used to study a wide range of weather events. However, this century-old conservation result is only valid in an idealized setting: an atmosphere devoid of clouds or an ocean without dissolved salt—i.e., without salinity. Here, the conservation law is extended to hold in a cloudy atmosphere and salty oceans. The generalized conservation statement presented here will aid in the wider applicability of potential vorticity for analyzing and understanding atmospheric and oceanic dynamics.

**1 Introduction**

In addition to conservation laws of mass, momentum, and energy, other conservation laws arise for fluids, such as conservation of circulation and potential vorticity. The circulation  $\Gamma$  is defined as the integral of the velocity  $\vec{u}$  along a closed curve  $C$ ,

$$\Gamma = \oint_C \vec{u} \cdot d\vec{l}, \quad (1)$$


and the circulation theorems of Kelvin and Bjerknes (Thomson, 1867; Bjerknes, 1898; Thorpe et al., 2003) show that, under certain assumptions, the circulation is conserved if the closed curve  $C(t)$  moves with the fluid. See Figure 1 for an illustration. The circulation is also related to the vorticity,  $\vec{\omega} = \nabla \times \vec{u}$ , since Eqn. 1 can be rewritten, using Stokes' theorem, as

$$\Gamma = \int_A \vec{\omega} \cdot d\vec{A}, \quad (2)$$

which is an area integral over any surface  $A$  whose boundary is the closed curve  $C$ , and where  $d\vec{A}$  is a surface area element whose vector direction is perpendicular to the surface. Helmholtz (Helmholtz, 1858) had earlier formulated some related conservation statements in terms of vorticity. An associated quantity is the potential vorticity (PV), which is the dot product of the vorticity vector and the entropy gradient,  $\nabla s$ , weighted by the fluid density  $\rho$ ; that is,

$$PV = \frac{\vec{\omega} \cdot \nabla s}{\rho}. \quad (3)$$

Ertel (Ertel, 1942) showed that PV is conserved following the trajectory of a fluid parcel, which builds on an earlier result of Rossby (Rossby, 1940) for the shallow water equations.



**Figure 1.** Geometry of the circulation theorem for a dry atmosphere or freshwater ocean. The curve  $C(t)$  may be any curve on a surface of constant entropy,  $s(\vec{x}, t) = c$ .

PV is a fundamental quantity of interest in geophysical flows (Müller, 1995; Salmon, 1998). It is extensively used in analysis of weather events (Davis & Emanuel, 1991; Lackmann, 2002), the general ocean circulation (Rhines, 1986; Pollard & Regier, 1990; Holland et al., 1984; Müller, 1995), magnetohydrodynamics (MHD) and astrophysical fluid dynamics (Webb & Mace, 2015). The wide-ranging applications of PV stem from its conservation properties and the notion of PV inversion (Hoskins et al., 1985; Smith & Stechmann, 2017; Wetzel et al., 2020).

Despite the importance of PV and the further knowledge gained over approximately a century, the conservation laws of circulation and PV are limited to fluids that are relatively idealized. In particular, these conservation laws apply to an ocean or atmosphere that is composed of a single fluid substance—water in the ocean, or dry air in the atmosphere. In nature, the ocean and atmosphere include other substances which are of fundamental significance. Moisture and clouds in the atmosphere are associated with weather events such as hurricanes, tornadoes, and monsoons. Salinity in the ocean affects the density and helps drive the thermohaline circulation and meridional overturning circulation that move heat from low latitudes to high latitudes (Marshall & Schott, 1999; Curry & Mauritzen, 2005; Peterson et al., 2006; Vallis, 2017). As one route to better understanding and predicting these phenomena, it is desirable to have knowledge of the associated conserved quantities.

The main goal of the present paper is to derive conservation principles for circulation and potential vorticity, for the case of an ocean with salinity or an atmosphere with moisture and clouds. The atmospheric case will be described first, and then the oceanic case will follow in a similar way. Also, circulation will be treated first, and then potential vorticity.

## 2 Circulation

The starting point is the evolution equation for the circulation  $\Gamma$ , which was defined above in Eqn. 1. Its evolution equation follows from the equations for conservation of mass and momentum (Thorpe et al., 2003; Bjerknes, 1898; Vallis, 2017) and it takes the form

$$\frac{D\Gamma}{Dt} = \oint_{C(t)} \frac{1}{\rho} \nabla p \cdot d\vec{l}, \quad (4)$$

where  $p$  is the pressure,  $C(t)$  is a closed curve that moves along with the fluid, and  $D/Dt = \partial/\partial t + \vec{u} \cdot \nabla$  is a material derivative—i.e., a derivative along the trajectory of a fluid parcel. See the Supplementary Materials for a derivation. The fluid is assumed to be compressible and inviscid.

Before considering the moist case, it is instructive to consider the dry case, for which it is known that circulation is conserved, under certain assumptions on the curve  $C(t)$ . To see this, one can rewrite (4) as

$$\frac{D\Gamma}{Dt} = \oint_{C(t)} \theta d\pi, \quad (5)$$

where the Exner function  $\pi(p)$  is a function of pressure, and where the potential temperature  $\theta$  is a function of entropy, as  $s = c_p \log \theta + \text{const.}$ , where  $c_p$ , commonly assumed constant for a dry atmosphere, is the specific heat at constant pressure. The details of this derivation are provided in the Supplementary Materials. From (5), one can deduce the scenarios when the right-hand side is zero and circulation  $\Gamma$  is conserved. In particular, if  $\theta$  (and hence also entropy  $s$ ) is a constant on the closed curve  $C(t)$ , then the right-hand side of (5) is zero, by the fundamental theorem of calculus for line integrals. Consequently, circulation is conserved,

$$\frac{D\Gamma}{Dt} = 0, \quad (6)$$

if the closed curve  $C(t)$  lies on a surface of constant entropy. See Figure 1 for an illustration. In order for the material curve  $C(t)$  to remain on a surface of constant entropy for all times, the fluid is assumed to be adiabatic, so that  $Ds/Dt = 0$ . Finally, from (6), the circulation  $\Gamma$  is then said to be a *material invariant*, since its material derivative is zero.

In the moist case, one could try to repeat the derivation from the dry case to arrive at a circulation theorem. However, this line of reasoning fails for a moist system, since the integrand that would arise in (5) would no longer be a material invariant, due to phase changes and cloud latent heating.

We now show that the challenges of the moist case can actually be overcome by referring to fundamental principles of moist thermodynamics. Two derivations will be shown: a derivation that directly refers to enthalpy, and a derivation that does not. In terms of the enthalpy  $h(s, p, q_t)$ , the fundamental thermodynamic relation (Landau & Lifshitz, 1980, 1987; Pauluis, 2008) states that

$$dh = T ds + \rho^{-1} dp + \mu dq_t, \quad (7)$$


which can be viewed as the differential of the function  $h(s, p, q_t)$  as

$$dh = \left( \frac{\partial h}{\partial s} \right)_{p, q_t} ds + \left( \frac{\partial h}{\partial p} \right)_{s, q_t} dp + \left( \frac{\partial h}{\partial q_t} \right)_{s, p} dq_t, \quad (8)$$

where  $h$  is enthalpy,  $T$  is temperature,  $q_t$  is the total water specific humidity, and  $\mu = \mu_v - \mu_d$  is the difference of the chemical potentials associated with water vapor and dry air, respectively. As a first important observation, notice that (7) includes the term  $\rho^{-1} dp$ , which is also the sole term that appears in the evolution equation for circulation in (4). Therefore, inserting the fundamental relation (7) into the evolution equation for circulation in (4), we have

$$\frac{D\Gamma}{Dt} = - \oint_{C(t)} T ds - \oint_{C(t)} \mu dq_t, \quad (9)$$

where it was used that  $\oint_{C(t)} dh = 0$  for a closed curve. The aim now is to identify scenarios when the right-hand side is zero. In this direction, note that the line integrals are zero if the curve  $C(t)$  lies on a surface of constant entropy  $s$  and a surface of constant



**Figure 2.** Geometry of the circulation theorem for an atmosphere with clouds or an ocean with salinity. The curve  $C(t)$  is a curve of constant entropy  $s$  and total water specific humidity  $q_t$ .

$q_t$ . Furthermore, in order for the material curve to have constant  $s$  and  $q_t$  for all times, it is assumed that  $Ds/Dt = 0$  and  $Dq_t/Dt = 0$ , so that  $s$  and  $q_t$  are both material invariants, if reversible phase changes are assumed between water vapor and liquid water, and precipitation is absent. Therefore, the right hand side of (9) vanishes and we have

$$\frac{D\Gamma}{Dt} = 0, \quad (10)$$

if  $C(t)$  is a closed curve of constant  $s$  and  $q_t$ . See Figure 2 for an illustration. This is the moist, cloudy analog of the circulation theorems of Kelvin and Bjerknes.

As an alternative derivation of (10) that does not directly refer to enthalpy, return again to the starting point of (4), where the integrand is  $\rho^{-1}$ . Now use a fundamental property of moist thermodynamics: any moist thermodynamic variable can be written as a function of  $s$ ,  $q_t$  and  $p$  (Pauluis, 2008). Hence,  $\rho^{-1} dp$  can be written as  $\rho^{-1}(s, q_t, p) dp$ , which, on a circuit  $C(t)$  of constant entropy  $s$  and total water mixing ratio  $q_t$ , is a function of pressure  $p$  alone. Consequently, (4) can furthermore be simplified to

$$\frac{D\Gamma}{Dt} = \oint_{C(t)} df, \quad (11)$$

where  $f(p)$  is an antiderivative that satisfies  $df = (df/dp) dp = \rho^{-1}(s, q_t, p) dp$ . By (7) and (8), since  $(dh/dp)_{s, q_t} = \rho^{-1}$ , the function  $f$  can be identified as the enthalpy  $h$ , which is insightful but not essential for this derivation. Finally, since  $C(t)$  is a closed curve, it follows from (11) that

$$\frac{D\Gamma}{Dt} = 0, \quad (12)$$

if  $C(t)$  is a closed curve of constant  $s$  and  $q_t$ .

As a comparison of the dry and moist cases, notice that both apply to material curves  $C(t)$  on surfaces of constant entropy  $s$ . However, while *any* such curve is valid for a dry atmosphere, the curve must also have constant  $q_t$  for a moist, cloudy atmosphere.

### 3 Potential vorticity

Now consider the second quantity of interest here, defined above in (3): potential vorticity. Its evolution equation follows from the equations for conservation of mass, momentum, and entropy (Müller, 1995; Schubert et al., 2001) and it takes the form

$$\rho \frac{D}{Dt} \left( \frac{\vec{\omega} \cdot \nabla s}{\rho} \right) = \nabla s \cdot \nabla \times \left( \frac{1}{\rho} \nabla p \right) \quad (13)$$

See the Supplementary Materials for a derivation. On the right-hand side, the term  $\nabla \times (\rho^{-1} \nabla p)$  arises from the curl of the pressure gradient in the statement of conservation of momentum.

For a dry atmosphere, PV is conserved for each fluid parcel, and it is helpful to review this case as a prelude to the moist case. To see how dry PV is conserved, note that the right-hand side of (13) can be written as

$$\nabla s \cdot \nabla \times \left( \frac{1}{\rho} \nabla p \right) = -\frac{1}{\rho^2} \nabla s \cdot \nabla p \times \nabla \rho, \quad (14)$$

which is a scalar triple product. Then one can use a fundamental property of thermodynamics for a dry atmosphere: the entropy can be expressed as a function of pressure and density, so that  $s = s(p, \rho)$ . Consequently, we have

$$\nabla s = \frac{\partial s}{\partial p} \nabla p + \frac{\partial s}{\partial \rho} \nabla \rho \quad (15)$$

and inserting this into (14) shows that the right-hand side of (13) is zero, and the PV is conserved for each fluid parcel:

$$\frac{D}{Dt} \left( \frac{\vec{\omega} \cdot \nabla s}{\rho} \right) = 0. \quad (16)$$

In essence, this proof works smoothly because the entropy  $s$  is a function of pressure  $p$  and density  $\rho$ , for a dry atmosphere. However, for a moist atmosphere, due to the effects of phase changes and cloud latent heating, it is not true that  $s = s(p, \rho)$ , and it is therefore difficult to see how to extend these proof ideas to apply to the moist case.

For a moist atmosphere with clouds, we now show that the difficulty of PV conservation can be resolved by appealing to fundamentals of moist thermodynamics. In particular, use the gradient form of the fundamental thermodynamic relation in (7), and insert into (13) for the  $\rho^{-1} \nabla p$  term. Then, after using vector calculus identities, (13) becomes


$$\rho \frac{D}{Dt} \left( \frac{\vec{\omega} \cdot \nabla s}{\rho} \right) = -\nabla s \cdot \nabla \times (\mu \nabla q_t). \quad (17)$$

The benefit of having replaced  $\rho^{-1} \nabla p$  with  $\mu \nabla q_t$  will be seen below and arises from the fact that the right-hand side of (17) now involves two quantities,  $s$  and  $q_t$ , which are both conserved for each fluid parcel, for the case of reversible phase changes between water vapor and cloud liquid water, with no precipitation.

To proceed with finding the conservation law for moist PV, we abandon the idea that moist PV can be conserved for each fluid parcel. In other words, we abandon hope of showing that the right-hand side of (17) is zero. Instead, consider the integral of moist PV over certain volumes. The evolution equation of volume-integrated PV is

$$\frac{D}{Dt} \int \int \int_{V(t)} \frac{\vec{\omega} \cdot \nabla s}{\rho} \rho dV = \int \int \int_{V(t)} \rho \frac{D}{Dt} \left( \frac{\vec{\omega} \cdot \nabla s}{\rho} \right) dV \quad (18)$$

$$= - \int \int \int_{V(t)} \nabla \cdot (\mu \nabla q_t \times \nabla s) dV \quad (19)$$



**Figure 3.** Geometry of the PV conservation law for an atmosphere with clouds or an ocean with salinity. The conserved quantity is not PV of individual parcels, but PV integrated over certain cylinder-like volumes.

$$= - \int \int_{A(t)} \mu \nabla q_t \times \nabla s \cdot d\vec{A}, \quad (20)$$

where the volume integral in (18) is mass-weighted using density  $\rho$ , and where the volume  $V(t)$  is a material volume that moves with the fluid but otherwise remains to be specified. See the Supplementary Materials for details of the derivation. In brief, (18) uses the transport theorem, (19) uses (17) and some vector calculus identities, and (20) follows from the divergence theorem, where  $A(t)$  is the surface that encloses the material volume  $V(t)$ .

Now it is time to identify the special volumes for which (20) is zero and integrated moist PV is conserved. The intuition is similar to the case of the moist circulation theorem in (9) where the appearance of the line elements  $ds$  and  $dq_t$  suggested the choice of curves of constant  $s$  and  $q_t$ . Here, it is a surface integral that arises in (20), and in order for the integral to vanish, the surface area element  $d\vec{A}$  must be perpendicular to  $\nabla s$  or  $\nabla q_t$ . Hence, we have

$$\frac{D}{Dt} \int \int \int_{V(t)} \frac{\vec{\omega} \cdot \nabla s}{\rho} \rho dV = 0, \quad (21)$$

if the volume is defined to be enclosed by surfaces of constant  $s$  and surfaces of constant  $q_t$ . An alternative derivation, which does not explicitly use the fundamental thermodynamic relation, is shown in the Supplementary Materials. Either derivation leads to the same result, (21), the conservation law for PV for a cloudy atmosphere, which applies to volume-integrated PV.

More generally, the surface could be composed piecewise, based on level surfaces of general functions  $F(s, q_t)$ , since any such level surface is perpendicular to  $\nabla s \times \nabla q_t$ . Nevertheless, it is convenient and simple to choose surfaces of constant  $s$  or constant  $q_t$ , as illustrated in Figure 3. The simplest such volume is shaped like a thin cylinder, or pill box, or pancake.

As perhaps the most special case, notice that the volume  $V(t)$  in Figure 3 will shrink to a point at a maximum or minimum value of  $q_t$  on a constant-entropy surface. As a result, at minima or maxima of  $q_t$ , the PV is conserved for the individual parcel.

	Cloudy Atmosphere	Salty Ocean
Circulation Theorem	$\frac{D}{Dt} \oint_{C(t)} \vec{u} \cdot d\vec{l} = 0$	$\frac{D}{Dt} \oint_{C(t)} \vec{u} \cdot d\vec{l} = 0$
Curve $C(t)$	Constant $s$ and $q_t$	Constant $\eta$ and $S$
PV Conservation Law	$\frac{D}{Dt} \int \int \int_{V(t)} \frac{\vec{\omega} \cdot \nabla s}{\rho} \rho dV = 0$	$\frac{D}{Dt} \int \int \int_{V(t)} \frac{\vec{\omega} \cdot \nabla \eta}{\rho} \rho dV = 0$
Surface of Volume $V(t)$	Constant $s$ or $q_t$	Constant $\eta$ or $S$

**Table 1. Summary of conservation laws** Conservation laws for an atmosphere with clouds and an ocean with salinity. Commonly used notation for a cloudy atmosphere is  $s$  and  $q_t$  for the entropy and total water specific humidity, respectively, and for a salty ocean is  $\eta$  and  $S$  for the entropy and salinity, respectively. Rotation and the Coriolis force could also be incorporated with modifications described in (23) and (24).

As another generalization, notice that the PV could be defined as  $\rho^{-1} \vec{\omega} \cdot \nabla q_t$  with  $q_t$  in the role that is normally played by entropy  $s$ . A similar conservation law also follows for such a  $q_t$ -based PV. Moreover, the PV could be defined as  $\rho^{-1} \vec{\omega} \cdot \nabla F(s, q_t)$  for any function  $F$  of  $s$  and  $q_t$ . In that case, the proof proceeds in the same way by writing the fundamental thermodynamic relation in a form that replaces  $s$  and  $q_t$  by  $F(s, q_t)$  and  $G(s, q_t)$ , where  $G$  is arbitrary but functionally independent of  $F$ .

For an ocean with salinity, the conservation laws for circulation and PV follow in the same way. Simply replace specific humidity  $q_t$  by salinity  $S$ , and to avoid confusion, change the notation of the entropy  $s$  to instead be  $\eta$ , as is common in oceanography. Table 1 summarizes the conservation laws for both the cloudy atmosphere and salty ocean.

#### 4 Symmetry–conservation relationship

A symmetry–conservation relationship is also associated with these conservation laws of circulation and PV. In particular, it is a particle-relabeling symmetry, which, as the name suggests, involves particles whose positions are  $\vec{x}(\vec{a}, t)$  or the relabeled positions  $\vec{x}(\vec{a}', t)$ , where the original label  $\vec{a}$  was replaced by the relabeling  $\vec{a}' = \vec{a}'(\vec{a}, t)$ . In a discrete setting, the relabeling of  $\vec{x}_n(t)$  to  $\vec{x}_{n'}(t)$  would simply involve a permutation of the labels,  $n = 1, 2, 3, \dots, N$ , but there is no guarantee of an associated conservation law for such a discrete symmetry. For the continuum setting of a cloudy atmosphere, the symmetry applies to the Lagrangian

$$\mathcal{L}(t) = \int \int \int \rho \left[ \frac{1}{2} |\vec{u}|^2 - E(\rho, s, q_t) - \Phi(\vec{x}) \right] dV_a, \quad (22)$$

where  $E(\rho, s, q_t)$  is the internal energy,  $\Phi(\vec{x})$  is the gravitational potential, and the integral is over the particle labels  $\vec{a}$ , as the continuum version of a sum over the particles.

A key observation is that only some special particle relabelings will leave the Lagrangian unchanged. For instance, for a dry atmosphere, the internal energy  $E(\rho, s)$  is a function of density and entropy, and it is well-known that  $E(\rho, s)$  and the Lagrangian remain unchanged for certain particle relabelings on surfaces of constant entropy  $s$  (Salmon, 1998); and it follows that the Kelvin–Bjerknes circulation theorem holds for curves  $C(t)$  that lie on surfaces of constant entropy, as mentioned in (6) and Figure 1 above. Now,

for a moist atmosphere, one can see the natural extension: the particle relabeling should keep both  $s$  and  $q_t$  unchanged in order to keep  $E(\rho, s, q_t)$  and the Lagrangian unchanged; consequently, the moist circulation theorem in (10) and Figure 2 holds for curves  $C(t)$  of constant  $s$  and  $q_t$ . These symmetry–conservation ideas have been derived in detail for a moist Boussinesq setting (Kooloth et al., 2022) and here provide motivation and further understanding of the moist conservation laws of circulation and PV. For instance, because the particle relabeling is constrained to curves  $C(t)$  of both constant  $s$  and  $q_t$ , it follows that moist PV is not conserved for individual parcels.

## 5 Concluding discussion

Rotation and the Coriolis term could also be incorporated here, in which case the circulation theorem becomes

$$\frac{D}{Dt} \oint_{C(t)} (\vec{u} + \vec{\Omega} \times \vec{x}) \cdot d\vec{l} = 0, \quad (23)$$

where  $\Omega$  is the angular rotation vector and  $\vec{x}$  is the position vector. The extension to include rotation follows standard procedures as in cases without clouds or salinity (Salmon, 1998; Vallis, 2017; Cotter & Holm, 2014). Furthermore, the PV conservation laws have the same form as in Table 1 except with  $\omega = \nabla \times \vec{u}$  replaced by absolute vorticity,

$$\vec{\omega}_a = \nabla \times \vec{u} + 2\vec{\Omega}, \quad (24)$$

which is the sum of the relative vorticity and the angular rotation vector.

Many applications are possible for the conservation laws presented here. One common use of PV conservation is to diagnose non-conservation, which is an indication of additional physical processes (Haynes & McIntyre, 1987). The PV conservation laws here could be used for similar purposes, with certain modifications. For instance, cloud latent heating is considered one of the sources of non-conservation for dry PV, but is incorporated into the conservation law here for moist PV. Another application of PV conservation is in numerical models of weather and climate. It is often desirable to design such numerical methods to accurately satisfy the PV conservation law (Thuburn, 2008; Taylor & Fournier, 2010). The conservation laws here provide additional targets for numerical conservation, now incorporating additional physical processes of salinity and cloud latent heating.

## 6 Open Research

Data sharing not applicable to this article as no data sets were generated or analysed during the current study.

### Acknowledgments

Leslie Smith and Sam Stechmann gratefully acknowledge grant support from the Division of Mathematical Sciences of the National Science Foundation, in conjunction with the award DMS-1907667.

### References

- Bjerknes, V. (1898). Über einen hydrodynamischen zirkulationssatz und seine anwendung auf die mechanik der atmosphäre und des weltmeeres. *Kongl. Svenska Vetenskapskad. Handl. Bd.*, 31, 97–106.
- Cotter, C., & Holm, D. (2014). Variational formulations of sound-proof models. *Quart. J. Roy. Met. Soc.*, 140(683), 1966–1973.

- Curry, R., & Mauritzen, C. (2005). Dilution of the northern North Atlantic Ocean in recent decades. *Science*, *308*(5729), 1772–1774.
- Davis, C. A., & Emanuel, K. A. (1991). Potential vorticity diagnostics of cyclogenesis. *Monthly Weather Review*, *119*(8), 1929–1953.
- Ertel, H. (1942). Ein neuer hydrodynamischer wirbelsatz. *Met. Zeitschr.*, *59*, 277–281.
- Haynes, P. H., & McIntyre, M. E. (1987). On the evolution of vorticity and potential vorticity in the presence of diabatic heating and frictional or other forces. *J. Atmos. Sci.*, *44*(5), 828–841.
- Helmholtz, H. v. (1858). Über integrale der hydrodynamischen gleichungen, welche den wirbelbewegungen entsprechen. *J. Angew. Math.*, *55*, 25–55.
- Holland, W., Keffer, T., & Rhines, P. (1984). Dynamics of the oceanic general circulation: the potential vorticity field. *Nature*, *308*(5961), 698–705.
- Hoskins, B. J., McIntyre, M. E., & Robertson, A. W. (1985). On the use and significance of isentropic potential vorticity maps. *Quart. J. Roy. Met. Soc.*, *111*(470), 877–946.
- Kooloth, P., Smith, L. M., & Stechmann, S. N. (2022). Hamilton’s principle with phase changes and conservation principles for moist potential vorticity. , submitted.
- Lackmann, G. M. (2002). Cold-frontal potential vorticity maxima, the low-level jet, and moisture transport in extratropical cyclones. *Monthly Weather Review*, *130*(1), 59–74.
- Landau, L. D., & Lifshitz, E. M. (1980). *Statistical physics, part 1* (3rd ed., Vol. 5). Elsevier.
- Landau, L. D., & Lifshitz, E. M. (1987). *Fluid mechanics* (2nd ed., Vol. 6). Elsevier.
- Marshall, J., & Schott, F. (1999). Open-ocean convection: Observations, theory, and models. *Reviews of Geophysics*, *37*(1), 1–64.
- Müller, P. (1995). Ertel’s potential vorticity theorem in physical oceanography. *Reviews of Geophysics*, *33*(1), 67–97.
- Pauluis, O. (2008). Thermodynamic consistency of the anelastic approximation for a moist atmosphere. *J. Atmos. Sci.*, *65*(8), 2719–2729.
- Peterson, B. J., McClelland, J., Curry, R., Holmes, R. M., Walsh, J. E., & Aagaard, K. (2006). Trajectory shifts in the Arctic and subarctic freshwater cycle. *Science*, *313*(5790), 1061–1066.
- Pollard, R., & Regier, L. (1990). Large variations in potential vorticity at small spatial scales in the upper ocean. *Nature*, *348*(6298), 227–229.
- Rhines, P. B. (1986). Vorticity dynamics of the oceanic general circulation. *Annual Review of Fluid Mechanics*, *18*(1), 433–497.
- Rossby, C. G. (1940). Planetary flow patterns in the atmosphere. *Quart. J. Roy. Met. Soc.*, *66* (Suppl.), 68–87.
- Salmon, R. (1998). *Lectures on Geophysical Fluid Dynamics*. Oxford University Press.
- Schubert, W. H., Hausman, S. A., Garcia, M., Ooyama, K. V., & Kuo, H.-C. (2001). Potential vorticity in a moist atmosphere. *J. Atmos. Sci.*, *58*(21), 3148–3157.
- Smith, L. M., & Stechmann, S. N. (2017). Precipitating quasigeostrophic equations and potential vorticity inversion with phase changes. *J. Atmos. Sci.*, *74*(10), 3285–3303.
- Taylor, M. A., & Fournier, A. (2010). A compatible and conservative spectral element method on unstructured grids. *Journal of Computational Physics*, *229*(17), 5879–5895.
- Thomson, W. (1867). 4. on vortex atoms. *Proceedings of the Royal Society of Edinburgh*, *6*, 94–105. doi: 10.1017/S0370164600045430
- Thorpe, A. J., Volkert, H., & Ziemiański, M. J. (2003). The Bjerknes’ circulation theorem: A historical perspective: A historical perspective. *Bull. Am. Meteorol.*

- rol. Soc.*, 84(4), 471–480.
- Thuburn, J. (2008). Some conservation issues for the dynamical cores of NWP and climate models. *Journal of Computational Physics*, 227(7), 3715–3730.
- Vallis, G. K. (2017). *Atmospheric and Oceanic Fluid Dynamics*. Cambridge University Press.
- Webb, G., & Mace, R. (2015). Potential vorticity in magnetohydrodynamics. *Journal of Plasma Physics*, 81(1).
- Wetzel, A. N., Smith, L. M., Stechmann, S. N., Martin, J. E., & Zhang, Y. (2020). Potential vorticity and balanced and unbalanced moisture. *J. Atmos. Sci.*, 77(6), 1913–1931.