

On Effective Field Theories with Celestial Duals

Lecheng Ren,^a Marcus Spradlin,^{a,b} Akshay Yelleshpur Srikant^a and Anastasia Volovich^a

^a*Department of Physics, Brown University, Providence, RI 02912, USA*

^a*Brown Theoretical Physics Center, Brown University, Providence, RI 02912, USA*

E-mail: lecheng_ren@brown.edu, marcus_spradlin@brown.edu,
akshay_yelleshpur_srikant@brown.edu, anastasia_volovich@brown.edu

ABSTRACT: We show that associativity of the tree-level OPE in a celestial CFT imposes constraints on the coupling constants of the corresponding bulk theory. These constraints are the same as those derived in arXiv:2111.11356 from the Jacobi identity of the algebra of soft modes. The constrained theories are interesting as apparently well-defined celestial CFTs with a deformed $w_{1+\infty}$ symmetry algebra. We explicitly work out the ramifications of these constraints on scattering amplitudes involving gluons, gravitons and scalars in these theories. We find that all four-point amplitudes constructible solely from holomorphic or anti-holomorphic three-point amplitudes vanish on the support of these constraints, which implies that all purely holomorphic or purely anti-holomorphic higher-point amplitudes vanish.

Contents

1	Introduction	1
2	Review of the coupling constants constraints	2
3	Tree-level OPE associativity from amplitudes	4
4	Amplitudes in EFTs with celestial dual	5
4.1	Four-point amplitudes in the graviton-scalar sector	6
4.2	Four-point amplitudes in the gluon-scalar sector	7
4.3	Amplitudes of arbitrary multiplicity	8
5	Discussion	9
A	All-line shift recursion relations	10

1 Introduction

The central tenet of celestial holography [1–3] is that a scattering amplitude of n massless particles, with momenta $\{p_i\}$ and helicities $\{s_i\}$, when recast in a basis of boost eigenstates, can be interpreted as a correlation function of n operators with conformal weights (h_i, \bar{h}_i) in a two-dimensional celestial conformal field theory (CCFT). The four-dimensional helicity s_i of particle i is the same as the two-dimensional spin $s_i = h_i - \bar{h}_i$, and we use $\Delta_i = h_i + \bar{h}_i$.

This dictionary implies that the OPE of two operators in CCFT is determined by the collinear limit of the corresponding scattering amplitudes [4–6]. Specifically, the OPE is given by

$$\mathcal{O}_{h_1, \bar{h}_1}(z_1, \bar{z}_1) \mathcal{O}_{h_2, \bar{h}_2}(z_2, \bar{z}_2) \sim \frac{1}{z_{12}} \sum_p \sum_{m=0}^{\infty} C_p^{(m)}(\bar{h}_1, \bar{h}_2) \bar{z}_{12}^{p+m} \bar{\partial}^m \mathcal{O}_{h_1+h_2-1, \bar{h}_1+\bar{h}_2+p}(z_2, \bar{z}_2) \quad (1.1)$$

where $z_{ij} = z_i - z_j$ and the OPE coefficient $C_p^{(m)}(\bar{h}_1, \bar{h}_2)$ corresponding to the contribution of an operator with weights $(h_1 + h_2 - 1, \bar{h}_1 + \bar{h}_2 + p + m)$ is given by [4–6]

$$C_p^{(m)}(\bar{h}_1, \bar{h}_2) = -\frac{1}{2} \kappa_{s_1, s_2, -s_I} \frac{1}{m!} B(2\bar{h}_1 + p + m, 2\bar{h}_2 + p), \quad (1.2)$$

where $B(a, b)$ is the Euler beta function and $\kappa_{s_1, s_2, -s_I}$ is the coupling constant appearing in the bulk three-point scattering amplitude of particles with helicities s_1, s_2 and $-s_I$, with $s_I = s_1 + s_2 - p - 1$. Thus there is a direct link between the bulk three-point couplings and the celestial OPE. Restricting to tree-level and excluding massless higher spins leaves a finite roster of bulk three-point amplitudes which could potentially contribute to the sum in (1.1). These have been tabulated in [5].

It was observed in [7] that the OPE of conformally soft graviton operators (those with $\Delta_i = 2, 1, 0, -1, -2, \dots$), in the absence of higher derivative interactions in the bulk, forms a symmetry algebra. This algebra was identified as the Kac Moody algebra of the wedge subalgebra of $w_{1+\infty}$

in [8]. The inclusion of higher derivative interactions in the bulk deforms the algebra as demonstrated in [9]. In particular, note that for a given value of p , only the soft currents with $\Delta \geq p-1$ are modified. This implies that there is short list of operators that can modify the subleading and subsubleading soft currents. This is consistent with the results of [10–12]. In [9] it was also shown that the algebra of soft modes violates the Jacobi identity unless the bulk couplings satisfy certain particular constraints (reviewed in Section 2 below). In particular, this implies that the OPE of the soft modes is not associative for arbitrary couplings. It is natural to wonder about the implications of celestial OPE associativity on bulk scattering amplitudes. The purpose of this paper is to answer this question and to investigate the properties of amplitudes in theories which have an associative celestial OPE at tree level.

The paper is organized as follows. In Section 2 we briefly review the constraints obtained by imposing the Jacobi identity on the algebra of soft modes [9]. We then rephrase the question of tree-level associativity directly at the level of bulk scattering amplitudes in Section 3 and show that they yield the same constraints on the bulk couplings. In Section 4 we demonstrate that these constraints lead to the vanishing of certain four-point amplitudes and we work out the consequences on amplitudes of higher multiplicity. We close with a discussion of various open questions.

2 Review of the coupling constants constraints

We begin by listing all relevant three-point amplitudes and briefly reviewing the constraints among their couplings found in [9]. The most generic scenario involves a graviton, gluons and scalars. However, there are subsets of these particles which yield an associative OPE.

The first subset is the graviton-scalar sector, where we consider the usual Einstein-Hilbert term with coupling proportional to $\kappa_{-2,2,2}$, an R^3 interaction proportional to $\kappa_{2,2,2}$, an $R^2\phi$ interaction proportional to $\kappa_{0,2,2}$ and an $R\phi^2$ interaction proportional to $\kappa_{0,0,2}$. To be more precise, we specify this sector by the anti-holomorphic¹ three-point amplitudes

$$\begin{aligned}
& \begin{array}{c} 2^{++} \\ \diagdown \\ \text{---} \bigcirc \kappa_{-2,2,2} \text{---} \\ \diagup \\ 1^{++} \end{array} \text{---} 3^{--} = \kappa_{-2,2,2} \frac{[12]^6}{[23]^2 [13]^2} \\
& \begin{array}{c} 2^{++} \\ \diagdown \\ \text{---} \bigcirc \kappa_{0,2,2} \text{---} \\ \diagup \\ 1^{++} \end{array} \text{---} 3^\phi = \kappa_{0,2,2} [12]^4 \\
& \begin{array}{c} 2^{++} \\ \diagdown \\ \text{---} \bigcirc \kappa_{2,2,2} \text{---} \\ \diagup \\ 1^{++} \end{array} \text{---} 3^{++} = \kappa_{2,2,2} [12]^2 [23]^2 [13]^2 \\
& \begin{array}{c} 2^\phi \\ \diagdown \\ \text{---} \bigcirc \kappa_{0,0,2} \text{---} \\ \diagup \\ 1^{++} \end{array} \text{---} 3^\phi = \kappa_{0,0,2} \frac{[12]^2 [13]^2}{[23]^2}
\end{aligned} \tag{2.1}$$

together with their holomorphic parity conjugates. Note that in this paper we don't equate the coupling constants of the parity conjugate amplitudes with those of the original amplitude. Namely we regard $\kappa_{s_1, s_2, s_3} \neq \kappa_{-s_1, -s_2, -s_3}$ in general. The OPE of conformally soft operators satisfies the Jacobi identity only if the couplings are related by

$$(\kappa_{-2,2,2} - \kappa_{0,0,2}) \kappa_{0,2,2} = 0, \quad (\kappa_{-2,2,2} - \kappa_{0,0,2}) \kappa_{0,0,2} = 0, \tag{2.2}$$

$$3\kappa_{0,2,2}^2 = 10 \kappa_{-2,2,2} \kappa_{2,2,2}. \tag{2.3}$$

¹We call $\mathcal{A}(1^{s_1}, 2^{s_2}, 3^{s_3})$ anti-holomorphic if $s_1 + s_2 + s_3 > 0$ and holomorphic if $s_1 + s_2 + s_3 < 0$; see (A.11).

Jacobi identity also requires

$$(\kappa_{-2,2,2} - \kappa_{-1,1,2}) \kappa_{-1,1,2} = 0, \quad (\kappa_{-2,2,2} - \kappa_{-1,1,2}) \kappa_{1,1,2} = 0, \quad (2.8)$$

$$\kappa_{-1,1,2} \kappa_{1,1,2} = \kappa_{0,2,2} \kappa_{0,1,1}, \quad (\kappa_{0,0,2} - \kappa_{-1,1,2}) \kappa_{0,1,1} = 0, \quad (2.9)$$

$$\kappa_{-1,1,2} \kappa_{1,1,1} = 3\kappa_{1,1,2} \kappa_{-1,1,1}. \quad (2.10)$$

Note that all of the constraints presented above are consistent with dimensional analysis since κ_{s_1, s_2, s_3} has mass dimension $1 - |s_1 + s_2 + s_3|$.

3 Tree-level OPE associativity from amplitudes

The standard way to check OPE associativity is to introduce a mode expansion of the participating operators and then compute their commutators using the OPE. If the OPE is associative, the commutator of these modes must satisfy the Jacobi identity. In the context of CCFT, the bulk interpretation of this procedure is opaque. Our goal in this paper is not just to check OPE associativity, but also to interpret the condition directly at the level of momentum space scattering amplitudes. To derive a more direct check of OPE associativity consider the following identity involving contour integrals²:

$$\oint_{|z_{13}|=2} dz_1 \oint_{|z_{23}|=1} dz_2 \langle \mathcal{O}_{\Delta_1, s_1}(z_1, \bar{z}_1) \dots \mathcal{O}_{\Delta_n, s_n}(z_n, \bar{z}_n) \rangle = \oint_{|z_{23}|=2} dz_2 \oint_{|z_{13}|=1} dz_1 \langle \mathcal{O}_{\Delta_1, s_1}(z_1, \bar{z}_1) \dots \mathcal{O}_{\Delta_n, s_n}(z_n, \bar{z}_n) \rangle + \oint_{|z_{23}|=2} dz_2 \oint_{|z_{12}|=1} dz_1 \langle \mathcal{O}_{\Delta_1, s_1}(z_1, \bar{z}_1) \dots \mathcal{O}_{\Delta_n, s_n}(z_n, \bar{z}_n) \rangle. \quad (3.1)$$

This is valid as long as all of the (Δ_i, s_i) are such that the correlation functions are single-valued functions of z_i . Practically, we evaluate these integrals using the OPE. Thus the identity serves as a check on the OPE associativity. We can write this in a condensed form as

$$\left[\text{Res}_{2 \rightarrow 3} \text{Res}_{1 \rightarrow 2} - \text{Res}_{1 \rightarrow 3} \text{Res}_{2 \rightarrow 3} + \text{Res}_{2 \rightarrow 3} \text{Res}_{1 \rightarrow 3} \right] \langle \mathcal{O}_{\Delta_1, s_1}(z_1, \bar{z}_1) \dots \mathcal{O}_{\Delta_n, s_n}(z_n, \bar{z}_n) \rangle = 0. \quad (3.2)$$

The correlator in (3.2) is nothing but the Mellin transform of the amplitude. If we parameterize the momenta of outgoing massless external particles as

$$p_i^\mu \sim \lambda_i \tilde{\lambda}_i \quad \text{with} \quad \lambda_i = \sqrt{2\omega_i} \begin{pmatrix} 1 \\ z_i \end{pmatrix} \quad \text{and} \quad \tilde{\lambda}_i = \sqrt{2\omega_i} \begin{pmatrix} 1 \\ \bar{z}_i \end{pmatrix}, \quad (3.3)$$

the transformation from momentum to boost eigenstates is implemented via the Mellin transform

$$\langle \mathcal{O}_{\Delta_1, s_1}(z_1, \bar{z}_1) \dots \mathcal{O}_{\Delta_n, s_n}(z_n, \bar{z}_n) \rangle = \int \prod_{i=1}^n \frac{d\omega_i}{\omega_i^{1-\Delta_i}} \mathcal{A}_n \left(\left\{ \lambda_1, \tilde{\lambda}_1 \right\}^{s_1}, \left\{ \lambda_2, \tilde{\lambda}_2 \right\}^{s_2}, \dots, \left\{ \lambda_n, \tilde{\lambda}_n \right\}^{s_n} \right). \quad (3.4)$$

Demanding

$$\left[\text{Res}_{2 \rightarrow 3} \text{Res}_{1 \rightarrow 2} - \text{Res}_{1 \rightarrow 3} \text{Res}_{2 \rightarrow 3} + \text{Res}_{2 \rightarrow 3} \text{Res}_{1 \rightarrow 3} \right] \mathcal{A}_n \left(\left\{ \lambda_1, \tilde{\lambda}_1 \right\}^{s_1}, \left\{ \lambda_2, \tilde{\lambda}_2 \right\}^{s_2}, \dots, \left\{ \lambda_n, \tilde{\lambda}_n \right\}^{s_n} \right) = 0 \quad (3.5)$$

²A similar analysis of the associativity of the one-loop OPE in self-dual Yang-Mills was performed in [13].

ensures that the same is true of the correlator. The upshot is that we can check associativity of the celestial OPE by evaluating double residues directly on the amplitude rather than dealing with the celestial correlators.

The first step is to isolate the collinear limits. Bearing in mind that we will later be interested in examining the effects of (3.5) on amplitudes with various higher derivative corrections, we will analyze the collinear limit using the all-line shift recursion relations [14, 15], reviewed briefly in Appendix A. We begin by using (A.10) to isolate the collinear channel as the momentum of particle one approaches that of particle two

$$\begin{aligned} & \mathcal{A}_n \left(\left\{ \lambda_1, \tilde{\lambda}_1 \right\}^{s_1}, \left\{ \lambda_2, \tilde{\lambda}_2 \right\}^{s_2}, \dots, \left\{ \lambda_n, \tilde{\lambda}_n \right\}^{s_n} \right) \\ &= \sum_{s_I} \hat{\mathcal{A}}_3 \left(\left\{ \hat{\lambda}_1, \tilde{\lambda}_1 \right\}^{s_1}, \left\{ \hat{\lambda}_2, \tilde{\lambda}_2 \right\}^{s_2}, \left\{ \hat{\lambda}_I, \hat{\lambda}_I \right\}^{-s_I} \right) \frac{1}{\langle 12 \rangle [12]} \hat{\mathcal{A}}_{n-1} \left(\left\{ \hat{\lambda}_I, \hat{\lambda}_I \right\}^{s_I}, \dots, \left\{ \hat{\lambda}_n, \tilde{\lambda}_n \right\}^{s_n} \right) \\ & \quad + \{\text{other channels}\}. \end{aligned} \quad (3.6)$$

In terms of these, the residue is

$$\text{Res}_{z_1 \rightarrow z_2} \mathcal{A}_n = \sum_{s_I} \mathcal{A}_3 \left(\tilde{\lambda}_1^{s_1}, \tilde{\lambda}_2^{s_2}, \tilde{\lambda}_I^{-s_I} \right) \frac{1}{2\sqrt{\omega_1 \omega_2} [12]} \mathcal{A}_{n-1} \left(\left\{ \lambda_I, \tilde{\lambda}_I \right\}^{s_I}, \dots, \left\{ \lambda_n, \tilde{\lambda}_n \right\}^{s_n} \right) \quad (3.7)$$

where, on the right-hand side, in the limit $z_{12} = 0$ we have

$$\lambda_1 = \sqrt{\frac{\omega_1}{\omega_1 + \omega_2}} \lambda_I, \quad \lambda_2 = \sqrt{\frac{\omega_2}{\omega_1 + \omega_2}} \lambda_I, \quad \tilde{\lambda}_I = \sqrt{\frac{\omega_1}{\omega_1 + \omega_2}} \tilde{\lambda}_1 + \sqrt{\frac{\omega_2}{\omega_1 + \omega_2}} \tilde{\lambda}_2. \quad (3.8)$$

In arriving at (3.7), we have made use of the fact that only anti-holomorphic three-point amplitudes contribute to the collinear limit (see Appendix A) and that the terms from other channels drop out in this limit. Furthermore, the subamplitude \mathcal{A}_{n-1} now depends only on the unshifted momenta as the deformation parameter α vanishes in the collinear limit (see (A.7)). Applying this formula a second time, to compute the double residue, therefore translates (3.5) into

$$\begin{aligned} & \sum_{s_{I_1}} \left[\frac{1}{\bar{z}_{12} \bar{z}_{I_1 3}} \frac{1}{\omega_1 + \omega_2} \mathcal{A}_3 \left(\tilde{\lambda}_1^{s_1}, \tilde{\lambda}_2^{s_2}, \tilde{\lambda}_{I_1}^{-s_{I_1}} \right) \mathcal{A}_3 \left(\tilde{\lambda}_{I_1}^{s_{I_1}}, \tilde{\lambda}_3^{s_3}, \tilde{\lambda}_{I_2}^{-s_{I_2}} \right) \right. \\ & \quad + \frac{1}{\bar{z}_{23} \bar{z}_{I_1 1}} \frac{1}{\omega_2 + \omega_3} \mathcal{A}_3 \left(\tilde{\lambda}_2^{s_2}, \tilde{\lambda}_3^{s_3}, \tilde{\lambda}_{I_1}^{-s_{I_1}} \right) \mathcal{A}_3 \left(\tilde{\lambda}_{I_1}^{s_{I_1}}, \tilde{\lambda}_1^{s_1}, \tilde{\lambda}_{I_2}^{-s_{I_2}} \right) \\ & \quad \left. + \frac{1}{\bar{z}_{31} \bar{z}_{I_1 2}} \frac{1}{\omega_1 + \omega_3} \mathcal{A}_3 \left(\tilde{\lambda}_3^{s_3}, \tilde{\lambda}_1^{s_1}, \tilde{\lambda}_{I_1}^{-s_{I_1}} \right) \mathcal{A}_3 \left(\tilde{\lambda}_{I_1}^{s_{I_1}}, \tilde{\lambda}_2^{s_2}, \tilde{\lambda}_{I_2}^{-s_{I_2}} \right) \right] = 0 \end{aligned} \quad (3.9)$$

with $\bar{z}_{I_1} = \frac{\omega_1 \bar{z}_1 + \omega_2 \bar{z}_2}{\omega_1 + \omega_2}$. (We emphasize that this equation must hold for arbitrary s_{I_2} .) Using the three-point amplitudes defined in the previous section, it is now a straightforward (if slightly tedious) exercise to show that (3.9) is true only when the constraints (2.2), (2.3), (2.5), (2.6) and (2.8)–(2.10) are satisfied.

4 Amplitudes in EFTs with celestial dual

In this section, we will examine the properties of amplitudes in effective theories that satisfy (3.9) or equivalently, the constraints reviewed in Section 2.

4.1 Four-point amplitudes in the graviton-scalar sector

We start with amplitudes involving external gravitons, which were computed in [16, 17]. The amplitude involving four positive helicity gravitons is³

$$\begin{aligned}
 \mathcal{A}_4(1^{++}, 2^{++}, 3^{++}, 4^{++}) = & \text{Diagram 1} + \text{Diagram 2} \\
 & + (\text{cyclic of } 1, 2, 3)
 \end{aligned}
 \tag{4.1}$$

where (cyclic of 1,2,3) refers to the diagrams obtained by making the stated replacements. Evaluating the diagrams results in

$$\begin{aligned}
 \mathcal{A}_4(1^{++}, 2^{++}, 3^{++}, 4^{++}) &= \kappa_{-2,2,2}\kappa_{2,2,2} \left(\frac{[12]^5[34]^2}{\langle 12 \rangle} \frac{\langle 1X \rangle^2 \langle 2X \rangle^2}{\langle 3X \rangle^2 \langle 4X \rangle^2} + \frac{[34]^5[12]^2}{\langle 34 \rangle} \frac{\langle 3X \rangle^2 \langle 4X \rangle^2}{\langle 1X \rangle^2 \langle 2X \rangle^2} \right) \\
 &+ \kappa_{0,2,2}^2 \frac{[12]^4[34]^4}{s_{12}} + (\text{cyclic of } 1, 2, 3) \\
 &= (10\kappa_{-2,2,2}\kappa_{2,2,2} - 3\kappa_{0,2,2}^2) s_{12}s_{13}s_{23} \frac{[12][23][34][41]}{\langle 12 \rangle \langle 23 \rangle \langle 34 \rangle \langle 41 \rangle} \\
 &= 0
 \end{aligned}
 \tag{4.2}$$

where we used $s_{ij} = \langle ij \rangle [ij]$. In the last line, we have imposed the constraint in (2.3) to conclude that the amplitude vanishes.

The remaining four-graviton amplitudes $\mathcal{A}(1^{++}, 2^{++}, 3^{++}, 4^{--})$, $\mathcal{A}(1^{++}, 2^{--}, 3^{--}, 4^{++})$, and their parity conjugates are not all-line shift constructible since they violate the condition (A.5). Instead, they were computed in [16–18]:

$$\begin{aligned}
 \mathcal{A}(1^{++}, 2^{++}, 3^{++}, 4^{--}) &= \kappa_{2,2,2}\kappa_{-2,-2,2} (\langle 14 \rangle [13] \langle 34 \rangle)^2 \frac{[12][23][31]}{\langle 12 \rangle \langle 23 \rangle \langle 31 \rangle}, \\
 \mathcal{A}(1^{++}, 2^{--}, 3^{--}, 4^{++}) &= \frac{(\langle 23 \rangle [14])^4}{s_{14}} \left(\kappa_{2,2,2}^2 s_{12} s_{13} - \kappa_{2,2,0}^2 + \kappa_{2,2,-2}^2 \frac{1}{s_{12} s_{13}} \right),
 \end{aligned}
 \tag{4.3}$$

and we note that these amplitudes are non-vanishing, even on the support of the associativity constraints.

Moving on to amplitudes involving external scalars in addition to gravitons, the only amplitude

³We will omit explicitly displaying the dependence of the amplitude on $\lambda, \tilde{\lambda}$ for brevity and only display the helicities.

that is all-line shift constructible is

$$\begin{aligned}
\mathcal{A}_4(1^{++}, 2^{++}, 3^{++}, 4^\phi) &= \text{Diagram 1} + \text{Diagram 2} \\
&\quad + (\text{cyclic of } 1,2,3) \\
&= \kappa_{-2,2,2}\kappa_{2,2,0} \frac{[12][34]^4}{\langle 12 \rangle} \frac{\langle 4X \rangle^4}{\langle 1X \rangle^2 \langle 2X \rangle^2} + \kappa_{2,2,0}\kappa_{0,0,2} \frac{[12]^3 [34]^2 \langle 4X \rangle^2}{\langle 12 \rangle \langle 3X \rangle^2} \\
&\quad + (\text{cyclic of } 1,2,3) \\
&= 0
\end{aligned} \tag{4.4}$$

and it vanishes on the support of the constraints (2.2).

4.2 Four-point amplitudes in the gluon-scalar sector

The situation in the gluon-scalar sector is similar: the constraints lead to the vanishing of all-line shift constructible amplitudes. In this sector the only all-line shift constructible all-gluon amplitude is

$$\begin{aligned}
\mathcal{A}_4(1^{+,a_1}, 2^{+,a_2}, 3^{+,a_3}, 4^{+,a_4}) &= \text{Diagram 1} \\
&\quad + \text{Diagram 2} + \text{Diagram 3} + (\text{cyclic of } 1,2,3)
\end{aligned} \tag{4.5}$$

A straightforward calculation yields

$$\begin{aligned}
&\mathcal{A}_4(1^{+,a_1}, 2^{+,a_2}, 3^{+,a_3}, 4^{+,a_4}) \\
&= -f^{a_1 a_2 e} f^{e a_3 a_4} \kappa_{-1,1,1} \kappa_{1,1,1} \left(\frac{[12]^2 [34]}{\langle 12 \rangle} \frac{\langle 1X \rangle \langle 2X \rangle}{\langle 3X \rangle \langle 4X \rangle} + \frac{[34]^2 [12]}{\langle 34 \rangle} \frac{\langle 3X \rangle \langle 4X \rangle}{\langle 1X \rangle \langle 2X \rangle} \right) \\
&\quad + \frac{2\kappa_{0,1,1}^2}{N_c} \left(\delta^{a_1 a_2} \delta^{a_3 a_4} \frac{[12]^2 [34]^2}{\langle 12 \rangle [12]} \right) + \kappa_{0,1,1}^2 \left(d^{a_1 a_2 b} d^{a_3 a_4 b} \frac{[12]^2 [34]^2}{\langle 12 \rangle [12]} \right) + (\text{cyclic of } 1,2,3) \\
&= (4\kappa_{-1,1,1} \kappa_{1,1,1} - 2\kappa_{0,1,1}^2) \frac{[13][24]^2}{\langle 13 \rangle} \text{Tr}[T^{a_1} T^{a_2} T^{a_3} T^{a_4}] + (\text{permutation of } 1,2,3).
\end{aligned} \tag{4.6}$$

Here we used properties of f^{abc} and d^{abc} , and in the last line the constraint $\kappa_{0,1,1}^2 = 2\kappa_{-1,1,1} \kappa_{1,1,1}$ from (2.6) to find that this amplitude vanishes as well. The remaining four-gluon amplitudes are not all-line shift constructible and are non-vanishing for generic couplings, even when all the constraints from Section 2 are imposed.

We also find that the four-point amplitude involving three positive helicity gluons and one scalar (either the adjoint or singlet) is vanishing. For the adjoint scalar, the full amplitude reads (here we

shift recursion relations to glue together two anti-holomorphic three-point amplitudes:

$$\begin{aligned}
& \mathcal{A}_4 \left(\left\{ \lambda_1, \tilde{\lambda}_1 \right\}^{s_1}, \left\{ \lambda_2, \tilde{\lambda}_2 \right\}^{s_2}, \left\{ \lambda_3, \tilde{\lambda}_3 \right\}^{s_3}, \left\{ \lambda_4, \tilde{\lambda}_4 \right\}^{s_4} \right) \\
&= \text{Diagram} + (\text{cyclic of } 1,2,3) \\
&= \sum_{s_I} \left[\mathcal{A}_3 \left(\tilde{\lambda}_1^{s_1}, \tilde{\lambda}_2^{s_2}, \hat{\lambda}_I^{-s_I} \right) \frac{1}{\langle 12 \rangle [12]} \mathcal{A}_3 \left(\hat{\lambda}_I^{s_I}, \tilde{\lambda}_3^{s_3}, \tilde{\lambda}_4^{s_4} \right) + (\text{cyclic of } 1,2,3) \right].
\end{aligned} \tag{4.9}$$

Using momentum conservation, we can rewrite this as

$$\begin{aligned}
& \frac{[34]}{\langle 12 \rangle} \sum_{s_I} \left[\mathcal{A}_3 \left(\tilde{\lambda}_1^{s_1}, \tilde{\lambda}_2^{s_2}, \hat{\lambda}_I^{-s_I} \right) \frac{1}{[12] [34]} \mathcal{A}_3 \left(\hat{\lambda}_I^{s_I}, \tilde{\lambda}_3^{s_3}, \tilde{\lambda}_4^{s_4} \right) \right. \\
& \quad + \mathcal{A}_3 \left(\tilde{\lambda}_1^{s_1}, \tilde{\lambda}_3^{s_3}, \hat{\lambda}_I^{-s_I} \right) \frac{1}{[31] [24]} \mathcal{A}_3 \left(\hat{\lambda}_I^{s_I}, \tilde{\lambda}_2^{s_2}, \tilde{\lambda}_4^{s_4} \right) \\
& \quad \left. + \mathcal{A}_3 \left(\tilde{\lambda}_1^{s_1}, \tilde{\lambda}_2^{s_2}, \hat{\lambda}_I^{-s_I} \right) \frac{1}{[14] [23]} \mathcal{A}_3 \left(\hat{\lambda}_I^{s_I}, \tilde{\lambda}_3^{s_3}, \tilde{\lambda}_4^{s_4} \right) \right],
\end{aligned} \tag{4.10}$$

which must hold for arbitrary s_4 . This is precisely the double residue condition (3.9) on the support of four-point momentum conservation! Hence we conclude that tree-level associativity of the celestial OPE forces any four-point amplitude constructed solely from anti-holomorphic vertices to vanish⁴.

The construction of higher-point amplitudes proceeds in a similar manner. However, unlike the four-particle case, we can encounter amplitudes that are all-line shift constructible but non-vanishing. This can occur because none of the non-zero four-point amplitudes are purely anti-holomorphic or purely holomorphic, so any higher-point amplitudes (when constructed via all-line shift recursion) that receive contributions from these amplitudes will also be non-vanishing. An example is the five-point all-plus amplitude $\mathcal{A}_5(1^+, 2^+, 3^+, 4^+, 5^+)$, which is all-line shift constructible but has a term involving the product of two non-zero amplitudes $\mathcal{A}_3(1^+, 2^+, 3^+) \times \mathcal{A}_4(1^-, 2^+, 3^+, 4^+)$, the second of which includes both holomorphic and anti-holomorphic vertices. However, constructible higher-point amplitudes will be vanishing at tree level if they do not receive contributions from any channels other than those involving only purely holomorphic or anti-holomorphic vertices as building blocks.

5 Discussion

The constraints (2.2), (2.3), (2.5), (2.6) and (2.8)–(2.10) are incredibly restrictive. Most apparently sensible theories fail to satisfy them. Heterotic string theory (compactified on a torus to 4D) ostensibly fails to satisfy them due to the presence of $R^2\phi$ terms and the absence of an R^3 term.⁵ However, the four-positive helicity graviton amplitude vanishes in this theory as can be seen from the double copy construction [19]. This is because $\kappa_{2,2,2}$ vanishes and the two scalars (axion and dilaton) give

⁴The apparent asymmetry between holomorphic and anti-holomorphic in our discussion arises from our choice of using the holomorphic all-line shift in Appendix A. Of course, all conclusions hold for parity conjugate amplitudes as well, and could be manifested by using an anti-holomorphic all-line shift instead. In constructing higher-point amplitudes recursively, one has the freedom to independently choose the holomorphic or anti-holomorphic shift term-by-term and in various levels of the recursion.

⁵We thank Mina Himwich for pointing this out.

contributions to the left-hand side of (2.3) that vanish when summed. While the vanishing of the four-positive helicity amplitude does not guarantee associativity, it serves as an easy (particularly when the spectrum involves multiple particles with the same helicity) but powerful check on OPE associativity. It is intriguing to wonder about other contributions (massive particles, bound states, resonances, extended objects) to the OPE of massless particles, see for e.g. [20]. If some or all of these contributions are indeed present, this suggests that the OPE in (1.1) is incomplete and these additional contributions might modify the analysis in this paper. Alternatively, theories which fail to satisfy these constraints simply do not have celestial duals.

Interestingly, one other example of a theory which does satisfy the constraints is the chiral higher-spin theory studied in [21–23]. Working in the light-cone approach, they derived the following solution for the coupling constants after requiring Poincaré symmetry:

$$\kappa_{s_1, s_2, s_3} \sim \frac{(l_P)^{s_1+s_2+s_3-1}}{\Gamma(s_1+s_2+s_3)} \quad s_1+s_2+s_3 > 0, \quad (5.1)$$

where l_P is a parameter with dimension of length. One can easily check that (5.1) satisfies all the constraints in (2.2), (2.3), (2.5), (2.6) and (2.8)–(2.10).

In light of [13, 24] it is natural to speculate about the prospect of a self-dual theory on the support of the associativity constraints, generalizing self-dual Yang-Mills and self-dual gravity with some higher derivative corrections. The key property of self-dual Yang-Mills and self-dual gravity is that only the anti-holomorphic three-point vertices are nonzero, and all higher-point amplitudes vanish at tree level. After including the higher-derivative interactions, if we continue to keep only the anti-holomorphic three-point vertices and set the holomorphic three-point vertices to zero, then all higher-point tree-level amplitudes will be vanishing only when the constraints are satisfied as we have shown. We leave the study of such possibilities for future work.

Acknowledgments

We are grateful to the organizers and participants of the workshop “Possible and Impossible in Effective Field Theory: From S-Matrix to the Swampland”, where part of this work was presented, for useful comments and discussion. We also thank Rishabh Bhardwaj, Shounak De, Mina Himwich, Yangrui Hu, Konstantinos Koutrolikos, Luke Lippstreu, Xianlong Liu, Jorge Mago, Oliver Schlotterer and Andy Strominger for stimulating comments and correspondence. This work was supported in part by the US Department of Energy under contract DE-SC0010010 Task F and by Simons Investigator Award #376208 (AV).

A All-line shift recursion relations

In this appendix we will review the all-line shift recursion relations following [16].

The all-line shift recursion relations are based on shifting all of the external momenta. For the purpose this appendix, we treat $\lambda, \tilde{\lambda}$ (consequently z, \bar{z}) as independent complex variables. We will refer to them as holomorphic and anti-holomorphic respectively. We denote a scattering amplitude involving n massless particles (all considered outgoing) with momenta $p_1 = \lambda_1 \tilde{\lambda}_1, \dots, p_n = \lambda_n \tilde{\lambda}_n$ and helicities s_1, \dots, s_n by $\mathcal{A}_n \left(\left\{ \lambda_1, \tilde{\lambda}_1 \right\}^{s_1}, \dots, \left\{ \lambda_n, \tilde{\lambda}_n \right\}^{s_n} \right)$. Consider an all-line holomorphic shift

$$\hat{\lambda}_i = \lambda_i + \alpha w_i X \quad i = 1, \dots, n \quad (\text{A.1})$$

where X is an arbitrary reference spinor, α is the deformation parameter and the w_i are chosen to satisfy momentum conservation

$$\sum_{i=1}^n w_i \tilde{\lambda}_i = 0. \quad (\text{A.2})$$

The deformed amplitude has the following large α behavior [16]

$$\hat{\mathcal{A}}_n(\alpha) \rightarrow \alpha^a \quad \text{as} \quad \alpha \rightarrow \infty \quad \text{with} \quad 2a = 4 - n - c - \sum_{i=1}^n s_i, \quad (\text{A.3})$$

where c is the mass dimension of the product of couplings in the amplitude. The undeformed amplitude can be related to its residues at non-zero values of α via the residue theorem


$$\mathcal{A}_n(\alpha = 0) = \oint_{\alpha=0} \frac{d\alpha}{\alpha} \mathcal{A}_n(\alpha) = - \sum_j \text{Res}_{\alpha=\alpha_j} \left[\frac{1}{\alpha} \mathcal{A}_n \right]. \quad (\text{A.4})$$

In order to use the all-line shift recursion to compute amplitudes recursively, it is crucial that there be no contribution to (A.4) from a residue at infinity. From (A.3) we see that this will be the case as long as

$$4 - n - c - \sum_{i=1}^n s_i < 0. \quad (\text{A.5})$$

When this holds, then the tree-level deformed amplitude only has simple poles which occur when the sum of a subset of the external momenta goes on shell. The amplitude factorizes into lower-point subamplitudes on the residue of the pole.

In the paper, our main interest lies in the collinear channel where we have



$$(\text{A.6})$$

The value of α that corresponds to this channel is the one that makes the intermediate momentum P_I go on shell:

$$\hat{P}_I^2 = \langle \hat{1} \hat{2} \rangle [12] = 0 \quad \implies \quad \alpha = - \frac{\langle 12 \rangle}{w_1 \langle X2 \rangle - w_2 \langle X1 \rangle}, \quad (\text{A.7})$$

and we can write

$$\hat{\lambda}_1 = \frac{\langle X1 \rangle}{w_1 \langle X2 \rangle - w_2 \langle X1 \rangle} (w_1 \lambda_2 - w_2 \lambda_1), \quad \hat{\lambda}_2 = \frac{\langle X2 \rangle}{w_1 \langle X2 \rangle - w_2 \langle X1 \rangle} (w_1 \lambda_2 - w_2 \lambda_1), \quad (\text{A.8})$$

$$\hat{\lambda}_I = w_1 \lambda_2 - w_2 \lambda_1, \quad \hat{\lambda}_I = \frac{\langle X1 \rangle \tilde{\lambda}_1 + \langle X2 \rangle \tilde{\lambda}_2}{w_1 \langle X2 \rangle - w_2 \langle X1 \rangle}. \quad (\text{A.9})$$

We therefore have

$$\begin{aligned} & \mathcal{A}_n \left(\left\{ \lambda_1, \tilde{\lambda}_1 \right\}^{s_1}, \left\{ \lambda_2, \tilde{\lambda}_2 \right\}^{s_2}, \dots, \left\{ \lambda_n, \tilde{\lambda}_n \right\}^{s_n} \right) \\ &= \sum_{s_I} \hat{\mathcal{A}}_3 \left(\left\{ \hat{\lambda}_1, \tilde{\lambda}_1 \right\}^{s_1}, \left\{ \hat{\lambda}_2, \tilde{\lambda}_2 \right\}^{s_2}, \left\{ \hat{\lambda}_I, \hat{\lambda}_I \right\}^{-s_I} \right) \frac{1}{\langle 12 \rangle [12]} \hat{\mathcal{A}}_{n-1} \left(\left\{ \hat{\lambda}_I, \hat{\lambda}_I \right\}^{s_I}, \dots, \left\{ \hat{\lambda}_n, \tilde{\lambda}_n \right\}^{s_n} \right) \\ & \quad + \{\text{other channels}\} \end{aligned} \quad (\text{A.10})$$

The other channels may include other collinear (for e.g. 34 collinear) as well as multiparticle poles.

The recursive computation of amplitudes is seeded by three-point amplitudes, which are completely fixed by Lorentz invariance and little group scaling to be⁶

$$A(1^{s_1}, 2^{s_2}, 3^{s_3}) = \begin{cases} \kappa_{s_1, s_2, s_3} [12]^{s_1+s_2-s_3} [23]^{s_2+s_3-s_1} [31]^{s_3+s_1-s_2}, & \text{if } s_1 + s_2 + s_3 > 0, \\ \kappa_{s_1, s_2, s_3} \langle 12 \rangle^{s_3-s_1-s_2} \langle 23 \rangle^{s_1-s_2-s_3} \langle 31 \rangle^{s_2-s_1-s_3}, & \text{if } s_1 + s_2 + s_3 < 0. \end{cases} \quad (\text{A.11})$$

A crucial aspect of the holomorphic shift is that the holomorphic three-point amplitudes vanish. This follows from the proportionality of the three holomorphic spinors $\hat{\lambda}_1$, $\hat{\lambda}_2$ and $\hat{\lambda}_I$ in (A.8), (A.9). Thus, only anti-holomorphic three-point amplitudes contribute in the collinear limit.

References

- [1] S. Pasterski, S.-H. Shao and A. Strominger, *Flat Space Amplitudes and Conformal Symmetry of the Celestial Sphere*, *Phys. Rev. D* **96** (2017) 065026 [[1701.00049](#)].
- [2] S. Pasterski and S.-H. Shao, *Conformal basis for flat space amplitudes*, *Phys. Rev. D* **96** (2017) 065022 [[1705.01027](#)].
- [3] S. Pasterski, S.-H. Shao and A. Strominger, *Gluon Amplitudes as 2d Conformal Correlators*, *Phys. Rev. D* **96** (2017) 085006 [[1706.03917](#)].
- [4] W. Fan, A. Fotopoulos and T.R. Taylor, *Soft Limits of Yang-Mills Amplitudes and Conformal Correlators*, *JHEP* **05** (2019) 121 [[1903.01676](#)].
- [5] M. Pate, A.-M. Raclariu, A. Strominger and E.Y. Yuan, *Celestial operator products of gluons and gravitons*, *Rev. Math. Phys.* **33** (2021) 2140003 [[1910.07424](#)].
- [6] E. Himwich, M. Pate and K. Singh, *Celestial Operator Product Expansions and $w_{1+\infty}$ Symmetry for All Spins*, [2108.07763](#).
- [7] A. Guevara, E. Himwich, M. Pate and A. Strominger, *Holographic symmetry algebras for gauge theory and gravity*, *JHEP* **11** (2021) 152 [[2103.03961](#)].
- [8] A. Strominger, *$w_{1+\infty}$ and the Celestial Sphere*, [2105.14346](#).
- [9] J. Mago, L. Ren, A.Y. Srikant and A. Volovich, *Deformed $w_{1+\infty}$ Algebras in the Celestial CFT*, [2111.11356](#).
- [10] S. He, Y.-t. Huang and C. Wen, *Loop Corrections to Soft Theorems in Gauge Theories and Gravity*, *JHEP* **12** (2014) 115 [[1405.1410](#)].
- [11] P. Di Vecchia, R. Marotta and M. Mojaza, *Subsubleading soft theorems of gravitons and dilatons in the bosonic string*, *JHEP* **06** (2016) 054 [[1604.03355](#)].
- [12] H. Elvang, C.R.T. Jones and S.G. Naculich, *Soft Photon and Graviton Theorems in Effective Field Theory*, *Phys. Rev. Lett.* **118** (2017) 231601 [[1611.07534](#)].
- [13] K. Costello and N.M. Paquette, *On the associativity of one-loop corrections to the celestial OPE*, [2204.05301](#).
- [14] F. Cachazo, P. Svrcek and E. Witten, *MHV vertices and tree amplitudes in gauge theory*, *JHEP* **09** (2004) 006 [[hep-th/0403047](#)].
- [15] H. Elvang, D.Z. Freedman and M. Kiermaier, *Proof of the MHV vertex expansion for all tree amplitudes in $N=4$ SYM theory*, *JHEP* **06** (2009) 068 [[0811.3624](#)].

⁶In (2.4) and (2.7) we define certain κ 's with different overall normalizations compared to this standard.

- [16] T. Cohen, H. Elvang and M. Kiermaier, *On-shell constructibility of tree amplitudes in general field theories*, *JHEP* **04** (2011) 053 [[1010.0257](#)].
- [17] J. Broedel and L.J. Dixon, *Color-kinematics duality and double-copy construction for amplitudes from higher-dimension operators*, *JHEP* **10** (2012) 091 [[1208.0876](#)].
- [18] D. Bai and Y.-H. Xing, *Higher Derivative Theories for Interacting Massless Gravitons in Minkowski Spacetime*, *Nucl. Phys. B* **932** (2018) 15 [[1610.00241](#)].
- [19] T. Azevedo, M. Chiodaroli, H. Johansson and O. Schlotterer, *Heterotic and bosonic string amplitudes via field theory*, *JHEP* **10** (2018) 012 [[1803.05452](#)].
- [20] D. García-Sepúlveda, A. Guevara, J. Kulp and J. Wu, *Notes on Resonances and Unitarity from Celestial Amplitudes*, [2205.14633](#).
- [21] R.R. Metsaev, *S matrix approach to massless higher spins theory. 2: The Case of internal symmetry*, *Mod. Phys. Lett. A* **6** (1991) 2411.
- [22] R.R. Metsaev, *Poincare invariant dynamics of massless higher spins: Fourth order analysis on mass shell*, *Mod. Phys. Lett. A* **6** (1991) 359.
- [23] D. Ponomarev and E.D. Skvortsov, *Light-Front Higher-Spin Theories in Flat Space*, *J. Phys. A* **50** (2017) 095401 [[1609.04655](#)].
- [24] A. Ball, S.A. Narayanan, J. Salzer and A. Strominger, *Perturbatively exact $w_{1+\infty}$ asymptotic symmetry of quantum self-dual gravity*, *JHEP* **01** (2022) 114 [[2111.10392](#)].