

# ANALYSIS OF CURVATURE APPROXIMATIONS VIA COVARIANT CURL AND INCOMPATIBILITY FOR REGGE METRICS

J. GOPALAKRISHNAN, M. NEUNTEUFEL, J. SCHÖBERL, AND M. WARDETZKY

**ABSTRACT.** The metric tensor of a Riemannian manifold can be approximated using Regge finite elements and such approximations can be used to compute approximations to the Gauss curvature and the Levi-Civita connection of the manifold. It is shown that certain Regge approximations yield curvature and connection approximations that converge at a higher rate than previously known. The analysis is based on covariant (distributional) curl and incompatibility operators which can be applied to piecewise smooth matrix fields whose tangential-tangential component is continuous across element interfaces. Using the properties of the canonical interpolant of the Regge space, we obtain superconvergence of approximations of these covariant operators. Numerical experiments further illustrate the results from the error analysis.

**Key words:** Gauss curvature, Regge calculus, finite element method, differential geometry

**MSC2020:** 65N30, 53A70, 83C27

## 1. INTRODUCTION

This paper is concerned with the finite element approximation of the Gauss curvature  $K$  of a two-dimensional Riemannian manifold. As shown by Gauss's Theorema Egregium,  $K$  is an intrinsic quantity of the manifold. It can be computed solely using the metric tensor of the manifold. Therefore, when a finite element approximation of the metric tensor is given, it is natural to ask if an approximation to  $K$  can be computed. The answer was given in the affirmative by the recent works of [27] and [11], assuming that the metric is approximated using Regge finite elements. This paper is heavily based on these works and is devoted to proving that the resulting curvature and connection approximations converge at a higher rate than previously known.

The Regge finite element takes its name from Regge calculus, originally developed for solving Einstein field equations in general relativity. It discretizes the metric tensor through edge-length specifications, allowing the curvature to be approximated by means of angle deficits [39]. Regge calculus was established in theoretical and numerical physics and routinely finds applications in relativity and quantum mechanics. In [49, 40, 10] a comprehensive overview of the development of Regge calculus over the last fifty years can be found. Just as Whitney forms [48] can be interpreted as finite elements, it was observed that Regge's approach of prescribing quantities on edges is equivalent to defining a piecewise constant metric tensor whose tangential-tangential components are continuous across element interfaces [43]. The first rigorous proof of

convergence of Regge's angle deficits to the scalar curvature, for a sequence of appropriate triangulations in the sense of measures, was accomplished in [16]. Later, it was also shown [20] that for a given metric in the lowest order Regge finite element space, a sequence of mollified metrics converges to the angle deficit in the sense of measures. Methods based on angle deficits for approximating the Gauss curvature on triangulations consisting of piecewise flat triangles is well-established in discrete differential geometry and computer graphics. On specific triangulations satisfying certain conditions, convergence in the  $L^\infty$ -norm up to quadratic order was proven, but for a general irregular grid there is no reason to expect convergence [13, 50, 51]. In [34], Regge's concept of angle deficits has been extended to quadrilateral meshes. Notable among the results applicable for higher dimensional manifolds is the proof of convergence for approximated Ricci curvatures of isometrically embedded hypersurfaces  $\subset \mathbb{R}^{n+1}$  presented in [24], and used later for Ricci flows [25].

Another natural perspective to place the modern developments on the Regge finite element is within the emergence of *finite element exterior calculus* (FEEC) [7, 6]. Finite element structures for Regge calculus were developed in [18, 19] and the resulting elements became popular in FEEC under the name Regge finite elements [33]. Regge elements approximating metric and strain tensors were extended to arbitrary polynomial order on triangles, tetrahedra, and higher dimensional simplices in [33], and for quadrilaterals, hexahedra, and prisms in [35]. The utility of Regge elements when discretizing parts of the Kröner complex, or the elasticity complex, was considered in [8, 19, 28]. Properties of Regge elements were exploited to construct a method avoiding membrane locking for general triangular shell elements [36].

In this backdrop, the recent work of [27] provides a new and interesting application of Regge elements by developing a high-order Gauss curvature approximation formula based on higher degree Regge elements (and applied to Ricci and Ricci-DeTurck flow [26]). The key is an integral formulation of the angle deficit, extendable to higher orders. Using it, rigorous proofs of convergence at specific rates were proved in [27]. Even more recently, in [11], this approach has been reformulated in terms of a nonlinear distributional curvature and connection 1-form (Levi-Civita connection), using the element-wise Gauss curvature, jump of geodesic curvature at edges, and angle deficits at vertices as sources of curvature [44]. The authors show that  $L^2$ -convergence of the approximated curvature can be obtained if Regge elements using polynomials of degree at least two are used to approximate the metric. This is in line with the rule of thumb that a second order differential operator approximated using polynomials of degree  $k$  leads to convergence rates of order  $k - 2$ . Nonetheless, convergence rates better than this rule of thumb have often been observed in compatible discretizations in FEEC. One of our goals in this paper is to establish such an improved rate for the curvature approximation.

In a later section, we extend the ideas in [27, 11] by exploiting certain orthogonality properties for the error in the canonical interpolation by Regge finite elements to obtain one extra order of convergence for the curvature approximation. This extra order is comparable to super-convergence properties of mixed methods [12, 22] and has

been observed for the Hellan–Herrmann–Johnson for the biharmonic plate and shell equation [9, 46].

Another significant difference in our analysis, in comparison to [27, 11], is the use of the intrinsic (or covariant) incompatibility operator (which we define using the covariant curl on the manifold). It is now well known that linearizing the curvature operator around the Euclidean metric gives a first order term involving the incompatibility operator [19] and we exploit this relationship in the analysis of the curvature approximation. On Euclidean manifolds, the incompatibility operator is well known to be the natural differential operator for Regge elements in any dimension. By showing that the curvature approximation can be analyzed via the incompatibility operator, we hope to generate new ideas for computing and analyzing approximations of the intrinsic curvature tensor of higher dimensional manifolds. The incompatibility operator also arises in modeling elastic materials with dislocations [2, 3], another potential area of application.

This paper can be read linearly, but we have structured it so a numerical analyst can also directly start with the error analysis in Section 6, referring back to the previous sections as needed. The next section (§2) establishes notation and introduces preliminaries on differential geometry and finite element spaces. Section 3 defines the curvature approximation formula and also provides formulas in coordinates that we used for numerical computations. In Section 4, covariant differential operators on piecewise smooth metric tensors are derived, concentrating on the covariant curl and incompatibility operator. The linearization of the curvature and the connection 1-form, presented in §5, reveals their relationships to the covariant inc and curl. Section 6 is devoted to the numerical analysis of the errors in the method. The analysis is performed by first proving optimal convergence rates for the distributional covariant curl and inc, and then for the approximations of the Gauss curvature and connection 1-form. Numerical examples illustrating the theoretical results are presented in §7.

## 2. NOTATION AND PRELIMINARIES

This section provides definitions that we use throughout. We give intrinsic definitions of quantities on a manifold, but in view of our computational goals, we also make extensive use of coordinate expressions. We use the Einstein summation convention, by which a term where the same integer index appears twice, as both an upper and a lower subscript, is tacitly assumed to be summed over the values of that index in  $\{1, 2\}$ . Summation convention does not apply when a repeated index is not an integer (such as when a subscript or a superscript represents a vector field or other non-integer quantities).

**2.1. Spaces on the manifold.** Let  $M$  denote a two-dimensional oriented manifold with or without boundary. Endowed with a smooth metric  $\bar{g}$ , let  $(M, \bar{g})$  be a Riemannian manifold. Let the unique Levi-Civita connection generated by  $\bar{g}$  be denoted by  $\bar{\nabla}$ . Let  $\mathfrak{X}(M)$ ,  $\Lambda^k(M)$ , and  $\mathcal{T}_l^k(M)$  denote, respectively, the sets of smooth vector fields on  $M$ ,  $k$ -form fields on  $M$ , and  $(k, l)$ -tensor fields on  $M$ . The value of a tensor  $\rho \in \mathcal{T}_l^k(M)$  acting on  $k$  vectors  $X_i \in \mathfrak{X}(M)$  and  $l$  covectors  $\mu_j \in \Lambda^1(M)$  is denoted by  $\rho(\mu_1, \dots, \mu_l, X_1, \dots, X_k)$ . Note that  $\Lambda^1(M) = \mathcal{T}_0^1(M)$  and  $\mathfrak{X}(M) = \mathcal{T}_1^0(M)$ . Note also

that it is standard to extend the Levi-Civita connection  $\bar{\nabla}$  from vector fields to tensor fields (see e.g., [31, Lemma 4.6]) so that Leibniz rule holds.

For coordinate computations, we use a chart to move locally to a Euclidean domain with coordinates  $x^1, x^2$ . Let the accompanying coordinate frame and coframe be denoted by  $\partial_i$  and  $dx^i$ . We assume these coordinates preserve orientation, so the orientation of  $M$  is given by the ordering  $(\partial_1, \partial_2)$ . Let  $\mathcal{S}(M) = \{\sigma \in \mathcal{T}_0^2(M) : \sigma(X, Y) = \sigma(Y, X) \text{ for } X, Y \in \mathfrak{X}(M)\}$  and  $\mathcal{S}^+(M) = \{\sigma \in \mathcal{S}(M) : \sigma(X, X) > 0 \text{ for } X \in \mathfrak{X}(M)\}$ . They represent the subspace of symmetric tensors in  $\mathcal{T}_0^2(M)$ , whose elements  $\sigma$  can be expressed in coordinates as  $\sigma = \sigma_{ij} dx^i \otimes dx^j$  with smoothly varying coefficients  $\sigma_{ij}$  satisfying  $\sigma_{ij} = \sigma_{ji}$  and are additionally positive definite, respectively.

We will use standard operations on 2-manifold spaces such as the Hodge star  $\star : \Lambda^k(M) \rightarrow \Lambda^{2-k}(M)$ , the exterior derivative  $d^k : \Lambda^k(M) \rightarrow \Lambda^{k+1}(M)$ , the tangent to cotangent isomorphism  $\flat : \mathfrak{X}(M) \rightarrow \Lambda^1(M)$ , and the reverse operation  $\sharp : \Lambda^1(M) \rightarrow \mathfrak{X}(M)$ . Their definitions can be found in standard texts [31, 37, 45].

**2.2. Curvature.** The exact metric  $\bar{g}$  is an element of  $\mathcal{S}^+(M)$ . We define the *Riemann curvature tensor*  $\bar{R} \in \mathcal{T}_0^4(M)$  of the manifold following [31],

$$\bar{R}(X, Y, Z, W) = \bar{g}(\bar{\nabla}_X \bar{\nabla}_Y Z - \bar{\nabla}_Y \bar{\nabla}_X Z - \bar{\nabla}_{[X, Y]} Z, W), \quad X, Y, Z, W \in \mathfrak{X}(M). \quad (2.1)$$

The *Gauss curvature* of  $M$  can be expressed using any basis  $u, v$  of the tangent space by

$$K(\bar{g}) = \frac{\bar{R}(u, v, v, u)}{\bar{g}(u, u)\bar{g}(v, v) - \bar{g}(u, v)^2} \quad (2.2)$$

whose value is well known to be independent of the choice of the basis (see, e.g., [31, p. 144] or [15, Ch. 4, Proposition 3.1]).

We will also need the *geodesic curvature* along a curve  $\Gamma$  in the manifold  $(M, \bar{g})$ . To recall its standard definition (see [45, p. 140] or [31]), we let  $0 < s < a$  be the  $\bar{g}$ -arclength parameter so that  $\Gamma$  is described by  $\mu(s)$  for some smooth  $\mu : [0, a] \rightarrow M$  and its  $\bar{g}$ -unit tangent vector is  $T(s) = d\mu/ds$ . Let  $N(s)$  be such that  $(T(s), N(s))$  is a  $\bar{g}$ -orthonormal set of two vectors in the tangent space whose orientation is the same as that of  $M$ , i.e.,  $dx^1 \wedge dx^2(T(s), N(s)) > 0$ . Then

$$\kappa(\bar{g}) = \bar{g}(\bar{\nabla}_{T(s)} T(s), N(s)) \quad (2.3)$$

gives the geodesic curvature at the point  $\mu(s)$  of  $\Gamma$ .

**2.3. Connection 1-forms.** Let  $\{E_i\}_{i=1}^2$  be a local frame on an open subset  $U$  of the 2-manifold  $M$ . Then there is a uniquely determined  $2 \times 2$  matrix of 1-forms  $\{\varpi_i^j\}_{i,j=1}^2$ ,  $\varpi_i^j \in \Lambda^1(U)$  such that

$$\bar{\nabla}_X E_i = \varpi_i^j(X) E_j, \quad \text{for all } X \in \mathfrak{X}(M), i = 1, 2. \quad (2.4)$$

The  $\varpi_i^j$  are called *connection 1-forms*. If the frame  $\{E_i\}_{i=1}^2$  is chosen to be  $\bar{g}$ -orthonormal then the skew-symmetric matrix  $\{\varpi_i^j\}_{i,j=1}^2$  is fully determined by exactly one connection 1-form  $\varpi \in \Lambda^1(U)$  given by

$$\varpi(X) = \bar{g}(E_1, \bar{\nabla}_X E_2) = -\bar{g}(\bar{\nabla}_X E_1, E_2) \quad (2.5)$$

for all  $X \in \mathfrak{X}(M)$ . In this case, we have

$$\begin{aligned}\bar{\nabla}_X E_1 &= -\varpi(X)E_2, & \bar{\nabla}_X E_2 &= \varpi(X)E_1, \\ d^1\varpi &= \bar{K}(\bar{g})\sqrt{\det \bar{g}} dx^1 \wedge dx^2.\end{aligned}$$

**2.4. Approximate metric.** We are interested in approximating  $\bar{K}(\bar{g})$  when the metric is given only approximately. We assume that  $M$  has been subdivided into a geometrically conforming triangulation  $\mathcal{T}$ . The edges of  $\mathcal{T}$  may be curved, but do not necessarily consist of geodesics.

On each element  $T \in \mathcal{T}$ , we are given an approximation  $g|_T \in \mathfrak{S}(T)$  of  $\bar{g}|_T$ . When the approximation is sufficiently good,  $g$  will also be positive definite since  $\bar{g}$  is positive definite. Then each  $T \in \mathcal{T}$  can be considered to be a Riemannian manifold  $(T, g|_T)$  with  $g|_T$  as its metric. Since  $g|_T$  is smooth within each element  $T$  (not across  $\partial T$ ), we use the unique Levi-Civita connection  $\nabla$  generated by  $g|_T$  to compute covariant derivatives within  $T$ . (Constraints on  $g$  across element boundaries are clarified below in (2.14).) We drop the accent  $\bar{\phantom{x}}$  in any previous definition to indicate that it pertains to the manifold  $(T, g|_T)$  instead of  $(M, \bar{g})$ , e.g.,  $R$  refers to the Riemann curvature tensor computed using  $g$  and  $\nabla$  in place of  $\bar{g}$  and  $\bar{\nabla}$  in (2.1).

A point  $p \in T$  can be viewed either as a point in the manifold  $M$  or as a point in the manifold  $T$ . Irrespective of the two viewpoints, the meanings of coordinate frame  $\partial_i$ , coframe  $dx^i$ , and the tangent space  $T_p M$  at  $p$  are unchanged. In coordinates,

$$g_{ij} = g(\partial_i, \partial_j), \quad g^{ij} = g^{-1}(dx^i, dx^j) \quad (2.6)$$

may be viewed as entries of symmetric positive definite matrices. *Christoffel symbols of the first kind* ( $\Gamma_{ijk}$ ) and the *second kind* ( $\Gamma_{ij}^k$ ) are defined by

$$\Gamma_{ijk} = g(\nabla_{\partial_i} \partial_j, \partial_k), \quad \nabla_{\partial_i} \partial_j = \Gamma_{ij}^k \partial_k. \quad (2.7a)$$

They can alternately be expressed, using (2.6), as

$$\Gamma_{ijl} = \frac{1}{2}(\partial_i g_{jl} + \partial_j g_{li} - \partial_l g_{ij}), \quad \Gamma_{ij}^k = g^{kl} \Gamma_{ijl}. \quad (2.7b)$$

**2.5. Tangents and normals on element boundaries.** Throughout this paper, we use  $\tau$  to denote a tangent vector (not  $g$ -normalized; cf. (3.14) later) along an element boundary  $\partial T$  for any  $T \in \mathcal{T}$ . The orientation of  $\tau$  is aligned with the boundary orientation of  $\partial T$  (inherited from the orientation of  $T$ , which is the same as the orientation of  $M$ ). For any  $p \in \partial T$ , define  $\tilde{\nu} \in T_p M$  by

$$g(\tilde{\nu}, X) = (dx^1 \wedge dx^2)(\tau, X), \quad \text{for all } X \in T_p M. \quad (2.8)$$

It is easy to see from (2.8) that the ordered basis  $(\tau, \tilde{\nu})$  has the same orientation as  $(\partial_1, \partial_2)$  since  $(dx^1 \wedge dx^2)(\tau, \tilde{\nu}) > 0$ , and moreover,

$$g(\tilde{\nu}, \tau) = 0, \quad \text{and} \quad g(\tilde{\nu}, \tilde{\nu}) \det(g) = g(\tau, \tau). \quad (2.9)$$

In particular, defining

$$\hat{\nu} = \frac{\tilde{\nu}}{\sqrt{g_{\tilde{\nu}\tilde{\nu}}}}, \quad \hat{\tau} = \frac{\tau}{\sqrt{g_{\tau\tau}}}, \quad (2.10)$$

we obtain a  $g$ -orthonormal basis  $(\hat{\tau}, \hat{\nu})$  of normal and tangent vectors along every element boundary  $\partial T$ , whose orientation matches the manifold's orientation. E.g., if  $M$  is the unit disc in  $\mathbb{R}^2$  with the Euclidean metric  $g = \delta$  and the standard orientation, then  $\hat{\tau}$  is oriented counterclockwise and  $\hat{\nu}$  points inward.

In (2.10) and throughout, we use  $\sigma_{uv}$  to denote  $\sigma(u, v)$  for any vectors  $u, v \in T_p M$  and  $\sigma \in \mathcal{S}(M)$ . Note that  $g_{uv}$  is not to be confused with the  $g_{ij}$  introduced in (2.6) where the indices are integers (which trigger the summation convention) rather than vectors.

To write  $\tilde{\nu}$  in coordinates, it is useful to introduce the alternating symbol  $\varepsilon^{ij}$  whose value is 1,  $-1$ , or 0 according to whether  $(i, j)$  is an even permutation, odd permutation, or not a permutation of  $(1, 2)$ , respectively. The value of the symbols  $\varepsilon_{ij}$ ,  $\varepsilon_i^j$ , and  $\varepsilon^i_j$  equal  $\varepsilon^{ij}$ . It is easy to see that (2.8) implies

$$\tilde{\nu}^k = -g^{kj} \varepsilon_{ji} \tau^i \quad (2.11)$$

and furthermore (e.g., using (2.11), (2.9), and the Hodge star in coordinates,  $\star dx^i = \sqrt{\det g} g^{ij} \varepsilon_{jk} dx^k$ ) that

$$(\star\omega)(\hat{\nu}) = \omega(\hat{\tau}), \quad (\star\omega)(\hat{\tau}) = -\omega(\hat{\nu}), \quad (2.12)$$

for any  $\omega \in \Lambda^1(M)$ .

**2.6. Finite element spaces.** Let  $C^\infty(\mathcal{T})$  denote the space of *piecewise smooth* functions on  $\mathcal{T}$ , by which we mean functions that are infinitely smooth within each mesh element and continuous up to (including) the boundary of each mesh element  $T$ . A notation in §2.1 with  $\mathcal{T}$  in place of  $M$  indicates the piecewise smooth analogue, e.g.,  $\mathfrak{X}(\mathcal{T}) = \{X = X^i \partial_i : X^i \in C^\infty(\mathcal{T})\}$ ,  $\mathcal{S}(\mathcal{T}) = \{\sigma_{ij} dx^i \otimes dx^j : \sigma_{ij} = \sigma_{ji} \in C^\infty(\mathcal{T})\}$ , etc. Note that a  $\sigma \in \mathcal{S}(\mathcal{T})$  need not be continuous across the element interfaces. Let  $E = \partial T_+ \cap \partial T_-$  denote an interior mesh edge (possibly curved) shared between elements  $T_\pm \in \mathcal{T}$ . Let  $T_p E$  denote the one-dimensional tangent space of the curve  $E$  at any one of its points  $p$ . (Note that the tangent space at  $p$  from either element  $T_\pm$  coincides with  $T_p M$  and  $T_p E \subset T_p M$ .) We say that a  $\sigma \in \mathcal{S}(\mathcal{T})$  has “tangential-tangential continuity” or that  $\sigma$  is *tt-continuous* if

$$\sigma|_{T_+}(X, X) = \sigma|_{T_-}(X, X), \quad \text{for all } X \in T_p E, \quad (2.13)$$

at all  $p \in E$ , and for every interior mesh edge  $E$ . Let

$$\mathcal{R}(\mathcal{T}) = \{\sigma \in \mathcal{S}(\mathcal{T}) : \sigma \text{ is } tt\text{-continuous}\}, \quad (2.14a)$$

$$\mathcal{R}^+(\mathcal{T}) = \{\sigma \in \mathcal{R}(\mathcal{T}) : \sigma(X, X) > 0 \text{ for all } X \in T_p M\}. \quad (2.14b)$$

The approximate metric  $g$  is assumed to be in  $\mathcal{R}^+(\mathcal{T})$ . For the numerical analysis later, we will additionally assume that it is in  $\mathcal{R}_h^k$  defined below.

In finite element computations, we use a reference element  $\hat{T}$ , the unit triangle/simplex, and the space  $\mathcal{P}^k(\hat{T})$  of polynomials of degree at most  $k$  on  $\hat{T}$ . Let  $\tilde{T}$  denote a Euclidean triangle with possibly curved edges that is diffeomorphic to  $\hat{T}$  via  $\hat{\Phi} : \hat{T} \rightarrow \tilde{T}$ . For finite element computations on manifolds, we need charts so that each whole element  $T \in \mathcal{T}$  of the manifold is covered by a single chart giving the coordinates  $x^i$  on  $T$ . The chart identifies the parameter domain of  $T$  as the (possibly curved) Euclidean

triangle  $\tilde{T}$  diffeomorphic to  $T$ . Let  $\Phi : T \rightarrow \tilde{T}$  denote the diffeomorphism. Then  $\Phi_T = \hat{\Phi}^{-1} \circ \Phi : T \rightarrow \hat{T}$  maps diffeomorphically to the reference element where  $\mathcal{P}^k(\hat{T})$  is defined. We use its pullback  $\Phi_T^*$  below.

Define the *Regge finite element space* of degree  $k$  on the manifold  $M$  by

$$\begin{aligned} \mathcal{R}_h^k &= \{\sigma \in \mathcal{R}(\mathcal{T}) : \text{for all } T \in \mathcal{T}, \sigma|_T = \sigma_{ij} dx^i \otimes dx^j \\ &\quad \text{with } \sigma_{ij} = \Phi_T^* \hat{q} \text{ for some } \hat{q} \in \mathcal{P}^k(\hat{T})\}. \end{aligned} \quad (2.15)$$

The subscript  $h$  indicates some mesh size parameter, e.g., on meshes whose elements are close to straight-edged triangles, one may set  $h = \max_{T \in \mathcal{T}} \text{diam}(T)$ . Let

$$\begin{aligned} \mathcal{V}(\mathcal{T}) &= \{u \in \Lambda^0(\mathcal{T}) : u \text{ is continuous on } M\}, \\ \mathring{\mathcal{V}}_T(\mathcal{T}) &= \{u \in \mathcal{V}(\mathcal{T}) : u|_T = 0\}, \end{aligned} \quad (2.16)$$

where  $\Gamma$  denote a subset of the boundary  $\partial M$  of positive boundary measure.

The *Lagrange finite element space* on  $M$  and its subspaces with essential boundary conditions are defined by

$$\begin{aligned} \mathcal{V}_h^k &= \{u \in \mathcal{V}(\mathcal{T}) : \text{for all } T \in \mathcal{T}, u|_T = \Phi_T^* \hat{u} \text{ for some } \hat{u} \in \mathcal{P}^k(\hat{T})\}, \\ \mathring{\mathcal{V}}_{h,\Gamma}^k &= \{u \in \mathcal{V}_h^k : u|_\Gamma = 0\} \quad \text{and} \quad \mathring{\mathcal{V}}_h^k = \mathring{\mathcal{V}}_{h,\partial M}^k. \end{aligned} \quad (2.17)$$

The previous definitions in this subsection were independent of the metric. We will now introduce a metric-dependent space of normal-continuous vector fields. First, we introduce the following notation surrounding an interior mesh edge  $E$  shared by two adjacent elements in  $\mathcal{T}$ ,

$$E = \partial T_- \cap \partial T_+, \quad T_\pm \in \mathcal{T}. \quad (2.18a)$$

In this context, the  $g$ -orthonormal tangent and normal vectors introduced above along  $\partial T_\pm$  are denoted by  $\hat{\tau}_\pm$  and  $\hat{\nu}_\pm$ , respectively. For a collection of scalar functions,  $\{f_{\partial T}(\hat{\nu}) : T \in \mathcal{T}\}$ , each depending on the normal  $\hat{\nu}$  at an element boundary, we define the jump on  $E$  by

$$\llbracket f(\hat{\nu}) \rrbracket = f_{\partial T_+}(\hat{\nu}_+) + f_{\partial T_-}(\hat{\nu}_-). \quad (2.18b)$$

Thus the jump function  $\llbracket f(\hat{\nu}) \rrbracket$  is well defined (and single-valued) on the union of all interior mesh edges, excluding the mesh vertices. The jump of an element-boundary function dependent on  $\hat{\tau}$  is defined similarly.

We say that a piecewise smooth vector field  $W \in \mathfrak{X}(\mathcal{T})$  has “ $g$ -normal continuity” across element interfaces if  $\llbracket g(W, \hat{\nu}) \rrbracket = 0$ . Define

$$\begin{aligned} \mathcal{W}_g(\mathcal{T}) &= \{W \in \mathfrak{X}(\mathcal{T}) : \llbracket g(W, \hat{\nu}) \rrbracket = 0\}, \\ \mathring{\mathcal{W}}_{g,\Gamma}(\mathcal{T}) &= \{W \in \mathcal{W}_g(\mathcal{T}) : g(W|_\Gamma, \hat{\nu}) = 0\}, \quad \mathring{\mathcal{W}}_g(\mathcal{T}) = \mathring{\mathcal{W}}_{g,\partial M}(\mathcal{T}). \end{aligned} \quad (2.19)$$

**2.7. Integrals over the manifold triangulation.** On every element  $T \in \mathcal{T}$ , in order to integrate a scalar function  $f \in \Lambda^0(T)$ , adopting the notation of [32], we tacitly use the unique Riemannian volume form  $\text{vol}_{(T,g)}$  to convert it to a 2-form and then pullback to integrate over the Euclidean parameter domain  $\tilde{T}$ , i.e.,

$$\int_{(T,g)} f \equiv \int_T f \text{vol}_{(T,g)} = \int_{\tilde{T}} (\Phi^{-1})^*(f \text{vol}_{(T,g)}) = \int_{\tilde{T}} f \circ \Phi^{-1} \frac{\sqrt{\det g}}{\det(D\Phi)} \text{da}, \quad (2.20)$$

where  $\det(D\Phi)$  denotes the Jacobian determinant of  $\Phi$ , we have used the standard extension of pullback to forms, and we have appended an area measure notation “da” to emphasize that the right most integral is a standard Lebesgue integral over the Euclidean domain  $\tilde{T}$ . For  $v, w \in \Lambda^0(\mathcal{T})$ , set

$$\int_{\mathcal{T}} w = \sum_{T \in \mathcal{T}} \int_{(T,g)} w, \quad (v, w)_{\mathcal{T}} = \sum_{T \in \mathcal{T}} \int_{(T,g)} v w, \quad (2.21a)$$

with the understanding that the right hand sides above must be evaluated using (2.20). In order to integrate along the boundary curve  $\partial T$ , we use the one-dimensional analogue of the formula in (2.20) to compute on the Euclidean domain  $\partial\tilde{T} = \Phi(\partial T)$ , namely

$$\int_{(\partial T, g)} f = \int_{\partial\tilde{T}} (\Phi^{-1})^*(f \operatorname{vol}_{(\partial T, g)}) = \int_{\partial\tilde{T}} f \circ \Phi^{-1} \sqrt{g_{tt}} \, dl, \quad (2.21b)$$

where  $t$  is a tangent vector along  $\partial\tilde{T}$  of unit Euclidean length—and to emphasize that the last integral is a standard Euclidean integral, we have appended the length measure “dl”. We use

$$\int_{\partial\mathcal{T}} w = \sum_{T \in \mathcal{T}} \int_{(\partial T, g)} w \quad (2.21c)$$

to simplify notation for sum of integrals over element boundaries.

### 3. CURVATURE APPROXIMATION

In this section we give the curvature approximation formula and discuss a few non-trivial computational details on curved elements. In order to approximate the Gauss curvature  $K(\bar{g})$ , one may consider computing  $K(g|_T)$  on each element  $T \in \mathcal{T}$  using the given approximation  $g$  of the exact metric  $\bar{g}$ . However, this alone cannot generally be a good approximation to  $K(\bar{g})$  because discontinuities of  $g$  across elements generate additional sources of curvature on the edges and vertices of the mesh. Below we provide a curvature approximation incorporating these extra sources. Since it coincides with the formula given in a recent work [11] for a specific case, we opt for a brief description, expanding only on aspects complementary to that work.

**3.1. Curvature approximation formula.** Given a metric  $g \in \mathcal{R}^+(\mathcal{T})$  approximating  $\bar{g}$  we identify three sources of curvature, modeled after similar terms in the Gauss-Bonnet formula, and define them as the following linear functionals acting on  $\varphi \in \mathcal{V}(\mathcal{T})$ :

$$\begin{aligned} \langle K_g^T, \varphi \rangle_{\mathcal{V}(\mathcal{T})} &= \int_{(T,g)} K(g) \varphi_T, & \langle K_{E,g}^T, \varphi \rangle_{\mathcal{V}(\mathcal{T})} &= \int_{(E,g)} \kappa(g) \varphi_T, \\ \langle K_{V,g}, \varphi \rangle_{\mathcal{V}(\mathcal{T})} &= \left(2\pi - \sum_{T \in \mathcal{T}_V} \angle_V^T(g)\right) \varphi(V), \end{aligned} \quad (3.1)$$

where  $\mathcal{T}_V$  denotes the set of all elements of  $\mathcal{T}$  which have  $V$  as a vertex. Here  $K(g)$  and  $\kappa(g)$  are defined by (2.2) and (2.3) after replacing  $\bar{g}$  by  $g|_T$ , and  $\varphi_T = \varphi|_T$  for any  $T \in \mathcal{T}$ . Throughout, we use  $\langle f, \varphi \rangle_H$  to denote duality pairing on a vector space  $H$  that gives the action of a linear functional  $f$  on a  $\varphi \in H$ . Moreover, we let  $\mathcal{V}$  and  $\mathcal{E}$



denote the set of mesh vertices and edges, respectively (so in (3.1),  $V \in \mathcal{V}$  and  $E \in \mathcal{E}$ ). Finally, we let  $\angle_V^T(\cdot)$  denote the interior angle at a vertex  $V$  of  $T$  determined using the metric in its argument (computable using (3.4) below).

In addition to  $g$ , suppose that we are also given boundary curvature data in essential (Dirichlet) or natural (Neumann) forms for manifolds with boundary. The former type of data arises when we know that  $M$  is a submanifold of a larger manifold whose Gauss curvature is known outside of  $M$ . To accommodate such information only on a part of the boundary of  $M$ , we split  $\partial M$  into two non-overlapping parts  $\Gamma_D$  and  $\Gamma_N$ . One of these can be empty, and in case one is nonempty, both must have positive length. On  $\Gamma_D$ , we assume that we are given  $K^D(\bar{g}) = K(\bar{g})|_{\Gamma_D}$  and that  $K^D(\bar{g})$  is in the trace of the Lagrange finite element space  $\mathcal{V}_h^k$ . E.g., when a manifold is flat around  $\Gamma_D$  (i.e.,  $K(\bar{g})$  vanishes in a neighborhood of  $\Gamma_D$ ), we may set homogeneous Dirichlet data  $K^D(\bar{g}) = 0$  on  $\Gamma_D$ .

The other type of boundary data, in the form of a natural (or Neumann) boundary condition, is motivated by the Gauss–Bonnet theorem, and provides geodesic curvature data at the boundary. These natural Neumann-type boundary data is given in the form of the functional

$$\langle \kappa^N(\bar{g}), \varphi \rangle_{\mathcal{V}(\mathcal{T})} = \int_{(\Gamma_N, \bar{g})} \kappa(\bar{g}) \varphi - \sum_{V \in \mathcal{V}^N} \tilde{\angle}_V^N(\bar{g}) \varphi(V), \quad (3.2)$$

for  $\varphi \in \mathcal{V}(\mathcal{T})$ , where  $\mathcal{V}^N \subset \mathcal{V} \cap \Gamma_N$  is the subset of the manifold's vertices contained in the interior of  $\Gamma_N$  and  $\tilde{\angle}_V^N$  denotes the exterior angle measured by the edges of  $\Gamma_N$  at  $V$ . If  $V$  is part of a smooth boundary  $\angle_V^N(\delta) = \pi$ , whereas at kinks of the boundary the angle has to be provided as input data. The action of functional (3.2) on the finite-dimensional subspace  $\mathcal{V}_h^k \subset \mathcal{V}(\mathcal{T})$  is computable if we are given the exact metric  $\bar{g}$  on and near  $\Gamma_N$ . In practice, if  $\bar{g}$  is unknown near the boundary, we require that approximations to  $\kappa^N(\bar{g})$  and  $K^D(\bar{g})$  be provided. For manifolds without boundary, there is no need to provide any boundary data.

**Definition 3.1** (Curvature approximation). The curvature approximation  $K_h(g)$  is the unique function in  $\mathcal{V}_h^k$  determined by requiring that  $K_h(g)|_{\Gamma_D} = K^D(\bar{g})$  on  $\Gamma_D$  and for all  $u_h \in \mathring{\mathcal{V}}_{h, \Gamma_D}^k$ ,

$$\int_{\mathcal{T}} K_h(g) u_h = \sum_{T \in \mathcal{T}} \left( \langle K_g^T, u_h \rangle_{\mathcal{V}(\mathcal{T})} + \sum_{E \in \mathcal{E}_T} \langle K_{E,g}^T, u_h \rangle_{\mathcal{V}(\mathcal{T})} + \sum_{V \in \mathcal{V}_T} \langle K_{V,g}^T, u_h \rangle_{\mathcal{V}(\mathcal{T})} \right) - \langle \kappa^N(\bar{g}), u_h \rangle_{\mathcal{V}(\mathcal{T})}. \quad (3.3)$$

Here  $\mathcal{V}_T$  denotes the set of three vertices of an element  $T \in \mathcal{T}$  and  $\mathcal{E}_T$  denotes the set of three edges of  $\partial T$ .

In the remainder of this subsection, we discuss how to compute the quantities in (3.3). Recall (from §2.5) that the tangent vector  $\tau$  along the boundary  $\partial T$  is aligned with the boundary orientation of  $\partial T$ . At any vertex  $V \in \mathcal{V}_T$ , the tangent  $\tau$  undergoes a change in direction, between an incoming and an outgoing tangent vector, which we denote by  $\tau_{\text{in}}$  and  $\tau_{\text{out}}$ , respectively. The angle at  $V$  with respect to the metric  $g$  is

then computed by

$$\angle_V^T(g) = \arccos \left( \frac{g(-\tau_{\text{in}}, \tau_{\text{out}})}{\|\tau_{\text{in}}\|_g \|\tau_{\text{out}}\|_g} \right), \quad (3.4)$$

where  $\|w\|_g = g(w, w)^{1/2}$ . This and the same formula with  $g$  replaced by  $\delta$  is what we use to calculate the angle deficit functional  $K_{V,g}^T$  (3.1).

Next, consider the interior source term  $K_g^T$ , defined using  $K(g)$ , and related to the Riemann curvature by (2.2). By (2.1),  $R_{ijkl} = R(\partial_i, \partial_j, \partial_k, \partial_l)$  simplifies to

$$\begin{aligned} R_{ijkl} &= (\partial_i \Gamma_{jk}^p + \Gamma_{jk}^q \Gamma_{iq}^p - \partial_j \Gamma_{ik}^p - \Gamma_{ik}^q \Gamma_{jq}^p) g_{pl} \\ &= \partial_i \Gamma_{jkl} - \partial_j \Gamma_{ikl} - \Gamma_{pil} \Gamma_{jk}^p + \Gamma_{pjl} \Gamma_{ik}^p, \end{aligned} \quad (3.5)$$

where  $\Gamma_{ij}^k$  and  $\Gamma_{ijk}$  are as in (2.7). For two-dimensional manifolds the Gauss curvature can be expressed by  $K(g) = R_{1221}/\det g$ . Hence by (2.20),

$$\langle K_g^T, \varphi \rangle_{\mathcal{V}(\mathcal{T})} = \int_T K(g) \varphi \sqrt{\det g} \, dx^1 \wedge dx^2 = \int_{\bar{T}} \frac{\Phi_*(R_{1221} \varphi)}{\sqrt{\det g} \det(D\Phi)} \, da. \quad (3.6)$$

It thus remains to discuss the computation of the edge sources  $K_{E,g}^T$  using the definition of the  $\kappa(g)$  in (2.3). In finite element computations, we usually do not have ready access to the  $g$ -arclength parameter  $s$  used there. But  $\kappa(g)$  can be computed using the readily accessible  $\tau$  and  $\tilde{\nu}$  of §2.5, as shown below. Let  $\gamma(t)$  be an orientation-preserving parametrization that gives an oriented mesh edge  $E \subset \partial T$  as  $E = \{\gamma(t) : t_0 \leq t \leq t_1\}$ . Parametrizing scalar functions  $a$  on  $E$  by  $t$ , we abbreviate  $da/dt$  to  $\dot{a}$ . Note that the components of  $\tau = \tau^k \partial_k$  are given by  $\tau^k = \dot{\gamma}^k$  and their derivatives by  $d^2\gamma^k/dt^2 = \dot{\tau}^k$ . Let

$$\dot{\tau} = \dot{\tau}^k \partial_k, \quad G_{uv}^w = g(u^i v^j \Gamma_{ij}^k \partial_k, w), \quad (3.7)$$

where  $u = u^i \partial_i, v = v^i \partial_i, w \in \mathfrak{X}(T)$ .

**Lemma 3.2.** *The geodesic curvature along each edge of an element boundary  $\partial T$  is given by*

$$\kappa(g) = \frac{\sqrt{\det g}}{g_{\tau\tau}^{3/2}} (g_{\hat{\tau}\tilde{\nu}} + G_{\tau\tau}^{\tilde{\nu}}). \quad (3.8)$$

*Proof.* By (2.3) and our definitions of  $\hat{\tau}, \hat{\nu}$  in §2.5,

$$\kappa(g) = g(\nabla_{\hat{\tau}(s)} \hat{\tau}(s), \hat{\nu}(s)) \quad (3.9)$$

where  $s$  is the  $g$ -arclength parameter. Inverting  $s(t) = \int_0^t g(\tau(t), \tau(t))^{1/2} dt$  to write  $t$  as a function of  $s$ , applying the chain rule to  $\mu(s) = \gamma(t(s))$ , and using  $dt/ds = g_{\tau\tau}^{-1/2}$ ,

$$\frac{d\hat{\tau}^k}{ds} = \frac{d^2\mu^k}{ds^2} = \frac{d^2\gamma^k}{dt^2} \left( \frac{dt}{ds} \right)^2 + \frac{d\gamma^k}{dt} \frac{d^2t}{ds^2} = \frac{\dot{\tau}^k}{g_{\tau\tau}} + \tau^k \frac{d^2t}{ds^2}.$$

Using the properties of the connection  $\nabla$  (see e.g. [31]),

$$\nabla_{\hat{\tau}} \hat{\tau} = \frac{d\hat{\tau}^k}{ds} \partial_k + \hat{\tau}^i \hat{\tau}^j \Gamma_{ij}^k \partial_k = \frac{\dot{\tau}^k}{g_{\tau\tau}} + \tau \frac{d^2t}{ds^2} + \hat{\tau}^i \hat{\tau}^j \Gamma_{ij}^k \partial_k. \quad (3.10)$$

Hence (3.9), (3.10), and the  $g$ -orthogonality of  $\tau$  with  $\hat{\nu}$ , implies that at the point  $\gamma(t)$ ,

$$\kappa(g) = \frac{g(\dot{\tau}, \hat{\nu})}{g_{\tau\tau}} + \frac{g(\tau^i \tau^j \Gamma_{ij}^k \partial_k, \hat{\nu})}{g_{\tau\tau}}. \quad (3.11)$$

Now, by (2.9) and (2.10),  $\hat{\nu} = \tilde{\nu}(\det g/g_{\tau\tau})^{1/2}$ , so (3.11) implies

$$\kappa(g) = \frac{(\det g)^{1/2}}{g_{\tau\tau}^{3/2}} [g(\dot{\tau}, \tilde{\nu}) + g(\tau^i \tau^j \Gamma_{ij}^k \partial_k, \tilde{\nu})] \quad (3.12)$$

which proves (3.8).  $\square$

Returning to the edge source term, using Lemma 3.2 and (2.21b),

$$K_{E,g}^T(\varphi, g) = \int_E \kappa(g) \varphi \sqrt{g_{\tau\tau}} = \int_{\Phi(E)} \Phi_* \left( \frac{\sqrt{\det g}}{g_{\tau\tau}} (g_{\tilde{\nu}\nu} + g_{\tau\tau}^\nu) \varphi \right) dl. \quad (3.13)$$

Thus, through (3.4), (3.6), and (3.13), we have shown it is possible to easily compute all the terms in the curvature approximation (3.3) using standard finite element tools.

**3.2. A model case for further analysis.** Having shown how curvature of general manifolds can be computed in the remainder of the paper, we focus our analysis on the model case described now. We assume that the manifold  $M$ , as a set, equals  $\Omega \subset \mathbb{R}^2$ , a bounded open connected domain. The set  $\Omega$  forms the full parameter domain of  $M$  for the single trivial chart  $\Phi : M \rightarrow \Omega$  with  $\Phi$  equaling the identity map. The triangulation  $\mathcal{T}$  of  $M$  is now a conforming finite element mesh in the planar domain  $\Omega$  and its elements are (possibly curved) triangles.

In this setting, an element  $T \in \mathcal{T}$  can be considered as either the Euclidean manifold  $(T, \delta)$ , equipped with the identity metric  $\delta$ , or the Riemannian manifold  $(T, g)$ . Let us reconsider the tangent vector  $\tau$  on an element boundary  $\partial T$  with the orientation described in §2.5, previously left un-normalized. Henceforth, we assume that

$$1 = \delta(\tau, \tau). \quad (3.14)$$

We computed the normal vector  $\tilde{\nu}$  from  $\tau$  by (2.8) for the manifold  $(T, g)$ . For the manifold  $(T, \delta)$ , a generally different Euclidean normal vector arises at any  $p$  in  $\partial T$  and we denote it by  $\nu$ . It can be computed by simply replacing  $g$  with  $\delta$  in (2.8), i.e.,  $\nu \in T_p M$  satisfies

$$\delta(\nu, X) = dx^1 \wedge dx^2(\tau, X) \quad \text{for all } X \in T_p M. \quad (3.15)$$

Analogous to (2.11), we now have the accompanying coordinate expression,

$$\nu^k = -\delta^{kj} \varepsilon_{ji} \tau^i = -\varepsilon^k{}_i \tau^i. \quad (3.16)$$

Note that (3.14) implies that  $\delta(\nu, \nu) = 1$ . These identities, together with (2.10) guide us move between the  $\delta$ -orthonormal tangent and normal vectors ( $\tau$  and  $\nu$ ) and  $g$ -orthonormal tangent and normal vectors ( $\hat{\tau}$  and  $\hat{\nu}$ ), while preserving the orientation. Jumps of functions of  $\nu$  and  $\tau$  on the Euclidean element boundaries  $(\partial T, \delta)$ ,  $T \in \mathcal{T}$ , are defined in analogy to (2.18b).

We proceed to display the curvature approximation formula in coordinates for this model case using the above-mentioned Euclidean normal and tangent ( $\nu$  and  $\tau$ ).

**Lemma 3.3.** *The geodesic curvature along any mesh edge  $E$  is given by*

$$\kappa(g) = \frac{\sqrt{\det g}}{g_{\tau\tau}^{3/2}} (\dot{\tau}_\nu + \Gamma_{\tau\tau}^\nu)$$

where  $\Gamma_{\tau\tau}^\nu = \tau^i \tau^j \Gamma_{ij}^k \delta_{kl} \nu^l$  and  $\dot{\tau}_\nu = \delta(\dot{\tau}, \nu)$ .

*Proof.* Comparing (3.15) with (2.8),

$$g(\tilde{\nu}, X) = \delta(\nu, X) \quad (3.17)$$

for any  $X \in T_p M$ . Using (3.17) in Lemma 3.2, the result follows.  $\square$

In this model case of a single parameter domain, since the identity metric  $\delta$  is well defined throughout, the angle deficit in (3.1) can be reformulated using  $\delta$  as a sum of element contributions at each vertex. Define

$$\langle K_{V,g}^T, u_h \rangle_{\mathcal{V}(\mathcal{T})} = (\triangleleft_V^T(\delta) - \triangleleft_V^T(g)) u_h(V).$$

When summing  $\triangleleft_V^T(\delta)$  over all  $V \in \mathcal{V}_T$ , we clearly obtain  $2\pi$ . Hence

$$\sum_{V \in \mathcal{V}} \langle K_{V,g}, \varphi \rangle_{\mathcal{V}(\mathcal{T})} = \sum_{T \in \mathcal{T}} \sum_{V \in \mathcal{V}_T} \langle K_{V,g}^T, u_h \rangle_{\mathcal{V}(\mathcal{T})}. \quad (3.18)$$

**Proposition 3.4.** *In this model case, equation (3.3) implies that for all  $u_h \in \mathring{\mathcal{V}}_{h,\Gamma_D}^k$ ,*

$$\begin{aligned} \int_{\mathcal{T}} K_h(g) u_h &= -\langle \kappa^N(\bar{g}), u_h \rangle_{\mathcal{V}(\mathcal{T})} + \sum_{T \in \mathcal{T}} \sum_{V \in \mathcal{V}_T} \langle K_{V,g}^T, u_h \rangle_{\mathcal{V}(\mathcal{T})} \\ &+ \sum_{T \in \mathcal{T}} \left( \int_T \frac{R_{1221} u_h}{\sqrt{\det g}} \, da + \int_{\partial T} \frac{\sqrt{\det g}}{g_{\tau\tau}} (\dot{\tau}_\nu + \Gamma_{\tau\tau}^\nu) u_h \, dl \right). \end{aligned} \quad (3.19)$$

*Proof.* This follows from (3.18), (3.6), (3.13), and Lemma 3.3, once  $\Phi$  is set to the identity.  $\square$

The next result shows that if  $g$  is globally smooth enough (twice continuous differentiability is sufficient) for  $K(g)$  to be computed globally (using derivatives of  $g$  and (2.2)), then  $K_h(g)$  is the closest approximation to  $K(g)$  from the finite element space in the sense of (3.20) below. For its proof and later, it is convenient to divide the set of mesh vertices  $\mathcal{V}$  into the set of boundary and interior vertices

$$\mathcal{V}^{\text{bnd}} = \mathcal{V} \cap \partial M, \quad \mathcal{V}^{\text{int}} = \mathcal{V} \setminus \mathcal{V}^{\text{bnd}},$$

and further subdivide  $\mathcal{V}^{\text{bnd}}$  into  $\mathcal{V}_D^{\text{bnd}} = \mathcal{V}^{\text{bnd}} \cap \bar{\Gamma}_D$  and  $\mathcal{V}_N^{\text{bnd}} = \mathcal{V}^{\text{bnd}} \setminus \mathcal{V}_D^{\text{bnd}}$ . Similarly, we partition the set of mesh edges  $\mathcal{E}$  into the set of boundary edges  $\mathcal{E}^{\text{bnd}} = \{E \in \mathcal{E} : E \subset \partial M\}$  and the set of interior edges  $\mathcal{E}^{\text{int}} = \mathcal{E} \setminus \mathcal{E}^{\text{bnd}}$ .

**Lemma 3.5** (Consistency). *Suppose  $g \in \mathcal{S}^+(M)$  and  $\kappa^N(\bar{g}) = \kappa^N(g)$ . Then for any  $u_h \in \mathring{\mathcal{V}}_{h,\Gamma_D}^k$ ,*

$$\int_{\mathcal{T}} K_h(g) u_h = \int_{\mathcal{T}} K(g) u_h. \quad (3.20)$$

*Proof.* At every interior vertex  $V \in \mathcal{V}^{\text{int}}$ , a rearrangement gives

$$\sum_{T \in \mathcal{T}} \sum_{V \in \mathcal{V}_T^{\text{int}}} \langle K_{V,g}^T, u_h \rangle_{\mathcal{V}(\mathcal{T})} = \sum_{V \in \mathcal{V}^{\text{int}}} \sum_{T \in \mathcal{T}_V} (\angle_V^T(\delta) - \angle_V^T(g)) u_h(V).$$

It vanishes because the smoothness of  $g$  implies  $\sum_{T \in \mathcal{T}_V} \angle_V^T(g) = 2\pi = \sum_{T \in \mathcal{T}_V} \angle_V^T(\delta)$ . Similarly, at a boundary vertex  $V \in \mathcal{V}_N^{\text{bnd}}$ , the sum of the interior angles  $\sum_{T \in \mathcal{T}_V} \angle_V^T(g)$  equals  $\angle_V^N(g)$  due to the smoothness of  $g$ . Hence, applying Proposition 3.4 and using  $\kappa^N(\bar{g}) = \kappa^N(g)$ , the integrals over  $\Gamma_N$  and the sum of all the angle deficit functionals on boundary vertices vanish, so we obtain

$$\begin{aligned} \int_{\mathcal{T}} (K_h(g) - K(g)) u_h &= \sum_{T \in \mathcal{T}} \int_{\partial T \setminus \Gamma_N} \frac{\sqrt{\det g}}{g_{\tau\tau}} (\dot{\tau}_\nu + \Gamma_{\tau\tau}^\nu) u_h \, dl \\ &= \sum_{E \in \mathcal{E}^{\text{int}}} \int_E \frac{\sqrt{\det g}}{g_{\tau\tau}} \llbracket \dot{\tau}_\nu + \Gamma_{\tau\tau}^\nu \rrbracket u_h \, dl. \end{aligned} \quad (3.21)$$

In the last step, we have used that  $u_h = 0$  on  $\Gamma_D$  and that the trace of  $\sqrt{\det g}/g_{\tau\tau}$  is well defined (single valued) on  $E$  due to the given smoothness of  $g$ . It is easy to see that  $\dot{\tau}$  has the same value from adjacent elements of  $E$  while  $\tau$  and  $\nu$  changes sign, so  $\llbracket \dot{\tau}_\nu \rrbracket = 0$  and  $\llbracket \Gamma_{\tau\tau}^\nu \rrbracket = 0$ . Hence (3.21) vanishes.  $\square$

#### 4. COVARIANT DERIVATIVES USING THE NONSMOOTH METRIC

The objective of this section is to formulate a covariant incompatibility operator that can be applied to our situation with piecewise smooth metrics. To this end, we first define several covariant derivatives in the smooth case, restricting ourselves to a single element, i.e., the smooth Riemannian manifold  $(T, g|_T)$ . Then we proceed to consider the changes needed due to the jumps of the metric across element interfaces.

**4.1. Covariant curl and incompatibility for the smooth metric.** For an  $\mu \in \Lambda^1(T)$ , the exterior derivative  $d^1\mu \in \Lambda^2(T)$  is characterized in terms of the connection [37] by

$$(d^1\mu)(X, Y) = (\nabla_X\mu)(Y) - (\nabla_Y\mu)(X), \quad (4.1)$$

for any  $X, Y \in \mathfrak{X}(T)$ . Next, for a  $\sigma \in \mathcal{T}_0^2(T)$  writing  $\sigma_Z(Y) = \sigma(Z, Y)$  for any  $X, Y, Z \in \mathfrak{X}(T)$ , we define an operation analogous to (4.1) on  $\sigma$  by

$$(d^1\sigma_Z)(X, Y) = (\nabla_X\sigma)(Z, Y) - (\nabla_Y\sigma)(Z, X). \quad (4.2)$$

Since the expressions in (4.1) and (4.2) are skew-symmetric in  $X, Y$ , they may be viewed as elements of  $\Lambda^2(T)$ . Then we may use the Hodge star  $\star$  operation to convert them to 0-forms, since  $T$  is two dimensional. Doing so, we define the covariant curls

$$\text{curl}_g \mu = \star(d^1\mu), \quad \mu \in \mathcal{T}_0^1(T) = \Lambda^1(T), \quad (4.3a)$$

$$(\text{curl}_g \sigma)(Z) = \star(d^1\sigma_Z), \quad \sigma \in \mathcal{T}_0^2(T), \quad Z \in \mathfrak{X}(T). \quad (4.3b)$$

Combining these operations in succession, we define the *covariant incompatibility*,

$$\text{inc}_g \sigma = \text{curl}_g \text{curl}_g \sigma, \quad \sigma \in \mathcal{T}_0^2(T). \quad (4.3c)$$

Note that while  $\text{curl}_g \mu$  and  $\text{inc}_g \sigma$ , as defined above, are in  $\Lambda^0(T)$ , the covariant curl of  $\sigma$  is in  $\Lambda^1(T)$ .

We will now quickly illustrate how to get coordinate expressions from such definitions. Consider the operator in (4.2). Expanding the right hand side of (4.2) using the Leibniz rule  $(\nabla_X \sigma)(Z, Y) = X \sigma(Z, Y) - \sigma(\nabla_X Z, Y) - \sigma(Z, \nabla_X Y)$  twice, substituting  $Z = \partial_i, X = \partial_j, Y = \partial_k$ , and simplifying using (2.7a),

$$(d^1 \sigma_{\partial_i})(\partial_j, \partial_k) = (\partial_j \sigma_{ik} - \Gamma_{ji}^m \sigma_{mk}) - (\partial_k \sigma_{ij} - \Gamma_{ki}^m \sigma_{mj}) \quad (4.4)$$

for  $\sigma = \sigma_{jk} dx^j \otimes dx^k \in \mathcal{T}_0^2(T)$ . Note that equation (4.4) can be rewritten as  $d^1 \sigma_{\partial_i} = \varepsilon^{jk} (\partial_j \sigma_{ik} - \Gamma_{ji}^m \sigma_{mk}) dx^1 \wedge dx^2$ . Next, since (4.3b) implies  $\text{curl}_g \sigma = (\star d^1 \sigma_{\partial_i}) dx^i$ , recalling that  $\star(f dx^1 \wedge dx^2) = f/\sqrt{\det g}$  for any scalar field  $f \in \Lambda^0(T)$ , we arrive at

$$\text{curl}_g(\sigma_{jk} dx^j \otimes dx^k) = \frac{1}{\sqrt{\det g}} \varepsilon^{jk} (\partial_j \sigma_{ik} - \Gamma_{ji}^m \sigma_{mk}) dx^i. \quad (4.5a)$$

Similarly, one obtains coordinate expressions for the remaining operators in (4.3), namely

$$\text{curl}_g(\mu_i dx^i) = \frac{1}{\sqrt{\det g}} \varepsilon^{ij} \partial_i \mu_j, \quad (4.5b)$$

$$\begin{aligned} \text{inc}_g(\sigma_{jk} dx^j \otimes dx^k) &= \frac{1}{\det g} \left( \varepsilon^{qi} \varepsilon^{jk} \partial_{jq} \sigma_{ik} - \varepsilon^{qi} \varepsilon^{jk} \partial_q (\Gamma_{ji}^m \sigma_{mk}) \right. \\ &\quad \left. - \Gamma_{lq}^l \varepsilon^{qi} \varepsilon^{jk} (\partial_j \sigma_{ik} - \Gamma_{ji}^m \sigma_{mk}) \right). \end{aligned} \quad (4.5c)$$

In the derivation of (4.5c), we have employed the useful formula

$$\Gamma_{lq}^l = \frac{\partial_q (\det g)}{2 \det g}. \quad (4.6)$$

It is useful to contrast the expressions in (4.5) with the standard Euclidean curl and inc. To this end we use matrix and vector proxies,  $[\sigma] \in \mathbb{R}^{2 \times 2}$  and  $[\mu] \in \mathbb{R}^2$ , of  $\mu \in \Lambda^1(T)$  and  $\sigma \in \mathcal{T}_0^2(T)$ , respectively [4]. These proxies are made up of coefficients in the coordinate basis expansion, specifically  $[\sigma]$  is the matrix whose  $(i, j)$ th entry is  $\sigma_{ij} = \sigma(\partial_i, \partial_j)$ , and  $[\mu]$  is the Euclidean vector whose  $i$ th component, denoted by  $[\mu]_i$ , equals  $\mu_i = \mu(\partial_i)$ . Then the standard two-dimensional curl operator applied to the vector  $[\mu]$  gives  $\text{curl}[\mu] = \varepsilon^{ij} \partial_i \mu_j$ . When this operator is repeated row-wise on a matrix, we get the standard row-wise matrix curl, namely  $[\text{curl}[\sigma]]_i = \varepsilon^{jk} \partial_j \sigma_{ik}$ . The standard Euclidean incompatibility operator [5, 2, 17] in two dimensions is  $\text{inc}[\sigma] = \varepsilon^{qi} \varepsilon^{jk} \partial_{jq} \sigma_{ik}$ . Using these, we can rewrite the formulas in (4.5) as

$$[\text{curl}_g \mu] = \frac{1}{\sqrt{\det g}} \text{curl}[\mu], \quad (4.7a)$$

$$[\text{curl}_g \sigma]_i = \frac{1}{\sqrt{\det g}} ([\text{curl}[\sigma]]_i - \varepsilon^{jk} \Gamma_{ji}^m \sigma_{mk}), \quad (4.7b)$$

$$[\text{inc}_g \sigma] = \frac{1}{\det g} \left( \text{inc}[\sigma] - \varepsilon^{qi} \varepsilon^{jk} \partial_q (\Gamma_{ji}^m \sigma_{mk}) - \Gamma_{lq}^l \varepsilon^{qi} (\text{curl}[\sigma]_i - \varepsilon^{jk} \Gamma_{ji}^m \sigma_{mk}) \right). \quad (4.7c)$$

Note how the expressions for covariant curl and covariant inc contain, but differ from their Euclidean analogues.

Other useful covariant operators include

$$\operatorname{rot}_g f = -(\star d^0 f)^\sharp, \quad f \in \Lambda^0(T), \quad (4.8a)$$

$$(\operatorname{rot}_g X)(\mu, \eta) = \mu(\nabla_{(\star\eta)^\sharp} X), \quad \mu, \eta \in \Lambda^1(T), \quad X \in \mathfrak{X}(T). \quad (4.8b)$$

Clearly,  $\operatorname{rot}_g f$  is in  $\mathfrak{X}(T)$ , while  $\operatorname{rot}_g X \in \mathcal{T}_2^0(T)$ . Their coordinate expressions are

$$\operatorname{rot}_g f = \frac{\varepsilon^{kp} \partial_p f}{\sqrt{\det g}} \partial_k = \frac{[\operatorname{rot} f]^k}{\sqrt{\det g}} \partial_k \quad (4.9a)$$

$$\operatorname{rot}_g X = \frac{\varepsilon^{pi} (\partial_i X^m + \Gamma_{ik}^m X^k)}{\sqrt{\det g}} \partial_m \otimes \partial_p = \frac{[\operatorname{rot}[X]]^{mp} + \varepsilon^{pi} \Gamma_{ik}^m X^k}{\sqrt{\det g}} \partial_m \otimes \partial_p, \quad (4.9b)$$

where, in the latter expressions, the proxy notation has been extended to  $\mathfrak{X}(T)$  and  $\mathcal{T}_2^0(T)$  in an obvious fashion to use the Euclidean matrix and vector  $\operatorname{rot}$ .

It is easy to see that the following integration by parts formula

$$\int_{(T,g)} (\operatorname{curl}_g \sigma)(Z) = \int_{(T,g)} \sigma(\operatorname{rot}_g Z) + \int_{(\partial T,g)} \sigma(Z, \hat{\tau}) \quad (4.10)$$

holds for all  $\sigma \in \mathcal{T}_0^2(T)$  and  $Z \in \mathfrak{X}(T)$ . (This can be seen, e.g., using the coordinate expressions (4.9b) and (4.7b) and standard integration by parts on the Euclidean parameter domain.) Here  $\hat{\tau}$  is the unit tangent defined in (2.10), the integrals are computed as indicated in (2.20), and  $\sigma(\operatorname{rot}_g Z)$  denotes the result obtained by acting the  $\mathcal{T}_0^2(T)$ -tensor  $\sigma$  on the  $\mathcal{T}_2^0(T)$ -tensor  $\operatorname{rot}_g Z$ . Equation (4.10) shows that  $\operatorname{rot}_g$  can be interpreted as the adjoint of  $\operatorname{curl}_g$ . A similar integration by parts formula for  $\phi \in \Lambda^0(T)$  and  $\mu \in \Lambda^1(T)$

$$\int_{(T,g)} \phi \operatorname{curl}_g \mu = \int_{(T,g)} \mu(\operatorname{rot}_g \phi) + \int_{(\partial T,g)} \phi \mu(\hat{\tau}) \quad (4.11)$$

connects the other  $\operatorname{curl}_g$  and  $\operatorname{rot}_g$  defined in (4.3a) and (4.8a).

**4.2. Covariant curl in the Regge metric.** We proceed to extend the definitions of the covariant operators to the case when the metric  $g$  is only tangential-tangential continuous (see (2.13)) across element interfaces. Let  $\check{M}$  denote the open set obtained from  $M$  by removing all the mesh vertices (after its triangulation by  $\mathcal{T}$ ). The topological manifold  $\check{M}$  can be endowed with a natural *glued smooth structure* based on the  $tt$ -continuity of  $g$ , as alluded to in the literature [16, 21, 29, 30], [33, Theorem 3.3] and [47]. This glued smooth structure is different from that given by the coordinates  $x^i$  that we have already been using (see §3.2) in which we plan to conduct all computations. A striking difference is that while  $g$  is only  $tt$ -continuous in  $x^i$ , it is fully continuous in the natural glued smooth structure.

The glued smooth structure can be understood using the following coordinate construction around any interior mesh edge  $E$ . Let  $z \in E$  and let  $U_z$  denote an open neighborhood of  $z$  not intersecting any other mesh edge or mesh vertex. Let  $d_g(\cdot, \cdot)$  denote the distance function generated by  $g$  on the manifold  $\check{M}$ . For any  $p \in U_z$ , let  $\pi(p) = \arg \min_{q \in E} d_g(q, p)$ . We use  $T_\pm, \hat{\nu}_\pm, \hat{\tau}_\pm$  introduced in (2.18). Let  $E_p$  denote the

submanifold of  $E$  connecting  $z$  to  $\pi(p)$  oriented in the  $\hat{\tau}_+$  direction. Then for any  $p \in U_z$ , define new coordinates

$$\check{x}^1(p) = \pm d_g(\pi(p), p) \quad \text{for } p \in T_{\pm}, \quad \check{x}^2(p) = \int_{(E_p, g)} 1. \quad (4.12)$$

Denote the coordinate frame of  $\check{x}^i$  by  $\check{\partial}_i$ . It can be shown [29] that  $\check{\partial}_1 = \hat{\nu}_+$  and  $\check{\partial}_2 = \hat{\tau}_+$  at points in  $U_z \cap E$ , and that  $g(\check{\partial}_i, \check{\partial}_j)$  is continuous across  $U_z \cap E$  for all  $i, j$ . Augmenting the set  $\check{M}$  with the maximal atlas giving such coordinates, we obtain a manifold with the glued smooth structure, which we continue to denote by  $\check{M}$ . Moreover,  $(\check{M}, g)$  is a Riemannian manifold with piecewise smooth and *globally continuous metric*  $g$ .

For the next result, we need the subspace of smooth vector fields

$$\mathfrak{X}_c(\check{M}) = \{X \in \mathfrak{X}(\check{M}) : \text{support of } X \text{ is relatively compact in } \check{M}\}.$$

Because the transformations  $\partial_i = (\partial \check{x}^j / \partial x^i) \check{\partial}_j$  and  $dx^i = (\partial x^i / \partial \check{x}^j) d\check{x}^j$  are smooth within mesh elements, previously defined piecewise smooth spaces like  $\mathcal{R}(\mathcal{T})$  carry over to the glued smooth structure.

**Proposition 4.1.** *For all  $\sigma \in \mathcal{R}(\mathcal{T})$  and  $\varphi \in \mathfrak{X}_c(\check{M})$ , we have*

$$\int_{(\check{M}, g)} \sigma(\text{rot}_g \varphi) = \int_{\mathcal{T}} (\text{curl}_g \sigma)(\varphi) - \int_{\partial \mathcal{T}} g(\varphi, \hat{\nu}) \sigma(\hat{\nu}, \hat{\tau}). \quad (4.13)$$

*Proof.* The integral on the left hand side of (4.13) may equally well be written as  $\int_{\mathcal{T}} \sigma(\text{rot}_g \varphi)$ , since the set of vertices excluded in  $\check{M}$  is of measure zero. Then, integrating by parts, element by element, using (4.10),

$$\int_{\mathcal{T}} \sigma(\text{rot}_g \varphi) = \int_{\mathcal{T}} (\text{curl}_g \sigma)(\varphi) - \int_{\partial \mathcal{T}} \sigma(\varphi, \hat{\tau}).$$

Now, using the  $g$ -orthogonal decomposition  $\varphi = g(\varphi, \hat{\tau})\hat{\tau} + g(\varphi, \hat{\nu})\hat{\nu}$ , we have

$$\int_{\partial \mathcal{T}} \sigma(\varphi, \hat{\tau}) = \int_{\partial \mathcal{T}} g(\varphi, \hat{\nu}) \sigma(\hat{\nu}, \hat{\tau}) + \int_{\partial \mathcal{T}} g(\varphi, \hat{\tau}) \sigma(\hat{\tau}, \hat{\tau}).$$

Boundary mesh edges do not contribute to the last integral since  $\varphi$  is compactly supported. Across an interior mesh edge, since  $g$  is continuous in the glued smooth structure of  $\check{M}$ , and since  $\sigma$  is  $tt$ -continuous, the contributions to the last integral from adjacent elements cancel each other. Hence the last integral is zero and (4.13) follows.  $\square$

We use  $(\check{M}, g)$  to extend the definition of covariant curl. Recall that the adjoint of  $\text{curl}_g$  is  $\text{rot}_g$ , as shown by (4.10). Hence, as in the theory of Schwartz distributions, a natural extension would be to consider  $\text{curl}_g \sigma$ , for  $\sigma \in \mathcal{R}(\mathcal{T})$ , as a linear functional on  $\mathfrak{X}_c(\check{M})$  defined by

$$\begin{aligned} \langle \text{curl}_g \sigma, \varphi \rangle_{\mathfrak{X}_c(\check{M})} &= \int_{(\check{M}, g)} \sigma(\text{rot}_g \varphi) \\ &= \int_{\mathcal{T}} (\text{curl}_g \sigma)(\varphi) - \int_{\partial \mathcal{T}} g(\varphi, \hat{\nu}) \sigma(\hat{\nu}, \hat{\tau}) \end{aligned} \quad (4.14)$$



for all  $\varphi \in \mathfrak{X}_c(\check{M})$ , where we have used Proposition 4.1 in the second equality. The next key observation is that we may extend the above functional to act on a piecewise smooth vector field  $W$  instead of the smooth  $\varphi$ , *provided  $W$  is  $g$ -normal continuous*:

**Definition 4.2.** For any  $\sigma \in \mathcal{R}(\mathcal{T})$ , extend  $\text{curl}_g \sigma$  as a linear functional on  $\mathring{W}_g(\mathcal{T})$  by

$$\langle \text{curl}_g \sigma, W \rangle_{\mathring{W}_g(\mathcal{T})} = \int_{\mathcal{T}} (\text{curl}_g \sigma)(W) - \int_{\partial \mathcal{T}} g(W, \hat{\nu}) \sigma(\hat{\nu}, \hat{\tau}) \quad (4.15)$$

for all  $W \in \mathring{W}_g(\mathcal{T})$ , the space defined in (2.19). The rationale for this definition is that there are functions  $\varphi$  in  $\mathfrak{X}_c(\check{M})$  approaching  $W \in \mathring{W}_g(\mathcal{T})$  in such a way that the right hand side of (4.14) converges to that of (4.15): see Proposition A.1. Furthermore, because of the  $g$ -normal continuity of  $W$ , the last term in (4.15) vanishes when  $\sigma$  is globally smooth, so (4.15) indeed extends  $\text{curl}_g$  on smooth functions.

We now develop a formula for computing the extended covariant curl in the  $x^i$  coordinates (not  $\check{x}^i$ ). We use the Euclidean parameter domain  $(\Omega, \delta)$  and the Euclidean  $\delta$ -orthonormal tangent and normal ( $\tau$  and  $\nu$ ) on element boundaries (see §3.2). Abbreviating  $\delta(w, X)$  to  $w_X$ , (2.19) for  $g = \delta$  reads

$$\mathring{W}(\Omega) = \{w \equiv [w^1, w^2] : \Omega \rightarrow \mathbb{R}^2 \mid w^i \in C^\infty(\mathcal{T}), \llbracket w_\nu \rrbracket = 0, w_\nu|_{\partial\Omega} = 0\}.$$

Since standard divergence-conforming finite elements, as Brezzi–Douglas–Marini (BDM) [14] and Raviart–Thomas (RT) [38], are contained in this space, it is more convenient for computations than (2.19). For any  $w = [w^1, w^2] \in \mathring{W}(\Omega)$ , let

$$Q_g w = \frac{w^1(x^1, x^2) \partial_1 + w^2(x^1, x^2) \partial_2}{\sqrt{\det g}}. \quad (4.16)$$

**Proposition 4.3** (Extended covariant curl in computational coordinates).

- (1) A vector field  $w \in \mathring{W}(\Omega)$  if and only if  $Q_g w$  is in  $\mathring{W}_g(\mathcal{T})$ .
- (2) For any  $\sigma \in \mathcal{R}(\mathcal{T})$  and  $w \in \mathring{W}(\Omega)$ ,

$$\begin{aligned} & \langle \text{curl}_g \sigma, Q_g w \rangle_{\mathring{W}_g(\mathcal{T})} \\ &= \sum_{T \in \mathcal{T}} \left( \int_T \frac{[\text{curl}[\sigma]]_i w^i - \sigma_{mk} \varepsilon^{jk} \Gamma_{ji}^m w^i}{\sqrt{\det g}} \, da - \int_{\partial T} \frac{g_{\tau\tau} \sigma_{\nu\tau} - g_{\nu\tau} \sigma_{\tau\tau}}{g_{\tau\tau} \sqrt{\det g}} w_\nu \, dl \right) \end{aligned} \quad (4.17a)$$

$$= \sum_{T \in \mathcal{T}} \left( \int_T \sigma_{mk} \frac{[\text{rot}[w]]^{mk} - \varepsilon^{kj} (\Gamma_{lj}^l w^m - \Gamma_{ji}^m w^i)}{\sqrt{\det g}} \, da + \int_{\partial T} \frac{\sigma_{\tau\tau} g_{i\tau} w^i}{g_{\tau\tau} \sqrt{\det g}} \, dl \right). \quad (4.17b)$$

*Proof.* By (3.17) and (2.9),

$$w_\nu = \delta(w, \nu) = g(w, \tilde{\nu}) = g(w, \hat{\nu} \sqrt{g_{\tilde{\nu}\tilde{\nu}}}) = g(w, \hat{\nu}) \frac{\sqrt{g_{\tau\tau}}}{\sqrt{\det g}} = g(Q_g w, \hat{\nu}) \sqrt{g_{\tau\tau}}. \quad (4.18)$$

Hence the continuity of  $g_{\tau\tau}$  across element interfaces implies that

$$\llbracket g(Q_g w, \hat{\nu}) \rrbracket = 0 \quad \text{if and only if} \quad \llbracket w_\nu \rrbracket = 0.$$

Similarly  $w_\nu|_{\partial\Omega} = 0$  if and only if  $g(Q_g w, \hat{\nu}) = 0$  vanishes on the boundary. Hence  $Q_g w \in \mathring{W}_g(\mathcal{T})$  if and only if  $w \in \mathring{W}(\Omega)$ .

To prove (4.17a), we write (4.15) in coordinates. Observe that the coordinate expression for covariant curl in (4.7b) and the integration formula (2.20) imply

$$\int_{\mathcal{T}} (\operatorname{curl}_g \sigma)(Q_g w) = \sum_{T \in \mathcal{T}} \int_T \frac{[\operatorname{curl}[\sigma]]_i [Q_g w]^i - \sigma_{mk} \varepsilon^{jk} \Gamma_{ji}^m [Q_g w]^i}{\sqrt{\det g}} \sqrt{\det g} \, da,$$

which coincides with the sum of integrals over elements  $T$  in (4.17a). For the element boundary integrals of  $g(Q_g w, \hat{\nu}) \sigma(\hat{\nu}, \hat{\tau})$  contributing to  $\langle \operatorname{curl}_g \sigma, Q_g w \rangle_{\dot{W}_g(\mathcal{T})}$ , first note that (4.18) implies

$$g(Q_g w, \hat{\nu}) \sigma(\hat{\nu}, \hat{\tau}) = \frac{w_\nu}{\sqrt{g_{\tau\tau}}} \frac{\sigma(\delta(\hat{\nu}, \nu) \nu + \delta(\hat{\nu}, \tau) \tau, \tau)}{\sqrt{g_{\tau\tau}}}. \quad (4.19)$$

To proceed, we observe

$$\begin{aligned} \delta(\hat{\nu}, \nu) &= g(\hat{\nu}, \tilde{\nu}) = g(\hat{\nu}, \hat{\nu} \sqrt{g_{\tilde{\nu}\tilde{\nu}}}) && \text{by (3.17) and (2.10)} \\ &= \frac{\sqrt{g_{\tau\tau}}}{\sqrt{\det g}} && \text{by (2.9),} \\ \delta(\hat{\nu}, \tau) &= \frac{\delta(\tilde{\nu}, \tau)}{\sqrt{g_{\tilde{\nu}\tilde{\nu}}}} = \frac{\sqrt{\det g}}{\sqrt{g_{\tau\tau}}} \delta(\tilde{\nu}, \tau) && \text{by (2.10) and (2.9),} \\ &= -\frac{\sqrt{\det g}}{\sqrt{g_{\tau\tau}}} g^{kj} \varepsilon_{ji} \tau^i \delta_{km} \tau^m && \text{by (2.11),} \\ &= \frac{-g(\nu, \tau)}{\sqrt{g_{\tau\tau} \det g}}, \end{aligned}$$

where, in the last step, we have simplified using the cofactor expansion of  $g^{kj}$  and (3.16). Hence (4.19) implies

$$g(Q_g w, \hat{\nu}) \sigma(\hat{\nu}, \hat{\tau}) = \frac{w_\nu}{g_{\tau\tau}} \left( \frac{\sqrt{g_{\tau\tau}}}{\sqrt{\det g}} \sigma_{\nu\tau} - \frac{g_{\nu\tau}}{\sqrt{g_{\tau\tau} \det g}} \sigma_{\tau\tau} \right) = \frac{w_\nu (\sigma_{\nu\tau} g_{\tau\tau} - g_{\nu\tau} \sigma_{\tau\tau})}{g_{\tau\tau} \sqrt{g_{\tau\tau} \det g}}.$$

Integrating this over each element boundary using the measure  $\sqrt{g_{\tau\tau}} \, dl$  and summing over elements, the right hand sides of (4.15) and (4.17a) are seen to be the same.

To prove the second identity (4.17b), consider any  $W \in \dot{W}_g(\mathcal{T})$ . We start by applying the integration by parts formula (4.10) to the first term on the right hand side of the definition (4.15):

$$\begin{aligned} \langle \operatorname{curl}_g \sigma, W \rangle_{\dot{W}_g(\mathcal{T})} &= \int_{\mathcal{T}} \sigma(\operatorname{rot}_g W) + \int_{\partial T} \sigma(W, \hat{\tau}) - \int_{\partial T} g(W, \hat{\tau}) \sigma(\hat{\nu}, \hat{\tau}) \\ &= \int_{\mathcal{T}} \sigma(\operatorname{rot}_g W) + \int_{\partial \mathcal{T}} g(W, \hat{\tau}) \sigma_{\hat{\tau}\hat{\tau}} \end{aligned}$$

after simplifying using  $g$ -orthogonal decomposition  $W = g(W, \hat{\tau}) \hat{\tau} + g(W, \hat{\nu}) \hat{\nu}$  on element boundaries. Substituting  $W = Q_g w$ , applying the quotient rule to compute  $\operatorname{rot}_g(Q_g w)$  using (4.6), and expressing the result in  $x^i$  coordinates, we obtain (4.17b).  $\square$



*Proof.* By (4.21), (4.15), (4.11), and (4.20),

$$\begin{aligned} \langle \text{inc}_g \sigma, u \rangle_{\hat{\mathcal{V}}(\mathcal{T})} &= \int_{\mathcal{T}} (\text{curl}_g \sigma)(\text{rot}_g u) - \int_{\partial \mathcal{T}} \sigma_{\hat{\nu}\hat{\tau}} g(\text{rot}_g u, \hat{\nu}) \\ &= \int_{\mathcal{T}} u \text{inc}_g \sigma - \int_{\partial \mathcal{T}} u (\text{curl}_g \sigma)(\hat{\tau}) + \int_{\partial \mathcal{T}} \sigma_{\hat{\nu}\hat{\tau}} (d^0 u)(\hat{\tau}). \end{aligned}$$

Now, on the edge  $E_i$  oriented from end point  $V_{i-1}$  to  $V_i$  (indexed mod 3; see Figure 1), by the one-dimensional integration by parts formula

$$\int_{(E_i, g)} \sigma_{\hat{\nu}\hat{\tau}} (d^0 u)(\hat{\tau}) = \sigma_{\hat{\nu}\hat{\tau}}(V_i) u(V_i) - \sigma_{\hat{\nu}\hat{\tau}}(V_{i-1}) u(V_{i-1}) - \int_{(E_i, g)} (d^0 \sigma_{\hat{\nu}\hat{\tau}})(\hat{\tau}) u.$$

When summing over the three edges  $E_i \subset \partial T$ , the above vertex values of  $\sigma_{\hat{\nu}\hat{\tau}}$  yield vertex jumps. Hence the first statement follows by summing over all  $T$  in  $\mathcal{T}$ .

To prove the second statement, note that the edge integrals from adjacent triangles cancel each other. To show that the last term with vertex contributions also vanish, let  $\mathcal{E}_V$  denote the set of vertices connected to  $V \in \mathcal{V}$  by an edge and let  $\hat{\tau}_V$  denote the  $g$ -unit tangent vector along an  $E \in \mathcal{E}_V$  pointing away from  $V$ . Then the jump  $\llbracket \sigma(\hat{\nu}, \hat{\tau}_V) \rrbracket$  on any edge  $E \in \mathcal{E}_V$  is defined as before, using (2.18). Its limit as we approach a vertex  $V$  along any edge  $E \in \mathcal{E}_V$ , is denoted by  $\llbracket \sigma \rrbracket_{\hat{\nu}\hat{\tau}}^V$ . Using it, the last sum can be rearranged to

$$\sum_{T \in \mathcal{T}} \sum_{V \in \mathcal{V}_T} \llbracket \sigma_{\hat{\nu}\hat{\tau}} \rrbracket_V^T u(V) = - \sum_{V \in \mathcal{V}^{\text{int}}} \sum_{E \in \mathcal{E}_V} \llbracket \sigma \rrbracket_{\hat{\nu}\hat{\tau}}^V u(V).$$

where  $\mathcal{V}^{\text{int}} \subset \mathcal{V}$  is the subset of mesh vertices in the interior of the domain. Each summand on the right hand side vanishes when  $\sigma$  is smooth.  $\square$

## 5. LINEARIZATION

**5.1. Connecting curvature with incompatibility.** The variational derivative of  $f : \mathcal{S}(T) \rightarrow \mathbb{R}$  at a  $\rho \in \mathcal{S}(T)$  in the direction of a  $\sigma \in \mathcal{S}(T)$ , denoted by  $D_\sigma(f)(\rho)$  or  $D_\sigma(f(\rho))$  is given by

$$D_\sigma(f)(\rho) \equiv D_\sigma(f(\rho)) = \lim_{t \rightarrow 0} \frac{f(\rho + t\sigma) - f(\rho)}{t} \quad (5.1)$$

when the limit exists. Note that the changes in  $\rho(p)$  and  $\sigma(p)$  as  $p$  varies in  $M$  are immaterial in the above definition. Hence, we may use Riemann normal coordinates [31] to prove the pointwise identities in the next result. Let  $\tilde{x}^i$  denote the Riemann normal coordinates given by a chart covering a point  $p \in T$ ,  $\tilde{\partial}_i$  denote the corresponding coordinate frame,  $\tilde{\sigma}_{ij} = \sigma(\tilde{\partial}_i, \tilde{\partial}_j)$  for any  $\sigma \in \mathcal{T}_0^2(T)$ ,  $\tilde{R}_{ijkl} = R(\tilde{\partial}_i, \tilde{\partial}_j, \tilde{\partial}_k, \tilde{\partial}_l)$ , and let  $\tilde{\Gamma}_{ijk}, \tilde{\Gamma}_{ij}^k$  be defined by (2.7) with  $\partial_i$  replaced by  $\tilde{\partial}_i$ . Then, by [31, Proposition 5.11],

$$\tilde{g}_{ij}|_p = \delta_{ij}, \quad \tilde{\partial}_k \tilde{g}_{ij}|_p = 0, \quad \tilde{\Gamma}_{ijk}|_p = 0, \quad (5.2)$$

which greatly simplify calculations. As an example, consider the expression for covariant incompatibility of a  $\sigma \in \mathcal{S}(T)$  in coordinates given by (4.7c). It simplifies in normal

coordinates, by virtue of (5.2), to  $[\text{inc}_g \sigma] = \text{inc}[\tilde{\sigma}] - \varepsilon^{ij} \varepsilon^{kl} \tilde{\sigma}_{ml} \tilde{\partial}_i \tilde{\Gamma}_{jk}^m$ . Expanding out the last term in terms of  $\tilde{g}_{ij}$  using (2.7b) and simplifying,

$$[\text{inc}_g \sigma] = \text{inc}[\tilde{\sigma}] - \frac{1}{2} \text{tr}[\tilde{\sigma}] \text{inc}[\tilde{g}]. \quad (5.3)$$

**Lemma 5.1** (Variation of curvature). *On any element manifold  $(T, g)$ ,  $T \in \mathcal{T}$ ,*

$$D_\sigma(K(g)\text{vol}_{(T,g)})(g) = -\frac{1}{2}\text{vol}_{(T,g)} \text{inc}_g \sigma, \quad \text{at every point in } T, \quad (5.4a)$$

$$D_\sigma(\kappa(g)\text{vol}_{(\partial T,g)})(g) = \frac{1}{2}(\text{curl}_g \sigma + d^0 \sigma_{\hat{\tau}\hat{\nu}})(\hat{\tau}), \quad \text{at points on edges of } \partial T, \quad (5.4b)$$

$$D_\sigma(\triangleleft(\delta)_V^T - \triangleleft(g)_V^T)(g) = -\frac{1}{2} \llbracket \sigma_{\hat{\nu}\hat{\tau}} \rrbracket_V^T \quad \text{at every vertex of } T. \quad (5.4c)$$

*Proof.* We provide an elementary proof of equation (5.4a) employing the readily verifiable identity  $D_\sigma(\text{vol}_{(T,g)})(g) = \frac{1}{2}\text{vol}_{(T,g)} \text{tr}(g^{-1}\sigma)$ . Since (2.2) implies  $K(g) = R_{1221}/\det g$ , the variation of the left-hand side reads

$$D_\sigma(K(g)\text{vol}_{(T,g)})(g) = \frac{1}{\sqrt{\det g}} \left( D_\sigma(R_{1221})(g) - \frac{1}{2} \text{tr}(g^{-1}\sigma) R_{1221} \right).$$

Next, since (5.2) implies that the last two terms of (3.5) vanish in normal coordinates,  $\tilde{R}_{1221} = \tilde{\partial}_1 \tilde{\Gamma}_{221} - \tilde{\partial}_2 \tilde{\Gamma}_{211} = -\frac{1}{2} \text{inc}[\tilde{g}]$ . Hence,

$$D_\sigma(K(g)\text{vol}_{(T,g)})(g) = -\frac{1}{2} \text{inc}[\tilde{\sigma}] + \frac{1}{4} \text{tr}[\tilde{\sigma}] \text{inc}[\tilde{g}],$$

so (5.4a) follows from (5.3).

To prove (5.4b), we observe that the expression for geodesic curvature from Lemma 3.3 reads as follows in normal coordinates

$$\begin{aligned} D_\sigma(\kappa_g(g)\text{vol}_{(\partial T,g)})(g) &= \frac{\sqrt{\det g}}{g_{\tau\tau}} \left[ \left( \frac{1}{2} \text{tr}_g \sigma - \frac{\sigma_{\tau\tau}}{g_{\tau\tau}} \right) (\Gamma_{\tau\tau}^\nu(g) + \dot{\tau}_\nu) \right. \\ &\quad \left. - \nu_i g^{ij} \sigma_{jk} g^{kl} \Gamma_{\tau\tau l}(g) + \nu_i g^{ij} \Gamma_{\tau\tau j}(\sigma) \right] \\ &= \left( \frac{1}{2} \text{tr}(\tilde{\sigma}) - \tilde{\sigma}_{\tau\tau} \right) \dot{\tau}_\nu + (\tilde{\partial}_\tau \tilde{\sigma})_{\nu\tau} - \frac{1}{2} (\tilde{\partial}_\nu \tilde{\sigma})_{\tau\tau}. \end{aligned}$$

Note that

$$\dot{\nu} = -\dot{\tau}_\nu \tau, \quad \dot{\tau} = \dot{\tau}_\nu \nu. \quad (5.5)$$

This follows by differentiating the equation  $\delta(\nu, \nu) = 1$  with respect to  $\tau$ , yielding  $\delta(\dot{\nu}, \nu) = 0$ . Hence there must be a scalar  $\alpha$  such that  $\dot{\nu} = \alpha\tau$ . Now differentiating  $\delta(\nu, \tau) = 0$ , we find that the  $\alpha$  must satisfy  $\delta(\alpha\tau, \tau) + \delta(\nu, \dot{\tau}) = 0$ , so  $\dot{\nu} = -\delta(\nu, \tau)\tau$  thus proving the first identity in (5.5). A similar argument using  $\delta(\tau, \tau) = 1$  shows that  $\dot{\tau} = \delta(\nu, \dot{\tau})\nu$ . As the distributional incompatibility operator is independent of the coordinate system we can also use normal coordinates leading for the edge boundary term,  $\hat{\tau} = \tau$  and  $\hat{\nu} = \nu$ ,

$$\begin{aligned} -(\text{curl}_g \sigma + d^0 \sigma_{\hat{\tau}\hat{\nu}})(\hat{\tau}) &= -(\text{curl } \tilde{\sigma}) \cdot \tau - (\tilde{\partial}_\tau \tilde{\sigma})_{\nu\tau} - \tilde{\sigma}_{\dot{\nu}\tau} - \tilde{\sigma}_{\nu\dot{\tau}} \\ &= -(\text{curl } \tilde{\sigma}) \cdot \tau - (\tilde{\partial}_\tau \tilde{\sigma})_{\nu\tau} - (\tilde{\sigma}_{\nu\nu} - \tilde{\sigma}_{\tau\tau}) \dot{\tau}_\nu, \end{aligned}$$

where we used (5.5). With  $-(\operatorname{curl} \tilde{\sigma}) \cdot \tau = -(\tilde{\partial}_\tau \tilde{\sigma})_{\nu\tau} + (\tilde{\partial}_\nu \tilde{\sigma})_{\tau\tau}$  the second identity follows.

Equation (5.4c) was proved in [11, Proposition 2.17] in terms of evolving metric tensors restricting the set of variational directions  $\sigma$ . The proof, however, can be directly adapted as there holds  $D_\sigma(\hat{\tau})(g) = -\frac{\sigma_{\tau\tau}}{2(g_{\tau\tau})^{3/2}}\tau$  which is still parallel to  $\hat{\tau} = (g_{\tau\tau})^{-1/2}\tau$ .  $\square$

By having the variational derivative of all sources of curvature the (distributional) Gauss curvature can be rewritten in the following integral representation. For ease of presentation we assume that homogeneous Dirichlet data are prescribed on the whole boundary,  $K(\bar{g})|_{\partial\Omega} = 0$ .

**Lemma 5.2.** *There holds for  $g \in \mathcal{R}^+(\mathcal{T})$ ,  $k \in \mathbb{N}_0$ ,  $K_h(g) \in \hat{\mathcal{V}}_h^{k+1}$  from Definition 3.1, the exact metric  $\bar{g}$  and its Gauss curvature  $K(\bar{g})$*

$$\int_{\mathcal{T}} K_h(g) u_h = \frac{1}{2} \int_0^1 b_h(\delta + t(g - \delta), g - \delta, u_h) dt, \quad \text{for all } u_h \in \hat{\mathcal{V}}_h^{k+1}, \quad (5.6)$$

$$\int_{\mathcal{T}} K(\bar{g}) u = \frac{1}{2} \int_0^1 b_h(\delta + t(\bar{g} - \delta), \bar{g} - \delta, u) dt, \quad \text{for all } u \in \hat{\mathcal{V}}(\mathcal{T}), \quad (5.7)$$

with the distributional covariant incompatibility operator (4.21)

$$b_h(g, \sigma, u) := -\langle \operatorname{inc}_g \sigma, u \rangle_{\hat{\mathcal{V}}(\mathcal{T})}. \quad (5.8)$$

*Proof.* The claim follows directly from consistency Lemma 3.5, variation Lemma 5.1, and the fundamental theorem of calculus in combination that the Euclidean metric  $\delta$  has zero curvature.  $\square$

*Remark 5.3.* The idea of using an integral representation of the distributional Gauss curvature has been introduced in [27, 11]. It is based on the computation

$$D_\sigma(K(g)\operatorname{vol}_{(T,g)})(g) = \frac{1}{2}(\operatorname{div}_g \operatorname{div}_g S_g \sigma) \operatorname{vol}_{(T,g)},$$

where a proof using [23] was suggested. A comparison between the distributional  $\operatorname{div}_g \operatorname{div}_g S_g$  and  $\operatorname{inc}_g$  is given in Appendix A.1. We prefer to use  $\operatorname{inc}$  as it is the natural differential operator for Regge elements in any dimension.

**5.2. Evolution of connection 1-form.** The variation of the connection 1-form  $\varpi$  is more involved than the Gauss curvature. It requires a  $g$ -orthonormal frame  $(e_1, e_2)$ , see (2.5), and depending on the chosen frame different representations of  $\varpi$  are obtained.

Following [11] a canonical connection 1-form will be fixed in terms of evolving metrics. Define for given metric tensor  $g$  the parameterized  $g(t) := \delta + t(g - \delta)$  such that  $g(0) = \delta$  and  $g(1) = g$ . Denote the derivative by  $\sigma := \frac{\partial}{\partial t} g(t) = g - \delta$ . At  $t = 0$  the Euclidean metric is considered and therefore define the Euclidean basis vectors  $E_1, E_2$  as orthonormal frame. By transforming these frame by  $e_1(t) = u(t)E_1$ ,  $e_2(t) = u(t)E_2$ , where  $u$  satisfies the ODE [11]

$$\dot{u}(t)u(t)^{-1} = -\frac{1}{2}(g - \delta)^\sharp, \quad u(0) = \operatorname{id} \quad (5.9)$$

and  $(g - \delta)^\sharp$  has to be understood with respect to the metric  $(1 - t)\delta + tg$ ,  $(e_1(t), e_2(t))$  build a  $g(t)$ -orthonormal frame for each  $t \in [0, 1]$ . In elementary form the transformation  $u(t)$ , needed later for the numerical experiments in §7, reads as follows.

**Lemma 5.4.** *The solution of ODE (5.9) is given by  $u(t) = ((1-t)\delta + tg)^{-\frac{1}{2}}$ . Especially, there holds  $u(1) = g^{-\frac{1}{2}}$ .*

*Proof.* The spectral decomposition of  $g = V\Lambda V^{-1}$  is independent of  $t$ . Use that  $(1 - t)\delta + tg = V((1 - t)\delta + t\Lambda)V^{-1}$  and thus  $u(t) = VD(t)V^{-1}$ , with the diagonal matrix  $D(t) = \text{diag}\left(\left((1 - t)1 + t\Lambda_{ii}\right)^{-\frac{1}{2}}\right)$ . Then

$$\dot{u}(t) = V\dot{D}(t)V^{-1} = -\frac{1}{2}V\tilde{D}(t)V^{-1}, \quad \tilde{D}(t) = \text{diag}\left(\left((1 - t)1 + t\Lambda_{ii}\right)^{-\frac{3}{2}}(\Lambda_{ii} - 1)\right).$$

On the other hand

$$\begin{aligned} -\frac{1}{2}(g - \delta)^\sharp u(t) &= -\frac{1}{2}((1 - t)\delta + tg)^{-1}(g - \delta)VD(t)V^{-1} \\ &= -\frac{1}{2}V\text{diag}\left(\left((1 - t) + t\Lambda_{ii}\right)^{-1}\right)V^{-1}V\text{diag}(\Lambda_{ii} - 1)V^{-1}VD(t)V^{-1} \\ &= -\frac{1}{2}V\text{diag}\left(\left((1 - t) + t\Lambda_{ii}\right)^{-1}\right)\text{diag}(\Lambda_{ii} - 1)D(t)V^{-1} \\ &= -\frac{1}{2}V\tilde{D}(t)V^{-1}. \end{aligned}$$

□

**Lemma 5.5.** *With the notation defined in this section there holds*

$$\frac{\partial}{\partial t}\varpi(g(t)) = -\frac{1}{2}\text{curl}_{g(t)}\sigma. \quad (5.10)$$

*Proof.* Using formula (6) and Proposition 2.14 from [11] (where the authors defined the connection 1-form with the opposite sign), where we can set  $f = 0$  due to the specific evolution equation  $u(t)$  described above, gives

$$\frac{\partial}{\partial t}\varpi(g(t)) = -\frac{1}{2}(\nabla_{e_1}\sigma e_2 - \nabla_{e_2}\sigma e_1) = \frac{1}{2}\star(\text{div}_{g(t)}S_{g(t)}\sigma)^\flat = -\frac{1}{2}\text{curl}_{g(t)}\sigma,$$

where we used Lemma A.2 for the last equality. □

As discussed in [11] the connection 1-form is not unique. Using the  $g$ -orthonormal frame  $e_1 = g^{-1/2}E_1$ ,  $e_2 = g^{-1/2}E_2$  defined above the canonical connection 1-form entails again an integral representation.

We will use normal continuous finite elements to approximate the connection 1-form  $\varpi(\bar{g})$ . They are sub-spaces of (2.19) by setting using  $g = \delta$  in the definition. The finite element space and its subspaces with essential boundary conditions are defined by,  $k \in \mathbb{N}_0$ ,

$$\begin{aligned} \mathcal{W}_h^k &= \{w \in \mathcal{W}_\delta(\mathcal{T}) : \text{for all } T \in \mathcal{T}, w|_T = \Phi_T^*\hat{w} \text{ for some } \hat{w} \in \mathcal{P}^k(\hat{T}, \mathbb{R}^2)\}, \\ \mathring{\mathcal{W}}_{h,\Gamma}^k &= \{w \in \mathcal{W}_h^k : \delta(w|_\Gamma, \nu) = 0\} \quad \text{and} \quad \mathring{\mathcal{W}}_h^k = \mathring{\mathcal{W}}_{h,\partial\Omega}^k, \end{aligned} \quad (5.11)$$

where the pull-back  $\Phi_T^* \hat{w}$  is the Piola transformation. For  $k > 0$  (5.11) coincides with *Brezzi–Douglas–Marini finite element space* (BDM) [14]. *Raviart–Thomas finite elements* (RT) [38] are defined similarly by requiring that  $\hat{w} \in \mathcal{P}^k(\hat{T}, \mathbb{R}^2) + \mathcal{P}^k(\hat{T}) (x_1 \ x_2)^\top$ .

**Definition 5.6.** Let  $g \in \mathcal{R}^+(\mathcal{T})$ ,  $k \in \mathbb{N}_0$ . The 1-form  $\varpi_h(g) \in \mathring{W}_h^k$  defined by

$$\int_{\mathcal{T}} \varpi_h(g) Q_g v_h = \frac{1}{2} \int_0^1 c_h(\delta + t(g - \delta), g - \delta, v_h) dt, \quad \text{for all } v_h \in \mathring{W}_h^k, \quad (5.12)$$

with  $Q_g$  from (4.16) and the distributional covariant curl (4.17)

$$c_h(g, \sigma, v) := -\langle \text{curl}_g \sigma, Q_g v \rangle_{\mathring{W}_g(\mathcal{T})}, \quad (5.13)$$

is called the *discrete canonical lifted connection 1-form*.

**Lemma 5.7.** *The exact canonical connection 1-form  $\varpi(\bar{g})$  fulfills*

$$\int_{\mathcal{T}} \varpi(\bar{g}) Q_g v = \frac{1}{2} \int_0^1 c_h(\delta + t(\bar{g} - \delta), \bar{g} - \delta, v) dt, \quad \text{for all } v \in \mathring{W}(\Omega). \quad (5.14)$$

*Proof.* The claim follows directly by the variation Lemma 5.5 and noting that for smooth  $\bar{g}$  the boundary terms of (4.17) vanish.  $\square$

*Remark 5.8.* There holds by inserting  $v_h := \text{rot}_\delta u_h$ ,  $u_h \in \mathring{V}_h^{k+1}$ , in (5.12)

$$\begin{aligned} \int_{\mathcal{T}} \varpi_h(g) Q_g \text{rot}_\delta u_h &= \frac{1}{2} \int_0^1 c_h(\delta + t(g - \delta), g - \delta, \text{rot}_\delta u_h) dt \\ &= \frac{1}{2} \int_0^1 b_h(\delta + t(g - \delta), g - \delta, u_h) dt = \int_{\mathcal{T}} K_h(g) u_h, \end{aligned}$$

where we used the identity

$$c_h(g, \sigma, \text{rot}_\delta u_h) = b_h(g, \sigma, u_h),$$

which can be proven directly with integration by parts. Theorem 6.2 will justify that  $\varpi_h(g)$  is the appropriate approximation of the exact connection 1-form  $\varpi(\bar{g})$ .

*Remark 5.9.* In [11] the dual (Hodge-star) of the connection 1-form is considered and its discrete lifting is approximated in the  $H(\text{curl})$ -conforming Nédélec finite element space, which is tangential continuous. In this work, however, we directly approximate the connection 1-form with the usage of  $H(\text{div})$ -conforming BDM elements.

## 6. ERROR ANALYSIS

**6.1. Statement of Theorems.** In the following we will assume that homogeneous Dirichlet data is prescribed on the whole boundary, i.e.,  $K(\bar{g})|_{\partial\Omega} = 0$  and  $\varpi(\bar{g})|_{\partial\Omega} \cdot \nu = 0$ , where  $\Omega \subset \mathbb{R}^2$  is a bounded domain with polygonal boundary. We note that the analysis can be extended to the case of non-homogeneous Dirichlet boundary conditions and for the Gauss curvature including Neumann data, compare Definition 3.1 and numerical examples in §7. Further, the triangulations  $\mathcal{T}$  of  $\Omega$  are assumed to be shape-regular and quasi-uniform such that we can define  $h := \max_{T \in \mathcal{T}} h_T$ , with



$h_T = \text{diam}(T)$ .

The main result of this work is stated in the following theorem. The involved function spaces and norms are defined in §6.2 below.

**Theorem 6.1.** *Let  $k \in \mathbb{N}_0$ ,  $\bar{g} \in W^{k+1,\infty}(\Omega)$  with  $K(\bar{g}) \in H^k(\Omega)$ , and  $g_h = \mathcal{J}_h^{\mathcal{R},k} \bar{g}$  the Regge interpolant. Then there holds for the lifted Gauss curvature  $K_h(g_h) \in \mathring{\mathcal{V}}_h^{k+1}$  defined in (3.19) for sufficiently small  $h$*

$$\begin{aligned} \|K_h(g_h) - K(\bar{g})\|_{H^{-1}(\Omega, \bar{g})} &\leq C \left( \|\bar{g} - g_h\|_{L^\infty(\Omega)} + h \|\bar{g} - g_h\|_{W_h^{1,\infty}(\Omega)} \right. \\ &\quad \left. + h \inf_{u_h \in \mathring{\mathcal{V}}_h^{k+1}} \|K(\bar{g}) - u_h\|_{L^2(\Omega)} \right) \quad (6.1) \\ &\leq Ch^{k+1} (\|\bar{g}\|_{W^{k+1,\infty}(\Omega)} + |K(\bar{g})|_{H^k(\Omega)}), \end{aligned}$$

where  $C = C(\|\bar{g}\|_{W^{1,\infty}}, \|\bar{g}^{-1}\|_{L^\infty}, K(\bar{g}), \Omega, \mathcal{T})$ .

Theorem 6.1 shows that convergence in the  $H^{-1}$ -norm is already obtained for lowest order Regge elements,  $g_h = \mathcal{J}_h^{\mathcal{R},0} \bar{g}$ . This looks surprising at the first glance as one would expect only convergence in the  $H^{-2}$ -norm if a second order derivative of a piecewise constant metric is approximated. Due to special orthogonality properties (presented below), which are valid for Regge finite elements, we however obtain one additional order of convergence, improving the results in [27, 11].

We obtain also the following improved convergence results for the connection 1-form.

**Theorem 6.2.** *Let  $k \in \mathbb{N}_0$ ,  $\bar{g} \in H^{k+1}(\Omega)$  with  $\varpi(\bar{g}) \in H^k(\Omega)$ , and  $g_h = \mathcal{J}_h^{\mathcal{R},k} \bar{g}$  the Regge interpolant. Then there holds for the lifted connection 1-form  $\varpi_h(g_h) \in \mathring{\mathcal{W}}_h^k$  defined in (5.12) for sufficiently small  $h$*

$$\begin{aligned} \|\varpi_h(g_h) - \varpi(\bar{g})\|_{L^2(\Omega)} &\leq C \left( \|\bar{g} - g_h\|_{L^2(\Omega)} + h \|\bar{g} - g_h\|_{H_h^1(\Omega)} + h \inf_{v_h \in \mathring{\mathcal{W}}_h^k} \|\varpi(\bar{g}) - v_h\|_{L^2(\Omega)} \right) \\ &\leq Ch^{k+1} (\|\bar{g}\|_{H^{k+1}(\Omega)} + |\varpi(\bar{g})|_{H^k}), \quad (6.2) \end{aligned}$$

where  $C = C(\|\bar{g}\|_{W^{1,\infty}}, \|\bar{g}^{-1}\|_{L^\infty}, \varpi(\bar{g}), \Omega, \mathcal{T})$ .

In the next subsections we summarize and present important properties and estimates, which are then used to proof optimal convergence rates of the distributional covariant curl, which is the main contribution of this paper. From these results we can then directly deduce the optimal convergence of the distributional covariant incompatibility operator. After this, we are in the position to present the proofs of the improved convergence rates for the lifted curvature and connection 1-form.

**6.2. Preliminaries.** For  $s \geq 0$  and  $p \in [1, \infty]$  we denote with  $W^{s,p}(\Omega)$  and  $\|\cdot\|_{W^{s,p}(\Omega)}$  the standard Sobolev spaces with corresponding norms. Further, we define the element-wise norms

$$\|u\|_{W_h^{s,p}(\Omega)}^p := \sum_{T \in \mathcal{T}} \|u\|_{W^{s,p}(T)}^p, \quad (6.3)$$

with the usual adaption for  $p = \infty$ , and for  $p = 2$  we write  $\|\cdot\|_{H_h^s(\Omega)} := \|\cdot\|_{W_h^{s,2}(\Omega)}$ . Further,  $|\cdot|_{W^{s,p}(\Omega)}$  denotes the usual Sobolev semi-norm summing only over derivatives

of order  $s$ . We might neglect the domain specification  $\Omega$  if no misunderstandings are possible.

Throughout what follows we assume that  $\bar{g} \in C^\infty(\Omega, \mathcal{S}^+)$  is the metric tensor of a smooth two-dimensional Riemannian manifold  $(\Omega, \bar{g})$ .

We denote for  $g \in L^\infty(\Omega, \mathcal{S}^+)$  with  $g^{-1} \in L^\infty(\Omega, \mathcal{S}^+)$  the  $g$  dependent norms for  $u, f \in L^2(\Omega)$

$$\|u\|_{L^2(\Omega, g)}^2 = (u, u)_g = \int_{\Omega} \sqrt{\det g} |u|^2 da, \quad \|f\|_{H^{-1}(\Omega, g)} = \sup_{v \in H_0^1(\Omega)} \frac{(f, v)_g}{\|v\|_{H^1(\Omega)}}. \quad (6.4)$$

**Definition 6.3.** The Regge interpolant  $\mathcal{J}_h^{\mathcal{R}, k} : C^0(\Omega, \mathcal{S}) \rightarrow \mathcal{R}_h^k$  of order  $k \in \mathbb{N}_0$  for  $g \in C^0(\Omega, \mathcal{S})$  is defined element-wise by the following equations

$$\int_E (\mathcal{J}_h^{\mathcal{R}, k} g)_{\tau\tau} q dl = \int_E g_{\tau\tau} q dl \text{ for all } q \in \mathcal{P}^k(E), \quad E \subset \partial T, \quad (6.5a)$$

$$\int_T \mathcal{J}_h^{\mathcal{R}, k} g : q da = \int_T g : q da \text{ for all } q \in \mathcal{P}^{k-1}(T, \mathcal{S}), \quad (6.5b)$$

where  $\tau$  denotes the tangential vector of the edge  $E$  and  $A : B = \sum_{i,j=1}^2 A_{ij} B_{ij}$  the Frobenius product between matrices  $A, B \in \mathbb{R}^{2 \times 2}$ .

Further, the  $L^2$ -projector  $\Pi^k : L^2(\Omega) \rightarrow \mathcal{P}^k(\mathcal{T})$  for  $u \in L^2(\Omega)$  is given by

$$\int_T \Pi^k u q da = \int_T u q da \text{ for all } q \in \mathcal{P}^k(T). \quad (6.6)$$

We will use the same symbol  $\Pi^k$  if the projection operator is applied on vector- or matrix-valued functions component-wise.

Additionally, the  $L^2$ -orthogonal projection with respect to the metric  $g$  into the Lagrange finite elements will be needed.

**Definition 6.4.** Let  $g \in L^\infty(\Omega, \mathcal{S}^+)$  with  $g^{-1} \in L^\infty(\Omega, \mathcal{S}^+)$ . The  $L^2(\Omega, g)$ -orthogonal projector  $P_h^k : L^2(\Omega) \rightarrow \hat{\mathcal{V}}_h^k$  is defined by the property

$$(P_h^k u - u, v_h)_g = 0 \quad \text{for all } u \in L^2(\Omega), \text{ for all } v_h \in \hat{\mathcal{V}}_h^k. \quad (6.7)$$

**6.3. Basic estimates.** Later, for the main proof of Lemma 6.18 the remaining terms will be identified to be related to the distributional Christoffel symbol of first kind acting on tangential-tangential continuous matrices. Note, that  $\nu$  points inwards the elements, see Figure 1.

**Lemma 6.5** (Distributional Christoffel symbol of first kind). *Let  $g \in W_h^{1,\infty}(\Omega, \mathcal{S})$  be tangential-tangential continuous and  $\Sigma \in \mathcal{Z}(\mathcal{T}) := \{\Lambda : \Omega \rightarrow \mathbb{R}^{2 \times 2 \times 2} \mid \Lambda|_T \in C^\infty(T, \mathbb{R}^{2 \times 2 \times 2}), [\Lambda^{\nu\nu i}]_E = 0 \text{ for all } E \in \mathcal{E}, i = 1, 2\}$  be a third-order tensor where its normal-normal component is continuous. Then the distributional Christoffel symbol of first kind reads*

$$\langle \Gamma_{ijk}(g), \Sigma^{ijk} \rangle_{\mathcal{Z}(\mathcal{T})} = \sum_{T \in \mathcal{T}} \int_T \Gamma_{ijk}(g) \Sigma^{ijk} da + \int_{\partial T} \Sigma^{\nu\nu i} (g_{\nu\tau} \tau_i + \frac{1}{2} g_{\nu\nu} \nu_i) dl. \quad (6.8)$$

*Proof.* For a smooth testfunction  $\Psi$  we apply the definition of the distributional derivative and then use integration by parts element-wise

$$\begin{aligned} \langle \Gamma_{ijk}(g), \Psi^{ijk} \rangle_{C^\infty} &= - \sum_{T \in \mathcal{T}} \int_T \frac{1}{2} g_{ij} (\partial_k \Psi^{kij} + \partial_k \Psi^{ikj} - \partial_k \Psi^{ijk}) da \\ &= \sum_{T \in \mathcal{T}} \int_T \Gamma_{ijk}(g) \Psi^{ijk} da + \int_{\partial T} g_{ij} \frac{1}{2} (\Psi^{\nu ij} + \Psi^{i \nu j} - \Psi^{ij \nu}) dl \\ &= \sum_{T \in \mathcal{T}} \int_T \Gamma_{ijk}(g) \Psi^{ijk} da + \int_{\partial T} \Psi^{\nu \nu i} (g_{\nu \tau} \tau_i + \frac{1}{2} g_{\nu \nu} \nu_i) dl, \end{aligned}$$

where we split the boundary term into  $\nu\nu$ ,  $\tau\nu$ ,  $\nu\tau$ , and  $\tau\tau$  components and used that  $g$  is tangential-tangential continuous in the last step. Now, by noting that only  $\Psi^{\nu\nu i}$  has to be continuous to obtain a well-defined boundary integral we can by density extend the definition of the distributional Christoffel symbol of first kind to  $\Psi \in \mathcal{Z}(\mathcal{T})$ .  $\square$

*Remark 6.6.* With  $\Sigma \in \mathcal{Z}(\mathcal{T})$  the distributional Christoffel symbol is continuous

$$|\langle \Gamma_{ijk}(g), \Sigma^{ijk} \rangle_{\mathcal{Z}(\mathcal{T})}| \leq c \|g\|_{\Gamma} \|\Sigma\|_{\mathcal{Z}}, \quad c > 0,$$

with the norms

$$\begin{aligned} \|g\|_{\Gamma}^2 &:= \sum_{T \in \mathcal{T}} \sum_{i,j,k=1}^2 \|\Gamma_{ijk}(g)\|_{L^2(T)}^2 + \sum_{E \in \mathcal{E}} \frac{1}{h} (\|[[g_{\nu\tau}]]\|_{L^2(E)}^2 + \|[[g_{\nu\nu}]]\|_{L^2(E)}^2), \\ \|\Sigma\|_{\mathcal{Z}}^2 &:= \sum_{T \in \mathcal{T}} \sum_{i,j,k=1}^2 \|\Sigma^{ijk}\|_{L^2(T)}^2 + \sum_{E \in \mathcal{E}} h (\|\Sigma^{\nu\nu\nu}\|_{L^2(E)}^2 + \|\Sigma^{\nu\nu\tau}\|_{L^2(E)}^2). \end{aligned}$$

For only piecewise smooth  $\Sigma \in C^\infty(\mathcal{T}, \mathbb{R}^{2 \times 2 \times 2})$  the right side of (6.8) can still be evaluated. Therefore, we allow also only piecewise smooth  $\Sigma$  (losing the continuity property), see also (6.11) of the following orthogonality lemma.

We start with the following simple but crucial orthogonality properties of the Regge interpolant  $\mathcal{J}_h^{\mathcal{R}}$ .

**Lemma 6.7** (Orthogonality). *Let  $T \in \mathcal{T}$  and  $k \in \mathbb{N}_0$ . There holds for the Regge interpolation operator  $\mathcal{J}_h^{\mathcal{R},k}$  (6.5)*

$$\int_E (g - \mathcal{J}_h^{\mathcal{R},k} g)_{\tau\tau} q dl = 0 \text{ for all } q \in \mathcal{P}^k(E), \quad E \subset \partial T, \quad g \in C^0(T, \mathcal{S}), \quad (6.9)$$

$$\int_T (g - \mathcal{J}_h^{\mathcal{R},k} g) : q da = 0 \text{ for all } q \in \mathcal{P}^{k-1}(T, \mathbb{R}^{2 \times 2}), \quad g \in C^0(T, \mathcal{S}), \quad (6.10)$$

$$\langle \Gamma_{ijl}(g - \mathcal{J}_h^{\mathcal{R},k} g), \Sigma_h^{ijl} \rangle_{\mathcal{Z}(T)} = 0 \text{ for all } \Sigma_h \in \mathcal{P}^k(T, \mathbb{R}^{2 \times 2 \times 2}), \quad g \in W^{1,\infty}(T, \mathcal{S}), \quad (6.11)$$

where  $\langle \Gamma_{ijl}(g - \mathcal{J}_h^{\mathcal{R},k} g), \Sigma_h^{ijl} \rangle_{\mathcal{Z}(T)}$  denotes the expression of the distributional Christoffel symbol of first kind on a single triangle  $T$ .

*Proof.* (6.9) follows directly from (6.5a), and (6.10) from (6.5b) together by noting that the Frobenius product of a symmetric and skew-symmetric matrix is zero. For (6.11)

we abbreviate  $g_h := \mathcal{J}_h^{\mathcal{R},k} g$  and start with (6.8) on  $T \in \mathcal{T}$  and integrate by parts

$$\begin{aligned} & \langle \Gamma_{ijl}(g - g_h), \Sigma_h^{ijl} \rangle_{z(T)} \\ &= \int_T \Gamma_{ijl}(g - g_h) \Sigma_h^{ijl} da + \int_{\partial T} \Sigma_h^{\nu\nu i} ((g - g_h)_{\nu\tau} \tau_i + \frac{1}{2}(g - g_h)_{\nu\nu} \nu_i) dl \\ &= - \int_T \frac{(g - g_h)_{ij}}{2} (\partial_l \Sigma_h^{ij} + \partial_l \Sigma_h^{ilj} - \partial_l \Sigma_h^{ijl}) da - \int_{\partial T} \frac{(g - g_h)_{\tau\tau}}{2} (\Sigma_h^{\nu\tau\tau} + \Sigma_h^{\tau\nu\tau} - \Sigma_h^{\tau\tau\nu}) dl. \end{aligned}$$

By noting that the  $\Sigma_h$  terms in the first integral are in  $\mathcal{P}^{k-1}(T, \mathbb{R}^{2 \times 2})$  and in the second integral in  $\mathcal{P}^k(E)$ ,  $E \subset \partial T$ , the claim follows with (6.9)–(6.10).  $\square$

**Lemma 6.8.** *There holds for the  $L^2(\Omega, g)$ -orthogonal projector from Definition 6.4*

$$\|P_h^k u - u\|_{H^{-1}(\Omega, g)} \leq Ch \inf_{v_h \in \mathcal{V}_h^k} \|v_h - u\|_{L^2(\Omega)} \quad \text{for all } u \in L^2(\Omega), \quad (6.12a)$$

$$\|P_h^k u\|_{H^1(\Omega)} \leq C \|u\|_{H^1(\Omega)} \quad \text{for all } u \in H_0^1(\Omega), \quad (6.12b)$$

where  $C = C(\|g\|_{L^\infty}, \|g^{-1}\|_{L^\infty}, \Omega, \mathcal{T}) > 0$ .

*Proof.* Follows from the equivalence of the  $L^2$ - and  $g$ -weighted  $L^2$ -norm, see e.g. [27, Lemma 4.7].  $\square$

We will make use several times of the approximation property of the lowest-order interpolation operators.

**Lemma 6.9.** *Denote the lowest-order Regge and element-wise  $L^2$ -projection by  $\mathcal{J}_h^{\mathcal{R},0}$  and  $\Pi^0$ , respectively. Then there holds for  $p \in [1, \infty]$*

$$\|(\text{id} - \mathcal{J}_h^{\mathcal{R},0})g\|_{L^p(T)} \leq hC |g|_{W^{1,p}(T)} \quad \text{for } g \in W^{1,p}(T, \mathcal{S}), \quad (6.13a)$$

$$\|(\text{id} - \Pi^0)f\|_{L^p(T)} \leq hC |f|_{W^{1,p}(T)} \quad \text{for } f \in W^{1,p}(T), \quad (6.13b)$$

where  $C = C(T, p, \mathcal{J}_h^{\mathcal{R},0}, \Pi^0) > 0$ .

*Proof.* Scaling arguments and Bramble–Hilbert.  $\square$

**Lemma 6.10** (Scaling/Trace inequality). *Let  $T \in \mathcal{T}$  and  $E \subset \partial T$  an edge of  $T$ . There holds for all  $u \in H^1(T)$*

$$\|u\|_{L^2(E)} \leq C \sqrt{h^{-1} \|u\|_{L^2(T)}^2 + h \|\nabla u\|_{L^2(T)}^2} \quad (6.14)$$

and for all  $u \in \mathcal{P}^k(T)$

$$\|u\|_{L^2(E)} \leq Ch^{-1/2} \|u\|_{L^2(T)}, \quad (6.15)$$

$$\|\nabla u\|_{L^2(T)} \leq Ch^{-1} \|u\|_{L^2(T)}, \quad (6.16)$$

where  $C = C(T, k) > 0$ .

*Proof.* See e.g. [1, Thm 3.10] for (6.14). (6.15)–(6.16) are simple scaling arguments.  $\square$

**Lemma 6.11.** *Let  $k \in \mathbb{N}_0$ ,  $p \in [1, \infty]$ ,  $g \in W^{1,\infty}(\Omega, \mathcal{S}^+)$  with  $g^{-1} \in L^\infty(\Omega, \mathcal{S}^+)$ , and  $g_h = \mathcal{J}_h^{\mathcal{R},k} g$ . There exists an  $\bar{h} > 0$  and a constant  $C = C(\|g\|_{W^{1,\infty}}, \|g^{-1}\|_{L^\infty}, \Omega, \mathcal{T}) > 0$  such that for all  $h \leq \bar{h}$*

$$\|g_h^{-1} - g^{-1}\|_{L^p(\Omega)} \leq C \|g_h - g\|_{L^p(\Omega)}, \quad (6.17)$$

$$\|g_h\|_{W_h^{1,\infty}(\Omega)} + \|g_h^{-1}\|_{L^\infty(\Omega)} \leq C, \quad (6.18)$$

$$C^{-1}u'u \leq u'g_h(x)u \leq Cu'u \quad \text{for all } x \in \overset{\circ}{T}, \text{ for all } T \in \mathcal{T}, \text{ for all } u \in \mathbb{R}^2, \quad (6.19)$$

$$\|\sqrt{\det g_h}\|_{L^\infty(\Omega)} + \|\sqrt{\det g_h}^{-1}\|_{L^\infty(\Omega)} \leq C, \quad (6.20)$$

where  $\overset{\circ}{T}$  denotes the interior of  $T$ . For  $l \in \{0, 1\}$  there further holds

$$\|\sqrt{\det g} - \sqrt{\det g_h}\|_{W_h^{l,p}(\Omega)} \leq C \|g_h - g\|_{W_h^{l,p}(\Omega)}. \quad (6.21)$$

*Proof.* See [27, Lemma 4.5 and Lemma 4.6]. The assumptions  $\sup_{h>0} \|g_h\|_{W_h^{1,\infty}(\Omega)} < \infty$  and  $\lim_{h \rightarrow 0} \|g_h - g\|_{L^\infty(\Omega)} = 0$  are both fulfilled for the Regge interpolant  $g_h = \mathcal{J}_h^{\mathcal{R},k} g$  for  $g \in W^{1,\infty}(\Omega, \mathcal{S}^+)$ .  $\square$

Note, that the condition of a sufficiently small mesh-size  $h$  is needed to guarantee that the approximated metric  $g_h$  is positive definite.

As an elementary but important preliminary work we split the boundary terms arising in the distributional covariant curl (4.17a) into three terms by using Taylor.

**Lemma 6.12.** *Let  $T \in \mathcal{T}$ ,  $k \in \mathbb{N}_0$ ,  $g \in C^0(T, \mathcal{S}^+)$  with  $g^{-1} \in L^\infty(T, \mathcal{S}^+)$ , and  $g_h = \mathcal{J}_h^{\mathcal{R},k} g$ . Then there exist smooth functions  $f_1, f_2 \in C^\infty(\mathcal{S}^+)$  such that there holds point-wise on  $\partial T$  for sufficiently small  $h$*

$$\begin{aligned} \frac{1}{\sqrt{\det g}} - \frac{1}{\sqrt{\det g_h}} &= (g - g_h)_{\tau\tau} f_1(g_h) + \frac{2(g_h)_{\nu\tau}(g - g_h)_{\nu\tau} - (g_h)_{\tau\tau}(g - g_h)_{\nu\nu}}{2\sqrt{\det g_h}^3} \\ &\quad + \mathcal{O}(C\|g - g_h\|^2), \end{aligned} \quad (6.22)$$

$$\begin{aligned} \frac{g_{\nu\tau}}{\sqrt{\det g} g_{\tau\tau}} - \frac{(g_h)_{\nu\tau}}{\sqrt{\det g_h} (g_h)_{\tau\tau}} &= (g - g_h)_{\tau\tau} f_2(g_h) + \frac{2(g_h)_{\nu\nu}(g - g_h)_{\nu\tau} - (g_h)_{\nu\tau}(g - g_h)_{\nu\nu}}{2\sqrt{\det g_h}^3} \\ &\quad + \mathcal{O}(C\|g - g_h\|^2), \end{aligned} \quad (6.23)$$

where  $C = C(\|g_h\|_{L^\infty(\partial T)}, \|g_h^{-1}\|_{L^\infty(\partial T)}) > 0$  and  $\nu, \tau$  are the normal and tangential vectors on  $\partial T$ .

*Proof.* There holds with Taylor and (6.18)

$$\begin{aligned} \frac{1}{\sqrt{\det g}} - \frac{1}{\sqrt{\det g_h}} &= -\frac{1}{2\sqrt{\det g_h}^3} \operatorname{cof}(g_h) : (g - g_h) + \mathcal{O}(C\|g - g_h\|^2) \\ &= -\frac{(g_h)_{\nu\nu}(g - g_h)_{\tau\tau} - 2(g_h)_{\nu\tau}(g - g_h)_{\nu\tau} + (g_h)_{\tau\tau}(g - g_h)_{\nu\nu}}{2\sqrt{\det g_h}^3} + \mathcal{O}(C\|g - g_h\|^2) \\ &= (g - g_h)_{\tau\tau} \frac{-(g_h)_{\nu\nu}}{2\sqrt{\det g_h}^3} + \frac{2(g_h)_{\nu\tau}(g - g_h)_{\nu\tau} - (g_h)_{\tau\tau}(g - g_h)_{\nu\nu}}{2\sqrt{\det g_h}^3} + \mathcal{O}(C\|g - g_h\|^2), \end{aligned}$$

where  $\text{cof}(\cdot)$  denotes the cofactor matrix. Defining  $f_1(g_h) := \frac{-(g_h)_{\nu\nu}}{2\sqrt{\det g_h}} \in C^\infty(\mathcal{S}^+)$  proves the first claim. For the second identity starting with Taylor gives with (6.18)

$$\frac{g_{\nu\tau}}{\sqrt{\det g g_{\tau\tau}}} - \frac{(g_h)_{\nu\tau}}{\sqrt{\det g_h (g_h)_{\tau\tau}}} = \frac{-(g - g_h)_{\tau\tau}}{\sqrt{\det g_h (g_h)_{\tau\tau}^2}} + \frac{2 \det g_h (g - g_h)_{\nu\tau} - (g_h)_{\nu\tau} \text{cof}(g_h) : (g - g_h)}{2\sqrt{\det g_h}^3 (g_h)_{\tau\tau}} + \mathcal{O}(C\|g - g_h\|^2).$$

A simple calculation reveals

$$2 \det g_h (g - g_h)_{\nu\tau} - (g_h)_{\nu\tau} \text{cof}(g_h) : (g - g_h) = (g_h)_{\tau\tau} (2(g_h)_{\nu\nu} (g - g_h)_{\nu\tau} - (g_h)_{\nu\tau} (g - g_h)_{\nu\nu}) - (g_h)_{\nu\tau} (g_h)_{\nu\nu} (g - g_h)_{\tau\tau}$$

and thus with an appropriate smooth function  $f_2 \in C^\infty(\mathcal{S}^+)$

$$\frac{g_{\nu\tau}}{\sqrt{\det g g_{\tau\tau}}} - \frac{(g_h)_{\nu\tau}}{\sqrt{\det g_h (g_h)_{\tau\tau}}} = (g - g_h)_{\tau\tau} f_2(g_h) + \frac{2(g_h)_{\nu\nu} (g - g_h)_{\nu\tau} - (g_h)_{\nu\tau} (g - g_h)_{\nu\nu}}{2\sqrt{\det g_h}^3} + \mathcal{O}(C\|g - g_h\|^2).$$

□

**Lemma 6.13.** *Let  $k \in \mathbb{N}_0$ ,  $T \in \mathcal{T}$ , and  $E \subset \partial T$  an edge of  $T$ . Further let either  $g \in H^1(T, \mathcal{S}) \cap C^0(T, \mathcal{S})$  and  $\Psi \in W^{1,\infty}(T)$  or  $g \in C^0(T, \mathcal{S})$  and  $\Psi \in H^2(T)$ . Then there holds for  $g_h = \mathcal{J}_h^{\mathfrak{R},k} g$  and all  $q \in \mathcal{P}^k(T)$  in the first setting*

$$\left| \int_E (g_h - g)_{\tau\tau} \Psi q \, dl \right| \leq C(\|g_h - g\|_{L^2(T)} + h|g_h - g|_{H^1(T)}) \|\Psi\|_{W^{1,\infty}(T)} \|q\|_{L^2(T)}, \quad (6.24)$$

where  $C = C(T) > 0$  and in the second setting

$$\left| \int_E (g_h - g)_{\tau\tau} \Psi q \, dl \right| \leq C\|g_h - g\|_{L^\infty(T)} (\|\Psi\|_{H^1(T)} + h|\Psi|_{H^2(T)}) \|q\|_{L^2(T)}. \quad (6.25)$$

*Proof.* We start by using the  $L^2$ -projection  $\Pi^0 : L^2(T) \rightarrow \mathcal{P}^0(T)$  and orthogonality (6.9) for  $q \Pi^0 \Psi \in \mathcal{P}^k(E)$

$$\begin{aligned} \left| \int_E (g_h - g)_{\tau\tau} \Psi q \, dl \right| &\leq \left| \int_E (g_h - g)_{\tau\tau} q \Pi^0 \Psi \, dl \right| + \left| \int_E (g_h - g)_{\tau\tau} q (\text{id} - \Pi^0) \Psi \, dl \right| \\ &= \left| \int_E (g_h - g)_{\tau\tau} q (\text{id} - \Pi^0) \Psi \, dl \right| \end{aligned}$$

For inequality (6.24) and (6.25) a Hölder inequality together with the trace-, triangle inequality, and (6.13b) is applied. As they differ merely in the used norms we present only (6.24) in detail

$$\begin{aligned} \left| \int_E (g_h - g)_{\tau\tau} \Psi q \, dl \right| &\leq \|(g_h - g)_{\tau\tau}\|_{L^2(E)} \|q\|_{L^2(E)} \|(\text{id} - \Pi^0) \Psi\|_{L^\infty(E)} \\ &\leq Ch^{-1} \sqrt{\|g_h - g\|_{L^2(T)}^2 + h^2 |g_h - g|_{H^1(T)}^2} \|q\|_{L^2(T)} \|(\text{id} - \Pi^0) \Psi\|_{L^\infty(T)} \\ &\leq C(\|g_h - g\|_{L^2(T)} + h|g_h - g|_{H^1(T)}) \|q\|_{L^2(T)} \|\Psi\|_{W^{1,\infty}(T)}. \end{aligned}$$

□

**6.4. Covariant curl convergence properties.** Now we are in position to proof optimal convergence results for the covariant curl. The first one investigates the convergence of the input of the covariant curl.

**Lemma 6.14.** *For  $k \in \mathbb{N}_0$ ,  $\sigma \in H^1(\Omega, \mathcal{S}) \cap C^0(\Omega, \mathcal{S})$ ,  $\sigma_h = \mathcal{J}_h^{\mathcal{R}, k} \sigma$ ,  $g \in W^{1, \infty}(\Omega, \mathcal{S}^+)$  with  $g^{-1} \in L^\infty(\Omega, \mathcal{S}^+)$ , and  $v_h \in \mathring{W}_h^k$  there holds*

$$|\langle \text{curl}_g(\sigma - \sigma_h), Q_g v_h \rangle_{\mathring{W}_g(\mathcal{T})}| \leq C(\|\sigma - \sigma_h\|_{L^2(\Omega)} + h|\sigma - \sigma_h|_{H_h^1(\Omega)})\|v_h\|_{L^2(\Omega)}, \quad (6.26)$$

where  $C = C(\|g\|_{W^{1, \infty}}, \|g^{-1}\|_{L^\infty}, \Omega, \mathcal{T}) > 0$  and  $Q_g$  defined as in (4.16).

*Proof.* The difference reads with (4.17b)

$$\begin{aligned} \langle \text{curl}_g(\sigma - \sigma_h), Q_g v_h \rangle_{\mathring{W}_g(\mathcal{T})} &= \sum_{T \in \mathcal{T}} \int_T \frac{1}{\sqrt{\det g}} (\sigma - \sigma_h)_{ij} ([\text{rot}[v_h]]^{ij} - \varepsilon^{jk} (\Gamma_{lk}^l v_h^i - \Gamma_{lk}^i v_h^l)) da \\ &\quad - \int_{\partial T} \frac{1}{\sqrt{\det g g_{\tau\tau}}} (\sigma - \sigma_h)_{\tau\tau} v_h^i g_{i\tau} dl. \end{aligned}$$

First, we consider the volume term on one single triangle and estimate by inserting the element-wise  $L^2$ -projection  $\Pi^0 : L^2(T) \rightarrow \mathcal{P}^0(T)$  onto piecewise constants and using the orthogonality (6.10) of Regge elements

$$\begin{aligned} &\left| \int_T \frac{1}{\sqrt{\det g}} (\sigma - \sigma_h)_{ij} ([\text{rot}[v_h]]^{ij} - \varepsilon^{jk} (\Gamma_{lk}^l v_h^i - \Gamma_{lk}^i v_h^l)) da \right| \\ &\leq \left| \int_T \frac{1}{\sqrt{\det g}} (\sigma - \sigma_h)_{ij} [\text{rot}[v_h]]^{ij} da \right| + C\|\sigma - \sigma_h\|_{L^2(T)}\|v_h\|_{L^2(T)} \\ &\leq \left| \int_T (\sigma - \sigma_h)_{ij} (\Pi^0 + (\text{id} - \Pi^0)) \left( \frac{1}{\sqrt{\det g}} \right) [\text{rot}[v_h]]^{ij} da \right| + C\|\sigma - \sigma_h\|_{L^2(T)}\|v_h\|_{L^2(T)} \\ &= \left| \int_T (\sigma - \sigma_h)_{ij} (\text{id} - \Pi^0) \left( \frac{1}{\sqrt{\det g}} \right) [\text{rot}[v_h]]^{ij} da \right| + C\|\sigma - \sigma_h\|_{L^2(T)}\|v_h\|_{L^2(T)} \\ &\leq C\|\sigma - \sigma_h\|_{L^2(T)}\|v_h\|_{L^2(T)}. \end{aligned}$$

In the last step we used (6.13b) and the inverse estimate (6.16) to obtain the  $L^2$ -norm of  $v_h$ . Note, that for  $v_h \in \mathcal{P}^0(\mathcal{T}, \mathbb{R}^2)$   $[\text{rot}[v_h]] = 0$  finishing the above estimate already after the first line.

For the boundary term we can directly use inequality (6.24) for each edge  $E \subset \partial T$  with  $\Psi := \frac{1}{\sqrt{\det g g_{\tau\tau}}} g_\tau \in W^{1, \infty}(T)$  by extending the constant tangent vector  $\tau$  from  $E$  into the element  $T$

$$\begin{aligned} \left| \int_E \frac{1}{\sqrt{\det g g_{\tau\tau}}} (\sigma - \sigma_h)_{\tau\tau} v_h^i g_{i\tau} dl \right| &\leq C(\|\sigma - \sigma_h\|_{L^2(T)} + h|\sigma - \sigma_h|_{H_h^1(T)})\|v_h\|_{L^2(T)}\|\Psi\|_{W^{1, \infty}(T)} \\ &\leq C(\|\sigma - \sigma_h\|_{L^2(T)} + h|\sigma - \sigma_h|_{H_h^1(T)})\|v_h\|_{L^2(T)} \end{aligned}$$

concluding the proof by summing over all triangles, Cauchy-Schwarz, and Young inequality

$$\begin{aligned} |\langle \operatorname{curl}_g(\sigma - \sigma_h), Q_g v_h \rangle_{\dot{W}_g(\mathcal{T})}| &\leq \sum_{T \in \mathcal{T}} C(\|\sigma - \sigma_h\|_{L^2(T)} + h|\sigma - \sigma_h|_{H_h^1(T)}) \|v_h\|_{L^2(T)} \\ &\leq C \sqrt{\sum_{T \in \mathcal{T}} (\|\sigma - \sigma_h\|_{L^2(T)} + h|\sigma - \sigma_h|_{H_h^1(T)})^2} \sqrt{\sum_{T \in \mathcal{T}} \|v_h\|_{L^2(T)}^2} \\ &\leq C(\|\sigma - \sigma_h\|_{L^2(\Omega)} + h|\sigma - \sigma_h|_{H_h^1(\Omega)}) \|v_h\|_{L^2(\Omega)}. \end{aligned}$$

□

*Remark 6.15.* In the Euclidean setting,  $g = \delta$ , there exactly holds  $\langle \operatorname{curl}(\sigma - \mathcal{J}_h^{\mathcal{R},k} \sigma), v_h \rangle_{\dot{W}(\Omega)} = 0$ . Due to the involved non-linearity for a general metric  $g$  Lemma 6.14 states that still the optimal order of convergence is obtained.

The second crucial result estimates the deviation of the covariant curl when using an approximated metric tensor  $g_h = \mathcal{J}_h^{\mathcal{R},k} g$  of  $g$ . As a first step we rewrite the difference as high-order terms and the distributional Christoffel symbol of first kind applied element-wise to the difference  $g - g_h$ .

**Lemma 6.16.** *Let  $k \in \mathbb{N}_0$ ,  $\sigma_h \in \mathcal{R}_h^k$ ,  $v_h \in \dot{W}_h^k$ ,  $g \in W^{1,\infty}(\Omega, \mathcal{S}^+)$  with  $g^{-1} \in L^\infty(\Omega, \mathcal{S}^+)$ , and  $g_h = \mathcal{J}_h^{\mathcal{R},k} g$ . Then for sufficiently small  $h$*

$$\begin{aligned} \langle \operatorname{curl}_g \sigma_h, Q_g v_h \rangle_{\dot{W}_g(\mathcal{T})} - \langle \operatorname{curl}_{g_h} \sigma_h, Q_{g_h} v_h \rangle_{\dot{W}_{g_h}(\mathcal{T})} &= \langle \Gamma_{ijk}(g - g_h), \Sigma_h^{ijl} \rangle_{\mathcal{Z}(\mathcal{T})} \quad (6.27) \\ &+ \mathcal{O}(C(\|g - g_h\|_{L^\infty(\Omega)} + h\|g - g_h\|_{W_h^{1,\infty}(\Omega)}) \|\sigma_h\|_{H_h^1(\Omega)} \|v_h\|_{L^2(\Omega)}), \end{aligned}$$

$$\begin{aligned} \langle \operatorname{curl}_g \sigma_h, Q_g v_h \rangle_{\dot{W}_g(\mathcal{T})} - \langle \operatorname{curl}_{g_h} \sigma_h, Q_{g_h} v_h \rangle_{\dot{W}_{g_h}(\mathcal{T})} &= \langle \Gamma_{ijl}(g - g_h), \Sigma_h^{ijl} \rangle_{\mathcal{Z}(\mathcal{T})} \quad (6.28) \\ &+ \mathcal{O}(C(\|g - g_h\|_{L^2(\Omega)} + h\|g - g_h\|_{H_h^1(\Omega)}) \|\sigma_h\|_{W^{1,\infty}(\Omega)} \|v_h\|_{L^2(\Omega)}), \end{aligned}$$

where  $C = C(\|g\|_{W^{1,\infty}}, \|g^{-1}\|_{L^\infty}, \Omega, \mathcal{T}) > 0$ ,  $\Sigma_h^{ijl} = \frac{(\sigma_h)_{mn}}{\sqrt{\det g_h}} \varepsilon^{jm} g_h^{nl} v_h^i$ ,  $\langle \Gamma_{ijl}(g - g_h), \Sigma_h^{ijl} \rangle_{\mathcal{Z}(\mathcal{T})}$  defined as in (6.8), and  $Q_g$  as in (4.16).

*Proof.* The difference reads split into volume and boundary terms by (4.17a)

$$\begin{aligned} &\langle \operatorname{curl}_g \sigma_h, Q_g v_h \rangle_{\dot{W}_g(\mathcal{T})} - \langle \operatorname{curl}_{g_h} \sigma_h, Q_{g_h} v_h \rangle_{\dot{W}_{g_h}(\mathcal{T})} = \\ &\sum_{T \in \mathcal{T}} \int_T \frac{[\operatorname{curl}[\sigma_h]]_i v_h^i - (\sigma_h)_{ij} \varepsilon^{ki} \Gamma_{lk}^j(g) v_h^l}{\sqrt{\det g}} - \frac{[\operatorname{curl}[\sigma_h]]_i v_h^i - (\sigma_h)_{ij} \varepsilon^{ki} \Gamma_{lk}^j(g_h) v_h^l}{\sqrt{\det g_h}} da \\ &+ \int_{\partial T} \left( \frac{g_{\tau\tau}(\sigma_h)_{\nu\tau} - g_{\nu\tau}(\sigma_h)_{\tau\tau}}{\sqrt{\det g} g_{\tau\tau}} - \frac{(g_h)_{\tau\tau}(\sigma_h)_{\nu\tau} - (g_h)_{\nu\tau}(\sigma_h)_{\tau\tau}}{\sqrt{\det g_h} (g_h)_{\tau\tau}} \right) v_{h,\nu} dl. \quad (6.29) \end{aligned}$$

First, let us consider the volume term on one single triangle  $T \in \mathcal{T}$ . We add and subtract  $\sqrt{\det g_h}^{-1}$  in the first term

$$\begin{aligned} &\int_T \frac{[\operatorname{curl}[\sigma_h]]_i v_h^i - (\sigma_h)_{ij} \varepsilon^{ki} \Gamma_{lk}^j(g) v_h^l}{\sqrt{\det g}} - \frac{[\operatorname{curl}[\sigma_h]]_i v_h^i - (\sigma_h)_{ij} \varepsilon^{ki} \Gamma_{lk}^j(g_h) v_h^l}{\sqrt{\det g_h}} da = \\ &\int_T \left( \frac{1}{\sqrt{\det g}} - \frac{1}{\sqrt{\det g_h}} \right) ([\operatorname{curl}[\sigma_h]]_i v_h^i - (\sigma_h)_{ij} \varepsilon^{ki} \Gamma_{lk}^j(g) v_h^l) + \frac{(\sigma_h)_{ij} \varepsilon^{ki} (\Gamma_{lk}^j(g_h) - \Gamma_{lk}^j(g)) v_h^l}{\sqrt{\det g_h}} da \end{aligned}$$



and estimate for the former term with Hölder and (6.21)

$$\begin{aligned} & \left| \int_T \left( \frac{1}{\sqrt{\det g}} - \frac{1}{\sqrt{\det g_h}} \right) ([\text{curl}[\sigma_h]]_i v_h^i - (\sigma_h)_{ij} \varepsilon^{ki} \Gamma_{lk}^j(g) v_h^l) da \right| \\ & \leq C \left\| \frac{1}{\sqrt{\det g}} - \frac{1}{\sqrt{\det g_h}} \right\|_{L^\infty(T)} \|\sigma_h\|_{H_h^1(T)} \|v_h\|_{L^2(T)} \\ & \leq C \|g - g_h\|_{L^\infty(T)} \|\sigma_h\|_{H_h^1(T)} \|v_h\|_{L^2(T)}. \end{aligned}$$

For the latter term we use the identity  $\Gamma_{ij}^k(g) = g^{kl} \Gamma_{ijl}$ , add and subtract  $g_h^{ij}$ , and use that the Christoffel symbol of first kind is linear to obtain

$$\int_T \frac{(\sigma_h)_{ij}}{\sqrt{\det g_h}} \varepsilon^{ki} g_h^{jm} v_h^l \Gamma_{lkm}(g_h - g) + \frac{(\sigma_h)_{ij}}{\sqrt{\det g_h}} \varepsilon^{ki} (g_h^{jm} - g^{jm}) \Gamma_{lkm}(g) v_h^l da.$$

We can estimate the second term with Hölder, (6.17), and (6.20)

$$\begin{aligned} \left| \int_T \frac{(\sigma_h)_{ij} \varepsilon^{ki} (g_h^{jm} - g^{jm}) \Gamma_{lkm}(g) v_h^l}{\sqrt{\det g_h}} da \right| & \leq C \|g - g_h\|_{L^\infty(T)} \|\sigma_h\|_{L^2(T)} \|v_h\|_{L^2(T)} \\ & \leq C \|g - g_h\|_{L^\infty(T)} \|\sigma_h\|_{L^2(T)} \|v_h\|_{L^2(T)} \end{aligned}$$

and thus the final expression for the volume term is

$$\begin{aligned} & \int_T \frac{[\text{curl}[\sigma_h]]_i v_h^i - (\sigma_h)_{ij} \varepsilon^{ki} \Gamma_{lk}^j(g) v_h^l}{\sqrt{\det g}} - \frac{[\text{curl}[\sigma_h]]_i v_h^i - (\sigma_h)_{ij} \varepsilon^{ki} \Gamma_{lk}^j(g_h) v_h^l}{\sqrt{\det g_h}} da = \\ & \int_T \Sigma_h^{ijk} \Gamma_{ijk}(g_h - g) da + \mathcal{O}(C(\|g - g_h\|_{L^\infty(T)} + h\|g - g_h\|_{W_h^{1,\infty}(T)}) \|\sigma_h\|_{H_h^1(T)} \|v_h\|_{L^2(T)}). \end{aligned}$$

Next, we focus on the boundary terms. We use Lemma 6.12 to manipulate them to obtain

$$\begin{aligned} & \int_{\partial T} \frac{(g_{\tau\tau}(\sigma_h)_{\nu\tau} - g_{\nu\tau}(\sigma_h)_{\tau\tau}) v_{h,\nu}}{\sqrt{\det g} g_{\tau\tau}} - \frac{((g_h)_{\tau\tau}(\sigma_h)_{\nu\tau} - (g_h)_{\nu\tau}(\sigma_h)_{\tau\tau}) v_{h,\nu}}{\sqrt{\det g_h} (g_h)_{\tau\tau}} dl = \\ & \int_{\partial T} \left( \sigma_{h,\tau\tau} \frac{(g - g_h)_{\nu\nu} g_{h,\nu\tau} - 2g_{h,\nu\nu}(g - g_h)_{\nu\tau}}{2\sqrt{\det g_h}^3} + \sigma_{h,\nu\tau} \frac{2g_{h,\nu\tau}(g - g_h)_{\nu\tau} - g_{\tau\tau}(g - g_h)_{\nu\nu}}{2\sqrt{\det g_h}^3} \right) v_{h,\nu} \\ & + (g - g_h)_{\tau\tau} (f_1(g_h) \sigma_{h,\nu\tau} + f_2(g_h) \sigma_{h,\tau\tau}) v_{h,\nu} + \mathcal{O}(C\|g - g_h\|^2) (\sigma_{h,\nu\tau} + \sigma_{h,\tau\tau}) v_{h,\nu} dl. \end{aligned}$$

We apply inequality (6.25) for the penultimate term for each edge  $E \subset \partial T$  by choosing  $\Psi := f_1(g_h) \sigma_{h,\nu\tau} + f_2(g_h) \sigma_{h,\tau\tau} \in W^{1,\infty}(T)$  and  $q := v_{h,\nu} \in \mathcal{P}^k(T)$  (we extend the constant vectors  $\nu$  and  $\tau$  from  $E$  to  $T$ )

$$\begin{aligned} \left| \int_E (g - g_h)_{\tau\tau} \Psi v_{h,\nu} dl \right| & \leq C \|g - g_h\|_{L^\infty(T)} \|v_h\|_{L^2(T)} (\|\Psi\|_{H_h^1(T)} + h|\Psi|_{H_h^2(T)}) \\ & \leq C \|g - g_h\|_{L^\infty(T)} \|v_h\|_{L^2(T)} (\|\sigma_h\|_{H_h^1(T)} + h|\sigma_h|_{H_h^2(T)}) \\ & \leq C \|g - g_h\|_{L^\infty(T)} \|v_h\|_{L^2(T)} \|\sigma_h\|_{H_h^1(T)}, \end{aligned}$$

where we used (6.18) and (6.16). Next, we estimate the last term by using that  $\|g - g_h\|_{L^\infty(\partial T)} \leq hC$  and the trace inequality

$$\begin{aligned} \left| \int_{\partial T} \|g - g_h\|^2 (\sigma_{h,\nu\tau} + \sigma_{h,\tau\tau}) v_{h,\nu} dl \right| &\leq C \|g - g_h\|_{L^\infty(\partial T)} \|g - g_h\|_{L^\infty(\partial T)} \|\sigma_h\|_{L^2(\partial T)} \|v_h\|_{L^2(\partial T)} \\ &\leq Ch \|g - g_h\|_{L^\infty(\partial T)} \|\sigma_h\|_{L^2(\partial T)} \|v_h\|_{L^2(\partial T)} \\ &\leq C \|g - g_h\|_{L^\infty(T)} \|\sigma_h\|_{L^2(T)} \|v_h\|_{L^2(T)}. \end{aligned}$$

We claim that for the remainder holds

$$\begin{aligned} & \left( \sigma_{h,\tau\tau} \frac{(g - g_h)_{\nu\nu} g_{h,\nu\tau} - 2g_{h,\nu\nu} (g - g_h)_{\nu\tau}}{2\sqrt{\det g_h}^3} + \sigma_{h,\nu\tau} \frac{2g_{h,\nu\tau} (g - g_h)_{\nu\tau} - g_{\tau\tau} (g - g_h)_{\nu\nu}}{2\sqrt{\det g_h}^3} \right) v_{h,\nu} \\ &= \Sigma_h^{\nu\nu i} \left( (g - g_h)_{\nu\tau} \tau_i + \frac{1}{2} (g - g_h)_{\nu\nu} \nu_i \right). \end{aligned}$$

We show first that the terms involving  $(g - g_h)_{\nu\tau}$  coincide. Therefore, we use that  $\varepsilon^{\nu\nu} = 0$ ,  $\varepsilon^{\nu\tau} = -1$ , and the cofactor representation of the inverse  $g^{-1} = \frac{1}{\det g} \text{cof}(g)$

$$\begin{aligned} \Sigma_h^{\nu\nu i} (g - g_h)_{\nu\tau} \tau_i &= \frac{(\sigma_h)_{lm} \varepsilon^{\nu l} g_h^{m\tau} v_h^\nu (g - g_h)_{\nu\tau}}{\sqrt{\det g_h}} = \frac{(\sigma_h)_{\tau m} \varepsilon^{\nu\tau} g_h^{m\tau} v_h^\nu (g - g_h)_{\nu\tau}}{\sqrt{\det g_h}} \\ &= - \left( \frac{(\sigma_h)_{\tau\tau}}{\sqrt{\det g_h}} g_h^{\tau\tau} v_h^\nu (g - g_h)_{\nu\tau} + \frac{(\sigma_h)_{\tau\nu}}{\sqrt{\det g_h}} g_h^{\nu\tau} v_h^\nu (g - g_h)_{\nu\tau} \right) \\ &= - \frac{1}{\sqrt{\det g_h}^3} \left( (\sigma_h)_{\tau\tau} (g_h)_{\nu\nu} (g - g_h)_{\nu\tau} - (\sigma_h)_{\nu\tau} (g_h)_{\nu\tau} (g - g_h)_{\nu\tau} \right) v_{h,\nu}. \end{aligned}$$

Analogously, it can be shown that the remaining terms involving  $(g - g_h)_{\nu\nu}$  also match.

Summing over all triangles and comparing with (6.8) finishes the proof for (6.27). Equation (6.28), however, follows exactly the same lines swapping the  $L^2$  and  $L^\infty$  norm between  $\|g - g_h\|$  and  $\|\sigma_h\|$ .  $\square$

Next, we show by using orthogonality condition (6.11) that the distributional Christoffel term is in fact of higher-order.

**Lemma 6.17.** *Adopt the assumptions of Lemma 6.16. Then for sufficiently small  $h$*

$$|\langle \Gamma_{ijk}(g - g_h), \Sigma_h^{ijl} \rangle_{z(\mathcal{T})}| \leq C (\|g - g_h\|_{L^\infty(\Omega)} + h \|g - g_h\|_{W_h^{1,\infty}(\Omega)}) \|\sigma_h\|_{H_h^1(\Omega)} \|v_h\|_{L^2(\Omega)}. \quad (6.30)$$

*The same result holds when swapping the  $L^\infty$  and  $L^2$ -norms between  $\|g - g_h\|$  and  $\|\sigma_h\|$ .*

*Proof.* We start with the volume term of  $\langle \Gamma_{ijk}(g - g_h), \Sigma_h^{ijl} \rangle_{z(\mathcal{T})}$ . If  $k > 0$  we add and subtract the projection into the lowest-order (constant) Regge elements  $g_{h0} := \mathcal{J}_h^{\mathcal{R},0} g_h$  and  $\sigma_{h0} := \mathcal{J}_h^{\mathcal{R},0} \sigma_h$ , and define  $\Sigma_{h,0}^{ijl} = \frac{(\sigma_{h0})_{mn} \varepsilon^{jm} g_{h0}^{nl} v_h^i}{\sqrt{\det g_{h0}}} \in \mathcal{P}^k(T, \mathbb{R}^{2 \times 2 \times 2})$

$$\int_T \Sigma_h^{ijl} \Gamma_{ijl}(g_h - g) da = \int_T \Sigma_{h,0}^{ijl} \Gamma_{ijl}(g_h - g) da + \mathcal{O}(hC \|g - g_h\|_{W_h^{1,\infty}(T)} \|\sigma_h\|_{H_h^1(T)} \|v_h\|_{L^2(T)}),$$

as with  $\sum_{i,j,l=1}^2 \|\Gamma_{ijl}(g - g_h)\|_{L^\infty(T)} \leq C\|g - g_h\|_{W_h^{1,\infty}(T)}$ , Hölder, (6.17), and (6.13a)

$$\begin{aligned} & \int_T \frac{(\sigma_h)_{mn} \varepsilon^{jm} v_h^i}{\sqrt{\det g_h}} (g_h^{nl} - g_{h0}^{nl}) \Gamma_{ijl}(g_h - g) da \\ & \leq C \|\sigma_h\|_{L^2(T)} \|g_h - g_{h0}\|_{L^\infty(T)} \sum_{i,j,l=1}^2 \|\Gamma_{ijl}(g - g_h)\|_{L^\infty(T)} \|v_h\|_{L^2(T)} \\ & \leq hC \|\sigma_h\|_{L^2(T)} \|g - g_h\|_{W_h^{1,\infty}(T)} \|v_h\|_{L^2(T)} \end{aligned}$$

(and analogously for  $\sqrt{\det g_h}$ ) and

$$\begin{aligned} \int_T \frac{(\sigma_h - \sigma_{h0})_{mn} \varepsilon^{jm} g_h^{nl} v_h^i}{\sqrt{\det g_h}} \Gamma_{ijl}(g_h - g) da & \leq C \|\sigma_h - \sigma_{h0}\|_{L^2(T)} \|v_h\|_{L^2(T)} \|g_h - g\|_{W_h^{1,\infty}(T)} \\ & \leq Ch \|\sigma_h\|_{H_h^1(T)} \|v_h\|_{L^2(T)} \|g_h - g\|_{W_h^{1,\infty}(T)}. \end{aligned}$$

For the boundary term in  $\langle \Gamma_{ijk}(g - g_h), \Sigma_h^{ijl} \rangle_{z(\mathcal{T})}$  we use the lowest-order Regge interpolation operator, analogously to the volume terms, to obtain

$$\begin{aligned} \int_{\partial T} \Sigma_h^{\nu\nu i} ((g - g_h)_{\nu\tau} \tau_i + \frac{1}{2}(g - g_h)_{\nu\nu} \nu_i) dl & = \int_{\partial T} \Sigma_{h,0}^{\nu\nu i} ((g - g_h)_{\nu\tau} \tau_i + \frac{1}{2}(g - g_h)_{\nu\nu} \nu_i) dl \\ & + \mathcal{O}(C(\|g - g_h\|_{L^\infty(T)} + h\|g - g_h\|_{W_h^{1,\infty}(T)}) \|\sigma_h\|_{H_h^1(T)} \|v_h\|_{L^2(T)}). \end{aligned}$$

Thus, by using orthogonality property (6.11) with  $\Sigma_{h,0}^{ijl} \in \mathcal{P}^k(T, \mathbb{R}^{2 \times 2 \times 2})$  we deduce

$$\begin{aligned} \langle \Gamma_{ijl}(g - g_h), \Sigma_h^{ijl} \rangle_{z(\mathcal{T})} & = \langle \Gamma_{ijl}(g - g_h), \Sigma_{h,0}^{ijl} \rangle_{z(\mathcal{T})} \\ & + \mathcal{O}(C(\|g - g_h\|_{L^\infty(T)} + h\|g - g_h\|_{W_h^{1,\infty}(T)}) \|\sigma_h\|_{H_h^1(T)} \|v_h\|_{L^2(T)}) \\ & = \mathcal{O}(C(\|g - g_h\|_{L^\infty(T)} + h\|g - g_h\|_{W_h^{1,\infty}(T)}) \|\sigma_h\|_{H_h^1(T)} \|v_h\|_{L^2(T)}) \end{aligned}$$

finishing the proof.

The proof for swapped norms follows exactly the same lines.  $\square$

Combining Lemma 6.16 and Lemma 6.17 now directly yields the optimal convergence for the distributional covariant curl.

**Lemma 6.18.** *Let  $k \in \mathbb{N}_0$ ,  $\sigma_h \in \mathcal{R}_h^k$ ,  $v_h \in \mathring{W}_h^k$ ,  $g \in W^{1,\infty}(\Omega, \mathcal{S}^+)$  with  $g^{-1} \in L^\infty(\Omega, \mathcal{S}^+)$ , and  $g_h = \mathcal{J}_h^{\mathcal{R},k} g$ . Then for sufficiently small  $h$*

$$\begin{aligned} |\langle \text{curl}_g \sigma_h, Q_g v_h \rangle_{\mathring{W}_g(\mathcal{T})} - \langle \text{curl}_{g_h} \sigma_h, Q_{g_h} v_h \rangle_{\mathring{W}_{g_h}(\mathcal{T})}| & \leq \\ & C(\|g - g_h\|_{L^\infty(\Omega)} + h\|g - g_h\|_{W_h^{1,\infty}(\Omega)}) \|\sigma_h\|_{H_h^1(\Omega)} \|v_h\|_{L^2(\Omega)}, \end{aligned} \quad (6.31)$$

$$\begin{aligned} |\langle \text{curl}_g \sigma_h, Q_g v_h \rangle_{\mathring{W}_g(\mathcal{T})} - \langle \text{curl}_{g_h} \sigma_h, Q_{g_h} v_h \rangle_{\mathring{W}_{g_h}(\mathcal{T})}| & \leq \\ & C(\|g - g_h\|_{L^2(\Omega)} + h\|g - g_h\|_{H_h^1(\Omega)}) \|\sigma_h\|_{W_h^{1,\infty}(\Omega)} \|v_h\|_{L^2(\Omega)}, \end{aligned} \quad (6.32)$$

where  $C = C(\|g\|_{W^{1,\infty}}, \|g^{-1}\|_{L^\infty}, \Omega, \mathcal{T}) > 0$  and  $Q_g$  defined as in (4.16).

*Remark 6.19.* The fundamental Lemma 6.18 states that when approximating the covariant  $\text{curl}_g$  by  $\text{curl}_{g_h}$  on a triangulation  $\mathcal{T}$  we obtain that the covariant differential operator converges optimally to its smooth pendant.

**6.5. Inc estimates.** After the distributional covariant curl has been analyzed we can turn to the incompatibility operator, where we obtain the same results, in a weaker norm, as for the curl.

**Corollary 6.20.** *Let  $k \in \mathbb{N}_0$ ,  $\sigma \in H^1(\Omega, \mathcal{S}) \cap C^0(\Omega, \mathcal{S})$ ,  $\sigma_h = \mathcal{J}_h^{\mathcal{R},k} \sigma$ ,  $g \in W^{1,\infty}(\Omega, \mathcal{S}^+)$  with  $g^{-1} \in L^\infty(\Omega, \mathcal{S}^+)$ , and  $u_h \in \mathring{V}_h^{k+1}$ . Then*

$$|\langle \text{inc}_g(\sigma - \sigma_h), u_h \rangle_{\mathring{V}(\mathcal{T})}| \leq C(\|\sigma - \sigma_h\|_{L^2(\Omega)} + h\|\sigma - \sigma_h\|_{H_h^1(\Omega)})\|\nabla u_h\|_{L^2(\Omega)}, \quad (6.33)$$

where  $C = C(\|g\|_{W^{1,\infty}}, \|g^{-1}\|_{L^\infty}, \Omega, \mathcal{T}) > 0$ .

*Proof.* By noting that  $\langle \text{inc}_g(\sigma - \sigma_h), u_h \rangle_{\mathring{V}(\mathcal{T})} = \langle \text{curl}_g(\sigma - \sigma_h), \text{rot } u_h \rangle_{\mathring{V}(\mathcal{T})}$  and  $\text{rot } u_h \in \mathring{W}_h^k$  we can directly use Lemma 6.14 with  $v_h := \text{rot } u_h$ .  $\square$

**Corollary 6.21.** *Let  $k \in \mathbb{N}_0$ ,  $\sigma_h \in \mathcal{R}_h^k$ ,  $u_h \in \mathring{V}_h^{k+1}$ ,  $g \in W^{1,\infty}(\Omega, \mathcal{S}^+)$  with  $g^{-1} \in L^\infty(\Omega, \mathcal{S}^+)$ , and  $g_h = \mathcal{J}_h^{\mathcal{R},k} g$ . Then for sufficiently small  $h$*

$$\begin{aligned} & |\langle \text{inc}_g \sigma_h, u_h \rangle_{\mathring{V}(\mathcal{T})} - \langle \text{inc}_{g_h} \sigma_h, u_h \rangle_{\mathring{V}(\mathcal{T})}| \\ & \leq C(\|g - g_h\|_{L^\infty(\Omega)} + h\|g - g_h\|_{W_h^{1,\infty}(\Omega)})\|\sigma_h\|_{H_h^1(\Omega)}\|\nabla u_h\|_{L^2(\Omega)}, \end{aligned} \quad (6.34)$$

$$\begin{aligned} & |\langle \text{inc}_g \sigma_h, u_h \rangle_{\mathring{V}(\mathcal{T})} - \langle \text{inc}_{g_h} \sigma_h, u_h \rangle_{\mathring{V}(\mathcal{T})}| \\ & \leq C(\|g - g_h\|_{L^2(\Omega)} + h\|g - g_h\|_{H_h^1(\Omega)})\|\sigma_h\|_{W_h^{1,\infty}(\Omega)}\|\nabla u_h\|_{L^2(\Omega)}, \end{aligned} \quad (6.35)$$

where  $C = C(\|g\|_{W^{1,\infty}}, \|g^{-1}\|_{L^\infty}, \Omega, \mathcal{T}) > 0$ .

*Proof.* Again, by choosing  $v_h := \text{rot } u_h$  we can directly apply Lemma 6.18.  $\square$

*Remark 6.22.* In the Euclidean setting there exactly holds  $\langle \text{inc}(\sigma - \mathcal{J}_h^{\mathcal{R},k} \sigma), u_h \rangle_{\mathring{V}(\mathcal{T})} = 0$  for all  $u_h \in \mathring{V}_h^{k+1}$ . Corollary 6.20 shows that for the nonlinear differential operator  $\text{inc}_g$  still optimal convergence is achieved. In Corollary 6.21 we obtain e.g. in the lowest order case,  $k = 0$ , linear convergence, although a second order differential operator relying on derivatives of the metric tensor is applied. The idea of splitting the terms into a high-order and a sub-optimal one, which then turns out to be zero, has been used in [46]. Therein, the convergence of the surface divdiv operator on an approximated triangulation is proven in an extrinsic manner to converge optimally.

*Remark 6.23.* Our analysis estimates the same terms as [27] by using the operator  $\text{inc}_g$  instead of  $\text{div}_g \text{div}_g S_g \sigma$ . In [27] the volume and jump terms are estimated independently by each other as a triangle inequality is applied first. We keep the volume and jump terms together and split it into terms of appropriate convergence order and the distributional Christoffel symbol of first kind as a sub-optimal remainder. This remainder, however, is zero due to the specific orthogonality condition (6.11).

**6.6. Proof for lifted curvature.** Before we come to the proof of our main theorem, we need two additional lemma involving the Gauss curvature.

**Lemma 6.24.** *Let  $k \in \mathbb{N}_0$ ,  $g_h = \mathcal{J}_h^{\mathcal{R},k} \bar{g}$ ,  $u \in H_0^1(\Omega)$ , and  $u_h = P_h^{k+1} u$  be the  $L^2(\Omega, \bar{g})$ -orthogonal projection (6.7). Then*

$$|(K_h(g_h) - K(\bar{g}), u - u_h)_{\bar{g}}| \leq hC\|u\|_{H^1(\Omega)} \inf_{v_h \in \mathring{V}_h^{k+1}} \|K(\bar{g}) - v_h\|_{L^2(\Omega)}, \quad (6.36)$$

where  $C = C(\|\bar{g}\|_{L^\infty}, \|\bar{g}^{-1}\|_{L^\infty}, \Omega, \mathcal{F})$ .

*Proof.* The claim follows with the property of the  $L^2$ -projection, (6.12a), and (6.12b)

$$\begin{aligned} |(K_h(g_h) - K(\bar{g}), u - u_h)_{\bar{g}}| &= |(P_h^{k+1}K(\bar{g}) - K(\bar{g}), u - u_h)_{\bar{g}}| \\ &\leq \|P_h^{k+1}K(\bar{g}) - K(\bar{g})\|_{H^{-1}(\Omega, \bar{g})} \|u - u_h\|_{H^1(\Omega)} \\ &\leq Ch \|u\|_{H^1(\Omega)} \inf_{v_h \in \mathring{V}_h^{k+1}} \|K(\bar{g}) - v_h\|_{L^2(\Omega)}. \end{aligned}$$

□

**Lemma 6.25.** *Let  $k \in \mathbb{N}_0$ ,  $g_h = \mathcal{J}_h^{\mathcal{R}, k} \bar{g}$ , and  $u_h \in \mathring{V}_h^{k+1}$ . Then*

$$|(K_h(g_h), u_h)_{\bar{g}} - (K_h(g_h), u_h)_{g_h}| \leq C \|\bar{g} - g_h\|_{L^\infty(\Omega)} \|K_h(g_h)\|_{L^2(\Omega)} \|u_h\|_{L^2(\Omega)}, \quad (6.37)$$

where  $C = C(\|\bar{g}\|_{W^{1,\infty}}, \|\bar{g}^{-1}\|_{L^\infty}, \Omega, \mathcal{F})$ .

*Proof.* With (6.21) we immediately obtain

$$\begin{aligned} (K_h(g_h), u_h)_{\bar{g}} - (K_h(g_h), u_h)_{g_h} &= \int_{\Omega} K_h(g_h) u_h (\sqrt{\det(\bar{g})} - \sqrt{\det(g_h)}) da \\ &\leq \|K_h(g_h)\|_{L^2(\Omega)} \|u_h\|_{L^2(\Omega)} \|\sqrt{\det(\bar{g})} - \sqrt{\det(g_h)}\|_{L^\infty(\Omega)} \\ &\leq C \|K_h(g_h)\|_{L^2(\Omega)} \|u_h\|_{L^2(\Omega)} \|\bar{g} - g_h\|_{L^\infty(\Omega)}. \end{aligned}$$

□

Now, we finally come to the proof of the theorem.

*Proof of Theorem 6.1.* The general structure of the proof follows [27].

Let  $u \in H_0^1(\Omega)$ . We start by splitting into different terms with  $u_h := P_h^{k+1}u \in \mathring{V}_h^{k+1}$  the  $L^2(\Omega, \bar{g})$ -orthogonal projector

$$\begin{aligned} (K_h(g_h) - K(\bar{g}), u)_{\bar{g}} &= (K_h(g_h) - K(\bar{g}), u - u_h + u_h)_{\bar{g}} = \\ &= (K_h(g_h) - K(\bar{g}), u - u_h)_{\bar{g}} + (K_h(g_h) - K(\bar{g}), u_h)_{\bar{g}-g_h+g_h} = \\ &= (K_h(g_h), u_h)_{g_h} - (K(\bar{g}), u_h)_{\bar{g}} + (K_h(g_h) - K(\bar{g}), u - u_h)_{\bar{g}} + (K_h(g_h), u_h)_{\bar{g}} - (K_h(g_h), u_h)_{g_h}. \end{aligned} \quad (6.38)$$

With the notation  $G(t) := \delta + t(\bar{g} - \delta)$  and  $G_h(t) := \delta + t(g_h - \delta)$  we can rewrite the first two terms in (6.38) as

$$\begin{aligned} (K_h(g_h), u_h)_{g_h} - (K(\bar{g}), u_h)_{\bar{g}} &= \frac{1}{2} \int_0^1 b_h(G_h(t), g_h - \delta, u_h) - b_h(G(t), \bar{g} - \delta, u_h) dt \\ &= \frac{1}{2} \int_0^1 b_h(G_h(t), g_h - \delta, u_h) - b_h(G(t), g_h - \delta, u_h) dt + \frac{1}{2} \int_0^1 b_h(G(t), g_h - \bar{g}, u_h) dt. \end{aligned}$$

By using  $g = G(t)$  and  $\sigma_h = g_h - \delta \in \mathcal{R}_h^k$  for (6.34) in Corollary 6.21 and  $g = G(t)$  and  $\sigma = \bar{g}$  for (6.33) in Corollary 6.20 we can estimate together with (6.18)

$$\begin{aligned} |(K_h(g_h), u_h)_{g_h} - (K(\bar{g}), u_h)_{\bar{g}}| &\leq C(\|\bar{g} - g_h\|_{L^\infty} + h\|\bar{g} - g_h\|_{W_h^{1,\infty}}) \|g_h - \delta\|_{H_h^1} \|\nabla u_h\|_{L^2} \\ &\leq C(\|\bar{g} - g_h\|_{L^\infty} + h\|\bar{g} - g_h\|_{W_h^{1,\infty}}) \|\nabla u_h\|_{L^2}. \end{aligned} \quad (6.39)$$

The next term in (6.38) can be directly estimated with Lemma 6.24

$$|(K_h(g_h) - K(\bar{g}), u - u_h)_{\bar{g}}| \leq hC \|u\|_{H^1(\Omega)} \inf_{v_h \in \check{V}_h^{k+1}} \|K(\bar{g}) - v_h\|_{L^2(\Omega)}. \quad (6.40)$$

For the last difference in (6.38) we use Lemma 6.25

$$|(K_h(g_h), u_h)_{\bar{g}} - (K_h(g_h), u_h)_{g_h}| \leq C \|\bar{g} - g_h\|_{L^\infty(\Omega)} \|K_h(g_h)\|_{L^2(\Omega)} \|u_h\|_{L^2(\Omega)}.$$

A direct computation shows that  $\|K_h(g_h)\|_{L^2(\Omega)}$  is bounded by using (6.19), (6.39), and (6.16)

$$\begin{aligned} \|K_h(g_h)\|_{L^2(\Omega)}^2 &\leq C \|K_h(g_h)\|_{L^2(\Omega, g_h)}^2 \\ &\leq C (|(K_h(g_h), K_h(g_h))_{g_h} - (K(\bar{g}), K_h(g_h))_{\bar{g}}| + |(K(\bar{g}), K_h(g_h))_{\bar{g}}|) \\ &\leq C \left( (\|\bar{g} - g_h\|_{L^\infty(\Omega)} + h \|\bar{g} - g_h\|_{W_h^{1,\infty}(\Omega)}) \|\nabla K_h(g_h)\|_{L^2(\Omega)} + \|K(\bar{g})\|_{L^2(\Omega)} \|K_h(g_h)\|_{L^2(\Omega)} \right) \\ &\leq C \left( \frac{1}{h} (\|\bar{g} - g_h\|_{L^\infty(\Omega)} + h \|\bar{g} - g_h\|_{W_h^{1,\infty}(\Omega)}) + \|K(\bar{g})\|_{L^2(\Omega)} \right) \|K_h(g_h)\|_{L^2(\Omega)} \\ &\leq C \|K_h(g_h)\|_{L^2(\Omega)}, \end{aligned}$$

as  $\frac{1}{h} (\|\bar{g} - g_h\|_{L^\infty(\Omega)} + h \|\bar{g} - g_h\|_{W_h^{1,\infty}(\Omega)}) \leq C$  is bounded by using  $\|\bar{g} - g_h\|_{L^\infty(\Omega)} \leq Ch$  and  $\|K_h(g_h)\|_{L^2(\Omega)} \leq C$ .

All together, we have with (6.12b)

$$\begin{aligned} |(K_h(g_h) - K(\bar{g}), u)_{\bar{g}}| &\leq C ((\|\bar{g} - g_h\|_{L^\infty(\Omega)} + h \|\bar{g} - g_h\|_{W_h^{1,\infty}(\Omega)}) \|u_h\|_{H^1(\Omega)} + h \|u\|_{H^1(\Omega)} \inf_{v_h \in \check{V}_h^{k+1}} \|K(\bar{g}) - v_h\|_{L^2(\Omega)}) \\ &\leq C (\|\bar{g} - g_h\|_{L^\infty(\Omega)} + h \|\bar{g} - g_h\|_{W_h^{1,\infty}(\Omega)} + h \inf_{v_h \in \check{V}_h^{k+1}} \|K(\bar{g}) - v_h\|_{L^2(\Omega)}) \|u\|_{H^1(\Omega)} \end{aligned}$$

and thus

$$\begin{aligned} \|K_h(g_h) - K(\bar{g})\|_{H^{-1}(\Omega, \bar{g})} &= \sup_{u \in H_0^1(\Omega)} \frac{(K_h(g_h) - K(\bar{g}), u)_{\bar{g}}}{\|u\|_{H^1(\Omega)}} \\ &\leq C (\|\bar{g} - g_h\|_{L^\infty(\Omega)} + h \|\bar{g} - g_h\|_{W_h^{1,\infty}(\Omega)} + h \inf_{v_h \in \check{V}_h^{k+1}} \|K(\bar{g}) - v_h\|_{L^2(\Omega)}). \end{aligned}$$

By standard interpolation theory we directly deduce the convergence rate

$$\|K_h(g_h) - K(\bar{g})\|_{H^{-1}(\Omega, g)} \leq Ch^{k+1} (\|\bar{g}\|_{W^{k+1,\infty}(\Omega)} + |K(\bar{g})|_{H^k}).$$

□

We obtain also optimal estimates in stronger norms. The proof follows exactly the same lines as in [27], where the error is compared to the  $H^{-1}$ -error. Due to the improved estimate derived in Theorem 6.1 we deduce one extra order of convergence.

**Corollary 6.26.** *Adopt the assumptions from Theorem 6.1. Then there holds for  $0 \leq l \leq k$*

$$\begin{aligned} \|K_h(g_h) - K(\bar{g})\|_{L^2} &\leq C(h^{-1}\|\bar{g} - g_h\|_{L^\infty} + \|\bar{g} - g_h\|_{W_h^{1,\infty}} + \inf_{v_h \in \mathring{V}_h^{k+1}} \|K(\bar{g}) - v_h\|_{L^2}) \\ &\leq Ch^k(\|\bar{g}\|_{W^{k+1,\infty}} + |K(\bar{g})|_{H^k}), \end{aligned} \quad (6.41)$$

$$\begin{aligned} |K_h(g_h) - K(\bar{g})|_{H_h^l} &\leq Ch^{-l}(h^{-1}\|\bar{g} - g_h\|_{L^\infty} + \|\bar{g} - g_h\|_{W_h^{1,\infty}} + \inf_{v_h \in \mathring{V}_h^{k+1}} \|K(\bar{g}) - v_h\|_{L^2}) \\ &\leq Ch^{k-l}(\|\bar{g}\|_{W^{k+1,\infty}} + |K(\bar{g})|_{H^k}). \end{aligned} \quad (6.42)$$

**6.7. Proof lifted connection 1-form.** The proof for the optimal convergence of the lifted connection 1-form follows the same ideas as for the Gauss curvature.

*Proof of Theorem 6.2.* For  $v \in L^2(\Omega, \mathbb{R}^2)$  we denote with  $v_h = P_h^{\mathcal{W},k}v \in \mathring{W}_h^k$  the  $L^2$ -orthogonal projector of  $v$  defined analogously to (6.7), with  $g = \delta$ , and compute

$$\begin{aligned} \|\varpi_h(g_h) - \varpi(\bar{g})\|_{L^2(\Omega)} &= \sup_{v \in L^2(\Omega, \mathbb{R}^2)} \frac{\int_\Omega (\varpi_h(g_h) - \varpi(\bar{g}))(v + v_h - v_h) da}{\|v\|_{L^2(\Omega)}} \\ &= \sup_{v \in L^2(\Omega, \mathbb{R}^2)} \frac{\int_\Omega (\varpi_h(g_h) - \varpi(\bar{g}))(v - v_h) da + \int_\Omega (\varpi_h(g_h) - \varpi(\bar{g}))v_h da}{\|v\|_{L^2(\Omega)}}. \end{aligned}$$

Due to the property of the  $L^2$ -orthogonal projector we obtain

$$\begin{aligned} \int_\Omega (\varpi_h(g_h) - \varpi(\bar{g}))(v - v_h) da &= \int_\Omega (P_h^{\mathcal{W},k}\varpi(\bar{g}) - \varpi(\bar{g}))(v - v_h) da \\ &\leq C\|v\|_{L^2(\Omega)}\|P_h^{\mathcal{W},k}\varpi(\bar{g}) - \varpi(\bar{g})\|_{L^2(\Omega)} \\ &\leq C\|v\|_{L^2(\Omega)} \inf_{w_h \in \mathring{W}_h^k} \|\varpi(\bar{g}) - w_h\|_{L^2(\Omega)}. \end{aligned}$$

With the notation  $G(t) := \delta + t(\bar{g} - \delta)$  and  $G_h(t) := \delta + t(g_h - \delta)$  we can rewrite

$$\begin{aligned} \int_\Omega (\varpi_h(g_h) - \varpi(\bar{g}))v_h da &= \frac{1}{2} \int_0^1 c_h(G_h(t), g_h - \delta, v_h) - c_h(G(t), g_h - \delta, v_h) dt \\ &\quad + \frac{1}{2} \int_0^1 c_h(G(t), g_h - \bar{g}, v_h) dt \end{aligned}$$

and estimate with Lemma 6.14, Lemma 6.18, and (6.18)

$$\begin{aligned} \left| \int_\Omega (\varpi_h(g_h) - \varpi(\bar{g}))v_h da \right| &\leq C(\|\bar{g} - g_h\|_{L^2(\Omega)} + h\|\bar{g} - g_h\|_{H_h^1(\Omega)})\|g_h - \delta\|_{W_h^{1,\infty}(\Omega)}\|v_h\|_{L^2(\Omega)} \\ &\leq C(\|\bar{g} - g_h\|_{L^2(\Omega)} + h\|\bar{g} - g_h\|_{H_h^1(\Omega)})\|v_h\|_{L^2(\Omega)}. \end{aligned}$$




FIGURE 2. Left: Embedded surface, color indicates to  $z$ -component. Right: Exact Gauss curvature as graph over the domain  $\Omega$ .

Thus, we obtain with  $\|v_h\|_{L^2(\Omega)} \leq C\|v\|_{L^2(\Omega)}$

$$\begin{aligned} & \|\varpi_h(g_h) - \varpi(\bar{g})\|_{L^2(\Omega)} \\ & \leq C \sup_{v \in L^2(\Omega, \mathbb{R}^2)} \frac{(\|\bar{g} - g_h\|_{L^2(\Omega)} + h\|\bar{g} - g_h\|_{H_h^1(\Omega)} + \inf_{w_h \in \dot{W}_h^k} \|\varpi(\bar{g}) - w_h\|_{L^2(\Omega)})\|v\|_{L^2(\Omega)}}{\|v\|_{L^2(\Omega)}} \\ & = C(\|\bar{g} - g_h\|_{L^2(\Omega)} + h\|\bar{g} - g_h\|_{H_h^1(\Omega)} + \inf_{w_h \in \dot{W}_h^k} \|\varpi(\bar{g}) - w_h\|_{L^2(\Omega)}) \end{aligned}$$

and further

$$\|\varpi_h(g_h) - \varpi(\bar{g})\|_{L^2(\Omega)} \leq Ch^{k+1}(\|\bar{g}\|_{H^{k+1}(\Omega)} + |\varpi(\bar{g})|_{H^k}).$$

□

## 7. NUMERICAL EXAMPLES

In this section we confirm by numerical examples that the theoretical convergence rates from Theorem 6.1 and Theorem 6.2 are sharp.

All experiments were performed in the open source finite element software NGSolve<sup>1</sup> [41, 42], where the Regge elements are available.

**7.1. Curvature approximation.** We consider the numerical example proposed in [27], where on the square  $\Omega = (-1, 1) \times (-1, 1)$  the smooth Riemannian metric tensor

$$\bar{g}(x, y) := \begin{pmatrix} 1 + (\partial_x f)^2 & \partial_x f \partial_y f \\ \partial_x f \partial_y f & 1 + (\partial_y f)^2 \end{pmatrix}$$

with  $f(x, y) := \frac{1}{2}(x^2 + y^2) - \frac{1}{12}(x^4 + y^4)$  is defined. This metric corresponds to the surface induced by the embedding  $(x, y) \mapsto (x, y, f(x, y))$  and its exact Gauss curvature is given by

$$K(\bar{g})(x, y) = \frac{81(1 - x^2)(1 - y^2)}{(9 + x^2(x^2 - 3))^2 + y^2(y^2 - 3)^2}.$$

The embedded surface and Gauss curvature are depicted in Figure 2.

To test also the case of non-homogeneous Dirichlet and Neumann boundary conditions we consider only one quarter  $\Omega = (0, 1) \times (0, 1)$  and define the right and bottom

<sup>1</sup>www.ngsolve.org






FIGURE 3. Left: Domain with Dirichlet and Neumann boundaries. Middle and right: perturbed unstructured triangular mesh grids.

boundaries as Dirichlet and the remaining parts as Neumann boundary. To avoid possible super-convergence properties due to a structured grid, we perturb all internal points of the triangular mesh by a uniform distribution in the range  $[-\frac{h}{4}, \frac{h}{4}]$ ,  $h$  denoting the maximal mesh-size of the originally generate mesh. The geodesic curvature on the left boundary is exactly zero, whereas on the top boundary we compute

$$\kappa_g(\bar{g})|_{\Gamma_{\text{left}}} = 0, \quad \kappa_g(\bar{g})|_{\Gamma_{\text{top}}} = \frac{-27(x^2 - 1)y(y^2 - 3)}{(x^2(x^2 - 3)^2 + 9)^{3/2} \sqrt{x^2(x^2 - 3)^2 + y^2(y^2 - 3)^2 + 9}}.$$

The vertex expressions  $K_V$  at the vertices of the Neumann boundary can directly be computed by measuring the angle  $\arccos(\frac{\tau'_1 \bar{g} \tau_2}{\sqrt{\tau'_1 \bar{g} \tau_1} \sqrt{\tau'_2 \bar{g} \tau_2}})$ .

To obtain the optimal convergence rates the metric tensor  $\bar{g}$  has to be interpolated into the Regge finite element space corresponding to the interpolation operator  $\mathcal{J}_h^{\mathcal{R},k}$ ,  $g_h = \mathcal{J}_h^{\mathcal{R},k} \bar{g}$ . Therefore, e.g., the moments on the edges have to coincide exactly,  $\int_E (g_h - \bar{g})_{\tau\tau} dl = 0$ . Thus, we use a high enough integration rule for interpolating  $\bar{g}$  minimizing the numerical deviation. Otherwise sub-optimal results or even reduced convergence rates might be observed numerically.

We compute the  $L^2$ -norm  $\|K(\bar{g}) - K_h(g_h)\|_{L^2(\Omega)}$  and the  $H^{-1}$ -norm, where we used that for  $w \in H_0^1(\Omega)$  solving the Poisson problem  $-\Delta w = K(\bar{g}) - K_h(g_h)$  there holds

$$\|K(\bar{g}) - K_h(g_h)\|_{H^{-1}(\Omega)} = \|w\|_{H^1(\Omega)}. \quad (7.1)$$

To minimize the discretization error by approximating  $w$  by  $w_h$  we increase the used polynomial degree by two compared to  $K_h(g_h)$ , i.e.,  $w_h \in \mathcal{V}_h^{k+3}$  for  $K_h(g_h) \in \mathcal{V}_h^{k+1}$ .

We start by approximating  $\bar{g}$  by lowest order piecewise constant Regge elements  $g_h \in \mathcal{R}_h^0$ . As shown in Figure 4 left we do not obtain convergence in the  $L^2$ -norm, however, linear convergence in the weaker  $H^{-1}$ -norm, which is in common with Theorem 6.1. When increasing the approximation order of Regge elements to linear and quadratic polynomials we observe the appropriate increase of the convergence rates, Figure 4




FIGURE 4. Convergence of Gauss curvature with respect to number of degrees of freedom (ndof) in different norms for Regge elements  $g_h \in \mathcal{R}_h^k$  of order  $k = 0, 1, 2$ .




FIGURE 5. Approximated Gauss curvature with Regge elements of order  $k = 0, 1, 2$ .

middle and right, confirming that the results stated in Theorem 6.1 and Corollary 6.26 are sharp. In Figure 5 the approximated Gauss curvature is shown.

**7.2. Connection 1-form approximation.** For testing the convergence of the connection 1-form we consider the same metric tensor  $\bar{g}$  as before, but on the whole domain  $\Omega = (-1, 1) \times (-1, 1)$ . The exact connection 1-form  $\varpi$  depicted in Figure 6 can be computed by

$$\varpi(\bar{g})(x, y) = -\frac{1}{2} \begin{pmatrix} (e^2)' \bar{g} \nabla_{E_1} e^1 - (e^1)' \bar{g} \nabla_{E_1} e^2 \\ (e^2)' \bar{g} \nabla_{E_2} e^1 - (e^1)' \bar{g} \nabla_{E_2} e^2 \end{pmatrix} = \quad (7.2)$$

$$\begin{pmatrix} -\frac{(9x^2-9)y(y^2-3)\sqrt{y^6-6y^4+9y^2+9+x^6-6x^4+9x^2+(3y^9-27y^7+81y^5-54y^3-81y)x^2-3y^9+27y^7-81y^5+54y^3+81y+(3x^8-21x^6+45x^4-27x^2)y(y^2-3)}}{\sqrt{y^6-6y^4+9y^2+9+x^2(x^2-3)^2}(y^6-6y^4+9y^2+18+x^6-6x^4+9x^2)\sqrt{y^6-6y^4+9y^2+9+x^6-6x^4+9x^2+6y^6-36y^4+54y^2+54+6x^6-36x^4+54x^2}} \\ \frac{(3y^6+3x^6y^2-21y^6-3x^6-18x^4y^2+45y^4+18x^4+27x^2y^2+9\sqrt{y^6-6y^4+9y^2+9+x^6-6x^4+9x^2}y^2-27x^2-27-9\sqrt{y^6-6y^4+9y^2+9+x^6-6x^4+9x^2})(x^2-3)x}{\sqrt{y^6-6y^4+9y^2+9+x^2(x^2-3)^2}(y^6-6y^4+9y^2+18+x^6-6x^4+9x^2)\sqrt{y^6-6y^4+9y^2+9+x^6-6x^4+9x^2+6y^6-36y^4+54y^2+54+6x^6-36x^4+54x^2}} \end{pmatrix},$$

where  $E_i \in \mathbb{R}^2$  are the Euclidean basis vectors,  $e^i := \bar{g}^{-\frac{1}{2}} E_i$  the  $\bar{g}$ -orthonormal basis in corresponding to the integral representation (5.12), compare Lemma 5.4, and the covariant derivative  $(\bar{\nabla}_X Y)^i = ((\bar{\nabla} X)Y)^i + \Gamma_{jk}^i X^j Y^k$ ,  $\Gamma_{ij}^k$  the Christoffel symbol of second kind with respect to  $\bar{g}$ . Note that there holds  $\text{curl } \varpi(\bar{g}) = \sqrt{\det \bar{g}} K(\bar{g})$ .

We use the integral representation (5.12) and approximate the parameter integral with a Gauss quadrature of order 20 to alleviate model errors. Note, that the exact 1-form has zero normal component at the boundary,  $\varpi(\bar{g}) \cdot \nu = 0$  on  $\partial\Omega$ .




FIGURE 6. Exact connection 1-form  $\varpi$ . Left:  $\|\varpi\|_2$ , middle:  $x$ -component, right:  $y$ -component.




FIGURE 7.  $L^2$ -convergence of connection 1-form error with BDM and RT elements for polynomial orders  $k = 1, 2$ .




FIGURE 8. Approximated connection 1-form. Left:  $\mathcal{W}_h^1$ , middle:  $\mathcal{W}_h^2$ , right: second order RT elements.

For the results shown in Figure 7 we use BDM and RT elements of order  $k = 1, 2$ . The optimal  $L^2$ -convergence rates stated in Theorem 6.2 are confirmed to be sharp when using BDM elements for  $k > 0$ . If we increase the test-space, however, to  $RT$  elements, which additionally include specific polynomials of one order higher than BDM, one order of convergence is lost, compare also Figure 8. Note, that the lowest order BDM finite space already includes linear elements.

## 8. CONCLUSION

In this paper we presented an improved convergence analysis for the Gauss curvature and connection 1-form. The proof is based on the covariant curl and incompatibility operators based on only tangential-tangential continuous metric fields. These distributional nonlinear differential operators appear when linearizing the Gauss curvature and connection 1-form enabling an integral representation. The second crucial ingredient for the numerical analysis were the orthogonality properties valid for the interpolation operator into the Regge finite element space. They enabled to extract one additional order of convergence compared to standard estimates. The theory predicts a super-convergence property for Regge finite elements known for mixed methods. Numerical experiments validated that the stated order of convergence is reached and sharp.

## ACKNOWLEDGMENTS

Michael Neunteufel and Joachim Schöberl acknowledge support by the Austrian Science Fund (FWF) project F65.

## APPENDIX A. RATIONALE FOR THE TEST SPACE IN COVARIANT CURL EXTENSION

This section briefly presents a justification for the use of  $g$ -normal continuous vector fields in Definition 4.2. For any mesh vertex  $V \in \mathcal{V}$ , let  $B_\varepsilon(V) = \{q \in M : d_g(q, V) \leq \varepsilon\}$ . Then put  $D_\varepsilon = \cup_{V \in \mathcal{V}} B_\varepsilon(V)$  and  $\check{M}_\varepsilon = M \setminus D_\varepsilon$ . Let  $U_i, \check{\Phi}_i : U_i \rightarrow \mathbb{R}^2$  denote a chart of the glued smooth structure. In the parameter domain  $\check{\Phi}_i(U_i)$ , using the Euclidean divergence operator, define  $W^p(\text{div}, \check{\Phi}_i(U_i)) = \{w \in L^p(\check{\Phi}_i(U_i)), \text{div}(w) \in L^p(\check{\Phi}_i(U_i))\}$  for any  $1 \leq p < \infty$  with its natural Euclidean norm. This norm and the duality pairings defined in (4.14) and (4.15) feature in the next result.

**Proposition A.1.** *Let  $\sigma \in \mathcal{R}(\mathcal{T})$  and  $W \in \mathring{W}_g(\mathcal{T})$ . For any given  $\varepsilon_1 > 0$ , there exists a  $p > 2$ , an  $\varepsilon_2 > 0$ , finitely many charts  $\{(U_i, \check{\Phi}_i) : i \in I\}$  covering  $\check{M}_{\varepsilon_2}$  in the glued smooth structure, a partition of unity  $\psi_i$  subordinate to  $U_i$ , and a smooth  $\varphi \in \mathfrak{X}_c(\check{M})$  such that  $\varphi = \sum_{i \in I} \varphi_i$  with support of  $\varphi_i$  contained in  $U_i$ , satisfies*

$$|\langle \text{curl}_g \sigma, \varphi \rangle_{\mathfrak{X}_c(\check{M})} - \langle \text{curl}_g \sigma, W \rangle_{\mathring{W}_g(\mathcal{T})}| \leq \varepsilon_1, \quad \text{and} \quad (\text{A.1})$$

$$\|(\check{\Phi}_i)_*(\varphi_i - \psi_i W)\|_{W^p(\text{div}, \check{\Phi}_i(U_i \cap \check{M}_{\varepsilon_2}))} \leq \varepsilon_1, \quad \text{for all } i \in I. \quad (\text{A.2})$$

*Proof.* As a first step, we zero out the vector field  $W$  near vertices. For any  $\varepsilon > 0$ , let  $0 \leq \chi_\varepsilon \leq 1$  be a smooth cutoff function satisfying  $\chi_\varepsilon \equiv 1$  in  $\check{M}_\varepsilon$  and  $\chi_\varepsilon \equiv 0$  in  $D_{\varepsilon/2}$ , and let

$$r_\varepsilon(W) = \sum_{T \in \mathcal{T}} \int_{(T \cap D_\varepsilon, g)} g(W, W)^{1/2} + \int_{(\partial T \cap D_\varepsilon, g)} g(W, W)^{1/2}.$$

Since  $\sigma$  is piecewise smooth, there is a constant  $C_\sigma$  depending only on  $\sigma$  (independent of  $\varepsilon$ ) such that  $|\langle \text{curl}_g \sigma, W - \chi_\varepsilon W \rangle_{\mathring{W}_g(\mathcal{T})}| \leq C_\sigma r_\varepsilon(W)$ . Since  $W$  is piecewise smooth,  $r_\varepsilon(W)$  approaches zero as  $\varepsilon \rightarrow 0$ . Hence for the given  $\varepsilon_1$ , there is a  $\varepsilon_2 > 0$  such that

$$\left| \langle \text{curl}_g \sigma, W - \chi_\varepsilon W \rangle_{\mathring{W}_g(\mathcal{T})} \right| \leq C_\sigma r_\varepsilon(W) \leq \frac{\varepsilon_1}{2} \quad \text{for all } \varepsilon \leq \varepsilon_2. \quad (\text{A.3})$$

Next, to approximate  $\chi_{\varepsilon_2}W$  by a smooth vector field, we use the precompactness of  $\check{M}_{\varepsilon_2/2}$  to extract a finite subcover from the maximal atlas of  $\check{M}$ . Denoting the resulting finitely many charts by  $U_i, \check{\Phi}_i$ , let  $\psi_i$  be a partition of unity subordinate to the cover  $U_i$ . We focus on a  $U_i$  that intersects an edge  $E$  (since the other cases are easier) and use the accompanying notation in (2.18). By the previous discussion of the coordinate construction (4.12),  $\check{\partial}^1 = \hat{\nu}_+$  and  $\check{\partial}^2 = \hat{\tau}_+$  along  $E$ , so the expansion  $W = W^i \check{\partial}_i$  is  $g$ -orthonormal and

$$g(W|_{T_{\pm}}, \hat{\nu}_{\pm}) = \pm W^1|_{T_{\pm}} \text{ on } E \cap U_i. \quad (\text{A.4})$$

The  $g$ -normal continuity  $\llbracket g(W, \hat{\nu}) \rrbracket = 0$  implies that  $W_i = \psi_i \chi_{\varepsilon_2} W$  pushed forward to the parameter domain, namely  $(\check{\Phi}_i)_* W_i$ , has continuous normal component across  $\check{\Phi}_i(E)$ , a subset of the axis  $\check{Y} = \{(\check{x}^1, \check{x}^2) \in \mathbb{R}^2 : \check{x}^1 = 0\}$ . Hence  $(\check{\Phi}_i)_* W_i$  is in  $\check{W}^p(\text{div}, \check{\Phi}_i(U_i))$ , the subspace of  $W^p(\text{div}, \check{\Phi}_i(U_i))$  with zero normal traces on the boundary of  $\check{\Phi}_i(U_i)$ . By a well known density result in the Euclidean domain, there exists an infinitely smooth compactly supported vector field  $\check{\varphi}_i$  on  $\check{\Phi}_i(U_i)$  that is arbitrarily close to  $(\check{\Phi}_i)_* W_i$ , so letting  $\varphi_i = (\check{\Phi}_i^{-1})_* \check{\varphi}_i$ , we have

$$\|(\check{\Phi}_i)_*(\varphi_i - W_i)\|_{W^p(\text{div}, \check{\Phi}_i(U_i))} = \|\check{\varphi}_i - (\check{\Phi}_i)_* W_i\|_{W^p(\text{div}, \check{\Phi}_i(U_i))} \leq \varepsilon_1. \quad (\text{A.5})$$

Moreover, since  $\chi_{\varepsilon_2} \equiv 1$  in  $U_i \cap \check{M}_{\varepsilon_2}$ , the functions  $W_i$  and  $\psi_i W$  coincide there, so (A.2) follows. Constructing such  $\varphi_i$  on every  $U_i$ , put  $\varphi = \sum_{i \in I} \varphi_i$ .

To prove (A.1), in view of (A.3), it suffices to show that

$$\left| \langle \text{curl}_g \sigma, \chi_{\varepsilon_2} W \rangle_{\check{W}_g(\mathcal{J})} - \langle \text{curl}_g \sigma, \varphi \rangle_{\mathfrak{X}_c(\check{M})} \right| \leq \frac{\varepsilon_1}{2}. \quad (\text{A.6})$$

Obviously, the element contributions in the difference above,  $\int_{\mathcal{J}} (\text{curl}_g \sigma)(\chi_{\varepsilon_2} W - \varphi)$ , can be made arbitrarily small by (A.5), revising the choice of  $\varphi_i$  if needed. For the element boundary terms,

$$\int_{\partial \mathcal{J}} g(\chi_{\varepsilon_2} W - \varphi, \hat{\nu}) \sigma(\hat{\nu}, \hat{\tau}) = \sum_{i \in I} \int_{\partial \mathcal{J}} g(W_i - \varphi_i, \hat{\nu}) \sigma(\hat{\nu}, \hat{\tau}), \quad (\text{A.7})$$

we focus, as before, on a neighborhood  $U_i$  intersecting an edge  $E = \partial T_- \cap \partial T_+$ . On  $\partial T^+$ , by (A.4),  $g(W_i - \varphi_i, \hat{\nu}) = W_i^1 - \varphi_i^1$  yields the normal component of the pushforward  $(\check{\Phi}_i)_*(W_i - \varphi_i)$  on  $\check{Y}$ -axis in the parameter domain. Since the latter converges to zero in  $W^p(\text{div}, \check{\Phi}_i(U_i))$  by (A.5), its normal trace converges to zero in  $W^{-1/p, p}(\check{\Phi}_i(E \cap U_i))$ . Choose  $p > 2$  and  $q$  such that  $1/p + 1/q = 1$ . When  $\sigma(\hat{\nu}, \hat{\tau})$  is mapped to  $\check{\Phi}_i(E \cap U_i)$  and extended to  $\check{Y}$ -axis by zero, is in  $W^{1-1/q, q}(\check{Y})$  since  $1 - 1/q < 1/2$ . Hence the contribution from  $U_i \cap E$  to the right hand side of (A.7) vanishes. Repeating this argument on other charts, (A.6) is proved.  $\square$

**A.1. Relation between covariant inc and covariant rotated divdiv.** In this section we show that the distributional covariant inc from Proposition 4.5 and the covariant distributional divdiv operator of a rotated sigma used in [11] coincide in the sense

$$\langle \text{div}_g \text{div}_g S_g \sigma, u \rangle_{\check{\mathcal{V}}(\mathcal{J})} = -\langle \text{inc}_g \sigma, u \rangle_{\check{\mathcal{V}}(\mathcal{J})} \quad \text{for all } u \in \check{\mathcal{V}}(\mathcal{J}). \quad (\text{A.8})$$

This is in common with the Euclidean identity  $\operatorname{div}\operatorname{div}S\sigma = \operatorname{div}\operatorname{div}g - \Delta \operatorname{tr}(\sigma) = -\operatorname{inc}\sigma$ ,  $\operatorname{tr}(\cdot)$  denoting the trace of a matrix. The distributional covariant  $\operatorname{div}\operatorname{div}$  reads

$$\begin{aligned} \langle \operatorname{div}_g \operatorname{div}_g S_g \sigma, u \rangle_{\dot{\mathcal{V}}(\mathcal{T})} &= \int_{\mathcal{T}} u \operatorname{div}_g \operatorname{div}_g S_g \sigma + \int_{\partial \mathcal{T}} u ((\operatorname{div}_g S_g \sigma)^\flat(\hat{\nu}) + (d^0 \sigma_{\hat{\nu}\hat{\tau}})(\hat{\tau})) \\ &\quad - \sum_{T \in \mathcal{T}} \sum_{V \in \mathcal{V}_T} \llbracket \sigma_{\hat{\nu}\hat{\tau}} \rrbracket_V^T u(V), \end{aligned} \quad (\text{A.9})$$

where  $S_g \sigma = \sigma - \operatorname{tr}_g(\sigma)g$  with  $\operatorname{tr}_g(\sigma) = \sigma_{ij}g^{ij}$ . Note, that the authors in [11] used  $(\hat{\nu}, \hat{\tau})$  as positively oriented frame, whereas we use  $(\hat{\tau}, \hat{\nu})$  such that the signs in the boundary and vertex terms differ.

The covariant divergence is defined as the  $L^2$ -adjoint of the covariant gradient. For  $f \in \Lambda^0(\Omega)$  its covariant gradient is given by the equation

$$g(\operatorname{grad}_g f, v) = d^0 f(v) \quad \text{for all } v \in \mathfrak{X}(\Omega)$$

and  $\operatorname{div}_g : \mathfrak{X}(\Omega) \rightarrow \Lambda^0(\Omega)$  by

$$\int_{\Omega} g(\operatorname{grad}_g f, v) = \int_{\Omega} f \operatorname{div}_g v, \quad \text{for all } v \in \mathring{\mathcal{W}}_g(\Omega), f \in \Lambda^0(\Omega),$$

in coordinates

$$\operatorname{div}_g v = \frac{1}{\sqrt{\det g}} \partial_i (\sqrt{\det g} v^i), \quad v \in \mathfrak{X}(\Omega).$$

The usual extension to tensor fields, see e.g. [27],  $\operatorname{div}_g : \mathcal{T}_0^2(\Omega) \rightarrow \mathfrak{X}(\Omega)$  reads in coordinates

$$\operatorname{div}_g \sigma = (\partial_j \sigma^{ij} + \Gamma_{lj}^i \sigma^{lj} + \Gamma_{jl}^j \sigma^{il}) \partial_i, \quad \sigma \in \mathcal{T}_0^2(\Omega).$$

Note, that for  $\sigma \in \mathcal{T}_2^0$  there holds  $\sigma^{ij} = g^{ik} \sigma_{kl} g^{lj}$ .

**Lemma A.2.** *There holds for  $g \in \mathcal{R}^+(\mathcal{T})$  and  $\sigma \in \mathcal{R}(\mathcal{T})$*

- (1)  $\star(\operatorname{div}_g S_g \sigma)^\flat = -\operatorname{curl}_g \sigma$  and  $(\operatorname{div}_g S_g \sigma)^\flat(\hat{\nu}) = (\operatorname{curl}_g \sigma)(\hat{\tau})$  on  $\partial \mathcal{T}$ ,
- (2)  $\operatorname{div}_g \operatorname{div}_g S_g \sigma = \operatorname{div}_g \operatorname{div}_g \sigma - \Delta_g \operatorname{tr}_g(\sigma) = -\operatorname{inc}_g \sigma$  on  $\mathcal{T}$ ,

where  $\Delta_g f := \operatorname{div}_g \operatorname{grad}_g f$ ,  $f \in \Lambda^0(\Omega)$  denotes the Laplace-Beltrami operator.

*Proof.* In the first identity only first order derivatives of  $g$  are involved. Thus, in normal coordinates,  $\tilde{x}^i$  (see §5.1),  $(\operatorname{div}_g S_g \sigma)^\flat$  becomes the Euclidean version  $\operatorname{div}(\sigma - \operatorname{tr}(\sigma)\delta)$ , which reads in components

$$(\operatorname{div}_g S_g \sigma)^\flat = \begin{pmatrix} \tilde{\partial}_2 \tilde{\sigma}_{12} - \tilde{\partial}_1 \tilde{\sigma}_{22} \\ \tilde{\partial}_1 \tilde{\sigma}_{12} - \tilde{\partial}_2 \tilde{\sigma}_{11} \end{pmatrix}.$$

The Hodge-star operator translates in a counter-clockwise 90-degree rotation,  $\tilde{\star} = -\varepsilon^{ij}$ , such that

$$\star(\operatorname{div}_g S_g \sigma)^\flat = - \begin{pmatrix} \tilde{\partial}_1 \tilde{\sigma}_{12} - \tilde{\partial}_2 \tilde{\sigma}_{11} \\ \tilde{\partial}_1 \tilde{\sigma}_{22} - \tilde{\partial}_2 \tilde{\sigma}_{12} \end{pmatrix} = -\operatorname{curl} \tilde{\sigma},$$

which coincides with  $-\operatorname{curl}_g \sigma$  in normal coordinates. The identity  $(\operatorname{div}_g S_g \sigma)^\flat(\hat{\nu}) = (\operatorname{curl}_g \sigma)(\hat{\tau})$  follows now by (2.12).

The identity  $\operatorname{div}_g \operatorname{div}_g S_g \sigma = \operatorname{div}_g \operatorname{div}_g \sigma - \Delta_g \operatorname{tr}_g(\sigma)$  [27] is used to treat the terms separately in normal coordinates. The Laplace-Beltrami operator becomes

$$[\Delta_g \operatorname{tr}_g(\sigma)] = [\operatorname{div}_g \nabla_g \operatorname{tr}_g(\sigma)] = [\operatorname{div}_g(g^{ij} \partial_j \operatorname{tr}_g(\sigma))] = \tilde{\partial}_i^2 \operatorname{tr}(\tilde{\sigma} \tilde{g}^{-1}) = \Delta \operatorname{tr}(\tilde{\sigma}) - \operatorname{tr}(\tilde{\sigma} \Delta \tilde{g})$$

and the divdiv part

$$[\operatorname{div}_g \operatorname{div}_g \sigma] = \operatorname{div} \operatorname{div}[\tilde{\sigma}] - 2\tilde{\partial}_{ij}^2 \tilde{g}_{ik} \tilde{\sigma}_{kj} + \tilde{\partial}_i \tilde{\Gamma}_{lji} \tilde{\sigma}_{lj} - \tilde{\partial}_i \tilde{\Gamma}_{jlj} \tilde{\sigma}_{il}.$$

Note, that we used in abuse of notation summation of repeated indices although all indices are subscripts due to the usage of normal coordinates.

Further, by inserting the definition of Christoffel symbols of first kind and a lengthy but elementary computation gives

$$\begin{aligned} -2\tilde{\partial}_{ij}^2 \tilde{g}_{ik} \tilde{\sigma}_{kj} + \tilde{\partial}_i \tilde{\Gamma}_{lji} \tilde{\sigma}_{lj} - \tilde{\partial}_i \tilde{\Gamma}_{jlj} \tilde{\sigma}_{il} &= \tilde{\sigma}_{11} \left( \frac{1}{2} \tilde{\partial}_1^2 \tilde{g}_{22} - \tilde{\partial}_1^2 \tilde{g}_{11} - \frac{1}{2} \tilde{\partial}_2^2 \tilde{g}_{11} - \tilde{\partial}_1 \tilde{\partial}_2 \tilde{g}_{12} \right) \\ &\quad - 2\tilde{\sigma}_{12} (\tilde{\partial}_1^2 \tilde{g}_{12} + \tilde{\partial}_2^2 \tilde{g}_{12}) - \tilde{\sigma}_{22} (\tilde{\partial}_2^2 \tilde{g}_{22} + \frac{1}{2} \tilde{\partial}_1^2 \tilde{g}_{22} + \tilde{\partial}_1 \tilde{\partial}_2 \tilde{g}_{12} - \frac{1}{2} \tilde{\partial}_2^2 \tilde{g}_{11}) \end{aligned}$$

and thus combining the results

$$[\operatorname{div}_g \operatorname{div}_g S_g \sigma] = -\operatorname{inc}[\tilde{\sigma}] + \frac{1}{2} \operatorname{tr}[\tilde{\sigma}] \operatorname{inc}[\tilde{g}],$$

which finishes the proof by comparing with (5.3).  $\square$

Using the results of Lemma A.2 and comparing the terms of (A.9) with Proposition 4.5 shows that identity (A.8) holds.

## REFERENCES

- [1] AGMON, S. *Lectures on Elliptic Boundary Value Problems*. Van Nostrand Math. Stud. 2, Van Nostrand, Princeton, Toronto, London, 1965.
- [2] AMSTUTZ, S., AND VAN GOETHEM, N. Analysis of the incompatibility operator and application in intrinsic elasticity with dislocations. *SIAM Journal on Mathematical Analysis* 48, 1 (2016), 320–348.
- [3] AMSTUTZ, S., AND VAN GOETHEM, N. Existence and asymptotic results for an intrinsic model of small-strain incompatible elasticity. *Discrete & Continuous Dynamical Systems - B* 25, 10 (2020), 3769–3805.
- [4] ARNOLD, D. N. *Finite element exterior calculus*. SIAM, Philadelphia, 2018.
- [5] ARNOLD, D. N., AWANOU, G., AND WINTHER, R. Finite elements for symmetric tensors in three dimensions. *Math. Comp.* 77, 263 (2008), 1229–1251.
- [6] ARNOLD, D. N., FALK, R., AND WINTHER, R. Finite element exterior calculus: from Hodge theory to numerical stability. *Bulletin of the American Mathematical Society* 47, 2 (2010), 281–354.
- [7] ARNOLD, D. N., FALK, R. S., AND WINTHER, R. Finite element exterior calculus, homological techniques, and applications. *Acta numerica* 15 (2006), 1–155.
- [8] ARNOLD, D. N., AND HU, K. Complexes from complexes. *Foundations of Computational Mathematics* 21, 6 (2021), 1739–1774.
- [9] ARNOLD, D. N., AND WALKER, S. W. The Hellan–Herrmann–Johnson method with curved elements. *SIAM Journal on Numerical Analysis* 58, 5 (2020), 2829–2855.
- [10] BARRETT, J. W., ORITI, D., AND WILLIAMS, R. M. Tullio Regge’s legacy: Regge calculus and discrete gravity. *arXiv preprint arXiv:1812.06193* (2018).
- [11] BERCHENKO-KOGAN, Y., AND GAWLIK, E. S. Finite element approximation of the Levi-Civita connection and its curvature in two dimensions, 2021.

- [12] BOFFI, D., BREZZI, F., AND FORTIN, M. *Mixed finite element methods and applications*, 1 ed., vol. 44. Springer-Verlag Berlin Heidelberg, Berlin, Heidelberg, 2013.
- [13] BORRELLI, V., CAZALS, F., AND MORVAN, J.-M. On the angular defect of triangulations and the pointwise approximation of curvatures. *Computer Aided Geometric Design* 20, 6 (2003), 319–341.
- [14] BREZZI, F., DOUGLAS, J., AND MARINI, L. D. Two families of mixed finite elements for second order elliptic problems. *Numerische Mathematik* 47, 2 (1985), 217–235.
- [15] CARMO, M. P. D. *Riemannian Geometry*, 1 ed. Birkhäuser Basel, Mannheim, 1992.
- [16] CHEEGER, J., MÜLLER, W., AND SCHRADER, R. On the curvature of piecewise flat spaces. *Communications in Mathematical Physics* 92, 3 (1984), 405–454.
- [17] CHRISTIANSEN, S., GOPALAKRISHNAN, J., GUZMÁN, J., AND HU, K. A discrete elasticity complex on three-dimensional Alfeld splits. *Preprint* (2021).
- [18] CHRISTIANSEN, S. H. A characterization of second-order differential operators on finite element spaces. *Mathematical Models and Methods in Applied Sciences* 14, 12 (2004), 1881–1892.
- [19] CHRISTIANSEN, S. H. On the linearization of Regge calculus. *Numerische Mathematik* 119, 4 (2011), 613–640.
- [20] CHRISTIANSEN, S. H. Exact formulas for the approximation of connections and curvature, 2015.
- [21] CLARKE, C. J. S., AND DRAY, T. Junction conditions for null hypersurfaces. *Classical and Quantum Gravity* 4, 2 (1987), 265–275.
- [22] COMODI, M. I. The Hellan–Herrmann–Johnson method: Some new error estimates and postprocessing. *Mathematics of Computation* 52, 185 (1989), 17–29.
- [23] FISCHER, A. E., AND MARSDEN, J. E. Deformations of the scalar curvature. *Duke Math. J.* (1975).
- [24] FRITZ, H. Isoparametric finite element approximation of Ricci curvature. *IMA Journal of Numerical Analysis* 33, 4 (2013), 1265–1290.
- [25] FRITZ, H. Numerical Ricci–DeTurck flow. *Numerische Mathematik* 131, 2 (2015), 241–271.
- [26] GAWLIK, E. S. Finite element methods for geometric evolution equations. In *Geometric Science of Information* (Cham, 2019), F. Nielsen and F. Barbaresco, Eds., Springer International Publishing, pp. 532–540.
- [27] GAWLIK, E. S. High-order approximation of Gaussian curvature with Regge finite elements. *SIAM Journal on Numerical Analysis* 58, 3 (2020), 1801–1821.
- [28] HAURET, P., AND HECHT, F. A discrete differential sequence for elasticity based upon continuous displacements. *SIAM Journal on Scientific Computing* 35, 1 (2013), B291–B314.
- [29] KOSOVSKIĬ, N. N. Gluing of Riemannian manifolds of curvature  $\geq \kappa$ . *Algebra i Analiz* 14, 3 (2002), 140–157.
- [30] KOSOVSKIĬ, N. N. Gluing with branching of Riemannian manifolds of curvature  $\leq \kappa$ . *Algebra i Analiz* 16, 4 (2004), 132–145.
- [31] LEE, J. M. *Riemannian manifolds: an introduction to curvature*, 1 ed. Springer, New York, NY, New York, 1997.
- [32] LEE, J. M. *Introduction to Smooth Manifolds*, 2 ed. Springer, 2012.
- [33] LI, L. *Regge Finite Elements with Applications in Solid Mechanics and Relativity*. PhD thesis, University of Minnesota, 2018.
- [34] LIU, D., AND XU, G. Angle deficit approximation of Gaussian curvature and its convergence over quadrilateral meshes. *Computer-Aided Design* 39, 6 (2007), 506–517.
- [35] NEUNTEUFEL, M. *Mixed Finite Element Methods for Nonlinear Continuum Mechanics and Shells*. PhD thesis, TU Wien, 2021.
- [36] NEUNTEUFEL, M., AND SCHÖBERL, J. Avoiding membrane locking with Regge interpolation. *Computer Methods in Applied Mechanics and Engineering* 373 (2021), 113524.
- [37] PETERSEN, P. *Riemannian Geometry*, third ed. Springer, 2016.
- [38] RAVIART, P.-A., AND THOMAS, J.-M. A mixed finite element method for 2-nd order elliptic problems. In *Mathematical Aspects of Finite Element Methods*, vol. 66. Springer, 1977, pp. 292–315.



- [39] REGGE, T. General relativity without coordinates. *Il Nuovo Cimento (1955-1965)* 19, 3 (1961), 558–571.
- [40] REGGE, T., AND WILLIAMS, R. M. Discrete structures in gravity. *Journal of Mathematical Physics* 41, 6 (2000), 3964–3984.
- [41] SCHÖBERL, J. NETGEN an advancing front 2D/3D-mesh generator based on abstract rules. *Computing and Visualization in Science* 1, 1 (1997), 41–52.
- [42] SCHÖBERL, J. C++ 11 implementation of finite elements in NGSolve. *Institute for Analysis and Scientific Computing, Vienna University of Technology* (2014).
- [43] SORKIN, R. Time-evolution problem in Regge calculus. *Phys. Rev. D* 12 (1975), 385–396.
- [44] SULLIVAN, J. M. *Curvatures of Smooth and Discrete Surfaces*. Birkhäuser Basel, Basel, 2008, pp. 175–188.
- [45] TU, L. W. *Differential Geometry: Connections, Curvature, Characteristic Classes*. Springer, 2017.
- [46] WALKER, S. W. The Kirchhoff plate equation on surfaces: the surface Hellan–Herrmann–Johnson method. *IMA Journal of Numerical Analysis* (2021).
- [47] WARDETZKY, M. *Discrete differential operators on polyhedral surfaces—convergence and approximation*. PhD thesis, Freie Universität Berlin, 2006.
- [48] WHITNEY, H. *Geometric integration theory*. Princeton University Press, Princeton, N. J, 1957.
- [49] WILLIAMS, R. M., AND TUCKEY, P. A. Regge calculus: a brief review and bibliography. *Classical and Quantum Gravity* 9, 5 (1992), 1409–1422.
- [50] XU, G. Convergence analysis of a discretization scheme for Gaussian curvature over triangular surfaces. *Computer Aided Geometric Design* 23, 2 (2006), 193–207.
- [51] XU, Z., AND XU, G. Discrete schemes for Gaussian curvature and their convergence. *Computers & Mathematics with Applications* 57, 7 (2009), 1187–1195.

PORTLAND STATE UNIVERSITY, PO BOX 751, PORTLAND OR 97207, USA  
Email address: gjay@pdx.edu

INSTITUTE FOR ANALYSIS AND SCIENTIFIC COMPUTING, TU WIEN, WIEDNER HAUPTSTR. 8-10,  
1040 WIEN, AUSTRIA  
Email address: michael.neunteufel@tuwien.ac.at

INSTITUTE FOR ANALYSIS AND SCIENTIFIC COMPUTING, TU WIEN, WIEDNER HAUPTSTR. 8-10,  
1040 WIEN, AUSTRIA  
Email address: joachim.schoeberl@tuwien.ac.at

INSTITUTE OF NUMERICAL AND APPLIED MATHEMATICS, UNIVERSITY OF GÖTTINGEN, LOTZESTR.  
16-18, 37083 GÖTTINGEN, GERMANY  
Email address: email: wardetzky@math.uni-goettingen.de