

CFT description of BH's and ECO's: QNMs, superradiance, echoes and tidal responses

Dario Consoli,^a Francesco Fucito,^a Jose Francisco Morales,^a Rubik Poghossian.^b

^a*Dipartimento di Fisica, Università di Roma "Tor Vergata" & Sezione INFN Roma2, Via della ricerca scientifica 1, 00133, Roma, Italy*

^b*Yerevan Physics Institute, Alikhanian Br. 2, 0036 Yerevan, Armenia*

ABSTRACT: Using conformal field theory and localization techniques we study the propagation of waves in gravity backgrounds described by Schrödinger like equations with Fuchsian singularities. Exact formulae for the connection matrices relating the asymptotic behaviour of the wave functions near the singularities are obtained in terms of braiding and fusion rules of the CFT. The results are applied to the study of quasi normal modes, absorption cross sections, amplification factors, echoes and tidal responses of black holes (BH) and exotic compact objects (ECO) in four and five dimensions. In particular, we propose a definition of dynamical Love numbers in gravity.

Contents

1	Introduction	2
2	SW Gravity correspondence	4
2.1	The CFT description	4
2.1.1	The conformal blocks	5
2.1.2	The BPZ equation	6
2.2	Braiding and fusion relations	7
2.3	Five-point conformal block	9
2.4	BPZ equation from the quantum deformed SW curve	12
2.5	Irregular singularities	14
3	Wave scattering: QNM's, cross sections and echoes	14
3.1	Absorption cross section, QNM's and superradiance	15
3.2	Echoes	17
4	Kerr-Newman geometries in four dimensions	18
4.1	The wave equation	18
4.2	Angular motion	18
4.3	Radial motion	20
4.4	Love numbers	20
4.5	Amplification factors	22
5	Solutions of the Einstein-Maxwell theory in five dimensions	23
5.1	The wave equation	23
5.2	Angular motion	24
5.3	Radial motion	24
5.4	Love numbers	25
5.5	Amplification factors	26
6	Summary of results	26
A	Instanton partition function	27
B	Hypergeometric identities	28

1 Introduction

Very recently we have entered the era of gravitational wave (GW) astronomy with the first measurements coming from coalescing binaries [1]. Can this experiments test fundamental physics? The coalescence of two astrophysical objects can be divided into three phases: the inspiral stage where the two objects are well separated and orbit around each other, the merger phase when they coalesce and the ringdown phase when the compact remnant relaxes emitting gravitational waves of very characteristic frequencies, the so called quasi-normal modes (QNM's) [2]. The inspiral and ring-down phases can be studied using space-time perturbation theory. Indeed, the tidal response of a neutron star can be traced from the GW signal emitted in the late inspiral stage and expressed in terms of Love numbers [3]. On the other hand, QNM's dominate the ring-down phase.

Can the GW signal be used to distinguish BH's from exotic objects more massive than neutron stars but without a horizon ? The list of alternatives is long: gravastars, wormholes, firewalls and fuzzballs, collectively known as Exotic Compact Objects (ECO) [4]. ECO's tend to be as black as BH's. If compact enough, they are typically surrounded by a photon sphere where both massive and massless probes orbit in unstable circular motion [5]. External perturbations excite the QNM's which appear as damped vibrations of the ECO response. A fraction of this signal, moving towards the ECO interior, is reflected back producing a sequences of echoes that can be viewed as the smoking gun for the existence of a horizonless object [6]. Several attempts to distinguish BH's from fuzzballs studying their photon sphere sizes and shapes, QNM's, multipolar structure and echoes has recently appeared in [7]-[16]. Hidden symmetries of black hole and fuzzball photon spheres have been explored in [17, 18]

In this paper, we study the wave response of BH's and ECO's to small deformations of the geometry. These studies were pioneered by Regge and Wheeler in the late 50's [19]. The wave propagation in the Schwarzschild geometry turns out to be a Schrödinger like equation describing the scattering of a particle in the presence of an effective potential. The results were generalized by Teukolski to the case of Kerr-Newman geometry [20–22]. In general, these equations cannot be solved analytically even for very simple BH's, so most of the available results rely on numerical analysis.

Quite recently, a new astonishing connection between the QNM spectral problem and the quantum periods of the Seiberg-Witten (SW) geometries was proposed in [23] and tested in the Kerr case. The correspondence was extended in [24, 25] to several gravity systems including BH's in higher dimensions, D-brane bound states and fuzzballs. Moreover, using the AGT correspondence [26], the solution of the wave equation in the Kerr case was identified with certain five-point conformal blocks of a two-dimensional conformal field theory (CFT). This provides a new tool to study other interesting observables such as Love numbers, absorption coefficients and grey body factors [27]. QNM's and connections matrices have been also determined in [30] using techniques based on integrability.

In this work, following [27], we apply CFT and localization techniques to the study of wave equations describing radial and angular motion in separable gravity backgrounds described by Schrödinger like equations with Fuchsian singularities. We relate the solutions

of a differential equation with N Fuchsian singularities to $(N + 1)$ -point conformal blocks of Liouville theory involving the insertion of N primaries and a degenerate field, i.e. a primary field containing a null state in its Verma module. These conformal blocks obey a differential equation known as (BPZ)[31] and are related via AGT correspondence [26] to $\mathcal{N} = 2$ supersymmetric quiver gauge theories living in curved space times. In a particular Ω -background (NS) [32], the gauge dynamics is given by a quantization of the SW curve [33, 34] and the BPZ equation reduces to an ordinary differential equation that can be identified with the wave equation in gravity.

Combining the two pictures, we will derive a combinatorial formula for the wave function and exact formulae for the connection matrices relating the asymptotic expansions of the solutions near any two singularities. These matrices are given by the "braiding" and "fusion" rules of the underlying CFT. Similar results have been recently obtained in [35]. Solutions of Fuchsian equations have been also studied in [36–43] using isomonodromic methods and Painlevé transcendents.

We discuss various applications of the gauge/CFT techniques in the context of BH's and ECO's in four and five dimensions. ECO's are represented by solutions of Einstein equations on space times ending on a membrane with reflectivity properties. The effect of the inner boundary reflectivity coefficient on QNM's and echo responses is discussed. Finally we compute the amplification factors and dynamical Love numbers for slowly oscillating (yet non-static) waves. In the astrophysical realm static Love numbers were first defined and studied in [44, 45]. For four dimensional Schwarzschild [44–46] and Kerr [47] BH's they were shown to vanish. They are generically non-trivial in higher dimensions [48, 49], non-asymptotically flat spaces [50] and alternative theories of gravity [51, 52]. In the dynamical case, there is no agreement on their definition. Here we propose a definition of the dynamical Love numbers describing the tidal response to a slowly varying (yet not static) source in gravity. The Love numbers are extracted from the asymptotic form of the scalar wave function in the region $r_+ - r_- \ll r \ll \omega^{-1}$ ¹. We derive a unique formula describing the Love and dissipation numbers of all BH's of Einstein-Maxwell theory in four and five dimensions in terms of few parameters specifying the gravity background. The entire non-triviality of the differential equation (the Heun equation) is codified into a single function: the quantum SW period $a(u)$ specifying the gauge theory prepotential $\mathcal{F}(a)$! This function can be derived from a continuous fraction equation representing a quantum deformed version of the SW curve [33, 34].

This is the plan of the paper: in Section 2 we describe the gauge-CFT-gravity dictionary. In Section 3 we study the QNM's and echo wave responses of gravity backgrounds on spacetimes ending on a membrane with non-trivial reflectivity properties. In Sections 4 and 5 we compute amplification factors and dynamical Love numbers for BH's in four and five dimensions using the CFT description of the wave functions and connection matrices. In Section 6 we summarise our results.

¹Different definitions of the dynamical Love numbers have been proposed in [27, 53].

2 SW Gravity correspondence

In this section, we review the SW gravity correspondence in its most general set up. We consider a gravity system whose wave equation can be separated into ordinary differential equations with Fuchsian singularities describing radial and angular motion. An ordinary differential equation of second order with N Fuchsian singularities can be always put into the normal form

$$\left[\frac{d^2}{dx^2} + Q(x) \right] \Psi(x) = 0 \quad (2.1)$$

with

$$Q(x) = \sum_{i=1}^N \left[\frac{\delta_i}{(x-x_i)^2} + \frac{c_i}{(x-x_i)} \right] \quad (2.2)$$

where δ_i, c_i, x_i are some constants parametrizing the positions x_i of the singularities and the behaviour of the equation in their neighborhood. Here we assume that infinity is a regular point, i.e. $Q(x)$ vanishes at x^{-4} , at large x , i.e.

$$\sum_{i=1}^N c_i = \sum_{i=1}^N (c_i x_i + \delta_i) = \sum_{i=1}^N (c_i x_i^2 + 2 \delta_i x_i) = 0 \quad (2.3)$$

There is a coordinate freedom in choosing the location of the singularities. We will always use this freedom to map the locations of three of the singularities x_0, x_1, x_∞ to $0, 1, \infty$. We take

$$x \rightarrow z(x) = \frac{x-x_0}{x-x_\infty} \frac{x_1-x_\infty}{x_1-x_0} \quad (2.4)$$

Under a general coordinate transformation $x \rightarrow z(x)$, a differential equation of the form (2.1) can be brought to its normal form by taking

$$Q(z) = Q(x(z)) x'(z)^2 + \frac{x'''(z)}{2x'(z)} - \frac{3}{4} \left[\frac{x''(z)}{x'(z)} \right]^2, \quad \Psi(z) = \frac{\Psi(x(z))}{\sqrt{x'(z)}} \quad (2.5)$$

where the prime stands for derivation. (2.3) can be solved for c_0, c_1 and c_∞ as

$$c_\infty = 0, \quad c_1 = \delta - \sum_{i \neq 0,1,\infty} z_i c_i, \quad c_0 = -\delta + \sum_{i \neq 0,1,\infty} (z_i - 1) c_i \quad (2.6)$$

with

$$\delta = 2\delta_\infty - \sum_{i=1}^N \delta_i \quad (2.7)$$

The aim of this section is to derive a combinatorial formula for the solutions $\Psi_\alpha(z)$ and the connection matrices relating its asymptotic behaviour near the singularities.

2.1 The CFT description

We consider a Liouville theory with background charge Q and central charge c given by

$$c = 1 + 6Q^2, \quad Q = b + \frac{1}{b} \quad (2.8)$$




Figure 1: (a) The diagram for the linear quiver $U(2)^n$ gauge theory. (b) The AGT dual conformal $n + 3$ -point block of the Toda field theory.

2.1.1 The conformal blocks

We consider the $(n + 3)$ -point correlator involving the insertion of chiral primary fields with momenta

$$P_s = \frac{1}{b}(p_0, k_0, k_1 \dots k_n, p_{n+1}) \quad (2.9)$$

and vertices $V_{P_s} = e^{(Q-2P_s)\phi}$. We denote by ²

$$\mathcal{F}_{p_0}^{k_0} \dots \mathcal{F}_{p_{n+1}}^{k_n}(x_i) = \langle p_0 | V_{k_0}(x_0) \dots V_{k_n}(x_n) | p_{n+1} \rangle_{p_1 \dots p_n} = \infty \frac{k_0}{p_0} \frac{k_1}{p_1} \dots \frac{k_n}{p_n} \frac{0}{p_{n+1}} \quad (2.10)$$

the corresponding conformal block with $p_1, p_2, \dots p_n$ specifying the intermediate momenta. The dimensions of the operators are related to momenta by

$$\Delta_{p_i} = \frac{Q^2}{4} - \frac{p_i^2}{b^2} \quad , \quad \Delta_{k_i} = \frac{Q^2}{4} - \frac{k_i^2}{b^2} \quad (2.11)$$

The conformal block $\mathcal{F}_{p_0}^{k_0} \dots \mathcal{F}_{p_{n+1}}^{k_n}$ is related via the AGT duality to the instanton partition function of a $U(2)^n$ quiver gauge theory. The gauge theory is specified by the scalar vevs a_{iu} , with $u = 1, 2$ a gauge index, $i = 0, \dots n + 1$ labelling the quiver nodes, the gauge couplings q_i and the Ω -background parameters ϵ_1, ϵ_2 . They are related to the CFT variables via the AGT dictionary

$$p_i = \frac{a_{i1} - a_{i2}}{2} \quad , \quad b \frac{Q}{2} - k_i = \sum_{u=1}^2 \frac{a_{i+1,u} - a_{i,u}}{2} \quad , \quad \frac{x_i}{x_{i-1}} = q_i \quad , \quad \epsilon_1 = b^2 \quad , \quad \epsilon_2 = 1 \quad (2.12)$$

Finally the conformal block is related to the instanton partition function. The latter can be written as a sum over a set of $2n$ Young tableaux $Y = \{Y_{iu}\}$, $i = 1, \dots n$, while $Y_{0u} = Y_{n+1,u} = \emptyset$ (see Appendix A for details). Translated into the CFT variables, the conformal block can be written as

$$\mathcal{F}_{p_0}^{k_0} \dots \mathcal{F}_{p_{n+1}}^{k_n}(x_i) = \prod_{i=0}^n x_i^{\Delta_{p_i} - \Delta_{p_{i+1}} - \Delta_{k_i}} \prod_{\substack{i,j=0 \\ i < j}}^2 \left(1 - \frac{x_j}{x_i} \right)^{\frac{(2k_i - bQ)(2k_j + bQ)}{2b^2}} Z_{\text{inst } p_0}^{k_0} \dots \mathcal{F}_{p_{n+1}}^{k_n}(x_i) \quad (2.13)$$

²As usual, we define $\langle p_0 | = \lim_{z_{-1} \rightarrow \infty} \langle 0 | z_{-1}^{2\Delta_{p_0}} V_{p_0}(z_{-1})$ and $| p_{n+1} \rangle = V_{p_{n+1}}(0) | 0 \rangle$

with

$$Z_{\text{inst}p_0}^{k_0} \dots_{p_{n+1}}^{k_n}(x_i) = \sum_Y \prod_{i=0}^n x_i^{|Y_i| - |Y_{i+1}|} \frac{\prod_{u,v}^2 \mathcal{Z}_{Y_{iu}, Y_{i+1}, v}(k_i + (-)^u p_i - (-)^v p_{i+1} - b \frac{Q}{2})}{\mathcal{Z}_{Y_{i1}, Y_{i2}}(2p_i) \mathcal{Z}_{Y_{i2}, Y_{i1}}(-2p_i)} \quad (2.14)$$

and

$$\mathcal{Z}_{Y_u, W_v}(y) = \prod_{i,j \in Y_u} [y + \epsilon_1(i - k'_{jv}) - \epsilon_2(j - k_{iu} - 1)] \prod_{i,j \in W_v} [y - \epsilon_1(i - k'_{ju} - 1) + \epsilon_2(j - k_{iv})] \quad (2.15)$$

Here i, j run over the columns and rows of the tableau and k_{iu}, k'_{ju} denote the height and lengths of the columns and rows of the tableau Y_u .

The first product in (2.13) in front of Z_{inst} can be viewed as the tree level contribution to the quiver partition function. The second one subtract the U(1) part. In principle one could also include a one-loop contribution but since it is z_i -independent, it is irrelevant for our analysis here. We notice that neither z_0 , that accounts for the freedom of an overall rescaling of the vertex field locations, nor the gauge theory center of mass $\sum_{i,u} a_{iu}$ are determined by the AGT dictionary. They can be fixed at will.

2.1.2 The BPZ equation

We are interested in the conformal blocks involving the insertion of a degenerate field. Degenerate fields correspond to primaries with momenta

$$\frac{k_{mm'}}{b} \equiv \frac{1}{2b}m + \frac{b}{2}m' \quad (2.16)$$

where m, m' are positive integers. To each degenerate field one can associate a null state with vertex

$$V_{\text{null}, mm'} = \mathbf{L}_{mm'} V_{\alpha_{mm'}} \quad (2.17)$$

where $\mathbf{L}_{mm'}$ is some polynomial of the Virasoro generators at level $N = mm'$. A correlator involving the insertion of a null state vanishes and so the corresponding conformal block will satisfy a differential equation of order mm' .

In order to get a second order differential equation, we consider the descendant at level 2 of $V_{k_{12}}$

$$V_{\text{null}, 12} = (L_{-1}^2 + b^2 L_{-2}) V_{k_{12}} \quad k_{12} = \frac{1}{2} + b^2 \quad \Delta_{12} = -\frac{1}{2} - \frac{3b^2}{4} \quad (2.18)$$

An important simplification comes from the fact that the OPE of a degenerated field $V_{k_{12}}$ with a primary of momentum p/b , produces only two operators with momenta [31]

$$\frac{p^\alpha}{b} = \frac{p}{b} + \alpha \frac{b}{2} \quad (2.19)$$

and $\alpha = \pm$. Consequently we denote by

$$\Psi_{i\alpha}(z_s) = \mathcal{F}_{p_0}^{k_0} \dots_{p_i}^{-\alpha} \dots_{p_i}^{k_{12}} \dots_{p_i}^{k_n} \dots_{p_{n+1}}(z_s) \quad (2.20)$$

the degenerate conformal block obtained by the insertion of a degenerate field at the i^{th} position, i.e. $k_i = k_{12}$. The function $\Psi_{i\alpha}(z_s)$ satisfies the BPZ equation

$$\left[\partial_{z_i}^2 + b^2 \sum_{s \neq i}^{n+3} \left(\frac{\Delta_{P_s}}{(z_i - z_s)^2} + \frac{1}{z_i - z_s} \partial_{z_s} \right) \right] \Psi_{i\alpha}(z_s) = 0 \quad (2.21)$$

Different choices of i correspond to different orderings of the vertex insertions and provide a different solution of the same equation. On the other hand, given “ i ”, $\alpha = \pm$ labels the two independent solutions of the differential equation.

To evaluate (2.21) one should first use the Ward identities

$$\sum_{s=1}^{n+3} \partial_{z_s} = \sum_{s=1}^{n+3} (z_s \partial_{z_s} + \Delta_{P_s}) = \sum_{s=1}^{n+3} (z_s^2 \partial_{z_s} + 2z_s \Delta_{P_s}) = 0 \quad (2.22)$$

to express ∂_{z_s} for three of the insertions in terms of the remaining ones, and then set the three positions to $\infty, 1, 0$. The resulting equation involves derivatives with respect to z and the remaining x_s . To reduce it to an ordinary differential equation in the variable z , one takes the Ω -background with $\epsilon_1 \neq 0, \epsilon_2 = 0$ [32]. This corresponds to the limit

$$b \rightarrow 0 \quad , \quad \Delta_s \rightarrow \infty \quad , \quad \delta_s = b^2 \Delta_s : \text{finite} \quad (2.23)$$

for $s \neq i$ while $b^2 \Delta_i \rightarrow 0$. Notice that in this limit p_s and k_s are finite. An explicit evaluation of the conformal blocks, denoted generically as $\Psi(z, x_s)$, shows that in this limit they behave as (see for example formula (2.45) for the 5-pt conformal block)

$$\Psi(z, x_s) = e^{\frac{f_0(x_s)}{b^2} + f_1(x_s, z) + \dots} \quad (2.24)$$

with the leading contribution independent of the z -variable³. Consequently $b^2 \partial_z$ -terms in (2.21) can be discarded while x_i -derivatives can be replaced by z -independent functions related to the Coulomb branch parameters of the quiver gauge theory with one node less. As a result the BPZ equation reduces to an ordinary differential equation of type (2.1) with $N = n + 2$ Fuchsian singularities.

2.2 Braiding and fusion relations

In this section we determine the relations between the different solutions $\Psi_{i\alpha}$ of the BPZ equations. Throughout this section we keep b finite.

To establish the braiding and fusion relations it is enough to consider the simplest $n = 1$ case, i.e. the case of a degenerate four-point conformal blocks. The corresponding BPZ equation in this case contains three Fuchsian singularities and therefore can be explicitly solved in terms of hypergeometric functions. The BPZ equation reads

$$\left(b^{-2} \partial_z^2 + \frac{1-2z}{z(z-1)} \partial_z + \frac{\Delta_{p_0} - \Delta_k - \Delta_{12} - \Delta_{p_2}}{z(z-1)} + \frac{\Delta_{p_2}}{z^2} + \frac{\Delta_k}{(z-1)^2} \right) \Psi(z) = 0 \quad (2.25)$$

³This is not surprising since the dimension $\Delta_{k_{12}}$ of the degenerate field is much smaller than the dimensions of the remaining operators, so its insertion in the four-point correlator cannot modify significantly its leading behaviour.

This equation can be brought to the standard hypergeometric form by writing

$$\Psi(z) = (1-z)^{\left(\frac{1}{2}+k+\frac{b^2}{2}\right)} z^{\left(\frac{1}{2}+p_2+\frac{b^2}{2}\right)} F(z) \quad (2.26)$$

with $F(z)$ a hypergeometric function. There are six natural choices for $F(z)$ denoted by w_i in [54]. The six functions w_i can be written as (up to constants)

$$\begin{aligned} \Psi_\alpha^R(z) &= (1-z)^{\frac{1}{2}+k+\frac{b^2}{2}} z^{\frac{1}{2}+\alpha p_2+\frac{b^2}{2}} {}_2F_1\left(\frac{1}{2}+k+p_0+\alpha p_2, \frac{1}{2}+k-p_0+\alpha p_2, 1+2\alpha p_2; z\right) \quad (2.27) \\ \tilde{\Psi}_\alpha^L(z) &= (z-1)^{\frac{1}{2}+\alpha k+\frac{b^2}{2}} z^{\frac{1}{2}+p_2+\frac{b^2}{2}} {}_2F_1\left(\frac{1}{2}+p_0+p_2+\alpha k, \frac{1}{2}-p_0+p_2+\alpha k, 1+2\alpha k, 1-z\right) \\ \Psi_\alpha^L(z) &= (z-1)^{\frac{1}{2}+k+\frac{b^2}{2}} z^{\alpha p_0-k+\frac{b^2}{2}} {}_2F_1\left(\frac{1}{2}+k+p_2-\alpha p_0, \frac{1}{2}+k-p_2-\alpha p_0, 1-2\alpha p_0; \frac{1}{z}\right) \end{aligned}$$

with $\alpha = \pm$. In addition, we introduce a fourth pair of solutions

$$\tilde{\Psi}_\alpha^R(z) = (-)^{\frac{1}{2}+\alpha k+\frac{b^2}{2}} \tilde{\Psi}_\alpha^L(z) \quad (2.28)$$

In graphical notation

$$\begin{aligned} \Psi_\alpha^R(z) &= \frac{\begin{array}{c} k \quad k_{12} \\ | \quad | \\ 1 \quad z \\ \hline p_0 \quad p_2^{-\alpha} \quad p_2 \end{array}}{\quad}, & \Psi_\alpha^L(z) &= \frac{\begin{array}{c} k_{12} \quad k \\ | \quad | \\ z \quad 1 \\ \hline p_0 \quad p_0^\alpha \quad p_2 \end{array}}{\quad}, \\ \tilde{\Psi}_\alpha^R(z) &= \frac{\begin{array}{c} k \quad k_{12} \\ \diagdown \quad \diagup \\ 1 \quad z \\ | \\ k_\alpha \\ \hline p_0 \quad p_2 \end{array}}{\quad}, & \tilde{\Psi}_\alpha^L(z) &= \frac{\begin{array}{c} k_{12} \quad k \\ \diagup \quad \diagdown \\ z \quad 1 \\ | \\ k_\alpha \\ \hline p_0 \quad p_2 \end{array}}{\quad} \quad (2.29) \end{aligned}$$

The four pairs of solutions are related via the hypergeometric identities, so which one to use depends on our convenience. In particular, using the hypergeometric identities (B.1) one finds the braiding and fusion relations

$$\begin{aligned} \frac{\begin{array}{c} k_{12} \quad k \\ | \quad | \\ z \quad 1 \\ \hline p_0 \quad p_0^\alpha \quad p_2 \end{array}}{\quad} &= \sum_\alpha B_{\alpha\alpha'} \begin{array}{c} [k_{12} \quad k] \\ [p_0 \quad p_2] \\ | \quad | \\ 1 \quad z \\ \hline p_0 \quad p_2^{-\alpha} \quad p_2 \end{array} \\ \frac{\begin{array}{c} k_{12} \quad k \\ | \quad | \\ z \quad 1 \\ \hline p_0 \quad p_0^\alpha \quad p_2 \end{array}}{\quad} &= \sum_\alpha F_{\alpha\alpha'} \begin{array}{c} [k_{12} \quad k] \\ [p_0 \quad p_2] \\ \diagdown \quad \diagup \\ z \quad 1 \\ | \\ k_{\alpha'} \\ \hline p_0 \quad p_2 \end{array} \\ \frac{\begin{array}{c} k \quad k_{12} \\ \diagdown \quad \diagup \\ 1 \quad z \\ | \\ k_{\alpha'} \\ \hline p_0 \quad p_2 \end{array}}{\quad} &= \sum_{\alpha'} F_{\alpha\alpha'}^{-1} \begin{array}{c} [k \quad k_{12}] \\ [p_0 \quad p_2] \\ | \quad | \\ 1 \quad z \\ \hline p_0 \quad p_2^{-\alpha'} \quad p_2 \end{array} \quad (2.30) \end{aligned}$$

with braiding and fusion matrices

$$\begin{aligned}
B_{\alpha\alpha'} \begin{bmatrix} k_{12} & k \\ p_0 & p_2 \end{bmatrix} &= \frac{e^{i\pi(\alpha p_0 - \alpha' p_2 + \frac{b^2}{2})} \Gamma(1 - 2\alpha p_0) \Gamma(-2\alpha' p_2)}{\Gamma(\frac{1}{2} - \alpha p_0 - \alpha' p_2 + k) \Gamma(\frac{1}{2} - \alpha p_0 - \alpha' p_2 - k)} \\
F_{\alpha\alpha'} \begin{bmatrix} k_{12} & k \\ p_0 & p_2 \end{bmatrix} &= \frac{\Gamma(1 - 2\alpha p_0) \Gamma(2\alpha' k)}{\Gamma(\frac{1}{2} - \alpha p_0 + \alpha' k + p_2) \Gamma(\frac{1}{2} - \alpha p_0 + \alpha' k - p_2)} \\
F_{\alpha\alpha'}^{-1} \begin{bmatrix} k & k_{12} \\ p_0 & p_2 \end{bmatrix} &= \frac{\Gamma(1 - 2\alpha k) \Gamma(-2\alpha' p_2)}{\Gamma(\frac{1}{2} - \alpha k + p_0 - \alpha' p_2) \Gamma(\frac{1}{2} - \alpha k - p_0 - \alpha' p_2)} \tag{2.31}
\end{aligned}$$

The crucial observation [55, 56] is that the braiding and fusion moves are local operations, independent of the positions of the vertex insertions, so they are still valid if we extend the conformal block by adding any number of vertices to the left or right of the diagram. In this way, one can derive exact connection formulae relating the asymptotics near any two singularities (different point insertions of the degenerate vertex) for a differential equation with an arbitrary number of Fuchsian singularities !

Confluent limits where two Fuchsian singularities collide can be studied in a similar way. The limit is obtained by taking for example k , p_2 and z large, keeping finite the combinations $c = k + p_2$ and $\hat{z} = z/(2k)$. In this limit, the conformal blocks reduce to

$$\begin{aligned}
\Psi_{\alpha}^{L,\text{conf}}(\hat{z}) &= z^{\frac{1}{2} + p_0 + b^2} e^{-\frac{1}{2z}} {}_1F_1\left(\frac{1}{2} + c - \alpha p_0, 1 - 2\alpha p_0, \frac{1}{z}\right) \\
\Psi_{\alpha}^{R,\text{conf}}(\hat{z}) &= z^{1 + \alpha c + b^2} e^{-\frac{\alpha}{2z}} {}_2F_0\left(\frac{1}{2} + \alpha c + p_0, \frac{1}{2} + \alpha c - p_0, -\alpha \hat{z}\right) \tag{2.32}
\end{aligned}$$

that are related to each other by the braiding relation

$$\Psi_{\alpha}^{L,\text{conf}}(\hat{z}) = \sum_{\alpha'} B_{\alpha\alpha'}^{\text{conf}} \begin{bmatrix} k_{12} & c \\ p_0 & c \end{bmatrix} \Psi_{\alpha'}^{R,\text{conf}}(\hat{z}) \tag{2.33}$$

with

$$B_{\alpha\alpha'}^{\text{conf}} \begin{bmatrix} k_{12} & c \\ p_0 & c \end{bmatrix} = e^{-i\pi\delta_{1\alpha''}(\frac{1}{2} + c - \alpha' p_0)} \frac{\Gamma(1 - 2\alpha' p_0)}{\Gamma(\frac{1}{2} - \alpha' p_0 - \alpha'' c)} \tag{2.34}$$

We remark that the second line of (2.32) is understood as a series expansion around $\hat{z} \approx 0$, since the ${}_2F_0$ series does not converge. In this paper we focus on the NS limit (2.23), so we can discard the b^2 -terms in all formulas above and the ∂_z -term in (2.25).

2.3 Five-point conformal block

Next we consider the $n = 2$ case (degenerate five-point conformal block) leading to a differential equation with four Fuchsian singularities (Heun equation). The connection matrices in this case have been obtained in section 3.1 of [27]. They have been also related to the Q-matrix of integrable systems in [28–30].

Taking the singularities at

$$z_0 = \infty \quad , \quad z_1 = 1 \quad , \quad z_2 = z \quad , \quad z_3 = x \quad , \quad z_4 = 0 \tag{2.35}$$

the Ward identities (2.22) can be solved as

$$\partial_{z_0} = 0 \quad , \quad \partial_{z_1} = \Delta - z\partial_z - x\partial_x \quad , \quad \partial_{z_4} = (z-1)\partial_z + (x-1)\partial_x - \Delta \tag{2.36}$$

with

$$\Delta = \Delta_{p_0} - \Delta_{p_3} - \Delta_{k_0} - \Delta_k - \Delta_{12} \quad (2.37)$$

Plugging this into (2.21), taking the limit (2.23) and using (2.24), the BPZ equation takes the normal form (2.1) with

$$Q(z, x) = \frac{\delta_{p_3}}{z^2} + \frac{\delta_{k_0}}{(z-1)^2} + \frac{\delta_k}{(z-x)^2} + \frac{\delta}{z(z-1)} + \frac{(1-x)U}{z(z-1)(z-x)} \quad (2.38)$$

and

$$U(x) = \lim_{b \rightarrow 0} b^2 x \partial_x \ln \Psi(z, x) \quad , \quad \delta = \delta_{p_0} - \delta_{p_3} - \delta_{k_0} - \delta_k \quad (2.39)$$

The first formula in (2.39) is nothing but the quantum version of the Matone relation [57, 58].

Again there are several basis of solutions one can use depending on the position of the degenerated field insertion. We will mainly use the following three representations [27]

$$\begin{aligned} \Psi_\alpha(z, x) &= \frac{\begin{array}{c} k_0 \\ \diagdown \\ 1 \\ \diagup \\ k_{0\alpha} \end{array} \begin{array}{c} k_{12} \\ \diagup \\ z \\ \diagdown \\ \end{array} \begin{array}{c} k \\ | \\ x \end{array}}{\begin{array}{c} p_0 \\ p_2 \\ p_3 \end{array}} = \sum_{\alpha'} F_{\alpha\alpha'}^{-1} \begin{array}{c} [k_0 \quad k_{12}] \\ p_0 \quad p_2 \end{array} \frac{\begin{array}{c} k_0 \\ | \\ 1 \end{array} \begin{array}{c} k_{12} \\ \vdots \\ z \end{array} \begin{array}{c} k \\ | \\ x \end{array}}{\begin{array}{c} p_0 \\ p_2^{-\alpha'} \\ p_2 \\ p_3 \end{array}} \\ &= \sum_{\alpha', \alpha''} F_{\alpha\alpha'}^{-1} \begin{array}{c} [k_0 \quad k_{12}] \\ p_0 \quad p_2 \end{array} B_{\alpha'\alpha''} \begin{array}{c} [k_{12} \quad k] \\ p_2^{-\alpha'} \quad p_3 \end{array} \frac{\begin{array}{c} k_0 \\ | \\ 1 \end{array} \begin{array}{c} k \\ | \\ x \end{array} \begin{array}{c} k_{12} \\ \vdots \\ z \end{array}}{\begin{array}{c} p_0 \\ p_2^{-\alpha'} \\ p_3^{-\alpha''} \\ p_3 \end{array}} \end{aligned} \quad (2.40)$$

To compute the asymptotics we use the degenerate three point correlator

$$\begin{array}{c} P \\ \diagdown \\ z_s \\ \diagup \\ P^\alpha \end{array} \begin{array}{c} k_{12} \\ \diagup \\ z \\ \diagdown \\ \end{array} \approx (z - z_s)^{\Delta_{P^\alpha} - \Delta_P - \Delta_{12}} \approx (z - z_s)^{\frac{1}{2} - \alpha P} \quad (2.41)$$

Here and below we will always discard b^2 -terms since we are interested in the $b \rightarrow 0$ limit. Plugging (2.41) into the left and right hand sides of (2.40) one finds the asymptotics of the solutions at the two boundaries

$$\begin{aligned} \Psi_\alpha(z, x) &\underset{z \rightarrow 1}{\approx} f_\alpha(z) \frac{\begin{array}{c} k_{0\alpha} \\ | \\ 1 \end{array} \begin{array}{c} k \\ | \\ x \end{array}}{\begin{array}{c} p_0 \\ p_2 \\ p_3 \end{array}} \\ \Psi_\alpha(z, x) &\underset{z \rightarrow 0}{\approx} \sum_{\alpha', \alpha''} \tilde{f}_{\alpha''}(z) F_{\alpha\alpha'}^{-1} \begin{array}{c} [k_0 \quad k_{12}] \\ p_0 \quad p_2 \end{array} B_{\alpha'\alpha''} \begin{array}{c} [k_{12} \quad k] \\ p_2^{-\alpha'} \quad p_3 \end{array} \frac{\begin{array}{c} k_0 \\ | \\ 1 \end{array} \begin{array}{c} k \\ | \\ x \end{array}}{\begin{array}{c} p_0 \\ p_2^{-\alpha'} \\ p_3^{-\alpha''} \end{array}} \end{aligned} \quad (2.42)$$

with

$$f_\alpha(z) = (1-z)^{\frac{1}{2} - \alpha k_0} \quad , \quad \tilde{f}_{\alpha''}(z) = z^{\frac{1}{2} + \alpha'' p_3} \quad (2.43)$$

The conformal blocks appearing in right hand sides of (2.40) can be written as instanton sums. For example

$$\begin{aligned} \mathcal{F}_{p_0}^{k_0}{}_{p_2^{-\alpha'}}{}^{k_{12}}{}_{p_2}{}^k{}_{p_3}(1, z, x) &= x^{\Delta_{p_2} - \Delta_{p_3} - \Delta_{12}} (1-x)^{\frac{(2k_0 - bQ)(2k_2 + bQ)}{2b^2}} \\ &\times z^{\frac{1}{2} + \alpha' p_2 + \frac{b^2}{2}} (1-z)^{\frac{(2k_0 - bQ)(1+b^2)}{b^2}} \left(1 - \frac{x}{z}\right)^{\frac{1}{2} + k + \frac{b^2}{2}} Z_{\text{inst} p_0}^{k_0}{}_{p_2^{-\alpha'}}{}^{k_{12}}{}_{p_2}{}^k{}_{p_3}(1, z, x) \end{aligned} \quad (2.44)$$

In applications to gravity, the terms in the first line can be discarded since they do not depend on z . The instanton part can be computed from (2.14) order by order in $q_1 = z$, $q_2 = x/z$. Up to one instanton order one finds

$$\begin{aligned} \ln \mathcal{F}_{p_0}^{k_0}{}_{p_2^{-\alpha'}}{}^{k_{12}}{}_{p_2}{}^k{}_{p_3}(1, z, x) &\approx \frac{x\rho_1\rho_2}{8b^2(1-4p_2^2)} + \left(\frac{p_3^2 - p_2^2}{b^2} - 1\right) \log x \\ &+ \left(\frac{1}{2} + \alpha' p_2\right) \log z - \frac{z\rho_1}{4(2\alpha' p_2 + 1)} + \frac{x\rho_2}{4z(2\alpha' p_2 - 1)} - x \frac{4\rho_1\alpha' p_2 + 8\rho_2 p_2^2 + \rho_1\rho_2 - 2\rho_2 - 2\rho_1}{8(1-2\alpha' p_2)(1-4p_2^2)} + \dots \end{aligned} \quad (2.45)$$

with

$$\rho_1 = 1 + 4p_0^2 - 4k_0^2 - 4p_2^2 \quad , \quad \rho_2 = 1 + 4p_3^2 - 4k^2 - 4p_2^2 \quad (2.46)$$

We notice that the terms in the first line of (2.45) grow as b^{-2} in the NS limit and depend only on x consistently with (2.24). For applications to gravity, we will mainly be interested on the limit $x, z \rightarrow 0$ keeping $\frac{x}{z}$ finite. In this limit, an explicit evaluation of the first few instanton corrections shows that Z_{inst} can be resummed into a hypergeometric function leading to

$$\mathcal{F}_{p_0}^{k_0}{}_{p_2^{-\alpha'}}{}^{k_{12}}{}_{p_2}{}^k{}_{p_3} \underset{z \rightarrow 0}{\sim} z^{\frac{1}{2} + \alpha' p_2} \left(1 - \frac{x}{z}\right)^{\frac{1}{2} + k} {}_2F_1\left(\frac{1}{2} + k + p_3 - \alpha' p_2, \frac{1}{2} + k - p_3 - \alpha' p_2, 1 - 2\alpha' p_2, \frac{x}{z}\right) \quad (2.47)$$

As before, one can consider the confluent limit of the five-point conformal block. This limit is obtained by taking k, p_3 and z large but $c = k + p_3$ and $\hat{z} = z/(2k)$ finite. Combining (2.40) and (2.47) one finds

$$\Psi_\alpha(\hat{z}) \underset{\hat{z} \rightarrow 0}{\approx} \sum_{\alpha'} \hat{z}^{\frac{1}{2} + \alpha' p_2} e^{-\frac{x}{2\hat{z}}} F_{\alpha\alpha'}^{-1} \left[\begin{matrix} k_0 & k_{12} \\ p_0 & p_2 \end{matrix} \right] {}_1F_1\left(\frac{1}{2} + c - \alpha' p_2, 1 - 2\alpha' p_2, \frac{x}{\hat{z}}\right) \quad (2.48)$$

On the other hand in the limit where $\hat{z} \ll x$ using the asymptotic expansion of the hypergeometric function one finds

$$\begin{aligned} \Psi_\alpha(\hat{z}) \underset{\hat{z} \rightarrow 0}{\approx} \sum_{\alpha', \alpha''} x^{\frac{1}{2} + \alpha' p_2} \left(\frac{\hat{z}}{x}\right)^{1 + \alpha'' c} e^{-\alpha'' \frac{x}{2\hat{z}}} F_{\alpha\alpha'}^{-1} \left[\begin{matrix} k_0 & k_{12} \\ p_0 & p_2 \end{matrix} \right] B_{\alpha'\alpha''}^{\text{conf}} \left[\begin{matrix} k_{12} \\ p_2 \end{matrix} \middle| c \right] \\ \times {}_2F_0\left(\frac{1}{2} + \alpha'' c + p_0, \frac{1}{2} + \alpha'' c - p_0, -\alpha'' \frac{\hat{z}}{x}\right) \end{aligned} \quad (2.49)$$

with

$$B_{\alpha'\alpha''}^{\text{conf}} \left[\begin{matrix} k_{12} \\ p_2 \end{matrix} \middle| c \right] = e^{-i\pi\delta_{1\alpha''}} \frac{\Gamma(1 - 2\alpha' p_2)}{\Gamma\left(\frac{1}{2} - \alpha' p_2 - \alpha'' c\right)} \quad (2.50)$$

2.4 BPZ equation from the quantum deformed SW curve

The BPZ equation can alternatively be obtained as a quantum deformation of the geometry describing the $U(2)$ gauge theory with four fundamental hypermultiplets resulting from omitting the node associated to the "degenerate" insertion, let us say node 1. In the limit $\epsilon_1 \rightarrow 0$ and $\epsilon_2 = \hbar$ the dynamics of this theory is described by the difference equation [34, 59]

$$y(v)y(v-\hbar) - P_2(v)y(v-\hbar) + qP_0(v-\hbar)P_3(v) = 0 \quad (2.51)$$

with

$$\begin{aligned} P_0(v) &= (v - a_{01})(v - a_{02}) \quad , \quad P_3(v) = (v - a_{31})(v - a_{32}); \\ P_2(v) &= v^2(1+q) - qv(a_{01}+a_{02}+a_{31}+a_{32}) - u + q[u + (a_{01}+a_{02})(a_{31}+a_{32}-\hbar) + a_{01}a_{02} + a_{31}a_{32}] \end{aligned} \quad (2.52)$$

Here a_{0u} and a_{3u} play the role of masses while u parametrizes the quantum Coulomb branch. $y(v)$ is the generating function of the expectation values $\langle \text{tr} \phi^J \rangle$ of the scalar ϕ in the vector multiplet

$$\frac{d \log y(v)}{dv} = \sum_{J=0}^{\infty} \frac{\langle \text{tr} \phi^J \rangle}{v^{J+1}} = \frac{2}{v} + \frac{2u}{v^3} + \dots \quad (2.53)$$

with

$$\langle \text{tr} 1 \rangle = 2 \quad , \quad \langle \text{tr} \phi \rangle = 0 \quad , \quad \langle \text{tr} \phi^2 \rangle = 2u \quad (2.54)$$

The form of $P_2(v)$ is determined by requiring that the expansion (2.53) satisfies the difference equation (2.51) at large v . In the $\hbar \rightarrow 0$ limit (2.51) reduces to the SW curve. This is why the case in which \hbar is kept finite is usually referred as the deformed (or "quantum") geometry. As in the undeformed case, the function $y(v)$ determines the differential

$$\lambda = v d \log y(v). \quad (2.55)$$

Solving (2.51) order by order in the limit of small q , one finds that at any instanton order the number of poles of λ is finite and is of the form $v_r = \pm\sqrt{u} + r\hbar$ with $r = 0, 1, \dots, k$ and k being the instanton order. The expectation value is given by

$$a = \oint_{\gamma} \frac{dv}{2\pi i} v \partial_v \log y(v) \quad (2.56)$$

where γ is the contour encircling all the poles around $\sqrt{u} + r\hbar$ with $r = 0, 1, \dots, k$. Then to find the gauge prepotential as a power series in q one can first invert (2.56) to express u as a function of a and make use of the relation [58]

$$\mathcal{F}^{U(2)} = - \int^a \frac{u(a, q')}{q'} dq' \quad (2.57)$$

There is an alternative, more direct way to find the function $u(a)$ (again based on (2.51)) by solving the equation [33]

$$\frac{qM(a+\hbar)}{P_2(a+\hbar) - \frac{qM(a+2\hbar)}{P_2(a+2\hbar)-\dots}} + \frac{qM(a)}{P_2(a-\hbar) - \frac{qM(a-\hbar)}{P_2(a-2\hbar)-\dots}} = P_2(a) \quad (2.58)$$

where, for the sake of simplicity, we used the notation

$$M(v) = P_0(v - \hbar)P_3(v) \quad (2.59)$$

Plugging u inside the P_2 function, one can solve (2.58) for $u(a)$ or $a(u)$ as a power series in q

$$\begin{aligned} u &= a^2 + c_1(a)q + c_2(a)q^2 + \dots \\ a &= \sqrt{u} + \tilde{c}_1(u)q + \tilde{c}_2(u)q^2 + \dots \end{aligned} \quad (2.60)$$

Equation (2.58) is a very convenient tool for the numerical evaluation of $u(a)$ for large values of q , where the series expansion is not useful. It is convenient to represent the y -function as the ratio

$$y(v) = \frac{P_0(v)Y(v)}{Y(v - \hbar)} \quad (2.61)$$

and denote the Fourier transform of $Y(v)$ as

$$F(z) = \sum_{v \in a_{1u} + \hbar\mathbb{Z}} Y(v)z^{-\frac{v}{\hbar}} \quad (2.62)$$

In these terms, the difference equation (2.51) translates into the differential equation [60]

$$\left[P_0 \left(-\hbar z \frac{d}{dz} \right) - P_2 \left(-\hbar z \frac{d}{dz} \right) z^{-1} + q P_3 \left(-\hbar z \frac{d}{dz} \right) z^{-2} \right] F(z) = 0 \quad (2.63)$$

This equation can be put into the normal form by writing

$$F(z) = z^{\frac{3}{2} - \frac{a_{21} + a_{22}}{2\hbar}} (1 - z)^{-1 - \frac{a_{01} + a_{02}}{2\hbar}} (z - q)^{-1 + \frac{a_{21} + a_{22}}{2\hbar}} \Psi(z) \quad (2.64)$$

Setting $\hbar = 1$ one finds (2.38) with $x = q$ and

$$\begin{aligned} \delta_{p_3} &= \frac{1}{4} - p_3^2 = \frac{1}{4} - \left(\frac{a_{31} - a_{32}}{2} \right)^2, & \delta_{p_0} &= \frac{1}{4} - p_0^2 = \frac{1}{4} - \left(\frac{a_{01} - a_{02}}{2} \right)^2 \\ \delta_{k_0} &= \frac{1}{4} - k_0^2 = \frac{1}{4} - \left(\frac{1 + a_{01} + a_{02}}{2} \right)^2, & \delta_k &= \frac{1}{4} - k^2 = \frac{1}{4} - \left(\frac{1 - a_{31} - a_{32}}{2} \right)^2 \\ U &= u - \frac{1}{4} + \delta_k + \delta_{p_3} + \frac{x(a_{01} + a_{02})(2 - a_{31} - a_{32})}{2(1 - x)} \approx u - \frac{1}{4} + \delta_k + \delta_{p_3} - \frac{x(1 - 2k_0)(1 + 2k)}{2(1 - x)} \end{aligned} \quad (2.65)$$

where in the right hand side of each equation we used the AGT dictionary (2.12) and set $k_1 = k_{12} \approx \frac{1}{2}$. This leads to the identification

$$\Psi(z) = \mathcal{F}_{p_0}^{k_0} p_2^{-\alpha} k_{12} k_{p_3}^k (1, z, x) = \frac{\begin{array}{c|c|c} k_0 & k_{12} & k \\ \hline 1 & z & x \end{array}}{\begin{array}{c|c|c} p_0 & p_2^{-\alpha} & p_2 & p_3 \end{array}} \quad (2.66)$$

2.5 Irregular singularities

Theories with a smaller number of hypermultiplets can be obtained from the generic $N_f = (2, 2)$ case by decoupling i.e. by sending some of the “masses” a_{0u} and/or a_{3u} to infinity keeping the products qa_{0u} or qa_{3u} finite. In addition, when sending a_{0u} to infinity one should also rescale $z \rightarrow z/a_{0u}$ and $Q \rightarrow Q/a_{0u}^2$. The general case is labeled by the two-value set $\{\nu_f\}$, $\nu_f = 0, 1$, $f = 1, \dots, 4$, that determines which flavours are decoupled. The Q -function is obtained by taking a_{0u} and/or a_{3u} large depending on whether $\nu_f = 0, 1$ and keeping the finite terms under the rescaling

$$Q_{2-\nu_1-\nu_2, 2-\nu_3-\nu_4} = \frac{Q_{22}}{a_{01}^{2\nu_1} a_{02}^{2\nu_2}} \quad , \quad z \rightarrow \frac{z}{a_{01}^{\nu_1} a_{02}^{\nu_2}} \quad , \quad x \rightarrow \frac{x}{a_{01}^{\nu_1} a_{02}^{\nu_2} a_{31}^{\nu_3} a_{32}^{\nu_4}} \quad (2.67)$$

The results are given by

$$\begin{aligned} Q_{2,2} &= \frac{\delta_{p_3}}{z^2} + \frac{\delta_{k_0}}{(z-1)^2} + \frac{\delta_k}{(z-x)^2} + \frac{\delta}{z(z-1)} + \frac{(1-x)U}{z(z-1)(z-x)} && \text{HE} \quad (2.68) \\ Q_{2,1} &= -\frac{x^2}{4z^4} + \frac{xc}{z^3} + \frac{u - \frac{1}{4} + x(\frac{1}{2} - k_0)}{(z-1)z^2} + \frac{\delta_{k_0}}{(z-1)^2 z} + \frac{\delta_{p_0}}{(z-1)z} && \text{CHE} \\ Q_{1,2} &= \frac{\delta_k}{(x-z)^2} + \frac{\delta_{p_3}}{z^2} - \frac{c'}{z} + \frac{u - \frac{1}{4} + \delta_k + \delta_{p_3} + x(k + \frac{1}{2})}{z(x-z)} - \frac{1}{4} && \text{CHE} \\ Q_{2,0} &= -\frac{x}{z^3} + \frac{u - \frac{1}{4}}{(z-1)z^2} + \frac{\delta_{k_0}}{(z-1)^2 z} + \frac{\delta_{p_0}}{(z-1)z} && \text{RCHE} \\ Q_{0,2} &= \frac{\delta_k}{(x-z)^2} + \frac{\delta_{p_3}}{z^2} - \frac{1}{z} + \frac{u - \frac{1}{4} + \delta_k + \delta_{p_3}}{z(x-z)} && \text{RCHE} \\ Q_{1,1} &= -\frac{x^2}{4z^4} + \frac{xc}{z^3} + \frac{\frac{1}{4} - u + \frac{x}{2}}{z^2} + \frac{c'}{z} - \frac{1}{4} && \text{DCHE} \\ Q_{1,0} &= -\frac{x}{z^3} + \frac{\frac{1}{4} - u}{z^2} + \frac{c'}{z} - \frac{1}{4} && \text{DRCHE} \\ Q_{0,1} &= -\frac{x^2}{4z^4} + \frac{xc}{z^3} + \frac{\frac{1}{4} - u}{z^2} - \frac{1}{z} && \text{DRCHE} \\ Q_{0,0} &= -\frac{x}{z^3} + \frac{\frac{1}{4} - u}{z^2} - \frac{1}{z} && \text{DRDCHE} \end{aligned}$$

with U given by (2.65) and

$$c = k + p_3 \quad , \quad c' = p_0 - k_0 \quad (2.69)$$

The differential equations with these Q function are known in the mathematical literature as Heun Equation (HE), Confluent Heun Equation (CHE), Reduced Confluent Heun Equation (RCHE), Double Confluent Heun Equation (DCHE), Double Reduced Confluent Heun Equation (DRCHE) and Double Reduced Double Confluent Heun Equation (DRDCHE) and have been recently studied with techniques similar to the ones employed here in [35].

3 Wave scattering: QNM's, cross sections and echoes

The general solution of the BPZ equation can be written as a linear combination of $\Psi_\alpha(z)$ given in (2.40). In this section we consider the scattering from $z = 0$ towards an inner

boundary at $z = 1$ characterised by some reflectivity properties. Near the two boundaries, we write

$$\Psi(z) \underset{z \rightarrow 1}{\approx} \sum_{\alpha} c_{\alpha} f_{\alpha}(z) \quad , \quad \Psi(z) \underset{z \rightarrow 0}{\approx} \sum_{\alpha''} \tilde{c}_{\alpha''} \tilde{f}_{\alpha''}(z) \quad (3.1)$$

with $f_{\alpha}(z)$ and $\tilde{f}_{\alpha''}(z)$, the local solutions (2.43) and

$$\tilde{c}_{\alpha''} = \sum_{\alpha} c_{\alpha} M_{\alpha\alpha''} \quad (3.2)$$

The connection matrix $M_{\alpha\alpha''}$ follows from (2.42) and reads [27]

$$M_{\alpha\alpha''}(x) = \sum_{\alpha'} F_{\alpha\alpha'}^{-1} \begin{bmatrix} k_0 & k_{12} \\ p_0 & p_2 \end{bmatrix} B_{\alpha'\alpha''} \begin{bmatrix} k_{12} & k \\ p_2 & p_3 \end{bmatrix} \frac{\mathcal{F}_{p_0}^{k_0} p_2^{-\alpha'} p_3^{-\alpha''}(1, x)}{\mathcal{F}_{p_0}^{k_0} p_2^k p_3(1, x)} \quad (3.3)$$

The reflectivity at the inner boundary is specified by the ratio

$$\mathfrak{R} = \frac{c_+}{c_-} \quad (3.4)$$

The physical properties of the membrane determine the dependence of \mathfrak{R} from the frequency, ω . A typical choice is $\mathfrak{R} \sim e^{-i\omega x_0}$ with x_0 specifying the size of the ECO interior. For simplicity we locate the membrane exactly at the position of the singularity (the would be horizon).

We will mainly consider the two extreme cases: perfectly absorbing ($c_+ = 0$) and perfectly reflecting ($c_- = 0$) boundary conditions. The perfectly absorbing boundary conditions represent a BH horizon from which nothing can escape. The opposite case, represents a mirror like ECO able to reflect all incoming light. Motivated by this physical distinction we will refer to the two extreme choices of boundary conditions at $z = 1$ as

$$\begin{aligned} \text{Black hole :} & \quad c_+ = 0 \\ \text{Perfectly reflecting ECO :} & \quad c_- = 0 \end{aligned} \quad (3.5)$$

3.1 Absorption cross section, QNM's and superradiance

We first consider the scattering of a scalar wave in a geometry described by the BPZ wave equation. The asymptotic forms of the BPZ solutions near the two boundaries are given by (2.43). Alternatively one can introduce a tortoise like coordinate

$$z_* = \ln(1 - z) - \ln(z) \quad (3.6)$$

that maps the z -interval boundaries 0 and 1 to ∞ and $-\infty$ respectively. In these variables, the asymptotic behaviour at the two boundaries are planar waves

$$\frac{f_{\alpha}(z)}{\sqrt{z(1-z)}} \underset{z \rightarrow 1}{\sim} e^{-ak_0 z_*} \quad , \quad \frac{\tilde{f}_{\alpha''}(z)}{\sqrt{z(1-z)}} \underset{z \rightarrow 0}{\sim} e^{-\alpha'' p_3 z_*} \quad (3.7)$$

We will consequently refer to the $-$ and $+$ components for the indices α, α'' as ingoing and outgoing waves respectively. Using the equation of motion (2.1), it is easy to show that

the flux \mathfrak{f} carried by the wave is constant. Using the asymptotic forms (2.43) near the two boundaries $z \approx 0, 1$ and assuming k_0, p_3 to be purely imaginary, one finds

$$\mathfrak{f} = \text{Im} [\Psi^*(z)\Psi'(z)] = ik_0 (|c_+|^2 - |c_-|^2) = ip_3 (|\tilde{c}_+|^2 - |\tilde{c}_-|^2) \quad (3.8)$$

We refer to the contributions

$$\mathfrak{f}_{\text{in}} = -ip_3|\tilde{c}_-|^2 \quad , \quad \mathfrak{f}_{\text{out}} = -ip_3|\tilde{c}_+|^2 \quad , \quad \mathfrak{f}_{\text{abs}} = -ik_0|c_-|^2 \quad , \quad \mathfrak{f}_{\text{ref}} = -ik_0|c_+|^2 \quad (3.9)$$

as the incoming/outgoing and absorbed/reflected fluxes. We define the amplification factor

$$Z = \frac{\mathfrak{f}_{\text{out}}}{\mathfrak{f}_{\text{in}}} - 1 = \frac{k_0}{p_3|\tilde{c}_-|^2} (|c_+|^2 - |c_-|^2) \quad (3.10)$$

that measures how much the incoming wave is amplified by the gravity solution. Super-radiance takes place when $Z > 0$, i.e. when the ratio k_0/p_3 is positive for a black hole and negative for a perfectly reflecting ECO.

The absorption cross section is defined by the ratio

$$\sigma_{\text{abs}} = \frac{\mathfrak{f}_{\text{abs}}}{\mathfrak{f}_{\text{in}}} = \frac{k_0|c_-|^2}{p_3|\tilde{c}_-|^2} \quad (3.11)$$

The QNM's are defined as the poles of the absorption cross section σ_{abs} . They are located at the zeros of $\tilde{c}_- = 0$. The QNM frequencies are then defined as the solutions in x_{QNM} of the equation

$$\mathfrak{R} = \frac{c_+}{c_-} = -\frac{M_{--}(x_{\text{QNM}})}{M_{+-}(x_{\text{QNM}})} \quad (3.12)$$

for a given \mathfrak{R} . For example the QNM's for a BH and a perfectly reflecting ECO are defined by

$$\begin{aligned} \text{Black hole :} \quad & M_{--} = 0 \\ \text{Perfectly reflecting ECO :} \quad & M_{+-} = 0 \end{aligned} \quad (3.13)$$

In the limit $b \rightarrow 0$, the BH condition $M_{--} = 0$ can be written as

$$\begin{aligned} e^{\partial_{p_2} \ln \mathcal{F}_{p_0}^{k_0} p_2^k p_3(1,x)} &= -\frac{F_{-+}^{-1} \begin{bmatrix} k_0 & k_{12} \\ p_0 & p_2 \end{bmatrix} B_{+-} \begin{bmatrix} k_{12} & k \\ p_2^+ & p_3 \end{bmatrix}}{F_{--}^{-1} \begin{bmatrix} k_0 & k_{12} \\ p_0 & p_2 \end{bmatrix} B_{--} \begin{bmatrix} k_{12} & k \\ p_2^- & p_3 \end{bmatrix}} \\ &= e^{2\pi i p_2} \frac{\Gamma(-2p_2)^2}{\Gamma(2p_2)^2} \prod_{\alpha=\pm} \frac{\Gamma(\frac{1}{2}+k_0+\alpha p_0+p_2)\Gamma(\frac{1}{2}+\alpha k_2+p_3+p_2)}{\Gamma(\frac{1}{2}+k_0+\alpha p_0-p_2)\Gamma(\frac{1}{2}+\alpha k_2+p_3-p_2)} \end{aligned} \quad (3.14)$$

There are infinite solutions to this equation parametrized by an integer, the overtone number. The results for the ECO can be obtained from those of the black holes after sending $k_0 \rightarrow -k_0$.

3.2 Echoes

ECO's are distinguished from black holes by their echo response to perturbations. We refer the readers to [61] for a general discussion about echoes. The radial wave response of a geometry to a localized perturbation is typically described (after Fourier transforming from time to the ω domain), by a non homogenous equation of the form

$$\left[\frac{d^2}{dz^2} + Q(z) \right] \Phi(z) = S(z) \quad (3.15)$$

with $S(z)$ charactering the $t = 0$ initial conditions (the shape of the perturbation) localized somehow inside the interval $z \in [0, 1]$ and vanishing at the boundaries, i.e. $S(0) = S(1) = 0$. We look for solutions to (3.15) with boundary conditions

$$\Phi(z) \underset{z \rightarrow 1}{\sim} f_-(z) + \mathfrak{R} f_+(z) \quad , \quad \Phi(z) \underset{z \rightarrow 0}{\sim} \tilde{f}_+(z) \quad (3.16)$$

where $f_{\pm}(z)$, $\tilde{f}_{\pm}(z)$ are local solutions and \mathfrak{R} the reflection coefficient.

We denote by $\psi_{\text{in}}(z)$, $\psi_{\text{out}}(z)$ the solutions of the homogenous equation satisfying the boundary conditions

$$\begin{aligned} \psi_{\text{in}}(z) &\approx \begin{cases} f_-(z) & z \approx 1 \\ A_+ \tilde{f}_+(z) + A_- \tilde{f}_-(z) & z \approx 0 \end{cases} \\ \psi_{\text{out}}(z) &\approx \begin{cases} B_+ f_+(z) + B_- f_-(z) & z \approx 1 \\ \tilde{f}_+(z) & z \approx 0 \end{cases} \end{aligned} \quad (3.17)$$

with

$$A_{\alpha} = M_{-\alpha} \quad , \quad B_{\alpha} = (M^{-1})_{+\alpha} \quad (3.18)$$

and Jacobian

$$W = \det \begin{pmatrix} \psi_{\text{in}}(z) & \psi_{\text{out}}(z) \\ \psi'_{\text{in}}(z) & \psi'_{\text{out}}(z) \end{pmatrix} = 2k_0 B_+ = 2p_3 A_- \quad , \quad (3.19)$$

Finally the Green function $G(z, z')$, is defined by the equation

$$\left[\frac{d^2}{dz^2} + Q(z) \right] G(z, z') = \delta(z - z') \quad (3.20)$$

with boundary conditions (3.16) in the z -variable. The Green function can be written as

$$G(z, z') = \theta(z - z') \frac{\psi_{\text{in}}(z) \psi_{\text{out}}(z')}{W} + \theta(z' - z) \frac{\psi_{\text{in}}(z') \psi_{\text{out}}(z)}{W} + \frac{\mathcal{K}}{W} \psi_{\text{out}}(z') \psi_{\text{out}}(z) \quad (3.21)$$

with

$$\mathcal{K} = \frac{\mathfrak{R}}{B_+ - \mathfrak{R} B_-} \quad (3.22)$$

The solution near $z \approx 0$ can therefore be written as

$$\Phi(z) = \int dz' G(z, z') S(z') \underset{z \rightarrow 0}{\approx} \frac{\tilde{f}_+(z)}{W} (S_{\text{in}} + \mathcal{K} S_{\text{out}}) \quad (3.23)$$

with

$$S_{\text{in}} = \int \psi_{\text{in}}(z') S(z') dz' \quad , \quad S_{\text{out}} = \int \psi_{\text{out}}(z') S(z') dz' \quad (3.24)$$

For a black hole, $\mathfrak{R} = 0$ and therefore the echo \mathcal{K} -component is missing. For \mathfrak{R} small, a monochromatic wave will therefore produce a sequence of echoes proportional to $(\mathfrak{R}B_-/B_+)^n$ obtained by expanding the \mathcal{K} -signal for small \mathfrak{R} . On the other hand for a multi-chromatic signal, the Fourier transform of the echo signal will pick up either the poles of W or the poles of \mathcal{K} . The former gives an echo signal with frequencies of the underlying black hole QNM's, while the latter gives an echo signal with the characteristic QNM's of the ECO given by the solutions to (3.12).

4 Kerr-Newman geometries in four dimensions

In this section we compute the amplification factors and Love numbers for asymptotically flat Kerr-Newman BH's (or its associated ECO geometry) in four dimensions. The wave functions describing the radial and angular motion will be related to degenerate five point conformal blocks in the confluent limit where two Fuchsian singularities collide.

4.1 The wave equation

The Kerr-Newman (KN) metric describes a solution to the Einstein-Maxwell equations in four dimensions with mass \mathcal{M} , charge \mathcal{Q} and angular momentum $J = \mathcal{M}a_{\mathcal{J}}$. The line element of the KN metric in Boyer-Lindquist coordinates reads [62]

$$ds^2 = -\frac{\Delta_r}{\rho^2} (dt - a_{\mathcal{J}} \sin^2 \theta d\phi)^2 + \frac{\sin^2 \theta}{\rho^2} [a_{\mathcal{J}} dt - (r^2 + a_{\mathcal{J}}^2) d\phi]^2 + \frac{\rho^2 dr^2}{\Delta_r} + \rho^2 d\theta^2 \quad (4.1)$$

where

$$\Delta_r = (r - r_+)(r - r_-) = r^2 - 2\mathcal{M}r + \mathcal{Q}^2 + a_{\mathcal{J}}^2, \quad \rho^2 = r^2 + C^2 \cos^2 \theta, \quad (4.2)$$

These BH's possess two generally distinct horizons located at the zeroes of Δ_r

$$r_{\pm} = \mathcal{M} \pm \sqrt{\mathcal{M}^2 - a_{\mathcal{J}}^2 - \mathcal{Q}^2} \quad (4.3)$$

The wave-equation can be separated using the ansatz

$$\Phi(t, r, \chi, \phi) = e^{i(-\omega t + m\phi)} R(r) S(\chi) = e^{i(-\omega t + m\phi)} \frac{\widehat{R}(r) \widehat{S}(\chi)}{\sqrt{(1 - \chi^2) \Delta_r(r)}} \quad (4.4)$$

with $\chi = \cos \theta$. In terms of these variables, the wave equation splits into two ordinary differential equations of the form (2.1) describing angular and radial motion.

4.2 Angular motion

In terms of the variables

$$\Psi(z) = \widehat{S}(\chi), \quad \chi = \frac{2z}{x} - 1 \quad (4.5)$$

The wave equation describing the angular motion takes the form (2.1) with

$$Q_\chi(\chi) = \frac{(1 - \chi^2)(a_{\mathcal{J}}^2 \omega^2 \chi^2 + A) - m^2 + 1}{(1 - \chi^2)^2} \quad (4.6)$$

where A is the separation constant and m the azimuthal angular momentum of the incoming wave. Matching (4.6) against the BPZ Q -function

$$Q_{1,2}^{\text{angular}}(z) = \frac{\delta_{k_2}}{(x-z)^2} + \frac{\delta_{p_3}}{z^2} - \frac{c^\chi}{z} + \frac{u - \frac{1}{4} + \delta_k + \delta_{p_3} + x(k_2 + \frac{1}{2})}{z(x-z)} - \frac{1}{4} \quad (4.7)$$

one finds the gauge/gravity dictionary

$$x = q^\chi = 4a_{\mathcal{J}}\omega, \quad c^\chi = 0, \quad k_2^\chi = p_3^\chi = \frac{m}{2}, \quad u^\chi = A + \frac{1}{4} + a_{\mathcal{J}}^2 \omega^2 - 2a_{\mathcal{J}}\omega(m+1) \quad (4.8)$$

where the superscript χ is to distinguish the angular variables from the radial ones in the next section. In the gauge theory language regularity of the wavefunction $\hat{S}(\chi)$ at the boundaries $\chi = \pm 1$ requires the a -cycle quantization condition [24, 63]

$$p_2^\chi = a = \ell + \frac{1}{2} \quad (4.9)$$

with ℓ a non-negative integer. When $a_{\mathcal{J}}\omega = 0$, the gauge theory is free ($q = 0$) and the equation can be explicitly solved in terms of hypergeometric functions (spherical harmonics)

$$\Psi(\chi) \Big|_{a_{\mathcal{J}}\omega=0} \approx \sqrt{1 - \chi^2} P_{\ell m}(\chi) = \frac{\sqrt{1 - \chi^2}}{\Gamma(1 - m)} \left(\frac{1 + \chi}{1 - \chi} \right)^{\frac{m}{2}} {}_2F_1 \left(-\ell, \ell + 1, 1 - m; \frac{1 - \chi}{2} \right) \quad (4.10)$$

with m an integer with $\ell > |m|$. The separation constant becomes

$$A \Big|_{a_{\mathcal{J}}\omega=0} \approx \ell(\ell + 1) \quad (4.11)$$

In this limit, (2.60) and (4.8) yield $u^\chi \approx (a^\chi)^2 \approx (\ell + \frac{1}{2})^2$. Turning on the contribution of $a_{\mathcal{J}}\omega$, the relation between u and a is determined order by order in q^χ from the difference equation (2.58) with

$$\begin{aligned} P_0 &= -(v - a_{02}) \quad , \quad P_3 = (v - a_{31})(v - a_{32}) \\ P_2(v) &= v^2 - u^\chi + q^\chi(a_{02} + a_{31} + a_{32} - \hbar - v) \end{aligned} \quad (4.12)$$

Plugging the ansatz (2.60) for $u^\chi(a)$ into (2.58), solving for the expansion coefficients and using the gauge gravity dictionary (4.8) one finds

$$A = \ell(\ell + 1) + \frac{a_{\mathcal{J}}^2 \omega^2 (1 + 2m^2 - 2\ell(\ell + 1))}{4\ell(\ell + 1) - 3} + \dots \quad (4.13)$$

(4.13) reproduces the result (12) in [64] in the scalar case after the identification ${}_0\lambda_l^{m_\phi} = A + a_{\mathcal{J}}^2 \omega^2$.

4.3 Radial motion

The radial equation can be written in the (2.1) form with $\widehat{R}(r) = z^{-1}\Psi(z)$ and

$$Q_r(r) = \frac{\left(\omega(r^2 + a_{\mathcal{J}}^2) - a_{\mathcal{J}}m\right)^2 - \Delta_r \left(a_{\mathcal{J}}^2\omega^2 - 2a_{\mathcal{J}}\omega m + A + 1\right) + \frac{1}{4}\Delta_r'^2}{\Delta_r^2} \quad (4.14)$$

This function displays two Fuchsian singularities at the horizons $r = r_{\pm}$ and an irregular singularity at $r = \infty$. Using the coordinates

$$z = \frac{r_+ - r_-}{r - r_-} \quad (4.15)$$

we map the region $r \in [r_+, \infty]$ into the interval $z \in [0, 1]$ with 1, 0 corresponding to the outer horizon and infinity respectively. Plugging (4.15) into (2.5) one finds that $Q(z)$ has the form of (2.68)

$$Q_{2,1}^{\text{radial}}(z) = \frac{\delta_{p_0}}{z(z-1)} + \frac{\delta_{k_0}}{(z-1)^2z} + \frac{u - \frac{1}{4} + x\left(\frac{1}{2} - k_0\right)}{z^2(z-1)} + \frac{cx}{z^3} - \frac{x^2}{4z^4} \quad (4.16)$$

with gauge and gravity variables identified according to

$$\begin{aligned} x = q = 2i\omega(r_+ - r_-), \quad k_0 = iQ_{\text{grav}}, \quad p_0 = iQ_{\text{grav}} + 2i\omega\mathcal{M}, \quad c = -2i\omega\mathcal{M} \\ u = A + \frac{1}{4} - i\omega(r_+ - r_-)(1 - 2iQ_{\text{grav}}) - \omega^2(a_{\mathcal{J}}^2 + 4\mathcal{M}^2 + 2r_+^2) \end{aligned} \quad (4.17)$$

with

$$Q_{\text{grav}} = \frac{a_{\mathcal{J}}m - \omega(r_+^2 + a_{\mathcal{J}}^2)}{r_+ - r_-} = (\Omega m - \omega) \frac{r_+^2 + a_{\mathcal{J}}^2}{r_+ - r_-} \quad (4.18)$$

where Ω is coordinate angular velocity. The solutions $\Psi_{\alpha}(z)$ of this equation are given in (2.40). In particular, imposing ingoing boundary conditions at $z \approx 1$ one finds that $\Psi(z) \sim \Psi_-(z)$. Again the non trivial task is to determine the relation between u and p_2 . As before, this can be done using (2.58) where now

$$\begin{aligned} P_0(a) &= (v - a_{01})(v - a_{02}), \quad P_3 = -(v - a_{32}) \\ P_2(v) &= v^2 - u + q(a_{01} + a_{02} + a_{32} - v) \end{aligned} \quad (4.19)$$

Solving (2.58) now for $a(u)$ order by order in q , using the gauge gravity dictionary (4.17) and formula (4.13) for the separation constant one finds

$$p_2 = \ell + \frac{1}{2} + \Delta\nu = \ell + \frac{1}{2} + 2\omega^2 \frac{\mathcal{M}^2 [11 - 15\ell(\ell+1)] + Q^2 [3\ell(\ell+1) - 2]}{(2\ell+1)(4\ell(\ell+1) - 3)} + \dots \quad (4.20)$$

4.4 Love numbers

Love and dissipation numbers parametrize the deformability of a geometry under an external perturbation. They can be extracted from the asymptotic behaviour at infinity of a scalar wave propagating in the gravity background. For a static wave $\omega = 0$, they are typically defined as the ratio between the term falling as $r^{-\ell-1}$ and the one growing as r^{ℓ} representing the response and the source terms respectively. Here, we adopted the

same definition for a extremely slowly rotating wave. We expand the wave function in the intermediate region

$$r_+ - r_- \ll r \ll \omega^{-1} \quad (4.21)$$

According to the gauge gravity dictionary (4.15), (4.17), this region describes the regime where both gauge couplings z and $\frac{x}{z}$ of the quiver gauge theory are small, i.e.

$$z \approx \frac{r_+ - r_-}{r} \ll 1 \quad , \quad \frac{x}{z} \approx \omega r \ll 1 \quad (4.22)$$

The solution with ingoing boundary conditions at $z \approx 1$, it is proportional to $\Psi_-(z, x)$ in (2.40). From (2.48) one finds

$$\Psi_-(z) \underset{z \rightarrow 0}{\approx} e^{-\frac{x}{z}} \sum_{\alpha'} z^{\frac{1}{2} + \alpha' p_2} F_{-\alpha'}^{-1} \left[\begin{matrix} k_0 & k_{12} \\ p_0 & p_2 \end{matrix} \right]_1 F_1 \left(\frac{1}{2} + c - \alpha' p_2, 1 - 2\alpha' p_2, \frac{2x}{z} \right) \quad (4.23)$$

with

$$F_{-\alpha'}^{-1} \left[\begin{matrix} k_0 & k_{12} \\ p_0 & p_2 \end{matrix} \right] = \frac{\Gamma(1 + 2k_0) \Gamma(-2\alpha' p_2)}{\Gamma\left(\frac{1}{2} - \alpha' p_2 + k_0 + p_0\right) \Gamma\left(\frac{1}{2} - \alpha' p_2 + k_0 - p_0\right)} \quad (4.24)$$

Using (4.20) one can write

$$\Psi(z) \underset{z \rightarrow 0}{\sim} z^{-\ell} + z^{\ell+1+2\Delta\nu} \mathfrak{L}_0 [1 + \dots] \quad (4.25)$$

with dots denoting z and $\frac{x}{z}$ corrections. We define the dynamical Love L and dissipation number Θ as the real and imaginary parts of the ratio \mathfrak{L}_0 . The result is given by the remarkable simple formula

$$\mathfrak{L}_0 = L + i\Theta = \frac{F_{-+}^{-1} \left[\begin{matrix} k_0 & k_{12} \\ p_0 & p_2 \end{matrix} \right]}{F_{--}^{-1} \left[\begin{matrix} k_0 & k_{12} \\ p_0 & p_2 \end{matrix} \right]} = \frac{\Gamma(-2p_2) \Gamma\left(\frac{1}{2} + k_0 + p_0 + p_2\right) \Gamma\left(\frac{1}{2} + k_0 - p_0 + p_2\right)}{\Gamma(2p_2) \Gamma\left(\frac{1}{2} + k_0 + p_0 - p_2\right) \Gamma\left(\frac{1}{2} + k_0 - p_0 - p_2\right)} \quad (4.26)$$

with p_0, k_0 given by (4.17) and p_2 by (4.20). (4.26) reproduces the results for the Love and dissipation number of Kerr-Newman BH's in the ‘‘strictly’’ static limit $\omega = 0$. In this limit $p_0 = k_0 = iQ_{\text{grav}}$, $p_2 = \ell + \frac{1}{2}$, with ℓ almost (but not exactly) an integer, leading to

$$\mathfrak{L}_0 \approx \frac{\Gamma(-2\ell-1) \Gamma(\ell+1) \Gamma(\ell+1+2iQ_{\text{grav}})}{\Gamma(2\ell+1) \Gamma(-\ell) \Gamma(-\ell+2iQ_{\text{grav}})} = -\frac{iQ_{\text{grav}} (\ell)!^2}{(2\ell)! (2\ell+1)!} \prod_{n=1}^{\ell} (n^2 + 4Q_{\text{grav}}^2) \quad (4.27)$$

and reproducing (3.55) of [53] (or (113) of [27]). The analytic continuation in ℓ is required to avoid a mixing between $1/r$ corrections to the $z^{-\ell}$ and $z^{\ell+1}$ terms in (4.25) when ω is exactly zero. On the other hand when $\omega \neq 0$, there is no mixing and one could set $\ell \rightarrow \mathbb{N}$ from the very beginning. This produces a different result ! The two limits $\ell \rightarrow \mathbb{N}$ and $\omega \rightarrow 0$ do not commute. Indeed, taking $\ell \rightarrow \mathbb{N}$ first and then $\omega \rightarrow 0$ one finds⁴

$$\mathfrak{L}_0 \approx \frac{2Q_{\text{grav}} \omega \mathcal{M}(\ell)!^2}{\Delta\nu (2\ell)! (2\ell+1)!} \prod_{n=1}^{\ell} (n^2 + 4Q_{\text{grav}}^2) \quad (4.28)$$

⁴Here we use the $\Gamma(-n+x) \underset{x \rightarrow 0}{\approx} \frac{(-)^n}{n!x}$ when $n \in \mathbb{N}$.

with $\Delta\nu$ given by (4.20). This yields to a non-trivial divergent result for the Love number, suggesting a breaking of the perturbative expansion. The physical interpretation of this result is not clear to us.

Finally, we observe that the limit $\omega r \ll 1$ can be relaxed and the exact dependence on ωr of the ratio between the response and the source terms can be explicitly determined from (4.23). Indeed, the wave function at infinity can be written as

$$\Psi(z) \underset{z \rightarrow 0}{\sim} z^{-\ell} + z^{\ell+1+2\Delta\nu} \mathfrak{L}\left(\frac{x}{z}\right) [1 + O(z)] \quad (4.29)$$

with $\mathfrak{L}\left(\frac{x}{z}\right)$ a sort of ‘‘Love function’’ given by

$$\mathfrak{L}\left(\frac{x}{z}\right) = \mathfrak{L}_0 \frac{{}_1F_1\left(\frac{1}{2} + c - p_2, 1 - 2p_2, \frac{2x}{z}\right)}{{}_1F_1\left(\frac{1}{2} + c + p_2, 1 + 2p_2, \frac{2x}{z}\right)} \quad (4.30)$$

It would be interesting to see whether the first expansion coefficients of this function in powers of ωr can be detected in the GW signal.

4.5 Amplification factors

In this section we compute the amplification factors for the KN BH in the slowly rotating limit $\omega\mathcal{M} \ll 1$. The asymptotic solution at infinity is given by (2.49)

$$\Psi_-(z) \underset{z \rightarrow 0}{\sim} Y_- z^{1-c} e^{-\frac{x}{z}} + Y_+ z^{1+c} e^{\frac{x}{z}} \quad (4.31)$$

with

$$Y_{\alpha''} = \sum_{\alpha'} \mathcal{Y}_{\alpha'\alpha''} = \sum_{\alpha'} x^{\alpha'p_2 - \alpha''c} F_{-\alpha'}^{-1} \begin{bmatrix} k_0 & k_{12} \\ p_0 & p_2 \end{bmatrix} B_{\alpha'\alpha''}^{\text{conf}} \begin{bmatrix} k_{12} \\ p_2 \ c \end{bmatrix} \quad (4.32)$$

and F^{-1} , B^{conf} given by (4.24) and (2.50). Explicitly

$$\mathcal{Y}_{\alpha'\alpha''} = \frac{x^{\alpha'p_2 - \alpha''c} e^{-i\pi\delta_{1\alpha''}\left(\frac{1}{2} + c - \alpha'p_2\right)} \Gamma(1 + 2k_0) \Gamma(-2\alpha'p_2) \Gamma(1 - 2\alpha'p_2)}{\Gamma\left(\frac{1}{2} + k_0 + p_0 - \alpha'p_2\right) \Gamma\left(\frac{1}{2} + k_0 - p_0 - \alpha'p_2\right) \Gamma\left(\frac{1}{2} - \alpha''c - \alpha'p_2\right)} \quad (4.33)$$

In terms of these variables the amplification factor becomes

$$Z = 1 - \left| \frac{Y_+}{Y_-} \right|^2 = 1 - \left| \frac{\mathcal{Y}_{-+} + \mathcal{Y}_{++}}{\mathcal{Y}_{--} + \mathcal{Y}_{+-}} \right|^2 \quad (4.34)$$

In the slowly rotating limit $x \rightarrow 0$, $p_0 \approx k_0 \approx iQ_{\text{grav}}$, $c \approx 0$, $p_2 \approx \ell + \frac{1}{2}$ with ℓ almost (but not exactly) an integer. One finds

$$Z \approx 1 - \left| \frac{1 - \Sigma}{1 + \Sigma} \right|^2 \approx 4\text{Re}\Sigma \quad (4.35)$$

with

$$\Sigma = \frac{\mathcal{Y}_{+-}}{\mathcal{Y}_{--}} \approx - \frac{x^{2\ell+1} \Gamma(-2\ell)^2 \Gamma(\ell+1)^2 \Gamma(\ell+1+2iQ_{\text{grav}})}{\Gamma(-\ell)^2 \Gamma(2\ell+2)^2 \Gamma(-\ell+2iQ_{\text{grav}})} \quad (4.36)$$

leading to

$$Z \underset{x \rightarrow 0}{\approx} \frac{2|x|^{2\ell+1} Q_{\text{grav}} \ell!^4}{(2\ell)!^2 (2\ell+1)!^2} \prod_{n=1}^{\ell} (n^2 + 4Q_{\text{grav}}^2) \quad (4.37)$$

in agreement with formula (26) in [64]. Superradiance takes place when $\omega < m_\phi\Omega$. For the perfectly reflecting ECO, one finds the same result with $k_0 \rightarrow -k_0$, i.e. $Z \rightarrow -Z$. Superradiance in this case is obtained when $Q_{\text{grav}} < 0$ i.e. when $\omega > m_\phi\Omega$!

We notice that once again the order of the limits $\omega \rightarrow 0$ and $\ell \rightarrow \mathbb{N}$ is crucial in producing the result above. Indeed, considering the opposite limit when ℓ is a natural number and $\omega \rightarrow 0$ one finds a different result !

5 Solutions of the Einstein-Maxwell theory in five dimensions

The results in the last section can be easily generalized to BH in five dimensions, since as shown in [17, 18, 24, 63], they can be described by different confluent limits of the same ordinary differential equation.

5.1 The wave equation

We consider the Einstein-Maxwell theory with Lagrangian

$$16\pi G_5 \mathcal{L}_{5d} = \sqrt{g_{5d}} \left(R_{5d} - \frac{1}{4} F^2 \right) - \frac{1}{3\sqrt{3}} F \wedge F \wedge A \quad (5.1)$$

We set units such that the five dimensional Newton constant⁵ is $G_5 = 3\pi/4$. The general solution obtained in [65] is specified by four parameters, the mass \mathcal{M} , the charge \mathcal{Q} and two angular momentum parameters ℓ_1, ℓ_2 . The line element reads

$$ds^2 = -dt^2 - \frac{2\mathcal{Q}\omega_2}{\Sigma}(dt - \omega_1) + \Delta_t(dt - \omega_1)^2 + \Sigma \left(d\theta^2 + \frac{r^2 dr^2}{\Delta_r} \right) + d\psi^2 \cos^2 \theta (r^2 + \ell_2^2) + d\phi^2 \sin^2 \theta (r^2 + \ell_1^2) \quad (5.2)$$

with

$$\omega_1 = \ell_2 \cos^2 \theta d\psi + \ell_1 \sin^2 \theta d\phi \quad , \quad \omega_2 = \ell_1 \cos^2 \theta d\psi + \ell_2 \sin^2 \theta d\phi$$

$$\Delta_r = (r^2 - r_+^2)(r^2 - r_-^2), \quad \Delta_t = \frac{2\mathcal{M}\Sigma - \mathcal{Q}^2}{\Sigma^2}, \quad \Sigma = r^2 + \ell_1^2 \cos^2 \theta + \ell_2^2 \sin^2 \theta$$

These geometries possess two horizons located at

$$y_\pm = r_\pm^2 = \widehat{\mathcal{M}} \pm \sqrt{\widehat{\mathcal{M}}^2 - \widehat{\mathcal{Q}}^2} \quad (5.3)$$

with

$$\widehat{\mathcal{M}} = \mathcal{M} - \frac{\ell_1^2 + \ell_2^2}{2} \quad , \quad \widehat{\mathcal{Q}} = \mathcal{Q} + \ell_1 \ell_2 \quad (5.4)$$

Introducing the variables $\chi = \cos \theta$, $y = r^2$, the scalar wave equation can be separated, setting

$$\Phi(t, y, \chi, \phi_1, \phi_2) = e^{i(-\omega t + m_\phi \phi + m_\psi \psi)} R(r) S(\cos \theta) = e^{i(-\omega t + m_\phi \phi + m_\psi \psi)} \frac{\widehat{R}(y) \widehat{S}(\chi)}{\sqrt{\Delta_\chi(\chi) \Delta_y(y)}} \quad (5.5)$$

with

$$\Delta_y = (y - y_+)(y - y_-). \quad , \quad \Delta_\chi(\chi) = \chi(1 - \chi^2) \quad (5.6)$$

⁵The constant G_5 has been chosen so that the expansion at large distance of the g_{tt} component of the metric reads $g_{tt} \sim -1 + \frac{2\mathcal{M}}{r^2}$.

5.2 Angular motion

The angular equation can be written in the normal form (2.1) for $\widehat{S}(\chi)$ with

$$Q_\chi(\chi) = \frac{1-4(1-\chi^2)m_\psi^2-4\chi^2m_\phi^2+4\chi^2(1-\chi^2)(A+\omega^2\chi^2[\ell_1^2+\ell_2^2(1-\chi^2)]+3\chi^2(1-\chi^2))}{4\chi^2(1-\chi^2)} \quad (5.7)$$

and A the separation constant. The Q_χ -function displays Fuchsian singularities at $\xi = 0, 1$ and an irregular singularity at infinity, so writing $\widehat{S}(\chi) = (4x - 4z)^{-\frac{1}{4}}\Psi(z)$ and

$$\chi^2 = 1 - \frac{z}{x} \quad (5.8)$$

the differential equation takes the normal form with

$$Q_{0,2} = \frac{\delta_k}{(x-z)^2} + \frac{\delta_{p_3}}{z^2} - \frac{1}{z} + \frac{u - \frac{1}{4} + \delta_k + \delta_{p_3}}{z(x-z)} \quad (5.9)$$

and

$$x = q^\chi = \frac{\omega^2}{4}(\ell_2^2 - \ell_1^2) \quad , \quad p_3^\chi = \frac{m_\phi}{2} \quad , \quad k^\chi = \frac{m_\psi}{2} \quad , \quad u^\chi = \frac{1}{4}(A + 1 + \omega^2 \ell_2^2) \quad (5.10)$$

We notice that for $\ell_1^2 = \ell_2^2$, the gauge theory is free and therefore the differential equation can be solved in terms of hypergeometric functions. For simplicity, we will restrict to this case, so we will write

$$p_2^\chi = \sqrt{u^\chi} = \sqrt{\frac{A+1}{4}} = \frac{\ell+1}{2} \quad (5.11)$$

i.e. $A = \ell(\ell+2)$. In terms of these variables, the two solutions read

$$\Psi_\alpha = (z-x)^{\frac{m_\psi+1}{2}} \left(\frac{iz}{x} \right)^{\frac{1+\alpha m_\phi}{2}} {}_2F_1\left(1+\frac{1}{2}(m_\psi+\alpha m_\phi+\ell), \frac{1}{2}(m_\psi+\alpha m_\phi-\ell), 1+\alpha m_\phi, \frac{z}{x}\right) \quad (5.12)$$

with $\alpha = \pm$. Finally requiring regularity at $z \approx x$, i.e. the cancelation of the $\log(z-x)$ -term, one finds a single regular solution (hyperspherical harmonics on S^3) assuming ℓ, m_ϕ, m_ψ are all quantized.

5.3 Radial motion

Now let us consider the radial motion. It is convenient to introduce the following combinations

$$\begin{aligned} m_+ &= (m_\phi + m_\psi)(\ell_1 + \ell_2) - \omega(2\mathcal{M} - \mathcal{Q}) \\ m_- &= (m_\phi - m_\psi)(\ell_1 - \ell_2) - \omega(2\mathcal{M} + \mathcal{Q}) \\ 2\kappa &= \widehat{\mathcal{M}}(m_+ - m_-)^2 + \widehat{\mathcal{Q}}(m_+^2 - m_-^2) \end{aligned} \quad (5.13)$$

parametrizing the frequencies ω , and the two angular quantum numbers m_ϕ, m_ψ . In terms of these variables, the radial wave equation can be written in the form (2.1) for $\widehat{R}(y)$ with

$$Q_y(y) = \frac{y m_+ m_- + \kappa - \Delta_y (A + 4 - \omega^2 (y + 2\mathcal{M})) + \Delta_y'^2}{4\Delta_y^2} \quad (5.14)$$

and A the separation constant. Introducing the variable

$$z = \frac{y_+ - y_-}{y - y_-} \quad (5.15)$$

one can write the radial wave equation in its normal form (2.1) with

$$Q_{2,0}(z) = \frac{\delta_{p_0}}{z(z-1)} + \frac{\delta_{k_0}}{(z-1)^2 z} + \frac{u - \frac{1}{4}}{z^2(z-1)} - \frac{x}{z^3} \quad (5.16)$$

and

$$x = q = \frac{\omega^2}{4}(r_+^2 - r_-^2) \quad , \quad u = \frac{1}{4}(A + 1 - 3\mathcal{M}\omega^2 + \omega^2(r_-^2 + 2\mathcal{M}))$$

$$k_0 = iQ_{\text{grav}} = i \frac{\sqrt{r_-^2 m_+ m_- + \kappa}}{2(r_+^2 - r_-^2)} \quad , \quad p_0 = i \frac{\sqrt{r_+^2 m_+ m_- + \kappa}}{2(r_+^2 - r_-^2)} = iQ_{\text{grav}} + \Delta p \quad (5.17)$$

In the limit $\omega\mathcal{M}$ small, the relation between p_2 and u is determined from (2.58) with

$$P_0(v) = (v - a_{01})(v - a_{02}) \quad , \quad P_3(v) = 1 \quad , \quad P_2(v) = v^2 - u + q \quad (5.18)$$

Solving for $a(u)$ order by order in q , and using the dictionary (5.17) one finds

$$p_2 = \frac{\ell + 1}{2} + \Delta\nu = \frac{\ell + 1}{2} - \frac{\omega^2(m_+ m_- + \ell(\ell + 2)(r_+^2 + r_-^2 + 4\mathcal{M}))}{8\ell(\ell + 1)(\ell + 2)} + \dots \quad (5.19)$$

5.4 Love numbers

The computation of Love and dissipation numbers in five dimensions follows *mutatis mutandis* the same steps as before. The asymptotic solution is given again by (4.23) leading to

$$\Psi(z) \underset{z \rightarrow 0}{\sim} z^{-\frac{\ell}{2}} + z^{\frac{\ell}{2} + 1 + 2\Delta\nu} \mathfrak{L}_0 [1 + \dots] \quad (5.20)$$

The Love number is given again by the universal formula (4.26) where now p_0 , k_0 and p_2 are given by (5.17) and (5.19). In the slowly rotating limit $p_2 \approx \frac{\ell+1}{2}$, $p_0 = k_0 = iQ_{\text{grav}}$ leading to

$$\mathfrak{L}_0 \approx - \frac{\Gamma(-\ell - 2\Delta\nu) \Gamma(\frac{\ell}{2} + 1) \Gamma(\frac{\ell}{2} + 1 + 2iQ_{\text{grav}})}{\Gamma(\ell + 2) \Gamma(-\frac{\ell}{2}) \Gamma(-\frac{\ell}{2} + 2iQ_{\text{grav}})} \quad (5.21)$$

For ℓ even one finds

$$\mathfrak{L}_0 = L + i\Theta \approx - \frac{iQ_{\text{grav}} (\frac{\ell}{2})!^2}{(\ell)! (\ell + 1)!} \prod_{n=1}^{\frac{\ell}{2}} (n^2 + 4Q_{\text{grav}}^2) \quad \ell \in 2\mathbb{N} \quad (5.22)$$

For ℓ odd, \mathfrak{L}_0 diverges, and the static limit is not well defined due to the mixing of the source and the response terms. The leading $\log z$ term in the expansion of (5.20) is instead finite

$$\Psi(z) \underset{z \rightarrow 0}{\sim} z^{-\ell} + z^{\ell+1} (\mathfrak{L}_0 + R \log z + \dots) [1 + O(z)] \quad (5.23)$$

with

$$R = 2\Delta\nu \mathfrak{L}_0 = \frac{(-)^\ell \Gamma(\frac{\ell}{2} + 1) \Gamma(\frac{\ell}{2} + 1 + 2iQ_{\text{grav}})}{\ell! \Gamma(\ell + 2) \Gamma(-\frac{\ell}{2}) \Gamma(-\frac{\ell}{2} + 2iQ_{\text{grav}})} \quad \ell \in 2\mathbb{N} + 1 \quad (5.24)$$

(5.22) and (5.24) reproduces (60) of [49] after setting $Q_{\text{grav}} \rightarrow 0$ and $\ell \rightarrow 2l$.

5.5 Amplification factors

The computation of the amplification factor follows the same steps that the one in four dimensions. The asymptotic solution is given again by (4.31), and the amplification factor by (4.34) and (4.32) with p_0 , k_0 and p_2 given by (5.17) and (5.19). Now it makes difference if ℓ is even or odd. For example, in the static limit $\omega \rightarrow 0$, for ℓ even the amplification factor is given again by (4.37) with $\ell \rightarrow \ell/2$

$$Z \underset{x \rightarrow 0}{\approx} \frac{2|x|^{\ell+1} Q_{\text{grav}}(\frac{\ell}{2})!^4}{(\ell)!^2(\ell+1)!^2} \prod_{n=1}^{\frac{\ell}{2}} (n^2 + 4Q_{\text{grav}}^2) \quad \ell \in 2\mathbb{N} \quad (5.25)$$

while for ℓ odd one finds $Z \underset{x \rightarrow 0}{\approx} 1$. Again an ambiguity on the order of the limits $\omega \rightarrow 0$ and $\ell \rightarrow \mathbb{N}$ is observed.

6 Summary of results

In this work, we have studied the scattering of waves from BH's and ECO's geometries using conformal field theory and localization techniques. Following [27], we related the wave functions to the conformal blocks of Liouville theory and the connection matrices to "braiding" and "fusion" moves of the CFT.

We applied the results to the study of waves propagating in the background of BH's and ECO's in four and five dimensions. ECO's are represented by BH geometries ending on a membrane with reflectivity properties parametrized by the coefficient \mathfrak{R} . The QNM's are defined as poles (in the space of frequencies) of the absorption cross section and they are given by the solutions of the equation

$$M_{--} + \mathfrak{R}M_{+-} = 0 \quad (6.1)$$

with $M_{\alpha\alpha'}$ the connection matrix relating the solution at the would be horizon and infinity. The same characteristic frequencies appear as poles in the echo response of the geometry.

We studied also the tidal response of BH's and ECO's to perturbations slowly varying with time. In four dimensions, the tidal response was extracted from the asymptotic form of the scalar wave function in the region $r_+ - r_- \ll r \ll \omega^{-1}$. We write

$$\Psi(r) \underset{r \rightarrow \infty}{\sim} 1 + \left(\frac{r}{r_+ - r_-} \right)^{-2\ell - 1 - 2\Delta\nu} [\mathfrak{L}(\omega r) + \dots] \quad (6.2)$$

with $\Delta\nu$ given by (4.20) and dots denoting $\frac{r_+ - r_-}{r}$ corrections. The ratio between the response and source terms is parametrized by a "Love function" $\mathfrak{L}(\omega r)$ (rather than a number). We obtained an exact formula for this function in terms of hypergeometric functions. A similar formula *mutatis mutandis* is found in five dimensions with $r \rightarrow r^2$, $\ell \rightarrow \ell/2$. We define the dynamical Love L and dissipation Θ numbers as the real and imaginary parts of $\mathfrak{L}_0 = \mathfrak{L}(0)$. We obtained the universal formula

$$\mathfrak{L}_0 = L + i\Theta = -\frac{\Gamma(-2p_2) \Gamma(\frac{1}{2} + k_0 + p_0 + p_2) \Gamma(\frac{1}{2} + k_0 - p_0 + p_2)}{\Gamma(2p_2) \Gamma(\frac{1}{2} + k_0 + p_0 - p_2) \Gamma(\frac{1}{2} + k_0 - p_0 - p_2)} \quad (6.3)$$

valid for all BH's in four and five dimensions. The parameters k_0 , p_0 and p_2 are given by (4.17) and (4.20) for Kerr-Newmann in four dimensions, and by (5.17) and (5.19) in five dimensions. The non-triviality of the differential equation is codified into a single function: the quantum SW period $p_2 = a(u)$ specifying the gauge theory prepotential ! This function can be derived from a continuous fraction version of the quantum deformed SW curve [33, 34].

(6.3) reproduces the results [53] for the Love number of Kerr BH's in the "strictly" static limit where $\omega = 0$ and ℓ is almost (but not exactly) an integer. This analytic continuation of ℓ is required in order to have an unambiguous split between the source and response terms when ω is exactly zero. The definition of the static limit is subtle since the two limits $\omega \rightarrow 0$ and $\ell \rightarrow \mathbb{N}$ do not commute ! Indeed sending $\ell \rightarrow \mathbb{N}$ and then $\omega \rightarrow 0$ yields a non-trivial result for the Love number that grows with ω^{-1} . This may points to a breaking of the perturbative expansion or a phase transition. A similar behaviour has been observed in superconductors and recently discussed in [66]⁶. Superconductors obey the London equation $j_i(\omega, k) \sim A_i(\omega, k)$ that relates the Fourier transform of the (superconducting) current density j_i to the vector potential A_i , with ω the frequency and k the momentum. The limits $\omega \rightarrow 0$ and $k \rightarrow 0$ do not commute: setting $\omega = 0$ and then sending $k \rightarrow 0$ leads to the Meissner effect explaining the expulsion of magnetic lines out of the conductor while the opposite limit yields a current growing with ω^{-1} explaining the superconductivity. It would be interesting to see whether a similar physical interpretation of the two limits exists for the BH tidal response.

Finally we computed the amplification factors for BH's and ECO's in four and five dimensions that are given again in closed forms in terms of k_0 , p_0 , p_2 . Again ambiguities related to the non-commutativity of the limits $\omega \rightarrow 0$ and $\ell \rightarrow \mathbb{N}$ are found.

Acknowledgments

We thank M. Bianchi, G. Bonelli, V. Cardoso, T. Damour, D. Fioravanti, P. Pani and A. Tanzini for useful discussions and A. Grillo for collaboration in the early stages of this work. R.Poghossian thanks the INFN Roma Tor Vergata and the framework of the Armenian SCS grants 20RF-142 and 21AG-1C062 for financial support. The work of F.Fucito and J.F.Morales is partially supported by the MIUR PRIN Grant 2020KR4KN2 "String Theory as a bridge between Gauge Theories and Quantum Gravity".

A Instanton partition function

The instanton partition function of a four-dimensional SQGT with $\mathcal{N} = 2$ supersymmetry, gauge group $U(2)^n$ living on an Ω -background is specified by the gauge couplings $q_i = e^{2\pi i \tau_i}$, $i = 1, \dots, n$ being a label of the gauge group, the vacuum expectation values a_{iu} of the scalars in the vector multiplets, $u = 1, 2$ a gauge index and the parameters ϵ_1, ϵ_2 specifying the Ω -background. Finally we denote by a_{0u} and $a_{n+1,u}$ the vevs parametrizing the masses of

⁶We thank P.Pani for pointing us this analogy.

the hypermultiplets transforming in the fundamental and anti-fundamental representations at the left and right ends of the quiver. The quiver is displayed in figure 1.

The instanton partition function can be written as the sum [67–71]

$$Z_{\text{inst}}^{U(2)^n} = \sum_Y \prod_{i=1}^n q_i^{|\vec{Y}_i|} \frac{\prod_{s=0}^n \prod_{u,v=1}^2 \mathcal{Z}_{Y_{su}, Y_{s+1,v}}(a_{su} - a_{s+1,v})}{\prod_{i=1}^n \prod_{u \neq v}^2 \mathcal{Z}_{Y_{iu}, Y_{iv}}(a_{iu} - a_{iv})} \quad (\text{A.1})$$

over an array $\vec{Y}_s = \{Y_{s1}, Y_{s2}\}$ of Young tableaux specifying the positions of the instantons associated to the s^{th} -gauge group with the tableaux at the two ends of the quiver empty, i.e. $Y_0 = Y_{n+1} = \{\emptyset, \emptyset\}$.

Each tableau $Y = \{k_1 \geq k_2 \geq \dots\}$ in (A.1) is labelled by a sequence of decreasing integers k_j specifying the height of the j -th column. Similarly, the transpose diagram $Y^T = \{k'_1 \geq k'_2 \geq \dots\}$ is labelled by k'_j specifying the length of the i -th row in the original diagram. The instanton number $|\vec{Y}_s| = \sum_u |Y_{su}|$ with $|Y|$ denoting the total number of boxes in the tableau $|Y| = \sum_j k_j$. Finally the contribution of each bifundamental is given by

$$\mathcal{Z}_{Y_u, W_v}(x) = \prod_{i,j \in Y_u} [x + \epsilon_1(i - k'_{jv}) - \epsilon_2(j - k_{iu} - 1)] \prod_{i,j \in W_v} [x - \epsilon_1(i - k'_{ju} - 1) + \epsilon_2(j - k_{iv})] \quad (\text{A.2})$$

The gauge prepotential is identified with the free energy of the statistical system

$$\mathcal{F}_{\text{inst}}(a_{su}, q_i, \epsilon_1, \epsilon_2) = \epsilon_1 \epsilon_2 \log Z_{\text{inst}}(a_{su}, q_i, \epsilon_1, \epsilon_2) \quad (\text{A.3})$$

B Hypergeometric identities

For convenience of the reader here we collect the identities used in establish braiding and fusion relations

$$\begin{aligned} {}_2F_1(a, b; c; z^{-1}) &= e^{-i\pi a} z^a \frac{\Gamma(c)\Gamma(b-a)}{\Gamma(b)\Gamma(c-a)} {}_2F_1(a, a-c+1; a-b+1; z) \\ &\quad + e^{-i\pi b} z^b \frac{\Gamma(c)\Gamma(a-b)}{\Gamma(a)\Gamma(c-b)} {}_2F_1(b, b-c+1; -a+b+1; z) \\ {}_2F_1(a, b; c; z^{-1}) &= z^a \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)} {}_2F_1(a, a-c+1; a+b-c+1; 1-z) \\ &\quad + e^{i\pi(c-a-b)} z^a (1-z)^{c-a-b} \frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)} {}_2F_1(1-b, c-b; c-a-b+1; 1-z) \\ {}_2F_1(a, b; c; z) &= \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)} {}_2F_1(a, b; a+b-c+1; 1-z) \\ &\quad + (1-z)^{c-a-b} \frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)} {}_2F_1(c-a, c-b; c-a-b+1; 1-z) \end{aligned} \quad (\text{B.1})$$

References

- [1] **LIGO Scientific, Virgo** Collaboration, B. P. Abbott et al., *Observation of Gravitational Waves from a Binary Black Hole Merger*, *Phys. Rev. Lett.* **116** (2016), no. 6 061102, [[arXiv:1602.03837](https://arxiv.org/abs/1602.03837)].

- [2] E. Berti, V. Cardoso, and A. O. Starinets, *Quasinormal modes of black holes and black branes*, *Class. Quant. Grav.* **26** (2009) 163001, [[arXiv:0905.2975](#)].
- [3] E. E. Flanagan and T. Hinderer, *Constraining neutron star tidal Love numbers with gravitational wave detectors*, *Phys. Rev. D* **77** (2008) 021502, [[arXiv:0709.1915](#)].
- [4] V. Cardoso and P. Pani, *Testing the nature of dark compact objects: a status report*, *Living Rev. Rel.* **22** (2019), no. 1 4, [[arXiv:1904.05363](#)].
- [5] C. M. Claudel, K. S. Virbhadra and G. F. R. Ellis, *J. Math. Phys.* **42** (2001), 818-838 doi:10.1063/1.1308507 [[arXiv:gr-qc/0005050](#) [gr-qc]].
- [6] V. Cardoso and P. Pani, *Tests for the existence of black holes through gravitational wave echoes*, *Nature Astron.* **1** (2017), no. 9 586–591, [[arXiv:1709.01525](#)].
- [7] M. Bianchi, D. Consoli, A. Grillo, and J. F. Morales, *The dark side of fuzzball geometries*, *JHEP* **05** (2019) 126, [[arXiv:1811.02397](#)].
- [8] M. Bianchi, A. Grillo, and J. F. Morales, *Chaos at the rim of black hole and fuzzball shadows*, *JHEP* **05** (2020) 078, [[arXiv:2002.05574](#)].
- [9] I. Bena and D. R. Mayerson, *Multipole Ratios: A New Window into Black Holes*, *Phys. Rev. Lett.* **125** (2020), no. 22 22, [[arXiv:2006.10750](#)].
- [10] M. Bianchi, D. Consoli, A. Grillo, J. F. Morales, P. Pani, and G. Raposo, *Distinguishing fuzzballs from black holes through their multipolar structure*, *Phys. Rev. Lett.* **125** (2020), no. 22 221601, [[arXiv:2007.01743](#)].
- [11] I. Bena and D. R. Mayerson, *Black Holes Lessons from Multipole Ratios*, *JHEP* **03** (2021) 114, [[arXiv:2007.09152](#)].
- [12] M. Bianchi, D. Consoli, A. Grillo, J. F. Morales, P. Pani, and G. Raposo, *The multipolar structure of fuzzballs*, *JHEP* **01** (2021) 003, [[arXiv:2008.01445](#)].
- [13] D. R. Mayerson, *Fuzzballs and Observations*, *Gen. Rel. Grav.* **52** (2020), no. 12 115, [[arXiv:2010.09736](#)].
- [14] M. Bianchi, D. Consoli, A. Grillo, and J. F. Morales, *Light rings of five-dimensional geometries*, *JHEP* **03** (2021) 210, [[arXiv:2011.04344](#)].
- [15] T. Ikeda, M. Bianchi, D. Consoli, A. Grillo, J. F. Morales, P. Pani, and G. Raposo, *Black-hole microstate spectroscopy: ringdown, quasinormal modes, and echoes*, [arXiv:2103.10960](#).
- [16] I. Bah, I. Bena, P. Heidmann, Y. Li, and D. R. Mayerson, *Gravitational Footprints of Black Holes and Their Microstate Geometries*, [arXiv:2104.10686](#).
- [17] M. Bianchi and G. Di Russo, *Turning black-holes and D-branes inside out their photon-spheres*, [arXiv:2110.09579](#).
- [18] M. Bianchi and G. Di Russo, *Turning rotating D-branes and BHs inside out their photon-halo*, [arXiv:2203.14900](#).
- [19] T. Regge and J. A. Wheeler, *Stability of a Schwarzschild singularity*, *Phys. Rev.* **108** (1957) 1063–1069.
- [20] J. M. Bardeen, W. H. Press, and S. A. Teukolsky, *Rotating Black Holes: Locally Nonrotating Frames, Energy Extraction, and Scalar Synchrotron Radiation*, *The Astrophysical Journal* **178** (Dec, 1972) 347–370.
- [21] S. Chandrasekhar, *The mathematical theory of black holes*. 1985.

- [22] I. D. Novikov and V. P. Frolov, *PHYSICS OF BLACK HOLES*. Kluwer Academic, Dordrecht, Netherlands, 1989.
- [23] G. Aminov, A. Grassi, and Y. Hatsuda, *Black Hole Quasinormal Modes and Seiberg-Witten Theory*, [arXiv:2006.06111](#).
- [24] M. Bianchi, D. Consoli, A. Grillo, and J. F. Morales, *QNMs of branes, BHs and fuzzballs from Quantum SW geometries*, [arXiv:2105.04245](#).
- [25] M. Bianchi, J. F. Morales, L. Pieri, and N. Zinnato, *More on microstate geometries of 4d black holes*, *JHEP* **05** (2017) 147, [[arXiv:1701.05520](#)].
- [26] L. F. Alday, D. Gaiotto, and Y. Tachikawa, *Liouville Correlation Functions from Four-dimensional Gauge Theories*, *Lett. Math. Phys.* **91** (2010) 167–197, [[arXiv:0906.3219](#)].
- [27] G. Bonelli, C. Iossa, D. P. Lichtig, and A. Tanzini, *Exact solution of Kerr black hole perturbations via CFT2 and instanton counting: Greybody factor, quasinormal modes, and Love numbers*, *Phys. Rev. D* **105** (2022), no. 4 044047, [[arXiv:2105.04483](#)].
- [28] D. Fioravanti, M. Rossi and H. Shu, *JHEP* **12** (2020), 086 doi:10.1007/JHEP12(2020)086 [[arXiv:2004.10722 \[hep-th\]](#)].
- [29] D. Fioravanti and M. Rossi, [[arXiv:2106.07600 \[hep-th\]](#)].
- [30] D. Fioravanti and D. Gregori, *A new method for exact results on Quasinormal Modes of Black Holes*, [arXiv:2112.11434](#).
- [31] A. A. Belavin, A. M. Polyakov, and A. B. Zamolodchikov, *Infinite Conformal Symmetry in Two-Dimensional Quantum Field Theory*, *Nucl. Phys. B* **241** (1984) 333–380.
- [32] N. A. Nekrasov and S. L. Shatashvili, *Quantization of Integrable Systems and Four Dimensional Gauge Theories*, in *16th International Congress on Mathematical Physics*, pp. 265–289, 8, 2009. [arXiv:0908.4052](#).
- [33] H. Poghosyan, *Recursion relation for instanton counting for $SU(2)$ $\mathcal{N} = 2$ SYM in NS limit of Ω background*, *JHEP* **05** (2021) 088, [[arXiv:2010.08498](#)].
- [34] F. Fucito, J. F. Morales, D. R. Pacifici, and R. Poghossian, *Gauge theories on Ω -backgrounds from non commutative Seiberg-Witten curves*, *JHEP* **05** (2011) 098, [[arXiv:1103.4495](#)].
- [35] G. Bonelli, C. Iossa, D. P. Lichtig, and A. Tanzini, *Irregular Liouville correlators and connection formulae for Heun functions*, [arXiv:2201.04491](#).
- [36] F. Novaes and B. Carneiro da Cunha, *Isomonodromy, Painlevé transcendents and scattering off of black holes*, *JHEP* **07** (2014) 132, [[arXiv:1404.5188](#)].
- [37] B. Carneiro da Cunha and F. Novaes, *Kerr Scattering Coefficients via Isomonodromy*, *JHEP* **11** (2015) 144, [[arXiv:1506.06588](#)].
- [38] B. Carneiro da Cunha and F. Novaes, *Kerr-de Sitter greybody factors via isomonodromy*, *Phys. Rev. D* **93** (2016), no. 2 024045, [[arXiv:1508.04046](#)].
- [39] J. B. Amado, B. Carneiro da Cunha, and E. Pallante, *On the Kerr-AdS/CFT correspondence*, *JHEP* **08** (2017) 094, [[arXiv:1702.01016](#)].
- [40] J. Barragán Amado, B. Carneiro Da Cunha, and E. Pallante, *Scalar quasinormal modes of Kerr-AdS₅*, *Phys. Rev. D* **99** (2019), no. 10 105006, [[arXiv:1812.08921](#)].
- [41] F. Novaes, C. Marinho, M. Lencsés, and M. Casals, *Kerr-de Sitter Quasinormal Modes via Accessory Parameter Expansion*, *JHEP* **05** (2019) 033, [[arXiv:1811.11912](#)].

- [42] B. Carneiro da Cunha and J. a. P. Cavalcante, *Confluent conformal blocks and the Teukolsky master equation*, *Phys. Rev. D* **102** (2020), no. 10 105013, [[arXiv:1906.10638](#)].
- [43] J. B. Amado, B. Carneiro da Cunha, and E. Pallante, *Vector perturbations of Kerr-AdS₅ and the Painlevé VI transcendent*, *JHEP* **04** (2020) 155, [[arXiv:2002.06108](#)].
- [44] T. Damour and A. Nagar, *Relativistic tidal properties of neutron stars*, *Phys. Rev. D* **80** (2009) 084035, [[arXiv:0906.0096](#)].
- [45] T. Binnington and E. Poisson, *Relativistic theory of tidal Love numbers*, *Phys. Rev. D* **80** (2009) 084018, [[arXiv:0906.1366](#)].
- [46] H. Fang and G. Lovelace, *Tidal coupling of a Schwarzschild black hole and circularly orbiting moon*, *Phys. Rev. D* **72** (2005) 124016, [[gr-qc/0505156](#)].
- [47] H. S. Chia, *Tidal deformation and dissipation of rotating black holes*, *Phys. Rev. D* **104** (2021), no. 2 024013, [[arXiv:2010.07300](#)].
- [48] L. Hui, A. Joyce, R. Penco, L. Santoni, and A. R. Solomon, *Static response and Love numbers of Schwarzschild black holes*, *JCAP* **04** (2021) 052, [[arXiv:2010.00593](#)].
- [49] D. Pereñiguez and V. Cardoso, *Love numbers and magnetic susceptibility of charged black holes*, *Phys. Rev. D* **105** (2022), no. 4 044026, [[arXiv:2112.08400](#)].
- [50] R. Emparan, A. Fernandez-Pique, and R. Luna, *Geometric polarization of plasmas and Love numbers of AdS black branes*, *JHEP* **09** (2017) 150, [[arXiv:1707.02777](#)].
- [51] V. Cardoso, E. Franzin, A. Maselli, P. Pani, and G. Raposo, *Testing strong-field gravity with tidal Love numbers*, *Phys. Rev. D* **95** (2017), no. 8 084014, [[arXiv:1701.01116](#)]. [Addendum: *Phys.Rev.D* 95, 089901 (2017)].
- [52] V. Cardoso, M. Kimura, A. Maselli, and L. Senatore, *Black Holes in an Effective Field Theory Extension of General Relativity*, *Phys. Rev. Lett.* **121** (2018), no. 25 251105, [[arXiv:1808.08962](#)].
- [53] P. Charalambous, S. Dubovsky, and M. M. Ivanov, *On the Vanishing of Love Numbers for Kerr Black Holes*, *JHEP* **05** (2021) 038, [[arXiv:2102.08917](#)].
- [54] F. Olver, D. Lozier, R. Boisvert, and C. Clark, *The NIST Handbook of Mathematical Functions*. Cambridge University Press, New York, NY, 2010-05-12 00:05:00, 2010.
- [55] G. W. Moore and N. Seiberg, *Classical and Quantum Conformal Field Theory*, *Commun. Math. Phys.* **123** (1989) 177.
- [56] G. W. Moore and N. Seiberg, *Polynomial Equations for Rational Conformal Field Theories*, *Phys. Lett. B* **212** (1988) 451–460.
- [57] M. Matone, *Instantons and recursion relations in N=2 SUSY gauge theory*, *Phys. Lett. B* **357** (1995) 342–348, [[hep-th/9506102](#)].
- [58] R. Flume, F. Fucito, J. F. Morales, and R. Poghossian, *Matone’s relation in the presence of gravitational couplings*, *JHEP* **04** (2004) 008, [[hep-th/0403057](#)].
- [59] R. Poghossian, *Deforming SW curve*, *JHEP* **04** (2011) 033, [[arXiv:1006.4822](#)].
- [60] R. Poghossian, *Deformed SW curve and the null vector decoupling equation in Toda field theory*, *JHEP* **04** (2016) 070, [[arXiv:1601.05096](#)].
- [61] Z. Mark, A. Zimmerman, S. M. Du, and Y. Chen, *A recipe for echoes from exotic compact objects*, *Phys. Rev. D* **96** (2017), no. 8 084002, [[arXiv:1706.06155](#)].

- [62] M. M. Caldarelli, G. Cognola, and D. Klemm, *Thermodynamics of Kerr-Newman-AdS black holes and conformal field theories*, *Class. Quant. Grav.* **17** (2000) 399–420, [[hep-th/9908022](#)].
- [63] M. Bianchi, D. Consoli, A. Grillo, and J. F. Morales, *More on the SW-QNM correspondence*, *JHEP* **01** (2022) 024, [[arXiv:2109.09804](#)].
- [64] A. A. Starobinskii and S. M. Churilov, *Amplification of electromagnetic and gravitational waves scattered by a rotating "black hole"*, *Sov. Phys. JETP* **65** (1974), no. 1 1–5.
- [65] Z. W. Chong, M. Cvetič, H. Lu, and C. N. Pope, *General non-extremal rotating black holes in minimal five-dimensional gauged supergravity*, *Phys. Rev. Lett.* **95** (2005) 161301, [[hep-th/0506029](#)].
- [66] S. A. Hartnoll, C. P. Herzog, and G. T. Horowitz, *Holographic Superconductors*, *JHEP* **12** (2008) 015, [[arXiv:0810.1563](#)].
- [67] N. A. Nekrasov, *Adv. Theor. Math. Phys.* **7** (2003) no.5, 831-864
doi:10.4310/ATMP.2003.v7.n5.a4 [[arXiv:hep-th/0206161](#) [hep-th]].
- [68] R. Flume and R. Poghossian, *Int. J. Mod. Phys. A* **18** (2003), 2541
doi:10.1142/S0217751X03013685 [[arXiv:hep-th/0208176](#) [hep-th]].
- [69] U. Bruzzo, F. Fucito, J. F. Morales, and A. Tanzini, *Multiinstanton calculus and equivariant cohomology*, *JHEP* **05** (2003) 054, [[hep-th/0211108](#)].
- [70] F. Fucito, J. F. Morales and D. Ricci Pacifici, *JHEP* **01** (2013), 091
doi:10.1007/JHEP01(2013)091 [[arXiv:1210.3580](#) [hep-th]].
- [71] N. Nekrasov and V. Pestun, [[arXiv:1211.2240](#) [hep-th]].