

Fast rotation and inviscid limits for the SQG equation with general ill-prepared initial data

Leonardo Kosloff¹, Gabriele Sbaiz²

¹ UNIVERSIDADE ESTADUAL DE CAMPINAS, *IMECC-Departamento de Matemática*,
Rua Sérgio Buarque de Holanda, CEP 13083-859, Campinas, SP, Brazil

² UNIVERSITÀ DEGLI STUDI DI TRIESTE,
Dipartimento di Scienze Economiche, Aziendali, Matematiche e Statistiche “Bruno de Finetti”,
Via Valerio 4/1, 34127 Trieste, Italy

kosloff@ime.unicamp.br, gabriele.sbaiz@phd.units.it

June 22, 2022

Abstract

In the present paper, we study the fast rotation and inviscid limits for the 2-D dissipative surface quasi-geostrophic equation with a dispersive forcing term $A\mathcal{R}_1\vartheta$, in the domain $\Omega = \mathbb{T}^1 \times \mathbb{R}$. In the case when we perform the fast rotation limit (keeping the viscosity fixed), in the context of general ill-prepared initial data, we prove that the limit dynamics is described by a linear equation. On the other hand, performing the combined fast rotation and inviscid limits, we show that the initial data $\bar{\vartheta}_0$ is transported along the motion. The proof of the convergence is based on an application of the Aubin-Lions lemma.

2020 Mathematics Subject Classification: 35Q86 (primary); 35B25, 76U60, 35B40, 76M45 (secondary).

Keywords: SQG equation; fast rotation limit; inviscid limit; singular perturbation problem; ill-prepared data.

1 Introduction

In this paper, we are interested in the description of the surface temperature ϑ on the ocean (see e.g. [12] and [24]). We consider the two-dimensional surface quasi-geostrophic (SQG) equation with dissipation determined by a fractional Laplacian and a dispersive forcing term in the domain $\Omega = \mathbb{T}^1 \times \mathbb{R}$ given by:

$$\begin{cases} \partial_t \vartheta + \operatorname{div}(\vartheta \mathbf{u}) + \nu \Lambda \vartheta + A \mathcal{R}_1 \vartheta = 0 \\ \mathbf{u} = \mathcal{R}^\perp \vartheta := (-\mathcal{R}_2 \vartheta, \mathcal{R}_1 \vartheta) \\ \vartheta|_{t=0} = \vartheta_0, \end{cases} \quad (1.1)$$

where ϑ is a real-valued scalar function, \mathbf{u} is the divergence-free velocity field and A represents the amplitude parameter for the dispersive forcing term. To define the fractional Laplacian operator $\Lambda := \sqrt{-\Delta}$ we can adapt the Fourier transform for functions $f(x_1, x_2) \in L^1(\Omega)$ with $(n_1, \xi_2) \in \mathbb{Z} \times \mathbb{R}$, so that

$$\hat{f}(n_1, \xi_2) = \int_{\mathbb{Z} \times \mathbb{R}} e^{-i(n_1 x_1 + \xi_2 x_2)} f(x_1, x_2) dx_1 dx_2,$$

and Λ is then defined by:

$$\widehat{\Lambda f}(n_1, \xi_2) = (n_1^2 + \xi_2^2)^{1/2} \hat{f}(n_1, \xi_2) := |\tilde{\xi}| \hat{f}(n_1, \xi_2).$$

Moreover, we have $\nu > 0$ and $\mathcal{R}_i := \partial_i \Lambda^{-1}$, for $i = 1, 2$, are the usual Riesz transforms. The non-local dissipative term $\Lambda \vartheta$ comes from the Ekman pumping mechanism (we refer to [17], [20] and [21] for details); while the presence of an environmental horizontal gradient $\mathcal{R}_1 \vartheta = \partial_1 \Lambda^{-1} \vartheta$ represents the advection of a large-scale buoyancy coming from the meridional variation of the Coriolis force. The parameter $A > 0$ is the analogue of the Rossby number which determines the typically large weight of the Coriolis term in geophysical fluid-dynamics. The mathematical analysis of (1.1) started with the work [18] by Kiselev and Nazarov, where the existence of smooth solutions in the torus was addressed.

In the case without the presence of the dispersive forcing term, the non-dissipative SQG equations ($\nu = 0$) are the two-dimensional analogue of the 3D Euler equations in vorticity form (see for e.g. [10]), while the SQG equations are analogous to the 3D Navier-Stokes system; due to this analogy the global regularity of the SQG equations has been intensively studied in recent decades (we refer, instead, to [1], [2] and [3] for the pioneering studies for the Navier-Stokes equations). For an overview about these equations, we refer to [11] and references therein. However, the presence of the dispersive forcing term makes equations (1.1) analogous to the Navier-Stokes-Coriolis (NSC) system, which is a fundamental geophysical model dealing with large-scale phenomena (we refer to [9] for a more detailed discussion). The main advantage of the NSC system is that in the limit of vanishing Rossby number, the fast rotating term “produces” a stabilization effect which ensures the global well-posedness of strong solutions with large initial data, unlike the case of the Navier-Stokes equations (see also [8] in this respect). In particular, in [9] this was proved by establishing Strichartz estimates, which show how the dispersive phenomena weaken the non-linearity and stabilize NSC towards a 2D Navier-Stokes type system. Following this analogy, a similar result for the supercritical dispersive SQG equation was shown in [8], where in (1.1) the dissipative term is represented by $\nu \Lambda^{2\alpha} \vartheta$, $\alpha < 1/2$. In this context, we mention also the work [19] (in the context of well-prepared initial data), where the main tool employed is the relative energy inequality, in order to treat the inviscid incompressible limit of the NSC system with large rotation (see [14] and [7]). Furthermore, the method used in [19] allows to obtain the inviscid limit with fixed or no dispersion, providing an alternative to previous results [5] and [26] for the unforced SQG equation.

Motivated by the previous discussion, our goal is to perform the fast rotation and inviscid limits in the more general framework of *ill-prepared* initial data, where strong convergence of the initial data for the limiting and target systems, or *a priori* structural conditions for the dispersive estimates, are *not* required. In this direction, we remark that, in the domain $\Omega = \mathbb{T}^1 \times \mathbb{R}$, the stability of the SQG system without dispersive forcing and with horizontal dissipation, was recently proved in [25]. Thus, this result shows that periodic boundary conditions allow for a decomposition into the mean flow and its oscillations, in which global existence, and indeed, stability, can be obtained in the Sobolev space $H^2(\Omega)$, without explicit recourse to dispersive effects.

In our work, first of all, we study the regime when the rotational effects are predominant in the dynamics, keeping fixed the viscous coefficient $\nu > 0$, i.e.

$$A = \frac{1}{\varepsilon}, \quad (1.2)$$

for a given $\varepsilon \in]0, 1]$. Next, we analyse the combined fast rotation and inviscid limits where

$$A = \frac{1}{\varepsilon} \quad \text{and} \quad \nu = \nu(\varepsilon) = \varepsilon^\alpha, \quad \alpha > 0. \quad (1.3)$$

The scaling for ν in (1.3) is motivated by the physical Stommel boundary layer model for the western intensification of oceanic currents, where the Ekman pumping dissipation must be taken into account. In the inviscid limit case, this scaling would typically be written with $-1 < \alpha < -2/3$, but relabeling the parameter ε leads to (1.3). Note also that we disregard the frictional sublayer needed to adjust the no-slip boundary conditions (for more details, we refer to [13] and [16]).

In order to prove our results and get the improvement to the more general ill-prepared data, on the one hand, we take advantage of the special form of test functions, belonging to the kernel of the singular perturbation operator; on the other hand, we employ the structure of the system to find compactness properties for the means of temperature $\langle \vartheta_\varepsilon \rangle$. The special form of the test functions combined with the *Aubin-Lions lemma* will allow us to pass to the limit in the equations. Moreover, we need to work with periodic boundary conditions in the domain Ω , so that non-trivial test functions are still allowed by the compact support condition (see remark 3.2 after Theorem 3.1).

The strategy employed is a standard matter in the context of singular perturbations problems and it consists in the following steps:

- (i) develop an existence theory, which holds for any value of $\varepsilon > 0$ fixed;
- (ii) state uniform bounds for the family of solutions in order to extract weak limit points;
- (iii) find the constraints that the limit points have to satisfy;
- (iv) pass to the limit for test functions in the kernel of the singular perturbation operator.

To conclude this part, let us mention that, due to the stabilization effects of the Coriolis force, we expect convergence towards a linear equation in the case of scaling (1.2), while taking the combined scaling (1.3) we will show that the limit dynamics “collapses”. This will tell us that the initial limit profile is transported along the motion.

Let us now give an overview of the paper. In Section 2 we collect our assumptions and we state our main results. In Section 3 we study the singular perturbation part of the equations, recalling the uniform bounds on our family of weak solutions and establishing constraints that the limit points have to satisfy. Section 4 is devoted to the proof of the convergence results for the fast rotation limit and the combined fast rotation and inviscid limits.

Some notation and conventions. Let $B \subset \mathbb{R}^2$. The symbol $C_c^\infty(B)$ denotes the space of ∞ -times continuously differentiable functions on \mathbb{R}^2 and having compact support in B . The dual space $\mathcal{D}'(B)$ is the space of distributions on B . Given $p \in [1, +\infty]$, by $L^p(B)$ we mean the classical space of Lebesgue measurable functions g , where $|g|^p$ is integrable over the set B (with the usual modifications for the case $p = +\infty$). We use also the notation $L_T^p(L^q)$ to indicate the space $L^p([0, T]; L^q(B))$, with $T > 0$. Given

$k \geq 0$, we denote by $W^{k,p}(B)$ the Sobolev space of functions which belongs to $L^p(B)$ together with all their derivatives up to order k . When $p = 2$, we alternately use the notation $W^{k,2}(B)$ and $H^k(B)$. We denote by $\dot{W}^{k,p}(B)$ the corresponding homogeneous Sobolev spaces, i.e. $\dot{W}^{k,p}(B) = \{g \in L^p_{\text{loc}}(B) : D^\alpha g \in L^p(B), |\alpha| = k\}$. Recall that $\dot{W}^{k,p}$ is the completion of $C_c^\infty(\overline{B})$ with respect to the L^p norm of the k -th order derivatives. For the sake of simplicity, we will omit from the notation the set B , that we will explicitly point out if needed.

We denote by \mathbb{T}^1 the one-dimensional flat torus $\mathbb{T}^1 := [-1, 1]/\sim$, where \sim denotes the equivalence relation which identifies -1 and 1 .

In the whole paper, the symbols c and C will denote generic multiplicative constants, which may change from line to line, and which do not depend on the small parameter ε . Sometimes, we will explicitly point out the quantities that these constants depend on, by putting them inside brackets.

Let $(f_\varepsilon)_{0 < \varepsilon \leq 1}$ be a sequence of functions in a normed space X . If this sequence is bounded in X , we use the notation $(f_\varepsilon)_\varepsilon \subset X$.

Acknowledgements

L. Kosloff was supported by FAPESP - Brazil grant 2019/16537-0. G. Sbaiz is member of the INdAM (Italian Institute for Advanced Mathematics) group.

Moreover, the authors acknowledge Francesco Fanelli for his careful remarks that improve a lot the presentation of this work.

2 Setting of the SQG problem and main statements

In this section, we formulate our working hypotheses (see Subsection 2.1) and we state our main results (in Subsection 2.2).

2.1 Formulation of the problem

In this subsection, we present the rescaled SQG system with the dispersive forcing term, which we are going to consider in our study, and we formulate the main working hypotheses. The material of this part is mostly classical: unless otherwise specified, we refer to [19] for details.

2.1.1 Primitive system

To begin with, let us introduce the “primitive system”, i.e. the rescaled SQG system, supplemented with the scaling (1.2) presented in the introduction, where $\varepsilon \in]0, 1]$ is a small parameter. Thus, the system consists of the momentum equation and the quasi-geostrophic balance: respectively,

$$\partial_t \vartheta_\varepsilon + \operatorname{div}(\vartheta_\varepsilon \mathbf{u}_\varepsilon) + \nu \Lambda \vartheta_\varepsilon + \frac{1}{\varepsilon} \mathcal{R}_1 \vartheta_\varepsilon = 0 \quad (2.1)$$

$$\mathbf{u}_\varepsilon = \mathcal{R}^\perp \vartheta_\varepsilon := (-\mathcal{R}_2 \vartheta_\varepsilon, \mathcal{R}_1 \vartheta_\varepsilon). \quad (2.2)$$

The unknown is the fluid surface temperature $\vartheta_\varepsilon = \vartheta_\varepsilon(t, x)$, with $t \in \mathbb{R}_+$ and $x \in \Omega$.

Remark 2.1 *In the case of scaling (1.3), we replace ν with ε^α , $\alpha > 0$.*

2.1.2 Initial data and finite energy weak solutions

We address the singular perturbation problem described in subsection 2.1.1, with scaling (1.2), for general *ill prepared initial data*, in the framework of *finite energy weak solutions* (see e.g. [8]). Since we work with weak solutions based on dissipation estimates, we need to assume that the initial data satisfy the following bound:

$$\sup_{\varepsilon \in]0,1]} \|\vartheta_{0,\varepsilon}\|_{L^2(\Omega)} \leq C. \quad (2.3)$$

Thanks to the previous uniform estimate, up to extraction, we can argue that

$$\overline{\vartheta}_0 := \lim_{\varepsilon \rightarrow 0} \vartheta_{0,\varepsilon}, \quad (2.4)$$

where we agree that the previous limit is taken in the weak topology of $L^2(\Omega)$.

Let us specify better what we mean for *finite energy weak solution* (see [9] for details).

Definition 2.2 *We say that ϑ_ε is a weak solution to the dispersive SQG system (2.1)-(2.2) in $[0, T[\times \Omega$ (for some time $T > 0$) with the initial condition $\vartheta_{0,\varepsilon}$, if:*

- (i) $\vartheta_\varepsilon \in L^\infty([0, T[; L^2(\Omega)) \cap L^2([0, T[; \dot{H}^{1/2}(\Omega))$;
- (ii) *the momentum equation is satisfied in a weak sense: for any $\varphi \in C_c^\infty([0, T[\times \Omega)$, one has*

$$- \int_0^T \int_\Omega \left(\vartheta_\varepsilon \partial_t \varphi + \vartheta_\varepsilon \mathbf{u}_\varepsilon \cdot \nabla \varphi + \nu \Lambda^{1/2} \vartheta_\varepsilon \Lambda^{1/2} \varphi - \frac{1}{\varepsilon} \mathcal{R}_1 \vartheta_\varepsilon \varphi \right) dx dt = \int_\Omega \vartheta_{0,\varepsilon} \varphi(0) dx; \quad (2.5)$$

- (iii) *the quasi-geostrophic balance is satisfied in $\mathcal{D}'([0, T[\times \Omega)$;*
- (iv) *the solution satisfies the following energy estimate:*

$$\|\vartheta_\varepsilon(T)\|_{L^2}^2 + 2\nu \int_0^T \|\Lambda^{1/2} \vartheta_\varepsilon(\tau)\|_{L^2}^2 d\tau \leq \|\vartheta_{0,\varepsilon}\|_{L^2}^2 \quad \text{for all } T > 0. \quad (2.6)$$

The solution is *global* if the previous conditions are satisfied for all $T > 0$.

Under the previous assumptions (collected in subsections 2.1.1 and here above), at any *fixed* value of the parameter $\varepsilon \in]0, 1]$, the existence of a global in time finite energy weak solution ϑ_ε to system SQG, related to the initial data $\vartheta_{0,\varepsilon}$, has been proved in e.g. [22] (see also [8] in this respect).

Remark 2.3 *The term involving the Riesz transform \mathcal{R}_1 does not contribute to the energy estimate (2.6), because for any $s \in \mathbb{R}$, it holds:*

$$\langle \Lambda^s \mathcal{R}_1 \theta, \Lambda^s \theta \rangle_{L^2} = \int_\Omega \Lambda^s \mathcal{R}_1 \theta \overline{\Lambda^s \theta} dx = - \int_\Omega i n_1 |\tilde{\xi}|^{s-1} \hat{\theta} \overline{|\tilde{\xi}|^s \hat{\theta}} d\tilde{\xi} = - \langle \Lambda^s \theta, \Lambda^s \mathcal{R}_1 \theta \rangle_{L^2}. \quad (2.7)$$

2.2 Main results

We can now state our main results. The first statement concerns the case when the rotational and viscosity effects, with fixed ν , are predominant in the dynamics.

Theorem 2.4 *For any fixed value of $\varepsilon \in]0, 1]$, assume the initial data $\vartheta_{0,\varepsilon}$ verifies the hypothesis in subsection 2.1.2 and let ϑ_ε be a corresponding weak solution to system (2.1)-(2.2). Then, one has the following convergence property, for any $T > 0$:*

$$\vartheta_\varepsilon \xrightarrow{*} \overline{\vartheta} \quad \text{weakly-* in} \quad L_T^\infty(L^2(\Omega)) \cap L_T^2(\dot{H}^{1/2}(\Omega)). \quad (2.8)$$

In addition, $\overline{\vartheta}$ is a weak solution to the following linear equation in $\mathbb{R}_+ \times \Omega$:

$$\partial_t \overline{\vartheta} + \nu \Lambda_2 \overline{\vartheta} = 0, \quad (2.9)$$

supplemented with the initial condition $\overline{\vartheta}|_{t=0} = \overline{\vartheta}_0$.

The previous theorem characterizes the limit dynamics of system (2.1)-(2.2) when one considers a fast rotation regime. In contrast with [8], in Theorem 2.4, we are able to consider data which are *ill-prepared*.

Instead, in the fast rotation and inviscid limits, we need more regularity on ϑ_ε since we have to control $\Lambda \vartheta_\varepsilon$. In this case, the limit dynamics is trivial and tells us that the limit temperature $\overline{\vartheta}$ is constant (in time).

Theorem 2.5 *For any fixed value of $\varepsilon \in]0, 1]$, let the initial data $\vartheta_{0,\varepsilon} \in H^s(\Omega)$ for $s > 2$. Let ϑ_ε be a corresponding solution to system (2.1)-(2.2). Then, one has the following convergence property, for $s > 2$:*

$$\vartheta_\varepsilon \xrightarrow{*} \overline{\vartheta} \quad \text{weakly-* in} \quad L_T^\infty(H^s(\Omega)). \quad (2.10)$$

Moreover, one deduces the relation (in the weak sense)

$$\partial_t \overline{\vartheta} = 0,$$

with the initial condition $\overline{\vartheta}|_{t=0} = \overline{\vartheta}_0$.

Remark 2.6 *We point out that the condition $s > 2$ is necessary to have the embedding of Sobolev spaces H^s in the space $W^{1,\infty}$ of globally Lipschitz functions (see the Appendix A for more details in this respect).*

3 Inspection of the singular perturbation

The purpose of this section is twofold. First of all, in Subsection 3.1 we recall the uniform bounds and further properties for our family of weak solutions. Then, we study the singular operator underlying the primitive SQG equations, and determine constraints that the limit points of our family of weak solutions have to satisfy (see Subsection 3.2).

3.1 Uniform bounds

In this section we will state the uniform bounds on the sequence $(\vartheta_\varepsilon)_\varepsilon$.

Indeed, with the energy estimate (2.6) at hand, we can derive uniform bounds for our family of weak solutions. To begin with, we point out that, owing to the assumption (2.3), the right-hand side of (2.6) is *uniformly bounded* for all $\varepsilon \in]0, 1]$. Then, one has

$$\sup_{T \in \mathbb{R}_+} \|\vartheta_\varepsilon(T)\|_{L^2(\Omega)} \leq c \quad (3.1)$$

$$\int_0^T \|\Lambda^{1/2} \vartheta_\varepsilon(\tau)\|_{L^2(\Omega)}^2 d\tau \leq c \quad \text{for all } T > 0. \quad (3.2)$$

Due to the relation (2.2) and the fact that the Riesz operator is a 0-th order operator, one gets also

$$\sup_{T \in \mathbb{R}_+} \|\mathbf{u}_\varepsilon(T)\|_{L^2} + \int_0^T \|\Lambda^{1/2} \mathbf{u}_\varepsilon(\tau)\|_{L^2}^2 d\tau \leq c, \quad (3.3)$$

where the generic constant c is independent of $\varepsilon > 0$.

In view of the previous properties, there exist $\overline{\vartheta}, \overline{\mathbf{u}} \in L_T^\infty(L^2) \cap L_T^2(\dot{H}^{1/2})$ such that (up to the extraction of a suitable subsequence) one has

$$\vartheta_\varepsilon \xrightarrow{*} \overline{\vartheta} \quad \text{and} \quad \mathbf{u}_\varepsilon \xrightarrow{*} \overline{\mathbf{u}}. \quad (3.4)$$

3.2 Constraints on the limit dynamics

In this section, we establish some properties that the limit points of the family $(\vartheta_\varepsilon)_\varepsilon$ have to satisfy. These are static relations, which do not characterise the limit dynamics yet.

Proposition 3.1 *Let $(\vartheta_\varepsilon)_\varepsilon$ be a family of weak solutions, related to initial data $(\vartheta_{0,\varepsilon})_\varepsilon$ verifying hypothesis of subsection 2.1.2. Let $\overline{\vartheta}$ be a limit point of the previous sequence as identified in Subsection 3.1. Then, one deduces the relation*

$$\partial_1 \overline{\vartheta} = 0 \quad \text{a.e. in } \mathbb{R}_+ \times \Omega. \quad (3.5)$$

Proof: Let us consider the weak formulation of (2.1). We test it against $\varepsilon\varphi$ where φ is a compactly supported test function in $C_c^\infty(\mathbb{R}_+ \times \Omega)$. Denoting $[0, T] \times K = \text{supp } \varphi$ with $\varphi(T, \cdot) = 0$ (and $T > 0$), we have

$$-\varepsilon \int_0^T \int_K \left(\vartheta_\varepsilon \partial_t \varphi + \vartheta_\varepsilon \mathbf{u}_\varepsilon \cdot \nabla \varphi + \nu \Lambda^{1/2} \vartheta_\varepsilon \Lambda^{1/2} \varphi \right) dx dt + \int_0^T \int_K \mathcal{R}_1 \vartheta_\varepsilon \varphi dx dt = \varepsilon \int_K \vartheta_{0,\varepsilon} \varphi(0) dx. \quad (3.6)$$

By uniform bounds (3.1) and (3.2), the first three integrals on the left-hand side converge to 0, when tested against any smooth compactly supported φ .

Analogously, thanks to the bound (2.3) on the initial data, also the term on the right-hand side of (3.6) vanishes.

Then, passing to the limit for $\varepsilon \rightarrow 0$, we find

$$\int_0^T \int_K \mathcal{R}_1 \overline{\vartheta} \varphi dx dt = 0,$$

for any test function $\varphi \in C_c^\infty(\mathbb{R}_+ \times \Omega)$, which in particular implies

$$\mathcal{R}_1 \overline{\vartheta} = 0 \quad \text{a.e. in } \mathbb{R}_+ \times \Omega.$$

At this point, employing the Fourier transform, one can deduce that

$$\partial_1 \overline{\vartheta} = 0 \quad \text{a.e. in } \mathbb{R}_+ \times \Omega. \quad (3.7)$$

This completes the proof of the proposition. \blacksquare

Remark 3.2 *Observe that the proof above relies on the fact that the compact support of test functions does not force them to be 0 everywhere, unlike the case of the domain \mathbb{R}^2 , where the condition that the test functions are only dependent on the vertical variable implies they must vanish inside the support as well.*

4 Limit dynamics

4.1 Convergence to the fast rotation limit dynamics

In this section, we will show the convergence of momentum equation (2.5) towards the linear equation (2.9) depicted in Theorem 2.4.

The uniform bounds of Subsection 3.1 are not enough for proving convergence in the weak formulation of the momentum equation: the main problem relies on identifying the weak limit of the convective term $\text{div}(\vartheta_\varepsilon \mathbf{u}_\varepsilon)$.

At this point, we rewrite the convective term in the weak formulation

$$-\int_0^T \int_\Omega \vartheta_\varepsilon \mathbf{u}_\varepsilon \cdot \nabla \psi \, dx \, dt = -\int_0^T \int_{\mathbb{T}^1 \times \mathbb{R}} \vartheta_\varepsilon (u_\varepsilon)_2 \partial_2 \psi \, dx \, dt = -2 \int_{\mathbb{R}} \langle \vartheta_\varepsilon (u_\varepsilon)_2 \rangle \partial_2 \psi \, dx \, dt, \quad (4.1)$$

where the test-function ψ is defined as

$$\psi \in C_c^\infty([0, T[\times \Omega; \mathbb{R}) \quad \text{such that} \quad \partial_1 \psi = 0. \quad (4.2)$$

The previous relation (4.1) pushes our attention to find compactness properties for $\langle \vartheta_\varepsilon \rangle$.

Taking the mean in equation (2.1) and thanks to the fact that the mean of the Riesz term $\langle \mathcal{R}_1 \vartheta_\varepsilon \rangle$ is zero, we get

$$\partial_t \langle \vartheta_\varepsilon \rangle = -\text{div} \langle \vartheta_\varepsilon \mathbf{u}_\varepsilon \rangle - \nu \Lambda \langle \vartheta_\varepsilon \rangle. \quad (4.3)$$

To analyse the former term in the right-hand side of (4.3), we employ the paradifferential calculus (see Chapter 2 of [4]). Using the Bony decomposition, one can find that the term $\vartheta_\varepsilon \mathbf{u}_\varepsilon$ is in $L_T^2 B_{2,1}^{-1/2}$ and thanks to the embedding $B_{2,1}^{-1/2} \hookrightarrow H^{-1/2}$, we argue that $\text{div} \langle \vartheta_\varepsilon \mathbf{u}_\varepsilon \rangle$ belongs to $L_T^2 H^{-3/2}$.

Then, recalling equation (4.3), we deduce that $(\partial_t \langle \vartheta_\varepsilon \rangle)_\varepsilon$ is uniformly bounded in $L_T^2 H_{\text{loc}}^{-3/2}$, which implies $(\langle \vartheta_\varepsilon \rangle)_\varepsilon \subset W_T^{1,2} H_{\text{loc}}^{-3/2}$. Moreover, we already know that $(\langle \vartheta_\varepsilon \rangle)_\varepsilon \subset L_T^\infty L^2 \cap L_T^2 \dot{H}^{1/2}$. Therefore, the Aubin-Lions lemma gives compactness of $(\langle \vartheta_\varepsilon \rangle)_\varepsilon$ in e.g. $L_T^2 L_{\text{loc}}^2$. Thus, we deduce the strong convergence (up to extraction) for $\varepsilon \rightarrow 0$:

$$\langle \vartheta_\varepsilon \rangle \rightarrow \langle \overline{\vartheta} \rangle \quad \text{in} \quad L_T^2 L_{\text{loc}}^2.$$

This implies that

$$\langle \vartheta_\varepsilon \mathbf{u}_\varepsilon \rangle \rightarrow \langle \overline{\vartheta} \overline{\mathbf{u}} \rangle \quad \text{in} \quad \mathcal{D}'(\mathbb{R}_+ \times \Omega). \quad (4.4)$$

4.1.1 Description of the limit system

With the convergence established in (4.4), we can pass to the limit in the equations.

To begin with, we take a test-function ψ (for $T > 0$) as in (4.2). For such a ψ , the Riesz term $\mathcal{R}_1 \vartheta_\varepsilon$ vanishes identically. Hence, the momentum equation in its weak formulation reads:

$$- \int_0^T \int_\Omega \left(\vartheta_\varepsilon \partial_t \psi + \vartheta_\varepsilon \mathbf{u}_\varepsilon \cdot \nabla \psi + \nu \Lambda^{1/2} \vartheta_\varepsilon \Lambda^{1/2} \psi \right) dx dt = \int_\Omega \theta_{0,\varepsilon} \psi(0) dx. \quad (4.5)$$

Making use of the uniform bounds of Subsection 3.1, we can pass to the limit in the ∂_t term and in the viscosity term. Moreover, our assumptions imply that $\vartheta_{0,\varepsilon} \rightharpoonup \bar{\vartheta}_0$ in e. g. L^2_{loc} . Thanks to the convergence identified in (4.4) for the convective term, letting $\varepsilon \rightarrow 0$, we may infer that

$$- \int_0^T \int_\Omega \left(\bar{\vartheta} \partial_t \psi + \nu \Lambda^{1/2} \bar{\vartheta} \Lambda^{1/2} \psi \right) dx dt = \int_\Omega \bar{\vartheta}_0 \psi(0) dx, \quad (4.6)$$

where we have also employed the constraint (3.5) and the special structure of the test functions (4.2). This concludes the proof of Theorem 2.4.

4.2 Proof of the convergence in the inviscid and fast rotation case

In this subsection, we consider the case when $\nu = \nu(\varepsilon) = \varepsilon^\alpha$, $\alpha > 0$. In that scaling, the energy estimate reads

$$\|\vartheta_\varepsilon(T)\|_{L^2}^2 + 2\varepsilon^\alpha \int_0^T \|\Lambda^{1/2} \vartheta_\varepsilon(\tau)\|_{L^2}^2 d\tau \leq \|\vartheta_{0,\varepsilon}\|_{L^2}^2 \quad \text{for all } T > 0. \quad (4.7)$$

Therefore, one completely loses the uniform control on $\Lambda^{1/2} \vartheta_\varepsilon$. In order to recover a similar bound to (3.2), we have to consider strong solutions with the following regularity properties:

$$\vartheta_\varepsilon \in L^\infty([0, T[; H^s(\Omega)) \quad \text{and} \quad s > 2.$$

The aim now is to show higher order estimates for the temperature. We have already presented in (4.7) the L^2 estimate. In order to obtain the H^s control, we employ the Littlewood-Paley decomposition (see the Appendix A). Applying the dyadic blocks Δ_j to equation (2.1) (see e.g. [23]), we obtain

$$\partial_t \Delta_j \vartheta_\varepsilon + \mathbf{u}_\varepsilon \cdot \nabla \Delta_j \vartheta_\varepsilon + \varepsilon^\alpha \Lambda \Delta_j \vartheta_\varepsilon + \frac{1}{\varepsilon} \mathcal{R}_1 \Delta_j \vartheta_\varepsilon = [\mathbf{u}_\varepsilon \cdot \nabla, \Delta_j] \vartheta_\varepsilon.$$

Multiplying by $\Delta_j \vartheta_\varepsilon$ and using the orthogonality of the Riesz term, it follows that

$$\partial_t \|\Delta_j \vartheta_\varepsilon\|_{L^2}^2 + 2\varepsilon^\alpha \|\Lambda^{1/2} \Delta_j \vartheta_\varepsilon\|_{L^2}^2 \leq 2 \|[\mathbf{u}_\varepsilon \cdot \nabla, \Delta_j] \vartheta_\varepsilon\|_{L^2} \|\Delta_j \vartheta_\varepsilon\|_{L^2}.$$

In particular, we get

$$\|\Delta_j \vartheta_\varepsilon\|_{L^2} \leq \|\Delta_j \vartheta_{0,\varepsilon}\|_{L^2} + C \int_0^t \|[\mathbf{u}_\varepsilon \cdot \nabla, \Delta_j] \vartheta_\varepsilon\|_{L^2} d\tau.$$

At this point, thanks to the commutator estimates (we refer e.g. to [23]) we have

$$\|[\mathbf{u}_\varepsilon \cdot \nabla, \Delta_j] \vartheta_\varepsilon\|_{L^2} \leq C c_j(t) (\|\vartheta_\varepsilon\|_{L^\infty} \|\mathbf{u}_\varepsilon\|_{H^s} + \|\vartheta_\varepsilon\|_{H^s} \|\mathbf{u}_\varepsilon\|_{L^\infty}) \leq C c_j(t) \|\vartheta_\varepsilon\|_{L^2}^2,$$

where $(c_j(t))_{j \geq -1}$ is a sequence in the unit ball of ℓ^2 .

After summing on indices $j \geq -1$, we finally derive for all $t \geq 0$:

$$\|\vartheta_\varepsilon(t)\|_{H^s} \leq \|\vartheta_{0,\varepsilon}\|_{H^s} + C \int_0^t \|\vartheta_\varepsilon(\tau)\|_{H^s}^2 d\tau. \quad (4.8)$$

The scope now is finding a time $T^* > 0$ for which the solutions are uniformly bounded (in ε) in the interval $[0, T^*]$.

We define $T_\varepsilon^* > 0$ such that

$$T_\varepsilon^* := \sup \left\{ t > 0 : \int_0^t \|\vartheta_\varepsilon\|_{H^s}^2 \leq \|\vartheta_{0,\varepsilon}\|_{H^s} \right\}. \quad (4.9)$$

Then, we deduce $\|\vartheta_\varepsilon(t)\|_{H^s} \leq C \|\vartheta_{0,\varepsilon}\|_{H^s}$ for all times $t \in [0, T_\varepsilon^*]$ and for some positive constant $C = C(s)$. Therefore, for all $t \in [0, T_\varepsilon^*]$ we gather

$$\int_0^t \|\vartheta_\varepsilon(\tau)\|_{H^s}^2 d\tau \leq Ct \|\vartheta_{0,\varepsilon}\|_{H^s}^2.$$

By using the definition (4.9) of T_ε^* , we finally argue that

$$T_\varepsilon^* \geq \frac{C}{\|\vartheta_{0,\varepsilon}\|_{H^s}}.$$

In particular, this implies that there exists a time $T^* > 0$ such that

$$T^* := \sup_{\varepsilon > 0} T_\varepsilon^* > 0.$$

From the definition of $T^* > 0$ and relation (4.8), one can obtain that

$$\sup_{\varepsilon \in]0,1]} \|\vartheta_\varepsilon\|_{L_{T^*}^\infty(H^s)} \leq C, \quad (4.10)$$

and this allows to control also the dissipative term in the relation (4.7).

At this point, since those strong solutions are in particular weak solutions, one can adapt the arguments developed in subsections 3.1 and 4.1 to conclude. Indeed, letting $\varepsilon \rightarrow 0$, one has

$$-\int_0^{T^*} \int_\Omega \bar{\vartheta} \partial_t \psi \, dx \, dt = \int_\Omega \bar{\theta}_0 \psi(0) \, dx, \quad (4.11)$$

for all test functions ψ defined as in (4.2). This proves the limit dynamics in Theorem 2.5.

A Appendix – Littlewood-Paley theory

In this appendix, we present some tools from Littlewood-Paley theory, which we have exploited in our analysis. We refer e.g. to Chapter 2 of [4] for details. For simplicity of exposition, we deal with the \mathbb{R}^d case, with $d \geq 1$; however, the whole construction can be adapted also to the d -dimensional torus \mathbb{T}^d , and to the “hybrid” case $\mathbb{R}^{d_1} \times \mathbb{T}^{d_2}$.

First of all, we introduce the *Littlewood-Paley decomposition*. For this, we fix a smooth radial function χ such that $\text{supp } \chi \subset B(0, 2)$, $\chi \equiv 1$ in a neighborhood of $B(0, 1)$ and the map $r \mapsto \chi(re)$ is non-increasing over \mathbb{R}_+ for all unitary vectors $e \in \mathbb{R}^d$. Set $\varphi(\xi) = \chi(\xi) - \chi(2\xi)$ and $\varphi_j(\xi) := \varphi(2^{-j}\xi)$ for all $j \geq 0$. The dyadic blocks $(\Delta_j)_{j \in \mathbb{Z}}$ are defined by¹

$$\Delta_j := 0 \quad \text{if } j \leq -2, \quad \Delta_{-1} := \chi(D) \quad \text{and} \quad \Delta_j := \varphi(2^{-j}D) \quad \text{if } j \geq 0.$$

For any $j \geq 0$ fixed, we also introduce the *low frequency cut-off operator*

$$S_j := \chi(2^{-j}D) = \sum_{k \leq j-1} \Delta_k. \quad (\text{A.1})$$

Note that S_j is a convolution operator. More precisely, after defining

$$K_0 := \mathcal{F}^{-1}\chi \quad \text{and} \quad K_j(x) := \mathcal{F}^{-1}[\chi(2^{-j}\cdot)](x) = 2^{jd}K_0(2^jx),$$

for all $j \in \mathbb{N}$ and all tempered distributions $u \in \mathcal{S}'$, we have that $S_j u = K_j * u$. Thus, the L^1 norm of K_j is independent of $j \geq 0$, hence S_j maps continuously L^p into itself, for any $1 \leq p \leq +\infty$.

The following property holds true: for any $u \in \mathcal{S}'$, then one has the equality $u = \sum_j \Delta_j u$ in the sense of \mathcal{S}' . Let us also recall the so-called *Bernstein inequalities*.

Lemma A.1 *Let $0 < r < R$. A constant C exists so that, for any non-negative integer k , any couple (p, q) in $[1, +\infty]^2$, with $p \leq q$, and any function $u \in L^p$, we have, for all $\lambda > 0$,*

$$\begin{aligned} \text{supp } \widehat{u} \subset B(0, \lambda R) &\implies \|\nabla^k u\|_{L^q} \leq C^{k+1} \lambda^{k+d(\frac{1}{p}-\frac{1}{q})} \|u\|_{L^p}; \\ \text{supp } \widehat{u} \subset \{\xi \in \mathbb{R}^d : \lambda r \leq |\xi| \leq \lambda R\} &\implies C^{-k-1} \lambda^k \|u\|_{L^p} \leq \|\nabla^k u\|_{L^p} \leq C^{k+1} \lambda^k \|u\|_{L^p}. \end{aligned}$$

By use of Littlewood-Paley decomposition, we can define the class of Besov spaces.

Definition A.2 *Let $s \in \mathbb{R}$ and $1 \leq p, r \leq +\infty$. The non-homogeneous Besov space $B_{p,r}^s$ is defined as the subset of tempered distributions u for which*

$$\|u\|_{B_{p,r}^s} := \left\| \left(2^{js} \|\Delta_j u\|_{L^p} \right)_{j \geq -1} \right\|_{\ell^r} < +\infty.$$

Besov spaces are interpolation spaces between Sobolev spaces. In fact, for any $k \in \mathbb{N}$ and $p \in [1, +\infty]$ we have the chain of continuous embeddings $B_{p,1}^k \hookrightarrow W^{k,p} \hookrightarrow B_{p,\infty}^k$, which, when $1 < p < +\infty$, can be refined to $B_{p,\min(p,2)}^k \hookrightarrow W^{k,p} \hookrightarrow B_{p,\max(p,2)}^k$. In particular, for all $s \in \mathbb{R}$ we deduce that $B_{2,2}^s \equiv H^s$, with equivalence of norms:

$$\|f\|_{H^s} \sim \left(\sum_{j \geq -1} 2^{2js} \|\Delta_j f\|_{L^2}^2 \right)^{1/2}. \quad (\text{A.2})$$

¹We agree that $f(D)$ stands for the pseudo-differential operator $u \mapsto \mathcal{F}^{-1}[f(\xi)\widehat{u}(\xi)]$.

As an immediate consequence of the first Bernstein inequality, one gets the following embedding result, which generalises Sobolev embeddings.

Proposition A.3 *The space $B_{p_1, r_1}^{s_1}$ is continuously embedded in the space $B_{p_2, r_2}^{s_2}$ for all indices satisfying $p_1 \leq p_2$ and either $s_2 < s_1 - d(1/p_1 - 1/p_2)$, or $s_2 = s_1 - d(1/p_1 - 1/p_2)$ and $r_1 \leq r_2$.*

In particular, we get the following chain of continuous embeddings:

$$B_{p, r}^s \hookrightarrow W^{1, \infty},$$

whenever the triplet $(s, p, r) \in \mathbb{R} \times [1, +\infty]^2$ satisfies

$$s > 1 + \frac{d}{p} \quad \text{or} \quad s = 1 + \frac{d}{p} \quad \text{and} \quad r = 1. \quad (\text{A.3})$$

References

- [1] A. Babin, A. Mahalov, B. Nicolaenko: *Global splitting, integrability and regularity of 3D Euler and Navier-Stokes equations for uniformly rotating fluids*. European J. Mech. B/Fluids, **15** (1996), n. 3, 291-300.
- [2] A. Babin, A. Mahalov, B. Nicolaenko: *Regularity and integrability of 3D Euler and Navier-Stokes equations for rotating fluids*. Asymptot. Anal., **15** (1997), n. 2, 103-150.
- [3] A. Babin, A. Mahalov, B. Nicolaenko: *Global regularity of 3D rotating Navier-Stokes equations for resonant domains*. Indiana Univ. Math. J., **48** (1999), n. 3, 1133-1176.
- [4] H. Bahouri, J.-Y. Chemin, R. Danchin: *“Fourier analysis and nonlinear partial differential equations”*. Grundlehren der Mathematischen Wissenschaften (Fundamental Principles of Mathematical Sciences), **343**, Springer, Heidelberg (2011).
- [5] L. C. Berselli: *Vanishing viscosity limit and long-time behavior for 2D quasi-geostrophic equations*. Indiana Univ. Math. J., **51** (2002), n. 4, 905-930.
- [6] A. Bracco: *Boundary layer separation in the surface quasi-geostrophic equations*. Nuovo Cimento della Società Italiana di Fisica C, **23** (2000), n. 5, 487-505.
- [7] M. Caggio, S. Necasová: *Inviscid incompressible limits for rotating fluids*. Nonlinear Anal., **163** (2017), 1-18.
- [8] M. Cannone, C. Miao, L. Xue: *Global regularity for the supercritical dissipative quasi-geostrophic equation with large dispersive forcing*. Proc. Lond. Math. Soc., **106** (2013), 650-674.
- [9] J.-Y. Chemin, B. Desjardins, I. Gallagher, E. Grenier: *“Mathematical geophysics. An introduction to rotating fluids and the Navier-Stokes equations”*. Oxford Lecture Series in Mathematics and its Applications, **32**, Oxford University Press, Oxford (2006).
- [10] P. Constantin, A. J. Majda, E. Tabak: *Formation of strong fronts in the 2-D quasigeostrophic thermal active scalar*. Nonlinearity, **7** (1994), 1495-1533.
- [11] P. Constantin, V. Vicol, A. Tarfulea: *Long time dynamics of forced critical SQG*. Comm. Math. Phys., **335** (2015), n. 1, 93-141.
- [12] B. Cushman-Roisin: *“Introduction to geophysical fluid dynamics”*. Prentice-Hall, Englewood Cliffs (1994).
- [13] B. Desjardins, E. Grenier: *On the homogeneous model of wind-driven ocean circulation*, SIAM J. Appl. Math., **60** (2000), n. 1, 43-60.
- [14] E. Feireisl, B. J. Jin, A. Novotný: *Relative entropies, suitable weak solutions, and weak-strong uniqueness for the compressible Navier-Stokes system* J. Math. Fluid Mech., **14** (2012), n. 4, 717-730.
- [15] E. Feireisl, A. Novotný: *“Singular limits in thermodynamics of viscous fluids”*. Advances in Mathematical Fluid Mechanics, Birkhäuser Verlag, Basel (2009).
- [16] D. Gérard-Varet, T. Paul: *Remarks on Boundary Layer Expansions*. Comm. Partial Differential Equations, **33** (2008), n. 1-3, 97-130.
- [17] I. Held, R. Pierrehumbert, S. Garner, K. Swanson: *Surface quasi-geostrophic dynamics*. J. Fluid Mech., **282** (1995), 1-20.

- [18] A. Kiselev, F. Nazarov: *Global regularity for the critical dispersive dissipative surface quasi-geostrophic equation*. Nonlinearity, **23** (2010), n. 3, 549-554.
- [19] L. Kosloff, C. J. Niche, G. Planas: *Inviscid limit for SQG equation in different dispersive regimes via relative energy inequality*. Appl. Math. Lett., **88** (2019), 243-249.
- [20] G. Lapeyre: *Surface quasi-geostrophy*. Fluids, **2** (2017), n. 1, Paper n. 7.
- [21] J. Pedlosky: *“Geophysical fluid dynamics”*. Springer-Verlag, New-York (1987).
- [22] S. G. Resnick: *“Dynamical problems in non-linear advective partial differential equations”*. Ph. D. Thesis, The University of Chicago (1995), 76 pp.
- [23] G. Sbaiz: *Fast rotation limit for the 2-D non-homogeneous incompressible Euler equations*. J. Math. Anal. Appl., **512** (2022), n. 1, Paper n. 126140, 41 pp.
- [24] G. K. Vallis: *“Atmospheric and oceanic fluid dynamics: fundamentals and large-scale circulation”*. Cambridge University Press, Cambridge (2006).
- [25] P. Wang: *Stability for the 2D anisotropic surface quasi-geostrophic equation with horizontal dissipation*. Appl. Math. Lett., **123** (2022), Paper n. 107577, 9 pp.
- [26] J. Wu: *Inviscid limits and regularity estimates for the solutions of the 2-D dissipative quasi-geostrophic equations*. Indiana Univ. Math. J., **46** (1997), n. 4, 1113-1124.