

An anisotropic gravity theory

A. García-Parrado ¹ E. Minguzzi ²

¹Departamento de Matemáticas, Universidad de Córdoba,
Campus de Rabanales, 14071, Córdoba, Spain
e-mail: agparrado@uco.es

²Dipartimento di Matematica e Informatica “U. Dini”,
Università degli Studi di Firenze,
Via S. Marta 3, I-50139 Firenze, Italy.
e-mail: ettore.minguzzi@unifi.it

Abstract

We study an action integral for Finsler gravity obtained by pulling back an Einstein-Cartan-like Lagrangian from the tangent bundle to the base manifold. The vacuum equations are obtained imposing stationarity with respect to any section (observer) and are well posed as they are independent of the section. They imply that in vacuum the metric is actually independent of the velocity variable so the dynamics becomes coincident with that of general relativity.

1 Introduction

In this work we explore some features of a possible dynamics for Finsler gravity. The main mathematical objects of Finsler geometry, from the metric to the linear Finsler connections, live on the slit tangent bundle $E = TM \setminus 0$ of the base manifold M . Typically, in order to construct a Finsler action, one would integrate over the indicatrix, namely over the locus $\{(x, y) : 2\mathcal{L}(x, y) = -1\}$, where \mathcal{L} is the Finsler Lagrangian. This approach is followed in [1, 3, 16, 17] particularly with reference to an action which is the integral of the Ricci scalar, see also [15, 6, 8] for further analysis of this type of action. To the best of our knowledge no investigation has been devoted to a different approach, namely to actions obtained via an integral over M . In general, one would need to pull back all relevant quantities to M via a section $s : M \rightarrow E$ of the vector bundle $\pi : E \rightarrow M$. The problem with this approach is that the action would depend on the section. We believe that a natural solution to this problem is to impose that the dynamical equations should be equations on E such that the action is stationary for every possible choice of section s . The idea is that a section could represent an observer and hence the action could be observer dependent while the dynamical equations would not. It turns out that more is true. We shall find that, at least in vacuum and on-shell, the action is actually independent of the section.

Let $\{x^\alpha\}$ be local coordinates, let $e_a = e_a^\mu(x)\partial/\partial x^\mu$, be a basis field on M , let $\{e^a\}$ be the cobasis, and let $\{x^\alpha, y^a\}$ be local coordinates induced on E . A Finsler connection ∇ is a Koszul connection on the pullback¹ (linear) bundle $\pi^*TM \rightarrow E$. It is well known that that it is possible to construct many canonical Finsler connections (e.g. Cartan, Chern, Berwald) but that all these connections induce the same non-linear (Barthel) connection, namely the same splitting $TE = VE \oplus HE$. In all these cases $HE = \ker \nabla y$ where $y = y^a e_a : E \rightarrow TM$ is the Liouville vector field. One says that all notable Finsler connections are *regular*. The regularity condition is equivalent to the fact that the forms $\omega^a = \pi^* e^a$ joined with² $\bar{\omega}^a = (\nabla y)^a$, form a basis on T^*M , cf. [13].

We stress that, although we make use of a linear Finsler connection, our theory should be more properly referred as an anisotropic theory, rather than as a Finslerian theory, as we do not assume that the metric is a vertical Hessian of some Finsler Lagrangian \mathcal{L} as in (pseudo-)Finsler geometry. Also the non-linear connection will not be treated as an independent variable, rather it will be deduced from the linear Finsler connection by the imposition of the regularity condition. In this way the variations of the connections will be totally unconstrained as in our purely metric-affine theory [5].³

One should not expect to determine the Finsler connection univocally. Indeed, already in the Palatini formulation of the Einstein-Hilbert Lagrangian one finds that the connection is determined up to projective transformations. In order to find a unique connection, say the Levi-Civita connection, it is necessary to impose some a priori constraint such as metricity as in the Einstein-Cartan formulation.

In a fully metric-affine approach, in which metric and connection are a priori independent, one is left with the conundrum of justifying why the material part of the action does not exhibit shear hypermomentum. In a previous work [5] we showed that one could solve this problem by enlarging the action symmetry to what we called the *amplified symmetry*. In short our approach showed that one can really work with equivalence classes of connections as long as the material part of the action shares the same symmetry that determines the classes.

In a Finslerian framework one is lead to reconsider these findings in light of specific features of Finsler geometry. One might expect the non-linear (Ehresmann) connection rather than the Finsler connection to be the physical ingredient of a Finslerian gravity theory. The theory developed in [16] does indeed share this feature as it is based on the Ricci scalar that indeed only depends on the non-linear curvature. Still there is also the chance that the true variable of a Finslerian action could be the Finsler connection, possibly constrained in some way. If that is the case one possibly natural constraint would be the metric

¹We use the pullback approach which is by now standard [2]. However, there are other approaches that give insights on the geometry of the indicatrix, see [10].

²Later we shall introduce the covariant exterior differential D of ∇ so that this equation can also be written $\bar{\omega}^a = Dy^a$.

³A Finsler connection is *strongly regular* if $\nabla y|_{VE} = Id_{VE}$. All notable connection share this property, but we do not impose it as this condition would constrain the variations.

constraint as in the Einstein-Cartan theory (with no assumption on the torsion, which in Finsler geometry leaves up much freedom).

Our work presents at once two equivalent theories. One theory, which is obtained removing all the ‘tildes’ in the following expressions, imposes a priori metricity for the Finsler connection, much as in the Einstein-Cartan theory. This is not a fully metric-affine theory. The other theory is obtained by implementing in the action the following *amplified symmetry* from our previous work [5]

$$\omega^a_b \rightarrow \omega^a_b + A^a_b, \quad A_{ab} = A_{ba}$$

where ω^a_b are the coefficients for the Finsler connection and the indices are lowered with the metric g_{ac} . In this theory one really works with equivalence classes of connections and the theory is fully metric-affine. A posteriori one can then select some specific connection from the solution class, such as the metric representative.

In this second approach we expect that the dependence on the linear connection would be mitigated and that the true variable of the theory could end up being the non-linear connection. Whether this is the case depends on the matter Lagrangian, which might have a restricted invariance. For instance, it might impose the additional condition $A_{ab}y^b = 0$. Then, due to the equation $HE = \ker \nabla y$, each element in a restricted amplified symmetry class would share the same non-linear connection, which shows that the action would really depend on the non-linear connection, see also [13] for some mathematical discussion on this restricted amplified symmetry.

We mention that the notable connections of Finsler geometry previously mentioned all belong to the same class of connections, two connections being equivalent iff connected by such transformation.

The implementation of such an amplified symmetry means modifying the Einstein-Cartan action on E to allow for non-metricity. The way we implement the amplified symmetry is by use of the following *metric representative Finsler connection* $\tilde{\nabla}$

$$\tilde{\omega}^a_b := \omega^a_b + \frac{1}{2}g^{ar}Dg_{rb}$$

where D is the covariant exterior differential induced by ∇ . Similarly, we shall denote with \tilde{D} the covariant exterior differential induced by $\tilde{\nabla}$, and in general any object dependent on $\tilde{\nabla}$ will carry a tilde. The connection $\tilde{\nabla}$ is indeed metric and its curvature 2-form \tilde{R} is related to that of ∇ , R , by

$$\tilde{R}_{ab} = R_{[ab]} - \frac{1}{4}Dg_{ac} \wedge g^{cs}Dg_{sb}.$$

When ∇ is one of the notable Finsler connections, $\tilde{\nabla}$ is the Cartan connection. The gravitational action studied in this work is

$$S_G(g_{ab}, \omega^a_b, e^a, s) := \int_M s^*(L_G), \quad L_G = \frac{1}{2}\eta_{abcd}\tilde{R}^{ab} \wedge \omega^c \wedge \omega^d \quad (1)$$

where $\omega^a = \pi^*e^a$ and where $\eta_{abcd} = \sqrt{|\det g(x, y)|}[abcd]$ is the Finslerian volume tensor. We shall restrict ourselves to the 4-dimensional spacetime case, but the generalization to different dimensions will be obvious.

As mentioned, although we shall work with the amplified symmetry, much in analogy with [5], the paper also includes the case in which this symmetry is not implemented which is recovered by removing all the ‘tildes’ and by assuming that ∇ is metric (but a priori not necessarily Cartan’s).

Save for the pullback s^* , and the use of the connection $\tilde{\nabla}$ to define the curvature, the Lagrangian is Einstein-Cartan’s. In the Einstein-Cartan theory one obtains that torsion can be present only inside matter, and in fact, perhaps not surprisingly, we shall obtain a similar result in the Finslerian case. However, all Finsler connections have some form of torsion unless the space is really non-Finslerian, so ultimately we shall obtain that in vacuum the spacetime is non-Finslerian, i.e. g does not depend on the fiber variables y .

1.1 Energy-momentum conservation

Independently of the previous discussion, there is another line of thought that brought us to consider a Lagrangian of the above form. A well known problem in Finsler geometry is that of establishing some form of conservation principle. The conservation of the stress-energy tensor was indeed one of the demands that led Einstein to the development of the general theory of relativity [14, 4]. Due to the abundance of potential field equations in Finsler gravity, a similar criteria could guide the selection of the correct ones.

We do not think, however, that in the Finslerian case one should search for a conserved stress-energy tensor but rather, as suggested in [11, Remark 19], for a conserved observer-dependent energy-momentum vector, or equivalently, 3-form (for other recent approaches and formalisms see [7, 9]).

In this regard we observe that the following modified Einstein’s vector-valued 3-form

$$\tilde{\tau}_d := \frac{1}{2}\eta_{abcd}\tilde{R}^{ab} \wedge \omega^c$$

satisfies $\tilde{\tau}_d \wedge \omega^d = L_G$ and

$$\tilde{D}\tilde{\tau}_d = \frac{1}{2}\eta_{abcd}\tilde{R}^{ab} \wedge \Psi^c \quad (2)$$

where, denoting with T^c the horizontal torsion for D ,

$$\Psi^c := \tilde{D}\omega^c = T^c + \frac{1}{2}g^{cr}Dg_{rb} \wedge \omega^b$$

is the horizontal torsion of the connection \tilde{D} . Recalling that $\bar{\omega}^c = Dy^c$ (which jointly with ω^c form a basis for T^*E) and similarly $\tilde{\omega}^c = \tilde{D}y^c$, we have, contracting (2) with the Liouville vector field

$$d(\tilde{\tau}_d y^d) = \frac{1}{2}\eta_{abcd}y^d \tilde{R}^{ab} \wedge \Psi^c - \tilde{\tau}_d \wedge \tilde{\omega}^d.$$

Every form can be expressed as a linear combination of wedge products of basis forms $\{\omega^c, \bar{\omega}^c\}$, so leading to various decompositions in horizontal-vertical parts. A similar statement applies if we take the basis $\{\omega^c, \tilde{\omega}^c\}$. We denote with Hor the operator that sends all the vertical forms to zero, in other words Hor

keeps only the terms that have a totally horizontal expansion. Similarly $\widetilde{\text{Hor}}$ does the same, but for the non-linear connection induced by $\widetilde{\nabla}$.

The nice fact is that for a notable connection $\widetilde{\Psi}^a = C^a_{bc}\tilde{\omega}^c \wedge \omega^b$ where $C_{abc} = \frac{1}{2}\frac{\partial}{\partial y^a}g_{bc}$ is the Cartan torsion, and hence $\widetilde{\text{Hor}}(\widetilde{\Psi}^a) = 0$. Whenever this equation holds the previous expression in display satisfies

$$\widetilde{\text{Hor}}(d(\tilde{\tau}_d y^d)) = 0.$$

When we pullback with a section $s : M \rightarrow E$, we can make use of the identity $s^*(\tilde{\omega}^c) =: \tilde{D}s^c$ where, with some abuse of notation, we denote with $\tilde{D}s^c$ the non-linear covariant derivative of the section. This non-linear covariant derivative is a 1-form that vanishes at a point precisely when the section represents a free falling non-rotating non-expanding reference frame (observer) at the point. Such observers exist at every point due to the fact that for every chosen point of E , linear coordinates can be chosen so that the connection coefficients vanish at the point [12].

We conclude that at that point and for such a frame the 3-form $s^*(\tilde{\tau}_d y^d)$ is closed

$$d(s^*(\tilde{\tau}_d y^d)) = 0. \quad (3)$$

Now assume dynamical equations of the form

$$\tilde{\tau}_d = t_d, \quad (4)$$

or more weakly $y^d \tilde{\tau}_d = y^d t_d$, where t_d is the energy-momentum 3-form and $y^d t_d$ is the energy-momentum of matter as viewed by an observer with velocity of direction y^d .

Observe that the equation (3), now written $d(s^*(t_d y^d)) = 0$, can be attained at any chosen point through a suitable choice of observer. However, it cannot be obtained locally or globally as, in general, a section such that $\tilde{D}s = 0$ does not exist locally. Still the conservation can be approximately expressed in integral form in a small neighborhood of the point where the previous equation holds true. So for a cylindrical neighborhood C , with basis B_1 , B_2 and lateral side S , we shall have

$$\int_{B_1} s^*(t_d y^d) + \int_{B_2} s^*(t_d y^d) + \int_S s^*(t_d y^d) = o(\text{size of cylinder}).$$

Using the orientation coming from the time orientation of the spacetime, one of the two first terms of the previous expression changes sign, and we get an expression which states that the energy-momentum $s^*(t_d)s^d$ entering C from the basis B_1 is equal to that escaping B_2 plus that entering the lateral side S . The contraction with s^d expresses the fact that the quantity $s^*(t_d)s^d$ is an energy-momentum 3-form (dually a vector) dependent on the chosen observer s rather than a tensor. This analysis is compatible and in the same spirit of that performed in [11, Remark 19], but here we do not require that the Landsberg tensor or mean Cartan torsion vanish.

In the next section we shall see that our action varied with respect to the vierbein indeed gives an equation of the form $\tilde{\tau}_d = t_d$. We stress that according to this theory conservation of energy-momentum can be expected only in free-falling non-rotating non-expanding reference frames and only locally. Both features are shared with classical general relativity (observe that in general relativity, if u^a is 4-velocity field, $(T^{ab}u_b)_{;a} = T^{ab}u_{a;b} \neq 0$).

Finally, we mention that in the non-Finslerian case, with $\tilde{\nabla}$ the Levi-Civita connection, we indeed recover from the previous formulas the expected results for general relativity, as we have

$$\tilde{\tau}_d = \frac{1}{6}G_d{}^r \eta_{ruvz} \omega^u \wedge \omega^v \wedge \omega^z$$

where $G_{ab} = R_{ab} - \frac{1}{2}Rg_{ab}$ is the Einstein tensor.

1.2 Action variation

In this section we shall investigate whether it is possible to obtain the dynamical equation (4) through the variation of an action. We shall be concerned in recovering just the left-hand side of (4) which has gravitational origin.

Let $T^a = D\omega^a$ and $\bar{T}^a = D\bar{\omega}^a$ be the horizontal and vertical torsions respectively, and let $\mathcal{G}_{ab} = Dg_{ab}$ be the non-metricity, all with reference to the same Finsler connection ∇ . It is useful to recall the next Bianchi identities which hold for any Finsler connection (including \tilde{D} which is metric, hence the last one establishes the antisymmetry of \tilde{R}_{ab})

$$\begin{aligned} DR^a{}_b &= 0, \\ DT^a &= R^a{}_b \wedge \omega^b, \\ D\bar{T}^a &= R^a{}_b \wedge \bar{\omega}^b, \\ D\mathcal{G}_{ab} &= -R_{ab} - R_{ba}. \end{aligned}$$

Let us consider the gravitational Einstein-Cartan-like action (1). We shall often use the following observation. By using normal coordinates at a point of E it is easy to check that given a non-vanishing k -vector on TM , and a k -vector on TE that projects on it, it is always possible to find a section $s : M \rightarrow E$ such that the former is sent to the latter (in normal coordinates it is sufficient to take $s^a = A^a_b x^b$ where A is a suitable matrix). Since any k -vector on TE can be approximated by a k -vector that projects to a non-vanishing k -vector we have, by continuity, that given any k -form ω on E , the equality $s^*\omega = 0$ holds for every s iff $\omega = 0$ holds (see the Appendix for a detailed derivation). This property shall explain why the pullback can be omitted from our equations, as they are supposed to hold for every section.

Another useful fact, that we shall systematically use in our calculations, is the commutativity of the exterior derivative with the pullback $ds^*\alpha = s^*d\alpha$, where α is a form on E . Ultimately, it will allow us to get rid of some exact terms.

The general variation for fixed section can be written, using $s^*\pi^*e^a = e^a$

$$\delta S_G = \int_M s^* \left(\frac{\delta L_G}{\delta \omega^a} \right) \wedge \delta e^a + s^* \left(\frac{\delta L_G}{\delta \omega^a_b} \wedge \delta \omega^a_b \right) + s^* \left(\frac{\delta L_G}{\delta g^{ab}} \delta g^{ab} \right).$$

Now, using the identity

$$\begin{aligned} \delta(\omega^c \wedge \tilde{\tau}_c) &= d \left(\eta^a_{bcd} \omega^c \wedge \left(\frac{1}{2} \omega^d \wedge \delta \omega^b_a - \frac{1}{4} \delta g_{ae} \omega^b \wedge Dg^{de} \right) \right) - \\ &- 2\tilde{\tau}_a \wedge \delta \omega^a + \eta^a_{bcd} \Psi^b \wedge \omega^d \wedge \delta \omega^c_a \\ &- \frac{1}{2} \left(g^{ab} \tilde{\tau}_c \wedge \omega^c + \eta^a_{cde} \left(\Psi^c \wedge \omega^d \wedge Dg^{be} + \omega^c \wedge \omega^d \wedge \tilde{R}^{eb} \right) \right) \delta g_{ab}, \end{aligned} \quad (5)$$

we obtain

$$\begin{aligned} \frac{\delta L_G}{\delta \omega^d} &= 2\tilde{\tau}_d = \eta_{abcd} \tilde{R}^{ab} \wedge \omega^c, \\ \frac{\delta L_G}{\delta \omega^a_r} &= -g^{rb} \eta_{abcd} \Psi^c \wedge \omega^d, \\ \frac{\delta L_G}{\delta g^{ab}} &= -\frac{\delta L_G}{\delta g_{mn}} g_{ma} g_{nb} = \\ &- \frac{1}{2} \eta_{ecd(a} Dg_{b)}^e \wedge \Psi^c \wedge \omega^d + \frac{1}{2} \eta_{rcd(b} \tilde{R}^r_a) \wedge \omega^c \wedge \omega^d - \frac{1}{2} g_{ab} L_G. \end{aligned}$$

In the second equation we used, for g^{ab} fixed, $\delta \tilde{\omega}_{ab} = \delta \omega_{[ab]}$. Actually these variational derivatives are not independent as the invariance under change of frame implies the identity

$$\frac{\delta L_G}{\delta \omega^a} \wedge \omega^b = 2 \frac{\delta L_G}{\delta g_{rb}} g_{ra} + D \frac{\delta L_G}{\delta \omega^a_b}. \quad (6)$$

As a consequence, it is sufficient to consider the dynamical equations for the connection and vierbeins.

Further, it can be observed that under a variation of section

$$\delta S_G = \int_M \delta s^*(L_G),$$

but $L_G = \tilde{\tau}_d \wedge \omega^d$ thus, as in vacuum $\tilde{\tau}_d = 0$, the action does not really depend on the section when the gravitational fields are on-shell. It remains to establish the consequences of the vacuum equations. Let

$$\tilde{\omega}^a_b = \tilde{H}^a_{bc} \omega^c + \tilde{V}^a_{bc} \tilde{\omega}^c,$$

where $\tilde{\omega}^c = \tilde{D}y^c$, be the expansion for the connection coefficients. Let $[e_a, e_b] = c^c_{ab}(x)e_c$ be the commutation relations for the chosen basis, so that $de^c = -\frac{1}{2}c^c_{ab}e^a \wedge e^b$. Then

$$\Psi^a = \tilde{D}\omega^a = (\tilde{H}^a_{bc} + \frac{1}{2}c^a_{bc})\omega^c \wedge \omega^b + \tilde{V}^a_{bc}\tilde{\omega}^c \wedge \omega^b.$$

The vacuum equation for the connection $\frac{\delta L_G}{\delta \omega^{ar}} = 0$ can be rewritten

$$\Psi^a \wedge \omega^b - \Psi^b \wedge \omega^a = 0 \quad (7)$$

which implies $\widetilde{\text{Hor}}(\Psi^a) = 0$ and hence $\tilde{H}^a{}_{[bc]} + \frac{1}{2}c^a{}_{[bc]} = 0$. But metricity of \tilde{D} implies, in particular, $\widetilde{\text{Hor}}(\tilde{D}g_{ab}) = 0$, that is, (in this expression $\frac{\tilde{\delta}}{\delta x^c} = \frac{\partial}{\partial x^c} - \tilde{N}_c^r(x, y) \frac{\partial}{\partial y^r}$ where \tilde{N}_c^r are the coefficients of the non-linear connection as determined by the regularity condition.)

$$0 = \frac{\tilde{\delta} g_{ab}}{\delta x^c} - \tilde{H}_{abc} - \tilde{H}_{bac}.$$

We arrive at

$$\tilde{H}^a{}_{bc} = g^{ar} \left\{ \frac{1}{2} \left(\frac{\tilde{\delta}}{\delta x^r} g_{ra} + \frac{\tilde{\delta}}{\delta x^a} g_{rb} - \frac{\tilde{\delta}}{\delta x^r} g_{ab} \right) - \frac{1}{2} (c_{acb} + c_{bac} - c_{cba}) \right\} \quad (8)$$

Let us consider the vertical information in Eq. (7). The contractions $i_{\tilde{\delta}/\delta x^c} i_Y$ of Eq. (7) where Y is vertical give

$$\Psi^a(Y) \delta_c^b - \Psi^b(Y) \delta_c^a = 0.$$

Taking the trace we get $\Psi^a(Y) = 0$ which means that Ψ^a is purely horizontal.

Since Ψ^a is purely horizontal $\tilde{V}^a{}_{bc} = 0$, which implies that \tilde{D} is strongly regular, cf. [13]. But \tilde{D} is metric, so a calculation of the vertical part of $\tilde{D}g_{ab} = 0$ gives $\frac{\partial g_{ab}}{\partial y^c} = 0$, namely the metric does not depend on the vertical variables. Finally, the expression (8) proves that in vacuum $\tilde{\nabla}$ does not depend on the vertical variables and it is actually the Levi-Civita connection of g_{ab} .

The original connection ∇ can only be determined up to a gauge because we implemented the amplified symmetry. The same Lagrangian with all the ‘tilde’ dropped would have established that the connection in vacuum is indeed Levi-Civita. It is indeed possible to impose that the connection is metric since the beginning. That gives a viable and physically equivalent approach, the only drawback being that the variation has to be constrained a priori since only metric connections should be considered (as in the Einstein-Cartan theory). Therefore, it is not an approach in which metric and connection are completely unconstrained (purely metric-affine).

In any case, the conclusion is that in vacuum the theory is equivalent general relativity, so according to the present theory in vacuum there is no torsion nor anisotropy.

2 Conclusions

We explored an anisotropic theory in which an Einstein-Cartan-like Lagrangian on the slit tangent bundle E is pulled back to M through a section $s : M \rightarrow E$. The dynamics was obtained by imposing stationarity for every possible section s , which results in equations on E independent of the section. This fact is

interpreted as independence of the equations from the observer. The dynamical equations imply that in vacuum the theory is in fact coincident with general relativity and hence that there is no torsion nor anisotropy. The theory might also implement the amplified symmetry of our previous work. This modification, motivated by the purpose of interpreting the physical field as the non-linear connection, can also be omitted in which case the theory is framed in terms of a metric connection, this time living on E rather than M , much as in the original Einstein-Cartan theory.

Appendix: Proof of $\forall s, s^*\omega = 0 \Rightarrow \omega = 0$

In this appendix we provide a detailed argument on why $s^*\omega = 0$ for every section $s : M \rightarrow E$ implies $\omega = 0$. Here ω is a k -form on E , $k \leq n$, n dimension of M .

Let $E = TM \setminus 0$ and let $\pi : E \rightarrow M$ be the projection on the base. The coordinates on E are (x^a, y^a) , those on TE are $(x^a, y^a, \dot{x}^a, \dot{y}^a)$. Elements in TE are denoted with capital letters, e.g. V , those in TM with lowercase letters, e.g. v .

Suppose $s^*\omega = 0$ for every s . In order to prove that $\omega = 0$ we just need to prove that ω vanishes over every k -vector $V_1 \wedge V_2 \wedge \dots \wedge V_k$ with $V_i \in TE$.

We can assume that $v_i = \pi(V_i)$ are linearly independent and hence that

$$\pi(V_1 \wedge V_2 \wedge \dots \wedge V_k) = v_1 \wedge v_2 \wedge \dots \wedge v_k \neq 0$$

indeed, if not, replace V_i with $V'_i(\epsilon) = V_i + \epsilon W_i$ for sufficiently small ϵ where $w_i = \pi(W_i)$ are linearly independent (if $t < k$ is the dimension of the space spanned by $\{v_i\}$ pick t vectors of $\{w_i\}$ as a basis of this space and $k - t$ vectors as a basis for a trasverse subspace). Then $\{V'_i\}$ would have linearly independent projections, and if we can prove for these type of k -vectors that

$$\omega(V'_1 \wedge V'_2 \wedge \dots \wedge V'_k) = 0,$$

then taking the limit $\epsilon \rightarrow 0$ we also get

$$\omega(V_1 \wedge V_2 \wedge \dots \wedge V_k) = 0.$$

Let $P \in E$, we want to prove that $\omega(P) = 0$. Let $p = \pi(P)$. We need only to try the form on $V_1 \wedge V_2 \wedge \dots \wedge V_k$ with projection $v_1 \wedge v_2 \wedge \dots \wedge v_k \neq 0$, that is, we can assume that v_i are linearly independent.

Introduce coordinates at the point $p \in M$ of interest such that $x^a(p) = 0$. Consider the section $s : M \rightarrow E$ having local components

$$s^a(x) = P^a + B^a_b x^b$$

where $P^a = y^a(P)$ and B is a constant $n \times n$ matrix. Note that

$$V_i = v_i^a \frac{\partial}{\partial x^a} + q_i^b \frac{\partial}{\partial y^b}$$

for some constants q_i^b . Choose B so that it maps $v_i^b \in \mathbb{R}^n$ to $q_i^b \in \mathbb{R}^n$, i.e. $q_i^b = B^b{}_a v_i^a$ (here linear independence of $\{v_i\}$ is used), then

$$s_*(v_i) = v_i^a \frac{\partial}{\partial x^a} + \frac{\partial s^a}{\partial x^c} v_i^c \frac{\partial}{\partial y^a} = V_i \in TE,$$

and hence

$$\omega(V_1 \wedge V_2 \wedge \cdots \wedge V_k) = s^* \omega(v_1 \wedge v_2 \wedge \cdots \wedge v_k) = 0.$$

Acknowledgments

A.G.-P. is supported by projects PY20-01391 and UCO-1380930 from the Regional Government of Andalusia (Spain) and ERDEF (UE). E.M. is partially supported by GNFM of INDAM.

Data Availability Statement. All data generated or analysed during this study are included in this published article.

References

- [1] H. Akbar-Zadeh. Generalized Einstein manifolds. *J. Geom. Phys.*, 17:342–380, 1995.
- [2] D. Bao, S.-S. Chern, and Z. Shen. *An Introduction to Riemann-Finsler Geometry*. Springer-Verlag, New York, 2000.
- [3] B. Chen and Y.B. Shen. On a class of critical Riemann-Finsler metrics. *Publ. Math. Debrecen*, 72:451–468, 2008.
- [4] A. Einstein. Die Grundlage der allgemeinen Relativitätstheorie. *Annalen der Physik*, 49:770–822, 1916.
- [5] A. García-Parrado and E. Minguzzi. Projective and amplified symmetries in metric-affine theories. *Class. Quantum Grav.*, 38:135001, 2021. arXiv:2006.04040.
- [6] M. Hohmann, C. Pfeifer, and N. Voicu. Finsler gravity action from variational completion. *Phys. Rev. D*, 100:064035, 2019.
- [7] M. Hohmann, C. Pfeifer, and N. Voicu. Mathematical foundations for field theories on Finsler spacetimes. *J. Math. Phys.*, 63:032503, 2022.
- [8] M. A. Javaloyes, M. Sánchez, and F.F. Villaseñor. The Einstein-Hilbert-Palatini formalism in pseudo-Finsler geometry. arXiv:2108.03197, 2021.
- [9] M. A. Javaloyes, M. Sánchez, and F.F. Villaseñor. On the significance of the stress-energy tensor in Finsler spacetimes. *Universe*, 8, 2021.

- [10] O. Makhmali. Differential geometric aspects of causal structures. *SIGMA*, 14:080, 2018.
- [11] E. Minguzzi. The connections of pseudo-Finsler spaces. *Int. J. Geom. Meth. Mod. Phys.*, 11:1460025, 2014. Erratum *ibid* 12 (2015) 1592001. arXiv:1405.0645.
- [12] E. Minguzzi. Special coordinate systems in pseudo-Finsler geometry and the equivalence principle. *J. Geom. Phys.*, 114:336–347, 2017. arXiv:1601.07952.
- [13] E. Minguzzi. A metrical approach to Finsler geometry. arXiv:2107.09227, 2021.
- [14] C. W. Misner, K. S. Thorne, and J. A. Wheeler. *Gravitation*. Freeman, San Francisco, 1973.
- [15] P. Ni and B. Shen. On variation of action integral in Finsler gravity. *Ann. Phys.*, 404:93–114, 2019.
- [16] C. Pfeifer and M. N. R. Wohlfarth. Finsler geometric extension of Einstein gravity. *Phys. Rev. D*, 85:064009, 2012.
- [17] Y.-B. Shen and Z. Shen. *Introduction to modern Finsler geometry*. Higher Education Press/World Scientific, Beijing/Singapore, 2016.