# Two results of *n*-exangulated categories

Jian He, Jing He and Panyue Zhou

#### Abstract

*n*-exangulated categories were introduced by Herschend-Liu-Nakaoka which are a simultaneous generalization of *n*-exact categories and (n + 2)-angulated categories. This paper consists of two results on *n*-exangulated categories: (1) we give an equivalent characterization of the axiom (EA2); (2) we provide a new way to construct a closed subfunctor of an *n*-exangulated category.

Keywords: *n*-exangulated categories; homotopy cartesian square; half exact functor 2020 Mathematics Subject Classification: 18G80; 18G50; 18G25.

## 1 Introduction

The notion of extriangulated categories was introduced in [7], which can be viewed as a simultaneous generalization of exact categories and triangulated categories. The data of such a category is a triplet  $(\mathscr{C}, \mathbb{E}, \mathfrak{s})$ , where  $\mathscr{C}$  is an additive category,  $\mathbb{E} \colon \mathscr{C}^{\text{op}} \times \mathscr{C} \to \mathsf{Ab}$  is an additive bifunctor and  $\mathfrak{s}$  assigns to each  $\delta \in \mathbb{E}(C, A)$  a class of 3-term sequences with end terms A and C such that certain axioms hold. Recently, Herschend–Liu–Nakaoka [1] introduced the notion of n-exangulated categories for any positive integer n. It should be noted that the case n = 1corresponds to extriangulated categories. As typical examples we know that n-exact categories and (n + 2)-angulated categories are n-exangulated categories, see [1, Proposition 4.34] and [1, Proposition 4.5]. However, there are some other examples of n-exangulated categories which are neither n-exact nor (n + 2)-angulated, see [1–3, 6].

Recall that homotopy cartesian squares in triangulated categories are the triangulated analogues of the pushout and pullback squares in abelian categories. It is well-known that the axiom (TR4) is equivalent to the homotopy cartesian axiom. Recently, Kong-Lin-Wang [4] introduced the notion of homotopy cartesian squares and proposed a new shifted octahedron in extriangulated categories. Our first main result show that Kong-Lin-Wang's result has a higher counterpart:

**Theorem 1.1.** (see Theorem 3.4 for details) Let  $(\mathscr{C}, \mathbb{E}, \mathfrak{s})$  be a pre-*n*-exangulated category. Then  $\mathscr{C}$  satisfies (EA2) if and only if  $\mathscr{C}$  satisfies (EA2-1).

Relative theories are explicated by use of closed subfunctors in *n*-exangulated categories [1]. An additive subfunctor  $\mathbb{F}$  of  $\mathbb{E}$  is a closed subfunctor if  $(\mathscr{C}, \mathbb{F}, \mathfrak{s}|_{\mathbb{F}})$  is an *n*-exangulated category,

Jian He was supported by the National Natural Science Foundation of China (Grant No. 12171230). Panyue Zhou was supported by the National Natural Science Foundation of China (Grant No. 11901190).

where  $\mathfrak{s}|_{\mathbb{F}}$  is a restriction of  $\mathfrak{s}$  to  $\mathbb{F}$ , see [1, Proposition 3.16]. Our second main result provide a new way to construct a closed subfunctor of an *n*-exangulated category. This is a higher counterpart of Sakai's result.

**Theorem 1.2.** (see Theorem 4.5 for details) Let  $(\mathscr{C}, \mathbb{E}, \mathfrak{s})$  be an *n*-exangulated category and  $F: \mathscr{C} \to \mathscr{A}$  a half exact functor to an abelian category  $\mathscr{A}$ . Then there exists a unique maximal closed subfunctor  $\mathbb{F}$  of  $\mathbb{E}$  such that F becomes a right exact functor from  $(\mathscr{C}, \mathbb{F}, \mathfrak{s}|_{\mathbb{F}})$  to  $\mathscr{A}$ .

This article is organized as follows. In Section 2, we review some elementary definitions and facts on n-exangulated categories. In Section 3, we prove our first main result. In Section 4, we prove our second main result.

### 2 Preliminaries

In this section, we briefly review basic concepts and results concerning n-exangulated categories.

Let  $\mathscr{C}$  be an additive category and  $\mathbb{E}: \mathscr{C}^{\operatorname{op}} \times \mathscr{C} \to \mathsf{Ab}$  (Ab is the category of abelian groups) an additive bifunctor. For any pair of objects  $A, C \in \mathscr{C}$ , an element  $\delta \in \mathbb{E}(C, A)$  is called an  $\mathbb{E}$ extension or simply an extension. We also write such  $\delta$  as  ${}_{A}\delta_{C}$  when we indicate A and C. The zero element  ${}_{A}0_{C} = 0 \in \mathbb{E}(C, A)$  is called the *split*  $\mathbb{E}$ -extension. For any pair of  $\mathbb{E}$ -extensions  ${}_{A}\delta_{C}$  and  ${}_{A'}\delta'_{C'}$ , let  $\delta \oplus \delta' \in \mathbb{E}(C \oplus C', A \oplus A')$  be the element corresponding to  $(\delta, 0, 0, \delta')$  through the natural isomorphism  $\mathbb{E}(C \oplus C', A \oplus A') \simeq \mathbb{E}(C, A) \oplus \mathbb{E}(C, A') \oplus \mathbb{E}(C', A) \oplus \mathbb{E}(C', A')$ .

For any  $a \in \mathscr{C}(A, A')$  and  $c \in \mathscr{C}(C', C)$ ,  $\mathbb{E}(C, a)(\delta) \in \mathbb{E}(C, A')$  and  $\mathbb{E}(c, A)(\delta) \in \mathbb{E}(C', A)$ are simply denoted by  $a_*\delta$  and  $c^*\delta$ , respectively.

Let  ${}_{A}\delta_{C}$  and  ${}_{A'}\delta'_{C'}$  be any pair of  $\mathbb{E}$ -extensions. A morphism  $(a, c): \delta \to \delta'$  of extensions is a pair of morphisms  $a \in \mathscr{C}(A, A')$  and  $c \in \mathscr{C}(C, C')$  in  $\mathscr{C}$ , satisfying the equality  $a_*\delta = c^*\delta'$ .

Let  $\mathscr{C}$  be an additive category as before, and let n be any positive integer.

**Definition 2.1.** [1, Definition 2.7] Let  $\mathbf{C}_{\mathscr{C}}$  be the category of complexes in  $\mathscr{C}$ . As its full subcategory, define  $\mathbf{C}_{\mathscr{C}}^{n+2}$  to be the category of complexes in  $\mathscr{C}$  whose components are zero in the degrees outside of  $\{0, 1, \ldots, n+1\}$ . Namely, an object in  $\mathbf{C}_{\mathscr{C}}^{n+2}$  is a complex  $X_{\bullet} = \{X_i, d_i^X\}$  of the form

$$X_0 \xrightarrow{d_0^X} X_1 \xrightarrow{d_1^X} \cdots \xrightarrow{d_{n-1}^X} X_n \xrightarrow{d_n^X} X_{n+1}.$$

We write a morphism  $f_{\bullet}: X_{\bullet} \to Y_{\bullet}$  simply  $f_{\bullet} = (f_0, f_1, \dots, f_{n+1})$ , only indicating the terms of degrees  $0, \dots, n+1$ .

**Definition 2.2.** [1, Definition 2.11] By Yoneda lemma, any extension  $\delta \in \mathbb{E}(C, A)$  induces natural transformations

$$\delta_{\sharp} \colon \mathscr{C}(-, C) \Rightarrow \mathbb{E}(-, A) \text{ and } \delta^{\sharp} \colon \mathscr{C}(A, -) \Rightarrow \mathbb{E}(C, -).$$

For any  $X \in \mathscr{C}$ , these  $(\delta_{\sharp})_X$  and  $\delta_X^{\sharp}$  are given as follows.

(1)  $(\delta_{\sharp})_X \colon \mathscr{C}(X, C) \to \mathbb{E}(X, A) : f \mapsto f^* \delta.$ 

(2)  $\delta_X^{\sharp} : \mathscr{C}(A, X) \to \mathbb{E}(C, X) : g \mapsto g_* \delta$ 

We simply denote  $(\delta_{\sharp})_X(f)$  and  $\delta_X^{\sharp}(g)$  by  $\delta_{\sharp}(f)$  and  $\delta^{\sharp}(g)$ , respectively.

**Definition 2.3.** [1, Definition 2.9] Let  $\mathscr{C}, \mathbb{E}, n$  be as before. Define a category  $\mathscr{E} := \mathscr{E}^{n+2}_{(\mathscr{C},\mathbb{E})}$  as follows.

(1) A object in  $\mathbb{R}^{n+2}_{(\mathscr{C},\mathbb{E})}$  is a pair  $\langle X_{\bullet}, \delta \rangle$  of  $X_{\bullet} \in \mathbb{C}^{n+2}_{\mathscr{C}}$  and  $\delta \in \mathbb{E}(X_{n+1}, X_0)$  satisfying

$$(d_0^X)_*\delta = 0$$
 and  $(d_n^X)^*\delta = 0.$ 

We call such a pair an  $\mathbb{E}$ -attached complex of length n + 2. We also denote it by

$$X_0 \xrightarrow{d_0^X} X_1 \xrightarrow{d_1^X} \cdots \xrightarrow{d_{n-2}^X} X_{n-1} \xrightarrow{d_{n-1}^X} X_n \xrightarrow{d_n^X} X_{n+1} \xrightarrow{\delta} .$$

(2) For such pairs  $\langle X_{\bullet}, \delta \rangle$  and  $\langle Y_{\bullet}, \rho \rangle$ , a morphism  $f_{\bullet} \colon \langle X_{\bullet}, \delta \rangle \to \langle Y_{\bullet}, \rho \rangle$  is defined to be a morphism  $f_{\bullet} \in \mathbf{C}^{n+2}_{\mathscr{C}}(X_{\bullet}, Y_{\bullet})$  satisfying  $(f_0)_* \delta = (f_{n+1})^* \rho$ .

We use the same composition and the identities as in  $\mathbf{C}_{\mathscr{C}}^{n+2}$ .

**Definition 2.4.** [1, Definition 2.13] An *n*-example is a pair  $\langle X_{\bullet}, \delta \rangle$  of  $X_{\bullet} \in \mathbf{C}_{\mathscr{C}}^{n+2}$  and  $\delta \in \mathbb{E}(X_{n+1}, X_0)$  which satisfies the following conditions.

(1) The following sequence of functors  $\mathscr{C}^{\mathrm{op}} \to \mathsf{Ab}$  is exact.

$$\mathscr{C}(-,X_0) \xrightarrow{\mathscr{C}(-, d_0^X)} \cdots \xrightarrow{\mathscr{C}(-, d_n^X)} \mathscr{C}(-,X_{n+1}) \xrightarrow{\delta_{\sharp}} \mathbb{E}(-,X_0)$$

(2) The following sequence of functors  $\mathscr{C} \to \mathsf{Ab}$  is exact.

$$\mathscr{C}(X_{n+1},-) \xrightarrow{\mathscr{C}(d_n^X,-)} \cdots \xrightarrow{\mathscr{C}(d_0^X,-)} \mathscr{C}(X_0,-) \xrightarrow{\delta^{\sharp}} \mathbb{E}(X_{n+1},-)$$

In particular any *n*-example is an object in  $\mathcal{E}$ . A morphism of *n*-examples simply means a morphism in  $\mathcal{E}$ . Thus *n*-examples form a full subcategory of  $\mathcal{E}$ .

**Definition 2.5.** [1, Definition 2.22] Let  $\mathfrak{s}$  be a correspondence which associates a homotopic equivalence class  $\mathfrak{s}(\delta) = [{}_{A}X_{\bullet C}]$  to each extension  $\delta = {}_{A}\delta_{C}$ . Such  $\mathfrak{s}$  is called a *realization* of  $\mathbb{E}$  if it satisfies the following condition for any  $\mathfrak{s}(\delta) = [X_{\bullet}]$  and any  $\mathfrak{s}(\rho) = [Y_{\bullet}]$ .

(R0) For any morphism of extensions  $(a, c): \delta \to \rho$ , there exists a morphism  $f_{\bullet} \in \mathbf{C}^{n+2}_{\mathscr{C}}(X_{\bullet}, Y_{\bullet})$ of the form  $f_{\bullet} = (a, f_1, \ldots, f_n, c)$ . Such  $f_{\bullet}$  is called a *lift* of (a, c).

In such a case, we simply say that "X<sub>•</sub> realizes  $\delta$ " whenever they satisfy  $\mathfrak{s}(\delta) = [X_{\bullet}]$ .

Moreover, a realization  $\mathfrak{s}$  of  $\mathbb{E}$  is said to be *exact* if it satisfies the following conditions.

- (R1) For any  $\mathfrak{s}(\delta) = [X_{\bullet}]$ , the pair  $\langle X_{\bullet}, \delta \rangle$  is an *n*-example.
- (R2) For any  $A \in \mathscr{C}$ , the zero element  ${}_A 0_0 = 0 \in \mathbb{E}(0, A)$  satisfies

$$\mathfrak{s}(A0_0) = [A \xrightarrow{\mathrm{id}_A} A \to 0 \to \dots \to 0 \to 0].$$

Dually,  $\mathfrak{s}(_00_A) = [0 \to 0 \to \cdots \to 0 \to A \xrightarrow{\mathrm{id}_A} A]$  holds for any  $A \in \mathscr{C}$ .

Note that the above condition (R1) does not depend on representatives of the class  $[X_{\bullet}]$ .

**Definition 2.6.** [1, Definition 2.23] Let  $\mathfrak{s}$  be an exact realization of  $\mathbb{E}$ .

- (1) An *n*-example  $\langle X_{\bullet}, \delta \rangle$  is called an  $\mathfrak{s}$ -distinguished *n*-example if it satisfies  $\mathfrak{s}(\delta) = [X_{\bullet}]$ . We often simply say distinguished *n*-example when  $\mathfrak{s}$  is clear from the context.
- (2) An object  $X_{\bullet} \in \mathbf{C}_{\mathscr{C}}^{n+2}$  is called an  $\mathfrak{s}$ -conflation or simply a conflation if it realizes some extension  $\delta \in \mathbb{E}(X_{n+1}, X_0)$ .
- (3) A morphism f in  $\mathscr{C}$  is called an  $\mathfrak{s}$ -inflation or simply an inflation if it admits some conflation  $X_{\bullet} \in \mathbf{C}_{\mathscr{C}}^{n+2}$  satisfying  $d_0^X = f$ .
- (4) A morphism g in  $\mathscr{C}$  is called an  $\mathfrak{s}$ -deflation or simply a deflation if it admits some conflation  $X_{\bullet} \in \mathbf{C}_{\mathscr{C}}^{n+2}$  satisfying  $d_n^X = g$ .

**Definition 2.7.** [1, Definition 2.27] For a morphism  $f_{\bullet} \in \mathbf{C}^{n+2}_{\mathscr{C}}(X_{\bullet}, Y_{\bullet})$  satisfying  $f_0 = \mathrm{id}_A$  for some  $A = X_0 = Y_0$ , its mapping cone  $M^f_{\bullet} \in \mathbf{C}^{n+2}_{\mathscr{C}}$  is defined to be the complex

$$X_1 \xrightarrow{d_0^{M_f}} X_2 \oplus Y_1 \xrightarrow{d_1^{M_f}} X_3 \oplus Y_2 \xrightarrow{d_2^{M_f}} \cdots \xrightarrow{d_{n-1}^{M_f}} X_{n+1} \oplus Y_n \xrightarrow{d_n^{M_f}} Y_{n+1}$$

where  $d_0^{M_f} = \begin{bmatrix} -d_1^X \\ f_1 \end{bmatrix}$ ,  $d_i^{M_f} = \begin{bmatrix} -d_{i+1}^X & 0 \\ f_{i+1} & d_i^Y \end{bmatrix}$   $(1 \le i \le n-1)$ ,  $d_n^{M_f} = \begin{bmatrix} f_{n+1} & d_n^Y \end{bmatrix}$ .

The mapping cocone is defined dually, for morphisms  $h_{\bullet}$  in  $\mathbf{C}_{\mathscr{C}}^{n+2}$  satisfying  $h_{n+1} = \mathrm{id}$ .

**Definition 2.8.** [1, Definition 2.32] An *pre-n-exangulated category* is a triplet  $(\mathscr{C}, \mathbb{E}, \mathfrak{s})$  of additive category  $\mathscr{C}$ , additive bifunctor  $\mathbb{E} \colon \mathscr{C}^{\mathrm{op}} \times \mathscr{C} \to \mathsf{Ab}$ , and its exact realization  $\mathfrak{s}$ , satisfying the following condition.

(EA1) Let  $A \xrightarrow{f} B \xrightarrow{g} C$  be any sequence of morphisms in  $\mathscr{C}$ . If both f and g are inflations, then so is  $g \circ f$ . Dually, if f and g are deflations, then so is  $g \circ f$ .

If the triplet  $(\mathscr{C}, \mathbb{E}, \mathfrak{s})$  moreover satisfies the following conditions, then it is called an *n*-exangulated category

- (EA2) For  $\rho \in \mathbb{E}(D, A)$  and  $c \in \mathscr{C}(C, D)$ , let  $_A\langle X_{\bullet}, c^*\rho\rangle_C$  and  $_A\langle Y_{\bullet}, \rho\rangle_D$  be distinguished *n*-exangles. Then  $(\mathrm{id}_A, c)$  has a good lift  $f_{\bullet}$ , in the sense that its mapping cone gives a distinguished *n*-exangle  $\langle M^f_{\bullet}, (d^X_0)_*\rho\rangle$ .
- $(EA2^{op})$  Dual of (EA2).

Note that in the case n = 1, a triplet  $(\mathscr{C}, \mathbb{E}, \mathfrak{s})$  is a 1-exangulated category if and only if it is an extriangulated category, see [1, Proposition 4.3].

**Example 2.9.** From [1, Proposition 4.34] and [1, Proposition 4.5], we know that *n*-exact categories and (n + 2)-angulated categories are *n*-exangulated categories. There are some other examples of *n*-exangulated categories which are neither *n*-exact nor (n + 2)-angulated, see [1-3, 6].

# 3 An equivalent characterization of the axiom (EA2)

In this section we introduce the definition of homotopy cartesian squares in pre-n-exangulated categories and provide an equivalent statement of the axiom (EA2).

Definition 3.1. The following commutative diagram

$X_0 \xrightarrow{f_0} X$	$K_1 \xrightarrow{f_1} \cdots$	$\xrightarrow{f_{n-2}} X_r$	$a - 1 \xrightarrow{f_{n-1}} X$	n
$arphi_0$	$\varphi_1 \\ \downarrow \\ \gamma_1 \xrightarrow{g_1} \cdots$		$\varphi_{n-1}$	$\varphi_n$
$Y_0 \xrightarrow{g_0} Y$	$\downarrow \qquad g_1 \\ \longrightarrow \cdots$	$\xrightarrow{g_{n-2}} Y_n$	$ \xrightarrow{g_{n-1}} Y $	'n

in a pre-n-exangulated category  ${\mathscr C}$  is called a  $homotopy\ cartesian\ diagram$  if the following sequence

$$X_{0} \xrightarrow{\begin{pmatrix} -f_{0} \\ \varphi_{0} \end{pmatrix}} X_{1} \oplus Y_{0} \xrightarrow{\begin{pmatrix} -f_{1} & 0 \\ \varphi_{1} & g_{0} \end{pmatrix}} X_{2} \oplus Y_{1} \xrightarrow{\begin{pmatrix} -f_{2} & 0 \\ \varphi_{2} & g_{1} \end{pmatrix}} \cdots$$
$$\cdots \xrightarrow{\begin{pmatrix} -f_{n-1} & 0 \\ \varphi_{n-1} & g_{n-2} \end{pmatrix}} X_{n} \oplus Y_{n-1} \xrightarrow{(\varphi_{n} & g_{n-1})} Y_{n} \xrightarrow{\partial}$$
(2.1)

is a distinguished *n*-example, where  $\partial$  is called a *differential*.

**Remark 3.2.** The distinguished n-example (2.1) in the definition of homotopy cartesian diagram can also be written as follows

$$X_{0} \xrightarrow{\begin{pmatrix} -f_{0} \\ \varphi_{0} \end{pmatrix}} X_{1} \oplus Y_{0} \xrightarrow{\begin{pmatrix} f_{1} & 0 \\ \varphi_{1} & g_{0} \end{pmatrix}} X_{2} \oplus Y_{1} \xrightarrow{\begin{pmatrix} f_{2} & 0 \\ -\varphi_{2} & g_{1} \end{pmatrix}} \cdots$$

$$\xrightarrow{\begin{pmatrix} f_{n-1} & 0 \\ (-1)^{n}\varphi_{n-1} & g_{n-2} \end{pmatrix}} X_{n} \oplus Y_{n-1} \xrightarrow{((-1)^{n+1}\varphi_{n} & g_{n-1})} Y_{n} \xrightarrow{\partial} .$$

$$(2.2)$$

In fact, the two distinguished *n*-examples are isomorphic:

**Lemma 3.3.** [1, Proposition 3.6] Let  $\mathscr{C}$  be an *n*-examplated category,  $_A\langle X_{\bullet}, \delta \rangle_C$  and  $_B\langle Y_{\bullet}, \rho \rangle_D$ 

be distinguished *n*-examples. Suppose that we are given a commutative square

$$\begin{array}{ccc} X_0 \xrightarrow{d_0^X} X_1 \\ a \\ \downarrow & \bigcirc & \downarrow b \\ Y_0 \xrightarrow{d_0^Y} Y_1 \end{array}$$

in  $\mathscr{C}$ . Then the following holds.

(1) There is a morphism  $f_{\bullet} \colon \langle X_{\bullet}, \delta \rangle \to \langle Y_{\bullet}, \rho \rangle$  which satisfies  $f_0 = a$  and  $f_1 = b$ .

(2) If  $X_0 = Y_0 = A$  and  $a = 1_A$  for some  $A \in \mathscr{C}$ , then the above  $f_{\bullet}$  can be taken to give a distinguished *n*-example  $\langle M_{\bullet}^f, (d_0^X)_* \rho \rangle$ .

**Theorem 3.4.** Let  $(\mathcal{C}, \mathbb{E}, \mathfrak{s})$  be a pre-*n*-exangulated category. Then  $\mathcal{C}$  satisfies (EA2) if and only if  $\mathcal{C}$  satisfies (EA2-1):

Let  $X_0 \xrightarrow{f_0} X_1 \xrightarrow{f_1} \cdots \xrightarrow{f_{n-2}} X_{n-1} \xrightarrow{f_{n-1}} X_n \xrightarrow{f_n} X_{n+1} \xrightarrow{\delta}$  and  $X_0 \xrightarrow{g_0} Y_1 \xrightarrow{g_1} \cdots \xrightarrow{g_{n-2}} Y_{n-1} \xrightarrow{g_{n-1}} Y_n \xrightarrow{g_n} Y_{n+1} \xrightarrow{\delta'}$  be distinguished n-exangles, and  $\varphi_1 : X_1 \to Y_1$  be a morphism such that  $\varphi_1 f_0 = g_0$ . Then there exist morphisms  $\varphi_i : X_i \to Y_i$  for  $2 \le i \le n+1$ , which give a morphism of distinguished n-exangles

and moreover, the following

$$X_{1} \xrightarrow{f_{1}} X_{2} \xrightarrow{f_{2}} \cdots \xrightarrow{f_{n-1}} X_{n} \xrightarrow{f_{n}} X_{n+1}$$

$$\downarrow \varphi_{1} \qquad \qquad \downarrow \varphi_{2} \qquad \qquad \qquad \downarrow \varphi_{n} \qquad \qquad \downarrow \varphi_{n+1}$$

$$Y_{1} \xrightarrow{g_{1}} Y_{2} \xrightarrow{g_{2}} \cdots \xrightarrow{g_{n-1}} Y_{n} \xrightarrow{g_{n}} Y_{n+1}$$

is a homotopy cartesian diagram and  $(f_0)_*\delta'$  is the differential.

*Proof.* (EA2) $\implies$  (EA2-1). It follows from Lemma 3.3.

(EA2-1)  $\Longrightarrow$  (EA2). Since  $(id, \varphi_{n+1}) : (\varphi_{n+1})^* \delta' \to \delta'$  is a morphism of  $\mathbb{E}$ -extensions, by (R0), there exist morphisms  $\varphi'_i : X_i \to Y_i$  for  $1 \le i \le n$  such that the following diagram commutative

$$\begin{array}{c} X_0 \xrightarrow{f_0} X_1 \xrightarrow{f_1} X_2 \xrightarrow{f_2} \cdots \longrightarrow X_n \xrightarrow{f_n} X_{n+1} \xrightarrow{(\varphi_{n+1})^* \delta'} \\ \parallel & \downarrow & \downarrow \\ & \downarrow & \varphi'_1 & \downarrow & \varphi'_2 & \downarrow \\ X_0 \xrightarrow{g_0} Y_1 \xrightarrow{g_1} Y_2 \xrightarrow{g_2} \cdots \longrightarrow Y_n \xrightarrow{g_n} Y_{n+1} \xrightarrow{\delta'} \cdot \end{array}$$

By (EA2-1), there exist morphisms  $\varphi_i'': X_i \to Y_i$  for  $2 \le i \le n+1$ , which give a morphism of distinguished *n*-examples

and moreover, the following

$$X_{1} \xrightarrow{f_{1}} X_{2} \xrightarrow{f_{2}} \cdots \xrightarrow{f_{n-1}} X_{n} \xrightarrow{f_{n}} X_{n+1}$$

$$\downarrow \varphi_{1}' \qquad \qquad \downarrow \varphi_{2}'' \qquad \qquad \qquad \downarrow \varphi_{n}'' \qquad \qquad \downarrow \varphi_{n+1}''$$

$$Y_{1} \xrightarrow{g_{1}} Y_{2} \xrightarrow{g_{2}} \cdots \xrightarrow{g_{n-1}} Y_{n} \xrightarrow{g_{n}} Y_{n+1}$$

is a homotopy cartesian diagram and  $(f_0)_*\delta'$  is the differential.

Since 
$$(\varphi_{n+1})^* \delta' = (\varphi_{n+1}'')^* \delta'$$
, that is,  $(\varphi_{n+1} - \varphi_{n+1}'')^* \delta' = 0$ , then by the exactness of  
 $\mathscr{C}(X_{n+1}, Y_n) \xrightarrow{\mathscr{C}(X_{n+1}, g_n)} \mathscr{C}(X_{n+1}, Y_{n+1}) \xrightarrow{\delta_{\sharp}'} \mathbb{E}(X_{n+1}, X_0),$ 

there is a morphism  $h_{n+1}: X_{n+1} \to Y_n$  which gives  $g_n h_{n+1} = \varphi_{n+1} - \varphi_{n+1}''$ .

Since  $g_n(\varphi_n'-\varphi_n''-h_{n+1}f_n) = g_n\varphi_n'-g_n\varphi_n''-g_nh_{n+1}f_n = g_n\varphi_n'-g_n\varphi_n''-\varphi_{n+1}f_n + \varphi_{n+1}''f_n = 0$ , then by the exactness of

$$\mathscr{C}(X_n, Y_{n-1}) \xrightarrow{\mathscr{C}(X_n, g_{n-1})} \mathscr{C}(X_n, Y_n) \xrightarrow{\mathscr{C}(X_n, g_n)} \mathscr{C}(X_n, Y_{n+1}),$$

there is a morphism  $h_n: X_n \to Y_{n-1}$  which gives  $g_{n-1}h_n = \varphi'_n - \varphi''_n - h_{n+1}f_n$ .

Inductively, for  $i = 2, 3, \dots, n-1$ , we obtain  $h_i : X_i \to Y_{i-1}$  such that  $g_{i-1}h_i = \varphi'_i - \varphi''_i - h_{i+1}f_i$ . In particular,  $g_1(\varphi'_1 - \varphi'_1 - h_2f_1) = -g_1h_2f_1 = -\varphi'_2f_1 + \varphi''_2f_1 + h_3f_2f_1 = -g_1\varphi'_1 + g_1\varphi'_1 = 0$ , and hence there exist  $h_1 : X_1 \to Y_0$  which gives  $g_0h_1 = -h_2f_1$ .

Set

$$\varphi_{i} = \begin{cases} \varphi_{i}^{'} = \varphi_{i}^{''} + h_{i+1}f_{i} + g_{i-1}h_{i} & \text{if } i = 2, 3, \cdots, n \\ \varphi_{i}^{'} + h_{i+1}f_{i} & \text{if } i = 1 \end{cases}$$

then  $\varphi_1 f_0 = (\varphi_1^{'} + h_2 f_1) f_0 = \varphi_1^{'} f_0 = g_0, \ \varphi_2 f_1 = (\varphi_2^{''} + h_3 f_2 + g_1 h_2) f_1 = \varphi_2^{''} f_1 + g_1 h_2 f_1 = g_1 \varphi_1^{'} + g_1 h_2 f_1 = g_1 \varphi_1^{'} + g_1 h_2 f_1 = g_1 \varphi_1^{'} + h_2 f_1) = g_1 \varphi_1.$  The following commutative diagram

implies that

$$X_{1} \xrightarrow{\begin{pmatrix} -f_{1} \\ \varphi_{1} \end{pmatrix}} X_{2} \oplus Y_{1} \xrightarrow{\begin{pmatrix} -f_{2} & 0 \\ \varphi_{2} & g_{1} \end{pmatrix}} X_{3} \oplus Y_{2} \xrightarrow{\begin{pmatrix} -f_{3} & 0 \\ \varphi_{3} & g_{2} \end{pmatrix}} \cdots$$
$$\cdots \xrightarrow{\begin{pmatrix} -f_{n-1} & 0 \\ \varphi_{n-1} & g_{n-2} \end{pmatrix}} X_{n} \oplus Y_{n-1} \xrightarrow{\begin{pmatrix} -f_{n} & 0 \\ \varphi_{n} & g_{n-1} \end{pmatrix}} X_{n+1} \oplus Y_{n} \xrightarrow{(\varphi_{n+1} & g_{n})} Y_{n+1} \xrightarrow{(f_{0})_{*}\delta'}$$

is a distinguished *n*-example. Therefore,  $(id_{X_0}, \varphi_{n+1})$  has a good lift  $(id_{X_0}, \varphi_1, \varphi_2, \cdots, \varphi_n, \varphi_{n+1})$ . That is to say (EA2) holds.

Corollary 3.5. In Theorem 3.4, when n = 1, it is just the dual of Theorem 3.3 in [4].

#### 4 Closed subfunctors arising from half exact functors

In this section, we introduce methods of constructing closed subfunctors of an *n*-exangulated category  $(\mathscr{C}, \mathbb{E}, \mathfrak{s})$  from half exact functors. Let's start with the following key lemma.

**Lemma 4.1.** [1, Proposition 3.16] Let  $(\mathscr{C}, \mathbb{E}, \mathfrak{s})$  be an *n*-exangulated category. For any additive subfunctor  $\mathbb{F} \subseteq \mathbb{E}$ , the following statements are equivalent.

- (1)  $(\mathscr{C}, \mathbb{F}, \mathfrak{s}|_{\mathbb{F}})$  is an *n*-exangulated category.
- (2)  $\mathfrak{s}|_{\mathbb{F}}$ -inflations are closed under composition.
- (3)  $\mathfrak{s}|_{\mathbb{F}}$ -deflations are closed under composition.

We call an additive subfunctor  $\mathbb{F} \subseteq \mathbb{E}$  a closed subfunctor if it satisfies the above equivalent conditions. In this case, we call  $(\mathscr{C}, \mathbb{F}, \mathfrak{s}|_{\mathbb{F}})$  a relative theory of  $(\mathscr{C}, \mathbb{E}, \mathfrak{s})$ . For a relative theory  $(\mathscr{C}, \mathbb{F}, \mathfrak{s}|_{\mathbb{F}})$ , we briefly denote it by  $(\mathscr{C}, \mathbb{F})$ .

**Definition 4.2.** Let  $\mathscr{A}$  be an abelian category. An additive functor  $F : \mathscr{C} \to \mathscr{A}$  is called a half exact functor if the sequence

$$FX_0 \xrightarrow{Ff_0} FX_1 \xrightarrow{Ff_1} \cdots \xrightarrow{Ff_{n-2}} FX_{n-1} \xrightarrow{Ff_{n-1}} FX_n \xrightarrow{Ff_n} FX_{n+1}$$

is exact for any distinguished *n*-example  $X_0 \xrightarrow{f_0} X_1 \xrightarrow{f_1} \cdots \xrightarrow{f_{n-2}} X_{n-1} \xrightarrow{f_{n-1}} X_n \xrightarrow{f_n} X_{n+1} \xrightarrow{\delta} X_{n-1}$ . Moreover, if  $Ff_n$  (resp.  $Ff_0$ ) is an epimorphism (resp. monomorphism), we call F a right exact functor (resp. left exact functor).

**Remark 4.3.** If the category  $\mathscr{C}$  are extriangulated, then Definition 4.2 coincides with the definition of half exact functor (homological functor) of extriangulated categories (cf. [5,8]).

**Definition 4.4.** Let  $F : \mathscr{C} \to \mathscr{A}$  be a half exact functor. We define a subset  $\mathbb{E}_R^F(X_{n+1}, X_0)$  of  $\mathbb{E}(X_{n+1}, X_0)$  consisting of  $\delta$  such that for any distinguished *n*-example  $X_0 \xrightarrow{f_0} X_1 \xrightarrow{f_1} \cdots \xrightarrow{f_{n-2}} X_{n-1} \xrightarrow{f_{n-1}} X_n \xrightarrow{f_n} X_{n+1} \xrightarrow{-\delta}$ , we have that  $Ff_n$  is an epimorphism in  $\mathscr{A}$ . Similarly, we define

a subset  $\mathbb{E}_{L}^{F}(X_{n+1}, X_0)$  of  $\mathbb{E}(X_{n+1}, X_0)$  consisting of  $\delta$  such that  $Ff_0$  is a monomorphism in  $\mathscr{A}$ .

Note that the above definition is well-defined, that is, it does not depend on the choice of a distinguished *n*-example of  $\delta$ . Moreover,  $\mathbb{E}_R^F(X_{n+1}, X_0)$  defines the maximum closed subfunctor such that F becomes a right exact functor.

**Theorem 4.5.** Let  $F : \mathscr{C} \to \mathscr{A}$  be a half exact functor. Then the following statements hold.

- (1)  $\mathbb{E}_{R}^{F}$  is a closed subfunctor of  $\mathbb{E}$ , hence  $(\mathscr{C}, \mathbb{E}_{R}^{F})$  is a relative theory of  $(\mathscr{C}, \mathbb{E}, \mathfrak{s})$ .
- (2) F restricts to a right functor  $F : (\mathscr{C}, \mathbb{E}_R^F) \to \mathscr{A}$ .

(3) Let  $(\mathscr{C}, \mathbb{F})$  be a relative theory of  $(\mathscr{C}, \mathbb{E}, \mathfrak{s})$ . If F restricts to a right functor  $F : (\mathscr{C}, \mathbb{F}) \to \mathscr{A}$ , then we have  $\mathbb{F} \subseteq \mathbb{E}_R^F$ .

*Proof.* (1) First of all, we claim that  $\mathbb{E}_R^F$  is a subfunctor of  $\mathbb{E}$ . In fact, for any  $\delta \in \mathbb{E}_R^F(X_{n+1}, X_0)$ and any morphism  $a_0 : X_0 \to Y_0$ , by (EA2<sup>op</sup>), we obtain the following commutative diagram

Applying F to the above diagram, we have the following commutative diagram with exact rows

in  $\mathscr{A}$ . Since  $Ff_n$  is epimorphic, then  $Fg_n$  is also epimorphic. So we have  $a_0\delta \in \mathbb{E}_R^F(X_{n+1}, Y_0)$ .

For any  $\delta \in \mathbb{E}_R^F(X_{n+1}, X_0)$  and any morphism  $a_{n+1} : Y_{n+1} \to X_{n+1}$ , by (EA2), we obtain the following commutative diagram

$$\begin{array}{c} X_0 \xrightarrow{g_0} Y_1 \xrightarrow{g_1} Y_2 \xrightarrow{g_2} \cdots \longrightarrow Y_n \xrightarrow{g_n} Y_{n+1} \xrightarrow{\delta a_{n+1}} \\ \parallel & \downarrow a_1 & \downarrow a_2 & \downarrow a_n & \downarrow a_{n+1} \\ X_0 \xrightarrow{f_0} X_1 \xrightarrow{f_1} X_2 \xrightarrow{f_2} \cdots \longrightarrow X_n \xrightarrow{f_n} X_{n+1} \xrightarrow{\delta} \end{array}$$

such that

$$Y_{1} \xrightarrow{\begin{pmatrix} -g_{1} \\ a_{1} \end{pmatrix}} Y_{2} \oplus X_{1} \xrightarrow{\begin{pmatrix} -g_{2} & 0 \\ a_{2} & f_{1} \end{pmatrix}} Y_{3} \oplus X_{2} \xrightarrow{\begin{pmatrix} -g_{3} & 0 \\ a_{3} & f_{2} \end{pmatrix}} \cdots$$
$$\cdots \xrightarrow{\begin{pmatrix} -g_{n-1} & 0 \\ a_{n-1} & f_{n-2} \end{pmatrix}} Y_{n} \oplus X_{n-1} \xrightarrow{\begin{pmatrix} -g_{n} & 0 \\ a_{n} & f_{n-1} \end{pmatrix}} Y_{n+1} \oplus X_{n} \xrightarrow{(a_{n+1} & f_{n})} X_{n+1} \xrightarrow{g_{0}\delta}$$

is a distinguished *n*-example. Note that  $g_0 \delta \in \mathbb{E}_R^F(X_{n+1}, Y_1)$  by the previous argument, we have the following exact sequence

$$FY_1 \xrightarrow{F\begin{pmatrix} -g_1 \\ a_1 \end{pmatrix}} FY_2 \oplus FX_1 \xrightarrow{F\begin{pmatrix} -g_2 & 0 \\ a_2 & f_1 \end{pmatrix}} FY_3 \oplus FX_2 \xrightarrow{F\begin{pmatrix} -g_3 & 0 \\ a_3 & f_2 \end{pmatrix}} \cdots$$

$$\cdots \xrightarrow{F\begin{pmatrix} -g_{n-1} & 0\\ a_{n-1} & f_{n-2} \end{pmatrix}} FY_n \oplus FX_{n-1} \xrightarrow{F\begin{pmatrix} -g_n & 0\\ a_n & f_{n-1} \end{pmatrix}} FY_{n+1} \oplus FX_n \xrightarrow{F(a_{n+1} & f_n)} FX_{n+1} \longrightarrow 0$$

in  $\mathscr{A}$ . In order to prove that  $Fg_n$  is an epimorphism in  $\mathscr{A}$ , suppose that  $k: FY_{n+1} \to M$ is a morphism in  $\mathscr{A}$  satisfying  $k \circ Fg_n = 0$ . Let  $(k,0) \in \mathscr{C}(FY_{n+1} \oplus FX_n, M)$ . Since  $k \circ Fg_n = 0$ , then  $(k,0) \begin{pmatrix} -Fg_n & 0 \\ Fa_n & Ff_{n-1} \end{pmatrix} = (-kFg_n, 0) = 0$ . Thus there exists unique a morphism  $s: FX_{n+1} \to M$ , such that  $s \circ (Fa_{n+1}, Ff_n) = (k, 0)$ . So  $s \circ Fa_{n+1} = k$  and  $s \circ Ff_n = 0$ . That is to say, we have the following commutative diagram

$$FY_n \oplus FX_{n-1} \xrightarrow{\begin{pmatrix} -Fg_n & 0 \\ Fa_n & Ff_{n-1} \end{pmatrix}} FY_{n+1} \oplus FX_n^{(Fa_{n+1} & Ff_n)} \xrightarrow{} FX_{n+1} \longrightarrow 0.$$

Note that  $Ff_n$  is an epimorphism, so we have s = 0. Then  $k = s \circ Fa_{n+1} = 0$ . This shows that  $Fg_n$  is an epimorphism in  $\mathscr{A}$ , hence we have  $\delta a_{n+1} \in \mathbb{E}_R^F$ . Thus  $\mathbb{E}_R^F$  is a subfunctor of  $\mathbb{E}$ .

Next, we need to show that  $\mathbb{E}_{R}^{F}(X_{n+1}, X_{0})$  is a subgroup of  $\mathbb{E}(X_{n+1}, X_{0})$ . Note that  $0 \in \mathbb{E}_{R}^{F}(X_{n+1}, X_{0})$ , we only need to show  $\delta' - \delta \in \mathbb{E}_{R}^{F}(X_{n+1}, X_{0})$  for any  $\delta', \delta \in \mathbb{E}_{R}^{F}(X_{n+1}, X_{0})$ . For  $(-\mathrm{id}, \mathrm{id}) : X_{0} \oplus X_{0} \to X_{0}$  and  $(\mathrm{id}_{\mathrm{id}}) : X_{n+1} \oplus X_{n+1} \to X_{n+1}$ , we have  $\delta' - \delta = (-\mathrm{id}, \mathrm{id})(\delta \oplus \delta')(\mathrm{id}_{\mathrm{id}})$ , it is enough to show that  $\delta \oplus \delta' \in \mathbb{E}_{R}^{F}(X_{n+1} \oplus X_{n+1}, X_{0} \oplus X_{0})$ , and this follows from Proposition 3.2 in [1]. Thus  $\mathbb{E}_{R}^{F}$  is an additive subfunctor of  $\mathbb{E}$ .

Finally, let  $X_n \xrightarrow{f_n} X_{n+1} \xrightarrow{g_n} Y_{n+1}$  be any sequence of morphisms in  $\mathscr{C}$ . Assume that  $f_n$  and  $g_n$  are  $\mathfrak{s}|_{\mathbb{E}_R^F}$ -deflations. By (EA1), we know that  $g_n \circ f_n$  is an  $\mathfrak{s}$ -deflation. Thus we assume

$$\mathfrak{s}(\delta) = [Z_0 \xrightarrow{h_0} Z_1 \xrightarrow{h_1} Z_2 \xrightarrow{h_2} \cdots \xrightarrow{h_{n-2}} Z_{n-1} \xrightarrow{h_{n-1}} X_n \xrightarrow{g_n f_n} Y_{n+1}].$$

Note that  $Fg_n$  and  $Ff_n$  are epimorphic, then  $F(g_nf_n)$  is an epimorphism, so  $\delta \in \mathbb{E}_R^F$ . This shows that  $\mathfrak{s}|_{\mathbb{E}_R^F}$ -deflations are closed under composition. Thus  $\mathbb{E}_R^F$  is a closed subfunctor of  $\mathbb{E}$ .

(2) and (3) follow immediately from the construction of  $\mathbb{E}_R^F$ .

**Remark 4.6.** Dually, we obtain the statement with respect to  $\mathbb{E}_L^F$  by applying the above proposition to a contravariant half exact functor  $F : \mathscr{C} \to \mathscr{A}^{\mathrm{op}}$ .

**Example 4.7.** Let  $\mathscr{H} \subseteq \mathscr{C}$  be a full subcategory. Define subfunctors  $\mathbb{E}_{\mathscr{H}}$  and  $\mathbb{E}^{\mathscr{H}}$  of  $\mathbb{E}$  by

$$\mathbb{E}_{\mathscr{H}}(X_{n+1}, X_0) = \{ \delta \in \mathbb{E}(X_{n+1}, X_0) | (\delta_{\sharp})_H = 0 \text{ for any } H \in \mathscr{H} \},\$$
$$\mathbb{E}^{\mathscr{H}}(X_{n+1}, X_0) = \{ \delta \in \mathbb{E}(X_{n+1}, X_0) | (\delta^{\sharp})_H = 0 \text{ for any } H \in \mathscr{H} \}.$$

In [1, Proposition 3.17], it is shown that  $\mathbb{E}_{\mathscr{H}}$  and  $\mathbb{E}^{\mathscr{H}}$  are closed subfunctors of  $\mathbb{E}$ . They are special cases of Theorem 4.5. Since the restricted Yoneda functors  $Y_{\mathscr{H}} : \mathscr{C} \to \operatorname{Mod}_{\mathscr{H}}$  and  $Y^{\mathscr{H}} : \mathscr{C} \to \operatorname{Mod}_{\mathscr{H}^{op}}$  are half exact functors, moreover,  $\mathbb{E}_{\mathscr{H}} = \mathbb{E}_{R}^{Y^{\mathscr{H}}}$  and  $\mathbb{E}^{\mathscr{H}} = \mathbb{E}_{L}^{Y^{\mathscr{H}}}$  hold. In fact, for any distinguished *n*-example

$$X_0 \xrightarrow{f_0} X_1 \xrightarrow{f_1} \cdots \xrightarrow{f_{n-2}} X_{n-1} \xrightarrow{f_{n-1}} X_n \xrightarrow{f_n} X_{n+1} \xrightarrow{\delta},$$

we have the following exact sequence

$$\mathscr{C}(-,X_0)|_{\mathscr{H}} \xrightarrow{Y_{\mathscr{H}}(f_0)} \cdots \xrightarrow{Y_{\mathscr{H}}(f_{n-1})} \mathscr{C}(-,X_n)|_{\mathscr{H}} \xrightarrow{Y_{\mathscr{H}}(f_n)} \mathscr{C}(-,X_{n+1})|_{\mathscr{H}} \xrightarrow{\delta_{\sharp}} \mathbb{E}(-,X_0)|_{\mathscr{H}}$$

Thus we have  $\mathbb{E}_{R}^{Y^{\mathscr{H}}}(X_{n+1}, X_0) = \mathbb{E}_{\mathscr{H}}(X_{n+1}, X_0).$ 

**Corollary 4.8.** In Theorem 4.5, when n = 1, it is just Proposition A in [9].

# References

- M. Herschend, Y. Liu, H. Nakaoka. n-exangulated categories (I): Definitions and fundamental properties. J. Algebra 570: 531–586, 2021.
- [2] M. Herschend, Y. Liu, H. Nakaoka. n-exangulated categories (II): Constructions from n-cluster tilting subcategories. J. Algebra, 594, 636–684, 2022.
- [3] J. Hu, D. Zhang, P. Zhou. Two new classes of n-exangulated categories. J. Algebra 568: 1–21, 2021.
- [4] X. Kong, Z. Lin, M. Wang. The (ET4) axiom for extriangulated categories. arXiv: 2112.06445v1, 2021.
- [5] Y. Liu, H. Nakaoka. Hearts of twin cotorsion pairs on extriangulated categories. J. Algebra 528: 96–149, 2019.
- [6] Y. Liu, P. Zhou. Frobenius n-exangulated categories. J. Algebra 559: 161–183, 2020.
- [7] H. Nakaoka, Y. Palu. Extriangulated categories, Hovey twin cotorsion pairs and model structures. Cah. Topol. Géom. Différ. Catég. 60(2): 117–193, 2019.
- [8] Y. Ogawa. Auslander's defects over extriangulated categories: an application for the general heart construction. arXiv: 1911.00259. 2019.
- [9] A. Sakai. Relative extriangulated categories arising from half exact functors. arXiv: 2111.14419v1, 2021.

#### Jian He

Department of Mathematics, Nanjing University, 210093 Nanjing, Jiangsu, P. R. China E-mail: jianhe30@163.com

#### Jing He

College of Science, Hunan University of Technology and Business, 410205 Changsha, Hunan P. R. China E-mail: jinghe1003@163.com

# Panyue Zhou

College of Mathematics, Hunan Institute of Science and Technology, 414006 Yueyang, Hunan, P. R. China. E-mail: panyuezhou@163.com