

# Chemical abundances in Seyfert galaxies– IX. Helium abundance estimates

O. L. Dors<sup>1</sup>★, M. Valerdi<sup>2</sup>, P. Freitas-Lemes<sup>1</sup>, A. C. Krabbe<sup>1</sup>, R. A. Riffel<sup>3</sup>, E. B. Amôres<sup>4</sup>, R. Riffel<sup>5</sup>, M. Armah<sup>5</sup>, A. F. Monteiro<sup>6</sup>, C. B. Oliveira<sup>1</sup>

<sup>1</sup>UNIVAP - Universidade do Vale do Paraíba. Av. Shishima Hifumi, 2911, CEP: 12244-000, São José dos Campos, SP, Brazil

<sup>2</sup>Instituto Nacional de Astrofísica, Óptica y Electrónica (INAOE), Luis E. Erro No. 1, Sta. Ma. Tonantzintla, Puebla, C.P. 72840, México.

<sup>3</sup>Departamento de Física, Centro de Ciências Naturais e Exatas, Universidade Federal de Santa Maria, 97105-900, Santa Maria, RS, Brazil

<sup>4</sup>Departamento de Física, Universidade Estadual de Feira de Santana, Av. Transnordenstina, S/N, CEP 44036-900 Feira de Santana, BA, Brazil

<sup>5</sup>Departamento de Astronomia, Universidade Federal do Rio Grande do Sul, Av. Bento Gonçalves 9500, Porto Alegre, RS, Brazil

<sup>6</sup>Instituto Federal do Maranhão. Av. Newton Bello s/n, CEP: 65906-335, Imperatriz, MA, Brazil

Accepted XXX. Received YYY; in original form ZZZ

## ABSTRACT

For the first time, the helium abundance relative to hydrogen (He/H), which relied on direct measurements of the electron temperature, has been derived in the narrow line regions (NLRs) from a local sample of Seyfert 2 nuclei. In view of this, optical emission line intensities [ $3000 < \lambda(\text{\AA}) < 7000$ ] of 65 local Seyfert 2 nuclei ( $z < 0.2$ ), taken from Sloan Digital Sky Survey Data Release 15 and additional compilation from the literature, were considered. We used photoionization model grid to derive an Ionization Correction Factor (ICF) for the neutral helium. The application of this ICF indicates that the NLRs of Seyfert 2 present a neutral helium fraction of  $\sim 50$  per cent in relation to the total helium abundance. We find that Seyfert 2 nuclei present helium abundance ranging from 0.60 to 2.50 times the solar value, while  $\sim 85$  per cent of the sample present over-solar abundance values. The derived (He/H)-(O/H) abundance relation from the Seyfert 2 is steeper than that of star-forming regions (SFs) and this difference could be due to excess of helium injected into the Interstellar Medium by the winds of Wolf Rayet stars. From a regression to zero metallicity, by using Seyfert 2 estimates combined with SFs estimates, we obtained a primordial helium mass fraction  $Y_p = 0.2441 \pm 0.0037$ , a value in good agreement with the one inferred from the temperature fluctuations of the cosmic microwave background by the Planck Collaboration, i.e.  $Y_p^{\text{Planck}} = 0.2471 \pm 0.0003$ .

**Key words:** galaxies: abundances; galaxies: active; galaxies: evolution; galaxies: formation; galaxies: ISM; galaxies: nuclei

## 1 INTRODUCTION

The helium abundance determination in the gas phase of Active Galactic Nuclei (AGNs) and gaseous nebulae (H II regions, H II galaxies, Planetary Nebulae) is essential to the characterization of the primordial stellar nucleosynthesis after the Big Bang as well as in the study of the Interstellar Medium (ISM) enrichment of galaxies along the Hubble time.

The first helium and heavy element determinations which relied on the direct measurements of the electron temperature ( $T_e$ -method<sup>1</sup>) in AGNs seems to have been carried out by Osterbrock & Miller (1975) for the radio galaxy 3C 405 (Cygnus A). These authors calculated the He abundance using the He<sup>+</sup> and He<sup>2+</sup> ionic abundances and derived a value of  $y \sim 0.10$ , where

$$y = N(\text{He})/N(\text{H}) \quad (1)$$

the ratio of the total number densities of helium to hydrogen. Numerical simulations by Shields (1974), which were carried out to reproduce observational data for the Seyfert galaxy 3C 120, inferred a value of  $y = 0.09$  (see also Koski 1978; Cruz-Gonzalez et al. 1991)

for this object. Based on the results above it seems that no abnormal helium abundance has been found in the local Seyferts in comparison to the solar value,  $y_\odot = 0.10$  (Grevesse et al. 2010), which is generally used as standard reference and scale factor (see Nicholls et al. 2017). However, helium abundance determination in a sample of AGNs is necessary to confirm this result.

On the other hand, quasars seem to exhibit a different and wider range of  $y$  values. For instance, Bahcall & Oke (1971), by using helium and hydrogen emission-lines observed in the optical and ultraviolet and assuming  $y = y^+ + y^{2+}$ , estimated  $y$  in the range 0.003–0.2 for 14 quasars. Baldwin (1975), assuming a similar approach by Bahcall & Oke (1971), derived the  $y$  for a sample of 14 low-redshift quasars ( $z \approx 0.2$ ) and found values from  $\sim 0.1$  to  $\sim 0.3$  (see also, Williams 1971; Jura 1973). The helium estimates by these authors were based on the  $T_e$ -method, i.e. a conventional and reliable method (Pilyugin 2003; Toribio San Cipriano et al. 2017).

Regarding gaseous nebulae, the first He abundance in this class of object was obtained by Aller & Menzel (1945) for 7 Galactic Planetary Nebulae by using optical emission lines measured by Wyse (1942) and, derived the mean abundance value of the helium relative to hydrogen as  $\sim 0.10$ . Thereafter this pioneering work, Mathis (1957) obtained the first helium abundance estimation from H II regions, in this case in the Orion Nebula, deriving values in the range  $\sim 0.09$ – $\sim 0.16$ , depending on the electron temperature assumed in

★ E-mail: olidors@univap.br

<sup>1</sup> For a review of the  $T_e$ -method see Peimbert et al. (2017) and Pérez-Montero (2017).

the calculations. Subsequent studies have been extended to extragalactic objects and the first helium abundance determination was obtained by [Johnson \(1959\)](#) for 30 Doradus (for other pioneering works see: [Aller & Faulkner 1962](#); [Mathis 1962, 1965](#); [Faulkner & Aller 1965](#); [Aller et al. 1968](#); [Peimbert & Spinrad 1970](#)). Thanks to the (relatively) recent spectroscopic surveys, as the Sloan Digital Sky Survey (SDSS, [York et al. 2000](#)), it has become possible to derive the helium abundance for thousands of star-forming regions (SFs; H II regions and star-forming galaxies) with a wide range of metallicity ( $Z$ , e.g. [Izotov et al. 2007, 2013](#); [Kurichin et al. 2021](#)), i.e.  $7.7 \lesssim 12 + \log[(N(O)/N(H))] \lesssim 8.7$  or  $0.10 \lesssim (Z/Z_{\odot}) \lesssim 1.0$ , assuming the solar oxygen value  $\log[(N(O)/N(H))_{\odot}] = -3.31$  ([Allende Prieto et al. 2001](#)), thus allowing the estimation of the primordial helium abundance.

The usual formalism ([Peimbert & Torres-Peimbert 1974, 1976](#)) establishes that the helium mass fraction, defined by

$$Y = \frac{m_{\text{He}}}{m_{\text{gas}}}, \quad (2)$$

has a relation with the metallicity ( $Z$ ), given by

$$Y = \frac{4y(1 - Z)}{1 + 4y}. \quad (3)$$

The  $Z$  can be traced by the  $N(O)/N(H)^2$  abundance in SFs (e.g. [Yates et al. 2012](#); [Kewley et al. 2019](#)) and in AGNs (e.g. [Storchi-Bergmann et al. 1998](#); [Dors 2021](#)) because the oxygen is the most abundant metal in the Universe and emission lines (e.g.  $[O II]\lambda 3726$ ,  $\lambda 3729$ ,  $[O III]\lambda 5007$ ) of its most abundant ions are ubiquitous measured in the rest-frame optical regime (e.g. [Berg et al. 2016](#)). Along this work  $Z$  and  $O/H$  are assumed interchangeably. Regarding the  $Z - (O/H)$  relation, [Peimbert et al. \(2007\)](#) pointed out that the oxygen by mass is in order of  $(55 \pm 10)$  per cent of the  $Z$  value. It is beyond the scope of the present work to discuss the  $Z - (O/H)$  relation. In any case, [Peimbert et al. \(2007\)](#) reported that the error in the  $O/Z$  ratio translates into an error slightly smaller than 0.0001 in the determination of the primordial  $Y$  value. We assumed the fixed relation  $Z = 20(O/H)$  proposed by [Pagel et al. \(1992\)](#) and used recently by [Kumari et al. \(2021\)](#). Thus, Equation 3 is given by

$$Y = \frac{4y[1 - 20(O/H)]}{1 + 4y}. \quad (4)$$

The primordial helium abundance  $Y_p$  is derived by extrapolating the  $Y - (O/H)$  relation to oxygen abundance (or  $Z$ ) equal to zero (for a review on uncertainties in  $Y$  determinations see [Peimbert et al. 2003, 2007](#); [Olive & Skillman 2004](#)). Recently, [Kurichin et al. \(2021\)](#), by using spectroscopic data of 100 SFs taken from the SDSS ([York et al. 2000](#)), derived the primordial helium mass  $Y_p = 0.2462 \pm 0.0022$ , which this result is in consonance with the value inferred from the temperature fluctuations of the cosmic microwave background by the [Planck Collaboration \(2020\)](#), i.e.  $Y_p^{\text{Planck}} = 0.2471 \pm 0.0003$  (see also [Izotov et al. 2014](#); [Aver et al. 2015](#); [Peimbert et al. 2016](#); [Cooke & Fumagalli 2018](#); [Fernández et al. 2019](#); [Valerdi et al. 2019, 2021a](#); [Hsyu et al. 2020](#)).

The extrapolation of the  $Y - (O/H)$  relation to derive  $Y_p$  is very dependent on the extreme oxygen abundance values, i.e. on the lowest and highest values derived for line emitter objects. Along decades, efforts have been made to derive He and O abundances in Extremely metal-poor galaxies (XMPs), in order to obtain  $Y$  values close to  $Y_p$  (e.g. [Kunth & Sargent 1983](#); [Skillman 1991](#); [Skillman et al. 1994](#); [Garnett & Kennicutt 1994](#); [Olive et al. 1997](#); [Izotov & Thuan 1998](#);

[Peimbert et al. 2000](#); [Izotov et al. 1999, 2007, 2009, 2014](#); [Annibali et al. 2019](#); [Valerdi et al. 2021b](#); [Aver et al. 2021](#)). However, despite efforts have been done to achieve direct oxygen estimation in SFs in the high abundance regime [ $12 + \log(O/H) \gtrsim 8.7$  or  $(Z/Z_{\odot}) \gtrsim 1.0$ ] (e.g., see [Kinkel & Rosa 1994](#); [Bresolin et al. 2004](#); [Garnett et al. 2004](#); [Castellanos et al. 2002](#); [Liang et al. 2007](#); [Andrews & Martini 2013](#); [Brown et al. 2016](#); [Berg et al. 2020](#)), helium abundance determinations in these objects are barely found in the literature.

The inclusion of objects with high metallicity, for which direct electron temperatures as well as  $\text{He}^+$  and  $\text{He}^{2+}$  abundances estimates are possible, will result in a significant improvement in the determination of the  $Y - O/H$  relation and, consequently, in the estimation of  $Y_p$ . In this context, AGNs are ideal objects because they have higher gas ionization degree (e.g. [Richardson et al. 2014](#); [Pérez-Díaz et al. 2021](#)) and higher  $Z$  (e.g. [Storchi-Bergmann et al. 1998](#); [Groves et al. 2006](#); [Revalski et al. 2018a, 2021](#); [Dors et al. 2020a](#)) in comparison with H II regions. These physical features make it possible to precisely measure the total helium abundance, since He II lines are stronger in AGN spectra than those in SFs and, they allow to constraint the  $Y$  value in the high metallicity regime.

Large spectroscopic surveys, such as Sloan Digital Sky Survey (SDSS, [York et al. 2000](#)), CALIFA ([Sánchez et al. 2012](#)), MaNGA ([Bundy et al. 2015](#)) and CHAOS ([Berg et al. 2015](#)) have made thousands of emission lines from SFs and AGNs available and these data have revolutionized our understanding of chemical evolution of galaxies (see, for instance, [Gómez et al. 2003](#); [Glazebrook et al. 2003](#); [Brinchmann et al. 2004](#); [Izotov et al. 2006](#); [Asari et al. 2007](#); [Kewley & Ellison 2008](#); [Pilyugin et al. 2012](#); [Sánchez et al. 2017](#); [Ellison et al. 2018](#); [Yates et al. 2020](#); [Berg et al. 2020](#), among others). However, majority of the data from these surveys have been used to derive abundances mainly in SFs, with AGNs still being understudied. In fact, most of the AGN studies have relied on large surveys just to address the oxygen abundance and/or physical properties (e.g., ionization degree, electron temperature, electron density) of these objects (e.g. [Groves et al. 2006](#); [Zhang et al. 2008](#); [Vaona et al. 2012](#); [Ludwig et al. 2012](#); [Zhang et al. 2013](#); [Richardson et al. 2014](#); [Coil et al. 2015](#); [Feltre et al. 2016](#); [Kawasaki et al. 2017](#); [Carvalho et al. 2020](#); [Dors et al. 2020a](#); [Dors 2021](#); [Agostino et al. 2021](#)). The abundances of other elements beyond the oxygen are poorly known in AGNs, in particular, the local Seyfert 2, for which large amount of spectroscopic data are available in the literature.

[Dors et al. \(2017\)](#), who built detailed photoionization models to reproduce optical narrow emission line ratio intensities of a sample of 44 Seyfert 2 nuclei, presented the first quantitative estimations of the nitrogen abundance for this class of object (see also [Pérez-Montero et al. 2019](#); [Ji et al. 2020](#); [Flury & Moran 2020](#); [Dors et al. 2021](#)). Recently, [Dors et al. \(2020b\)](#) proposed a new methodology of the  $T_e$ -method for AGNs which produce reliable  $O/H$  abundances, slightly lower ( $\approx 0.2$  dex) than those derived from detailed photoionization models. [Armah et al. \(2021\)](#) and [Monteiro & Dors \(2021\)](#), motivated by this new methodology and by the availability of spectroscopic data in the literature, developed Ionization Correction Factors (ICFs) as well as electron temperature relations based on photoionization model results and, for the first time, derived the neon and argon abundances in a sample of Seyfert 2 nuclei, respectively.

As a further study, in the present work, we apply the  $T_e$ -method to a combined spectroscopy data taken from the SDSS and additional data from distinct authors with the goal to derive the  $y$  values in a sample of local Seyfert 2 galaxies. These estimates allow access to the helium abundance in a higher metallicity regime than that in SFs and produce important constraints to primordial helium abundance as well as to the studies of stellar nucleosynthesis. The present study is

<sup>2</sup> For simplicity, along the paper, this ratio is defined by  $O/H$ .

organized as follows. In Sections 2 and 3 the observational data and the methodology used to estimate the helium and oxygen abundances are presented, respectively. The results and discussion are presented in Sect. 4. Finally, the conclusion of the outcome is given in Sect. 5.

## 2 OBSERVATIONAL DATA

### 2.1 Seyfert 2 nuclei

In the present work we studied a subsample of the sample presented by Dors et al. (2020a). These authors used the measurements of the SDSS-DR7 made available by MPA/JHU<sup>3</sup> group to obtain a sample of Seyfert 2 nuclei. Initially, by using the MPA/JHU data, Dors et al. (2020a) carried out a cross-correlation between the galaxy identification provided by the SDSS-DR7 and by the NED/IPAC<sup>4</sup> and obtained the classification of each object in Seyfert 1 and 2 nuclei. After applying the criteria to separate SF and AGN objects proposed by Kewley et al. (2001, 2006) and Pérez-Montero et al. (2013), which consider a set of emission line intensity ratios in diagnostic diagrams (Baldwin et al. 1981), 463 Seyfert 2 AGNs ( $z \lesssim 0.4$ ) with reliable optical emission lines in the optical range  $3000 \lesssim \lambda(\text{\AA}) \lesssim 7500$  were selected.

The data from the MPA/JHU group do not contain measurements for the He II  $\lambda 4686\text{\AA}$ . Therefore, we downloaded from SDSS-DR15 database<sup>5</sup> the spectra of the 463 objects selected by Dors et al. (2020a) in order to obtain a sub-sample of Seyfert 2 nuclei in which it is possible to determine the parameters necessary to calculate the He and O abundances through the  $T_e$ -method. Firstly, in each spectrum, we performed the extinction correction using the Cardelli et al. (1989) law assuming the parameterized extinction coefficient  $R_V = 3.1$ , a standard value for the diffuse interstellar medium. The Galactic interstellar extinction values provided by Schlegel et al. (1998) and calibrated by Schlafly & Finkbeiner (2011) obtained by using the VOTool for interstellar extinction called GALEX<sup>6</sup> (Amôres et al. 2021) were also assumed. We shifted the spectra for the rest frame wavelength using the redshifts available for each object and binned them with bin size equal to  $1\text{\AA}$ .

The stellar population continuum was subtracted from the spectra to obtain the pure nebular spectra using the stellar population synthesis STARLIGHT code (Cid Fernandes et al. 2005; Mateus et al. 2006; Asari et al. 2007). This code fits the observed spectrum of a galaxy using a combination of Simple Stellar Populations (SSPs), in different proportions and excluding the emission lines. We used a basis of 45 synthetic SSP spectra with three metallicities  $Z = 0.004$ ,  $0.02$  ( $Z_\odot$ ), and  $0.05$ , assuming 15 ages ranging from 1 Myr to 13 Gyr, taken from the evolutionary synthesis models of Bruzual & Charlot (2003). Prior to the fitting the synthetic spectra to the observational ones, we convoluted the SSP stellar spectra adopting a Gaussian function to achieve the same spectral resolution of the observational data. A detailed description of the SSP spectra fitting to spectroscopic observational data is given by Krabbe et al. (2011, 2017).

After the pure nebular spectra for the sample were obtained by subtracting the contribution of the stellar component from the observed spectra, the emission-lines were fitted using the publicly available IFSCUBE package (Ruschel-Dutra & de Oliveira 2020; Ruschel-Dutra

et al. 2021). Each emission line was fitted by a single Gaussian function and the kinematics (line width and centroid velocity) of lines from the same parent ion was kept tied. We also included in the fit a third order polynomial to account for any residual continuum emission. The uncertainties on the fluxes were obtained by performing 100 iterations of Monte Carlo simulations of the emission-line fits as the standard deviation of the fluxes obtained from these simulations. The fluxes were corrected for extinction following the procedure described by Riffel et al. (2021b), using the  $(H\alpha/H\beta)$  line ratio to estimate the visual extinction, assuming the theoretical value for the  $(H\alpha/H\beta) = 2.86$  proposed by Hummer & Storey (1987) at a temperature of  $10\,000\text{ K}$  and an electron density of  $100\text{ cm}^{-3}$ . A detailed analysis justifying the assumption for considering  $(H\alpha/H\beta) = 2.86$  for reddening correction in AGNs was presented by Armah et al. (2021). We found values of the extinction coefficient  $A_V$  ranging from 0.02 to 0.4 mag.

It is clear hitherto that the most widely used method for estimating the dust content is based on the relative strengths of the lower Balmer H I lines i.e.  $H\alpha/H\beta$ . When using the gas emission lines as a dust tracer, the canonical assumption is that the gas emission comes from the same position as the emission from the ionization source, which is usually not the case, because the gas in a spiral galaxy for instance is contained in a plane. Generally, little is known about the nature and real distribution of dust in AGNs (e.g. Heard & Gaskell 2016; Baron et al. 2016; Gaskell 2017, and references therein). However, neglecting these uncertainties at first order and considering the fact that  $H\alpha$  and  $H\beta$  are the strongest recombination lines of hydrogen in the optical spectrum, the reddening effect can be written on the reliance of the ratio  $H\alpha/H\beta$ . Assigning a single extinction value is just a rough first approximation, therefore, irrespective of the dereddening method used, it is good practice to verify that the  $H\alpha/H\beta$ ,  $H\gamma/H\beta$  and  $H\delta/H\beta$  emission line ratios have the expected values in comparison with the theoretical Case B ratios of 2.86 or 3.1, 0.468 and 0.259 (Halpern 1982; Halpern & Steiner 1983; Hummer & Storey 1987; Osterbrock & Ferland 2006), respectively. On the other hand, H7 and H8 can not be considered in reddening correction because they are usually blended with other emission lines. Additionally, the signal-to-noise ratio for lines such as H9 and H10 is often too low to allow for their detection with high precision in order to constrain the reddening and underlying stellar absorption. Hence, as part of the emission-line flux processing, the underlying continuum and absorption-line spectra were fitted and eliminated, removing the effects of stellar Balmer absorption on the  $H\alpha$  and  $H\beta$  (e.g. Moustakas & Kennicutt 2006). Finally, the amount of interstellar extinction obtained using  $H\alpha/H\beta$  is relatively in agreement with the amount of extinction determined considering other line ratios, i.e.  $[S\text{ II}]\lambda 4071/[S\text{ II}]\lambda 10320$  (Wampler 1971; Shields & Oke 1975; Koski 1978) or  $\text{He II}\lambda 3203/\text{He II}\lambda 4686$  (Shuder & Osterbrock 1981).

From the resulting sample, we selected only the objects that present the emission-lines  $[\text{O II}]\lambda 3726, \lambda 3729$  (hereafter  $[\text{O II}]\lambda 3727$ ),  $[\text{O III}]\lambda 4363$ ,  $\text{He II}\lambda 4686$ ,  $H\beta$ ,  $[\text{O III}]\lambda 5007$ ,  $\text{He I}\lambda 5876$ ,  $H\alpha$ , and  $[S\text{ II}]\lambda 6716, \lambda 6731$  with a signal/noise ratio (S/N) higher than 2.0. The final sample resulted in 9 objects with redshift  $z \lesssim 0.2$ . In Figure 1, an example of a pure Seyfert 2 nebular spectrum (in black) and the fitting (in red) produced by the IFSCUBE software (Ruschel-Dutra & de Oliveira 2020) are shown.


In addition to the SDSS data, we compiled from the literature fluxes of emission-lines from Seyfert 2 nuclei obtained by different authors. We applied the same selection criteria used for the SDSS data to these additional selected objects. This sample consists of 94 Seyfert 2 nuclei whose emission-line intensities were reddening corrected by the authors from which the data were taken. However, in the cases

<sup>3</sup> <https://wwwmpa.mpa-garching.mpg.de/SDSS/DR7/>

<sup>4</sup> [ned.ipac.caltech.edu](http://ned.ipac.caltech.edu)

<sup>5</sup> <https://dr15.sdss.org/optical/spectrum/search>

<sup>6</sup> <http://www.galexextin.org>



**Figure 1.** Optical spectrum of one of the Seyfert 2 nucleus in our sample (see Sect. 2.1) taken from SDSS DR15 and represented in black colour. The fitting to the emission-line profiles using the IFSCUBE code (Ruschel-Dutra & de Oliveira 2020) is represented in red colour. The measured emission lines and corresponding wavelength are indicated. Boxes show a zoom in regions of some weak lines, as indicated.

where the reddening correction was not performed in the original works, the same procedure applied to the SDSS data was considered. Since several measurements for the emission lines compiled from the literature do not have their uncertainties listed in the original papers where the data were compiled, we adopted a typical error of 10 per cent for strong emission-lines (e.g. [O III] $\lambda$ 5007) and 20 per cent for weak emission lines (i.e. [O III] $\lambda$ 4363, He II $\lambda$ 4686 and He I $\lambda$ 5876), as derived, for instance, by Kraemer et al. (1994).

Finally, we applied the criterion proposed by Kewley et al. (2001),

$$\log([\text{O III}]\lambda 5007/\text{H}\beta) > \frac{0.61}{\log([\text{N II}]\lambda 6584/\text{H}\alpha) - 0.47} + 1.19 \quad (5)$$

to separate SF and AGN objects. Additionally, the criterion proposed by Cid Fernandes et al. (2010) to separate AGN-like and Low-ionization nuclear emission-line region (LINER) objects, given


by

$$\log([\text{O III}]\lambda 5007/\text{H}\beta) > 0.47 + \log([\text{N II}]\lambda 6584/\text{H}\alpha) \times 1.10, \quad (6)$$

was also considered.

The final sample consists of 65 Seyfert 2 with redshift  $z \lesssim 0.2$  whose reddening-correction line intensities (in relation to  $\text{H}\beta = 1.0$ ) are listed in Table 1 in the Appendix. In Fig. 2, a diagnostic diagram  $\log([\text{O III}]\lambda 5007/\text{H}\beta)$  versus  $\log([\text{N II}]\lambda 6584/\text{H}\alpha)$ , the observational data for the objects as well as the curve representing the criteria above are shown. It can be seen that the objects of our sample cover a large range of ionization degree and metallicity, hence a wide range of [O III]/H $\beta$  and [N II]/H $\alpha$  line ratio intensities are noted (e.g. Groves et al. 2006; Feltre et al. 2016; Carvalho et al. 2020).

Dors et al. (2015) and Castro et al. (2017) presented a complete discussion on the use of heterogeneous sample and its possible implications on abundance estimates. Effects of aperture, electron density variation along the AGN radius, X-Ray dominated regions, shock,




**Figure 2.** Diagnostic diagram  $\log([O \text{ III}]\lambda 5007/H\beta)$  versus  $\log([N \text{ II}]\lambda 6584/H\alpha)$

). Blue points represent Seyfert 2 nuclei of our sample (see Sect. 2.1) while black points are for SFs (see Sect. 2.2). Solid and dashed curves represent the criteria proposed by Kewley et al. (2001) and Kauffmann et al. (2003), given by Equations 5 and 7, respectively, to separate SF and AGN objects. The red line represents the criterion proposed by Cid Fernandes et al. (2010) to separate AGN-like and Low-ionization nuclear emission-line region (LINER) objects, given by Eq. 6.

and electron temperature fluctuations in abundance determinations have also been presented by Dors et al. (2020a, 2021) and these are not repeated here. Basically, the effects of these parameters on abundance estimates produce uncertainties of  $\sim 0.1$  dex, i.e. in order of those derived by applying the  $T_e$ -method (e.g. Izotov & Thuan 2008) and strong-line methods (e.g. Storch-Bergmann et al. 1998). Since it is not possible to estimate, for each AGN of our sample, the aforementioned uncertainty is not considered in the resulting abundance values.

We present a comparison between the logarithm of the  $[O \text{ II}]\lambda 3727/H\beta$  intensity line ratio by adopting our extinction correction procedure with that assumed by the authors from which the data were compiled. This comparison was possible only for 46 objects (19 AGNs and 27 SFs) due to the fact that most of the original works present only the reddening corrected intensity lines. In Fig. 3, bottom panel, this comparison is shown where there exist a good agreement between them. In top panel of Fig. 3 the difference ( $D=y-x$ ) is presented, where one can see the uncertainty due the distinct approaches to extinction correction is in order of that produced by the error measurements ( $\sim 0.1$  dex, e.g. Kennicutt et al. 2003), with the mean difference ( $\langle D \rangle = 0.04$  dex) being about null. Thus, distinct methodologies employed for the extinction correction of intensity line ratios introduce a minimal uncertainty in our abundance estimates.



**Figure 3.** Bottom panel: Comparison between the  $[O \text{ II}]\lambda 3727/H\beta$  intensity line ratio, for part of the observational sample (AGNs and SFs), whose reddening correction were carried out following the methodology described in Sect. 2 and performed by the authors from which the data were compiled. A one-to-one correlation is plotted as a solid line. Top panel: Difference ( $D=y-x$ ) between the intensities. Solid line represent the null difference while dashed lines are the uncertainty of 0.1 dex in the line ratio measurements. The mean difference ( $\langle D \rangle$ ) in the measurements is indicated.


## 2.2 Star forming regions

We compiled emission-line intensities of SFs from the literature with the goal of comparing their estimates with our AGN results. We applied the same criteria above in the selection of SFs, i.e. we selected only objects whose emission-lines  $[O \text{ II}]\lambda 3727$ ,  $[O \text{ III}]\lambda 4363$ ,  $\text{He II}\lambda 4686$ ,  $H\beta$ ,  $[O \text{ III}]\lambda 5007$ ,  $\text{He I}\lambda 5876$ ,  $H\alpha$ , and  $[S \text{ II}]\lambda 6716$ ,  $\lambda 6731$  were measured. The measurements of these lines make it possible to calculate the He and O abundances by using the  $T_e$ -method following a similar methodology applied to the Seyfert 2 sample. Furthermore, we considered only the objects which satisfy the empirical criterion proposed by Kauffmann et al. (2003)

$$\log([O \text{ III}]\lambda 5007/H\beta) < \frac{0.61}{\log([N \text{ II}]\lambda 6584/H\alpha) - 0.05} + 1.3. \quad (7)$$

In Table 2, the emission line intensities (in relation to  $H\beta=1$ ) and the original works from which data were taken are listed. The data consist of reddening corrected emission lines of 85 H II regions and star-forming galaxies with redshift  $z < 0.2$ . In Fig. 2, the SF sample is represented by black points, where we can see the very known sequence (e.g. Baldwin et al. 1981; Kauffmann et al. 2003) formed by this object class.

In Fig. 4, we plotted the logarithm of  $[O \text{ III}]\lambda 5007/[O \text{ II}]\lambda 3727$  versus  $\text{He II}\lambda 4686/\text{He I}\lambda 5876$  line ratios for the AGN sample (blue points) and for the SF sample (black points). Despite the scattering produced mainly for the SF data, it can be seen the expected result that; a clear correlation is derived (see also, for instance, Deharveng et al. 2000) hence both line ratios are dependent on the ionization degree of the gas. Moreover, a distinction between Seyfert and SFs



**Figure 4.** Logarithm of  $[\text{O III}]\lambda 5007/[\text{O II}]\lambda 3727$  versus  $\text{He II}\lambda 4686/\text{He I}\lambda 5876$ . Blue and black points represent Seyfert 2 nuclei (see Sect. 2.1) and SFs (see Sect. 2.2) of our samples, respectively, as indicated. Red line indicates the separation criterion between Seyfert nuclei and SFs given by Eq. 8.

can be observed in Fig. 4 due to the high excitation of Seyfert in comparison with SFs. In Fig. 4, we observe a clear separation criterion, where objects with

$$\log\left(\frac{\text{He II}\lambda 4686}{\text{He I}\lambda 5876}\right) \gtrsim -0.4 \quad (8)$$

are classified as Seyfert in this diagram, otherwise, as SF (see also Nakajima & Maiolino 2022).

### 3 ABUNDANCE DETERMINATION

We calculated helium and oxygen abundances by using the observational data sample of Seyfert 2 nuclei and SFs described in Sect. 2. Therefore, the  $[\text{O III}](1.33 \times \lambda 5007)/\lambda 4363$  and  $[\text{S II}]\lambda 6716/\lambda 6731$  line ratios were used to derive electron temperature and density, respectively, thus obtaining the ionic abundances of these elements. Thereafter, we used photoionization models built with the CLOUDY code (Ferland et al. 2013) to produce a correction for the presence of  $\text{He}^0$  in the calculation of the total He. In what follows, a description of the methodology employed is presented for both object classes.

#### 3.1 Seyfert

##### 3.1.1 Temperature and density

To determine the abundance of an element it is necessary to estimate a representative value for the electron temperature and density of the gas phase occupied by the ions of this element.

Since the elements of interest are the He and O, which are in most part once and twice ionized, following Valerdi et al. (2021a), we adopted the two zone models, where the temperature for the high

ionization zone, defined as  $T_{\text{high}}$  is derived from the relation between this parameter and the  $[\text{O III}](1.33 \times \lambda 5007)/\lambda 4363$  line ratio, where we assumed, as usual,  $[\text{O III}](\lambda 4959)/(\lambda 5007)$  to be 0.33<sup>7</sup> (Rosa 1985). To calculate  $T_{\text{high}}$ , and the electron density ( $N_e$ ) from the  $[\text{S II}]\lambda 6716/\lambda 6731$  line ratio, we used the version 1.1.13 of PyNEB code (Luridiana et al. 2015), which permits an interactive procedure in the derivation of these parameters. We assumed the atomic data of recombination and collisionally excited lines listed in Table 1 for the temperature, density and ionic abundance calculations.

For almost all the data in our sample it was not possible to obtain the direct determination of  $T_{\text{low}}$  due to the lack of the  $[\text{N II}]\lambda 5755$  and  $[\text{O II}]\lambda 7319, \lambda 7330$  auroral line measurements. Thus, for consistence, we used the theoretical relation between  $T_{\text{low}}-T_{\text{high}}$  proposed by Dors et al. (2020b), obtained through a grid of photoionization models simulating Narrow Line Regions (NLRs) of AGNs built with the CLOUDY code (Ferland et al. 2013) by Carvalho et al. (2020), given by

$$t_{\text{low}} = (a \times t_{\text{high}}^3) + (b \times t_{\text{high}}^2) + (c \times t_{\text{high}}) + d, \quad (9)$$

where  $a = 0.17$ ,  $b = -1.07$ ,  $c = 2.07$  and  $d = -0.33$ , while  $t_{\text{low}}$  and  $t_{\text{high}}$  represent  $T_{\text{low}}$  and  $T_{\text{high}}$ , respectively, in units of  $10^4$  K. These temperatures predicted by the models correspond to the mean temperature for  $\text{O}^+$  ( $T_{\text{low}}$ ) and  $\text{O}^{2+}$  ( $T_{\text{high}}$ ) over the nebular AGN radius times the electron density.

In Riffel et al. (2021a), the relation represented by the Eq. 9 was compared with electron temperature values derived from observational auroral emission lines for a sample of 12 local AGNs, and a good agreement was found between them. However, as pointed out by these authors, some cautions must be taken into account in the use of Eq. 9 for AGNs with strong outflowing gas.

##### 3.1.2 Helium abundance

The total helium abundance in relation to hydrogen  $y = N(\text{He})/N(\text{H})$  was considered to be

$$y = y^0 + y^+ + y^{2+}, \quad (10)$$

where  $y^0 = N(\text{He}^0)/N(\text{H}^+ + \text{H}^0)$ ,  $y^+ = N(\text{He}^+)/N(\text{H}^+)$  and  $y^{2+} = N(\text{He}^{2+})/N(\text{H}^+)$ . The ionic abundance ratio  $y^+$  and  $y^{2+}$  were derived from the  $\text{He I}\lambda 5876/\text{H}\beta$  and  $\text{He II}\lambda 4696/\text{H}\beta$  intensity line ratios, respectively. We used the two-zone ionization model (e.g. Valerdi et al. 2019), which  $T_{\text{low}}$  (from Eq. 9) and  $T_{\text{high}}$  (from direct estimation) were assumed in the  $y^+$  and  $y^{2+}$  calculation, respectively. Recently, Berg et al. (2021) proposed a more detailed ionization model for H II regions, i.e. the 4-zones model, where the  $\text{He}^+$  is located in the low, intermediate, and high zones, i.e. in the zones occupied for the  $\text{O}^+$  and  $\text{O}^{2+}$  ions. Thus, in principle,  $T_{\text{low}}$  can be assumed as a representative temperature value for the region occupied by the  $\text{He}^+$ . It is worth noting that this is not a conventional approach, therefore some previous studies assume  $T_{\text{high}}$  in the  $\text{He}^+$  abundance calculations. Concerning  $\text{He}^{2+}$ , Berg et al. (2021) showed that it is located mainly in the very-high-ionization zone. To derive the temperature of the very-high-ionization zone the measurement of the  $[\text{Ne III}](\lambda 3342/\lambda 3868)$  ratio is required, which is not available in our data sample. Therefore, we assumed  $T_{\text{high}}$  in the abundance calculation of this ion. Both ionic abundances were computed using the PyNEB code (Luridiana et al. 2015). We consider the presence

<sup>7</sup> For a detailed discussion on the value of this ratio see Storey & Zeippen (2000).

**Table 1.** Atomic dataset used for recombination and collisionally excited lines of selected element ions.

| Ion              | Transition probabilities and energy levels               | Collisional strengths    | Recombination coefficients |
|------------------|----------------------------------------------------------|--------------------------|----------------------------|
| He <sup>+</sup>  | Porter et al. (2012)                                     | —                        | Porter et al. (2012)       |
| He <sup>2+</sup> | Storey & Hummer (1995)                                   | —                        | Storey & Hummer (1995)     |
| O <sup>+</sup>   | Wiese et al. (1996)                                      | Kisieliuss et al. (2009) | —                          |
| O <sup>2+</sup>  | Froese Fischer & Tachiev (2004), Storey & Zeippen (2000) | Storey et al. (2014)     | —                          |

of He<sup>0</sup> in the neutral and ionized gas since this is expected in low ionization objects (Peña 1986; Dinerstein & Shields 1986).

The fraction of y<sup>0</sup> in H II regions and Planetary Nebulae has been estimated to be in order of ~ 3 per cent of the total helium (e.g. Peña 1986; Peimbert et al. 1992) or even larger values could be predicted (e.g. Deharveng et al. 2000). In fact, Méndez-Delgado et al. (2022) found that only 9/42 Galactic H II regions have negligible contributions of y<sup>0</sup> (see also Delgado-Inglada et al. 2014). Since AGNs present a large amount of molecular gas (e.g., Rodríguez-Ardila et al. 2004; Davies et al. 2005; Rodríguez-Ardila et al. 2005; Mazzalay et al. 2013; Riffel et al. 2013, 2018, 2021c; Ramakrishnan et al. 2019; Riffel et al. 2019; Jarvis et al. 2020; Alonso-Herrero et al. 2020) and have similar ionization parameter (*U*) other than SFs (Pérez-Montero et al. 2019), a higher abundance of neutral and molecular gas are expected in AGNs as compared to H II regions, which favors the presence of larger contribution of y<sup>0</sup> in the total helium abundance. In fact, radio observations have shown the existence of a gas neutral reservoir in the central parts of galaxies containing AGNs (e.g. Dressel et al. 1982; Hutchings et al. 1987; Bertram et al. 2007; Ho et al. 2008; Bradford et al. 2018; Ellison et al. 2019; Combes et al. 2019) which can coexist with the ionized gas. García-Burillo et al. (2014), who used the Atacama Large Millimeter Array (ALMA) to map the emission of dense molecular gas in the Seyfert 2 NGC 1068, showed an overlay of the CO(3–2) (a tracer of H<sub>2</sub>) intensity contours on the Pa $\alpha$  emission obtained with the Hubble Space Telescope. In addition, recently, do Nascimento et al. (2022) derived the O/H abundance in NLR of 108 Seyfert nuclei and found that these values are lower (by 0.16 to 0.30 dex) than those inferred by the radial gradients along their galaxy discs and those from a matched control sample of no active galaxies. This discrepancy could be due to the accretion of a metal-poor gas to the AGN that feeds the nuclear supermassive black hole (SMBH), coming from the neutral reservoir.

To estimate the abundance of y<sup>0</sup>, we used a grid of photoionization models similar to that of Carvalho et al. (2020), built with the CLOUDY code version 17.00 (Ferland et al. 2013). These models were also used to derive neon and argon ICFs by Armah et al. (2021) and Monteiro & Dors (2021), respectively, considering a wide range of nebular parameters:

- Metallicity: ( $Z/Z_{\odot}$ ) = 3.0, 2.0, 1.0, 0.75, 0.5, and 0.2.
- Electron density:  $N_e$  (cm<sup>-3</sup>) = 3000, 1500, 500, 100.
- Ionization parameter (*U*): log *U* ranging from -1.5 to -3.5, with step of 0.5 dex.
- Spectra Electron Distribution (SED): The SED is parametrized by the continuum between 2 keV and 2500 Å (Tananbaum et al. 1979) and it is described by a power law with a spectral index  $\alpha_{ox}$  = -0.8, -1.1 and -1.4.

The outermost nebular radius was considered by Carvalho et al. (2020) to be the one where the electron temperature reaches 4000 K (the default value of the CLOUDY code), which produces a region with

hydrogen almost completely ionized. Since we are also interested in the neutral region, we rerun the grid of models by Carvalho et al. (2020) but using the version 17.02 of the CLOUDY code (Ferland et al. 2013) and assuming the outermost radius where the electron temperature reaches 1000 K; as was done by Dors et al. (2012), who analysed the dominant excitation mechanism of [Fe II] and H<sub>2</sub> emission lines in AGNs. These new models allow us to consider a larger part of the neutral gas instead of stopping the calculations at outermost radius of the modelled nebula where the temperature falls below 4000 K. We added cosmic-ray background emission as a second ionizing source with a value of H<sub>2</sub> ionization rate of 10<sup>-15</sup> s<sup>-1</sup>, which is about the same rate found by McCall et al. (2003) for a Galactic line of sight. The cosmic-ray background emission has little influence on the formation of emission lines located in the ionized gas and/or in the ionization of elements with ionization potential higher than the hydrogen one. In the models the abundance of each ion was considered as the average over the nebular radius times the electron density. For a detailed description of the photoionization models see Carvalho et al. (2020).


Based on the model results, we assumed the ICF for the y<sup>0</sup> to be

$$\text{ICF}(y^0) = \frac{y}{y^+ + y^{2+}}. \quad (11)$$

In Fig. 5, the photoionization model predictions for ICF(y<sup>0</sup>) versus  $x = y^+ / y^{2+}$  are shown, where the results are discriminated according to distinct nebular parameters, as indicated. Any dependence between the ICF(y<sup>0</sup>)-*x* relation and  $N_e$  and *Z* can be seen. However, a clear dependence of this relation with the ionization parameter *U* and  $\alpha_{ox}$  is noted, in the sense that a steeper ICF(y<sup>0</sup>)-*x* relation is derived for higher *U* and  $\alpha_{ox}$  values. The dependence of this relation on these parameters is due to the increase in the gas ionization degree, i.e. driven by *U* and the hardness of the SED ( $\alpha_{ox}$ ), produces lower values of *x*.

Estimations of  $\alpha_{ox}$  require observations of integrated nuclear flux at optical and UV wavelengths (e.g. Worrall et al. 1987; Ho 1999; Miller et al. 2011; Zhu et al. 2019, 2021), and values of this parameter are often times unavailable for AGNs. Moreover, detailed photoionization models by Dors et al. (2017) and Bayesian-like approach by Pérez-Montero et al. (2019) have predicted  $\alpha_{ox}$  to be higher than -1.2 for Seyfert 2 nuclei. Conversely, observational estimations of  $\alpha_{ox}$  by Miller et al. (2011) indicate that most AGNs have  $\alpha_{ox}$  in the order of -1.4 and even lower values (~ -2.0) are derived for these objects. This discrepancy is probably due to an extra physical process is missing in the models, probably shocks (e.g. Contini 2019). Therefore, as a result of few AGNs estimations of  $\alpha_{ox}$  in comparison to those with emission-line measurements, and also due to the existence of the discrepancy between model predictions and observational estimations of  $\alpha_{ox}$ , this parameter was not directly taken into account in the expression for the helium ICF derived here. In any case, the gas ionization degree, traced by *U*, probably has a direct relation with  $\alpha_{ox}$ .

We derive a bi-parametric calibration  $\text{ICF}(y^0) = f(x, \log U)$



**Figure 5.** Photoionization model predictions (see Sect 3.1.2) for  $ICF(y^0)$  versus  $y^+/y^{2+}$ . In each plot, model results assuming distinct nebular parameters are shown by different colours, as indicated.  $ICF(y^0)$  is defined as in Eq. 11.

shown in Fig. 6 and given by

$$z = [(0.14 \pm 0.01) \times \ln(x) \times w] + [(0.09 \pm 0.01) \times w^2] + [(0.03 \pm 0.01) \times x] + [(0.18 \pm 0.07) \times w] + [(1.71 \pm 0.08)], \quad (12)$$

where  $z = ICF(y^0)$ ,  $x = y^+/y^{2+}$  and  $w = \log U$ . The ionization parameter can be obtained using the semi-empirical calibration proposed by [Carvalho et al. \(2020\)](#)

$$\log U = (0.57 \pm 0.01 r^2) + (1.38 \pm 0.01 r) - (3.14 \pm 0.01), \quad (13)$$


where  $r = \log([O III] \lambda 5007 / [O II] \lambda 3727)$ . This calibration was obtained through a comparison of observational optical narrow emission line ratios of a sample of local Seyfert 2 nuclei with those predicted by photoionization models. An expression for deriving the helium ICF taking into account the ionization parameter but for H II regions was also derived by [Sauer & Jedamzik \(2002\)](#), by using a grid of photoionization models.

### 3.1.3 Oxygen abundance

The total oxygen abundance in relation to hydrogen was derived assuming

$$\frac{N(O)}{N(H)} = ICF(O) \times \left[ \frac{N(O^{2+})}{N(H^+)} + \frac{N(O^+)}{N(H^+)} \right], \quad (14)$$

where  $ICF(O)$  is the Ionization Correction Factor for the oxygen which takes into account the contribution of unobserved oxygen ions (e.g.  $O^{3+}$ ). In fact, several studies have found a strong intensity of the  $[O IV] \lambda 25.89 \mu m$  emission line in AGN spectra (e.g. [Diamond-Stanic & Rieke 2012](#); [Fernández-Ontiveros et al. 2016](#)), indicating a non-negligible fraction of the  $O^{3+}$  abundance.



**Figure 6.** Bi-parametric calibration among the  $ICF(y^0)$ ,  $x = y^+/y^{2+}$  and logarithm of the ionization parameter ( $\log U$ ). Points represent the photoionization model results (see Sect 3.1.2),  $ICF(y^0)$  is defined by Eq. 11. The surface is given by Eq. 12.

To calculate the  $ICF(O)$  the empirical expression proposed by [Torres-Peimbert & Peimbert \(1977\)](#):

$$ICF(O) = \frac{y^+ + y^{2+}}{y^+}, \quad (15)$$

used in SF (e.g. [Izotov et al. 2006](#); [Valerdi et al. 2021a](#)) and AGN (e.g. [Flury & Moran 2020](#); [Dors et al. 2020b](#)) abundances studies, was considered. To calculate the  $ICF(O)$  for each object of the sample, we consider our helium abundance results derived using the `PyNEB` code ([Luridiana et al. 2015](#)).

In Table 3 presented in the Appendix, we show the oxygen (ionic and total) and helium abundances. To estimate the abundance uncertainties, due to few objects present observational error in the line measurements, for all objects of the AGN sample, we use a Monte Carlo simulations. For each diagnostic line, we generate 1000 random values assuming a Gaussian distribution with a standard deviation equal to the associated uncertainty of the line intensity involved in the diagnostic.

### 3.2 Star-forming regions

To determine the He and O abundances for the SFs, we adopted a similar procedure considered for the Seyfert sample. The ionic helium abundances were also computed using the `PyNEB` code ([Luridiana et al. 2015](#)) which,  $y^{2+}$  and  $O^{2+}$  were derived assuming the  $T_{high}$  and  $N_e$ , calculated for each object through the  $[O III] (1.33 \times \lambda 5007) / \lambda 4363$  and  $[S II] \lambda 6716 / \lambda 6731$  line ratios, respectively.

The  $y^+$  and  $O^+$  were calculated assuming the  $T_{low}$  derived from the theoretical relation

$$t_{low}^{-1} = 0.693 t_{high}^{-1} + 0.281 \quad (16)$$

proposed by [Pérez-Montero & Díaz \(2003\)](#). We assume  $T_{low} \equiv T_e(N II) \approx T_e(O II)$ .

The total oxygen abundance is obtained as described previously:

$$\frac{O}{H} = ICF(O) \times \frac{O^+}{H^+} + \frac{O^{2+}}{H^+}, \quad (17)$$

where  $\text{ICF}(\text{O})$  is same as Eq. 15.


In the same way, we assume the expression

$$y = \text{ICF}(y^0) \times (y^+ + y^{2+}), \quad (18)$$

in the total helium abundance calculation. The  $\text{ICF}(y^0)$  expression (Eq. 11) proposed for Seyferts can not be applied to SFs due to the distinct ionization structure of these objects. Thus, we build a grid of photoionization models simulating SFs by using the CLOUDY code. The nebular parameters are similar to those adopted by Dors et al. (2018) and they are summarised in what follows.

- Metallicity:  $(Z/Z_\odot) = 0.03, 0.2, 0.5$  and  $1.0$ . This metallicity range includes values derived from the disk H II regions (e.g. Berg et al. 2020) and in XPMs (e.g. Izotov et al. 1999).
- Electron density:  $N_e \text{ (cm}^{-3}\text{)} = 100, 500$  and  $1000$ , the same range derived by Zhang et al. (2013) for SFs and calculated from  $[\text{S II}]\lambda 6716/\lambda 6731$  line ratio.
- Ionization parameter ( $U$ ):  $\log U$  ranging from  $-1.5$  to  $-3.5$ , with step of  $0.5$  dex. Similar range of  $\log U$  was derived by Dors et al. (2011) for local H II regions (see also Ji & Yan 2022; Zinchenko et al. 2019; Pérez-Montero 2014).
- SED: synthetic spectra of stellar clusters formed by an instantaneous burst and with ages equal to  $0.01, 1.0, 2.5, 4.0$  and  $5.0$  Myr, built with the STARBURST99 (Leitherer et al. 1999) were considered as ionizing source. The value  $0.01$  Myr corresponds to the lowest age considered in the STARBURST99. In the interval  $2.5 - 4.0$  Myr OB stars become Wolf-Rayet stars producing an increase in the number of ionizing photons (e.g. Leitherer & Heckman 1995; Dopita et al. 2006; Vale Asari et al. 2016). Nebulae older than about  $5$  Myr are difficult to observe because their original massive stars have cooled emitting little ionizing photons (e.g., Garcia-Vargas et al. 1995; Mayya & Prabhu 1996). We assumed the WM-basic stellar atmosphere models by Pauldrach et al. (2001), and the GENEVA tracks with stellar rotation (Levesque et al. 2012).

For these models the calculations stopped at outermost nebular radius where the temperature falls below  $4000$  K. We note that the photoionization model results predict a very low  $\text{He}^{2+}$  abundance in comparison to those derived from our observational data sample. This indicates that real SEDs of the ionizing source of SFs are harder than the theoretical ones assumed here. The Binary Population and Spectral Synthesis code (BPASS, Stanway & Eldridge 2018) produces somewhat harder SEDs than the STARBURST99 (e.g. D’Agostino et al. 2019), however, the former might be more appropriate for moderate/low metallicity SFs and less appropriate for high metallicity systems (Nakajima & Maiolino 2022). Due to the problem pointed out above, we derive a relation between the  $\text{ICF}(\text{He}^0)$  and the  $\text{O}^+/\text{O}$  abundance ratio instead of  $y^+/y^{2+}$  as previously. In Fig. 7, the model results for the relation  $\text{ICF}(\text{He}^0)$ -( $\text{O}^+/\text{O}$ ) are shown, where models assuming stellar clusters with distinct ages are indicated. It can be seen that, for  $(\text{O}^+/\text{O}) \lesssim 0.6$ , higher ICFs are obtained for older models ( $4$  and  $5$  Myr). We did not find any dependence of the  $\text{ICF}(\text{He}^0)$ -( $\text{O}^+/\text{O}$ ) relation with other nebular parameters. Dottori (1981) and Copetti et al. (1986) proposed that the equivalent width of H $\beta$ , the relative volume of  $\text{He}^+$  and  $\text{H}^+$  zones and the  $[\text{O III}](\lambda 4959, \lambda 5007)/\text{H}\beta$  line ratio are good age H II region indicators. However, the change of the value of these indicators along the spiral disk galaxies can be due to effects of variation of the hottest effective temperature of ionizing stars (Shields & Tinsley 1976) rather than evolution effects. Since our SF sample is mostly composed of disk H II regions, the age estimation, in principle, is not correct and it was not carried out.



**Figure 7.** Calibration between the  $\text{ICF}(y^0)$  and  $(\text{O}^+/\text{O})$ . Points represent SF photoionization model results (see Sect 3.2) assuming different ages for the ionizing source, as indicated.  $\text{ICF}(y^0)$  and the curve are defined by Eq. 19.

Thus, a fit taken into account all the points in Fig. 7 results in

$$\text{ICF}(y^0) = a \times (\text{O}^+/\text{O})^b + c, \quad (19)$$

where  $a = 0.385 \pm 0.017$ ,  $b = 4.742 \pm 0.275$  and  $c = 1.020 \pm 0.002$ .


The abundance uncertainties for this sample were derived taken into account the error in the observational emission-line intensities published by the authors which the data were compiled.

## 4 RESULTS AND DISCUSSION

### 4.1 Density and temperature

The electron density values derived from the  $[\text{S II}]\lambda 6716/\lambda 6731$  line ratio for the Seyfert 2 sample are in the range of  $20 \lesssim N_e(\text{cm}^{-3}) \lesssim 1200$ , with a median value of  $\sim 500 \text{ cm}^{-3}$ , which is in agreement with the estimation obtained by Dors et al. (2014). However, the derived mean value from our AGN sample is a factor of about two lower than the one derived by Vaona et al. (2012), who used optical spectra of  $\sim 2300$  AGNs obtained with the SDSS-DR7 (York et al. 2000). Thus, the discrepancy between our mean electron density result and the one derived by Vaona et al. (2012) is probably due to the distinct sample of objects. In any case, the  $N_e$  values are lower than the critical density for the emission lines involved in this work (see, for instance, Vaona et al. 2012), therefore, de-excitation effect has no influence on our abundance calculations (e.g. Osterbrock & Ferland 2006).

The values derived for the SF sample are in the range of  $20 \lesssim N_e(\text{cm}^{-3}) \lesssim 600$ , with a mean value of  $\sim 100 \text{ cm}^{-3}$ . This reflects the known discrepancy where NLRs of AGNs present higher electron density than SFs (e.g. Zhang et al. 2013). In Fig. 8, panel (a), the distribution of values for the  $N_e$  from both Seyfert and SF samples are shown. From this figure it is clear that AGNs have  $N_e \lesssim 1200 \text{ cm}^{-3}$  while most SF ( $\sim 90$  percent) have  $N_e < 200 \text{ cm}^{-3}$ .



**Figure 8.** Panel (a): Electron density distributions, calculated from  $[\text{S II}]\lambda 6716/\lambda 6731$  line ratio (see Sect. 3.1.1) by using the PYNEB (Luridiana et al. 2015), for distinct samples of objects (see Sect. 2), as indicated. Panel (b): As left panel but for  $T_{\text{high}}$  calculated through the observational values of the  $[\text{O III}](1.33 \times \lambda 5007)/\lambda 4363$  line ratio.

Regarding the electron temperature results, we derive values for  $T_{\text{high}}$  from the Seyferts ranging from  $\sim 9000$  K to  $\sim 25000$  K while values for SFs are in the range of  $\sim 7300$  K– $20000$  K. In panel (b) of Fig. 8, the distributions of these values for our Seyfert and SF samples are shown. The Seyfert mean value is  $\sim 15000$  K, similar to the ones derived by Revalski et al. (2018a,b) for the quasar type 2 Mrk 34 and for the Seyfert 2 Mrk 573, i.e. about  $13000$  K, respectively. Heckman & Balick (1979) suggested that AGNs with electron temperature values higher than  $20000$  K, probably, present a secondary source of energy in addition to photoionization, possibly the presence of shocks (see also Dors et al. 2021 and references therein). It can be seen that majority ( $\sim 92$  per cent) of the AGNs present  $T_{\text{high}}$  values lower than  $20000$  K. Therefore, the objects of our sample, possibly, are mainly (photo)ionized by the radiation from the AGN accretion disk (e.g. shocks can be neglected). For SFs, a somewhat lower mean value of  $\sim 11500$  K was obtained.


Peimbert (1967) proposed the presence of temperature fluctuations to explain the considerable differences found between the temperature estimations relied on distinct methods in H II regions. Recently, Riffel et al. (2021d), for the first time, by using Gemini GMOS-IFU observations of three luminous nearby Seyfert galaxies (Mrk 79, Mrk 348 and Mrk 607), quantified temperature variations in this class of object. These authors derived a lower limit for the temperature fluctuation parameter ( $t^2$ ) to be from  $\sim 0.04$  to  $\sim 0.1$ , in order of those derived for H II regions and Planetary Nebulae. Although temperature fluctuations have little effect on helium abundance estimates (e.g. Peimbert et al. 2012), the supposition of  $t^2 > 0$  leads to the derivation of higher O/H abundances and, consequently, lower ( $\sim 3$  per cent) values of  $Y_p$  (Peimbert et al. 2002). Since estimations of temperature fluctuation in AGNs are barely found in the literature, we assume the temperature values based on [O III] line ratio as fiducial values for the high ionization zone of Seyfert nuclei. However, we emphasise that temperature fluctuation can be present.

In Fig. 9, we assumed the range of observational  $\text{He II}\lambda 5876/\text{H}\beta$

and  $\text{He II}\lambda 4686/\text{H}\beta$  line ratio values for our sample of objects (listed in Table 1) and calculated the  $y^+$  and  $y^{2+}$  abundances considering temperatures from  $5000$  K to  $25000$  K and a fixed value of  $N_e = 100 \text{ cm}^{-3}$ . It can be seen that the effect of varying the temperature is in order of the uncertainty produced by the observational error for the line measurements. Therefore, we emphasize that, if temperature fluctuations exist in the objects of our sample, shocks or electron temperature fluctuations affect only the oxygen abundance estimations, while the helium estimations are reliable values.

## 4.2 Ionization Correction Factor

Delgado-Inglada et al. (2014) computed a large grid of photoionization models that covers a wide range of physical parameters of planetary nebulae (PNe). They reported that the derivation of an ICF for neutral helium based on other ions is not recommended because the relative populations of helium ions depend essentially on the effective temperature ( $T_{\text{eff}}$ ) of the central star ( $T_{\text{eff}}$  drives the flux of the ionizing photons), whereas those from metal ions also depend on the ionization parameter. These authors also pointed out that the correction for neutral helium is important only in models with ionizing sources at  $T_{\text{eff}} \lesssim 50000$  K. In the bi-parametric expression for the Seyfert ICF( $\text{He}^0$ ) proposed here (Eq. 12), both hardness of the ionizing radiation and the gas ionization degree are taken into account by the  $y^+/y^{2+}$  ratio and by the ionization parameter  $U$ . For SFs, we notice a dependence between ICF( $\text{He}^0$ ) and the age of the ionization source, while  $T_{\text{eff}}$  is similar to the age, since the temperature of the hottest stars of the ionizing stellar cluster of an H II region drives the flux of the ionizing photons (Zastrow et al. 2013) and it decreases with the time. However, the stellar evolution could have a minimal effect on our helium neutral fraction estimation in SFs because we considered the presence of  $\text{He II}\lambda 4876$  line as a selection criterion of the sample, which led to the selection of H II regions with young cluster ages. In fact, Bresolin et al. (1999) showed that only



**Figure 9.** Lower panel: Abundances of  $y^+$  versus the He II 5876/H $\beta$  line ratio for distinct electron temperature values (in units of  $10^4$  K), as indicated, assuming a fixed value of  $N_e = 100 \text{ cm}^{-3}$ . Top panel: Same as lower panel but for  $y^{2+}$  versus He II 4686/H $\beta$ . In both plots, the range of emission-line ratio values (x-axes) represents the range of observational values of our sample of objects listed in Table 1. The error bars represent the mean error in the emission-line measurements of the sample of objects.

photoionization models with clusters younger than 3 Myr are able to reproduce optical emission line intensities of H II regions.

The application of the ICF(He<sup>0</sup>), i.e. Eq. 12, to our AGN data yields ICFs values ranging from 1.3 to 1.7, with a mean value of  $\sim 1.5$ . This result indicates that in NLRs of Seyfert 2 the helium is  $\sim 50$  per cent in a neutral stage. As reported previously, radio and near-infrared (e.g. Rodríguez-Ardila et al. 2004; Storchi-Bergmann et al. 2009; Riffel et al. 2010, 2021c; Mezcuca et al. 2015; Schönell et al. 2017; Bianchin et al. 2022, among others) observations have shown that AGNs harbor a neutral and warm (1000-3000 K) molecular gas reservoir. Therefore, a substantial proportion of neutral helium is expected in Seyfert nuclei.

On the other hand, the application of Eq. 19 indicates that SFs have ICF(He<sup>0</sup>) ranging from 1.0 to 1.2, implying 70 per cent of the objects have ICF(He<sup>0</sup>)=1.0. The mean ICF(He<sup>0</sup>) value for SFs is 1.03, i.e. the helium is only  $\sim 3$  per cent in a neutral stage. Pagel et al. (1992), following the methodology from Mathis (1982), proposed the use of the abundance ratio

$$\eta = (\text{O}^+/\text{S}^+)/(\text{S}^{2+}/\text{O}^{2+}), \quad (20)$$

known by radiation softness parameter (Vilchez & Pagel 1988), as a means of estimating the significance of neutral helium in SFs. Pagel et al. (1992) pointed out that, for H II regions with  $\log \eta < 0.9$ , the fraction of neutral helium into Strömgren sphere is negligible. Recently, by using observational data from the MaNGA survey (Bundy et al. 2015), Kumari et al. (2021) determined through the  $T_e$ -method values of the  $\eta$  parameter for 67 star-forming galaxies. These authors found values in the range of  $-0.4 \lesssim \log \eta \lesssim 0.6$ . Bresolin et al. (2009) obtained spectrophotometric data for 28 H II regions in the spiral galaxy NGC 300 and, also by using the  $T_e$ -method, found that the majority of the objects present  $\log \eta$  around 0.7. Otherwise, most part of Galactic H II region observed by Méndez-Delgado et al. (2022) present  $\log \eta > 0.9$ , i.e. a non-negligible neutral helium

abundance. Our results are in consonance with those derived by Méndez-Delgado et al. (2022), who posited that, like Seyfert nuclei, SFs can have a significant neutral helium fraction.

One of the major sources of the uncertainty in abundance estimates is the use of ICFs, producing discrepancies by a factor of up to  $\sim 4$  according to the suppositions considered to derived them (e.g. Dors et al. 2013; Méndez-Delgado et al. 2020). Unfortunately, AGN ICFs for the helium are not found in the literature, therefore, it is not possible to estimate the ICF error in our estimates. In any case, detailed models built with the goal of reproducing line intensities of a large sample of AGNs (e.g. Dors et al. 2017) and SFs are needed to validate the neutral ICFs proposed here.

### 4.3 Helium and oxygen abundances


Initially, we compare the  $y$  and O/H range of values derived from our AGN sample with those derived from SFs. This is significant since the helium abundance at any ionization stage has hitherto been unknown for a large sample of AGNs.

In Fig. 10, panel (a), the distribution of  $12+\log(y)$  for the Seyfert 2 and SF samples are shown. Also in this plot, a line indicating the solar value of the helium abundance, i.e.  $12+\log(y)_\odot = 11.0$  derived by Grevesse et al. (2010) is indicated. We can see that most Seyfert 2 present higher helium abundance in comparison with those of SFs, with the Seyfert range being  $10.80 \lesssim [12 + \log(y)] \lesssim 11.40$  or  $0.60 \lesssim (y/y_\odot) \lesssim 2.50$ , and a mean value of  $\langle 12+\log(y) \rangle = 11.11 \pm 0.11$  or  $\langle (y/y_\odot) \rangle \sim 1.30$ . We can also note that the majority ( $\sim 84$  per cent) of AGNs present oversolar helium abundance. Concerning the SFs, we derived the range  $10.70 \lesssim [12 + \log(y)] \lesssim 11.10$  or  $0.50 \lesssim (y/y_\odot) \lesssim 1.20$ , with a mean value  $\langle 12+\log(y) \rangle = 10.92 \pm 0.05$  or  $\langle (y/y_\odot) \rangle \sim 0.80$ .

In Fig. 10, panel (b), the distribution of  $12+\log(\text{O}/\text{H})$  for the Seyfert 2 and SFs samples are shown. For the Seyfert sample, we derived the range  $8.20 \lesssim [12 + \log(\text{O}/\text{H})] \lesssim 9.10$ , or  $0.30 \lesssim (Z/Z_\odot) \lesssim 2.6$ , with most part ( $\sim 63$  per cent) having oversolar abundance. The mean value is  $\langle 12 + \log(\text{O}/\text{H}) \rangle = 8.74 \pm 0.16$  or  $\langle (Z/Z_\odot) \rangle \sim 1.10$ . This is a known result, i.e. subsolar metallicities are derived in few AGNs in the local universe (e.g., Groves et al. 2006), independently of the method used to estimate this parameter (Dors et al. 2020a). For SFs, the range of  $7.20 \lesssim [12 + \log(\text{O}/\text{H})] \lesssim 8.80$  or  $0.03 \lesssim (Z/Z_\odot) \lesssim 1.30$  is derived, being  $\sim 8$  per cent of the objects presenting oversolar oxygen abundance.

The lower helium and oxygen abundances in SFs in comparison with those in AGNs is due to the fact that the latter are located in the central parts of galaxies and, according to the inside-out scenario of galaxy formation (see Mollá & Díaz 2005 and references therein) have experimented a longer time-scale of chemical enrichment of the ISM in comparison with disk H II regions (most part of our SF sample). It is worth to mention that, our direct helium estimates, based on the  $y^+$  and  $y^{2+}$  ionic abundances of AGNs, seem to be the unique for the very high metallicity regime, i.e.  $(Z/Z_\odot) > 1.0$ . This fact is very important in the derivation of the  $y$ -O/H relation and of  $Y_p$  because our estimates, combined those for XMPs, produce a wide range of metallicities likely not previously considered.

In Fig. 11, the total helium abundance [in units of  $12+\log(y)$ ] versus the oxygen abundance [in units of  $12+\log(\text{O}/\text{H})$ ] from our Seyfert (blue points) and SF (black points) samples are shown. Other than in AGNs, we note that the SFs with the highest metallicities present near  $y$  values. Similar results were derived for nitrogen and argon by Carvalho et al. (2020) and Monteiro & Dors (2021), respectively. Moreover, we notice two clear behaviours, a smooth  $y$ -(O/H)



**Figure 10.** As Fig. 8 but for  $12+\log(y)$  [panel (a)] and  $12+\log(\text{O}/\text{H})$  [panel (b)]. The blue lines in panels (a) and (b) represent the solar values derived by [Allende Prieto et al. \(2001\)](#) and [Grevesse et al. \(2010\)](#), respectively. These values correspond to  $12+\log(y)_{\odot} = 11.0$  and  $12+\log(\text{O}/\text{H})_{\odot} = 8.69$ .

relation for the low metallicity regime [ $12 + \log(\text{O}/\text{H}) \lesssim 8.5$  or  $(Z/Z_{\odot}) \lesssim 0.6$ ] and a stepper relation for high metallicity regime. This difference could be due to an excess of helium injected into the ISM by winds of Wolf Rayet stars, which are more common in the high metallicity environment (e.g. [Maeder & Meynet 1994](#); [Leitherer et al. 1999](#); [Pindao et al. 2002](#); [Crowther et al. 2004](#); [Bresolin et al. 2005](#)), containing hydrogen burning products ([Pagel et al. 1992](#); [Crowther & Dessart 1998](#)). The scattering of the points observed in both kind of objects in Fig. 11 is, probably, due to effects of different star formation rate in them (e.g. [Berg et al. 2020](#); [Hayden-Pawson et al. 2022](#)) and/or due to radial migration of stars ([Vincenzo et al. 2021](#); [Johnson et al. 2022](#)) rather than evolutionary processes, by virtue of the fact that majority of the AGNs and SFs of our sample are located in the local universe.

In order to derive an expression for the  $y$ -O/H relation, we performed 1000 bootstrap realisations ([Davison & Hinkley 1997](#)) with Huber Regressor model ([Owen 2007](#)). The resulting fitting to the points obtained is given by

$$w = (0.1215 \pm 0.0422) \times x^2 - (1.8183 \pm 0.6977) \times x + (17.6732 \pm 2.8798) \quad (21)$$


where  $w=12+\log(y)$  and  $x=12+\log(\text{O}/\text{H})$ . [Dopita et al. \(2006\)](#), by using abundance estimates of SFs obtained by [Pagel et al. \(1992\)](#) and [Russell & Dopita \(1992\)](#), derived the relation

$$y = 0.0737 + 0.024 \times (Z/Z_{\odot}). \quad (22)$$

We converted the  $y$  and  $Z/Z_{\odot}$  values of this expression into the same units assumed previously as well as extrapolated it for the high metallicity regime. In Fig. 11, we compare the [Dopita et al. \(2006\)](#) relation with our relation (Eq. 21), which it can be seen a good agreement between both.


#### 4.4 Primordial helium abundance

[Searle & Sargent \(1972\)](#) presented abundance analysis of the two



**Figure 11.** Abundances of  $12+\log(y)$  versus  $12+\log(\text{O}/\text{H})$ . Blue and black points represent the values for our sample of Seyfert 2 and SFs, respectively, listed in Table 3. The red curve represents the fit to the points given by Eq. 21. The green curve represents Eq. 22 derived by [Dopita et al. \(2006\)](#), but with  $y$  and  $Z/Z_{\odot}$  values converted into  $12+\log(y)$  and  $12+\log(\text{O}/\text{H})$ , respectively.

compact dwarf galaxies IZw18 and II Zw40, and for the first time, suggested that metal-poor objects, such as these, would be used to estimate the primordial helium abundance. After this pioneering work, several studies have been undertaken in an effort to address



**Figure 12.** Left panel:  $Y$  versus the abundance ratio  $N(O)/N(H)$  derived by using the  $T_e$ -method.  $Y$  values are calculated as in Eq. 4. Blue and black points represent estimates for our Seyfert 2 and SF samples, respectively. Red line represents the linear regression to the points given by Eq. 23. The primordial helium abundance derived from this linear regression,  $Y_p = 0.2441 \pm 0.0037$ , is indicated. Right panel: Estimates of the primordial helium abundance ( $Y_p$ ) derived by different authors. The red line represents the estimation by Planck Collaboration (2020), i.e.  $Y_p = 0.2471 \pm 0.0003$ . Other estimates correspond to: (1) This work ( $Y_p = 0.2441 \pm 0.0037$ ), (2) Izotov et al. (2014), (3) Aver et al. (2015), (4) Peimbert et al. (2016), (5) Cooke & Fumagalli (2018), (6) Fernández et al. (2019), (7) Valerdi et al. (2019), (8) Hsyu et al. (2020), (9) Kurichin et al. (2021), (10) and (11) estimates by Méndez-Delgado et al. (2022) for  $t^2 = 0$  and  $t^2 > 0$ , respectively. The  $Y_p$  value obtained by each author is listed by Kurichin et al. (2021), with exception of the one derived by Méndez-Delgado et al. (2022).

the observation of XMPs and objects in a wide range of metallicities in order to derive  $Y_p$  (for a review see Peimbert 2008; Ferland et al. 2010; Skillman et al. 2012; Peimbert et al. 2017).

Peimbert et al. (2007) listed the main error sources in the  $Y_p$  determination, which produce uncertainties from  $\pm 0.0005$  to  $\pm 0.0015$  in the final value of  $Y_p$ , where the collisional excitation of the H I lines and temperature structure were considered as the most important physical conditions in the gaseous nebulae. Also, Méndez-Delgado et al. (2020) pointed out that an important source of error in the  $Y_p$  estimates is related to the ionization structure of the nebulae, i.e. the use of ICFs in the estimation of the total helium abundance, since spectra of typical SF only show recombination lines of  $He^+$ .

In our case, probably, the main source of uncertainty in the  $y$  estimation and, consequently of  $Y_p$ , is the fact that observational data of our AGN sample, in opposite to the SF sample, do not have errors in the line measurements for most of the objects and typical values for these were considered (see Sect. 2). This procedure, combined with the uncertainty produced by the use of ICF( $He^0$ ), certainly produced larger errors than those discussed by Peimbert et al. (2007) and Méndez-Delgado et al. (2020). Despite this drawback, our AGN estimates combined with estimations from SFs are useful in the derivation of  $Y_p$  taking objects with very high metallicity into account.

We calculated  $Y$  values using our Seyfert 2 and SFs results, listed in Table 3 together with Eq. 4 for the purpose of estimating  $Y_p$  which rely on a wide range of metallicities. These values are shown in panel (a) of Fig. 12. It can be seen that Seyfert 2s show higher  $Y$  and O/H values in comparison with those in SFs. As previously, we performed 1000 bootstrap realisations to obtain a linear fit that

yielded the following relation

$$Y = (94.0946 \pm 9.9534) \times N(O)/N(H) + (0.2441 \pm 0.0037) \quad (23)$$

with a Pearson correlation coefficient of  $R = 0.57$ . Thus, we derived the value of  $Y_p = 0.2441 \pm 0.0037$ .

In Fig. 12, panel (b), estimates of  $Y_p$  obtained by different authors and methods are compared to our estimate. Also in this figure the line representing the value derived by the Planck Collaboration (2020) is shown. It is worthwhile to note that our  $Y_p$  value is in a good agreement than the others, with exception of the estimation by Cooke & Fumagalli (2018) which presents the highest error bars. We performed a linear fit to the points considering only our SF estimates and found the expression:

$$Y = (36.5264 \pm 7.1944) \times N(O)/N(H) + (0.2459 \pm 0.0024), \quad (24)$$

with  $Y_p = 0.2459 \pm 0.0024$ , i.e. a value somewhat higher than the one obtained by assuming AGN and SF estimates. However, this value is in consonance with the one inferred by the Planck Collaboration (2020) and by other authors.

## 5 CONCLUSION

We used optical emission line intensities [ $3000 < \lambda(\text{\AA}) < 7000$ ] taken from SDSS DR15 and additional compilations from the literature to derive, through  $T_e$ -method, the helium and oxygen abundances relative to hydrogen in the narrow line regions (NLRs) of a local ( $z < 0.2$ ) sample of 65 Seyfert 2 nuclei and 85 star-forming regions (SFs; i.e. H II region, star-forming galaxies). Photoionization model grids, built with the CLOUDY code and simulating NLRs and

SFs, were used to obtain expressions for the Ionization Correction Factor (ICF) of the neutral helium. The application of these ICFs indicates that the NLRs of Seyfert 2 present a neutral helium fraction in relation to the total helium abundance ranging from  $\sim 20$  to  $\sim 70$  per cent (mean value 50 per cent) and SFs from  $\sim 0$  to  $\sim 20$  per cent (mean value 3 per cent). The high neutral helium abundance in AGNs it is due to these objects harbour a neutral and warm (1000–3000 K) molecular gas reservoir. We found that Seyfert 2 nuclei present helium abundance ranging from 0.60 to 2.50 times the solar value, which implies that for  $\sim 85$  per cent of the sample an oversolar helium abundance was derived. Our results indicate that NLR of Seyfert 2 nuclei have a steeper (He/H)-(O/H) relation than SFs. This difference could be due to excess of helium injected into the ISM by winds of Wolf Rayet stars, which are more common in the high metallicity environment, i.e.  $12 + \log(\text{O}/\text{H}) \gtrsim 8.7$  [ $(Z/Z_{\odot}) \gtrsim 1.0$ ]. From a regression to zero metallicity, by using Seyfert 2 and SFs estimates, we derived a primordial helium mass fraction  $Y_p = 0.2441 \pm 0.0037$ , a value in good agreement with the one inferred from the temperature fluctuations of the cosmic microwave background by the Planck Collaboration, i.e.  $Y_p^{\text{Planck}} = 0.2471 \pm 0.0003$ .

## DATA AVAILABILITY

The data underlying this article will be shared on reasonable request to the corresponding author.

## ACKNOWLEDGEMENTS

We are grateful to the referee for his/her dedicated work in reviewing our paper. OLD and ACK are grateful to the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) and to Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for the financial support. CBO is grateful to the FAPESP for the support under grant 2019/11934-0 and to the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES). Funding for the Sloan Digital Sky Survey has been provided by the Alfred P. Sloan Foundation, the U.S. Department of Energy Office of Science, and the Participating Institutions. SDSS acknowledges support and resources from the Center for High-Performance Computing at the University of Utah. The SDSS web site is [www.sdss.org](http://www.sdss.org).

## REFERENCES

- Agostino C. J., et al., 2021, arXiv e-prints, [p. arXiv:2108.07812](https://arxiv.org/abs/2108.07812)
- Allende Prieto C., Lambert D. L., Asplund M., 2001, *ApJ*, **556**, L63
- Aller L. H., Faulkner D. J., 1962, *PASP*, **74**, 219
- Aller L. H., Menzel D. H., 1945, *ApJ*, **102**, 239
- Aller L. H., Czyzak S. J., Walker M. F., 1968, *ApJ*, **151**, 491
- Alloin D., Bica E., Bonatto C., Prugniel P., 1992, *A&A*, **266**, 117
- Alonso-Herrero A., et al., 2020, *A&A*, **639**, A43
- Amôres E. B., et al., 2021, *MNRAS*, **508**, 1788
- Andrews B. H., Martini P., 2013, *ApJ*, **765**, 140
- Annibali F., et al., 2019, *MNRAS*, **482**, 3892
- Armah M., et al., 2021, *MNRAS*.
- Asari N. V., Cid Fernandes R., Stasińska G., Torres-Papaqui J. P., Mateus A., Sodré L., Schoenell W., Gomes J. M., 2007, *MNRAS*, **381**, 263
- Aver E., Olive K. A., Skillman E. D., 2015, *J. Cosmology Astropart. Phys.*, **2015**, 011
- Aver E., Berg D. A., Hirschauer A. S., Olive K. A., Pogge R. W., Rogers N. S. J., Salzer J. J., Skillman E. D., 2021, arXiv e-prints, [p. arXiv:2109.00178](https://arxiv.org/abs/2109.00178)
- Bahcall J. N., Oke J. B., 1971, *ApJ*, **163**, 235
- Baldwin J. A., 1975, *ApJ*, **201**, 26
- Baldwin J. A., Phillips M. M., Terlevich R., 1981, *PASP*, **93**, 5
- Baron D., Stern J., Poznanski D., Netzer H., 2016, *ApJ*, **832**, 8
- Berg D. A., Skillman E. D., Croxall K. V., Pogge R. W., Moustakas J., Johnson-Groh M., 2015, *ApJ*, **806**, 16
- Berg D. A., Skillman E. D., Henry R. B. C., Erb D. K., Carigi L., 2016, *ApJ*, **827**, 126
- Berg D. A., Pogge R. W., Skillman E. D., Croxall K. V., Moustakas J., Rogers N. S. J., Sun J., 2020, *ApJ*, **893**, 96
- Berg D. A., Chisholm J., Erb D. K., Skillman E. D., Pogge R. W., Olivier G. M., 2021, *ApJ*, **922**, 170
- Bergvall N., Johansson L., Olofsson K., 1986, *A&A*, **166**, 92
- Bertram T., Eckart A., Fischer S., Zuther J., Straubmeier C., Wisotzki L., Krips M., 2007, *A&A*, **470**, 571
- Bianchin M., et al., 2022, *MNRAS*, **510**, 639
- Bradford J. D., Geha M. C., Greene J. E., Reines A. E., Dickey C. M., 2018, *ApJ*, **861**, 50
- Bresolin F., Kennicutt Jr. R. C., Garnett D. R., 1999, *ApJ*, **510**, 104
- Bresolin F., Garnett D. R., Kennicutt Robert C. J., 2004, *ApJ*, **615**, 228
- Bresolin F., Schaerer D., González Delgado R. M., Stasińska G., 2005, *A&A*, **441**, 981
- Bresolin F., Gieren W., Kudritzki R.-P., Pietrzyński G., Urbaneja M. A., Carraro G., 2009, *ApJ*, **700**, 309
- Brinchmann J., Charlot S., White S. D. M., Tremonti C., Kauffmann G., Heckman T., Brinkmann J., 2004, *MNRAS*, **351**, 1151
- Brown J. S., Martini P., Andrews B. H., 2016, *MNRAS*, **458**, 1529
- Bruzual G., Charlot S., 2003, *MNRAS*, **344**, 1000
- Bundy et al. K., 2015, *ApJ*, **798**, 7
- Cardelli J. A., Clayton G. C., Mathis J. S., 1989, *ApJ*, **345**, 245
- Carvalho S. P., et al., 2020, *MNRAS*, **492**, 5675
- Castellanos M., Díaz A. I., Terlevich E., 2002, *MNRAS*, **329**, 315
- Castro C. S., Dors O. L., Cardaci M. V., Hägele G. F., 2017, *MNRAS*, **467**, 1507
- Cid Fernandes R., Mateus A., Sodré L., Stasińska G., Gomes J. M., 2005, *MNRAS*, **358**, 363
- Cid Fernandes R., Stasińska G., Schlickmann M. S., Mateus A., Vale Asari N., Schoenell W., Sodré L., 2010, *MNRAS*, **403**, 1036
- Coil A. L., et al., 2015, *ApJ*, **801**, 35
- Combes F., et al., 2019, *A&A*, **623**, A79
- Contini M., 2019, *MNRAS*, **488**, 4487
- Cooke R. J., Fumagalli M., 2018, *Nature Astronomy*, **2**, 957
- Copetti M. V. F., Pastoriza M. G., Dottori H. A., 1986, *A&A*, **156**, 111
- Crowther P. A., Dessart L., 1998, *MNRAS*, **296**, 622
- Crowther P. A., Hadfield L. J., Schild H., Schmutz W., 2004, *A&A*, **419**, L17
- Croxall K. V., Pogge R. W., Berg D. A., Skillman E. D., Moustakas J., 2015, *ApJ*, **808**, 42
- Croxall K. V., Pogge R. W., Berg D. A., Skillman E. D., Moustakas J., 2016, *ApJ*, **830**, 4
- Cruz-Gonzalez I., Guichard J., Serrano A., Carrasco L., 1991, *PASP*, **103**, 888
- D’Agostino J. J., Kewley L. J., Groves B., Byler N., Sutherland R. S., Nicholls D., Leitherer C., Stanway E. R., 2019, *ApJ*, **878**, 2
- Davies R. I., Sternberg A., Lehnert M. D., Tacconi-Garman L. E., 2005, *ApJ*, **633**, 105
- Davison A. C., Hinkley D. V., 1997, Bootstrap methods and their application. No. 1, Cambridge university press
- Deharveng L., Peña M., Caplan J., Costero R., 2000, *MNRAS*, **311**, 329
- Delgado-Inglada G., Morisset C., Stasińska G., 2014, *MNRAS*, **440**, 536
- Diamond-Stanic A. M., Rieke G. H., 2012, *ApJ*, **746**, 168
- Dinerstein H. L., Shields G. A., 1986, *ApJ*, **311**, 45
- Dopita M. A., et al., 2006, *ApJS*, **167**, 177
- Dopita M. A., et al., 2015, *ApJS*, **217**, 12
- Dors O. L., 2021, *MNRAS*, **507**, 466
- Dors Jr. O. L., Krabbe A., Hägele G. F., Pérez-Montero E., 2011, *MNRAS*, **415**, 3616
- Dors Oli L. J., Riffel R. A., Cardaci M. V., Hägele G. F., Krabbe Á. C., Pérez-Montero E., Rodrigues I., 2012, *MNRAS*, **422**, 252

- Dors O. L., et al., 2013, *MNRAS*, **432**, 2512
- Dors O. L., Cardaci M. V., Hägele G. F., Krabbe A. C., 2014, *MNRAS*, **443**, 1291
- Dors O. L., Cardaci M. V., Hägele G. F., Rodrigues I., Grebel E. K., Pilyugin L. S., Freitas-Lemes P., Krabbe A. C., 2015, *MNRAS*, **453**, 4102
- Dors O. L. J., Arellano-Córdova K. Z., Cardaci M. V., Hägele G. F., 2017, *MNRAS*, **468**, L113
- Dors O. L., Agarwal B., Hägele G. F., Cardaci M. V., Rydberg C.-E., Riffel R. A., Oliveira A. S., Krabbe A. C., 2018, *MNRAS*, **479**, 2294
- Dors O. L., et al., 2020a, *MNRAS*, **492**, 468
- Dors O. L., Maiolino R., Cardaci M. V., Hägele G. F., Krabbe A. C., Pérez-Montero E., Armah M., 2020b, *MNRAS*, **496**, 3209
- Dors O. L., Contini M., Riffel R. A., Pérez-Montero E., Krabbe A. C., Cardaci M. V., Hägele G. F., 2021, *MNRAS*, **501**, 1370
- Dottori H. A., 1981, *Ap&SS*, **80**, 267
- Dressel L. L., Bania T. M., Oconnell R. W., 1982, *ApJ*, **259**, 55
- Durret F., 1994, *A&AS*, **105**, 57
- Ellison S. L., Sánchez S. F., Ibarra-Medel H., Antonio B., Mendel J. T., Barrera-Ballesteros J., 2018, *MNRAS*, **474**, 2039
- Ellison S. L., Brown T., Catinaella B., Cortese L., 2019, *MNRAS*, **482**, 5694
- Faulkner D. J., Aller L. H., 1965, *MNRAS*, **130**, 393
- Feltre A., Charlot S., Gutkin J., 2016, *MNRAS*, **456**, 3354
- Ferland G. J., Izotov Y., Peimbert A., Peimbert M., Porter R. L., Skillman E., Steigman G., 2010, in Charbonnel C., Tosi M., Primas F., Chiappini C., eds, *Proceedings of the International Astronomical Union Vol. 268, Light Elements in the Universe*. pp 163–167, doi:10.1017/S1743921310004011
- Ferland G. J., et al., 2013, *RMXAA*, **49**, 137
- Fernández-Ontiveros J. A., Spinoglio L., Pereira-Santaella M., Malkan M. A., Andreani P., Dasyra K. M., 2016, *ApJS*, **226**, 19
- Fernández V., Terlevich E., Díaz A. I., Terlevich R., 2019, *MNRAS*, **487**, 3221
- Flury S. R., Moran E. C., 2020, *MNRAS*, **496**, 2191
- Froese Fischer C., Tachiev G., 2004, *Atomic Data and Nuclear Data Tables*, **87**, 1
- García-Burillo S., et al., 2014, *A&A*, **567**, A125
- García-Vargas M. L., Bressan A., Díaz A. I., 1995, *A&AS*, **112**, 35
- Garnett D. R., Kennicutt Robert C. J., 1994, *ApJ*, **426**, 123
- Garnett D. R., Kennicutt Robert C. J., Bresolin F., 2004, *ApJ*, **607**, L21
- Gaskell C. M., 2017, *MNRAS*, **467**, 226
- Glazebrook K., et al., 2003, *ApJ*, **587**, 55
- Gómez P. L., et al., 2003, *ApJ*, **584**, 210
- Grevesse N., Asplund M., Sauval A. J., Scott P., 2010, *Ap&SS*, **328**, 179
- Groves B. A., Heckman T. M., Kauffmann G., 2006, *MNRAS*, **371**, 1559
- Hägele G. F., Díaz Á. I., Terlevich E., Terlevich R., Pérez-Montero E., Cardaci M. V., 2008, *MNRAS*, **383**, 209
- Halpern J. P., 1982, PhD thesis, Harvard University, Cambridge, MA.
- Halpern J. P., Steiner J. E., 1983, *ApJ*, **269**, L37
- Hayden-Pawson C., et al., 2022, *MNRAS*, **512**, 2867
- Heard C. Z. P., Gaskell C. M., 2016, *MNRAS*, **461**, 4227
- Heckman T. M., Balick B., 1979, *A&A*, **79**, 350
- Ho L. C., 1999, *ApJ*, **516**, 672
- Ho L. C., Darling J., Greene J. E., 2008, *ApJS*, **177**, 103
- Hsyu T., Cooke R. J., Prochaska J. X., Bolte M., 2020, *ApJ*, **896**, 77
- Hummer D. G., Storey P. J., 1987, *MNRAS*, **224**, 801
- Hutchings J. B., Gower A. C., Price R., 1987, *AJ*, **93**, 6
- Izotov Y. I., Thuan T. X., 1998, *ApJ*, **500**, 188
- Izotov Y. I., Thuan T. X., 2008, *ApJ*, **687**, 133
- Izotov Y. I., Thuan T. X., Lipovetsky V. A., 1997, *ApJS*, **108**, 1
- Izotov Y. I., Chaffee F. H., Foltz C. B., Green R. F., Guseva N. G., Thuan T. X., 1999, *ApJ*, **527**, 757
- Izotov Y. I., Stasińska G., Meynet G., Guseva N. G., Thuan T. X., 2006, *A&A*, **448**, 955
- Izotov Y. I., Thuan T. X., Stasińska G., 2007, *ApJ*, **662**, 15
- Izotov Y. I., Guseva N. G., Fricke K. J., Papaderos P., 2009, *A&A*, **503**, 61
- Izotov Y. I., Stasińska G., Guseva N. G., 2013, *A&A*, **558**, A57
- Izotov Y. I., Thuan T. X., Guseva N. G., 2014, *MNRAS*, **445**, 778
- Jarvis M. E., et al., 2020, *MNRAS*, **498**, 1560
- Ji X., Yan R., 2022, *A&A*, **659**, A112
- Ji X., Yan R., Riffel R., Drory N., Zhang K., 2020, *MNRAS*, **496**, 1262
- Johnson H. M., 1959, *PASP*, **71**, 301
- Johnson J. W., Weinberg D. H., Vincenzo F., Bird J. C., Griffith E. J., 2022, arXiv e-prints, p. arXiv:2202.04666
- Jura M., 1973, *ApJ*, **181**, 627
- Kauffmann G., et al., 2003, *MNRAS*, **346**, 1055
- Kawasaki K., Nagao T., Toba Y., Terao K., Matsuoka K., 2017, *ApJ*, **842**, 44
- Kennicutt Robert C. J., Bresolin F., Garnett D. R., 2003, *ApJ*, **591**, 801
- Kewley L. J., Ellison S. L., 2008, *ApJ*, **681**, 1183
- Kewley L. J., Dopita M. A., Sutherland R. S., Heisler C. A., Trevena J., 2001, *ApJ*, **556**, 121
- Kewley L. J., Groves B., Kauffmann G., Heckman T., 2006, *MNRAS*, **372**, 961
- Kewley L. J., Nicholls D. C., Sutherland R. S., 2019, *ARA&A*, **57**, 511
- Kinkel U., Rosa M. R., 1994, *A&A*, **282**, L37
- Kisielius R., Storey P. J., Ferland G. J., Keenan F. P., 2009, *MNRAS*, **397**, 903
- Koski A. T., 1978, *ApJ*, **223**, 56
- Krabbe A. C., Pastoriza M. G., Winge C., Rodrigues I., Dors O. L., Ferreira D. L., 2011, *MNRAS*, **416**, 38
- Krabbe A. C., Rosa D. A., Pastoriza M. G., Hägele G. F., Cardaci M. V., Dors O. L. J., Winge C., 2017, *MNRAS*, **467**, 27
- Kraemer S. B., Wu C.-C., Crenshaw D. M., Harrington J. P., 1994, *ApJ*, **435**, 171
- Kumari N., Amorín R., Pérez-Montero E., Vílchez J., Maiolino R., 2021, *MNRAS*, **508**, 1084
- Kunth D., Sargent W. L. W., 1983, *ApJ*, **273**, 81
- Kurichin O. A., Kisilitsyn P. A., Klimenko V. V., Balashev S. A., Ivanchik A. V., 2021, *MNRAS*, **502**, 3045
- Leitherer C., Heckman T. M., 1995, *ApJS*, **96**, 9
- Leitherer C., et al., 1999, *ApJS*, **123**, 3
- Levesque E. M., Leitherer C., Ekstrom S., Meynet G., Schaerer D., 2012, *ApJ*, **751**, 67
- Liang Y. C., Hammer F., Yin S. Y., Flores H., Rodrigues M., Yang Y. B., 2007, *A&A*, **473**, 411
- Ludwig R. R., Greene J. E., Barth A. J., Ho L. C., 2012, *ApJ*, **756**, 51
- Luridiana V., Morisset C., Shaw R. A., 2015, *A&A*, **573**
- Maeder A., Meynet G., 1994, *A&A*, **287**, 803
- Mateus A., Sodré L., Cid Fernandes R., Stasińska G., Schoenell W., Gomes J. M., 2006, *MNRAS*, **370**, 721
- Mathis J. S., 1957, *ApJ*, **125**, 328
- Mathis J. S., 1962, *ApJ*, **136**, 374
- Mathis J. S., 1965, *PASP*, **77**, 90
- Mathis J. S., 1982, *ApJ*, **261**, 195
- Mayya Y. D., Prabhu T. P., 1996, *AJ*, **111**, 1252
- Mazzalay X., et al., 2013, *MNRAS*, **428**, 2389
- McCall B. J., et al., 2003, *Nature*, **422**, 500
- Méndez-Delgado J. E., Esteban C., García-Rojas J., Arellano-Córdova K. Z., Valerdi M., 2020, *MNRAS*, **496**, 2726
- Méndez-Delgado J. E., Amayo A., Arellano-Córdova K. Z., Esteban C., García-Rojas J., Carigi L., Delgado-Inglada G., 2022, *MNRAS*, **510**, 4436
- Mezcua M., Prieto M. A., Fernández-Ontiveros J. A., Tristram K., Neumayer N., Kotilainen J. K., 2015, *MNRAS*, **452**, 4128
- Miller B. P., Brandt W. N., Schneider D. P., Gibson R. R., Steffen A. T., Wu J., 2011, *ApJ*, **726**, 20
- Mollá M., Díaz A. I., 2005, *MNRAS*, **358**, 521
- Monteiro A. F., Dors O. L., 2021, *MNRAS*, **508**, 3023
- Moustakas J., Kennicutt Robert C. J., 2006, *ApJS*, **164**, 81
- Nakajima K., Maiolino R., 2022, *MNRAS*, **513**, 5134
- Nicholls D. C., Sutherland R. S., Dopita M. A., Kewley L. J., Groves B. A., 2017, *MNRAS*, **466**, 4403
- Olive K. A., Skillman E. D., 2004, *ApJ*, **617**, 29
- Olive K. A., Steigman G., Skillman E. D., 1997, *ApJ*, **483**, 788
- Osterbrock D. E., Ferland G. J., 2006, *Astrophysics of gaseous nebulae and active galactic nuclei*. UNIVERSITY SCIENCE BOOKS
- Osterbrock D. E., Miller J. S., 1975, *ApJ*, **197**, 535
- Owen A. B., 2007, *Contemporary Mathematics*, **443**, 59

- Pagel B. E. J., Simonson E. A., Terlevich R. J., Edmunds M. G., 1992, *MNRAS*, **255**, 325  
 Pauldrach A. W. A., Hoffmann T. L., Lennon M., 2001, *A&A*, **375**, 161  
 Peña M., 1986, *PASP*, **98**, 1061  
 Peimbert M., 1967, *ApJ*, **150**, 825  
 Peimbert M., 2008, arXiv e-prints, p. arXiv:0811.2980  
 Peimbert M., Spinrad H., 1970, *ApJ*, **159**, 809  
 Peimbert M., Torres-Peimbert S., 1974, *ApJ*, **193**, 327  
 Peimbert M., Torres-Peimbert S., 1976, *ApJ*, **203**, 581  
 Peimbert M., Torres-Peimbert S., Ruiz M. T., 1992, *RMXAA*, **24**, 155  
 Peimbert M., Peimbert A., Ruiz M. T., 2000, *ApJ*, **541**, 688  
 Peimbert A., Peimbert M., Luridiana V., 2002, *ApJ*, **565**, 668  
 Peimbert M., Peimbert A., Luridiana V., Ruiz M. T., 2003, in Perez E., Gonzalez Delgado R. M., Tenorio-Tagle G., eds, *Astronomical Society of the Pacific Conference Series Vol. 297, Star Formation Through Time*. p. 81 (arXiv:astro-ph/0211497)  
 Peimbert M., Luridiana V., Peimbert A., 2007, *ApJ*, **666**, 636  
 Peimbert A., Peña-Guerrero M. A., Peimbert M., 2012, *ApJ*, **753**, 39  
 Peimbert A., Peimbert M., Luridiana V., 2016, *RMXAA*, **52**, 419  
 Peimbert M., Peimbert A., Delgado-Inglada G., 2017, *PASP*, **129**, 082001  
 Pérez-Díaz B., Masegosa J., Márquez I., Pérez-Montero E., 2021, *MNRAS*, **505**, 4289  
 Pérez-Montero E., 2014, *MNRAS*, **441**, 2663  
 Pérez-Montero E., 2017, *PASP*, **129**, 043001  
 Pérez-Montero E., Díaz A. I., 2003, *MNRAS*, **346**, 105  
 Pérez-Montero et al. E., 2013, *A&A*, **549**, A25  
 Pérez-Montero E., Dors O. L., Vilchez J. M., García-Benito R., Cardaci M. V., Hägele G. F., 2019, *MNRAS*, **489**, 2652  
 Phillips M. M., Charles P. A., Baldwin J. A., 1983, *ApJ*, **266**, 485  
 Pilyugin L. S., 2003, *A&A*, **399**, 1003  
 Pilyugin L. S., Vilchez J. M., Mattsson L., Thuan T. X., 2012, *MNRAS*, **421**, 1624  
 Pindao M., Schaerer D., González Delgado R. M., Stasińska G., 2002, *A&A*, **394**, 443  
 Planck Collaboration 2020, *A&A*, **641**, A6  
 Porter R. L., Ferland G. J., Storey P. J., Detisch M. J., 2012, *MNRAS*, **425**, L28  
 Ramakrishnan V., et al., 2019, *MNRAS*, **487**, 444  
 Revalski M., Crenshaw D. M., Kraemer 2018a, *ApJ*, **856**, 46  
 Revalski M., et al., 2018b, *ApJ*, **867**, 88  
 Revalski M., et al., 2021, *ApJ*, **910**, 139  
 Richardson C. T., Allen J. T., Baldwin J. A., Hewett P. C., Ferland G. J., 2014, *MNRAS*, **437**, 2376  
 Riffel R. A., Storch-Bergmann T., Nagar N. M., 2010, *MNRAS*, **404**, 166  
 Riffel R., Rodríguez-Ardila A., Aleman I., Brotherton M. S., Pastoriza M. G., Bonatto C., Dors O. L., 2013, *MNRAS*, **430**, 2002  
 Riffel R. A., et al., 2018, *MNRAS*, **474**, 1373  
 Riffel R., et al., 2019, *MNRAS*, **486**, 3228  
 Riffel R. A., et al., 2021a, *MNRAS*, **501**, L54  
 Riffel R., et al., 2021b, *MNRAS*, **501**, 4064  
 Riffel R. A., et al., 2021c, *MNRAS*, **504**, 3265  
 Riffel R. A., Dors O. L., Krabbe A. C., Esteban C., 2021d, *MNRAS*, **506**, L11  
 Rodríguez-Ardila A., Pastoriza M. G., Viegas S., Sigut T. A. A., Pradhan A. K., 2004, *A&A*, **425**, 457  
 Rodríguez-Ardila A., Riffel R., Pastoriza M. G., 2005, *MNRAS*, **364**, 1041  
 Rogers N. S. J., Skillman E. D., Pogge R. W., Berg D. A., Moustakas J., Croxall K. V., Sun J., 2021, *ApJ*, **915**, 21  
 Rosa M., 1985, *The Messenger*, **39**, 15  
 Ruschel-Dutra D., de Oliveira B. D., 2020, danielrd6/ifscube v1.0, doi:10.5281/zenodo.3945237, <https://doi.org/10.5281/zenodo.3945237>  
 Ruschel-Dutra D., et al., 2021, *MNRAS*, **507**, 74  
 Russell S. C., Dopita M. A., 1992, *ApJ*, **384**, 508  
 Sánchez et al. S. F., 2012, *A&A*, **538**, A8  
 Sánchez S. F., et al., 2017, *MNRAS*, **469**, 2121  
 Sauer D., Jedamzik K., 2002, *A&A*, **381**, 361  
 Schlafly E. F., Finkbeiner D. P., 2011, *ApJ*, **737**, 103  
 Schlegel D. J., Finkbeiner D. P., Davis M., 1998, *ApJ*, **500**, 525  
 Schönnell Astor J. J., Storch-Bergmann T., Riffel R. A., Riffel R., 2017, *MNRAS*, **464**, 1771  
 Searle L., Sargent W. L. W., 1972, *ApJ*, **173**, 25  
 Shields G. A., 1974, *ApJ*, **191**, 309  
 Shields G. A., Oke J. B., 1975, *ApJ*, **197**, 5  
 Shields G. A., Tinsley B. M., 1976, *ApJ*, **203**, 66  
 Shuder J. M., 1980, *ApJ*, **240**, 32  
 Shuder J. M., Osterbrock D. E., 1981, *ApJ*, **250**, 55  
 Skillman E. D., 1991, *PASP*, **103**, 919  
 Skillman E. D., Terlevich R. J., Kennicutt Robert C. J., Garnett D. R., Terlevich E., 1994, *ApJ*, **431**, 172  
 Skillman E., Aver E., Olive K., 2012, *Memorie della Societa Astronomica Italiana Supplementi*, **22**, 164  
 Stanway E. R., Eldridge J. J., 2018, *MNRAS*, **479**, 75  
 Storch-Bergmann T., Schmitt H. R., Calzetti D., Kinney A. L., 1998, *AJ*, **115**, 909  
 Storch-Bergmann T., McGregor P. J., Riffel R. A., Simões Lopes R., Beck T., Dopita M., 2009, *MNRAS*, **394**, 1148  
 Storey P. J., Hummer D. G., 1995, *MNRAS*, **272**, 41  
 Storey P. J., Zeippen C. J., 2000, *MNRAS*, **312**, 813  
 Storey P. J., Sochi T., Badnell N. R., 2014, *MNRAS*, **441**, 3028  
 Tananbaum H., et al., 1979, *ApJ*, **234**, L9  
 Thomas A. D., et al., 2017, *ApJS*, **232**, 11  
 Toribio San Cipriano L., Domínguez-Guzmán G., Esteban C., García-Rojas J., Mesa-Delgado A., Bresolin F., Rodríguez M., Simón-Díaz S., 2017, *MNRAS*, **467**, 3759  
 Torres-Peimbert S., Peimbert M., 1977, *RMXAA*, **2**, 181  
 Vale Asari N., Stasińska G., Morisset C., Cid Fernandes R., 2016, *MNRAS*, **460**, 1739  
 Valerdi M., Peimbert A., Peimbert M., Sixtos A., 2019, *ApJ*, **876**, 98  
 Valerdi M., Peimbert A., Peimbert M., 2021a, *MNRAS*, **505**, 3624  
 Valerdi M., Barrera-Ballesteros J. K., Sánchez S. F., Espinosa-Ponce C., Carigi L., Mejía-Narváez A., 2021b, *MNRAS*, **505**, 5460  
 Vaona L., Ciroi S., Di Mille F., Cracco V., La Mura G., Rafanelli P., 2012, *MNRAS*, **427**, 1266  
 Vilchez J. M., Pagel B. E. J., 1988, *MNRAS*, **231**, 257  
 Vincenzo F., et al., 2021, arXiv e-prints, p. arXiv:2106.03912  
 Wampler E. J., 1971, *ApJ*, **164**, 1  
 Wiese W. L., Fuhr J. R., Deters T. M., 1996, *Atomic transition probabilities of carbon, nitrogen, and oxygen : a critical data compilation*  
 Williams R. E., 1971, *ApJ*, **167**, L27  
 Worrall D. M., Giommi P., Tananbaum H., Zamorani G., 1987, *ApJ*, **313**, 596  
 Wyse A. B., 1942, *ApJ*, **95**, 356  
 Yates R. M., Kauffmann G., Guo Q., 2012, *MNRAS*, **422**, 215  
 Yates R. M., Schady P., Chen T. W., Schweyer T., Wiseman P., 2020, *A&A*, **634**, A107  
 York D. G., Adelman J., Anderson John E. J., Anderson S. F., Annis J., Bahcall 2000, *AJ*, **120**, 1579  
 Zastrow J., Oey M. S., Veilleux S., McDonald M., 2013, *ApJ*, **779**, 76  
 Zhang K., Wang T., Dong X., Lu H., 2008, *ApJ*, **685**, L109  
 Zhang Z. T., Liang Y. C., Hammer F., 2013, *MNRAS*, **430**, 2605  
 Zhu S. F., Brandt W. N., Wu J., Garmire G. P., Miller B. P., 2019, *MNRAS*, **482**, 2016  
 Zhu S. F., Timlin J. D., Brandt W. N., 2021, *MNRAS*, **505**, 1954  
 Zinchenko I. A., Dors O. L., Hägele G. F., Cardaci M. V., Krabbe A. C., 2019, *MNRAS*, **483**, 1901  
 do Nascimento J. C., et al., 2022, *MNRAS*, **513**, 807

**Table 1.** Reddening corrected emission line intensities (in relation to  $H\beta = 1.0$ ) of the Seyfert 2 sample described in Sect. 2.1.

| ObjID                 | [O II]<br>$\lambda 3727$ | [O III]<br>$\lambda 4363$ | He II<br>$\lambda 4686$ | [O III]<br>$\lambda 5007$ | He I<br>$\lambda 5876$ | H $\alpha$<br>$\lambda 6563$ | [N II]<br>$\lambda 6583$ | [S II]<br>$\lambda 6716$ | [S II]<br>$\lambda 6731$ | Ref. |
|-----------------------|--------------------------|---------------------------|-------------------------|---------------------------|------------------------|------------------------------|--------------------------|--------------------------|--------------------------|------|
| J013957.81-004504.2   | 2.31±0.22                | 0.06±0.01                 | 0.16±0.01               | 7.45±0.11                 | 0.06±0.02              | 2.86±0.04                    | 2.51±0.04                | 0.79±0.02                | 0.67±0.02                | 1    |
| J033923.14-054841.5   | 2.63±0.20                | 0.09±0.01                 | 0.26±0.01               | 7.90±0.02                 | 0.09±0.01              | 2.86±0.01                    | 1.52±0.01                | 0.53±0.01                | 0.45±0.01                | 1    |
| J074257.23+333217.9   | 1.93±0.09                | 0.04±0.01                 | 0.08±0.01               | 3.03±0.01                 | 0.10±0.01              | 2.86±0.01                    | 1.63±0.01                | 0.49±0.01                | 0.43±0.01                | 1    |
| J093509.12+002557.4   | 2.58±0.10                | 0.07±0.01                 | 0.11±0.01               | 3.86±0.02                 | 0.07±0.01              | 2.86±0.01                    | 2.12±0.01                | 0.59±0.01                | 0.54±0.01                | 1    |
| J095759.45+022810.5   | 3.17±0.23                | 0.08±0.01                 | 0.26±0.01               | 7.59±0.07                 | 0.08±0.01              | 2.86±0.03                    | 2.12±0.02                | 0.84±0.01                | 0.69±0.01                | 1    |
| J100602.50+071131.8   | 6.21±0.04                | 0.15±0.01                 | 0.27±0.01               | 7.61±0.01                 | 0.09±0.01              | 2.86±0.01                    | 1.85±0.01                | 0.62±0.01                | 0.51±0.01                | 1    |
| J100921.26+013334.5   | 3.31±0.28                | 0.14±0.01                 | 0.31±0.01               | 12.29±0.07                | 0.11±0.01              | 2.86±0.02                    | 2.89±0.02                | 0.72±0.01                | 0.74±0.01                | 1    |
| J101754.72-002811.9   | 2.34±0.14                | 0.12±0.01                 | 0.27±0.01               | 6.05±0.03                 | 0.11±0.01              | 2.86±0.01                    | 1.75±0.01                | 0.53±0.01                | 0.43±0.01                | 1    |
| J102039.81+642435.8   | 4.38±1.00                | 0.17±0.01                 | 0.25±0.01               | 8.82±0.06                 | 0.07±0.01              | 2.86±0.02                    | 2.72±0.02                | 0.76±0.01                | 0.58±0.01                | 1    |
| J112850.39+021016.2   | 2.97±0.20                | 0.28±0.02                 | 0.28±0.02               | 6.75±0.15                 | 0.16±0.03              | 2.86±0.07                    | 1.79±0.05                | 0.64±0.03                | 0.51±0.03                | 1    |
| Mrk 176               | 3.54                     | 0.32                      | 0.43                    | 14.36                     | 0.10                   | 2.81                         | 2.99                     | 0.56                     | 0.54                     | 2    |
| 3C 33                 | 4.93                     | 0.32                      | 0.26                    | 12.68                     | 0.096                  | 2.63                         | 1.76                     | 0.87                     | 0.73                     | 2    |
| Mrk 3                 | 3.52                     | 0.24                      | 0.18                    | 12.67                     | 0.084                  | 3.10                         | 3.18                     | 0.73                     | 0.82                     | 2    |
| Mrk 573               | 2.92                     | 0.18                      | 0.36                    | 12.12                     | 0.10                   | 2.95                         | 2.57                     | 0.75                     | 0.80                     | 2    |
| Mrk 78                | 4.96                     | 0.14                      | 0.35                    | 11.94                     | 0.10                   | 2.46                         | 2.32                     | 0.68                     | 0.61                     | 2    |
| Mrk 348               | 4.45                     | 0.26                      | 0.22                    | 11.74                     | 0.12                   | 2.76                         | 2.28                     | 1.09                     | 1.25                     | 2    |
| Mrk 34                | 3.43                     | 0.15                      | 0.28                    | 11.46                     | 0.12                   | 2.99                         | 2.18                     | 0.82                     | 0.80                     | 2    |
| Mrk 1                 | 2.79                     | 0.21                      | 0.30                    | 10.95                     | 0.10                   | 2.66                         | 2.21                     | 0.49                     | 0.52                     | 2    |
| Mrk 270               | 5.64                     | 0.28                      | 0.22                    | 8.71                      | 0.19                   | 3.14                         | 2.93                     | 1.21                     | 1.39                     | 2    |
| III Zw 55             | 3.19                     | 0.25                      | 0.20                    | 6.92                      | 0.090                  | 2.86                         | 3.87                     | 0.66                     | 0.74                     | 2    |
| 3C 452                | 4.81                     | 0.18                      | 0.059                   | 6.85                      | 0.17                   | 2.98                         | 3.58                     | 1.10                     | 0.77                     | 2    |
| Mrk 198               | 2.51                     | 0.12                      | 0.075                   | 5.56                      | 0.11                   | 3.02                         | 2.26                     | 0.89                     | 0.68                     | 2    |
| Mrk 268               | 3.75                     | 0.25                      | 0.078                   | 4.82                      | 0.095                  | 3.38                         | 4.94                     | 1.28                     | 1.08                     | 2    |
| NGC 2110              | 4.38                     | 0.17                      | 0.06                    | 4.76                      | 0.14                   | 2.66                         | 3.76                     | 1.52                     | 1.42                     | 3    |
| ESO 138 G1            | 2.35 ± 0.08              | 0.34 ± 0.02               | 0.30 ± 0.02             | 8.71 ± 0.36               | 0.11 ± 0.02            | 3.00 ± 0.13                  | 0.68 ± 0.03              | 0.47 ± 0.03              | 0.48 ± 0.03              | 4    |
| NGC 3081              | 2.19                     | 0.24                      | 0.44                    | 12.62                     | 0.10                   | 2.84                         | 2.47                     | 0.60                     | 0.66                     | 5    |
| NGC 4388              | 2.73                     | 0.16                      | 0.21                    | 10.55                     | 0.11                   | 2.84                         | 1.50                     | 0.72                     | 0.62                     | 5    |
| NGC 5135              | 2.06                     | 0.11                      | 0.20                    | 4.42                      | 0.10                   | 2.83                         | 2.50                     | 0.40                     | 0.38                     | 5    |
| NGC 5728              | 3.50                     | 0.47                      | 0.32                    | 10.85                     | 0.15                   | 2.83                         | 3.93                     | 0.44                     | 0.435                    | 5    |
| Akn 347 <sup>a</sup>  | 2.72                     | 0.42                      | 0.33                    | 15.15                     | 0.12                   | 3.18                         | 3.93                     | 0.93                     | 0.89                     | 6    |
| UM 16 <sup>a</sup>    | 3.05                     | 0.25                      | 0.37                    | 13.8                      | 0.18                   | 2.77                         | 1.66                     | 0.47                     | 0.48                     | 6    |
| Mrk 612 <sup>a</sup>  | 1.90                     | 0.20                      | 0.16                    | 9.09                      | 0.20                   | 2.81                         | 3.81                     | 0.78                     | 0.57                     | 6    |
| Mrk 573               | 2.13 ± 0.05              | 0.14 ± 0.011              | 0.29 ± 0.02             | 10.26 ± 0.05              | 0.09 ± 0.01            | 2.86 ± 0.05                  | 2.34 ± 0.04              | 0.79 ± 0.01              | 0.73 ± 0.02              | 7    |
| NGC 2992              | 3.38 ± 0.11              | 0.21 ± 0.027              | 0.07 ± 0.02             | 5.25 ± 0.09               | 0.08 ± 0.01            | 2.86 ± 0.02                  | 2.70 ± 0.03              | 0.52 ± 0.01              | 0.47 ± 0.01              | 7    |
| IC 2560               | 1.65 ± 0.02              | 0.16 ± 0.017              | 0.26 ± 0.06             | 10.69 ± 0.10              | 0.10 ± 0.01            | 2.86 ± 0.08                  | 2.88 ± 0.07              | 0.74 ± 0.02              | 0.79 ± 0.01              | 7    |
| NGC 5664              | 1.83 ± 0.05              | 0.02 ± 0.006              | 0.12 ± 0.01             | 2.95 ± 0.07               | 0.09 ± 0.01            | 2.86 ± 0.12                  | 1.70 ± 0.11              | 0.58 ± 0.03              | 0.43 ± 0.07              | 7    |
| NGC 5728              | 2.32 ± 0.03              | 0.11 ± 0.016              | 0.17 ± 0.01             | 9.10 ± 0.11               | 0.08 ± 0.01            | 2.86 ± 0.13                  | 3.53 ± 0.12              | 0.97 ± 0.06              | 0.66 ± 0.03              | 7    |
| ESO 339-G11           | 2.39 ± 0.12              | 0.14 ± 0.074              | 0.13 ± 0.01             | 7.86 ± 0.05               | 0.09 ± 0.01            | 2.86 ± 0.12                  | 3.97 ± 0.09              | 0.69 ± 0.02              | 0.64 ± 0.02              | 7    |
| NGC 6890              | 1.66 ± 0.04              | 0.23 ± 0.037              | 0.19 ± 0.01             | 10.90 ± 0.03              | 0.06 ± 0.02            | 2.86 ± 0.10                  | 2.85 ± 0.09              | 0.43 ± 0.01              | 0.39 ± 0.01              | 7    |
| IC 5063               | 2.88 ± 0.06              | 0.12 ± 0.008              | 0.09 ± 0.01             | 7.83 ± 0.07               | 0.08 ± 0.01            | 2.86 ± 0.06                  | 1.81 ± 0.06              | 0.77 ± 0.06              | 0.77 ± 0.02              | 7    |
| NGC 7130              | 1.75 ± 0.03              | 0.09 ± 0.005              | 0.15 ± 0.01             | 4.62 ± 0.05               | 0.05 ± 0.01            | 2.86 ± 0.01                  | 3.64 ± 0.00              | 0.50 ± 0.05              | 0.50 ± 0.01              | 7    |
| NGC 7582              | 1.24 ± 0.02              | 0.03 ± 0.002              | 0.11 ± 0.05             | 2.14 ± 0.10               | 0.10 ± 0.01            | 2.86 ± 0.02                  | 1.86 ± 0.02              | 0.40 ± 0.01              | 0.38 ± 0.01              | 7    |
| NGC 7590              | 3.32 ± 0.19              | 0.05 ± 0.027              | 0.07 ± 0.03             | 3.47 ± 0.16               | 0.06 ± 0.02            | 2.86 ± 0.11                  | 2.39 ± 0.18              | 0.98 ± 0.08              | 0.78 ± 0.04              | 7    |
| Cygnus A              | 5.01                     | 0.20                      | 0.27                    | 12.30                     | 0.07                   | 3.09                         | 6.16                     | 1.65                     | 1.51                     | 8    |
| Mrk 1157 <sup>a</sup> | 7.32                     | 0.25                      | 0.24                    | 9.81                      | 0.12                   | 2.82                         | 3.07                     | 0.65                     | 0.69                     | 9    |
| ESO 428-G14           | 2.22                     | 0.24                      | 0.19                    | 11.2                      | 0.16                   | 2.90                         | 3.13                     | 0.83                     | 0.88                     | 10   |
| ESO 137-G34           | 3.07                     | 0.11                      | 0.21                    | 9.35                      | 0.11                   | 3.10                         | 3.35                     | 0.98                     | 1.20                     | 11   |
| IC 3639               | 1.24                     | 0.09                      | 0.12                    | 5.68                      | 0.10                   | 3.10                         | 2.52                     | 0.68                     | 0.73                     | 11   |
| IC 4777               | 2.32                     | 0.07                      | 0.17                    | 6.34                      | 0.10                   | 3.10                         | 3.82                     | 0.92                     | 0.87                     | 11   |
| IC 4995               | 1.32                     | 0.20                      | 0.27                    | 11.87                     | 0.08                   | 3.10                         | 2.49                     | 0.68                     | 0.76                     | 11   |
| IRAS 11215-2806       | 2.48                     | 0.14                      | 0.25                    | 7.25                      | 0.11                   | 3.10                         | 1.74                     | 0.84                     | 0.79                     | 11   |

**Table 1** – *continued*

| ObjID         | [O II]<br>$\lambda 3727$ | [O III]<br>$\lambda 4363$ | He II<br>$\lambda 4686$ | [O III]<br>$\lambda 5007$ | He I<br>$\lambda 5876$ | H $\alpha$<br>$\lambda 6563$ | [N II]<br>$\lambda 6583$ | [S II]<br>$\lambda 6716$ | [S II]<br>$\lambda 6731$ | Ref. |
|---------------|--------------------------|---------------------------|-------------------------|---------------------------|------------------------|------------------------------|--------------------------|--------------------------|--------------------------|------|
| MCG-01-24-012 | 3.31                     | 0.16                      | 0.20                    | 8.51                      | 0.09                   | 3.10                         | 1.77                     | 0.97                     | 0.86                     | 11   |
| MCG-02-51-008 | 2.35                     | 0.05                      | 0.09                    | 4.23                      | 0.10                   | 3.10                         | 2.10                     | 0.84                     | 0.63                     | 11   |
| NGC 1125      | 3.13                     | 0.08                      | 0.14                    | 5.86                      | 0.10                   | 3.10                         | 2.30                     | 0.79                     | 0.78                     | 11   |
| NGC 1194      | 1.64                     | 0.08                      | 0.06                    | 6.21                      | 0.11                   | 3.10                         | 1.61                     | 1.01                     | 0.77                     | 11   |
| NGC 1320      | 0.98                     | 0.13                      | 0.37                    | 9.34                      | 0.11                   | 3.10                         | 2.21                     | 0.65                     | 0.62                     | 11   |
| NGC 3281      | 2.06                     | 0.09                      | 0.29                    | 7.71                      | 0.11                   | 3.10                         | 2.85                     | 0.88                     | 0.76                     | 11   |
| NGC 3393      | 2.23                     | 0.10                      | 0.26                    | 9.428                     | 0.12                   | 3.10                         | 3.13                     | 0.92                     | 0.93                     | 11   |
| NGC 4939      | 2.05                     | 0.12                      | 0.30                    | 10.03                     | 0.12                   | 3.10                         | 4.16                     | 0.91                     | 0.99                     | 11   |
| NGC 4968      | 1.64                     | 0.22                      | 0.27                    | 9.45                      | 0.12                   | 3.10                         | 3.28                     | 0.62                     | 0.65                     | 11   |
| NGC 5427      | 2.15                     | 0.13                      | 0.18                    | 7.96                      | 0.14                   | 3.10                         | 3.95                     | 0.80                     | 0.88                     | 11   |
| NGC 5643      | 2.74                     | 0.12                      | 0.24                    | 8.60                      | 0.04                   | 3.10                         | 2.98                     | 0.97                     | 0.91                     | 11   |
| NGC 5990      | 1.88                     | 0.01                      | 0.07                    | 2.16                      | 0.09                   | 3.10                         | 2.06                     | 0.48                     | 0.42                     | 11   |
| NGC 7682      | 2.85                     | 0.16                      | 0.19                    | 9.34                      | 0.09                   | 3.10                         | 3.03                     | 1.09                     | 1.17                     | 11   |
| PKS 1306-241  | 3.47                     | 0.04                      | 0.05                    | 3.82                      | 0.10                   | 3.10                         | 1.25                     | 0.69                     | 0.61                     | 11   |

References: (1) SDSS-DR7 (York et al. 2000), (2) Koski (1978), (3) Shuder (1980), (4) Alloin et al. (1992), (5) Phillips et al. (1983), (6) Shuder & Osterbrock (1981), (7) Dopita et al. (2015), (8) Osterbrock & Miller (1975), (9) Durret (1994), (10) Bergvall et al. (1986), (11) Thomas et al. (2017).

**Table 2.** Reddening corrected emission line intensities (in relation to  $H\beta = 1.0$ ) of the SF sample described in Sect. 2.2.

| ObjID              | [O II]<br>$\lambda 3727$ | [O III]<br>$\lambda 4363$ | He II<br>$\lambda 4686$ | [O III]<br>$\lambda 5007$ | He I<br>$\lambda 5876$ | H $\alpha$<br>$\lambda 6563$ | [N II]<br>$\lambda 6583$ | [S II]<br>$\lambda 6716$ | [S II]<br>$\lambda 6731$ | Ref. |
|--------------------|--------------------------|---------------------------|-------------------------|---------------------------|------------------------|------------------------------|--------------------------|--------------------------|--------------------------|------|
| NGC3184-72.8+91.3  | $1.82 \pm 0.01$          | $0.002 \pm 0.0010$        | $0.011 \pm 0.0010$      | $0.384 \pm 0.002$         | $0.083 \pm 0.001$      | $2.801 \pm 0.027$            | $0.923 \pm 0.008$        | $0.352 \pm 0.003$        | $0.253 \pm 0.002$        | 1    |
| NGC3184+14.9-139.6 | $2.47 \pm 0.04$          | $0.008 \pm 0.0010$        | $0.024 \pm 0.0030$      | $1.445 \pm 0.016$         | $0.081 \pm 0.006$      | $2.883 \pm 0.046$            | $0.789 \pm 0.012$        | $0.424 \pm 0.006$        | $0.303 \pm 0.005$        | 1    |
| NGC3184+80.0-148.2 | $2.88 \pm 0.03$          | $0.004 \pm 0.0010$        | $0.007 \pm 0.0010$      | $0.860 \pm 0.018$         | $0.110 \pm 0.010$      | $2.937 \pm 0.056$            | $0.828 \pm 0.016$        | $0.439 \pm 0.009$        | $0.321 \pm 0.007$        | 1    |
| NGC3184-93.3-142.3 | $2.20 \pm 0.03$          | $0.008 \pm 0.0020$        | $0.001 \pm 0.0010$      | $1.902 \pm 0.031$         | $0.129 \pm 0.010$      | $2.840 \pm 0.067$            | $0.543 \pm 0.013$        | $0.385 \pm 0.009$        | $0.263 \pm 0.006$        | 1    |
| NGC3184-172.5-30.2 | $2.32 \pm 0.01$          | $0.014 \pm 0.0010$        | $0.003 \pm 0.0010$      | $2.434 \pm 0.017$         | $0.124 \pm 0.004$      | $2.801 \pm 0.023$            | $0.485 \pm 0.005$        | $0.251 \pm 0.002$        | $0.183 \pm 0.001$        | 1    |
| +164.6+9.9         | $1.94 \pm 0.04$          | $0.002 \pm 0.0002$        | $0.009 \pm 0.0010$      | $0.980 \pm 0.020$         | $0.116 \pm 0.002$      | $3.015 \pm 0.060$            | $0.650 \pm 0.027$        | $0.161 \pm 0.006$        | $0.116 \pm 0.005$        | 2    |
| +17.3-235.4        | $2.43 \pm 0.03$          | $0.004 \pm 0.0005$        | $0.001 \pm 0.0004$      | $1.170 \pm 0.023$         | $0.110 \pm 0.002$      | $2.865 \pm 0.057$            | $0.650 \pm 0.020$        | $0.240 \pm 0.007$        | $0.170 \pm 0.005$        | 2    |
| +189.2-136.3       | $1.60 \pm 0.03$          | $0.006 \pm 0.0002$        | $0.010 \pm 0.0017$      | $2.603 \pm 0.052$         | $0.140 \pm 0.002$      | $2.815 \pm 0.056$            | $0.400 \pm 0.014$        | $0.113 \pm 0.004$        | $0.090 \pm 0.003$        | 2    |
| -183.9-179.0       | $1.82 \pm 0.03$          | $0.003 \pm 0.0007$        | $0.001 \pm 0.0004$      | $1.534 \pm 0.031$         | $0.110 \pm 0.002$      | $2.955 \pm 0.059$            | $0.510 \pm 0.020$        | $0.223 \pm 0.009$        | $0.155 \pm 0.007$        | 2    |
| +225.6-124.1       | $2.49 \pm 0.04$          | $0.006 \pm 0.0004$        | $0.002 \pm 0.0004$      | $1.747 \pm 0.035$         | $0.118 \pm 0.002$      | $2.874 \pm 0.057$            | $0.530 \pm 0.011$        | $0.252 \pm 0.005$        | $0.179 \pm 0.003$        | 2    |
| +117.9-235.0       | $2.57 \pm 0.06$          | $0.006 \pm 0.0008$        | $0.018 \pm 0.0017$      | $1.381 \pm 0.028$         | $0.115 \pm 0.003$      | $2.850 \pm 0.057$            | $0.420 \pm 0.015$        | $0.202 \pm 0.007$        | $0.146 \pm 0.005$        | 2    |
| -200.3-193.6       | $2.00 \pm 0.04$          | $0.006 \pm 0.0006$        | $0.010 \pm 0.0017$      | $1.966 \pm 0.039$         | $0.114 \pm 0.003$      | $2.697 \pm 0.054$            | $0.380 \pm 0.015$        | $0.230 \pm 0.009$        | $0.161 \pm 0.007$        | 2    |
| +96.7+266.9        | $2.50 \pm 0.05$          | $0.003 \pm 0.0006$        | $0.003 \pm 0.0004$      | $0.820 \pm 0.016$         | $0.091 \pm 0.002$      | $2.976 \pm 0.060$            | $0.710 \pm 0.026$        | $0.430 \pm 0.016$        | $0.300 \pm 0.012$        | 2    |
| +252.2-109.8       | $2.44 \pm 0.05$          | $0.007 \pm 0.0005$        | $0.004 \pm 0.0003$      | $1.623 \pm 0.032$         | $0.099 \pm 0.002$      | $2.794 \pm 0.056$            | $0.600 \pm 0.023$        | $0.370 \pm 0.014$        | $0.260 \pm 0.010$        | 2    |
| +254.6-107.2       | $1.60 \pm 0.05$          | $0.013 \pm 0.0003$        | $0.008 \pm 0.0010$      | $3.455 \pm 0.069$         | $0.144 \pm 0.004$      | $2.421 \pm 0.048$            | $0.370 \pm 0.020$        | $0.144 \pm 0.008$        | $0.120 \pm 0.006$        | 2    |
| +281.4-71.8        | $2.84 \pm 0.04$          | $0.007 \pm 0.0005$        | $0.011 \pm 0.0013$      | $1.621 \pm 0.032$         | $0.120 \pm 0.002$      | $2.885 \pm 0.058$            | $0.430 \pm 0.015$        | $0.191 \pm 0.007$        | $0.130 \pm 0.005$        | 2    |
| -243.0+159.6       | $2.81 \pm 0.05$          | $0.002 \pm 0.0004$        | $0.003 \pm 0.0005$      | $0.860 \pm 0.017$         | $0.089 \pm 0.003$      | $2.798 \pm 0.056$            | $0.670 \pm 0.027$        | $0.400 \pm 0.016$        | $0.280 \pm 0.012$        | 2    |
| -297.7+87.1        | $3.01 \pm 0.08$          | $0.005 \pm 0.0005$        | $0.010 \pm 0.0013$      | $1.197 \pm 0.024$         | $0.109 \pm 0.004$      | $2.811 \pm 0.056$            | $0.630 \pm 0.029$        | $0.370 \pm 0.018$        | $0.270 \pm 0.015$        | 2    |
| -309.4+56.9        | $2.45 \pm 0.06$          | $0.004 \pm 0.0005$        | $0.008 \pm 0.0013$      | $1.094 \pm 0.022$         | $0.073 \pm 0.003$      | $2.785 \pm 0.056$            | $0.640 \pm 0.028$        | $0.400 \pm 0.017$        | $0.280 \pm 0.013$        | 2    |
| +354.1+71.2        | $2.43 \pm 0.06$          | $0.019 \pm 0.0005$        | $0.001 \pm 0.0002$      | $3.520 \pm 0.070$         | $0.113 \pm 0.002$      | $2.887 \pm 0.058$            | $0.330 \pm 0.014$        | $0.218 \pm 0.009$        | $0.159 \pm 0.007$        | 2    |
| -164.9-333.9       | $2.28 \pm 0.05$          | $0.008 \pm 0.0002$        | $0.007 \pm 0.0011$      | $2.441 \pm 0.049$         | $0.125 \pm 0.002$      | $2.796 \pm 0.056$            | $0.290 \pm 0.013$        | $0.152 \pm 0.006$        | $0.108 \pm 0.004$        | 2    |
| +360.9+75.3        | $1.75 \pm 0.04$          | $0.017 \pm 0.0005$        | $0.002 \pm 0.0002$      | $3.537 \pm 0.071$         | $0.114 \pm 0.003$      | $2.884 \pm 0.058$            | $0.250 \pm 0.012$        | $0.168 \pm 0.008$        | $0.120 \pm 0.005$        | 2    |
| -377.9-64.9        | $2.87 \pm 0.07$          | $0.007 \pm 0.0003$        | $0.004 \pm 0.0003$      | $1.324 \pm 0.026$         | $0.095 \pm 0.002$      | $2.797 \pm 0.056$            | $0.570 \pm 0.026$        | $0.430 \pm 0.020$        | $0.300 \pm 0.017$        | 2    |
| -99.6-388.0        | $1.87 \pm 0.04$          | $0.020 \pm 0.0004$        | $0.005 \pm 0.0008$      | $3.882 \pm 0.078$         | $0.125 \pm 0.003$      | $2.663 \pm 0.053$            | $0.260 \pm 0.012$        | $0.150 \pm 0.007$        | $0.122 \pm 0.005$        | 2    |
| -397.4-71.7        | $2.41 \pm 0.05$          | $0.010 \pm 0.0016$        | $0.014 \pm 0.0031$      | $1.847 \pm 0.037$         | $0.065 \pm 0.003$      | $2.691 \pm 0.054$            | $0.440 \pm 0.016$        | $0.310 \pm 0.012$        | $0.212 \pm 0.008$        | 2    |
| -226.9-366.4       | $3.94 \pm 0.09$          | $0.008 \pm 0.0006$        | $0.006 \pm 0.0007$      | $1.514 \pm 0.030$         | $0.109 \pm 0.002$      | $2.752 \pm 0.055$            | $0.400 \pm 0.016$        | $0.290 \pm 0.012$        | $0.204 \pm 0.008$        | 2    |
| -405.5-157.7       | $3.14 \pm 0.06$          | $0.010 \pm 0.0009$        | $0.011 \pm 0.0020$      | $1.299 \pm 0.026$         | $0.114 \pm 0.005$      | $2.801 \pm 0.056$            | $0.510 \pm 0.021$        | $0.360 \pm 0.015$        | $0.250 \pm 0.011$        | 2    |
| -345.5+273.8       | $3.52 \pm 0.07$          | $0.039 \pm 0.0014$        | $0.008 \pm 0.0016$      | $4.450 \pm 0.100$         | $0.100 \pm 0.017$      | $2.810 \pm 0.056$            | $0.380 \pm 0.021$        | $0.550 \pm 0.030$        | $0.400 \pm 0.022$        | 2    |
| -410.3-206.3       | $3.18 \pm 0.08$          | $0.006 \pm 0.0007$        | $0.004 \pm 0.0010$      | $1.462 \pm 0.029$         | $0.111 \pm 0.003$      | $2.736 \pm 0.055$            | $0.460 \pm 0.021$        | $0.250 \pm 0.012$        | $0.173 \pm 0.008$        | 2    |
| -371.1-280.0       | $2.52 \pm 0.04$          | $0.016 \pm 0.0003$        | $0.004 \pm 0.0005$      | $2.665 \pm 0.053$         | $0.111 \pm 0.002$      | $2.865 \pm 0.057$            | $0.360 \pm 0.014$        | $0.270 \pm 0.010$        | $0.195 \pm 0.008$        | 2    |
| -368.3-285.6       | $2.36 \pm 0.04$          | $0.016 \pm 0.0003$        | $0.003 \pm 0.0004$      | $3.231 \pm 0.065$         | $0.118 \pm 0.002$      | $2.932 \pm 0.059$            | $0.350 \pm 0.013$        | $0.222 \pm 0.008$        | $0.166 \pm 0.006$        | 2    |
| -392.0-270.1       | $1.55 \pm 0.03$          | $0.021 \pm 0.0004$        | $0.008 \pm 0.0012$      | $3.721 \pm 0.074$         | $0.117 \pm 0.002$      | $2.879 \pm 0.058$            | $0.223 \pm 0.009$        | $0.138 \pm 0.005$        | $0.105 \pm 0.004$        | 2    |
| -481.4-0.5         | $3.17 \pm 0.07$          | $0.039 \pm 0.0007$        | $0.009 \pm 0.0013$      | $2.843 \pm 0.057$         | $0.091 \pm 0.002$      | $2.443 \pm 0.049$            | $0.190 \pm 0.011$        | $0.250 \pm 0.015$        | $0.180 \pm 0.011$        | 2    |
| -453.8-191.8       | $2.65 \pm 0.06$          | $0.036 \pm 0.0009$        | $0.003 \pm 0.0004$      | $3.747 \pm 0.075$         | $0.103 \pm 0.002$      | $2.620 \pm 0.052$            | $0.185 \pm 0.007$        | $0.204 \pm 0.008$        | $0.142 \pm 0.005$        | 2    |
| +331.9+401.0       | $2.05 \pm 0.04$          | $0.029 \pm 0.0015$        | $0.003 \pm 0.0009$      | $3.610 \pm 0.072$         | $0.098 \pm 0.006$      | $2.812 \pm 0.056$            | $0.157 \pm 0.007$        | $0.128 \pm 0.005$        | $0.089 \pm 0.004$        | 2    |
| +509.5+264.1       | $2.04 \pm 0.04$          | $0.033 \pm 0.0006$        | $0.004 \pm 0.0001$      | $4.173 \pm 0.083$         | $0.107 \pm 0.002$      | $2.832 \pm 0.057$            | $0.153 \pm 0.006$        | $0.157 \pm 0.005$        | $0.110 \pm 0.004$        | 2    |
| +266.0+534.1       | $2.85 \pm 0.07$          | $0.053 \pm 0.0024$        | $0.030 \pm 0.0054$      | $4.707 \pm 0.094$         | $0.099 \pm 0.014$      | $2.759 \pm 0.055$            | $0.190 \pm 0.012$        | $0.230 \pm 0.014$        | $0.160 \pm 0.010$        | 2    |
| +667.9+174.1       | $1.21 \pm 0.03$          | $0.088 \pm 0.0027$        | $0.006 \pm 0.0002$      | $6.740 \pm 0.100$         | $0.110 \pm 0.002$      | $2.714 \pm 0.054$            | $0.082 \pm 0.004$        | $0.105 \pm 0.004$        | $0.086 \pm 0.003$        | 2    |
| +1.0+885.8         | $2.56 \pm 0.06$          | $0.027 \pm 0.0007$        | $0.017 \pm 0.0008$      | $1.839 \pm 0.037$         | $0.097 \pm 0.007$      | $2.654 \pm 0.053$            | $0.118 \pm 0.006$        | $0.167 \pm 0.007$        | $0.115 \pm 0.005$        | 2    |
| +6.6+886.3         | $2.37 \pm 0.05$          | $0.039 \pm 0.0010$        | $0.009 \pm 0.0008$      | $2.688 \pm 0.054$         | $0.103 \pm 0.004$      | $2.682 \pm 0.054$            | $0.111 \pm 0.005$        | $0.163 \pm 0.006$        | $0.112 \pm 0.004$        | 2    |
| NGC5194+30.2+2.2   | $1.51 \pm 0.03$          | $0.013 \pm 0.0035$        | $0.020 \pm 0.0023$      | $0.341 \pm 0.006$         | $0.062 \pm 0.001$      | $2.960 \pm 0.059$            | $1.394 \pm 0.028$        | $0.266 \pm 0.005$        | $0.246 \pm 0.004$        | 3    |
| NGC 2403-38+51     | $2.03 \pm 0.13$          | $0.006 \pm 0.0010$        | $0.018 \pm 0.0110$      | $1.446 \pm 0.114$         | $0.110 \pm 0.008$      | $2.949 \pm 0.182$            | $0.478 \pm 0.028$        | $0.173 \pm 0.013$        | $0.135 \pm 0.011$        | 4    |
| NGC 2403+7+37      | $2.12 \pm 0.14$          | $0.007 \pm 0.0010$        | $0.006 \pm 0.0040$      | $1.429 \pm 0.121$         | $0.114 \pm 0.008$      | $2.952 \pm 0.187$            | $0.455 \pm 0.034$        | $0.178 \pm 0.013$        | $0.131 \pm 0.010$        | 4    |
| NGC 2403+119-28    | $2.47 \pm 0.10$          | $0.006 \pm 0.0020$        | $0.008 \pm 0.0050$      | $2.098 \pm 0.094$         | $0.123 \pm 0.006$      | $3.017 \pm 0.126$            | $0.437 \pm 0.027$        | $0.221 \pm 0.014$        | $0.167 \pm 0.014$        | 4    |
| NGC 2403-59+118    | $1.47 \pm 0.10$          | $0.005 \pm 0.0010$        | $0.020 \pm 0.0130$      | $2.230 \pm 0.184$         | $0.130 \pm 0.009$      | $2.939 \pm 0.182$            | $0.266 \pm 0.182$        | $0.147 \pm 0.012$        | $0.104 \pm 0.009$        | 4    |
| NGC 2403+96+30     | $2.38 \pm 0.05$          | $0.008 \pm 0.0010$        | $0.008 \pm 0.0040$      | $2.159 \pm 0.070$         | $0.120 \pm 0.005$      | $2.909 \pm 0.092$            | $0.361 \pm 0.020$        | $0.190 \pm 0.011$        | $0.151 \pm 0.009$        | 4    |
| NGC 2403+44+82     | $2.99 \pm 0.21$          | $0.005 \pm 0.0010$        | $0.014 \pm 0.0090$      | $1.383 \pm 0.117$         | $0.114 \pm 0.007$      | $3.067 \pm 0.189$            | $0.453 \pm 0.033$        | $0.204 \pm 0.014$        | $0.144 \pm 0.014$        | 4    |
| NGC 2403+166-140   | $1.99 \pm 0.14$          | $0.010 \pm 0.0010$        | $0.014 \pm 0.0090$      | $2.132 \pm 0.180$         | $0.129 \pm 0.010$      | $2.923 \pm 0.183$            | $0.260 \pm 0.021$        | $0.182 \pm 0.015$        | $0.133 \pm 0.011$        | 4    |
| NGC 2403-99-59     | $3.31 \pm 0.18$          | $0.007 \pm 0.0010$        | $0.003 \pm 0.0020$      | $1.699 \pm 0.108$         | $0.109 \pm 0.005$      | $2.992 \pm 0.145$            | $0.482 \pm 0.034$        | $0.345 \pm 0.017$        | $0.247 \pm 0.013$        | 4    |
| NGC 2403-196+58    | $1.90 \pm 0.07$          | $0.007 \pm 0.0010$        | $0.006 \pm 0.0040$      | $2.160 \pm 0.084$         | $0.117 \pm 0.005$      | $2.902 \pm 0.102$            | $0.315 \pm 0.023$        | $0.150 \pm 0.008$        | $0.110 \pm 0.006$        | 4    |
| NGC 2403-22-162    | $3.63 \pm 0.15$          | $0.009 \pm 0.0020$        | $0.020 \pm 0.0160$      | $1.792 \pm 0.083$         | $0.116 \pm 0.007$      | $2.881 \pm 0.108$            | $0.394 \pm 0.024$        | $0.331 \pm 0.019$        | $0.241 \pm 0.016$        | 4    |
| NGC 2403+160-251   | $2.83 \pm 0.20$          | $0.026 \pm 0.0030$        | $0.015 \pm 0.0030$      | $3.160 \pm 0.261$         | $0.109 \pm 0.007$      | $2.976 \pm 0.178$            | $0.204 \pm 0.017$        | $0.201 \pm 0.014$        | $0.143 \pm 0.011$        | 4    |
| SDSS J1455         | $1.11 \pm 0.01$          | $0.102 \pm 0.0030$        | $0.007 \pm 0.0009$      | $6.135 \pm 0.033$         | $0.114 \pm 0.003$      | $2.775 \pm 0.018$            | $0.079 \pm 0.002$        | $0.100 \pm 0.002$        | $0.008 \pm 0.002$        | 5    |
| SDSS J1657         | $0.84 \pm 0.02$          | $0.085 \pm 0.2690$        | $0.032 \pm 0.0043$      |                           |                        |                              |                          |                          |                          |      |

**Table 2** – *continued*

| ObjID           | [O II]<br>λ3727 | [O III]<br>λ4363 | He II<br>λ4686 | [O III]<br>λ5007 | He I<br>λ5876 | Hα<br>λ6563   | [N II]<br>λ6583 | [S II]<br>λ6716 | [S II]<br>λ6731 | Ref. |
|-----------------|-----------------|------------------|----------------|------------------|---------------|---------------|-----------------|-----------------|-----------------|------|
| J0118+3512      | 0.84 ± 0.01     | 0.064 ± 0.0016   | 0.032 ± 0.0020 | 3.062 ± 0.005    | 0.114 ± 0.004 | 3.349 ± 0.005 | 0.039 ± 0.002   | 0.110 ± 0.003   | 0.085 ± 0.002   | 7    |
| J1322+5425      | 0.43 ± 0.01     | 0.075 ± 0.0013   | 0.010 ± 0.0008 | 2.938 ± 0.005    | 0.084 ± 0.001 | 2.670 ± 0.008 | 0.015 ± 0.001   | 0.044 ± 0.001   | 0.031 ± 0.001   | 7    |
| 0723+692A       | 0.61 ± 0.01     | 0.143 ± 0.0010   | 0.009 ± 0.0010 | 6.591 ± 0.007    | 0.106 ± 0.001 | 2.788 ± 0.004 | 0.024 ± 0.001   | 0.049 ± 0.001   | 0.037 ± 0.001   | 8    |
| 0723+692B       | 1.57 ± 0.01     | 0.082 ± 0.0040   | 0.014 ± 0.0040 | 4.303 ± 0.031    | 0.102 ± 0.003 | 2.794 ± 0.023 | 0.055 ± 0.003   | 0.114 ± 0.003   | 0.082 ± 0.003   | 8    |
| 0749+568        | 1.66 ± 0.04     | 0.098 ± 0.0110   | 0.018 ± 0.0080 | 4.880 ± 0.099    | 0.111 ± 0.009 | 2.797 ± 0.064 | 0.076 ± 0.007   | 0.178 ± 0.011   | 0.114 ± 0.008   | 8    |
| 0907+543        | 0.94 ± 0.02     | 0.121 ± 0.0080   | 0.029 ± 0.0010 | 6.838 ± 0.109    | 0.106 ± 0.007 | 2.803 ± 0.053 | 0.033 ± 0.005   | 0.065 ± 0.006   | 0.050 ± 0.005   | 8    |
| 0917+527        | 1.88 ± 0.01     | 0.092 ± 0.0040   | 0.023 ± 0.0030 | 4.680 ± 0.003    | 0.103 ± 0.003 | 2.797 ± 0.019 | 0.058 ± 0.002   | 0.164 ± 0.003   | 0.114 ± 0.003   | 8    |
| 0926+606        | 1.78 ± 0.01     | 0.083 ± 0.0030   | 0.016 ± 0.0020 | 4.772 ± 0.026    | 0.104 ± 0.003 | 2.804 ± 0.017 | 0.083 ± 0.002   | 0.182 ± 0.003   | 0.146 ± 0.003   | 8    |
| 0930+554        | 0.40 ± 0.00     | 0.060 ± 0.0020   | 0.027 ± 0.0020 | 1.961 ± 0.008    | 0.081 ± 0.001 | 2.930 ± 0.012 | 0.013 ± 0.001   | 0.031 ± 0.001   | 0.023 ± 0.001   | 8    |
| 1030+583        | 0.96 ± 0.00     | 0.104 ± 0.0020   | 0.024 ± 0.0020 | 5.028 ± 0.021    | 0.099 ± 0.002 | 2.786 ± 0.013 | 0.031 ± 0.001   | 0.096 ± 0.002   | 0.067 ± 0.001   | 8    |
| 1116+583B       | 0.58 ± 0.02     | 0.117 ± 0.0120   | 0.025 ± 0.0100 | 4.842 ± 0.111    | 0.101 ± 0.009 | 2.777 ± 0.072 | 0.027 ± 0.007   | 0.068 ± 0.007   | 0.065 ± 0.008   | 8    |
| 1205+557        | 2.13 ± 0.02     | 0.083 ± 0.0070   | 0.018 ± 0.0070 | 3.714 ± 0.043    | 0.099 ± 0.005 | 2.714 ± 0.035 | 0.083 ± 0.005   | 0.183 ± 0.007   | 0.130 ± 0.007   | 8    |
| 1222+614        | 1.16 ± 0.00     | 0.102 ± 0.0020   | 0.017 ± 0.0020 | 5.955 ± 0.022    | 0.100 ± 0.002 | 2.761 ± 0.012 | 0.038 ± 0.001   | 0.090 ± 0.001   | 0.065 ± 0.001   | 8    |
| 1223+487        | 0.71 ± 0.00     | 0.127 ± 0.0010   | 0.012 ± 0.0010 | 5.543 ± 0.008    | 0.102 ± 0.001 | 2.777 ± 0.005 | 0.029 ± 0.001   | 0.061 ± 0.001   | 0.045 ± 0.001   | 8    |
| 1256+351        | 1.10 ± 0.00     | 0.089 ± 0.0010   | 0.011 ± 0.0010 | 5.802 ± 0.009    | 0.106 ± 0.001 | 2.819 ± 0.005 | 0.046 ± 0.001   | 0.095 ± 0.001   | 0.071 ± 0.001   | 8    |
| 1319+579A       | 1.02 ± 0.00     | 0.093 ± 0.0020   | 0.008 ± 0.0010 | 6.700 ± 0.028    | 0.113 ± 0.002 | 2.827 ± 0.014 | 0.050 ± 0.001   | 0.101 ± 0.001   | 0.079 ± 0.001   | 8    |
| 1319+579B4      | 2.44 ± 0.06     | 0.050 ± 0.0150   | 0.038 ± 0.0230 | 3.282 ± 0.073    | 0.094 ± 0.011 | 2.815 ± 0.069 | 0.161 ± 0.013   | 0.374 ± 0.017   | 0.273 ± 0.015   | 8    |
| 1319+579C       | 2.38 ± 0.01     | 0.033 ± 0.0030   | 0.014 ± 0.0040 | 3.520 ± 0.025    | 0.106 ± 0.003 | 2.852 ± 0.022 | 0.150 ± 0.003   | 0.269 ± 0.004   | 0.192 ± 0.004   | 8    |
| 1358+576        | 1.67 ± 0.01     | 0.089 ± 0.0030   | 0.012 ± 0.0020 | 4.874 ± 0.026    | 0.112 ± 0.002 | 2.803 ± 0.017 | 0.108 ± 0.003   | 0.153 ± 0.003   | 0.111 ± 0.002   | 8    |
| 1441+294        | 1.50 ± 0.03     | 0.069 ± 0.0100   | 0.019 ± 0.0130 | 4.927 ± 0.099    | 0.116 ± 0.009 | 2.824 ± 0.064 | 0.083 ± 0.008   | 0.204 ± 0.011   | 0.138 ± 0.010   | 8    |
| 1533+574B       | 2.03 ± 0.01     | 0.065 ± 0.0040   | 0.011 ± 0.0030 | 5.329 ± 0.035    | 0.108 ± 0.003 | 2.849 ± 0.021 | 0.087 ± 0.002   | 0.167 ± 0.003   | 0.117 ± 0.002   | 8    |
| Pox 105         | 1.25 ± 0.00     | 0.113 ± 0.0000   | 0.015 ± 0.0000 | 5.540 ± 0.000    | 0.096 ± 0.000 | 2.800 ± 0.000 | 0.064 ± 0.000   | 0.084 ± 0.000   | 0.058 ± 0.000   | 9    |
| Pox 120         | 0.79 ± 0.00     | 0.135 ± 0.0000   | 0.016 ± 0.0000 | 6.290 ± 0.000    | 0.101 ± 0.000 | 2.780 ± 0.000 | 0.048 ± 0.000   | 0.074 ± 0.000   | 0.047 ± 0.000   | 9    |
| Pox 139         | 1.47 ± 0.00     | 0.096 ± 0.0000   | 0.023 ± 0.0000 | 5.710 ± 0.000    | 0.101 ± 0.000 | 2.800 ± 0.000 | 0.019 ± 0.000   | 0.117 ± 0.000   | 0.077 ± 0.000   | 9    |
| UM 160 A        | 1.33 ± 0.03     | 0.085 ± 0.0068   | 0.004 ± 0.0012 | 5.240 ± 0.524    | 0.100 ± 0.006 | 2.800 ± 0.028 | 0.048 ± 0.003   | 0.111 ± 0.006   | 0.085 ± 0.005   | 10   |
| UM 160 B        | 1.48 ± 0.03     | 0.066 ± 0.0066   | 0.023 ± 0.0035 | 4.740 ± 0.474    | 0.096 ± 0.007 | 2.780 ± 0.055 | 0.071 ± 0.005   | 0.152 ± 0.009   | 0.102 ± 0.006   | 10   |
| UM 420 B        | 2.57 ± 0.05     | 0.061 ± 0.0073   | 0.023 ± 0.0058 | 4.260 ± 4.260    | 0.113 ± 0.006 | 2.810 ± 0.056 | 0.245 ± 0.009   | 0.250 ± 0.010   | 0.196 ± 0.009   | 10   |
| TOL 0513-393    | 0.53 ± 0.01     | 0.155 ± 0.0061   | 0.010 ± 0.0015 | 7.760 ± 0.776    | 0.128 ± 0.005 | 2.820 ± 0.028 | 0.045 ± 0.002   | 0.062 ± 0.003   | 0.053 ± 0.003   | 10   |
| TOL 2146-391-C  | 0.61 ± 0.00     | 0.127 ± 0.0026   | 0.017 ± 0.0009 | 5.941 ± 0.059    | 0.109 ± 0.002 | 2.820 ± 0.028 | 0.027 ± 0.001   | 0.066 ± 0.001   | 0.052 ± 0.001   | 11   |
| TOL 2146-391-E  | 0.64 ± 0.00     | 0.127 ± 0.0038   | 0.018 ± 0.0013 | 5.897 ± 0.058    | 0.106 ± 0.003 | 2.770 ± 0.027 | 0.028 ± 0.001   | 0.067 ± 0.002   | 0.052 ± 0.002   | 11   |
| TOL 0357-3915-C | 0.81 ± 0.01     | 0.123 ± 0.0043   | 0.019 ± 0.0016 | 6.549 ± 0.065    | 0.111 ± 0.003 | 2.800 ± 0.028 | 0.055 ± 0.002   | 0.065 ± 0.002   | 0.049 ± 0.002   | 11   |
| TOL 0357-3915-E | 0.85 ± 0.01     | 0.116 ± 0.0058   | 0.013 ± 0.0019 | 6.300 ± 0.063    | 0.107 ± 0.005 | 2.810 ± 0.042 | 0.058 ± 0.004   | 0.070 ± 0.004   | 0.054 ± 0.003   | 11   |
| NGC 346         | 0.80 ± 0.01     | 0.070 ± 0.0014   | 0.002 ± 0.0002 | 5.220 ± 0.044    | 0.108 ± 0.001 | 2.820 ± 0.023 | 0.038 ± 0.001   | 0.074 ± 0.001   | 0.053 ± 0.001   | 12   |

References: (1) [Berg et al. \(2020\)](#), (2) [Croxall et al. \(2016\)](#), (3) [Croxall et al. \(2015\)](#), (4) [Rogers et al. \(2021\)](#), (5) [Hägele et al. \(2008\)](#), (6) [Annibali et al. \(2019\)](#), (7) [Hsyu et al. \(2020\)](#), (8) [Izotov et al. \(1997\)](#), (9) [Kunth & Sargent \(1983\)](#), (10) [Valerdi et al. \(2021a\)](#), (11) [Peimbert et al. \(2012\)](#), (12) [Valerdi et al. \(2019\)](#).

**Table 3.** Chemical abundances for the Seyfert 2 and Star-forming samples.

| ObjID               | <sup>a</sup> O <sup>+</sup> /H <sup>+</sup> | <sup>a</sup> O <sup>2+</sup> /H <sup>+</sup> | ICF(O) | <sup>b</sup> O/H | <sup>a</sup> He <sup>+</sup> /H <sup>+</sup> | <sup>a</sup> He <sup>2+</sup> /H <sup>+</sup> | ICF(He) | <sup>b</sup> He/H |
|---------------------|---------------------------------------------|----------------------------------------------|--------|------------------|----------------------------------------------|-----------------------------------------------|---------|-------------------|
| Seyfert 2           |                                             |                                              |        |                  |                                              |                                               |         |                   |
| J013957.81-004504.2 | 8.564 ± 0.041                               | 8.324 ± 0.006                                | 1.3    | 8.901 ± 0.129    | 10.626 ± 0.144                               | 10.120 ± 0.027                                | 1.5     | 10.918 ± 0.075    |
| J033923.14-054841.5 | 8.522 ± 0.033                               | 8.185 ± 0.001                                | 1.3    | 8.815 ± 0.026    | 10.805 ± 0.048                               | 10.339 ± 0.017                                | 1.5     | 11.114 ± 0.023    |
| J074257.23+333217.9 | 8.366 ± 0.021                               | 7.700 ± 0.001                                | 1.1    | 8.491 ± 0.018    | 10.852 ± 0.044                               | 9.833 ± 0.055                                 | 1.3     | 11.007 ± 0.031    |
| J093509.12+002557.4 | 8.438 ± 0.017                               | 7.647 ± 0.002                                | 1.2    | 8.580 ± 0.019    | 10.700 ± 0.061                               | 9.981 ± 0.039                                 | 1.4     | 10.931 ± 0.037    |
| J095759.45+022810.5 | 8.631 ± 0.033                               | 8.208 ± 0.004                                | 1.4    | 8.912 ± 0.029    | 10.757 ± 0.056                               | 10.338 ± 0.017                                | 1.6     | 11.092 ± 0.026    |
| J100602.50+071131.8 | 8.825 ± 0.003                               | 7.904 ± 0.001                                | 1.4    | 9.012 ± 0.014    | 10.807 ± 0.046                               | 10.370 ± 0.015                                | 1.6     | 11.158 ± 0.021    |
| J100921.26+013334.5 | 8.599 ± 0.036                               | 8.377 ± 0.002                                | 1.3    | 8.931 ± 0.025    | 10.888 ± 0.038                               | 10.416 ± 0.014                                | 1.5     | 11.191 ± 0.020    |
| J101754.72-002811.9 | 8.404 ± 0.026                               | 7.801 ± 0.002                                | 1.3    | 8.615 ± 0.022    | 10.895 ± 0.040                               | 10.370 ± 0.016                                | 1.5     | 11.183 ± 0.021    |
| J102039.81+642435.8 | 8.681 ± 0.099                               | 7.979 ± 0.003                                | 1.4    | 8.914 ± 0.083    | 10.699 ± 0.061                               | 10.334 ± 0.017                                | 1.6     | 11.066 ± 0.027    |
| J112850.39+021016.2 | 8.734 ± 0.029                               | 7.456 ± 0.010                                | 1.2    | 8.850 ± 0.033    | 11.049 ± 0.082                               | 10.409 ± 0.031                                | 1.4     | 11.298 ± 0.046    |
| Mrk176              | 8.564 ± 0.008                               | 8.115 ± 0.004                                | 1.5    | 8.882 ± 0.012    | 10.849 ± 0.028                               | 10.576 ± 0.007                                | 1.6     | 11.238 ± 0.012    |
| 3C33                | 8.740 ± 0.006                               | 7.998 ± 0.005                                | 1.3    | 8.939 ± 0.010    | 10.835 ± 0.028                               | 10.361 ± 0.011                                | 1.5     | 11.144 ± 0.014    |
| Mrk3                | 8.549 ± 0.008                               | 8.144 ± 0.005                                | 1.3    | 8.796 ± 0.010    | 10.771 ± 0.032                               | 10.194 ± 0.015                                | 1.5     | 11.035 ± 0.018    |
| Mrk573              | 8.492 ± 0.010                               | 8.244 ± 0.005                                | 1.4    | 8.844 ± 0.011    | 10.848 ± 0.028                               | 10.488 ± 0.008                                | 1.6     | 11.196 ± 0.012    |
| Mrk78               | 8.782 ± 0.006                               | 8.351 ± 0.005                                | 1.4    | 9.071 ± 0.010    | 10.851 ± 0.027                               | 10.469 ± 0.008                                | 1.6     | 11.204 ± 0.012    |
| Mrk348              | 8.654 ± 0.007                               | 8.032 ± 0.005                                | 1.2    | 8.836 ± 0.007    | 10.927 ± 0.023                               | 10.286 ± 0.013                                | 1.4     | 11.173 ± 0.013    |
| Mrk34               | 8.590 ± 0.008                               | 8.280 ± 0.005                                | 1.3    | 8.871 ± 0.008    | 10.929 ± 0.022                               | 10.376 ± 0.010                                | 1.5     | 11.202 ± 0.012    |
| Mrk1                | 8.450 ± 0.010                               | 8.073 ± 0.005                                | 1.4    | 8.740 ± 0.011    | 10.849 ± 0.027                               | 10.416 ± 0.009                                | 1.5     | 11.167 ± 0.013    |
| Mrk270              | 8.844 ± 0.005                               | 7.707 ± 0.007                                | 1.2    | 8.936 ± 0.006    | 11.123 ± 0.014                               | 10.298 ± 0.013                                | 1.4     | 11.320 ± 0.009    |
| IIIZw55             | 8.655 ± 0.009                               | 7.543 ± 0.009                                | 1.3    | 8.799 ± 0.011    | 10.799 ± 0.031                               | 10.260 ± 0.014                                | 1.5     | 11.086 ± 0.016    |
| 3C452               | 8.771 ± 0.006                               | 7.710 ± 0.009                                | 1.0    | 8.825 ± 0.006    | 11.088 ± 0.017                               | 9.719 ± 0.048                                 | 1.4     | 11.242 ± 0.012    |
| Mrk198              | 8.446 ± 0.012                               | 7.722 ± 0.011                                | 1.1    | 8.556 ± 0.011    | 10.898 ± 0.025                               | 9.815 ± 0.037                                 | 1.3     | 11.049 ± 0.018    |
| Mrk268              | 8.930 ± 0.007                               | 7.241 ± 0.012                                | 1.1    | 8.983 ± 0.008    | 10.824 ± 0.027                               | 9.861 ± 0.033                                 | 1.3     | 10.991 ± 0.019    |
| NGC2110             | 8.796 ± 0.006                               | 7.387 ± 0.012                                | 1.1    | 8.836 ± 0.007    | 10.992 ± 0.018                               | 9.736 ± 0.043                                 | 1.3     | 11.131 ± 0.013    |
| ESO138G1            | 8.572 ± 0.015                               | 7.596 ± 0.018                                | 1.4    | 8.752 ± 0.027    | 10.885 ± 0.079                               | 10.439 ± 0.029                                | 1.5     | 11.198 ± 0.037    |
| NGC3081             | 8.344 ± 0.013                               | 8.139 ± 0.005                                | 1.5    | 8.743 ± 0.013    | 10.848 ± 0.027                               | 10.582 ± 0.006                                | 1.6     | 11.234 ± 0.012    |
| NGC4388             | 8.481 ± 0.011                               | 8.173 ± 0.006                                | 1.2    | 8.745 ± 0.009    | 10.894 ± 0.025                               | 10.254 ± 0.013                                | 1.4     | 11.137 ± 0.015    |
| NGC5135             | 8.344 ± 0.014                               | 7.546 ± 0.014                                | 1.3    | 8.505 ± 0.014    | 10.850 ± 0.028                               | 10.247 ± 0.014                                | 1.5     | 11.113 ± 0.015    |
| NGC5728             | 8.813 ± 0.009                               | 7.634 ± 0.006                                | 1.3    | 8.950 ± 0.009    | 11.020 ± 0.019                               | 10.471 ± 0.009                                | 1.5     | 11.296 ± 0.010    |
| Akn347a             | 8.485 ± 0.011                               | 8.025 ± 0.004                                | 1.3    | 8.744 ± 0.010    | 10.927 ± 0.023                               | 10.467 ± 0.008                                | 1.5     | 11.232 ± 0.012    |
| UM16a               | 8.493 ± 0.010                               | 8.202 ± 0.004                                | 1.3    | 8.770 ± 0.007    | 11.103 ± 0.015                               | 10.505 ± 0.008                                | 1.4     | 11.360 ± 0.008    |
| Mrk612a             | 8.334 ± 0.016                               | 7.925 ± 0.007                                | 1.1    | 8.517 ± 0.012    | 11.159 ± 0.014                               | 10.145 ± 0.018                                | 1.4     | 11.338 ± 0.010    |
| Mrk573              | 8.382 ± 0.011                               | 8.213 ± 0.002                                | 1.4    | 8.751 ± 0.018    | 10.802 ± 0.050                               | 10.390 ± 0.031                                | 1.5     | 11.126 ± 0.024    |
| NGC2992             | 8.743 ± 0.014                               | 7.375 ± 0.008                                | 1.1    | 8.808 ± 0.020    | 10.752 ± 0.055                               | 9.813 ± 0.125                                 | 1.3     | 10.922 ± 0.037    |
| IC2560              | 8.245 ± 0.005                               | 8.185 ± 0.004                                | 1.3    | 8.638 ± 0.028    | 10.847 ± 0.045                               | 10.344 ± 0.105                                | 1.5     | 11.138 ± 0.028    |
| NGC5664             | 8.538 ± 0.012                               | 8.006 ± 0.010                                | 1.2    | 8.713 ± 0.014    | 10.801 ± 0.049                               | 9.991 ± 0.037                                 | 1.4     | 11.002 ± 0.033    |
| NGC5728             | 8.482 ± 0.006                               | 8.217 ± 0.005                                | 1.2    | 8.768 ± 0.013    | 10.762 ± 0.054                               | 10.157 ± 0.025                                | 1.4     | 11.017 ± 0.030    |
| ESO339-G11          | 8.480 ± 0.023                               | 7.957 ± 0.003                                | 1.2    | 8.666 ± 0.021    | 10.802 ± 0.050                               | 10.050 ± 0.035                                | 1.4     | 11.014 ± 0.030    |
| NGC6890             | 8.246 ± 0.010                               | 8.025 ± 0.001                                | 1.4    | 8.605 ± 0.209    | 10.626 ± 0.146                               | 10.219 ± 0.023                                | 1.5     | 10.951 ± 0.068    |
| IC5063              | 8.486 ± 0.009                               | 8.038 ± 0.004                                | 1.1    | 8.675 ± 0.011    | 10.751 ± 0.055                               | 9.886 ± 0.049                                 | 1.3     | 10.936 ± 0.035    |
| NGC7130             | 8.250 ± 0.008                               | 7.690 ± 0.005                                | 1.4    | 8.497 ± 0.031    | 10.546 ± 0.090                               | 10.115 ± 0.030                                | 1.6     | 10.874 ± 0.042    |
| NGC7582             | 8.139 ± 0.007                               | 7.517 ± 0.021                                | 1.1    | 8.285 ± 0.026    | 10.848 ± 0.045                               | 9.967 ± 0.206                                 | 1.3     | 11.030 ± 0.034    |
| NGC7590             | 8.650 ± 0.024                               | 7.717 ± 0.020                                | 1.1    | 8.763 ± 0.054    | 10.633 ± 0.141                               | 9.780 ± 0.180                                 | 1.4     | 10.830 ± 0.092    |
| CygnusA             | 8.728 ± 0.006                               | 8.206 ± 0.005                                | 1.5    | 9.010 ± 0.014    | 10.696 ± 0.040                               | 10.365 ± 0.011                                | 1.6     | 11.072 ± 0.017    |
| Mrk1157a            | 8.889 ± 0.004                               | 7.882 ± 0.006                                | 1.3    | 9.027 ± 0.006    | 10.927 ± 0.023                               | 10.327 ± 0.011                                | 1.5     | 11.204 ± 0.012    |
| ESO428-G14          | 8.351 ± 0.014                               | 8.028 ± 0.006                                | 1.1    | 8.580 ± 0.010    | 11.051 ± 0.018                               | 10.220 ± 0.015                                | 1.4     | 11.253 ± 0.011    |
| ESO137-G34          | 8.554 ± 0.010                               | 8.242 ± 0.007                                | 1.2    | 8.817 ± 0.009    | 10.886 ± 0.025                               | 10.249 ± 0.013                                | 1.4     | 11.132 ± 0.014    |
| IC3639              | 8.111 ± 0.024                               | 7.882 ± 0.011                                | 1.1    | 8.372 ± 0.017    | 10.847 ± 0.028                               | 10.011 ± 0.024                                | 1.4     | 11.045 ± 0.018    |
| IC4777              | 8.461 ± 0.012                               | 8.103 ± 0.009                                | 1.2    | 8.699 ± 0.010    | 10.849 ± 0.026                               | 10.154 ± 0.016                                | 1.4     | 11.077 ± 0.016    |
| IC4995              | 8.130 ± 0.022                               | 8.173 ± 0.005                                | 1.4    | 8.604 ± 0.015    | 10.750 ± 0.033                               | 10.366 ± 0.010                                | 1.5     | 11.087 ± 0.016    |
| IRAS11215-2806      | 8.409 ± 0.011                               | 7.892 ± 0.008                                | 1.3    | 8.631 ± 0.011    | 10.891 ± 0.024                               | 10.336 ± 0.011                                | 1.5     | 11.166 ± 0.013    |
| MCG-01-24-012       | 8.542 ± 0.009                               | 7.975 ± 0.007                                | 1.3    | 8.750 ± 0.010    | 10.804 ± 0.031                               | 10.238 ± 0.014                                | 1.5     | 11.077 ± 0.016    |
| MCG-02-51-008       | 8.486 ± 0.012                               | 7.900 ± 0.014                                | 1.1    | 8.631 ± 0.011    | 10.856 ± 0.026                               | 9.879 ± 0.029                                 | 1.3     | 11.018 ± 0.018    |
| NGC1125             | 8.542 ± 0.009                               | 7.968 ± 0.010                                | 1.2    | 8.713 ± 0.009    | 10.850 ± 0.027                               | 10.076 ± 0.020                                | 1.4     | 11.059 ± 0.017    |
| NGC1194             | 8.306 ± 0.018                               | 8.023 ± 0.010                                | 1.1    | 8.515 ± 0.012    | 10.896 ± 0.024                               | 9.703 ± 0.045                                 | 1.4     | 11.063 ± 0.016    |

Table 3 – *continued*

| ObjID                | <sup>a</sup> O <sup>+</sup> /H <sup>+</sup> | <sup>a</sup> O <sup>2+</sup> /H <sup>+</sup> | ICF(O) | <sup>b</sup> O/H | <sup>a</sup> He <sup>+</sup> /H <sup>+</sup> | <sup>a</sup> He <sup>2+</sup> /H <sup>+</sup> | ICF(He) | <sup>b</sup> He/H |
|----------------------|---------------------------------------------|----------------------------------------------|--------|------------------|----------------------------------------------|-----------------------------------------------|---------|-------------------|
| NGC1320              | 8.036 ± 0.029                               | 8.160 ± 0.006                                | 1.4    | 8.552 ± 0.015    | 10.890 ± 0.024                               | 10.498 ± 0.007                                | 1.5     | 11.226 ± 0.011    |
| NGC3281              | 8.407 ± 0.014                               | 8.163 ± 0.008                                | 1.3    | 8.722 ± 0.012    | 10.892 ± 0.025                               | 10.387 ± 0.009                                | 1.5     | 11.182 ± 0.013    |
| NGC3393              | 8.446 ± 0.013                               | 8.294 ± 0.006                                | 1.3    | 8.777 ± 0.010    | 10.926 ± 0.023                               | 10.338 ± 0.011                                | 1.4     | 11.186 ± 0.013    |
| NGC4939              | 8.377 ± 0.014                               | 8.265 ± 0.006                                | 1.3    | 8.740 ± 0.010    | 10.926 ± 0.023                               | 10.403 ± 0.009                                | 1.5     | 11.208 ± 0.012    |
| NGC4968              | 8.227 ± 0.017                               | 7.911 ± 0.006                                | 1.3    | 8.506 ± 0.013    | 10.927 ± 0.022                               | 10.375 ± 0.010                                | 1.5     | 11.200 ± 0.012    |
| NGC5427              | 8.346 ± 0.014                               | 8.014 ± 0.008                                | 1.2    | 8.576 ± 0.010    | 10.993 ± 0.019                               | 10.189 ± 0.015                                | 1.4     | 11.194 ± 0.012    |
| NGC5643              | 8.485 ± 0.011                               | 8.124 ± 0.007                                | 1.7    | 8.886 ± 0.031    | 10.449 ± 0.067                               | 10.309 ± 0.011                                | 1.7     | 10.919 ± 0.022    |
| NGC5990              | 8.721 ± 0.015                               | 8.059 ± 0.027                                | 1.1    | 8.844 ± 0.014    | 10.794 ± 0.029                               | 9.749 ± 0.038                                 | 1.3     | 10.947 ± 0.022    |
| NGC7682              | 8.465 ± 0.010                               | 8.059 ± 0.006                                | 1.3    | 8.709 ± 0.010    | 10.802 ± 0.030                               | 10.214 ± 0.014                                | 1.5     | 11.064 ± 0.016    |
| PKS1306-241          | 8.659 ± 0.008                               | 7.908 ± 0.016                                | 1.1    | 8.755 ± 0.009    | 10.849 ± 0.027                               | 9.620 ± 0.055                                 | 1.3     | 10.987 ± 0.020    |
| Star-forming regions |                                             |                                              |        |                  |                                              |                                               |         |                   |
| NGC3184-72.8-91.3    | 8.258±0.002                                 | 7.241±0.002                                  | 1.015  | 8.304±0.002      | 10.780±0.005                                 | 8.947±0.039                                   | 1.253   | 10.885±0.007      |
| NGC3184-14.9-139.6   | 8.350±0.007                                 | 7.784±0.005                                  | 1.033  | 8.469±0.006      | 10.771±0.033                                 | 9.286±0.056                                   | 1.127   | 10.837±0.031      |
| NGC3184-80.0-148.2   | 8.494±0.005                                 | 7.639±0.009                                  | 1.007  | 8.554±0.004      | 10.901±0.040                                 | 8.749±0.063                                   | 1.221   | 10.991±0.039      |
| NGC3184-93.3-142.3   | 8.414±0.006                                 | 8.032±0.007                                  | 1.001  | 8.565±0.005      | 10.968±0.034                                 | 7.892±0.452                                   | 1.094   | 11.007±0.035      |
| NGC3184-172.5-30.2   | 8.304±0.002                                 | 7.992±0.003                                  | 1.003  | 8.478±0.002      | 10.958±0.014                                 | 8.387±0.146                                   | 1.078   | 10.992±0.014      |
| 164.6-9.9            | 8.626±0.009                                 | 8.052±0.009                                  | 1.009  | 8.732±0.007      | 10.910±0.008                                 | 8.845±0.049                                   | 1.142   | 10.972±0.010      |
| 17.3-235.4           | 8.532±0.005                                 | 7.907±0.009                                  | 1.001  | 8.625±0.005      | 10.896±0.008                                 | 7.899±0.175                                   | 1.161   | 10.961±0.010      |
| 189.2-136.3          | 8.483±0.008                                 | 8.423±0.009                                  | 1.008  | 8.759±0.006      | 10.993±0.006                                 | 8.894±0.074                                   | 1.039   | 11.014±0.007      |
| 183.9-179.0          | 8.621±0.007                                 | 8.268±0.009                                  | 1.001  | 8.781±0.006      | 10.886±0.008                                 | 7.891±0.173                                   | 1.088   | 10.923±0.009      |
| 225.6-124.1          | 8.539±0.007                                 | 8.077±0.009                                  | 1.002  | 8.669±0.006      | 10.926±0.007                                 | 8.199±0.087                                   | 1.115   | 10.974±0.009      |
| 117.9-235.0          | 8.457±0.010                                 | 7.873±0.009                                  | 1.017  | 8.565±0.008      | 10.920±0.011                                 | 9.158±0.041                                   | 1.140   | 10.984±0.014      |
| 200.3-193.6          | 8.489±0.009                                 | 8.182±0.009                                  | 1.010  | 8.667±0.006      | 10.909±0.012                                 | 8.897±0.074                                   | 1.075   | 10.945±0.012      |
| 96.7-266.9           | 8.518±0.009                                 | 7.726±0.009                                  | 1.004  | 8.585±0.008      | 10.814±0.010                                 | 8.374±0.060                                   | 1.209   | 10.898±0.014      |
| 252.2-109.8          | 8.440±0.009                                 | 7.946±0.009                                  | 1.004  | 8.563±0.007      | 10.854±0.009                                 | 8.503±0.033                                   | 1.122   | 10.906±0.011      |
| 254.6-107.2          | 8.288±0.013                                 | 8.330±0.009                                  | 1.006  | 8.613±0.008      | 11.010±0.012                                 | 8.804±0.054                                   | 1.031   | 11.027±0.012      |
| 281.4-71.8           | 8.510±0.006                                 | 7.946±0.009                                  | 1.010  | 8.619±0.005      | 10.938±0.007                                 | 8.944±0.051                                   | 1.137   | 10.998±0.009      |
| 243.0-159.6          | 8.744±0.008                                 | 7.945±0.009                                  | 1.004  | 8.809±0.007      | 10.797±0.015                                 | 8.370±0.073                                   | 1.209   | 10.881±0.017      |
| 297.7-87.1           | 8.541±0.012                                 | 7.827±0.009                                  | 1.010  | 8.622±0.010      | 10.896±0.016                                 | 8.903±0.057                                   | 1.180   | 10.972±0.019      |
| 309.4-56.9           | 8.509±0.011                                 | 7.849±0.009                                  | 1.012  | 8.600±0.009      | 10.718±0.018                                 | 8.801±0.071                                   | 1.163   | 10.789±0.020      |
| 354.1-71.2           | 8.345±0.011                                 | 8.178±0.009                                  | 1.001  | 8.571±0.007      | 10.916±0.008                                 | 7.905±0.088                                   | 1.053   | 10.939±0.008      |
| 164.9-333.9          | 8.515±0.009                                 | 8.243±0.009                                  | 1.006  | 8.704±0.007      | 10.951±0.007                                 | 8.743±0.068                                   | 1.070   | 10.983±0.008      |
| 360.9-75.3           | 8.250±0.010                                 | 8.233±0.009                                  | 1.002  | 8.543±0.007      | 10.918±0.011                                 | 8.205±0.043                                   | 1.036   | 10.934±0.011      |
| 377.9-64.9           | 8.428±0.010                                 | 7.764±0.008                                  | 1.005  | 8.515±0.009      | 10.840±0.009                                 | 8.508±0.032                                   | 1.171   | 10.911±0.013      |
| 99.6-388.0           | 8.238±0.009                                 | 8.242±0.009                                  | 1.004  | 8.543±0.006      | 10.955±0.010                                 | 8.605±0.070                                   | 1.034   | 10.972±0.011      |
| 397.4-71.7           | 8.356±0.009                                 | 7.906±0.009                                  | 1.024  | 8.498±0.008      | 10.677±0.021                                 | 9.056±0.098                                   | 1.102   | 10.730±0.021      |
| 226.9-366.4          | 8.571±0.010                                 | 7.826±0.009                                  | 1.006  | 8.646±0.009      | 10.900±0.008                                 | 8.685±0.051                                   | 1.193   | 10.979±0.013      |
| 405.5-157.7          | 8.320±0.008                                 | 7.585±0.009                                  | 1.011  | 8.398±0.007      | 10.926±0.019                                 | 8.955±0.079                                   | 1.186   | 11.005±0.021      |
| 345.5-273.8          | 8.314±0.009                                 | 8.060±0.010                                  | 1.009  | 8.510±0.007      | 10.872±0.075                                 | 8.821±0.089                                   | 1.066   | 10.902±0.075      |
| 410.3-206.3          | 8.574±0.011                                 | 7.921±0.009                                  | 1.004  | 8.663±0.009      | 10.903±0.012                                 | 8.503±0.109                                   | 1.167   | 10.972±0.015      |
| 371.1-280.0          | 8.322±0.007                                 | 8.011±0.009                                  | 1.004  | 8.497±0.005      | 10.911±0.008                                 | 8.512±0.054                                   | 1.078   | 10.945±0.009      |
| 368.3-285.6          | 8.364±0.007                                 | 8.181±0.009                                  | 1.003  | 8.584±0.006      | 10.933±0.007                                 | 8.382±0.058                                   | 1.055   | 10.958±0.008      |
| 392.0-270.1          | 8.129±0.009                                 | 8.183±0.009                                  | 1.008  | 8.461±0.006      | 10.932±0.008                                 | 8.811±0.066                                   | 1.030   | 10.948±0.008      |
| 481.4-0.5            | 8.088±0.009                                 | 7.654±0.009                                  | 1.011  | 8.229±0.007      | 10.840±0.009                                 | 8.884±0.061                                   | 1.103   | 10.887±0.011      |
| 453.8-191.8          | 8.160±0.010                                 | 7.943±0.008                                  | 1.003  | 8.367±0.007      | 10.887±0.008                                 | 8.396±0.056                                   | 1.061   | 10.914±0.009      |
| 331.9-401.0          | 8.119±0.008                                 | 8.009±0.009                                  | 1.003  | 8.370±0.006      | 10.862±0.026                                 | 8.393±0.129                                   | 1.045   | 10.883±0.027      |
| 509.5-264.1          | 8.124±0.009                                 | 8.080±0.009                                  | 1.004  | 8.405±0.006      | 10.900±0.008                                 | 8.517±0.011                                   | 1.038   | 10.918±0.008      |
| 266.0-534.1          | 8.124±0.010                                 | 7.967±0.008                                  | 1.034  | 8.368±0.008      | 10.873±0.059                                 | 9.401±0.076                                   | 1.047   | 10.909±0.057      |
| 667.9-174.1          | 7.677±0.011                                 | 8.051±0.006                                  | 1.006  | 8.207±0.005      | 10.908±0.008                                 | 8.705±0.014                                   | 1.021   | 10.920±0.008      |
| 1.0-885.8            | 7.973±0.010                                 | 7.432±0.009                                  | 1.020  | 8.092±0.008      | 10.870±0.031                                 | 9.160±0.020                                   | 1.127   | 10.930±0.031      |
| 6.6-886.3            | 7.945±0.009                                 | 7.602±0.009                                  | 1.010  | 8.112±0.007      | 10.897±0.017                                 | 8.884±0.038                                   | 1.083   | 10.935±0.017      |
| NGC5194-30.2-2.2     | 7.364±0.009                                 | 6.264±0.008                                  | 1.042  | 7.415±0.008      | 10.632±0.007                                 | 9.258±0.050                                   | 1.242   | 10.743±0.013      |
| NGC2403-38-51        | 8.356±0.028                                 | 7.909±0.034                                  | 1.018  | 8.495±0.023      | 10.895±0.032                                 | 9.153±0.267                                   | 1.110   | 10.947±0.034      |
| NGC2403-7-37         | 8.319±0.028                                 | 7.831±0.036                                  | 1.006  | 8.444±0.023      | 10.917±0.030                                 | 8.683±0.287                                   | 1.126   | 10.971±0.035      |
| NGC2403-119-28       | 8.610±0.018                                 | 8.252±0.019                                  | 1.007  | 8.771±0.014      | 10.940±0.021                                 | 8.800±0.270                                   | 1.088   | 10.980±0.023      |
| NGC2403-59-118       | 8.470±0.029                                 | 8.378±0.035                                  | 1.016  | 8.734±0.023      | 10.961±0.029                                 | 9.189±0.275                                   | 1.044   | 10.987±0.030      |
| NGC2403-96-30        | 8.476±0.009                                 | 8.139±0.014                                  | 1.007  | 8.643±0.008      | 10.933±0.018                                 | 8.806±0.214                                   | 1.082   | 10.970±0.018      |
| NGC2403-44-82        | 8.587±0.031                                 | 7.948±0.037                                  | 1.014  | 8.683±0.026      | 10.913±0.027                                 | 9.048±0.279                                   | 1.169   | 10.985±0.038      |

Table 3 – *continued*

| ObjID           | <sup>a</sup> O <sup>+</sup> /H <sup>+</sup> | <sup>a</sup> O <sup>2+</sup> /H <sup>+</sup> | ICF(O) | <sup>b</sup> O/H | <sup>a</sup> He <sup>+</sup> /H <sup>+</sup> | <sup>a</sup> He <sup>2+</sup> /H <sup>+</sup> | ICF(He) | <sup>b</sup> He/H |
|-----------------|---------------------------------------------|----------------------------------------------|--------|------------------|----------------------------------------------|-----------------------------------------------|---------|-------------------|
| NGC2403-166-140 | 8.304±0.030                                 | 8.020±0.036                                  | 1.012  | 8.491±0.024      | 10.970±0.033                                 | 9.057±0.269                                   | 1.075   | 11.007±0.035      |
| NGC2403-99-59   | 8.583±0.024                                 | 7.982±0.028                                  | 1.003  | 8.682±0.020      | 10.895±0.020                                 | 8.376±0.291                                   | 1.159   | 10.961±0.027      |
| NGC2403-196-58  | 8.435±0.016                                 | 8.197±0.017                                  | 1.006  | 8.636±0.012      | 10.922±0.019                                 | 8.687±0.284                                   | 1.064   | 10.952±0.019      |
| NGC2403-22-162  | 8.549±0.018                                 | 7.926±0.020                                  | 1.019  | 8.649±0.016      | 10.926±0.026                                 | 9.211±0.344                                   | 1.154   | 10.996±0.030      |
| NGC2403-160-251 | 8.248±0.031                                 | 7.943±0.036                                  | 1.015  | 8.431±0.024      | 10.908±0.028                                 | 9.093±0.088                                   | 1.078   | 10.947±0.030      |
| Reg-1           | 6.672±0.016                                 | 7.037±0.019                                  | 1.029  | 7.205±0.014      | 10.861±0.014                                 | 9.316±0.131                                   | 1.021   | 10.883±0.014      |
| J0118-3512      | 7.334±0.005                                 | 7.479±0.001                                  | 1.032  | 7.727±0.002      | 10.937±0.015                                 | 9.443±0.027                                   | 1.025   | 10.962±0.015      |
| J1322-5425      | 6.969±0.010                                 | 7.358±0.001                                  | 1.013  | 7.512±0.003      | 10.820±0.005                                 | 8.946±0.035                                   | 1.021   | 10.835±0.005      |
| 0723-692A       | 7.182±0.007                                 | 7.792±0.001                                  | 1.010  | 7.892±0.002      | 10.912±0.004                                 | 8.895±0.048                                   | 1.020   | 10.925±0.004      |
| 0723-692B       | 7.650±0.003                                 | 7.672±0.003                                  | 1.015  | 7.969±0.003      | 10.896±0.013                                 | 9.084±0.123                                   | 1.032   | 10.917±0.013      |
| 0907-543        | 7.441±0.009                                 | 7.909±0.007                                  | 1.032  | 8.050±0.006      | 10.895±0.029                                 | 9.398±0.015                                   | 1.021   | 10.918±0.028      |
| 0917-527        | 7.722±0.002                                 | 7.695±0.001                                  | 1.025  | 8.021±0.002      | 10.903±0.012                                 | 9.300±0.056                                   | 1.035   | 10.929±0.012      |
| 0926-606        | 7.730±0.002                                 | 7.762±0.002                                  | 1.018  | 8.055±0.002      | 10.889±0.013                                 | 9.139±0.055                                   | 1.031   | 10.910±0.013      |
| 0930-554        | 6.853±0.001                                 | 7.086±0.002                                  | 1.038  | 7.302±0.002      | 10.801±0.005                                 | 9.382±0.032                                   | 1.023   | 10.828±0.005      |
| 1030-583        | 7.409±0.001                                 | 7.700±0.002                                  | 1.027  | 7.891±0.001      | 10.887±0.009                                 | 9.320±0.036                                   | 1.022   | 10.908±0.009      |
| 1116-583B       | 7.086±0.015                                 | 7.602±0.010                                  | 1.032  | 7.731±0.010      | 10.845±0.038                                 | 9.340±0.172                                   | 1.020   | 10.866±0.037      |
| 1205-557        | 7.716±0.004                                 | 7.526±0.005                                  | 1.020  | 7.941±0.005      | 10.886±0.022                                 | 9.194±0.170                                   | 1.053   | 10.917±0.022      |
| 1222-614        | 7.566±0.001                                 | 7.867±0.002                                  | 1.019  | 8.051±0.001      | 10.886±0.009                                 | 9.163±0.051                                   | 1.022   | 10.904±0.008      |
| 1223-487        | 7.229±0.001                                 | 7.689±0.001                                  | 1.013  | 7.824±0.001      | 10.900±0.004                                 | 9.021±0.036                                   | 1.021   | 10.914±0.004      |
| 1256-351        | 7.583±0.001                                 | 7.909±0.001                                  | 1.012  | 8.082±0.001      | 10.908±0.004                                 | 8.972±0.039                                   | 1.022   | 10.923±0.004      |
| 1319-579A       | 7.584±0.001                                 | 8.020±0.002                                  | 1.008  | 8.159±0.001      | 10.926±0.008                                 | 8.831±0.055                                   | 1.021   | 10.938±0.008      |
| 1319-579B4      | 7.933±0.011                                 | 7.663±0.010                                  | 1.047  | 8.139±0.014      | 10.856±0.050                                 | 9.512±0.259                                   | 1.062   | 10.901±0.050      |
| 1319-579C       | 8.125±0.002                                 | 7.929±0.003                                  | 1.015  | 8.345±0.002      | 10.899±0.012                                 | 9.065±0.122                                   | 1.055   | 10.928±0.012      |
| 1358-576        | 7.697±0.003                                 | 7.748±0.002                                  | 1.012  | 8.030±0.002      | 10.936±0.008                                 | 9.015±0.074                                   | 1.030   | 10.955±0.008      |
| 1441-294        | 7.758±0.009                                 | 7.882±0.009                                  | 1.018  | 8.133±0.009      | 10.945±0.035                                 | 9.194±0.318                                   | 1.027   | 10.963±0.034      |
| 1533-574B       | 7.953±0.002                                 | 7.985±0.003                                  | 1.011  | 8.275±0.002      | 10.913±0.012                                 | 8.966±0.120                                   | 1.032   | 10.931±0.012      |
| Pox105          | 7.502±0.001                                 | 7.747±0.010                                  | 1.019  | 7.951±0.006      | 10.846±0.019                                 | 9.120±0.083                                   | 1.023   | 10.864±0.019      |
| Pox120          | 7.276±0.001                                 | 7.777±0.009                                  | 1.019  | 7.905±0.007      | 10.859±0.018                                 | 9.144±0.079                                   | 1.020   | 10.876±0.018      |
| Pox139          | 7.663±0.001                                 | 7.857±0.010                                  | 1.026  | 8.083±0.006      | 10.880±0.018                                 | 9.297±0.054                                   | 1.024   | 10.902±0.018      |
| UM160-A         | 7.638±0.010                                 | 7.833±0.045                                  | 1.005  | 8.050±0.028      | 10.875±0.027                                 | 8.533±0.135                                   | 1.025   | 10.889±0.027      |
| UM160-B         | 7.753±0.009                                 | 7.868±0.042                                  | 1.027  | 8.127±0.024      | 10.865±0.031                                 | 9.291±0.064                                   | 1.027   | 10.887±0.030      |
| UM420-B         | 8.019±0.008                                 | 7.845±0.426                                  | 1.023  | 8.257±0.168      | 10.926±0.023                                 | 9.292±0.109                                   | 1.052   | 10.955±0.022      |
| TOL0513-393     | 7.134±0.008                                 | 7.900±0.043                                  | 1.009  | 7.972±0.036      | 10.971±0.017                                 | 8.941±0.065                                   | 1.020   | 10.983±0.017      |
| TOL2146-391-C   | 7.182±0.001                                 | 7.754±0.004                                  | 1.018  | 7.865±0.003      | 10.914±0.008                                 | 9.171±0.023                                   | 1.020   | 10.931±0.008      |
| TOL2146-391-E   | 7.202±0.001                                 | 7.747±0.004                                  | 1.020  | 7.864±0.003      | 10.906±0.012                                 | 9.196±0.032                                   | 1.020   | 10.922±0.012      |
| TOL0357-3915-C  | 7.364±0.005                                 | 7.861±0.004                                  | 1.019  | 7.990±0.004      | 10.929±0.012                                 | 9.215±0.037                                   | 1.020   | 10.946±0.012      |
| TOL0357-3915-E  | 7.387±0.005                                 | 7.855±0.004                                  | 1.014  | 7.988±0.004      | 10.902±0.021                                 | 9.050±0.065                                   | 1.021   | 10.917±0.020      |
| NGC346          | 7.506±0.005                                 | 7.930±0.004                                  | 1.002  | 8.069±0.003      | 10.915±0.004                                 | 8.229±0.042                                   | 1.021   | 10.925±0.004      |

<sup>a</sup> Ionic abundance in units of  $12 + \log_{10}(X^{n+}/H^+)$ <sup>b</sup> Total abundance in units of  $12 + \log_{10}(X/H)$