

POLYNOMIAL VOTING RULES

WENPIN TANG AND DAVID D. YAO

ABSTRACT. We propose and study a new class of polynomial voting rules for a general decentralized decision/consensus system, and more specifically for the PoS (Proof of Stake) protocol. The main idea, inspired by the Penrose square-root law and the more recent quadratic voting rule, is to differentiate a voter’s voting power and the voter’s share (fraction of the total in the system). We show that while voter shares form a martingale process, their voting powers follow a super-martingale that decays to zero over time. This prevents any voter from controlling the voting process, and thus enhances decentralization. When the initial total volume of votes (or stakes) is large, we show a phase transition in share stability (or the lack thereof), corresponding to the voter’s initial share relative to the total. We also study the scenario in which trading (of votes/stakes) among the voters is allowed, and quantify the level of risk sensitivity (or risk averse) that will remove any incentive for a voter to trade.

Key words: voting rules, PoS protocol, phase transition, stability, large deviations, concentration inequalities.

1. INTRODUCTION

Voting, in the traditional sense, refers to a set of rules for a community of individuals or groups (“voters”) to reach an agreement, or to make a collective decision on some choices and ranking problems. In today’s world, “voting” has become a ubiquitous notion that includes any decentralized decision-making protocol or system, where the voters are often abstract entities (“virtual”), and the voting process automated; and the purpose of reaching consensus is often non-social and non-political, such as to enhance the overall security of an industrial operation or infrastructure (Garcia-Molina (1982); Lamport et al. (1982)). Examples include cloud computing (Armbrust et al. (2009); Dean and Ghemawat (2008)), smart power grids (Huang and Baliga (2009)), and more recently, trading or payment platforms and exchanges built upon the blockchain technology (Nakamoto (2008); Wood (2014)).

At the core of a blockchain is the *consensus protocol*, which specifies a set of voting rules for the participants (“miners” or validators) to agree on an ever-growing log of transactions (the “longest chain”) so as to form a distributed ledger. There are several existing blockchain protocols, among which the most popular are Proof of Work (PoW, Nakamoto (2008)) and Proof of Stake (PoS, King and Nadal (2012); Wood (2014)). In the PoW protocol, miners compete with each other by solving a hashing puzzle. The miner who solves the puzzle first receives a reward (a number of coins) and whose work validates a new block’s addition to the blockchain. Hence, while the competition is open to everyone, the chance of winning is proportional to a miner’s computing power.

Date: July 13, 2022.

In the PoS protocol, there is a bidding mechanism to select a miner to do the work of validating a new block. Participants who choose to join the bidding are required to commit some “stakes” (coins they own), and the winning probability is proportional to the number of stakes committed. Hence, a participant in a PoS blockchain is actually a “bidder,” as opposed to a “miner” — only the winning bidder becomes the miner validating the block. (Any participant who chooses not to join the bidding can be viewed as a bidder who commits zero stakes.) Needless to add, bidding exists long before the PoS protocol, and has been widely used in many applications, such as auctions and initial public offerings (IPOs).

Let’s explore the PoS bidding mechanism a bit more formally. Suppose a voter (or bidder) k is in possession of $n_{k,t}$ votes (or stakes) at time t , an index that counts the rounds of voting or bidding in the protocol; and $N_t := \sum_k n_{k,t}$ is the total number of votes over all voters. Hence, voter k ’s share, fraction of the total, is $\pi_{k,t} := n_{k,t}/N_t$. Following a traditional voting rule, voter k ’s chance or probability of winning, which we call *voting power*, will be equal to $\pi_{k,t}$, voter k ’s share. Yet, this doesn’t have to be the case. That is, any voter’s voting power needs not be equal to the voter’s share (of the system total). Indeed, there are often good reasons for the two to be different.

Historically, the English scholar Lionel Penrose famously proposed a square-root voting rule (Penrose (1946)), around the time when the United Nations was founded shortly after WWII. According to Penrose, a world assembly such as the UN should designate each country a number of votes that is proportional to the square root of its population. The obvious idea is to limit the voting power of nations with very large populations. In the same spirit, the quadratic voting rule has attracted much attention in recent years (Lalley and Weyl (2018)). The idea is that each voter be given a budget (in dollars, for instance); the voter can cast multiple votes on any single or subset of choices or candidates on the ballot, with x votes (for any choice) costing x^2 dollars. Under both voting rules, the voting power is different from the voter’s share or representation in the system, population in the first case, and the budget in the second case.

Inspired by these ideas, we propose a new class of polynomial voting rules, denoted $\text{Poly}(\alpha)$, which grant every voter k a voting power that scales the voter’s share $\pi_{k,t}$ by a factor $N_t^{-\alpha}$ for $\alpha \geq 0$. When $\alpha = 0$, this reduces to the traditional voting case of power=share, which is a linear rule. When $\alpha = 1$, the rule is similar in spirit to the square-root and the quadratic voting rules mentioned above.

Here’s an overview of our main findings and results. We prove that under the $\text{Poly}(\alpha)$ voting rule, voter shares form a martingale process, which converges to some non-degenerate probability distribution as $t \rightarrow \infty$; while their voting powers follow a super-martingale process that decays to zero over time (Theorem 1). Thus, the $\text{Poly}(\alpha)$ voting enhances decentralization, preventing any voter or any group of voters from controlling the voting process and overpowering the system.

We further group the voters into three categories: large, medium and small, according to the initial (time zero) votes they own relative to the total (N_0). When N_0 is large, which is the case in most applications, we show a phase transition in the stability of voter shares across the three categories (Theorem 5). Notably, the same phenomenon is demonstrated under the traditional voting rule ($\alpha = 0$), refer to Roşu and Saleh (2021); Tang (2022), where the analysis is based on the well-known Pólya urn model. Our result establishes that this phase transition is in fact *universal*, in the sense that it applies to all values of $\alpha(\geq 0)$.

We also study the scenario in which trading (of votes/stakes) among the voters is allowed, motivated by PoS applications in cryptocurrency. For $\alpha = 0$, the trading scenario has been recently studied in Roşu and Saleh (2021). Not only our model is more general, in allowing any $\alpha \geq 0$, our results are also richer and sharper (Theorem 7). For instance, we quantify the level of risk sensitivity (or risk-averse) that will remove any incentive for a voter to trade, or even not to participate at all (in the bidding)—the notion of *non-participation*, which is not considered in Roşu and Saleh (2021).

The key to our analysis relies on the study of the random process N_t (the total volume of votes/stakes at time t), which is a time-homogeneous Markov chain. We develop some asymptotic results for this Markov chain, including large-deviation bounds (Theorem 2) and a fluid limit (Proposition 4).

In the remainder of this paper there are two main sections. Section 2 studies the $\text{PoLy}(\alpha)$ voting model from a general perspective, focusing on the associated stochastic processes, such as N_t , voter shares, and voting powers; their long-term behavior and limits, some of which are further characterized by concentration inequalities or large-deviation bounds. Section 3 concerns two aspects of the $\text{PoLy}(\alpha)$ voting rule that are more closely associated with the application of POS in cryptocurrency: (a) the evolution of bidder shares over time and the phase transition phenomenon mentioned above; (b) the issue of incentive and risk-sensitivity when trading is allowed. The more technical proofs in the two sections are deferred to Section 4. Concluding remarks and suggestions for further research are collected in Section 5.

2. THE $\text{PoLy}(\alpha)$ VOTING MODEL

In this section, we develop a formal model for the $\text{PoLy}(\alpha)$ voting rule, focusing on the stochastic processes associated with the model, their properties and limiting behavior.

First, here is a list of some of the notation used throughout below.

- \mathbb{N}_+ denotes the set of positive integers, and \mathbb{R} denotes the set of real numbers.
- \xrightarrow{d} denotes the convergence in distribution.
- $a = \mathcal{O}(b)$ means $\frac{a}{b}$ is bounded from above as $b \rightarrow \infty$; $a = \Theta(b)$ means $\frac{a}{b}$ is bounded from below and above as $b \rightarrow \infty$; and $a = o(b)$ or $b \gg a$ means $\frac{a}{b}$ decays towards zero as $b \rightarrow \infty$.

We use C, C', C'' etc to denote generic constants (which may change from line to line).

The voters, referred to as *bidders* below, are the participants in the decentralized system, where they engage in rounds of bidding following a pre-specified voting rule (the “consensus protocol”) so as to win more votes, or *stakes*. (The PoS protocol described in the Introduction provides a concrete instance to motivate the model here.) Let $K \in \mathbb{N}_+$ be the total number of bidders, which will stay fixed throughout the paper; and let $[K] := \{1, \dots, K\}$ denote the set of all bidders.

Time is discrete, indexed by $t = 0, 1, 2, \dots$, and corresponds to the rounds of bidding mentioned above. Bidder k initially owns $n_{k,0}$ stakes. Let $N := \sum_{k=1}^K n_{k,0}$ denote the total number of initial stakes owned by all K bidders. The term *bidder share* refers to the fraction

of stakes each bidder owns. So the initial bidder shares $(\pi_{k,0}, k \in [K])$ are given by

$$\pi_{k,0} := \frac{n_{k,0}}{N}, \quad k \in [K]. \quad (1)$$

Similarly, $n_{k,t}$ denotes the number of stakes owned by bidder k at time $t \in \mathbb{N}_+$, and the corresponding share is

$$\pi_{k,t} := \frac{n_{k,t}}{N_t}, \quad k \in [K], \quad \text{with } N_t := \sum_{k=1}^K n_{k,t}. \quad (2)$$

Here N_t is the total number of stakes at time t , and thus $N_0 = N$. (We shall often refer to N_t as the ‘‘volume of stakes’’ or, simply, ‘‘volume.’’) Clearly, for each $t \geq 0$, $(\pi_{k,t}, k \in [K])$ forms a probability distribution on $[K]$.

In each period t , a single stake (or ‘‘reward’’) is distributed as follows: each bidder k receives the reward with probability

$$\theta_{k,t} := \frac{n_{k,t}}{N_t^{1+\alpha}} = \frac{\pi_{k,t}}{N_t^\alpha}, \quad (3)$$

and receives nothing with probability $1 - \theta_{k,t}$. We call $\theta_{k,t}$ bidder k 's *voting power* at time t . When $\alpha = 0$, the voting power $\theta_{k,t}$ coincides with the bidder share $\pi_{k,t}$; this is the Pólya urn framework in Roşu and Saleh (2021); Tang (2022). On the other hand, when $\alpha = 1$, the $\text{Poly}(\alpha)$ model shares the same spirit of the Penrose square-root rule and the quadratic voting rule motivated in the Introduction.

Let $S_{k,t}$ be the random event that bidder k receives one unit of reward in period t . Thus, the number of stakes owned by each bidder evolves as follows,

$$n_{k,t} = n_{k,t-1} + 1_{S_{k,t}}, \quad k \in [K]; \quad (4)$$

or simply,

$$n_{k,t} = \begin{cases} n_{k,t-1} & \text{with probability } 1 - \theta_{k,t-1}, \\ n_{k,t-1} + 1 & \text{with probability } \theta_{k,t-1}. \end{cases} \quad (5)$$

Accordingly, the total number of stakes N_t evolves as follows, taking into account $\sum_{k=1}^K \theta_{k,t} = N_t$,

$$N_t = \begin{cases} N_{t-1} & \text{with probability } 1 - 1/N_{t-1}^\alpha, \\ N_{t-1} + 1 & \text{with probability } 1/N_{t-1}^\alpha. \end{cases} \quad (6)$$

In contrast with the Pólya urn (with a constant reward) in which N_t grows deterministically and linearly in t , here N_t as specified in (6) is a nondecreasing time-homogeneous Markov chain on $\{N, N+1, \dots\}$. As we will see in Theorem 1, the $\text{Poly}(\alpha)$ voting rule slows down the distribution of rewards, so the volume of stakes grows sublinearly. (This is consistent with the volume growth in many cryptocurrencies such as Bitcoin and Ethereum.)

2.1. Convergence and Large Deviations. Let \mathcal{F}_t be the filtration as the σ -algebra generated by the random events $(S_{k,r} : k \in [K], r \leq t)$. Our first result specifies the limiting behavior of N_t , as well as the martingale properties of $(\pi_{k,t}, k \in [K])$ and $(\theta_{k,t}, k \in [K])$.

Theorem 1 (Long-time behavior). *Under the $\text{Poly}(\alpha)$ voting rule, we have*

(i) *Volume of stakes: there is the convergence in probability*

$$\frac{N_t^{1+\alpha}}{t} \xrightarrow{d} 1 + \alpha \quad \text{as } t \rightarrow \infty. \quad (7)$$

(ii) Bidder shares: the process $(\pi_{k,t}, t \geq 0)$ is an \mathcal{F}_t -martingale, and with probability one,

$$(\pi_{1,t}, \dots, \pi_{K,t}) \longrightarrow (\pi_{1,\infty}, \dots, \pi_{K,\infty}) \quad \text{as } t \rightarrow \infty, \quad (8)$$

where $(\pi_{1,\infty}, \dots, \pi_{K,\infty})$ is some random probability distribution on $[K]$.

(iii) Voting power: the process $(\theta_{k,t}, t \geq 0)$ is an \mathcal{F}_t -super-martingale, and with probability one,

$$\theta_{k,t} \rightarrow 0 \quad \text{as } t \rightarrow \infty, k \in [K]. \quad (9)$$

Deferring the proof of Theorem 1 (to the end of this subsection), we first make a few remarks here. Part (i) of the theorem gives the growth rate of the volume of stakes: N_t grows as $((\alpha + 1)t)^{\frac{1}{1+\alpha}}$ as $t \rightarrow \infty$. When $\alpha = 0$, N_t follows the (deterministic) linear growth of the Pólya urn model (with a constant reward). For $\alpha = 1$, N_t grows as \sqrt{t} . Part (ii) shows that the bidder shares form a martingale, and converges in long term. See §3.1 for further discussions on the stability of the bidder shares when the initial stakes $N := N_0$ is large. Part (iii) reveals that each bidder's voting power decays to zero in the long run. This means that the power is diluted over time, and no individual can manipulate or control the voting process. In other words, decentralization is enhanced.

The next important question is how N_t “fluctuates” around its growth trajectory $((\alpha + 1)t)^{\frac{1}{1+\alpha}}$; specifically, how to establish large-deviation bounds on N_t . These bounds are provided in the next theorem, along with a corollary that confirms N_t 's concentration around its growth trajectory, for large t . (These results are useful to the proof of Theorem 1 (iii), and also to the study of the stability of the bidder shares in Section 3.)

Theorem 2 (Large deviations for N_t). *Define a function $f_\alpha(\cdot)$,*

$$f_\alpha : \lambda \in (0, \infty) \mapsto (1 + \alpha)\lambda \log \lambda - (1 + \alpha)\lambda + \frac{1}{\lambda^\alpha} \in \mathbb{R}.$$

Let $\lambda_-(\alpha) < \lambda_+(\alpha)$ be the two roots of $f_\alpha(\cdot)$ on $(-\infty, \infty)$. Under the *Pol $\mathbf{y}(\alpha)$* voting rule, we have

(i) For each $\lambda < \lambda_-(\alpha)$, we have for any $\varepsilon > 0$,

$$\mathbb{P}(N_t < \lambda t^{\frac{1}{1+\alpha}}) \leq \exp\left(-(1 - \varepsilon)f_\alpha(\lambda) t^{\frac{1}{1+\alpha}}\right) \quad \text{as } t \rightarrow \infty. \quad (10)$$

(ii) For each $\lambda > \lambda_+(\alpha)$, we have for any $\varepsilon > 0$,

$$\mathbb{P}(N_t > \lambda t^{\frac{1}{1+\alpha}}) \leq \exp\left(-(1 - \varepsilon)f_\alpha(\lambda) t^{\frac{1}{1+\alpha}}\right) \quad \text{as } t \rightarrow \infty. \quad (11)$$

As a corollary, the volume of stakes N_t concentrates around $((1 + \alpha)t)^{\frac{1}{1+\alpha}}$ for large t .

Corollary 3. *Under the *Pol $\mathbf{y}(\alpha)$* urn model, we have for each $\delta > 0$,*

$$\mathbb{P}\left(|N_t - ((1 + \alpha)t)^{\frac{1}{1+\alpha}}| > \delta t^{\frac{1}{1+\alpha}}\right) \rightarrow 0 \quad \text{as } t \rightarrow \infty. \quad (12)$$

The proofs of the theorem and the corollary above are deferred to §4.1 and §4.2.

Theorem 2 gives exponential deviation bounds for N_t when it is either sufficiently small (below $\lambda_-(\alpha)t^{\frac{1}{1+\alpha}}$), or sufficiently large (above $\lambda_+(\alpha)t^{\frac{1}{1+\alpha}}$). Note that there is a gap between the two bounding curves, since for each $\alpha > 0$, $\lambda_-(\alpha) < (1 + \alpha)^{\frac{1}{1+\alpha}} < \lambda_+(\alpha)$. (For instance, for $\alpha = 1$, $\lambda_-(1) \approx 0.56 < \sqrt{2} < 2.51 \approx \lambda_+(1)$.) The gap is due to the combinatorial

estimates in our proof, which may very well be improved. Refer to the Appendix for a numerical procedure that shrinks the gap.

Proof of Theorem 1. (i) We use the method of moments by computing $\mathbb{E}(N_t^{(1+\alpha)^j})$ for $j = 1, 2, \dots$. For $j = 1$, we have by definition

$$\begin{aligned} \mathbb{E}(N_{t+1}^{1+\alpha} - N_t^{1+\alpha} | N_t = x) &= (1+x)^{1+\alpha} \frac{1}{x^\alpha} + x^{1+\alpha} \left(1 - \frac{1}{x^\alpha}\right) - x^{1+\alpha} \\ &= 1 + \alpha + \mathcal{O}(x^{-1}) \quad \text{as } x \rightarrow \infty. \end{aligned}$$

The discussion after Theorem 1 implies that with probability one $N_t \rightarrow \infty$ as $t \rightarrow \infty$. As a result, $\mathbb{E}(N_{t+1}^{1+\alpha} - N_t^{1+\alpha}) \rightarrow 1 + \alpha$ as $t \rightarrow \infty$ which yields

$$\mathbb{E}N_t^{1+\alpha} \sim (1 + \alpha)t \quad \text{as } t \rightarrow \infty. \quad (13)$$

Next for $j = 2$, we have

$$\mathbb{E}(N_{t+1}^{2(1+\alpha)} - N_t^{2(1+\alpha)} | N_t = x) = 2(1 + \alpha)x^{1+\alpha} + \mathcal{O}(x^\alpha) \quad \text{as } x \rightarrow \infty.$$

Thus, $\mathbb{E}(N_{t+1}^{2(1+\alpha)} - N_t^{2(1+\alpha)}) = (2(1 + \alpha) + o(1))\mathbb{E}N_t^{1+\alpha} \sim 2(1 + \alpha)^2t$ by (13). Then we get $\mathbb{E}(N_t^{2(1+\alpha)}) \sim (1 + \alpha)^2t^2$ as $t \rightarrow \infty$. Proceeding by induction, we obtain

$$\mathbb{E}(N_t^{(1+\alpha)^j}) \sim (1 + \alpha)^j t^j \quad \text{as } t \rightarrow \infty, \quad j = 1, 2, \dots$$

By the method of moments (see e.g. (Billingsley, 1995, Section 30)), $N_t^{(1+\alpha)}/t$ converges in distribution, and thus in probability to $1 + \alpha$.

(ii) By (2) and (5), it is easily seen that for each $k \in [K]$ and $t \geq 0$,

$$\mathbb{E}(\pi_{k,t+1} | \mathcal{F}_t) = \frac{n_{k,t}}{N_t} \left(1 - \frac{1}{N_t^\alpha}\right) + \frac{n_{k,t}}{N_t + 1} \frac{N_t - n_{k,t}}{N_t^{1+\alpha}} + \frac{n_{k,t} + 1}{N_t + 1} \frac{n_{k,t}}{N_t^{1+\alpha}}. \quad (14)$$

Recognizing the first term on the right side of (14), $\frac{n_{k,t}}{N_t} = \pi_{k,t}$, while all other terms sum up to zero, we conclude that $(\pi_{k,t}, t \geq 0)$ is an \mathcal{F}_t -martingale. The convergence in (8) follows from the martingale convergence theorem (see e.g. (Durrett, 2019, Section 4.2)).

(iii) Applying the same derivation as in (14) but to $\theta_{k,t}$ instead, we have

$$\mathbb{E}(\theta_{k,t+1} | \mathcal{F}_t) = \frac{\pi_{k,t}}{N_t^\alpha} \left(1 - \frac{1}{N_t^\alpha}\right) + \frac{N_t \pi_{k,t}}{(N_t + 1)^{1+\alpha}} \cdot \frac{1 - \pi_{k,t}}{N_t^\alpha} + \frac{N_t \pi_{k,t} + 1}{(N_t + 1)^{1+\alpha}} \cdot \frac{\pi_{k,t}}{N_t^\alpha}. \quad (15)$$

The last two terms add up to $\frac{\pi_{k,t}}{N_t^\alpha(N_t+1)^\alpha} = \frac{\theta_{k,t}}{(N_t+1)^\alpha}$. Thus, we have

$$\mathbb{E}(\theta_{k,t+1} | \mathcal{F}_t) = \theta_{k,t} \left(1 - \frac{1}{N_t^\alpha} + \frac{1}{(N_t + 1)^\alpha}\right) \leq \theta_{k,t}, \quad (16)$$

i.e. $(\theta_{k,t}, t \geq 0)$ is an \mathcal{F}_t -super-martingale for each k . Again by the martingale convergence theorem, $\theta_{k,t} \rightarrow \theta_{k,\infty}$ as $t \rightarrow \infty$ with probability one. Fix $L > 0$ to be sufficiently large, and let $\Omega_{L,t} := \{N_t > Lt^{\frac{1}{1+\alpha}}\}$. By (16),

$$\begin{aligned} \mathbb{E}(\theta_{k,t+1}) &= \mathbb{E}\left(\theta_{k,t} \left(1 - \frac{1}{N_t^\alpha} + \frac{1}{(N_t + 1)^\alpha}\right) 1_{\Omega_{L,t}^c}\right) + \mathbb{E}\left(\theta_{k,t} \left(1 - \frac{1}{N_t^\alpha} + \frac{1}{(N_t + 1)^\alpha}\right) 1_{\Omega_{L,t}}\right) \\ &\leq \mathbb{E}(\theta_{k,t}) (1 - Ct^{-1}) + \mathbb{P}(\Omega_{L,t}) \quad \text{for some } C > 0. \end{aligned} \quad (17)$$

By Theorem 2 (2), we have $\mathbb{P}(\Omega_{L,t}) = \mathcal{O}\left(\exp\left(-C't^{\frac{1}{1+\alpha}}\right)\right)$ for some $C' > 0$. Combining with (17) yields $\mathbb{E}(\theta_{k,t}) = \mathcal{O}\left(t^{-C''+1}\right)$ for $C'' > 0$ arbitrarily large. By the dominated convergence theorem, we get $\theta_{k,\infty} = 0$ with probability one. \square

2.2. A Fluid Limit. Recall that the process of the volume of stakes $(N_t, t \geq 0)$ is a time-homogenous Markov chain. The path properties of a general time-homogenous Markov chain $(Z_t, t \geq 0)$ has long been studied since the work of Lamperti (1960, 1962, 1963). The basic idea is to study the recurrence, or transience of $(Z_t, t \geq 0)$ based on

$$m_1(x) = \mathbb{E}(Z_{t+1} - Z_t | Z_t = x) \quad \text{and} \quad m_2(x) = \mathbb{E}((Z_{t+1} - Z_t)^2 | Z_t = x).$$

For instance, if $\limsup_{x \rightarrow \infty} 2xm_1(x) + m_2(x) \leq 0$ then $(Z_t, t \geq 0)$ is recurrent; and if $\liminf_{x \rightarrow \infty} 2xm_1(x) + m_2(x) > 0$ then $(Z_t, t \geq 0)$ is transient. The regime corresponding to $m_1(x) = o(1)$ is called the *Markov chain with asymptotic zero drift*, and features active research (see e.g. Denisov et al. (2016); Menshikov et al. (2017)). Specializing to the process $(N_t, t \geq 0)$, we have

$$m_1(x) = m_2(x) = \frac{1}{x^\alpha}.$$

Interestingly, there seem to be few results on the Lamperti's problem where both $m_1(x)$ and $m_2(x)$ decreases to zero as $x \rightarrow \infty$, except that $(N_t, t \geq 0)$ is transient. Observe that $(N_t, t \geq 0)$ is nondecreasing, and the ‘‘upward’’ variance is larger than the ‘‘downward’’ variance. In the similar spirit to Lamperti (1962), the asymptotic growth (7) hinges on a degenerate fluid approximation of the process $(N_t, t \geq 0)$, as stated in the following proposition.

Proposition 4 (Fluid limit of N_t). *Under the $\text{Poly}(\alpha)$ voting rule, we have*

$$\left(\frac{N_{nu}}{n^{\frac{1}{1+\alpha}}}, u \geq 0\right) \xrightarrow{d} (X_u, u \geq 0) \quad \text{as } n \rightarrow \infty \text{ in } \mathcal{C}[0, \infty), \quad (18)$$

where N_s for non-integer s is defined by the linear interpolation of the chain $(N_t, t \geq 0)$, and $X_u = ((1 + \alpha)u)^{\frac{1}{1+\alpha}}$, $u \geq 0$ is the solution to the ordinary differential equation $dX_u = X_u^{-\alpha} dt$ with $X_0 = 0$.

The proof of Proposition 4 is given in §4.3. Note that the fluid limit in the proposition is different from the fluid limit in the literature of stochastic networks, where it usually takes the form of a FSLLN (functional strong law of large numbers) concerning a renewal process and the associated counting process. In that setting, the convergence (to a deterministic function of time) is stronger – the almost sure convergence, and uniformly on $[0, T]$. Refer to (Chen and Yao, 2001, §6.1). Here, the process $(N_t, t \geq 0)$ is non-renewal, thus the FSLLN limit does not apply; yet there is still the *weak* convergence, and the limit is still a deterministic function of time (X_u) , explicitly characterized above. Another notable point is, in the FSLLN setting, both time and space are scaled by the same scaling factor n whereas in (18) the time scaling remains the same, and the space scaling is by $n^{\frac{1}{1+\alpha}}$. But this is only because $(N_t, t \geq 0)$ grows in the order of $t^{\frac{1}{1+\alpha}}$ (Theorem 1(i)), whereas a renewal (counting) process grows linearly in t .

3. OTHER RESULTS, WITH PoS-CRYPTO APPLICATIONS

In this section, we present more results associated with the $\text{Poly}(\alpha)$ model that are largely motivated by the application of POS in cryptocurrency. There are two subsections: In the first one, §3.1, we study the evolution of bidder shares when $N := N_0$, the volume of initial stakes, is large. In §3.2, we study the additional feature of allowing the bidders to trade stakes among themselves, focusing on the issue of trading incentive (or the lack thereof).

It is important, as well as interesting, to note that all the results in both subsections are independent of the parametric value of α , and in this sense, *universal*.

3.1. Evolution of bidder shares and phase transitions. As explained in the introduction, one key feature of the $\text{Poly}(\alpha)$ voting rule is that the voting power $\theta_{k,t}$ of any bidder k is different from k 's share $\pi_{k,t}$ of the total volume of stakes at t ; specifically, $\theta_{k,t} = \pi_{k,t}/N_t^\alpha$. We have seen from Theorem 1 (iii) that the voting power is decreasing over time, which facilitates decentralization. On the other hand, the evolution of the share $\theta_{k,t}$ over time, from its initial value $\theta_{k,t}$, in both absolute and relative terms, is an important issue for any bidder k .

In the classical Pólya urn setting, it is shown in Roşu and Saleh (2021) that for a large bidder with initial stake $n_{k,0} = \Theta(N)$, there is the stability in bidder share, in the sense that

$$\mathbb{P}(|\pi_{k,\infty} - \pi_{k,0}| > \varepsilon) \rightarrow 0 \quad \text{as } N \rightarrow \infty.$$

Furthermore, similar, albeit qualitatively different, results are revealed in Tang (2022), for medium and small bidders (following the definitions in parts (ii) and (iii) of the theorem below),

Here, focusing on the ratio $\pi_{k,t}/\pi_{k,0}$, we establish results for the stability of bidder shares (or the lack thereof) for the $\text{Poly}(\alpha)$ model.

Theorem 5 (Phase transitions of $\pi_{k,t}$). *Let $N_0 = N$ be the total number of initial stakes. Under the $\text{Poly}(\alpha)$ voting rule, we have*

- (i) *For $n_{k,0} = f(N)$ such that $f(N) \rightarrow \infty$ as $N \rightarrow \infty$ (i.e. $\pi_{k,0} \gg 1/N$), there is a constant $C > 0$ independent of N , such that for each $\varepsilon > 0$ sufficiently small and each $t \geq 1$ or $t = \infty$,*

$$\mathbb{P}\left(\left|\frac{\pi_{k,t}}{\pi_{k,0}} - 1\right| > \varepsilon\right) \leq \frac{C}{\varepsilon^2 f(N)}, \quad (19)$$

which converges to 0 as $N \rightarrow \infty$.

- (ii) *For $n_{k,0} = \Theta(1)$ (i.e. $\pi_{k,0} = \Theta(1/N)$), we have $\text{Var}\left(\frac{\pi_{k,\infty}}{\pi_{k,0}}\right) = \Theta(1)$. Moreover, there is a constant $c > 0$ independent of N , such that for $\varepsilon > 0$ sufficiently small,*

$$\mathbb{P}\left(\left|\frac{\pi_{k,\infty}}{\pi_{k,0}} - 1\right| > \varepsilon\right) > c. \quad (20)$$

- (iii) *For $n_{k,0} = o(1)$ (i.e. $\pi_{k,0} = o(1/N)$), we have $\text{Var}\left(\frac{\pi_{k,\infty}}{\pi_{k,0}}\right) \rightarrow \infty$ as $N \rightarrow \infty$.*

The proof of Theorem 5 is given in Section 4.4. The theorem reveals a phase transition in the stability of shares, and identifies large, medium and small bidders, in terms of the size of their stakes, according to the categories in the three parts. A large bidder k is guaranteed to have stability, in the precise sense characterized in (19), that the share ratio

$\pi_{k,t}/\pi_{k,0}$ concentrates at 1, and converges to 1 in probability when $N \rightarrow \infty$, for any $t \geq 1$ (including $t = \infty$). For medium bidders, this is reversed: the concentration inequality in (19) becomes the anti-concentration inequality in (20), implying volatility; and more specifically, the variance of $\pi_{k,\infty}/\pi_{k,0}$ is of order $\Theta(1)$. For small bidders, this variance will increase to infinity as $N \rightarrow \infty$.

Notably, the phase transition demonstrated above for the $\text{Poly}(\alpha)$ model is very similar to the phase transition in the Pólya urn model (corresponding to $\alpha = 0$) revealed in (Tang, 2022, Theorem 1 and Theorem 2 (1))). Yet, this is perhaps not surprising, given all three parts of Theorem 5 are independent of α .

3.2. Participation and trading. So far, we have not considered the possibility of allowing the bidders to trade stakes (among themselves). In the classical Pólya urn model ($\alpha = 0$), it is shown in Roşu and Saleh (2021) that under certain conditions (which enforce some notion of “risk neutrality”), there will be no incentive for any bidder to trade. Here, we extend that to the $\text{Poly}(\alpha)$ model, allowing α to take any non-negative values. Furthermore, we allow a bidder-dependent risk-sensitivity (or risk-averse) parameter δ_k , and study the issue of incentive as it relates to δ_k .

In the new setting of allowing trading, we need to modify the problem formulation presented at the beginning of §2. First, for each $k \in [K]$, let $\nu_{k,t}$ be the number of stakes that bidder k will trade at time t . Then, instead of (4), the number of stakes $n_{k,t}$ evolves as

$$n_{k,t} = \underbrace{n_{k,t-1} + 1}_{n'_{k,t}} S_{k,t} + \nu_{k,t}, \quad (21)$$

i.e. $n'_{k,t}$ denotes the number of stakes bidder k owns in between time $t - 1$ and t , excluding those traded in period t .

Note that $\nu_{k,t}$ will be up to bidder k to decide, as opposed to the random event $S_{k,t}$ which is exogenous; in particular, $\nu_{k,t}$ can be negative (as well as positive or zero). We will elaborate more on this below, but note that $\nu_{k,t}$ will be constrained such that after the updating in (21) $n_{k,t}$ will remain nonnegative. This way, the dynamics of the voting power $\theta_{k,t}$ in (3) and of the volume N_t in (6) will stay the same.

Let $\{P_t, t \geq 0\}$ be the price process of each (unit of) stake, which is a stochastic process assumed to be independent of the randomness induced by the $\text{Poly}(\alpha)$ voting rule (specifically, the process $\{S_{k,t}\}$). Hence, we augment the filtration $\{\mathcal{F}_t\}_{t \geq 0}$ with that of the exogenous price process $\{P_t, t \geq 0\}$ to a new filtration denoted $\{\mathcal{G}_t\}_{t \geq 0}$.

Let $b_{k,t}$ denote (units of) the risk-free asset that bidder k holds at time t , and $r_{\text{free}} > 0$ the risk-free (interest) rate. (Here, the risk-free asset is naturally the one that underlies the above price process.) As we are mainly concerned with the effect of exchanging stakes to each individual, we only allow bidders to trade stakes, but not risk-free assets between them. Hence, each bidder has to trade risk-free asset with a third-party instead of trading that with another bidder.

The decision for each bidder k at t is hence a tuple, $(\nu_{k,t}, b_{k,t})$. Moreover, there is a terminal time, denoted $T_k \in \mathbb{N}_+$ (i.e., $T_k \geq 1$ is integer valued), by which time bidder k has to sell all assets, including both any risk-free asset and any stakes owned at that time, and leave the system. T_k can either be deterministic or random. In the latter case, assume it has a finite expectation, and is either adapted to $\{\mathcal{G}_t\}_{t \geq 0}$, or independent of all other randomness

(in which case augment $\{\mathcal{G}_t\}$ accordingly). We also allow bidder k to leave the system and liquidate prior to T_k at a stopping time τ_k relative to $\{\mathcal{G}_t\}_{t \geq 0}$. Thus, bidder k will also decide at which time τ_k to stop and exit. To simplify the notation, we abuse τ_k for $\tau_k \wedge T_k$, the minimum of τ_k and T_k .

Let $c_{k,t}$ denote the (free) cash flow (or, ‘‘consumption’’) of bidder k at time t , i.e.,

$$c_{k,t} = (1 + r_{\text{free}})b_{k,t-1} - b_{k,t} - \nu_{k,t}P_t, \quad \forall 1 \leq t < \tau_k; \quad (\text{C1})$$

with

$$b_{k,0} = 0, \quad b_{k,t} \geq 0, \quad 0 \leq n_{k,t} = n'_{k,t} + \nu_{k,t} \leq N_t, \quad \forall 1 \leq t < \tau_k; \quad (\text{C2})$$

and

$$c_{k,\tau_k} = (1 + r_{\text{free}})b_{k,\tau_k-1} + n'_{k,\tau_k}P_{\tau_k}, \quad \text{and } \nu_{k,\tau_k} = b_{k,\tau_k} = 0. \quad (\text{C3})$$

Observe that the equation in (C1) simply defines what’s available for ‘‘consumption’’ in period t . It is simply an accounting or budget constraint on the cash flow. The requirements in (C2) are all in the spirit of disallowing shorting, on both components of the decision, the free asset $b_{k,t}$ and the traded stakes $\nu_{k,t}$. In particular the latter is constrained such that $\nu_{k,t} \geq -n'_{k,t}$ (following $n_{k,t} \geq 0$) i.e., bidder k cannot sell more than what’s in possession at t ; it also ensures that no bidder can own a number of stakes beyond the current total volume ($n_{k,t} \leq N_t$). (C3) specifies how the assets are liquidated at the exit time τ_k : both ν_{k,τ_k} and b_{k,τ_k} will be set at zero, and all remaining stakes n'_{k,τ_k} liquidated (cashed out at P_{τ_k} per unit).

Denote bidder k ’s decision (process) or ‘‘strategy’’ as τ_k and $(\nu, b) := \{(\nu_{k,t}, b_{k,t}), 1 \leq t \leq \tau_k\}$. The objective of bidder k is

$$U_k^* := \max_{\tau_k, (\nu, b)} U_k := \mathbb{E} \left(\sum_{t=1}^{\tau_k} \delta_k^t c_{k,t} \right), \quad \text{subject to (C1), (C2), (C3);} \quad (22)$$

where $\delta_k \in (0, 1]$ is a discount factor, a given parameter measuring the risk sensitivity of bidder k . Clearly, bidder k ’s objective is to maximize a utility that is just the present value of k ’s total cash flow cumulated up to T_k .

We need to introduce two more processes that are related and central to understanding the dynamics of the system in the presence of trading. The first one is $\{M_t, t \geq 1\}$, where $M_t := N_t P_t$ denotes the market value of the volume of stakes at time t . The second one is $\{\Pi_{k,t}, t \geq 0\}$, for each bidder k , defined as follows:

$$\Pi_{k,0} := n_{k,0}P_0, \quad \text{and } \Pi_{k,t} := \delta_k^t n'_{k,t}P_t - \sum_{j=1}^{t-1} \delta_k^j \nu_{k,j}P_j, \quad t \geq 1; \quad (23)$$

where $n'_{k,t+1}$ follows (21). Note the two terms that define $\Pi_{k,t}$ are the discounted present values, respectively, of k ’s pre-trading stakes ($n'_{k,t}$) and of the return from k ’s trading (cumulated up to $t - 1$).

The connection between $\{M_t\}$ and $\{\Pi_{k,t}\}$ is presented in the following lemma, which reveals that their incremental gains (per time period) are proportional: each increment of $\Pi_{k,t}$ is a $\pi_{k,t}$ fraction of the corresponding increment of M_t . In other words, $\pi_{k,t}$ not only represents bidder k ’s share of the total volume of stakes, it also represents k ’s share of the system’s market value, with or without trading.

Lemma 6. *Under the $\text{Poly}(\alpha)$ voting rule, along with the trading specified above, we have*

$$\mathbb{E}(\Pi_{k,t+1} | \mathcal{G}_t) - \Pi_{k,t} = \delta_k^{t+1} \pi_{k,t} \mathbb{E}(M_{t+1} | \mathcal{G}_t) - \delta_k^t \pi_{k,t} M_t. \quad (24)$$

Proof. First, by (21) and (5), along with $\pi_{k,t} = n_{k,t}/N_t$, we have

$$\mathbb{E}(n'_{k,t+1} | \mathcal{F}_t) = n_{k,t}(1 + N_t^{-(1+\alpha)}) = \frac{n_{k,t}}{N_t}(N_t + N_t^{-\alpha}) = \pi_{k,t} \mathbb{E}(N_{t+1} | \mathcal{F}_t). \quad (25)$$

Next, from (23), we have

$$\Pi_{k,t+1} - \Pi_{k,t} = \delta_k^{t+1} n'_{k,t+1} P_{t+1} - \delta_k^t n'_{k,t} P_t - \delta_k^t \nu_{k,t} P_t, \quad t \geq 1. \quad (26)$$

Furthermore, as the price process $(P_t, t \geq 0)$ is independent of \mathcal{F}_t , we have

$$\begin{aligned} \mathbb{E}(n'_{k,t+1} P_{t+1} | \mathcal{G}_t) &= \mathbb{E}(\mathbb{E}(n'_{k,t+1} | \mathcal{F}_t) P_{t+1} | \mathcal{G}_t) \\ &\stackrel{(25)}{=} \pi_{k,t} \mathbb{E}(N_{t+1} P_{t+1} | \mathcal{G}_t) = \pi_{k,t} \mathbb{E}(M_{t+1} | \mathcal{G}_t). \end{aligned}$$

This, along with (26) yields the desired expression in (24), along with $n_{k,t} = n'_{k,t} + \nu_{k,t}$, $n_{k,t} = \pi_{k,t} N_t$, and $M_t = N_t P_t$. \square

The process $\{\Pi_{k,t}\}$ also connects to the utility U_k in (22). To see this, summing up both sides of (C1) and (C3) over t (along with $b_{k,0} = 0$ in (C2)), we have

$$\sum_{t \leq \tau_k} \delta_k^t c_{k,t} = \sum_{t \leq \tau_k} \delta_k^t c_{k,t} = \delta_k^{\tau_k} n'_{\tau_k} P_{\tau_k} - \sum_{t=1}^{\tau_k-1} \delta_k^t \nu_{k,t} P_t + \sum_{t=1}^{\tau_k-1} \delta_k^t [(1 + r_{\text{free}}) \delta_k - 1] b_{k,t}. \quad (27)$$

Observe that the first two terms on the RHS are equal to Π_{k,τ_k} , so we can rewrite the above as follows (after taking expectations on both sides), emphasizing the exit time τ_k and the strategy (ν, b) ,

$$U_k(\tau_k, \nu, b) = \mathbb{E}[\Pi_{k,\tau_k}(\nu)] + \mathbb{E}\left(\sum_{t=1}^{\tau_k-1} \delta_k^t [(1 + r_{\text{free}}) \delta_k - 1] b_{k,t}\right); \quad (28)$$

hence, the RHS above is *separable*: the first term depends on (ν) only while the second term, the summation, on (b) only. Furthermore, the second term is ≤ 0 provided $(1 + r_{\text{free}}) \delta_k \leq 1$ (which is the condition (a) assumed in Theorem 7 below), along with b being non-negative, part of the feasibility in (C2). In this case, we will have $U_k \leq \mathbb{E}(\Pi_{k,\tau_k}(\nu))$, which implies $U_k^* \leq \max_{\tau_k, \nu} \mathbb{E}(\Pi_{k,\tau_k}(\nu))$, with equality holding when $b_{k,t} = 0$ for all $t = 1, \dots, \tau_k$.

We are now ready to present the main result regarding the utility maximization problem in (22). A quick word on the parameter r_{cryp} that will appear prominently in Theorem 7 below. Simply put, it is the rate (expected rate of return) associated with each stake (e.g., a unit of some cryptocurrency), i.e., it is the counterpart of r_{free} , the rate for the risk-free asset. We will elaborate more on the two rates after proving the theorem.

In the theorem, two strategies are singled out: the “no-trading” strategy, $\tau_k \geq 1$ and $\nu_{k,t} = b_{k,t} = 0$ for all $t \geq 1$, which is one instance of the class of feasible strategies; and the “non-participation” strategy, under which bidder k turns all $n_{k,0}$ stakes into cash, and then never participates in either bidding or trading for all $t \geq 1$. Also note that the non-participation strategy and the class of feasible strategies are mutually exclusive (and complementary): the former concerns $\tau_k = 0$ by abuse of language, whereas the latter are for $\tau_k \geq 1$; and any feasible strategy (ν, b) presumes participation.

Theorem 7 (Trading versus no-trading/non-participation). *Assume the following two conditions:*

$$(a) \delta_k(1 + r_{free}) \leq 1 \quad \text{and} \quad (b) \mathbf{E}(M_{t+1} | \mathcal{G}_t) = (1 + r_{cryp})M_t. \quad (29)$$

Then, under the Poly(α) voting rule, the following results will hold.

First, with condition (a), the maximal utility U_k^ is achieved by setting $b_{k,t} = 0$ for all $t = 1, \dots, T_k$; i.e., $U_k^* = \max_{\nu} \mathbf{E}(\Pi_{k,T_k})$.*

Furthermore, all three parts of the following will hold.

- (i) *If $\delta_k(1 + r_{cryp}) \leq 1$, then any feasible strategy will provide no greater utility for bidder k than the non-participation strategy, i.e., $U_k^* \leq n_{k,0}P_0$.*
- (ii) *If $\delta_k(1 + r_{cryp}) \geq 1$, then the non-participation strategy will provide no greater utility for bidder k than the no-trading strategy. In this case, bidder k will participate in the bidding process until the terminal time T_k , i.e., $\tau_k = T_k$.*
- (iii) *If $\delta_k(1 + r_{cryp}) = 1$, then, bidder k is indifferent between the non-participation and the no-trading strategy with any exit time, both of which will provide no less utility than any feasible strategy (including the optimal strategy). In other words, all strategies achieve the same utility (which is $\Pi_{k,0} = n_0P_{k,0}$).*

Moreover, when $\delta_k = \delta := (1 + r_{cryp})^{-1}$ for all k , then no bidder will have any incentive to trade. Consequently, the long-term behaviors (of N_t , $\pi_{k,t}$ and $\theta_{k,t}$) characterized in Theorems 1 and 5 will hold.

Proof. That $U_k^* = \max_{\tau_k, \nu} \mathbf{E}(\Pi_{k,\tau_k})$ (with $b_{k,t}$ being set to 0 for all t), under condition (a) in (29), has already been established in the discussions following (28). So, it suffices to prove the three parts (i)-(iii).

(i) Applying the given condition (b) in (29), along with the assumed inequality $\delta_k(1 + r_{cryp}) \leq 1$, to the RHS of the equation in (24) will make it ≤ 0 ; i.e., $\{\Pi_{k,t}\}$ is a \mathcal{G}_t -super-martingale, implying $\mathbf{E}(\Pi_{k,\tau_k}) \leq \Pi_{k,0}$. Since $\Pi_{k,0}$ is independent of ν , we have

$$U_k^* = \max_{\tau_k, \nu} \mathbf{E}(\Pi_{k,\tau_k}) \leq \Pi_{k,0} = n_{k,0}P_0, \quad (30)$$

(ii) With the assumed inequality $\delta_k(1 + r_{cryp}) \geq 1$, $\{\Pi_{k,t}\}$ now becomes a \mathcal{G}_t -sub-martingale; and hence, the inequality below,

$$\mathbf{E}(\Pi_{k,T_k}) \geq \mathbf{E}(\Pi_{k,\tau_k}) \geq \Pi_{k,0} = n_{k,0}P_0. \quad (31)$$

(Note in this case we still have $U_k^* = \max_{\nu} \mathbf{E}(\Pi_{k,T_k})$, but of course $\max_{\nu} \mathbf{E}(\Pi_{k,T_k}) \geq \mathbf{E}(\Pi_{k,T_k})$; and the optimal ν is in general not easily identified.)

(iii) Under the assumed equality $\delta_k(1 + r_{cryp}) = 1$, $\{\Pi_{k,t}\}$ is a \mathcal{G}_t -martingale; hence, the inequality in (30) now holds as equality, i.e., $U_k^* = \Pi_{k,0} = n_{k,0}P_0$. Thus, all strategies lead to the optimal utility, including any feasible strategy (in particular, the no-trading strategy) and the non-participation strategy.

The “moreover” part of the theorem is immediate. \square

In what remains of this subsection, we make a few remarks on Theorem 7, in particular, to motivate and explain its required conditions. First, the rate r_{cryp} is determined by condition (b), the second equation in (29). As such, it should be distinct from r_{free} , the latter is associated with the risk-free asset that underlies the price $\{P_t\}$. Since $M_t = N_tP_t$, and N_t is driven by very different dynamics than P_t , r_{cryp} should take a different value. For all practical

purpose, we can assume $r_{\text{crupt}} \geq r_{\text{free}}$, even though this is not assumed in the theorem. When this relation does hold, then condition (a) will become superfluous in cases (i) and (iii).

Second, the discount factor δ_k in the utility objective in (22), a parameter that measures bidder k 's sensitivity towards risk, plays a key role in characterizing the three cases. In case (i), the inequality $\delta_k \leq 1/(1 + r_{\text{cryp}})$ implies bidder k is seriously risk-averse; and this is reflected in k 's non-participation strategy. In case (ii), the inequality holds in the opposite direction, implying bidder k is lightly risk-averse or even a risk taker. Accordingly, k 's strategy is to participate until the terminal time and to pursue the optimal strategy, or short of that to follow the no-trading strategy. (Note, however, the current analysis cannot conclude whether the no-trading strategy is as good as any feasible trading strategy.) In case (iii), the inequality becomes an equality $\delta_k = 1/(1 + r_{\text{cryp}})$, and $\{\Pi_{k,t}\}$ becomes a martingale. Consequently, bidder k is indifferent between non-participation and participation, and in the latter case, indifferent to all (feasible) strategies, including the no-trading strategy. The equality also has the effect to force all participating bidders to have the same risk sensitivity.

In Roşu and Saleh (2021), there is a single rate r_{free} , or equivalently. $r_{\text{cryp}} = r_{\text{free}}$ is assumed, which seems difficult to justify, since in most applications (cryptocurrency in particular) r_{cryp} will be significantly larger than r_{free} . Moreover, there is also a single risk sensitivity for all bidders, which is set at $\delta = 1/(1 + r_{\text{free}})$. Thus, Roşu and Saleh (2021) is limited to the martingale case only, reaching the same conclusion as our case (iii), that all feasible strategies, no-trading included, yields the same (expected) utility. As there's no stopping decision and super- or sub-martingale cases in Roşu and Saleh (2021), non-anticipation does not come up at all, neither do notions like risk-averse or risk-seeking.

The third (and last) point we want to emphasize is that the two conditions in (29) play very different roles. As evident from the proof of Theorem 7, condition (b) makes $\{\Pi_{k,t}\}$ a super- or sub-martingale or a martingale, according to bidder k 's risk sensitivity as specified by the inequalities and equality applied to δ_k (along with r_{free}) in the three cases. Yet, to solve the maximization problem in (22), $\{\Pi_{k,t}\}$ needs to be connected to the utility; and this is the role played by condition (a), under which it is necessary (for optimality) to set $b_{k,t} = 0$ for all $t \geq 1$, and applicable to all three cases in Theorem (7). In this sense, condition (a) alone solves half of the maximization problem, the $b_{k,t}$ half of the strategy. In fact, it's more than half, as the optimal ν strategy is only needed in the sub-martingale case; and even there, condition (a) pins down the fact that to participate (even without trading) is better than non-participation.

Finally, let us point out that Theorem 7 can be readily extended. For instance, the rates $r_{\text{cryp}}(t)$ and $r_{\text{free}}(t)$ can vary over the time. In this case, it suffices to modify the conditions in case (i) to

$$\left(1 + \sup_{t < T_k} r_{\text{cryp}}(t)\right) \delta_k \leq 1 \quad \text{and} \quad \left(1 + \sup_{t < T_k} r_{\text{free}}(t)\right) \delta_k \leq 1,$$

the conditions in case (ii) to

$$\left(1 + \inf_{t < T_k} r_{\text{cryp}}(t)\right) \delta_k \geq 1 \quad \text{and} \quad \left(1 + \sup_{t < T_k} r_{\text{free}}(t)\right) \delta_k \leq 1,$$

and the conditions in case (iii) to

$$\delta_k = (1 + r_{\text{cryp}})^{-1} \quad \text{and} \quad \sup_{t < T_k} r_{\text{free}}(t) \leq r_{\text{cryp}}.$$

Then, Theorem 7 will continue to hold. We can also include a processing cost $\kappa > 0$ that any bidder selected by the $\text{PoLy}(\alpha)$ mechanism will pay to receive the reward. (This corresponds to the “mining” cost to validate the block.) In this case, the budget constraint (C1) is modified by adding a term $-\kappa 1_{S_{k,t}}$ to the right side of the equation, and the same applies to the liquidation constraint (with t replaced by T_k). Condition (b) in (29) is modified to $\mathbb{E}(M_{t+1} | \mathcal{G}_t) = (1 + r_{\text{cryp}})M_t + \kappa$.

4. PROOF OF TECHNICAL RESULTS

This section collects all the deferred proofs from the last two sections.

4.1. Proof of Theorem 2.

Proof. Without loss of generality, assume that $N_0 = 1$. Note that $(N_t, t \geq 0)$ has increments $\{0, 1\}$, so there are $\binom{t}{k}$ paths ending at $N_t = k + 1$ (one has to choose k upward steps “1” out of t steps). Moreover, the probability of each path ending at $N_t = k + 1$ is upper bounded by

$$\frac{1}{(k!)^\alpha} \left(1 - \frac{1}{(k+1)^\alpha}\right)^{t-k},$$

since the k upward “1” steps contribute $1/k!$, and the remaining $t - k$ flat “0” steps have at most probability $\left(1 - \frac{1}{(k+1)^\alpha}\right)^{t-k}$. Thus,

$$\mathbb{P}(N_t \leq m + 1) \leq \sum_{k \leq m} a_k \quad \text{and} \quad \mathbb{P}(N_t > m) \leq \sum_{k \geq m} a_k$$

where

$$a_k := \binom{t}{k} \frac{1}{(k!)^\alpha} \left(1 - \frac{1}{(k+1)^\alpha}\right)^{t-k}. \quad (32)$$

As we will see, a_k is nondecreasing on $[1, \lambda_-(\alpha) t^{\frac{1}{1+\alpha}}]$ and nonincreasing on $(\lambda_+(\alpha) t^{\frac{1}{1+\alpha}}, t]$. The idea is to study the term $a_{\lfloor \lambda t^{\frac{1}{1+\alpha}} \rfloor}$ for $\lambda > 0$ as $t \rightarrow \infty$. By Stirling’s formula,

$$\binom{t}{\lfloor \lambda t^{\frac{1}{1+\alpha}} \rfloor} \sim \frac{1}{\sqrt{2\pi\lambda}} t^{-\frac{1}{2(1+\alpha)}} \exp\left(\frac{\lambda\alpha}{1+\alpha} t^{\frac{1}{1+\alpha}} \log t + (\lambda - \lambda \log \lambda) t^{\frac{1}{1+\alpha}} + o\left(t^{\frac{1}{1+\alpha}}\right)\right),$$

$$\left(\lfloor \lambda t^{\frac{1}{1+\alpha}} \rfloor\right)! \sim \sqrt{2\pi\lambda} t^{\frac{1}{2(1+\alpha)}} \exp\left(\frac{\lambda}{1+\alpha} t^{\frac{1}{1+\alpha}} \log t + (\lambda \log \lambda - \lambda) t^{\frac{1}{1+\alpha}}\right),$$

and

$$\left(1 - \frac{1}{\lambda^\alpha t^{\frac{\alpha}{1+\alpha}}}\right)^{t - \lambda t^{\frac{1}{1+\alpha}}} = \exp\left(-\frac{t^{\frac{1}{1+\alpha}}}{\lambda^\alpha} + o\left(t^{\frac{1}{1+\alpha}}\right)\right).$$

Combining the above estimates yields

$$a_{\lfloor \lambda \sqrt{t} \rfloor} \sim (2\pi\lambda)^{-\frac{1+\alpha}{2}} t^{-\frac{1}{2}} \exp\left(-f_\alpha(\lambda) t^{\frac{1}{1+\alpha}}\right). \quad (33)$$

Note that the function $\lambda \rightarrow f_\alpha(\lambda)$ has two roots $\lambda_-(\alpha) < \lambda_+(\alpha)$ on $[0, \infty)$, and $f_\alpha > 0$ on $(0, \lambda_-(\alpha)) \cup (\lambda_+(\alpha), \infty)$. As a result, for each $\lambda < \lambda_-(\alpha)$ we have

$$\mathbb{P}(N_t < \lambda t^{\frac{1}{1+\alpha}}) \leq C_\lambda t^{-\frac{1}{2} + \frac{1}{1+\alpha}} \exp\left(-f_\alpha(\lambda) t^{\frac{1}{1+\alpha}}\right) \leq \exp\left(-(1-\varepsilon)f_\alpha(\lambda) t^{\frac{1}{1+\alpha}}\right) \text{ as } t \rightarrow \infty,$$

and each $\lambda > \lambda_+(\alpha)$ we have

$$\mathbb{P}(N_t > \lambda t^{\frac{1}{1+\alpha}}) \leq C'_\lambda t^{\frac{1}{2}} \exp\left(-f_\alpha(\lambda) t^{\frac{1}{1+\alpha}}\right) \leq \exp\left(-(1-\varepsilon)f_\alpha(\lambda) t^{\frac{1}{1+\alpha}}\right) \text{ as } t \rightarrow \infty.$$

□

4.2. Proof of Corollary 3.

Proof. By Theorem 1 (i), we get $((1+\alpha)t)^{\frac{\alpha}{1+\alpha}} N_t^{-\alpha}$ converges in probability to 1. Fix $\ell > 0$ to be sufficiently small. By Proposition 2 (1), we get for $r > 1$,

$$\begin{aligned} & \mathbb{E} \left(\left(((1+\alpha)t)^{\frac{\alpha}{1+\alpha}} N_t^{-\alpha} \right)^{1+r} \right) \\ &= \mathbb{E} \left(\left(((1+\alpha)t)^{\frac{\alpha}{1+\alpha}} N_t^{-\alpha} \right)^{1+r} 1_{\{N_t \leq \ell t^{\frac{1}{1+\alpha}}\}} \right) + \mathbb{E} \left(\left(((1+\alpha)t)^{\frac{\alpha}{1+\alpha}} N_t^{-\alpha} \right)^{1+r} 1_{\{N_t > \ell t^{\frac{1}{1+\alpha}}\}} \right) \\ &\leq ((1+\alpha)t)^{\frac{(1+r)\alpha}{1+\alpha}} e^{-Ct^{\frac{1}{1+\alpha}}} + ((1+\alpha)t)^{\frac{\alpha}{1+\alpha}} \ell^{-\alpha} t^{1+r}, \end{aligned}$$

which implies that $\limsup_t \mathbb{E} \left(((1+\alpha)t)^{\frac{\alpha}{1+\alpha}} N_t^{-\alpha} \right) < \infty$, and hence, $((1+\alpha)t)^{\frac{\alpha}{1+\alpha}} N_t^{-\alpha}$ is uniformly integrable. Consequently, $\lim_{t \rightarrow \infty} \mathbb{E} \left(((1+\alpha)t)^{\frac{\alpha}{1+\alpha}} N_t^{-\alpha} \right) = 1$ and

$$\mathbb{E}(N_t^{-\alpha}) \sim ((1+\alpha)t)^{-\frac{\alpha}{1+\alpha}} \text{ as } t \rightarrow \infty. \quad (34)$$

Similarly, we have

$$\mathbb{E}(N_t^{1-\alpha}) \sim ((1+\alpha)t)^{\frac{1-\alpha}{1+\alpha}} \text{ as } t \rightarrow \infty. \quad (35)$$

Note that

$$\mathbb{E}(N_t) = \sum_{k \leq t} \mathbb{E}(N_k^{-\alpha}) \quad \text{and} \quad \mathbb{E}(N_t^2) = 2 \sum_{k \leq t} \mathbb{E}(N_k^{1-\alpha}) + \sum_{k \leq t} \mathbb{E}(N_k^{-\alpha}). \quad (36)$$

Combining (34), (35) and (36) yields $\text{Var}(N_t) = o\left(t^{\frac{2}{1+\alpha}}\right)$ as $t \rightarrow \infty$. The bound (12) then follows from Chebyshev's inequality. □

4.3. Proof of Proposition 4.

Proof. Fix $T > 0$. It suffices to prove the weak convergence (18) on $[0, T]$. (Refer to (Billingsley, 1999, Chapter 2) for background on the weak convergence in $\mathcal{C}[0, \infty)$.) By Theorem 1 (i), there is the convergence in probability $N_{[nT]}/n^{\frac{1}{1+\alpha}} \rightarrow X_T$ as $n \rightarrow \infty$. Given $\varepsilon > 0$, there is $n(\varepsilon) > 0$ such that for any $n > n(\varepsilon)$,

$$\mathbb{P} \left(N_{[nT]}/n^{\frac{1}{1+\alpha}} < 2X_T \right) > 1 - \varepsilon.$$

Let $K(\varepsilon) := \max(2X_T, \max_{n \leq n(\varepsilon)} (N_0 + [nT])/n^{\frac{1}{1+\alpha}})$. We have $\mathbb{P}(N_{[nT]}/n^{\frac{1}{1+\alpha}} < K(\varepsilon)) > 1 - \varepsilon$ for each $n \in \mathbb{N}_+$. Note that for each $n \in \mathbb{N}_+$, the process $N^{n,T} := \left(N_{[nT]}/n^{\frac{1}{1+\alpha}}, 0 \leq t \leq T \right)$ is nondecreasing. Thus,

$$\mathbb{P}(N^{n,T} \in [0, T] \times [0, K(\varepsilon)]) > 1 - \varepsilon.$$

So the sequence of processes $(N^{n,T}, n \in \mathbb{N}_+)$ is tight. Moreover, for each $t \in [0, T]$, $N_{[nt]}/n^{\frac{1}{1+\alpha}}$ converges in probability to X_t as $n \rightarrow \infty$. Then for $0 \leq t_1 < \dots < t_k$, the vector $(N_{[nt_1]}/n^{\frac{1}{1+\alpha}}, \dots, N_{[nt_k]}/n^{\frac{1}{1+\alpha}})$ converges in probability to $(X_{t_1}, \dots, X_{t_k})$, i.e. the convergence in finite-dimensional distributions. The weak convergence follows readily from the tightness and the convergence in finite-dimensional distributions (see e.g. (Billingsley, 1999, Chapter 2)). \square

4.4. Proof of Theorem 5. The proof breaks into a couple of lemmas. The first one (below) establishes algebraic expressions for the moments of $\pi_{k,t}$.

Lemma 8. *Under the $\text{Poly}(\alpha)$ voting rule, we have:*

(i) For each $t \geq 0$,

$$\text{Var}(\pi_{k,t+1}) = \text{Var}(\pi_{k,t}) + \mathbb{E} \left(\frac{\pi_{k,t}(1 - \pi_{k,t})}{N_t^\alpha (N_t + 1)^2} \right). \quad (37)$$

(ii) Let $\mu_3(\pi_{k,t}) := \mathbb{E}(\pi_{k,t} - \pi_{k,0})^3$ be the third central moment of $\pi_{k,t}$. For each $t \geq 0$,

$$\mu_3(\pi_{k,t+1}) = \underbrace{3 \mathbb{E} \left(\frac{\pi_{k,t}(1 - \pi_{k,t})(1 - 2\pi_{k,t})}{(1 + N_t)^3 N_t^\alpha} \right)}_{(a)} + \underbrace{3 \mathbb{E} \left(\frac{(\pi_{k,t} - \pi_{k,0})\pi_{k,t}(1 - \pi_{k,t})}{(1 + N_t)^2 N_t^\alpha} \right)}_{(b)}. \quad (38)$$

(iii) Let $\mu_4(\pi_{k,t}) := \mathbb{E}(\pi_{k,t} - \pi_{k,0})^4$ be the fourth central moment of $\pi_{k,t}$. For each $t \geq 0$,

$$\begin{aligned} \mu_4(\pi_{k,t+1}) &= \mu_4(\pi_{k,t}) + \underbrace{\mathbb{E} \left(\frac{\pi_{k,t}^4(1 - \pi_{k,t})}{(1 + N_t)^4 N_t^\alpha} \right)}_{(a')} + \underbrace{\mathbb{E} \left(\frac{\pi_{k,t}(1 - \pi_{k,t})^4}{(1 + N_t)^4 N_t^\alpha} \right)}_{(b')} \\ &\quad + \underbrace{6 \mathbb{E} \left((\pi_{k,t} - \pi_{k,0})^2 \frac{\pi_{k,t}^2(1 - \pi_{k,t})}{(1 + N_t)^2 N_t^\alpha} \right)}_{(c')} + \underbrace{6 \mathbb{E} \left((\pi_{k,t} - \pi_{k,0})^2 \frac{\pi_{k,t}(1 - \pi_{k,t})^2}{(1 + N_t)^2 N_t^\alpha} \right)}_{(d')} \\ &\quad - \underbrace{4 \mathbb{E} \left((\pi_{k,t} - \pi_{k,0}) \frac{\pi_{k,t}^3(1 - \pi_{k,t})}{(1 + N_t)^3 N_t^\alpha} \right)}_{(e')} + \underbrace{4 \mathbb{E} \left((\pi_{k,t} - \pi_{k,0}) \frac{\pi_{k,t}(1 - \pi_{k,t})^3}{(1 + N_t)^3 N_t^\alpha} \right)}_{(f')}. \end{aligned} \quad (39)$$

Proof. By the definitions in (2) and (5), we can write

$$\mathbb{E}(\pi_{k,t+1} | \mathcal{F}_t) = \frac{n_{k,t}}{N_t} 1_{S_1} + \frac{n_{k,t}}{1 + N_t} 1_{S_2} + \frac{1 + n_{k,t}}{1 + N_t} 1_{S_3},$$

where S_1, S_2, S_3 are mutually exclusive events with probabilities $P(S_1) = 1 - N_t^{-\alpha}$, $P(S_2) = \frac{1 - \pi_{k,t}}{N_t^\alpha}$ and $P(S_3) = \frac{\pi_{k,t}}{N_t^\alpha}$. Replacing 1_{S_1} with $1 - 1_{S_2} - 1_{S_3}$, we get

$$\mathbb{E}(\pi_{k,t+1} | \mathcal{F}_t) = \pi_{k,t} - \frac{\pi_{k,t}}{1 + N_t} 1_{S_2} + \frac{1 - \pi_{k,t}}{1 + N_t} 1_{S_3}.$$

Noting that $\text{Var}(\pi_{k,t+1}) = \mathbb{E} \left((\mathbb{E}(\pi_{k,t} | \mathcal{F}_t) - \pi_{k,0})^2 \right)$ and $\mu_m(\pi_{k,t+1}) = \mathbb{E} \left((\mathbb{E}(\pi_{k,t} | \mathcal{F}_t) - \pi_{k,0})^m \right)$ ($m = 3, 4$), and using the multinomial expansion yields the formulas (37) – (39). \square

Relying on the recursions in Lemma 8, we give some estimates for the moments of $\pi_{k,t}$ in the next lemma.

Lemma 9. *Under the $\text{Poly}(\alpha)$ voting rule, we have:*

(i) *There is $C > 0$ such that*

$$\text{Var}(\pi_{k,t}) \leq \frac{C\pi_{k,0}}{N} \quad \text{for all } t \geq 1. \quad (40)$$

Assume further that $\pi_{k,0} = \mathcal{O}(N^{-1})$. There is $c > 0$ such that

$$\text{Var}(\pi_{k,\infty}) \geq \frac{c\pi_{k,0}}{N} \quad \text{for all } t \geq 1. \quad (41)$$

(ii) *Assume that $\pi_{k,0} = \mathcal{O}(N^{-1})$. There is $C > 0$ such that*

$$\mu_3(\pi_{k,t}) \leq \frac{C\pi_{k,0}}{N^2} \quad \text{for all } t \geq 1. \quad (42)$$

(iii) *Assume that $\pi_{k,0} = \mathcal{O}(N^{-1})$. There is $C > 0$ such that*

$$\mu_4(\pi_{k,t}) \leq \frac{C\pi_{k,0}}{N^3} \quad \text{for all } t \geq 1. \quad (43)$$

Proof. (i) Since $0 \leq \pi_{k,t} \leq 1$, we have

$$\mathbb{E} \left(\frac{\pi_{k,t}(1 - \pi_{k,t})}{N_t^\alpha (N_t + 1)^2} \right) \leq \mathbb{E} \left(\frac{\pi_{k,t}}{N_t^{2+\alpha}} \right). \quad (44)$$

The same argument as in Proposition 2 shows that there are $\ell, \theta > 0$ such that for N sufficiently large,

$$\mathbb{P} \left(N_t \leq \ell t^{\frac{1}{1+\alpha}} \right) \leq \exp(-Ct^{\frac{1}{1+\alpha}}) \quad \text{for some } C > 0 \text{ and } t > \theta N^{1+\alpha}. \quad (45)$$

Let $\Omega_{\ell,t} := \{N_t > \ell t^{\frac{1}{1+\alpha}}\}$. By (45), we have for $t > \theta N^{1+\alpha}$, $\mathbb{P}(\Omega_{\ell,t}^c) \leq \exp(-Ct^{\frac{1}{1+\alpha}})$ and hence,

$$\begin{aligned} \mathbb{E} \left(\frac{\pi_{k,t}}{N_t^{2+\alpha}} \right) &= \mathbb{E} \left(\frac{\pi_{k,t}}{N_t^{2+\alpha}} 1_{\Omega_{\ell,t}} \right) + \mathbb{E} \left(\frac{\pi_{k,t}}{N_t^{2+\alpha}} 1_{\Omega_{\ell,t}^c} \right) \\ &\leq C' \pi_{k,0} t^{-\frac{2+\alpha}{1+\alpha}} + \exp(-Ct^{\frac{1}{1+\alpha}}). \end{aligned} \quad (46)$$

By (37) and (44), we get

$$\begin{aligned} \text{Var}(\pi_{k,t}) &\leq \sum_{j<t} \mathbb{E} \left(\frac{\pi_{k,j}}{N_j^{2+\alpha}} \right) \\ &= \sum_{j \leq \theta N^{1+\alpha}} \mathbb{E} \left(\frac{\pi_{k,j}}{N_j^{2+\alpha}} \right) + \sum_{\theta N^{1+\alpha} < j < t} \mathbb{E} \left(\frac{\pi_{k,j}}{N_j^{2+\alpha}} \right). \end{aligned} \quad (47)$$

The first term in (47) is easily bounded from above by $\theta\pi_{k,0}/N$. By (46) and the sum-integral trick, the second term in (47) is bounded by $C''\pi_{k,0}(N^{1+\alpha})^{-\frac{1}{1+\alpha}} = C''\pi_{k,0}/N$. We have proved (40). Assuming further that $\pi_{k,0} = \mathcal{O}(N^{-1})$, we have

$$\mathbb{E}\pi_{k,t}^2 = \pi_{k,0}^2 + \text{Var}(\pi_{k,t}) \leq \pi_{k,0}^2 + \frac{C'\pi_{k,0}}{N} \leq \frac{C'\pi_{k,0}}{N}. \quad (48)$$

As a result, $\mathbb{E}(\pi_{k,t}(1 - \pi_{k,t})) \geq c\pi_{k,0}$ for some $c > 0$. Again the same argument as in Proposition 2 shows that there are $L, \eta > 0$ such that for N sufficiently large,

$$\mathbb{P} \left(N_t \geq Lt^{\frac{1}{1+\alpha}} \right) \leq \exp(-Ct^{\frac{1}{1+\alpha}}) \quad \text{for some } C > 0 \text{ and } t > \eta N^{1+\alpha}. \quad (49)$$

Let $\Omega'_{L,t} := \{N_t \leq Lt^{\frac{1}{1+\alpha}}\}$. By (49), we have for $t > \eta N^{1+\alpha}$, $\mathbb{P}(\Omega'_{L,t}) \geq 1 - \exp(-Ct^{\frac{1}{1+\alpha}})$ and hence,

$$\mathbb{E} \left(\frac{\pi_{k,t}(1 - \pi_{k,t})}{N_t^\alpha (N_t + 1)^2} \right) \geq \mathbb{E} \left(\frac{\pi_{k,t}(1 - \pi_{k,t})}{N_t^\alpha (N_t + 1)^2} 1_{\Omega'_{L,t}} \right) \geq c'\pi_{k,0}t^{-\frac{2+\alpha}{1+\alpha}}$$

Therefore, $\text{Var}(\pi_{k,\infty}) \geq \sum_{j > \eta N^{1+\alpha}} \mathbb{E} \left(\frac{\pi_{k,j}(1 - \pi_{k,j})}{N_j^\alpha (N_j + 1)^2} \right) \geq c''\pi_{k,0}(N^{1+\alpha})^{-\frac{1}{1+\alpha}} = c''\pi_{k,0}/N$.

(ii) Recall the terms (a)–(b) in (38). It is easy to see that

$$(a) \leq 3\mathbb{E} \left(\frac{\pi_{k,t}}{N_t^{3+\alpha}} \right) \quad \text{and} \quad (b) \leq 3\mathbb{E} \left(\frac{\pi_{k,t}^2}{N_t^{2+\alpha}} \right).$$

The same truncation and sum-integral trick as in (1) shows that

$$\sum_{j<t} \mathbb{E} \left(\frac{\pi_{k,j}}{N_j^{3+\alpha}} \right) \leq \frac{C\pi_{k,0}}{N^2} \quad \text{and} \quad \sum_{j<t} \mathbb{E} \left(\frac{\pi_{k,t}^2}{N_t^{2+\alpha}} \right) \stackrel{(48)}{\leq} \frac{C'\pi_{k,0}}{N^2},$$

which leads to the bound (42).

(iii) By (42) and the fact that $\pi_{k,0} = \mathcal{O}(N^{-1})$, we have

$$\mathbb{E}\pi_{k,t}^3 \leq C \left(\frac{\pi_{k,0}}{N^2} + \frac{\pi_{k,0}^2}{N} + \pi_{k,0}^3 \right) \leq \frac{C'\pi_{k,0}}{N^2}. \quad (50)$$

Recall the terms (a')–(f') in (39). We have

$$(a') + (b') \leq C\mathbb{E} \left(\frac{\pi_{k,t}}{N_t^{4+\alpha}} \right), \quad (c') + (d') \leq C'\mathbb{E} \left(\frac{\pi_{k,t}^3}{N_t^{2+\alpha}} \right), \quad (e') + (f') \leq C''\mathbb{E} \left(\frac{\pi_{k,t}^2}{N_t^{3+\alpha}} \right).$$

Again using the truncation and the sum integral trick, we obtain

$$\sum_{j<t} \mathbb{E} \left(\frac{\pi_{k,j}}{N_j^{4+\alpha}} \right) \leq \frac{C\pi_{k,0}}{N^3}, \quad \sum_{j<t} \mathbb{E} \left(\frac{\pi_{k,j}^3}{N_j^{2+\alpha}} \right) \stackrel{(50)}{\leq} \frac{C'\pi_{k,0}}{N^3}, \quad \sum_{j<t} \mathbb{E} \left(\frac{\pi_{k,j}^2}{N_j^{3+\alpha}} \right) \stackrel{(48)}{\leq} \frac{C''\pi_{k,0}}{N^3}.$$

The above estimates yield the bound (43). \square

Proof of Theorem 5. (i) By (40), we have $\text{Var}(\pi_{k,t}) \leq C\pi_{k,0}/N$ for some $C > 0$. By Chebyshev's inequality, we get

$$\mathbb{P}\left(\left|\frac{\pi_{k,t}}{\pi_{k,0}} - 1\right| > \varepsilon\right) \leq \frac{\text{Var}(\pi_{k,t})}{\varepsilon^2\pi_{k,0}^2} \leq \frac{C}{\varepsilon^2 N \pi_{k,0}},$$

which yields (19).

(ii) & (iii) Again by (40), we get

$$\text{Var}\left(\frac{\pi_{k,\infty}}{\pi_{k,0}}\right) = \frac{\text{Var}(\pi_{k,\infty})}{\pi_{k,0}^2} \leq \frac{C}{N\pi_{k,0}} = \frac{C}{n_{k,0}},$$

which is of order $\Theta(1)$ if $n_{k,0} = \Theta(1)$, and explodes if $n_{k,0} = o(1)$. The proof of (20) is the same as (Tang, 2022, (2.13)) given the estimates in Lemma 9. For the sake of completeness, we include the proof. For $n_{k,0} = \Theta(1)$, we deduce from the lower bound (41) that there exists $c > 0$ independent of N such that $\text{Var}\left(\frac{\pi_{k,\infty}}{\pi_{k,0}}\right) \geq c$. For $\varepsilon > 0$, we have

$$\begin{aligned} c \leq \text{Var}\left(\frac{\pi_{k,\infty}}{\pi_{k,0}}\right) &= \mathbb{E}\left(\left(\frac{\pi_{k,\infty}}{\pi_{k,0}} - 1\right)^2 \mathbb{1}_{\left\{\left|\frac{\pi_{k,\infty}}{\pi_{k,0}} - 1\right| > \varepsilon\right\}}\right) + \mathbb{E}\left(\left(\frac{\pi_{k,\infty}}{\pi_{k,0}} - 1\right)^2 \mathbb{1}_{\left\{\left|\frac{\pi_{k,\infty}}{\pi_{k,0}} - 1\right| \leq \varepsilon\right\}}\right) \\ &\leq \sqrt{\mu_4\left(\frac{\pi_{k,\infty}}{\pi_{k,0}}\right)} \mathbb{P}\left(\left|\frac{\pi_{k,\infty}}{\pi_{k,0}} - 1\right| > \varepsilon\right) + \varepsilon^2. \end{aligned}$$

By the upper bound (43), we get $\mu_4\left(\frac{\pi_{k,\infty}}{\pi_{k,0}}\right) \leq \frac{C_4}{n_{k,0}^3} := C$. Thus, for $\varepsilon < \sqrt{c}$,

$$\mathbb{P}\left(\left|\frac{\pi_{k,\infty}}{\pi_{k,0}} - 1\right| > \varepsilon\right) \geq \frac{(c - \varepsilon^2)^2}{C},$$

which leads to the desired result. \square

5. CONCLUSION

We have proposed in this study a new $\text{Poly}(\alpha)$ voting rule that is more general than the traditional voting rule (which is linear, corresponding to $\alpha = 0$). More importantly, the $\text{Poly}(\alpha)$ voting rule distinguishes voting power from voter share, and hence decouples the two.

Applying the $\text{Poly}(\alpha)$ voting rule to the PoS protocol, where the voters are the bidders (competing for the validation of a new block), we show this decoupling will enhance decentralization, a central tenet of the PoS protocol. Specifically, we prove that while bidder shares form a martingale process, each bidder's voting power is a super-martingale that decreases to zero over time. Furthermore, we show a phase transition in the stability of bidder shares in terms of each bidder's initial share relative to the total in the system. We also study the issue of a bidder's risk sensitivity when trading is allowed, and provide conditions under which a bidder will have no incentive to participate in the bidding process, or, if participate, will forgo trading.

Throughout this paper we have assumed there is a constant (one unit) reward every period. Hence, a natural next step is to consider more general reward schemes. Another possible extension is to allow the number of voters/bidders (which is a constant K here) to grow over time. In fact, in the case of $\alpha = 0$, i.e., in the setting of Pólya's urn, Tang (2022) has studied both features. For instance, the phase transition phenomenon brought forth in Tang (2022) (which is also present under the $\text{Poly}(\alpha)$ voting rule as we have shown in §3.2 here) holds not only for constant reward but also for rewards that are decreasing over time.

Acknowledgement: W. Tang gratefully acknowledges financial support through an NSF grant DMS-2113779 and through a start-up grant at Columbia University.

APPENDIX. IMPROVEMENT OF $\lambda_{\pm}(\alpha)$ IN THEOREM 2

Here we provide a way to slightly improve the values of $\lambda_{\pm}(\alpha)$ in Proposition 2. To simplify the presentation, we consider $\alpha = 1$ (quadratic voting rules) with $\lambda_-(1) \approx 0.56$ and $\lambda_+(1) \approx 2.51$.

The idea relies on a multi-scale analysis by splitting the interval $[0, t]$ into $[0, t/2]$ and $[t/2, t]$. Our goal is to upper bound $\mathbb{P}(N_t = \lambda\sqrt{t})$ for $\lambda > 0$. In the sequel, we neglect the polynomial factors and only focus on the exponential terms. Note that

$$\mathbb{P}(N_t = \lambda\sqrt{t}) = \sum_{k \leq \lambda\sqrt{t}} \binom{t/2}{k} \binom{t/2}{\lambda\sqrt{t} - k} \frac{1}{(\lambda\sqrt{t})!} \underbrace{\left(1 - \frac{1}{k}\right)^{t/2-k} \left(1 - \frac{1}{\lambda\sqrt{t}}\right)^{t/2+k-\lambda\sqrt{t}}}_{(a'')}.$$

Next we split the range of $k \leq \lambda\sqrt{t}$ into $S_1 := \{k \leq a\sqrt{t}\} \cup \{k \geq (\lambda - a)\sqrt{t}\}$, and $S_2 := \{a\sqrt{t} < k < (\lambda - a)\sqrt{t}\}$ with $a < \frac{\lambda}{2}$. For $k \in S_1$, we simply bound the term (a) by $\left(1 - \frac{1}{\lambda\sqrt{t}}\right)^{t/2-\lambda\sqrt{t}}$ while for $k \in S_2$ we bound the term (a'') by

$$\left(1 - \frac{1}{(\lambda - a)\sqrt{t}}\right)^{t/2-(\lambda-a)\sqrt{t}} \left(1 - \frac{1}{\lambda\sqrt{t}}\right)^{t/2-a\sqrt{t}}.$$

Consequently,

$$\begin{aligned} \mathbb{P}(N_t = \lambda\sqrt{t}) &\leq \underbrace{\left(\sum_{k \in S_1} \binom{t/2}{k} \binom{t/2}{\lambda\sqrt{t} - k}\right) \frac{1}{(\lambda\sqrt{t})!} \left(1 - \frac{1}{\lambda\sqrt{t}}\right)^{t/2-\lambda\sqrt{t}}}_{(b'')} \\ &+ \underbrace{\left(\sum_{k \in S_2} \binom{t/2}{k} \binom{t/2}{\lambda\sqrt{t} - k}\right) \frac{1}{(\lambda\sqrt{t})!} \left(1 - \frac{1}{(\lambda - a)\sqrt{t}}\right)^{t/2-(\lambda-a)\sqrt{t}} \left(1 - \frac{1}{\lambda\sqrt{t}}\right)^{t/2-a\sqrt{t}}}_{(c'')}. \end{aligned}$$

Recall that

$$\binom{\alpha t}{\beta\sqrt{t}} \sim \exp\left(\frac{\beta}{2}\sqrt{t} \log t + (\beta \log \alpha + \beta - \beta \log \beta)\sqrt{t}\right).$$

Thus,

$$\sum_{k \in \mathcal{S}_1} \binom{t/2}{k} \binom{t/2}{\lambda\sqrt{t}-k} \sim \exp\left(\frac{\lambda}{2}\sqrt{t}\log t + (-\lambda\log 2 + \lambda - a\log a - (\lambda - a)\log(\lambda - a))\sqrt{t}\right),$$

$$\sum_{k \in \mathcal{S}_2} \binom{t/2}{k} \binom{t/2}{\lambda\sqrt{t}-k} \sim \exp\left(\frac{\lambda}{2}\sqrt{t}\log t + (\lambda - \lambda\log \lambda)\sqrt{t}\right).$$

As a result, we get exponential bounds for the terms (b'') and (c''):

$$(b'') \sim \exp\left(\left(-\lambda\log 2 + 2\lambda - a\log a - (\lambda - a)\log(\lambda - a) - \lambda\log \lambda - \frac{1}{\lambda}\right)\sqrt{t}\right),$$

$$(c'') \sim \exp\left(\left(2\lambda - 2\lambda\log \lambda - \frac{1}{2\lambda} - \frac{1}{2(\lambda - a)}\right)\sqrt{t}\right).$$
(51)

By equating the two coefficients before \sqrt{t} in (51) we have

$$-\lambda\log 2 + 2\lambda - a\log a - (\lambda - a)\log(\lambda - a) - \lambda\log \lambda - \frac{1}{\lambda} = 2\lambda - 2\lambda\log \lambda - \frac{1}{2\lambda} - \frac{1}{2(\lambda - a)}.$$

By letting $a = \theta\lambda$ with $\theta < \frac{1}{2}$, the above equation yields

$$\lambda = \sqrt{\frac{\theta}{2(1-\theta)(\log 2 + \theta\log \theta + (1-\theta)\log(1-\theta))}}.$$
(52)

and the coefficient before \sqrt{t} is

$$f(\lambda) = 2\lambda\log \lambda - 2\lambda + \frac{1}{2\lambda} + \frac{1}{2(1-\theta)\lambda},$$
(53)

where θ is specified by (52). By injecting the expression (52) into (53), f is a function of θ . It is easy to see that $f(\theta)$ has only one root on $(0, 1/2)$ which is approximately 0.1575, and $\lambda_-(1)$ is improved numerically from 0.56 to 0.60. Similarly, the value of $\lambda_+(1)$ is improved numerically from 2.51 to 2.44.

We can continue this procedure, for instance to split $[0, t]$ into $[0, t/3]$, $[t/3, 2t/3]$ and $[2t/3, t]$...to get better and better numerical values of $\lambda_-(1)$ and $\lambda_+(1)$. However, it is not clear whether this approach will eventually get all the way to the threshold $\sqrt{2} \approx 1.41$.

We conjecture that the exponential deviation holds right off the threshold $(1 + \alpha)^{\frac{1}{1+\alpha}}$, which is supported by the numerical experiments; refer to Figure 1.

REFERENCES

- M. Armbrust, A. Fox, R. Griffith, A. D. Joseph, R. H. Katz, A. Konwinski, G. Lee, D. A. Patterson, A. Rabkin, and I. Stoica. Above the clouds: A Berkeley view of cloud computing. 2009. Technical Report UCB/EECS-2009-28. Available at <https://www2.eecs.berkeley.edu/Pubs/TechRpts/2009/EECS-2009-28.pdf>.
- P. Billingsley. *Probability and measure*. Wiley Series in Probability and Mathematical Statistics. John Wiley & Sons, Inc., New York, third edition, 1995.
- P. Billingsley. *Convergence of probability measures*. Wiley Series in Probability and Statistics: Probability and Statistics. John Wiley & Sons, Inc., New York, second edition, 1999.




FIGURE 1. Volume of stakes N_t with $N_0 = 5$ and $\alpha = 1$ (quadratic voting).

- H. Chen and D. D. Yao. *Fundamentals of queueing networks*, volume 46 of *Applications of Mathematics*. Springer-Verlag, 2001.
- J. Dean and S. Ghemawat. Mapreduce: simplified data processing on large clusters. *Commun. ACM*, 51(1):107–113, 2008.
- D. Denisov, D. Korshunov, and V. Wachtel. At the edge of criticality: Markov chains with asymptotically zero drift. 2016. arXiv:1612.01592.
- R. Durrett. *Probability—theory and examples*. Cambridge University Press, 2019.
- H. Garcia-Molina. Elections in a distributed computing system. *IEEE Trans. Comput.*, 31(01):48–59, 1982.
- A. Q. Huang and J. Baliga. FREEDM System: Role of power electronics and power semiconductor devices in developing an energy internet. In *21st International Symposium on Power Semiconductor Devices & IC's*, pages 9–12, 2009.
- S. King and S. Nadal. Ppcoin: Peer-to-peer crypto-currency with proof-of-stake. 2012. Available at <https://decred.org/research/king2012.pdf>.
- S. P. Lalley and E. G. Weyl. Quadratic voting: How mechanism design can radicalize democracy. In *AEA Papers and Proceedings*, volume 108, pages 33–37, 2018.
- J. Lamperti. Criteria for the recurrence or transience of stochastic process. I. *J. Math. Anal. Appl.*, 1:314–330, 1960.
- J. Lamperti. A new class of probability limit theorems. *J. Math. Mech.*, 11:749–772, 1962.
- J. Lamperti. Criteria for stochastic processes. II. Passage-time moments. *J. Math. Anal. Appl.*, 7:127–145, 1963.
- L. Lamport, R. Shostak, and M. Pease. The Byzantine Generals Problem. *ACM Transactions on Programming Languages and Systems*, 4(3):382–401, 1982.
- M. Menshikov, S. Popov, and A. Wade. *Non-homogeneous random walks*, volume 209 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2017. Lyapunov function methods for near-critical stochastic systems.
- S. Nakamoto. Bitcoin: A peer-to-peer electronic cash system. *Decentralized Business Review*, page 21260, 2008.
- L. S. Penrose. The elementary statistics of majority voting. *J. R. Stat. Soc.*, 109(1):53–57, 1946.

- I. Roşu and F. Saleh. Evolution of shares in a proof-of-stake cryptocurrency. *Manag. Sci.*, 67(2):661–672, 2021.
- W. Tang. Stability of shares in the Proof of Stake protocol – concentration and phase transitions. 2022. arXiv:2206.02227.
- G. Wood. Ethereum: A secure decentralised generalised transaction ledger. *Ethereum Project Yellow Paper*, 151:1–32, 2014.

DEPARTMENT OF INDUSTRIAL ENGINEER AND OPERATIONS RESEARCH, COLUMBIA UNIVERSITY.
Email address: wt2319@columbia.edu

DEPARTMENT OF INDUSTRIAL ENGINEER AND OPERATIONS RESEARCH, COLUMBIA UNIVERSITY.
Email address: yao@columbia.edu