

Quantum foam Cosmology

-using a class of smooth 3+1 spacetimes to measure the immeasurable

Claes Cramer

YBC—Young Business Creatives, upper secondary school, Experimentgatan 7, 131 54 Nacka, Sweden

E-mail: claes.cramer@utb.nacka.se

Abstract

To understand the fabric of spacetime we can search for inspiration in Wheeler's layman's analogy of the Quantum foam using an ocean surface observed from an aeroplane at different heights. From a 3+1 spacetime perspective, the ocean surface represents the hypersurfaces of spacetime at different times. Therefore, the ocean surface observed from a great height, when it appears smooth, homogenous, isotropic and flat is like looking at the current universe. When we get closer and see the waves, then we are moving backwards in time, and when we start to see waves breaking and coarse foam-like structures emerge, then we are at the boundary of the Planck scale. This foamy structure is called the Quantum foam and to understand it - would it then not make sense to create a model that mimics Wheeler's analogy?

Here a 3+1 Quantum foam model is constructed with a dynamical shift vector. The shift vector is a fundamental solution to a linearized wave equation. The existence of a sequence of smooth and compactly supported shift vectors that converge to the fundamental solution as distributions is used to define a Planck scale 3+1 Quantum foam as the formal limit of a sequence of 3+1 smooth spacetimes, each with a smooth and compactly supported shift vector.

Using basic topology and the existence of a partition of unity of test functions it is shown that a 'sum of sets' Quantum foam spacetime can be constructed by glueing together the sequence of smooth spacetimes.

It is proved that such Quantum foam spacetime necessarily has an inflation period.

It is shown, no matter how volatile the extrinsic curvature is in an early universe, that it will be smeared as time goes on. This might provide support to Carlip's view of an absorbed and nearly invisible cosmological constant.

Keywords: Quantum foam, 3+1 spacetime, dynamic shift vector, cosmology, inflation, initial data set glueing, cosmological constant

Introduction

The standard cosmology models for the observable universe presented in undergraduate or graduate textbooks on General Relativity or Quantum Fields in Curved Spacetime are homogeneous and isotropic solutions to Einstein's field equations, see e.g. [1],[2] or [3]. These are models where spacetime is foliated by a one-parameter family of hypersurfaces such that for each instant of time and for any two points on the corresponding hypersurface there exist an isometry of the metric field such that the points are indistinguishable. Moreover, the property that the universe is isotropic is a statement about rotational symmetry in the hypersurfaces for the timelike observers whose tangents necessarily are orthogonal to each hypersurface. Furthermore, the flow of time in any such model is decomposed by a lapse function and shift vector that are not considered dynamical since the view is that they only provide information about the global time evolution and the shift of the space coordinates between neighbouring hypersurfaces in the foliated spacetime.

But is it really possible that the universe remains homogeneous and isotropic as we trace time backwards into the high frequency epoch of the early universe? Is it not more likely that the early universe would be aligned with Wheeler's Quantum foam and its "wiggly business" [4]?

To understand the meaning of this question, let us use Wheeler's ocean surface analogy [4]. In this analogy the ocean surface is observed from an aeroplane but at different heights. An interpretation of this, from the perspective of a 3+1 cosmological model, is that the ocean, when seen from different heights, represents hypersurfaces at different times in the foliated universe. Therefore, the ocean surface seen from a great height when it appears smooth, flat and homogenous is like looking at the current hypersurface in our universe. When we get closer to the ocean and see the waves then we are moving backwards in time, and as we move even further closer and start to see waves breaking and a wiggly and coarse foam like structure, then we are getting closer to the physics at the Planck scale and the beginning of time. It is this foamy structure that is called the Quantum foam and to understand it - would it then not make sense to create a model that mimics Wheeler's analogy?

Here this is attempted using a 3+1 spacetime with a dynamic shift vector. The shift vector is taken to be a fundamental solution to the linearized wave equation that is zero at the beginning of time with a particle velocity that equals the Dirac measure. The existence of a sequence of smooth and compactly supported shift vector fields that converge as distributions to a fundamental solution of the linearized wave equation and hence a distribution in the

hypersurfaces suggests that the Quantum foam spacetime at the Planck scale should be defined as a 3+1 spacetime that is the formal limit of a sequence of spacetimes. Here each spacetime in the sequence is a 3+1 spacetime with a smooth and compactly supported test shift vector from the sequence of smooth and compactly supported shift vector fields.

This definition is motivated by two reasons besides that the 'kinematics' of a shift vector field that is a distribution should be sufficient to generate the coarse and wiggly foam-like structure in the hypersurfaces. The first reason, that has been mentioned but is repeated here, is the basic result from distribution theory; that for a given distribution there exists a sequence of compactly supported test functions such that the sequence converges to the given distribution as distributions. Therefore, a class of smooth spacetimes can be defined where each class element is a 3+1 spacetime with a shift vector field from the corresponding element in the sequence of compactly supported shift vector fields that converge to the fundamental solution. Consequently, there exists, at least formally, a good motivation for a Planck scale definition of a Quantum foam as a limit of a sequence of smooth spacetimes where the limit is in the sense of distributions.

Each class member is also a spacetime with a constant propagation phase number. This leads us to the second reason, that each class member represents a given period in time and thus an epoch of the universe. In conjunction with this and since each class member in the sequence of spacetimes is a globally hyperbolic smooth 3+1 spacetime, then it is reasonable that the class can be said to be a single globally hyperbolic spacetime in some well-defined meaning, provided that we can use some 'initial data engineering procedure' to glue together each class member to a single spacetime. It will be shown that this is possible and hence that a spacetime that is a 'sum of sets' can be constructed by glueing together the topological spaces in the sequence by forming a cover and use the existence of partition of unity in test functions to extend the shift vector field to the cover with a support that essentially is the sum of supports for each shift vector field in the sequence.

It should here be mentioned that there exists a well-defined theoretical framework for glueing initial data sets in General Relativity developed by Chruściel, Pollack and Isenberg [5]. However, here their framework is not used. Instead, the methodology outlined above where basic topology and the theory for test functions is applied to do the engineering of a representation of a 3+1 Quantum foam based cosmology.

With this 'set of sums' Quantum foam cosmology in place, it will be shown that it necessarily has an inflation period. It will also be shown that even if the spacetime in a neighbourhood of the Planck scale exhibits volatile extrinsic

curvature, will it be smeared out as the universe becomes homogeneous, isotropic and flat as time goes on.

These are two interesting ‘observations’ since the first provides a path forward to a generic and ad-hoc free resolution to the horizon problem while the latter a resolution to the problem with the prediction of a cosmological constant that is some 120 orders of magnitude too large at the Planck scale to be aligned with the currently observed accelerating universe [6]. The latter is also similar to the one proposed by Carlip [7]. That is, if the universe is really a ‘set of sums’ of smooth 3+1 proxy spacetimes that are glued together continuously, in the sense of initial data sets, then it provides for a mechanism where the Planck scale cosmological constant gets absorbed as time goes on.

Before we close this section we need to address “the elephant in the room” namely that the class of spacetimes converges to a linear distribution and hence that the metric field that is depending on such a shift vector is inconsistent with Einstein’s field equations. Or to put it precisely, the existence of a smooth sequence that converges to the fundamental solution as distributions is restricted to a linear theory but General relativity is inherently a non-linear theory where the curvature requires the connection to have a weak derivative and also to be locally square integrable [8]. To handle this we would perhaps be forced to use a non-linear distribution theory based on e.g. Colombeau algebra, see e.g. [8].

However, and this is a crucial remark, the manifold of the spacetime is countable; it then follows that every coordinate system has measure zero. This measure zero can thus be given an observer-independent notion in the manifold. Moreover, the universe is essentially the measure zero at the beginning of time. Expressed slightly differently; the sum of volumes of the open cubes that the image of any local coordinate system is a subset of can be made arbitrarily small and this is the measure zero. Practically this means that the less than Planck scale universe can be viewed as the measure zero set and hence a set with zero support. Perhaps and somewhat remarkable, this signifies that the potential problem of having a metric field where the shift vector in the limit is a distribution likely is not a problem since the limit has support zero and is not directly (and simultaneously) measurable but we should still be able to measure the ‘immeasurable’ with a smooth sequence of spacetimes. This also captures the essence of applied distribution theory. That is, to measure any physical quantity we must necessarily use a class of measurement devices where each class member has a well-defined size and hence non-zero support. In consequence, since any measurement device occupies a non-zero volume element it follows that any measurement necessarily is a functional (average) over the volume element. Particularly, and in the context of the Quantum

foam, no other choice exists than using test functions to understand the Planck scale Quantum foam ‘hidden’ in the measure zero set and the later evolution of the universe as time goes on.

Returning to the perception that the shift vector typicality is either zero or constant, there is actually an ongoing change in this. In fact, there is a growing number of studies e.g. work on binary black evolution and merger, see [9], [10] or [11] for models and a summary of models where the view is that the shift vector is dynamical. Also, I am working on a semi-classical analysis for a massless scalar field on an explicit class of spacetimes with a dynamic shift vector field [12].

Finally, this work is closed with a discussion and a summary of the results.

Constructing a Quantum foam Cosmology - measuring the immeasurable

In this section the construction of the Quantum foam cosmology is presented. Informally, a Planck scale Quantum foam cosmology is a 3+1 spacetime with a linear shift vector field that is a fundamental solution to the linearized wave equation and hence a distribution on the hypersurfaces such that it is zero at the beginning of time and with a Quantum foam particle velocity that equals the Dirac measure.

To be more precise, any given linear distribution can be approximated by a sequence of smooth and compactly supported test functions that converge to the distribution as distributions. Thus, if a vector field is a fundamental solution to the linearized wave equation then the solution is the convergence limit of a sequence of smooth and compactly supported vector fields. It then suggests, that a Planck scale Quantum foam spacetime with a dynamic shift vector that is a fundamental solution to the linearized wave equation can be defined as a 3+1 spacetime that is the formal limit of a sequence of smooth 3+1 spacetimes, each with a metric field that depends on a smooth and compactly supported shift vector field that is a solution to the linearized wave equation.

This is the definition that will be applied here. It is also shown directly, and as an example of initial data engineering, that a ‘sum of sets’ cosmology can be constructed from the sequence of spacetimes. Given this it will be proved that such a cosmology is an inflation model.

Finally, it is shown that even if the Quantum foam in a neighbourhood of the Planck region has a highly volatile extrinsic curvature it will be smeared as time goes on.

Before I start the construction of the model let us recall some basic preliminaries from topology, differential geometry and distribution theory that will be needed in the construction. Here I will follow standard texts on General

Relativity and distribution theory in [1], [2], [3] and [13] and [14].

To start, any manifold M in a spacetime, that is a pair (M, g) where g is a metric field on M , is a 4-dimensional Hausdorff topological space with a C^∞ structure. This means that M is covered by countable basis $\{O_i\}$ (that generates the topology) and that for each point $p \in M$ in the countable basis there exists a neighbourhood of p that is homeomorphic to some open set in \mathbb{R}^4 . That is, for each open subset O_i in the cover there exists a homeomorphism

$$\kappa_i: O_i \rightarrow U_i \subset \mathbb{R}^4 \quad (1)$$

where U_i is open in \mathbb{R}^4 . Moreover, if the intersection of any two homeomorphisms κ_i and κ_j is not empty then the transformation between κ_i and κ_j is homeomorphic and infinitely and continuously differentiable. The homeomorphism is interchangeable in the literature called a chart or local coordinate system, see e.g., [2], [14]. Here the homeomorphism will be called a local coordinate system.

The 3+1 spacetime model that will be used throughout the article is a globally hyperbolic spacetime that hence can be sliced or foliated by a one-parameter family of hypersurfaces. Recall that a hypersurface in a manifold M is the image $\Sigma = \phi(\hat{\Sigma})$ of a homeomorphic embedding $\phi: \hat{\Sigma} \rightarrow M$ where $\hat{\Sigma}$ is a 3-dimensional manifold, see e.g. [1], [2] or [11].

Thus the topology of the manifold M is $\mathbb{R} \times \Sigma$ where Σ is any Cauchy surface. In more details, since a globally hyperbolic spacetime is causally stable, meaning that no closed time-like curves exists, there exist a regular function $f: M \rightarrow \mathbb{R}$ such that each level surface of constant f is a Cauchy surface:

$$\Sigma_t = \{p \in M: f(p) = t\} \quad (2)$$

This generates a one-parameter family of Cauchy surfaces as time t goes on. Therefore,

$$M = \bigcup_{t \in \mathbb{R}} \Sigma_t. \quad (3)$$

This notation will be abused and for the union we will instead write

$$\Sigma_t \equiv \bigcup_{t \in \mathbb{R}} \Sigma_t \quad (4)$$

Now let us endow the manifold M with a structure of distributions by making use of the local coordinate systems and convolution as follows. For each coordinate system $\kappa_i: O_i \rightarrow U_i$ let $C_0^\infty(U_i)$ be the set of infinitely differentiable compactly supported test functions compactly supported on U_i . This space is denoted $\mathcal{D}(U_i)$ going forward. If for each such coordinate system κ_i we have a local distribution $u_{\kappa_i} \in \mathcal{D}'(U_i)$ such that

$$u_{\kappa_j} = (\kappa_i \circ \kappa_j^{-1}) * u_{\kappa_i} \quad (5)$$

in

$$\kappa_j(U_i \cap U_j) \quad (6)$$

then we say that u_{κ_i} is a distribution u_i in $O_i \subset M$. The set of all such distributions is denoted $\mathcal{D}'(M)$.

Next, we will make use of the notion of 'cut-off functions' from distribution theory as they are defined implicitly in Theorem 1 below:

Theorem 1 If U is in an open set in \mathbb{R}^n and K is a compact subset of U then there exists a compactly supported positive cut-off test function ϕ in U such that $0 \leq \phi \leq 1$ and where $\phi = 1$ in a neighbourhood of K . For a proof see [13] or theorem 1.4.1 in [14].

This theorem can be used to prove that a test function with compact support on a union of open sets can be written as a sum of test functions, each with support in the open sets of the union. Moreover, this and the cut-off test functions prove that a partition of unity exists, for proofs of these statements see [13].

The following theorem will turn out to be crucial in the construction of the proxy for the Quantum foam Cosmology:

Theorem 2 For any distribution $u \in \mathcal{D}'(U)$, there exists a sequence of $\{\phi_k\}$ of test functions in $\mathcal{D}(U)$ such that $\phi_k \rightarrow u$ as distributions. See theorem 2.2.1 in [13] or 4.1.5 in [14] for a proof.

We are now ready to construct a 3+1 Quantum foam cosmology with a dynamical shift vector field and will start with the notion of a Quantum foam shift vector field.

Definition 1 Quantum foam shift vector field β

Let β be shift vector on any hypersurface Σ_t with components β^i where $i = 1, 2, 3$ with $\beta^i \in \mathcal{D}'(\Sigma_t)$ for each i and such that each β^i is a fundamental solution to the linearized wave equation on Σ_t . That is

$$\Delta\beta - \frac{1}{N^2} \partial_t^2 \beta = 0 \quad (7)$$

with the initial conditions $\beta(\mathbf{x}, 0) = 0$ and $\partial_t \beta(\mathbf{x}, t)|_{t=0} = \delta(\mathbf{x})$, $\mathbf{x} \in \Sigma_t$ and where N is the lapse function. Then β is said to be a Quantum foam shift vector field.

Remark 1:

In local Cartesian coordinates is (7) expressed as

$$\frac{\partial^2 \beta_i}{\partial x^2} + \frac{\partial^2 \beta_i}{\partial y^2} + \frac{\partial^2 \beta_i}{\partial z^2} = \frac{1}{N^2} \frac{\partial^2 \beta_i}{\partial t^2}, i = x, y, z \quad (8)$$

Furthermore and to reinforce, with a fundamental solution to the linearized wave equation we mean any distribution β on any hyper surface Σ_t that is parametrized by time and is a solution to the linearized wave equation (7). The linearization of the wave equation is clearly an

approximation but required to avoid any problems having products of distributions that are not well-defined.

Example 1 Opposite Moving Heaviside disturbances

In one dimension we have that a fundamental solution is

$$\beta(t, x) = \frac{1}{2}(H(x+t) - H(x-t)) \quad (9)$$

with $N = 1$ and where H is the Heaviside function. Clearly

$$\beta(0, x) = 0 \quad (10)$$

and since the derivative of the Heaviside function is a Dirac measure

$$\partial_t \beta(t, x)|_{t=0} = \delta(x) \quad (11)$$

Given the Quantum foam shift vector field then there exist a smooth and a compactly supported sequence of shift vector fields that converge as distributions to the Quantum foam shift vector, see theorem 2. This sequence will now be introduced.

Definition 2 Smooth Quantum foam shift vector field β_k

Let $\{\beta_k\}$ be a sequence of compactly supported vector fields $\beta_k \in \mathcal{D}(\Sigma)$ where $k \in \mathbb{N}$ such that $\beta_k = (\beta_k^1, \beta_k^2, \beta_k^3)$ is a smooth solution to the linearized wave equation (7)

$$\Delta \beta_k - \frac{1}{N_k^2} \partial_t^2 \beta_k = 0 \quad (12)$$

If the sequence converge as distributions to the Quantum foam vector field β in definition 1 then it is said to be a sequence of Quantum foam shift vector fields and an element β_k is said to be a smooth Quantum foam shift vector field with a lapse N_k and a propagation phase number k .

Example 2 The logistic representation

In 1+1 dimension we have that one representation of a Quantum foam shift vector field sequence can be constructed from a sigmoid with suitable support:

$$\beta_k(x, t) = \frac{1}{2}(\tanh(v_k(x+t)) - \tanh(v_k(x-t))) \quad (13)$$

$$v_k = kv \quad (14)$$

where v_k is the indexed frequency and v is said to be the background frequency.

Remark 2:

Here it is worthwhile to notice the resemblance of the shift vector in (13) with Alcubierre's warp drive shift vector [8].

Furthermore, to get some insight into the indexed frequency, v_k in (14),

$$v_k = kv,$$

we introduce a regular function $K_t: \Sigma_t \rightarrow \mathbb{R}$ and choose a coordinate system κ . Then a local level surface for each hypersurface Σ_t can be defined using the map

$$K_\kappa = K_t \circ \kappa^{-1} \quad (15)$$

Hence, in local coordinates we have

$$K_\kappa(t, x) = C_t \quad (16)$$

where C_t is the height at a given time t . Consequently we have the local level surface

$$loc(\Sigma_t) = K^{-1}_\kappa(C_t) \quad (17)$$

To continue for a period dt/k when the shift has completed a period, then the height C_t has decreased with one unit and as a plane wave approximation we define the indexed frequency by

$$v_k = \partial_t K = k/dt = kv \quad (18)$$

where

$$v = 1/dt \quad (19)$$

is said to be the background frequency.

Next, the class of smooth Quantum foam spacetimes is given by definition 3.

Definition 3 Class of smooth Quantum foam Cosmologies

Let $\{\beta_k\}$ be a sequence of Quantum foam shift vector field and $\{g_{\beta_k}\}$ a sequence of metric fields such that

$$g_{\beta_k} = -N_k^2 dt^2 + \eta_{ij}(dx^i + \beta_k^i dt)(dx^j + \beta_k^j dt) \quad (20)$$

$$\eta = \text{diag}(1, 1, 1) \quad (21)$$

$$N_k = \frac{1}{k}, k^2 = k_1^2 + k_2^2 + k_3^2, k_1, k_2, k_3 \in \mathbb{N} \quad (22)$$

where N_k is the lapse function. Then the sequence

$$\{(\Sigma_t, g_{\beta_k})\} \quad (23)$$

of smooth 3+1 spacetimes is said to be a class of smooth Quantum foam cosmologies and an element (Σ_t, g_{β_k}) is said to be an Quantum foam Cosmology with a propagation phase number k .

Remark 3:

Here any expansion factor or scale factor a as a function of the global time has been left out from the metric field. Also, any discussion about the initial hypersurface has been deliberately left out until definition 4 below has been given.

Example 3 A 1+1 toy Heaviside Universe

In 1+1 dimension we can use the sigmoid sequence $\{\beta_k^1\}$ in Example 2 to construct a Quantum foam spacetime with an early inflation period but that becomes flat as time goes on:

$$g_{\beta_k} = -\frac{1}{k^2} dt^2 + (dx + \beta_k^1 dt)(dx + \beta_k^1 dt) \quad (24)$$

where

$$\beta_k^1(x, t) = \frac{1}{2}(\tanh(v_k(x+t)) - \tanh(v_k(x-t))) \quad (25)$$

$$v_k = kv, k \in \mathbb{N} \quad (26)$$

We are now ready for the definition of a 3+1 Quantum foam cosmology at the Planck scale and hence to provide meaning to the notion of the initial hypersurface.

Definition 4 Planck scale 3+1 Quantum foam spacetime

Given the class of Quantum foam spacetimes, then we define an initial Planck scale Quantum foam spacetime as the *formal limit*

$$(\Sigma_0, g_\beta) = \lim_{k \rightarrow \infty} (\Sigma_{1/k}, g_{\beta_k}) \quad (27)$$

Remark 4:

It is important here to notice that there do exist many possible classes of smooth spacetimes that converge to this initial hypersurface and metric field. In terms of measurement this is a statement of the existence of different measurement devices that can be used to measure a physical quantity. Nevertheless, observations in our current universe should uniquely determine a class of spacetimes that fits the observations.

Remark 5:

In definition 4 it is not a requirement that the metric field should satisfy the nonlinear field equations of General relativity even if every single member of the sequence will, since the limit is the measure zero. However, the notion of measure zero on a manifold needs to be given a precise meaning and it needs to be shown that it is observer independent for this statement to make sense. Thus let us use the standard definition of measure zero on a manifold and show that it is observer independent.

Definition 5 Measure zero on a Hausdorff manifold M with a C^∞ structure

Let O be a subset of M then O is said to be a measure zero in M if for any local coordinate system

$$\kappa_i: O_i \rightarrow U_i \quad (28)$$

the image of $O \cap O_i$ is a measure zero in U_i .

The following result should be a standard (graduate) result, but will be proved here for clarity.

Theorem 3 Let M be a countable Hausdorff manifold with a C^∞ structure then M has a measure zero for any coordinate system.

Proof

The manifold M is generated by a countable basis $\{O_i\}$ such that for each O_i a homeomorphism exists in the family

$$\mathfrak{S} = \{\kappa_i\} \quad (29)$$

of coordinate systems where

$$\kappa_i: O_i \rightarrow U_i \subset \mathbb{R}^4, O_i \subset M \quad (30)$$

To each local coordinate system and hence U_i we can define an open cube, C_{ij} , using an open interval (a_{ij}, b_{ij}) where j is an index for the interval. That is

$$C_{ij} = (a_{ij}, b_{ij}) \times (a_{ij}, b_{ij}) \times (a_{ij}, b_{ij}) \times (a_{ij}, b_{ij}) \quad (31)$$

and form a union of cubes so that,

$$U_i \subset C_i = \cup_{j=1}^{\infty} C_{ij} \quad (32)$$

The volume of this can be made arbitrarily small,

$$Vol(C_i) = \sum_{j=1}^{\infty} (b_{ij} - a_{ij})(b_{ij} - a_{ij})(b_{ij} - a_{ij})(b_{ij} - a_{ij}) < \varepsilon \quad (33)$$

for every

$$\varepsilon > 0$$

since if we take

$$a_{ij} = x_{ij} - \frac{\varepsilon^{1/4}}{2^{j/4+5/4}}, b_{ij} = x_{ij} + \frac{\varepsilon^{1/4}}{2^{j/4+5/4}} \quad (34)$$

then

$$Vol(C_i) = \frac{\varepsilon}{2} < \varepsilon \quad (35)$$

Thus for each local coordinate system κ_i there is a measure zero (in M) that is observer independent.

From an ‘engineering perspective’ the universe at scales less than a cube with sides equal or less than the Planck length can be identified with measure zero and hence represent a physical quantity with zero support and as such necessarily is a distribution that can only be measured by measurement devices with a well-defined size, that is test functions. In this context it means a sequence of smooth Quantum foam spacetimes or for that matter a ‘sum of sets’ spacetime that will be introduced next:

Theorem 4 A globally hyperbolic Quantum foam spacetime can be constructed by glueing together the elements in the class of smooth Quantum foam spacetimes

Proof

Each member in the class of smooth Quantum foam cosmologies in definition 3 is a smooth 3+1 spacetime and hence globally hyperbolic. Use the sequence to create a cover. This cover is a countable union of manifolds since the class is countable. Create a partition of unity in test functions for the cover that extends the shift vector field to the cover. This forms a sum of sets in the sense of the support of the shift vector fields with support in the union. Construct the metric field from this sum of sets shift vector field with the lapse function already given. This completes the initial data engineering

I will now prove that the Quantum foam cosmology inherently has a period of inflation.

Theorem 5 The Quantum foam cosmology necessarily has an inflation period

Proof

Let β_k be a Quantum foam shift vector field for a propagation phase number k then since β_k is constructed from cut-off functions (see theorem 1) and converge as distributions to a linear distribution it follows that $\text{supp}(\beta_k) \rightarrow \{0\}$ as $k \rightarrow \infty$ but since β_k is smooth and compactly supported, that is $\beta_k \in C^\infty_0$, there exist a $N_0 \in \mathbb{N}$ such that for all $k > N_0$ β_k increases from 0 to 1 (or decreases from 1 to 0) faster than any polynomial. Finally, this change applied to any scale factor a shows the existence of an inflation period.

It will now be proved that the cosmology is homogeneous and isotropic in the low wavenumber limit

Theorem 6 The Quantum foam cosmology is homogeneous, isotropic in the propagation phase number limit $k_i \rightarrow 1$ for all i .

Proof

Form a cube with side L_t such that the observable universe at any time t is contained in a cube with side L_t then we have that the indexed background frequency ν_{k_i} (wave number) is equal to

$$\nu_{k_i} = \frac{2\pi}{L_t} k_i, k_i \in \mathbb{N}, i = 1,2,3 \quad (36)$$

Now when $k_i \rightarrow 1$ then the indexed background frequency vanishes since the side $L_t \rightarrow \infty$ and it follows that $\beta_k^i \rightarrow 0$ and hence that

$$g_{\beta_k} = -\frac{1}{k^2} dt^2 + \eta_{ij}(dx^i + \beta_k^i dt)(dx^j + \beta_k^j dt) \rightarrow g = -dt^2 + \eta_{ij} dx^i dx^j \quad (37)$$

Remark 6:

A consequence of this is that independently of whatever volatility that is exhibited in the extrinsic curvature of the spacetime and in a neighbourhood of the Planck region, it follows that the volatility will average out as time goes on.

Discussion

In this final section I will summarise the developed ‘engineering’ framework for the Quantum foam cosmology that has been introduced. I would also like to take the opportunity to comment on Carlip’s view of the Quantum foam in his discussion of whether the cosmological constant is hidden [7]. Finally, I would also like to point out future directions of research.

This framework has been truly inspired by the interview with the late Professor Wheeler [4] and his use of the ocean surface analogy to describe the Quantum foam. A description I have interpreted from the perspective of a foliated universe with the ocean surface as the hypersurfaces. I then concluded that it would be reasonable to assume that the shift vector, tangent to the hypersurfaces, should not be an unchanging and static quantity but rather a dynamic entity with wavelike properties and not necessarily smooth. If this is true then assuming that the shift vector is a fundamental solution to the linearized wave equation is reasonable since not only do we allow for wigglyness and coarse hypersurfaces but we also maintain time as a parameter. In this study I have avoided working directly with this fundamental solution since it is a distribution with zero support. Instead I have used the existence of a sequence of well-behaved smooth test functions that converge to the solution as distributions to define a class of smooth spacetimes. This means that each spacetime in the class is a globally hyperbolic spacetime, but

it could well be that these spacetimes will stack up and generate a Cauchy horizon as we move closer to the beginning of time. But it is not clear if this really is a problem given the distributional limit and hence that the Planck scale spacetime will be the measure zero.

To continue, the framework that has been introduced here, in order to create a model of the Quantum foam, can clearly be criticised to be coordinate dependent given the explicit form of the metric field in the definition of the Quantum foam as the formal limit of the class of smooth spacetimes. Nevertheless, the classic theorem that any globally hyperbolic spacetime can be foliated by the hyper surfaces of a regular function with a height that is the time, see e.g. [2] for a proof, suggests that the model as such is not void of observable physics. Or to express it slightly differently - even if a choice of coordinates may lead straight into the abyss if we are not careful, we can still make sense of results in coordinates if we choose to interpret the result in the frames that shares some property e.g. that the frequency is very high as we typically do when we construct and interpret the frequency four vector (wave vector) that is the derivative of a scalar function in the hypersurfaces parameterized by time.

Consequently, I believe one needs to be somewhat cautious with statements of void physics because a selection of a coordinate system is made. Further to this, we need to remember that the existence of a measure zero is a coordinate system independent statement and thus in the convergence limit of the Planck scale Quantum foam there do not exist any coordinate system selection problem; particularly the class of smooth spacetimes presented here should be perfectly valid approximations to the Quantum foam.

Furthermore, I should mention that since having a dynamic shift vector that model’s Wheeler’s Quantum foam where the dynamic shift cause volatile curvature on the hypersurfaces suggests that Carlip’s high level analysis [7] that relies on a zero a shift vector most likely is not sufficient to support a highly volatile cosmological constant at the Planck scale that is smeared as time goes on. Instead, I would argue that his model should be changed to allow for a highly dynamic shift vector that in the beginning and in the direction of time is represented by a Dirac measure. The class of spacetimes that approximates the Planck scale Quantum foam spacetime then provides a resolution to the dilemma with a cosmological constant that is out of range by an order of 120 since if the curvature is highly volatile for phases in the neighbourhood of the Planck scale, it will be smeared as time goes on and the propagation phase number goes to zero.

In summary, this work has hence shown that it is possible to construct a self-consistent Planck scale Quantum foam cosmology where the metric field is wiggly in the sense that the linear vector shift field is a fundamental solution to the

linearized wave equation but that linear distribution theory can be used to create a class of spacetimes and hence a sum of sets glued spacetime that represent all the different epochs of the universe. Using this spacetime, it has been proved from distribution theory that the spacetime has an inflation period. It has also been shown that no matter how volatile the extrinsic curvature is in an early universe, it will be averaged out as time goes on. Finally, I am working on the analysis for a quantum field theory for a massless scalar field on an explicit 3+1 model where the fundamental solution is a time-scaled Dirac measure. The motivation behind that study is to understand the effects the volatility has on the scalar field in relation to regions that would be causally disconnected if it were not for the inflation mechanism inherent in the Quantum foam cosmology [12].

- [11]ourgoulhon E 2012 3+1 Formalism in General Relativity Bases of Numerical Relativity (Berlin and Heidelberg: Springer-Verlag)
- [12]Cramer C 2022 The time-scaled Dirac 3+1 Quantum foam spacetime To be Published
- [13]Strichartz R S 2003 A Guide to Distribution Theory and Fourier Transforms (World Scientific)
- [14]Hörmander L 1990 The Analysis of Linear Partial Differential Operators I Distribution Theory and Fourier Analysis (Berlin and Heidelberg: Springer-Verlag)

References

- [1] [1]Hawking S W, Ellis G F R 1973 The large scale structure of space-time (Cambridge: Cambridge University Press)
- [2] Wald R M 1984 General Relativity (Chicago: Chicago University Press)
- [3] Birrell N D, Davies P C W 1982 Quantum fields in curved space (Cambridge: Cambridge University Press)
- [4] Web of Stories - Life Stories of Remarkable People 2017 Interview with John Wheeler Quantum ideas. Quantum foam. Max Planck and Karl PopperLike a story <https://www.webofstories.com/playAll/john.wheeler?sId=9542>, <https://www.youtube.com/watch?v=iNxioquoK0g>
- [5] Chruściel P T, Pollack D, Isenberg J 2004 Gluing Initial Data Sets for General Relativity Phys. Rev. Lett. 93, 081101
- [6] Hobson M P, Efstathiou G P, Lasenby A N 2012 General Relativity An Introduction for Physicists (Cambridge: Cambridge University Press), p187
- [7] Carlip S 2019 Hiding the Cosmological Constant Phys. Rev Lett 123. 13102
- [8] Steinbauer R, Vickers J A 2006 The use of generalized functions and distributions in general relativity Class.Quant.Grav. 23 (2006) R91-R114
- [9] Alcubierre M 1994 The warp drive: hyper-fast travel within general relativity Class.Quant.Grav.11:L73-L77,1994
- [10] Alcubierre M, Brügmann B, Diener P , Guzmán F. S, Hawke I, Hawley S, Herrmann F, Koppitz M, Pollney D , Seidel E, and Thornburg J 2005 Dynamical evolution of quasicircular binary black hole data Phys. Rev. D 72, 044004