

Quantum foam Cosmology

-using a class of smooth 3+1 spacetimes to measure the immeasurable

Claes Cramer

YBC—Young Business Creatives, Upper Secondary School, Experimentgatan 7, 131 54 Nacka, Sweden

Abstract

To understand the fabric of spacetime we can search for inspiration in Wheeler's layman's analogy of the Quantum foam using an ocean surface observed from an aeroplane at different heights. From a 3+1 spacetime perspective, the ocean surface represents the hypersurfaces of spacetime at different times. Therefore, the ocean surface observed from a great height, when it appears smooth, homogenous, isotropic and flat is like looking at the current universe. When we get closer and see the waves, then we are moving backwards in time, and when we start to see waves breaking and coarse foam-like structures emerge, then we are at the boundary of the Planck scale. This foamy structure is called the Quantum foam and to understand it - would it then not make sense to create a model that mimics Wheeler's analogy?

Here a 3+1 Quantum foam model is constructed with a dynamical shift vector. The shift vector is a fundamental solution to a linearised wave equation. The existence of a sequence of smooth and compactly supported shift vectors that converge to the fundamental solution as distributions is used to define a Planck scale 3+1 Quantum foam as the formal limit of a sequence of 3+1 smooth spacetimes, each with a smooth and compactly supported shift vector.

Using basic topology and the existence of a partition of unity of test functions it is shown that a sum-of-sets Quantum foam spacetime can be constructed by gluing together the sequence of smooth spacetimes.

It is proved that such Quantum foam spacetime necessarily has an inflation period.

It is shown, no matter how volatile the extrinsic curvature is in an early universe, that it will be smeared as time goes on. This might provide support to Carlip's view of an absorbed and nearly invisible cosmological constant.

Keywords: Quantum foam, 3+1 spacetime, dynamic shift vector, cosmology, inflation, initial data set gluing, cosmological constant

Introduction

The standard cosmology models for the observable universe presented in undergraduate or graduate textbooks on General Relativity or Quantum Fields in Curved Spacetime are homogeneous and isotropic solutions to Einstein's field equations, see e.g. [1],[2] or [3]. These are models where spacetime is foliated by a one-parameter family of hypersurfaces such that for each instant of time and for any two points on the corresponding hypersurface there exist an isometry of the metric field such that the points are indistinguishable. Moreover, the property that the universe is isotropic is a statement about rotational symmetry in the hypersurfaces for the timelike observers whose tangents necessarily are orthogonal to each hypersurface. The flow of time in any such model is decomposed by a lapse function and a shift vector that are not considered dynamical since the view is that they only provide information about the time evolution and the shift of the space coordinates between neighbouring hypersurfaces in the foliated spacetime.

But is it really possible that the universe remains homogeneous and isotropic as we trace time backwards into the high frequency epoch of the early universe? Is it not more likely that the early universe would be aligned with Wheeler's Quantum foam and its "wiggly business" [4]?

To understand the meaning of this question, let us use Wheeler's ocean surface analogy [4]. In this analogy the ocean surface is observed from an aeroplane but at different heights. An interpretation of this, from the perspective of a 3+1 cosmological model, is that the ocean, when seen from different heights, represents hypersurfaces at different times in the foliated universe. Therefore, the ocean surface seen from a great height when it appears smooth, flat and homogenous is like looking at the current hypersurface in our universe. When we get closer to the ocean and see the waves then we are moving backwards in time, and as we move even further closer and start to see waves breaking and a wiggly and coarse foam like structure, then we are getting closer to the physics at the Planck scale and the beginning of time. It is this foamy and hence bubbly structure that is called the Quantum foam and to understand it - would it then not make sense to create a model that mimics Wheeler's analogy?

Here this is attempted using a 3+1 spacetime with a dynamic shift vector. The shift vector is taken to be a fundamental solution to a linearised wave equation that is zero at the beginning of time with a particle velocity that equals the Dirac measure. The existence of a sequence of smooth and compactly supported shift vector fields that converge as distributions to a fundamental solution of a linearised wave equation and hence a distribution in the hypersurfaces suggests that the Quantum foam spacetime at

the Planck scale should be defined as a 3+1 spacetime that is the formal limit of a sequence of spacetimes. Here each spacetime in the sequence is a 3+1 spacetime with a smooth and compactly supported test shift vector from the sequence of smooth and compactly supported shift vector fields.

This definition is motivated by two reasons besides that the 'kinematics' of a shift vector field that is a fundamental solution the wave equation and as such a distribution should be sufficient to generate the coarse and wiggly foam-like structure in the hypersurfaces. The first reason, that has been mentioned but is repeated here, is the basic result from distribution theory; that for a given distribution there exists a sequence of compactly supported test functions such that the sequence converges to the given distribution as distributions, see e.g. [5] or [6] for proof of this statement. Therefore, a class of smooth spacetimes can be defined where each class element is a 3+1 spacetime with a shift vector field from the corresponding element in the sequence of compactly supported shift vector fields that converge to the fundamental solution. Consequently, there exists, at least formally, a good motivation for a Planck scale definition of a Quantum foam as a limit of a sequence of smooth spacetimes where the limit is in the sense of distributions.

Each class member is also a spacetime with a constant propagation number. This leads us to the second reason, that each class member represents a given period in time and thus an epoch of the universe with a constant lapse that is the reciprocal of the propagation number. In conjunction with this and since each class member in the sequence is a globally hyperbolic smooth 3+1 spacetime, then it is reasonable that the class can be said to be a single globally hyperbolic spacetime if it is possible to glue together the elements. It will be shown that this is possible and hence that a spacetime that is a sum-of-sets can be constructed by forming a disjoint union of the elements, showing that a surjection exists that generates an equivalence relation that adhere the elements and provides for the quotient topology. To define a metric on this sum-of-sets is the shift vector extended to the disjoint union with a support that essentially is the sum of supports for each shift vector field.

It should here be mentioned that there exists a well-defined theoretical framework for gluing initial data sets in general relativity developed by Chruściel, Pollack and Isenberg [7]. However, here their framework is not used directly. Instead, to perform the engineering work to glue together a sum-of-sets 3+1 Quantum foam cosmology is the process previously outlined used.

Next it will be shown that a Quantum foam cosmology has a period when its expansion is more rapidly than any polynomial. To make this statement more comprehensive

will it also be shown that the volume expansion for the Eulerian observers that are freely falling is more rapidly than any polynomial. Also, the trace of the extrinsic curvature displays the same characteristics. All together, this provides support for the idea that the Quantum cosmology has an period of inflation.

Next, it will also be shown that even if the spacetime in a neighbourhood of the Planck scale exhibits volatile extrinsic curvature, will it be smeared out since the shift is compactly supported and hence vanishes as time goes on and consequently the universe is necessarily becoming homogeneous, isotropic and flat as time goes on. These are two noteworthy ‘observations’ since the first provides a path forward to a generic and ad-hoc free resolution to the horizon problem while the latter a resolution to the problem with the prediction of a cosmological constant that is some 120 orders of magnitude too large at the Planck scale to be aligned with the currently observed accelerating universe [8]. The latter is also similar to the one proposed by Carlip [9]. That is, if the universe is really a sum-of-sets of smooth 3+1 proxy spacetimes that are glued together continuously, in the sense of initial data sets, then it provides for a mechanism where the Planck scale cosmological constant gets absorbed as time goes on.

Before this section is closed we need to address “the elephant in the room” namely that the shift vector in the class of spacetimes converges to a linear distribution with zero support and hence to a linear combination of the Dirac measure and its derivatives. This follows from the standard structure theorem that a distribution with point support necessarily has the form of a linear combination of the Dirac measure and its derivatives, see [5] or [6] for a proof.

This implies that the metric field depending on such a shift vector is inconsistent with Einstein’s field equations. Or to put it precisely, the existence of a smooth sequence that converges to the fundamental solution as distributions is restricted to a linear theory but general relativity is inherently a non-linear theory. Therefore, general relativity requires non-linear operations to be well-defined and this is problematic in relation to linear distribution theory since the product of two arbitrary distributions is undefined. To handle this we would perhaps be forced to use a non-linear distribution theory based on e.g. Colombeau algebra [10].

Nonetheless, the discrete measure, that is a measure that is a sum of Dirac measures, on each spacetime in the sequence is scaled by the propagation number in such a way that it has a zero limit as the propagation number goes to infinity. Perhaps and somewhat remarkable, this signifies that the potential issue with a metric field where the shift vector in the limit is a distribution is not that problematic and that we should be able to measure the “immeasurable” with a smooth

sequence of globally hyperbolic 3+1 spacetimes. This also captures the essence of applied distribution theory. That is, to measure any physical quantity we must necessarily use a class of measurement devices (test functions) where each class member has a well-defined size and hence non-zero support. In consequence, since any measurement device occupies a non-zero volume element it follows that any measurement necessarily is a functional (average) on the space of the test functions over the volume element or a sequence of test functions that converge as distributions to the physical quantity. Particularly and to summarise, in the context of the Quantum foam, no other choice exists than using a sequence spacetimes, each with a compactly supported shift vector field, to understand the Planck scale Quantum foam, hidden in a measure space that scales to almost nothing at the beginning of time, and the later evolution of the universe as time goes on.

In relation to the discussion about the field equations we should also say something about the energy conditions for the Quantum cosmology. Using that Alcubierre’s warp drive [11], or rather that a warp bubble with a constant velocity is a special case of this model if we allow ourself to extend the model to the space that the shift vectors is dense in, that is to shift vectors in the space of rapidly decreasing functions and its dual space of tempered distributions. The warp bubble is then a representation of a Quantum foam and it has been shown (as expected) that it violates the weak energy condition [12]. Also, the null energy condition is violated [12]. Given this, we conclude that the Quantum foam will manifest at least the same violations even though the severity will vary due to a lapse functions reciprocal relation to the propagation number.

Returning to the issue that the shift vector usually is assumed to be a non-dynamical quantity, then it should be noted that there is an ongoing change in this perception and there exist a number studies, see [11], [13], [14] or [15] for models and a summary of models where the view is that the shift vector is dynamical. Also, I am working on a semi-classical analysis for a massless scalar field on an explicit class of spacetimes with a dynamic shift vector field [16].

Finally, this work is closed with a discussion and a summary of the results.

Constructing a Quantum foam Cosmology - measuring the immeasurable

In this section the construction of the Quantum foam cosmology is presented. Informally, a Planck scale Quantum foam cosmology is a 3+1 spacetime with a linear shift vector field that is a fundamental solution to a linearised wave equation and hence a distribution on the hypersurfaces such that it is zero at the beginning of time and with a Quantum foam particle velocity that equals the Dirac measure.

To be more precise, any given linear distribution can be approximated by a sequence of smooth and compactly supported test functions that converge to the distribution as distributions. Thus, if a vector field is a fundamental solution to a linearised wave equation then the solution is the convergence limit of a sequence of smooth and compactly supported vector fields. It then suggests, that a Planck scale Quantum foam spacetime with a dynamic shift vector that is a fundamental solution to the wave equation can be defined as a 3+1 spacetime that is the formal limit of a sequence of smooth 3+1 spacetimes, each with a metric field that depends on a smooth and compactly supported shift vector field that is a solution to the wave equation. This is the definition that will be applied here. This limit will have point support and hence it follows from a basic structure theorem, see [5] or theorem 2.3.4 in [6], in distribution theory that the limit necessarily has the form of a linear combination of the Dirac measure and its derivatives. However, it will be shown that the discrete measure, constructed from the σ -algebra of the power set for the manifold scales to zero in the convergence limit when the propagation number goes to infinity. This means that the Quantum foam only can be understood by the sequence and in fact arise out of almost nothing.

It is also shown as an example of manifold gluing, that a sum-of-sets cosmology can be constructed from the sequence of spacetimes.

It will be proved that a Quantum cosmology has a period when the expansion is more rapid than any polynomial. In relation to this is it also shown that freely falling Eulerian observers experience a volume expansion that is more rapid than any polynomial. It is also shown that the trace of the extrinsic curvature manifest the same result.

Finally, it is shown that even if the Quantum foam in a neighbourhood of the Planck region has a highly volatile extrinsic curvature it will be smeared as time goes on.

Before the model is constructed let us recall some basic preliminaries from topology, differential geometry and distribution theory that will be needed in the construction. Here the presentation will follow standard texts on general

relativity and distribution theory in [1], [2], [3] and [5] and [6].

To start, any manifold M in a spacetime, that is a pair (M, g) where g is a metric field on M , is a 4-dimensional Hausdorff topological space with a C^∞ structure. This means that M is covered by countable basis $\{O_i\}$ (that generates the topology) and that for each point $p \in M$ in the countable basis there exists a neighbourhood of p that is homeomorphic to some open subset in \mathbb{R}^4 . That is, for each open subset O_i in the cover there exists a homeomorphism

$$\kappa_i: O_i \rightarrow U_i \subset \mathbb{R}^4 \quad (1)$$

where U_i is an open subset in \mathbb{R}^4 . Moreover, if the intersection of any two homeomorphisms κ_i and κ_j is not empty then the transformation between κ_i and κ_j is homeomorphic and infinitely and continuously differentiable. The homeomorphism is interchangeable in the literature called a chart or local coordinate system, see e.g., [2], [6]. Here the homeomorphism will be called a local coordinate system.

The 3+1 spacetime model that will be used throughout the article is a globally hyperbolic spacetime that hence can be sliced or foliated by a one-parameter family of hypersurfaces. Recall that a hypersurface in a manifold M is the image $\Sigma = \phi(\hat{\Sigma})$ of a homeomorphic embedding $\phi: \hat{\Sigma} \rightarrow M$ where $\hat{\Sigma}$ is a 3-dimensional manifold, see e.g. [1], [2] or [15].

Thus the topology of the manifold M is $\mathbb{R} \times \Sigma$ where Σ is any Cauchy surface. In more details, since a globally hyperbolic spacetime is causally stable, meaning that no closed time-like curves exists, there exist a regular function $f: M \rightarrow \mathbb{R}$ such that each level surface of constant f is a Cauchy surface:

$$\Sigma_t = \{p \in M: f(p) = t\}. \quad (2)$$

This generates a one-parameter family of Cauchy surfaces as time t goes on. Therefore,

$$M = \bigcup_{t \in \mathbb{R}} \Sigma_t. \quad (3)$$

This notation will be abused and for the union we will instead write

$$\Sigma_t \equiv \bigcup_{t \in \mathbb{R}} \Sigma_t \quad (4)$$

whenever it is needed.

Now let us endow the manifold M with a structure of distributions by making use of the local coordinate systems and convolution as follows. For each coordinate system $\kappa_i: O_i \rightarrow U_i$ let $C_0^\infty(U_i)$ be the set of infinitely differentiable compactly supported test functions compactly supported on U_i . This space is denoted $\mathcal{D}(U_i)$ going forward. If for each

such coordinate system κ_i we have a local distribution $u_{\kappa_i} \in \mathcal{D}'(U_i)$ such that

$$u_{\kappa_j} = (\kappa_i \circ \kappa_j^{-1}) * u_{\kappa_i} \quad (5)$$

in

$$\kappa_j(U_i \cap U_j) \quad (6)$$

then we say that u_{κ_i} is a distribution u_i in $O_i \subset M$. The set of all such distributions is denoted $\mathcal{D}'(M)$.

Next, we will make use of the notion of cut-off test functions from distribution theory as they are defined implicitly in Theorem 1 below:

Theorem 1 If U is in an open set in \mathbb{R}^n and K is a compact subset of U then there exists a compactly supported positive cut-off test function ϕ in U such that $0 \leq \phi \leq 1$ and where $\phi = 1$ in a neighbourhood of K . For a proof see [5] or theorem 1.4.1 in [6].

This theorem can be used to prove that a test function with compact support on a union of open sets can be written as a sum of test functions, where each test function has support in one open set of the union. Moreover, this and the cut-off test functions prove that a partition of unity exists, for proofs of these statements see theorem 1.4.4 and 1.4.5 in [6].

The following theorem will turn out to be crucial in the construction of the proxy for the Quantum foam Cosmology:

Theorem 2 For any distribution $u \in \mathcal{D}'(U)$, there exists a sequence $\{\phi_k\}_{k \in \mathbb{N}}$ of test functions in $\mathcal{D}(U)$ such that $\phi_k \rightarrow u$ as distributions. See theorem 2.2.1 in [5] or 4.1.5 in [6] for a proof.

We are now ready to construct a 3+1 Quantum foam cosmology with a dynamical shift vector field and will start with the notion of a Quantum foam shift vector field.

Definition 1 Quantum foam shift vector field β

Let β be shift vector on any hypersurface Σ_t with components β^i where $i = 1, 2, 3$ with $\beta^i \in \mathcal{D}'(\Sigma_t)$ for each i and such that each β^i is a fundamental solution to the linearised wave equation on Σ_t . That is

$$\Delta \beta - \frac{1}{N^2} \partial_t^2 \beta = 0 \quad (7)$$

with the initial conditions $\beta(\mathbf{x}, 0) = 0$ and $\partial_t \beta(\mathbf{x}, t)|_{t=0} = \delta(\mathbf{x})$, $\mathbf{x} \in \Sigma_t$ and where N is a constant lapse. Then β is said to be a Quantum foam shift vector field.

Remark 1:

In local Cartesian coordinates is (7) expressed as

$$\frac{\partial^2 \beta^i}{\partial x^2} + \frac{\partial^2 \beta^i}{\partial y^2} + \frac{\partial^2 \beta^i}{\partial z^2} = \frac{1}{N^2} \frac{\partial^2 \beta^i}{\partial t^2}, i = x, y, z. \quad (8)$$

Furthermore and to reinforce, with a fundamental solution to the linearised wave equation we mean any distribution β on any hyper surface Σ_t that is parametrized by time and is a solution to the linearised wave equation (7). The linearisation, that is the linear part of the wave equation in a foliated 3+1 spacetime is clearly an approximation but required to avoid any problems having products of distributions that are not well-defined.

Example 1 Opposite Moving Heaviside disturbances

In one dimension we have that a fundamental solution is

$$\beta(t, x) = \frac{1}{2} (H(x+t) - H(x-t)) \quad (9)$$

where H is the Heaviside function.

Clearly,

$$\beta(0, x) = 0 \quad (10)$$

and since the derivative of the Heaviside function is a Dirac measure

$$\partial_t \beta(t, x)|_{t=0} = \delta(x). \quad (11)$$

Given the Quantum foam shift vector field then there exist a smooth and a compactly supported sequence of shift vector fields that converge as distributions to the Quantum foam shift vector in definition 1, see theorem 2. This sequence will now be introduced.

Definition 2 Smooth Quantum foam shift vector field β_k

Let $\{\beta_k\}_{k \in \mathbb{N}}$ be a sequence of compactly supported vector fields $\beta_k \in \mathcal{D}(\Sigma_t)$ such that each $\beta_k = (\beta_k^1, \beta_k^2, \beta_k^3)$ is a smooth solution to the wave equation (7),

$$\Delta \beta_k - \frac{1}{N_k^2} \partial_t^2 \beta_k = 0. \quad (12)$$

If the sequence converge as distributions to the Quantum foam vector field β in definition 1 then it is said to be a sequence of Quantum foam shift vector fields and an element β_k is said to be a smooth Quantum foam shift vector field with a propagation phase number k and an associated constant lapse N_k .

Example 2 The logistic representation

In 1+1 dimension we have that one representation of a Quantum foam shift vector field sequence can be constructed from a sigmoid with suitable support:

$$\beta_k(x, t) = \frac{1}{2} \left(\tanh(v_k(x+t)) - \tanh(v_k(x-t)) \right), \quad (13)$$

$$v_k = kv \quad (14)$$

where v_k is the indexed frequency and v is said to be the background frequency.

Remark 2:

Here it is worthwhile to notice the resemblance of the shift vector (13) with Alcubierre's warp drive shift vector [11] when the velocity is constant.

Furthermore, to get some insight into the indexed frequency, v_k (14),

$$v_k = kv,$$

we introduce a regular function $K_t: \Sigma_t \rightarrow \mathbb{R}$. Then a local level surface for each hypersurface Σ_t can be defined using the map

$$K_\kappa = K_t \circ \kappa^{-1}. \quad (15)$$

Hence, in local coordinates we have

$$K_\kappa(t, x, y, z) = C_t \quad (16)$$

where C_t is the height at a given global time t . Consequently we have the local level surface

$$loc(\Sigma_t) = K^{-1}_\kappa(C_t). \quad (17)$$

To continue for a freely falling observer whose worldline is orthogonal to the hypersurfaces then the proper period is $d\tau = dt/k$ when the shift has completed a period and hence when then the height C_t has decreased with one unit. As a plane wave and linear approximation we then can define the indexed frequency by

$$v_k = \partial_\tau K_\kappa = \frac{k}{dt} = kv \quad (18)$$

where

$$v = \frac{1}{dt} \quad (19)$$

is said to be the background frequency.

Next, the class of smooth Quantum foam spacetimes is given by definition 3.

Definition 3 Class of smooth Quantum foam Cosmologies

Let $\{\beta_k\}_{k \in \mathbb{N}}$ be a sequence of Quantum foam shift vector fields as in definition 2 and $\{g_{\beta_k}\}_{k \in \mathbb{N}}$ a sequence of metric fields such that

$$g_{\beta_k} = -N_k^2 dt^2 + \eta_{ij}(dx^i + \beta_k^i dt)(dx^j + \beta_k^j dt) \quad (20)$$

where

$$\eta = \text{diag}(1, 1, 1) \quad (21)$$

and

$$N_k = \frac{1}{k}, k^2 = k_1^2 + k_2^2 + k_3^2, k_1, k_2, k_3 \in \mathbb{N} \quad (22)$$

where N_k is a constant lapse function. Then the sequence

$$\{(\Sigma_{t/k}, g_{\beta_k})\}_{k \in \mathbb{N}} \quad (23)$$

of smooth 3+1 spacetimes is said to be a class of smooth Quantum foam cosmologies and an element $(\Sigma_{t/k}, g_{\beta_k})$ is said to be an Quantum foam Cosmology with a propagation phase number k .

Remark 3:

Here any expansion factor or scale factor a as a function of the global time has been left out from the metric field. Also, any discussion about the initial hypersurface has been deliberately left out until definition 4 below has been given.

Example 3 A 1+1 toy Heaviside Universe

In 1+1 dimension we can use the sigmoid sequence in example 2 to construct a Quantum foam spacetime with an early inflation period but that becomes flat as time goes on:


$$g_{\beta_k} = -\frac{1}{k^2} dt^2 + (dx + \beta_k^1 dt)(dx + \beta_k^1 dt) \quad (24)$$

where from (13)

$$\beta_k^1(x, t) = \frac{1}{2}(\tanh(v_k(x+t)) - \tanh(v_k(x-t))) \quad (25)$$

$$v_k = kv, k \in \mathbb{N}. \quad (26)$$

It is straightforward to extend this example to more realistic 3+1 spacetime with a bump surface that is a compactly supported test function. Also, an illustration of a bumpy shift



vector that gives an idea of a single bubble or foam like structure of the Quantum foam is provided in figure 1 below.

Figure 1. Quantum foam illustration with a bumpy and bubbly shift vector element.

We are now ready for the definition of a 3+1 Quantum foam cosmology at the Planck scale and hence to provide meaning to the notion of the initial hypersurface.

Definition 4 Planck scale 3+1 Quantum foam spacetime

Given the class of Quantum foam spacetimes, then we define an initial Planck scale Quantum foam spacetime by the formal limit

$$(\Sigma_0, g_\beta) = \lim_{k \rightarrow \infty} (\Sigma_{t/k}, g_{\beta_k}). \quad (27)$$

Remark 4:

It is important to notice that there do exist many possible classes of smooth spacetimes that converge to this initial hypersurface and metric field. In terms of measurement this is a statement of the existence of different measurement devices that can be used to measure a physical quantity. Nevertheless, observations in our current universe should uniquely determine a class of spacetimes that fits the observations.

Remark 5:

In definition 4 the limit is as distributions for the shift vector and hence a distribution with zero support and hence it follows from a structure theorem in distribution theory, see [5] or theorem 2.3.4 in [6], that the distribution necessarily is

a linear combination of the Dirac measure and its distributional derivatives. To get a deeper understanding of this limit it then makes sense to construct the discrete measure on each element in the class of Quantum foam spacetimes. In relation to that we also need to define a notion of measure zero on a manifold. We will start with the latter and then construct the measure.

Definition 5 Measure zero in a Hausdorff manifold M with a C^∞ structure

Let O be a subset of M then O is said to be a measure zero in M if for any local coordinate system

$$\kappa_i: O_i \rightarrow U_i \quad (28)$$

the image of $O \cap O_i$ is a measure zero in U_i .

Let us now introduce the class of discrete measures (spaces) for the class of Quantum foam cosmologies in definition 3.

Definition 6 Discrete class of measure spaces for the class of Quantum Cosmologies

Take \mathcal{F}_k to be the power set of $\Sigma_{t/k}$ in the spacetime $(\Sigma_{t/k}, g_{\beta_k})$ constructed from the countable basis. Then this generates the needed σ -algebra. To construct the measure μ_k for $(\Sigma_{t/k}, \mathcal{F}_k)$ take

$$\mu_k(U) = \sqrt{|g_{\beta_k}|} \sum_j m_j \delta_{x_j}(U) = \frac{1}{k} \sum_j m_j \delta_{x_j}(U), U \in \mathcal{F}_k \quad (29)$$

where

$$m_j = \frac{1}{2^j} \quad (30)$$

is the natural mass for a power set and for the atom at x_j with the Dirac measure $\delta_{x_j}(U)$ then $(\Sigma_{t/k}, \mathcal{F}_k, \mu_k)$ is said to be a discrete measure space on the Quantum foam cosmology $(\Sigma_{t/k}, g_{\beta_k})$ and

$$\{(\Sigma_{t/k}, \mathcal{F}_k, \mu_k)\}_{k \in \mathbb{N}} \quad (31)$$

is said to be the class of discrete Quantum foam measure spaces.

Remark 6:

From (29) we immediately see that the discrete measure scales to zero as the propagation number goes to infinity. This scaling to zero is likely to be expected for any Radon measure in this setting. Consequently, the quantum foam that arise out of almost nothing and its evolution can only be measured by a sequence of smooth Quantum foam spacetimes or for that matter a sum-of-sets spacetime that will be introduced next.

Theorem 4 A globally hyperbolic Quantum foam spacetime can be constructed by gluing together the elements in the class of smooth Quantum foam spacetimes.

Proof

Each member in the class of smooth Quantum foam cosmologies in definition 3 is a smooth 3+1 spacetime and hence globally hyperbolic. Use the sequence (23) in definition 3 to create a disjoint union or topological sum of manifolds:

$$M_\sqcup = \bigsqcup_k \Sigma_{t/k} \quad (32)$$

and extend the shift vector $\beta_\sqcup \in C_0^\infty(M_\sqcup)$ to the union by

$$\beta_\sqcup = \sum_k \beta_k \quad (33)$$

for each $\beta_k \in \Sigma_{t/k}$. Next, define a surjection

$$\tilde{\beta}_k: \Sigma_{loc(k)} \subset \Sigma_{t/k} \rightarrow \Sigma_{t/(k+1)} \quad (34)$$

Now, each shift vector is a cut-off function and as such $\beta_k = 1$ on a compact subset $K_k \subset \Sigma_{t/k}$ for any k and hence it is always possible to construct a surjection (38) such that $\tilde{\beta}_k(p)$ for all $p \in \Sigma_{loc(k)}$ are identified and therefore is an equivalence relation. It is here important to understand that when the shift vector equals one then the space is flat on the compact set and hence this makes the continuation of the initial data trivial. Thus, this cuts out a (flat) ball from $\Sigma_{loc(k)}$ and attach it to a region in $\Sigma_{t/(k+1)}$. Repeated throughout the disjoint union this generates a sum-of-sets (adjunction space) and thus a quotient topology. Finally, the metric field is constructed from the extended sum-of-sets shift vector field (42). This completes the gluing and the engineering exercise.

It will now be proved that the Quantum foam cosmology has a period of inflation.

Theorem 5 The Quantum foam cosmology necessarily has an inflation period.

Proof

From the glued spacetime consider the open set $\Sigma_{t/k}$ with the Quantum foam shift vector β_k . Now since the shift vector is constructed from cut-off functions and converge as distributions to a linear distribution it follows that for the proper period $d\tau = (0 + dt)/k$ that $supp(\beta_k) \rightarrow \{0\}$ as $k \rightarrow \infty$ but since β_k is a smooth and compactly supported test function, that is $\beta_k \in C_0^\infty(\Sigma_{t/k})$ it follows that there exist a positive integer $K_0 \in \mathbb{N}$ such that for all $k > K_0$ β_k increases from 0 to 1 (decrease from 1 to 0) more rapidly than any polynomial and hence there exists an inflation period in the proper period

$$d\tau < \frac{dt}{k} \text{ when } k > K_0. \quad (35)$$

This statement can be made more comprehensible by considering the volume element expansion for any Eulerian observer that follows the flow in the cosmology in the very same proper period:

Theorem 6 A proper period exists in the Quantum foam cosmology when the rate of change of the volume element expansion for any Eulerian observer, whose worldline is a geodesic, is more rapid than any polynomial.

Proof

We can without any loss of generality consider an observer moving with a 4-velocity U_k given by

$$U_k^\alpha = n_k^\alpha \quad (36)$$

where n_k is defined as the lapse scaled gradient of the regular function (2) that is

$$n_{k\alpha} = -N_k \nabla_\alpha t = (-N_k, 0, 0, 0) \quad (37)$$

and hence $n_{k\alpha}$ is normal to any hypersurface Σ_t . Consequently, we have

$$n_k^\alpha = \left(\frac{1}{N_k}, -\frac{\beta_k^1}{N_k}, -\frac{\beta_k^2}{N_k}, -\frac{\beta_k^3}{N_k} \right) \quad (38)$$

and therefore the worldline of the observer is orthogonal to any hypersurface. Moreover, the observer is falling freely and hence following the flow of evolution of the universe since the acceleration A_k is zero:

$$\begin{aligned} A_k^\alpha &= U_k^\gamma \nabla_\gamma U_k^\alpha = g^{\alpha\delta} U_k^\gamma \nabla_\gamma (-N_k \nabla_\delta t) \\ &= g^{\alpha\delta} U_k^\gamma \nabla_\delta (-N_k \nabla_\gamma t) = \frac{1}{2} g^{\alpha\delta} \nabla_\delta (U_k^\gamma U_{k\gamma}) = 0. \end{aligned} \quad (39)$$

The lapse of proper time $d\tau$ for any two neighbouring events for the observer can be determined using the metric (20) and that the relative velocity β^i for the Eulerian observer and the lines of constant space coordinates for any k is given by

$$\frac{dx^i}{dt} = -\beta_k^i, \quad i = 1, 2, 3 \quad (40)$$

we then get that

$$d\tau = N_k dt = \frac{dt}{k}. \quad (41)$$

Thus, the proper time for Eulerian observers equals the coordinate time scaled by the constant propagation number. Here we remark that this relation between the proper time and the coordinate time has already been used in the discussion of the indexed frequency (14) without any detailed motivation except an implicit assumption of a freely falling observer.

Next, the volume expansion for the observer, that is a scalar, is given by

$$\theta = \nabla_\alpha U_k^\alpha = -\frac{1}{N_k} \partial_i \beta_k^i. \quad (42)$$

Here and again the last equality follows from that the lapse is constant. Notice also that the expansion is a linear sum of partial derivatives of the shift vector field.

Finally, using distribution theory and similar arguments as in theorem 5 for (42) and thus on the distributional partial derivatives and hence the partial derivative for each direction of the shift vector proves the theorem.

Notice that that there exist a proper period when the rate of change of the trace of the extrinsic curvature K_k is faster than any polynomial. This follows since

$$\text{Tr}(K_k) = -\frac{1}{N_k} \nabla_\alpha U_k^\alpha. \quad (43)$$

Thus we have the following corollary:

Corollary 1 There exists a proper period when the rate of change of the trace of the extrinsic curvature K_k is more rapid than any polynomial.

Remark 7:

At this point it is worthwhile to mention that the results from Alcubierre's warp drive [11] (when the bubble has a constant speed) can be reproduced by this model and one can therefore argue that the bubble in [11] is simply a manifestation of a Quantum foam bubble. In fact any warp drive model that is constructed from compactly supported test functions that is a solution to the wave equation (7) is a representation of the Quantum foam. Furthermore, the compactly supported test functions is dense in the space of rapidly decreasing test functions. The dual to this space is the space of tempered distributions. Thus, by closing the space and hence by including the limit points we include the Alcubierre warp bubble [11]. However, this comes with a price since tempered distributions are somewhat "nicer and less rough" than the distributions on the space of compactly supported test functions, for more details on tempered distributions see [5] or [6].

It will now be proved that the cosmology is homogeneous, isotropic and flat in the low wavenumber limit.

Theorem 7 The Quantum foam cosmology is homogeneous, isotropic and flat in the propagation phase number limit $k_i \rightarrow 1, i = 1, 2, 3$.

Proof

This follows immediately from that the support of the extended shift vector field is a subset of the union of all the supports and hence as time goes on that

$$\beta_k \rightarrow 0 \text{ as } k_i \rightarrow 1, i = 1, 2, 3 \quad (43)$$

and hence that

$$\begin{aligned} g_{\beta_k} &= -\frac{1}{k^2} dt^2 + \eta_{ij} (dx^i + \beta_k^i dt) (dx^j + \beta_k^j dt) \rightarrow \\ g &= -\frac{1}{3} dt^2 + \eta_{ij} dx^i dx^j. \end{aligned} \quad (44)$$

Remark 8:

A consequence of this is that independently of whatever volatility that is exhibited in the extrinsic curvature of the spacetime and in a neighbourhood of the Planck region, it follows that the volatility will average out as time goes on.

Discussion

In this final section I will summarise the developed engineering framework for the Quantum foam cosmology that has been introduced. I would also like to take the opportunity to comment on Carlip's view of the Quantum foam in his discussion of whether the cosmological constant is hidden [9]. Finally, I would also like to point out future directions of research.

This framework has been inspired by the interview with the late Professor Wheeler [4] and his use of the ocean surface analogy to describe the Quantum foam. A description I have interpreted from the perspective of a foliated universe with the ocean surface as the hypersurfaces. I then concluded that it would be reasonable to assume that the shift vector, tangent to the hypersurfaces, should not be an unchanging and static quantity but rather a dynamic entity with wavelike properties and not even necessarily smooth. If this is true then assuming that the shift vector is a fundamental solution to the linear part of the wave equation in a 3+1 spacetime is reasonable since not only do we allow for wiggleness and coarse hypersurfaces but we also maintain time as a parameter. In this study I have avoided working directly with this fundamental solution since it is a distribution with the zero support. Instead I have used the existence of a sequence of well-behaved smooth test functions that converge to the solution as distributions to define a class of smooth spacetimes. This means that each spacetime in the class is a globally hyperbolic spacetime, but it could well be that these spacetimes will stack up and generate a Cauchy horizon as we move closer to the beginning of time. But it is not clear if this really is a problem given the distributional limit and hence that the Planck scale Quantum foam spacetime will be an almost measure zero set since the discrete measure on the spacetime is scaled by the propagation number. To be more precise the discrete measure has a zero limit as the propagation number goes to infinity. Thus the sequence of smooth spacetimes should be a valid way to model the evolution of the universe.

To continue, the framework that has been introduced here, in order to create a model of the Quantum foam, can clearly be criticised to be coordinate dependent given the explicit form of the metric field in the definition of the Quantum foam as the formal limit of the class of smooth spacetimes. Nevertheless, the classic theorem that any globally hyperbolic spacetime, and hence a causally stable spacetime with no closed time like curves, can be foliated by the hyper surfaces of a regular function with a height that is the time, see e.g. [2] for a proof, strongly suggests that the model as such is not void of observable physics. Or to express it slightly differently – if the shift vector is a fundamental solution to the wave equation then there is a wide class of causally reasonable solutions to Einstein's field equations that has a natural inflation mechanism. Furthermore, even if

a choice of coordinates may lead straight into the abyss if we are not careful, we can still make sense of results in coordinates if we choose to interpret the result in the frames that shares some property e.g. that the frequency is very high as we typically do when we construct and interpret the frequency four vector (wave vector) that is the derivative of a scalar function in the hypersurfaces parameterised by time. .

Consequently, I believe one needs to be somewhat cautious with statements of void physics because a selection of a coordinate system is made. Further to this, we need to remember that the existence of a measure zero is a coordinate system independent statement and thus in the convergence limit of the Planck scale Quantum foam with a discrete and single-point atomic measure there do not really exist any coordinate system selection problem; particularly the class of smooth spacetimes presented here should be perfectly valid approximations to the Quantum foam.

Furthermore, I should mention that since having a dynamic shift vector that model's Wheeler's Quantum foam where the dynamic shift cause volatile curvature on the hypersurfaces suggests that Carlip's high level analysis [9] that relies on a zero a shift vector most likely is not sufficient to support a highly volatile cosmological constant at the Planck scale that is smeared as time goes on. Instead, I would argue that his model should be changed to allow for a highly dynamic shift vector that in the beginning and in the direction of time is represented by a Dirac measure. The class of spacetimes that approximates the Planck scale Quantum foam spacetime then provides a resolution to the dilemma with a cosmological constant that is out of range by an order of 120 since if the curvature is highly volatile for phases in the neighbourhood of the Planck scale, it will be smeared as time goes on and the propagation phase number goes to zero.

In summary, this work has hence shown that it is possible to construct a self-consistent Planck scale Quantum foam cosmology where the metric field is wiggly in the sense that the linear vector shift field is a fundamental solution to the linearised wave equation but where linear distribution theory can be used to create a class of spacetimes and hence a sum of sets glued spacetime that represent all the different epochs of the universe. Using this spacetime, it has been proved from distribution theory that the spacetime has an inflation period. It has also been shown that no matter how volatile the extrinsic curvature is in an early universe, it will be averaged out as time goes on. Finally, I am working on the analysis for a quantum field theory for a massless scalar field on an explicit 3+1 model where the fundamental solution is a time-scaled Dirac measure. The motivation behind that study is to understand the effects the volatility has on the scalar field in relation to regions that would be causally disconnected if it were not for the inflation mechanism inherent in the Quantum foam cosmology [16].

References

- [1] Hawking S W, Ellis G F R 1973 The large scale structure of space-time (Cambridge: Cambridge University Press)
- [2] Wald R M 1984 General Relativity (Chicago: Chicago University Press)
- [3] Birrell N D, Davies P C W 1982 Quantum fields in curved space (Cambridge: Cambridge University Press)
- [4] Web of Stories - Life Stories of Remarkable People 2017 Interview with John Wheeler Quantum ideas. Quantum foam. Max Planck and Karl Popper Like a story <https://www.webofstories.com/playAll/john.wheeler?sId=9542>, <https://www.youtube.com/watch?v=iNxioquoK0g>
- [5] Strichartz R S 2003 A Guide to Distribution Theory and Fourier Transforms (World Scientific)
- [6] Hörmander L 1990 The Analysis of Linear Partial Differential Operators I Distribution Theory and Fourier Analysis (Berlin and Heidelberg: Springer-Verlag)
- [7] Chruściel P T, Pollack D, Isenberg J 2004 Gluing Initial Data Sets for General Relativity Phys. Rev. Lett. 93, 081101
- [8] Hobson M P, Efstathiou G P, Lasenby A N 2012 General Relativity An Introduction for Physicists (Cambridge: Cambridge University Press), p187
- [9] Carlip S 2019 Hiding the Cosmological Constant Phys. Rev Lett 123. 13102
- [10] Steinbauer R, Vickers J A 2006 The use of generalized functions and distributions in general relativity Class.Quant.Grav. 23 (2006) R91-R114
- [11] Alcubierre M 1994 The warp drive: hyper-fast travel within general relativity Class.Quant.Grav.11:L73-L77,1994
- [12] Alcubierre, M., Lobo, F.S.N. (2017). Warp Drive Basics. In: Lobo, F. (eds) Wormholes, Warp Drives and Energy Conditions. Fundamental Theories of Physics, vol 189 pp 257-279. Springer, Cham.
- [13] Alcubierre M, Brügmann B, Diener P, Guzmán F. S, Hawke I, Hawley S, Herrmann F, Koppitz M, Pollney D, Seidel E, and Thornburg J 2005 Dynamical evolution of quasicircular binary black hole data Phys. Rev. D 72, 044004
- [14] Alcubierre M 2008 Introduction to 3+1 Numerical Relativity (Oxford: Oxford University Press)
- [15]ourgoulhon E 2012 3+1 Formalism in General Relativity Bases of Numerical Relativity (Berlin and Heidelberg: Springer-Verlag)
- [16] Cramer C 2022 The time-scaled Dirac 3+1 Quantum foam spacetime To be Published